

4.5 Worksheet: The Fundamental Theorem of Calculus

Purpose: The Fundamental Theorem of Calculus can look like it comes out of nowhere. In this worksheet we will try to explain why it is true through examples.

1. Area functions:

- a) Let $f(x)$ be the graph to the right and let $A(x) = \int_0^x f(x) dx$ be its *area function*. For this problem, use geometry to calculate the areas under the curve to compute the following.

$$A(1) = \underline{\hspace{2cm}} \quad A(3) = \underline{\hspace{2cm}}$$


$$A(2) = \underline{\hspace{2cm}} \quad A(4) = \underline{\hspace{2cm}}$$

and, in general, $A(x) = \underline{\hspace{2cm}}$
(hint: think triangle plus rectangle)

Take the derivative to compute

$$A'(x) = \underline{\hspace{2cm}}$$

How does $A'(x)$ compare to $f(x)$?




- b) Let $f(x)$ be the function with the given graph to the right and let $A(x) = \int_0^x f(x) dx$ be its area function. Compute the following.

$$A(1) = \underline{\hspace{2cm}} \quad A'(1) = \underline{\hspace{2cm}}$$

$$A(2) = \underline{\hspace{2cm}} \quad A'(2) = \underline{\hspace{2cm}}$$

$$A(3) = \underline{\hspace{2cm}} \quad A'(3) = \underline{\hspace{2cm}}$$

$$A(4) = \underline{\hspace{2cm}} \quad A'(4) = \underline{\hspace{2cm}}$$



The maximum value of $A(x)$ on the interval $[0, 5]$ is $\underline{\hspace{2cm}}$.

The maximum value of $A'(x)$ on the interval $[0, 5]$ is $\underline{\hspace{2cm}}$.


2. **Velocity and Position:** A toy car is travelling on a stright track. Its velocity $v(t)$, in m/s , is given by the graph to the right. Define $s(t)$ to be the position of the car in meters. Choose coordinates so that $s(0) = 0$. Compute the following.

$$s(2) = \underline{\hspace{2cm}} \qquad v(2) = \underline{\hspace{2cm}}$$

$$s(4) = \underline{\hspace{2cm}} \qquad v(4) = \underline{\hspace{2cm}}$$


$$s(6) = \underline{\hspace{2cm}} \qquad v(6) = \underline{\hspace{2cm}}$$

- (a) What is the maximum value of $s(t)$ on the interval $[0, 7]$?
- (b) What is the minimum value of $s(t)$ on the interval $[0, 7]$?
- (c) What is the maximum value of $v(t)$ on the interval $[0, 7]$?
- (d) What is the minimum value of $v(t)$ on the interval $[0, 7]$?



Sketch a graph of $s(t)$ on the empty graph to the right.

- (a) What is the net change in position from $t = 0$ to $t = 3$?
- (b) What is area under the velocity graph from $t = 0$ to $t = 3$?
- (c) What is the net change in position from $t = 0$ to $t = 7$?
- (d) Use the Fundamental theorem of calculus to compute the integral $\int_0^7 v(t) dt$.



3. **Extra Problems:** Use the Fundamental Theorem of Calculus to compute the definite integrals:

a) $\int_0^2 2x - 3 dx$

b) $\int_0^1 6e^{-3x} + 4 dx$

c) $\int_0^{\pi/4} \sec(t) \tan(t) dt$

d) $\int_0^{1/2} \frac{3}{\sqrt{1-x^2}} dx$