
a
+ sign-flip test (p-value only)
Occasions per cluster
2
3
4 or more
GEE = MMM
usable 10/4%
GEE = MMM
usable 32/19%
GEE = MMM
usable 91/68%
GEE
usable 35/21%
GEE = MMM
usable 69/46%
GEE = MMM
usable 99/91%
GEE
usable 77/59%
GEE = MMM
usable 94/84%
GEE = MMM
usable >99/99%
10
15-25
> 25
Clusters (e.g. patients)
b
Mean absolute departure from 5% (pp), lower = better
LRT
MMM, t
MD, exch., t
FG, AR(1), t
FG, exch., t
Occasions per cluster
c
Design points with false positives clearly above 7.5% (%), lower = better
LRT
MMM, t
MD, exch., t
FG, exch., t
FG, AR(1), t
Clusters (e.g. patients)
d
Mean absolute departure from 5% (pp), lower = better
FG, T = 2
MMM, T ≥ 3
MMM, T = 2
FG, T ≥ 3
Clusters (e.g. patients)
e
Datasets with a usable result (%), higher = better
Sign-flip
FG, AR(1), t
MMM, t
Occasions per cluster
f
Effect detected among attempted datasets (%)
MMM, t
Sign-flip
FG, AR(1), t
Occasions per cluster