

2012.12.09

This report describes data from 2012.12.09 experiment (1 day of D and / or T).

## day 1 body weight (g)

Treatment	Average (SD; n)
V	21.5 (0.393; n = 5)
D	23.6 (0.437; n = 5)
T	22.2 (0.736; n = 5)
C	23.1 (0.502; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.0867

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.287
<b>1</b>	D vs DT	0.01228

Kruskal-Wallis p value for the three-way comparison is 0.0536

	Comparison	P value	Direction
<b>3</b>	V vs D	0.2617	V < D
<b>1</b>	D vs DT	0.009649	D > DT

## day 2 body weight (g)

Treatment	Average (SD; n)
V	22.1 (0.402; n = 5)
D	24.4 (0.395; n = 5)
T	23 (0.645; n = 5)
C	23.7 (0.485; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.0528

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.1747
<b>1</b>	D vs DT	0.004062

Kruskal-Wallis p value for the three-way comparison is 0.0305

	Comparison	P value	Direction
<b>3</b>	V vs D	0.1611	V < D
<b>1</b>	D vs DT	0.004444	D > DT

## body weight gain after 1 days (g)

Treatment	Average (SD; n)
V	0.66 (0.051; n = 5)
D	0.82 (0.26; n = 5)
T	0.86 (0.268; n = 5)
C	0.54 (0.051; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.786

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.1881
<b>1</b>	D vs DT	0.4255

Kruskal-Wallis p value for the three-way comparison is 0.539

	Comparison	P value	Direction
<b>3</b>	V vs D	0.1594	V < D
<b>1</b>	D vs DT	0.4716	D > DT

## body weight gain after 2 days (g)

Treatment	Average (SD; n)
V	3.07 (0.24; n = 5)
D	3.52 (1.12; n = 5)
T	3.97 (1.28; n = 5)
C	2.34 (0.238; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.708

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.168
<b>1</b>	D vs DT	0.5

Kruskal-Wallis p value for the three-way comparison is 0.403

	Comparison	P value	Direction
<b>3</b>	V vs D	0.1289	V < D
<b>1</b>	D vs DT	0.4718	D > DT

levator (mg)

Treatment	Average (SD; n)
V	51.4 (4.28; n = 5)
D	54.1 (3.62; n = 5)
T	54.1 (4.45; n = 5)
C	50.8 (3.95; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.907

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.3344
<b>1</b>	D vs DT	0.3442

Kruskal-Wallis p value for the three-way comparison is 0.854

	Comparison	P value	Direction
<b>3</b>	V vs D	0.3355	V < D
<b>1</b>	D vs DT	0.2978	D > DT

tibialis (mg)

Treatment	Average (SD; n)
V	47.7 (2.37; n = 5)
D	49 (4.62; n = 5)
T	55 (4.31; n = 5)
C	48.3 (5.73; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.386

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.3345
<b>1</b>	D vs DT	0.2606

Kruskal-Wallis p value for the three-way comparison is 0.756

	Comparison	P value	Direction
<b>3</b>	V vs D	0.2858	V < D
<b>1</b>	D vs DT	0.2398	D > DT

## gastrocnemius (mg)

Treatment	Average (SD; n)
V	121 (5.13; n = 5)
D	117 (4.67; n = 5)
T	122 (6.95; n = 5)
C	118 (2.78; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.886

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.4787
<b>1</b>	D vs DT	0.2436

Kruskal-Wallis p value for the three-way comparison is 0.763

	Comparison	P value	Direction
<b>3</b>	V vs D	0.5	V > D
<b>1</b>	D vs DT	0.2623	D < DT

## quadriceps (mg)

Treatment	Average (SD; n)
V	161 (3.91; n = 5)
D	176 (4.62; n = 5)
T	172 (8.89; n = 5)
C	166 (7.8; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.374

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.1308
<b>1</b>	D vs DT	0.04359

Kruskal-Wallis p value for the three-way comparison is 0.151

	Comparison	P value	Direction
<b>3</b>	V vs D	0.1015	V < D
<b>1</b>	D vs DT	0.02812	D > DT

triceps (mg)

Treatment	Average (SD; n)
V	99.1 (7.57; n = 5)
D	99 (2.09; n = 5)
T	104 (3.57; n = 5)
C	100 (4.62; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.708

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.3345
<b>1</b>	D vs DT	0.4574

Kruskal-Wallis p value for the three-way comparison is 0.932

	Comparison	P value	Direction
<b>3</b>	V vs D	0.3886	V > D
<b>1</b>	D vs DT	0.4718	D < DT

levator (permille)

Treatment	Average (SD; n)
V	2.31 (0.157; n = 5)
D	2.22 (0.168; n = 5)
T	2.34 (0.168; n = 5)
C	2.15 (0.168; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.722

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.5
<b>1</b>	D vs DT	0.2436

Kruskal-Wallis p value for the three-way comparison is 0.756

	Comparison	P value	Direction
<b>3</b>	V vs D	0.4438	V > D
<b>1</b>	D vs DT	0.2398	D > DT

## tibialis (permille)

Treatment	Average (SD; n)
V	2.16 (0.138; n = 5)
D	2.01 (0.195; n = 5)
T	2.38 (0.161; n = 5)
C	2.04 (0.243; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.354

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.4153
<b>1</b>	D vs DT	0.2783

Kruskal-Wallis p value for the three-way comparison is 0.827

	Comparison	P value	Direction
<b>3</b>	V vs D	0.4718	V > D
<b>1</b>	D vs DT	0.2858	D < DT

## gastrocnemius (permille)

Treatment	Average (SD; n)
V	5.46 (0.146; n = 5)
D	4.81 (0.171; n = 5)
T	5.31 (0.255; n = 5)
C	5 (0.0876; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.0436

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.2271
<b>1</b>	D vs DT	0.004407

Kruskal-Wallis p value for the three-way comparison is 0.0271

	Comparison	P value	Direction
<b>3</b>	V vs D	0.2183	V > D
<b>1</b>	D vs DT	0.004444	D < DT

## quadriceps (permille)

Treatment	Average (SD; n)
V	7.27 (0.258; n = 5)
D	7.22 (0.165; n = 5)
T	7.46 (0.285; n = 5)
C	7.01 (0.189; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.679

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.1818
<b>1</b>	D vs DT	0.3152

Kruskal-Wallis p value for the three-way comparison is 0.566

	Comparison	P value	Direction
<b>3</b>	V vs D	0.1444	V > D
<b>1</b>	D vs DT	0.3357	D > DT

## triceps (permille)

Treatment	Average (SD; n)
V	4.48 (0.343; n = 5)
D	4.06 (0.0997; n = 5)
T	4.54 (0.174; n = 5)
C	4.22 (0.147; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.277

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.1962
<b>1</b>	D vs DT	0.06724

Kruskal-Wallis p value for the three-way comparison is 0.368

	Comparison	P value	Direction
<b>3</b>	V vs D	0.2398	V > D
<b>1</b>	D vs DT	0.07865	D < DT

fat mass before (g)

Treatment	Average (SD; n)
V	2.78 (0.103; n = 5)
D	2.94 (0.189; n = 5)
T	2.77 (0.104; n = 5)
C	2.24 (0.175; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.0633

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.005148
<b>1</b>	D vs DT	0.2606

Kruskal-Wallis p value for the three-way comparison is 0.0545

	Comparison	P value	Direction
<b>3</b>	V vs D	0.009812	V < D
<b>1</b>	D vs DT	0.2623	D > DT

lean mass before (g)

Treatment	Average (SD; n)
V	17.4 (0.382; n = 5)
D	19 (0.378; n = 5)
T	18.2 (0.655; n = 5)
C	19.6 (0.421; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.031

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.2113
<b>1</b>	D vs DT	0.02398

Kruskal-Wallis p value for the three-way comparison is 0.0185



	Comparison	P value	Direction
<b>3</b>	V vs D	0.1981	V < D
<b>1</b>	D vs DT	0.02812	D < DT

total water before (g)

Treatment	Average (SD; n)
V	14.2 (0.946; n = 5)
D	14.9 (0.337; n = 5)
T	13.8 (0.502; n = 5)
C	14.9 (0.418; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.151

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.3541
<b>1</b>	D vs DT	0.03068

Kruskal-Wallis p value for the three-way comparison is 0.174

	Comparison	P value	Direction
<b>3</b>	V vs D	0.3618	V < D
<b>1</b>	D vs DT	0.03855	D > DT

fat mass after (g)

Treatment	Average (SD; n)
V	2.91 (0.136; n = 5)
D	3.29 (0.18; n = 5)
T	2.85 (0.133; n = 5)
C	2.71 (0.19; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.136

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.01238
<b>1</b>	D vs DT	0.06724

Kruskal-Wallis p value for the three-way comparison is 0.0846

	Comparison	P value	Direction
<b>3</b>	V vs D	0.01419	V < D
<b>1</b>	D vs DT	0.07865	D > DT

lean mass after (g)

Treatment	Average (SD; n)
V	17.6 (0.323; n = 5)
D	18.5 (0.306; n = 5)
T	18.6 (0.546; n = 5)
C	18.7 (0.496; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.169

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.4787
<b>1</b>	D vs DT	0.03068

Kruskal-Wallis p value for the three-way comparison is 0.114

	Comparison	P value	Direction
<b>3</b>	V vs D	0.4718	V < D
<b>1</b>	D vs DT	0.03855	D < DT

total water after (g)

Treatment	Average (SD; n)
V	13.6 (0.488; n = 5)
D	14.8 (0.684; n = 5)
T	15.1 (0.752; n = 5)
C	14.7 (0.703; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.541

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.4787
<b>1</b>	D vs DT	0.1425

Kruskal-Wallis p value for the three-way comparison is 0.468

	Comparison	P value	Direction
<b>3</b>	V vs D	0.4438	V < D
<b>1</b>	D vs DT	0.1611	D > DT

## fat mass before (percent of BW)

Treatment	Average (SD; n)
V	12.9 (0.422; n = 5)
D	12.4 (0.737; n = 5)
T	12.5 (0.418; n = 5)
C	9.67 (0.732; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.0177

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.009339
<b>1</b>	D vs DT	0.2783

Kruskal-Wallis p value for the three-way comparison is 0.0167

	Comparison	P value	Direction
<b>3</b>	V vs D	0.01419	V > D
<b>1</b>	D vs DT	0.3103	D > DT

## lean mass before (percent of BW)

Treatment	Average (SD; n)
V	81 (1.07; n = 5)
D	80.6 (1.85; n = 5)
T	82.2 (1.69; n = 5)
C	84.9 (3.2; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.536

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.1095
<b>1</b>	D vs DT	0.4574

Kruskal-Wallis p value for the three-way comparison is 0.472

	Comparison	P value	Direction
<b>3</b>	V vs D	0.1444	V > D
<b>1</b>	D vs DT	0.5	D < DT

## total water before (percent of BW)

Treatment	Average (SD; n)
V	66.1 (4.72; n = 5)
D	63.4 (1.31; n = 5)
T	62.2 (2.23; n = 5)
C	64.5 (2.89; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.997

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.4153
<b>1</b>	D vs DT	0.4363

Kruskal-Wallis p value for the three-way comparison is 0.932

	Comparison	P value	Direction
<b>3</b>	V vs D	0.3618	V > D
<b>1</b>	D vs DT	0.3886	D < DT

## fat mass after (percent of BW)

Treatment	Average (SD; n)
V	13.2 (0.564; n = 5)
D	13.4 (0.559; n = 5)
T	12.4 (0.495; n = 5)
C	11.5 (0.952; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.166

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.01855
<b>1</b>	D vs DT	0.3345

Kruskal-Wallis p value for the three-way comparison is 0.141

	Comparison	P value	Direction
<b>3</b>	V vs D	0.033	V < D
<b>1</b>	D vs DT	0.3886	D > DT

## lean mass after (percent of BW)

Treatment	Average (SD; n)
V	79.5 (0.591; n = 5)
D	75.7 (0.572; n = 5)
T	80.6 (0.513; n = 5)
C	79 (2.85; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.0423

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.03068
<b>1</b>	D vs DT	0.03068

Kruskal-Wallis p value for the three-way comparison is 0.0872

	Comparison	P value	Direction
<b>3</b>	V vs D	0.033	V > D
<b>1</b>	D vs DT	0.02386	D < DT

## total water after (percent of BW)

Treatment	Average (SD; n)
V	61.5 (1.6; n = 5)
D	60.7 (3.03; n = 5)
T	65.7 (3.58; n = 5)
C	62.3 (4.19; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.741

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.3742
<b>1</b>	D vs DT	0.2606

Kruskal-Wallis p value for the three-way comparison is 0.811

	Comparison	P value	Direction
<b>3</b>	V vs D	0.416	V > D
<b>1</b>	D vs DT	0.2623	D < DT

## fat mass gain (g)

Treatment	Average (SD; n)
V	0.138 (0.0866; n = 5)
D	0.35 (0.108; n = 5)
T	0.0864 (0.0904; n = 5)
C	0.469 (0.271; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.261

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.3541
<b>1</b>	D vs DT	0.1198

Kruskal-Wallis p value for the three-way comparison is 0.196

	Comparison	P value	Direction
<b>3</b>	V vs D	0.2858	V < D
<b>1</b>	D vs DT	0.1147	D < DT

## lean mass gain (g)

Treatment	Average (SD; n)
V	0.197 (0.301; n = 5)
D	-0.505 (0.44; n = 5)
T	0.341 (0.393; n = 5)
C	-0.904 (0.567; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.183

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.5
<b>1</b>	D vs DT	0.07448

Kruskal-Wallis p value for the three-way comparison is 0.34

	Comparison	P value	Direction
<b>3</b>	V vs D	0.5	V > D
<b>1</b>	D vs DT	0.1015	D > DT

total water gain (g)

Treatment	Average (SD; n)
V	-0.542 (1.14; n = 5)
D	-0.126 (0.888; n = 5)
T	1.32 (1.01; n = 5)
C	-0.189 (0.369; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.608

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.3541
<b>1</b>	D vs DT	0.2113

Kruskal-Wallis p value for the three-way comparison is 0.698

	Comparison	P value	Direction
<b>3</b>	V vs D	0.3357	V < D
<b>1</b>	D vs DT	0.1981	D > DT

fat mass gain (percent of BW)

Treatment	Average (SD; n)
V	0.239 (0.365; n = 5)
D	1.01 (0.413; n = 5)
T	-0.105 (0.32; n = 5)
C	1.82 (1.2; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.179

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.2783
<b>1</b>	D vs DT	0.1425

Kruskal-Wallis p value for the three-way comparison is 0.208

	Comparison	P value	Direction
<b>3</b>	V vs D	0.2183	V < D
<b>1</b>	D vs DT	0.1611	D < DT

## lean mass gain (percent of BW)

Treatment	Average (SD; n)
V	-1.52 (1.42; n = 5)
D	-4.85 (1.82; n = 5)
T	-1.63 (2.07; n = 5)
C	-5.82 (2.51; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.488

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.4787
<b>1</b>	D vs DT	0.09977

Kruskal-Wallis p value for the three-way comparison is 0.34

	Comparison	P value	Direction
<b>3</b>	V vs D	0.5	V > D
<b>1</b>	D vs DT	0.1015	D > DT

## total water gain (percent of BW)

Treatment	Average (SD; n)
V	-4.55 (5.32; n = 5)
D	-2.61 (4.06; n = 5)
T	3.43 (4.87; n = 5)
C	-2.15 (1.43; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.7

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.3152
<b>1</b>	D vs DT	0.1962

Kruskal-Wallis p value for the three-way comparison is 0.613



	Comparison	P value	Direction
<b>3</b>	V vs D	0.3103	V < D
<b>1</b>	D vs DT	0.1611	D < DT

## fat mass gain (percent)

Treatment	Average (SD; n)
V	5 (3; n = 5)
D	12.6 (4.17; n = 5)
T	3.17 (3.3; n = 5)
C	24.8 (14.2; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.22

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.2436
<b>1</b>	D vs DT	0.1818

Kruskal-Wallis p value for the three-way comparison is 0.237

	Comparison	P value	Direction
<b>3</b>	V vs D	0.1981	V < D
<b>1</b>	D vs DT	0.1981	D < DT

## lean mass gain (percent)

Treatment	Average (SD; n)
V	1.22 (1.72; n = 5)
D	-2.53 (2.35; n = 5)
T	2.06 (2.35; n = 5)
C	-4.5 (2.78; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.185

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.4574
<b>1</b>	D vs DT	0.06056

Kruskal-Wallis p value for the three-way comparison is 0.281

	Comparison	P value	Direction
<b>3</b>	V vs D	0.4718	V > D
<b>1</b>	D vs DT	0.07865	D > DT

## total water gain (percent)

Treatment	Average (SD; n)
V	-2.18 (6.75; n = 5)
D	-0.463 (6.04; n = 5)
T	10.5 (8.28; n = 5)
C	-1.43 (2.5; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.586

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.3742
<b>1</b>	D vs DT	0.1962

Kruskal-Wallis p value for the three-way comparison is 0.651

	Comparison	P value	Direction
<b>3</b>	V vs D	0.3618	V < D
<b>1</b>	D vs DT	0.179	D > DT

## quadriceps calpain activity (RU)

Treatment	Average (SD; n)
V	31500 (5720; n = 5)
D	20000 (3770; n = 5)
T	29900 (3590; n = 5)
C	19400 (2610; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.182

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.4153
<b>1</b>	D vs DT	0.0823

Kruskal-Wallis p value for the three-way comparison is 0.336

	Comparison	P value	Direction
<b>3</b>	V vs D	0.4438	V > D
<b>1</b>	D vs DT	0.1147	D > DT

## gastrocnemius calpain activity (RU)

Treatment	Average (SD; n)
V	30300 (2530; n = 5)
D	18300 (2190; n = 5)
T	26700 (1570; n = 5)
C	18300 (2190; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.01

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.5
<b>1</b>	D vs DT	0.004698

Kruskal-Wallis p value for the three-way comparison is 0.0355

	Comparison	P value	Direction
<b>3</b>	V vs D	0.5	V > D
<b>1</b>	D vs DT	0.01263	D < DT

## gastrocnemius proteasome activity (RU)

Treatment	Average (SD; n)
V	6890 (949; n = 5)
D	5140 (380; n = 5)
T	12400 (1210; n = 5)
C	7550 (605; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.00346

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.01855
<b>1</b>	D vs DT	0.07448

Kruskal-Wallis p value for the three-way comparison is 0.024

	Comparison	P value	Direction
<b>3</b>	V vs D	0.003605	V > D
<b>1</b>	D vs DT	0.03855	D < DT

## quadriceps proteasome activity (RU)

Treatment	Average (SD; n)
V	10500 (1000; n = 5)
D	7620 (1380; n = 5)
T	12500 (1660; n = 5)
C	7630 (1250; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.0621

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.4153
<b>1</b>	D vs DT	0.04359

Kruskal-Wallis p value for the three-way comparison is 0.141

	Comparison	P value	Direction
<b>3</b>	V vs D	0.3886	V > D
<b>1</b>	D vs DT	0.033	D < DT

## triceps proteasome activity (RU)

Treatment	Average (SD; n)
V	15200 (1360; n = 5)
D	12000 (698; n = 5)
T	14300 (1110; n = 5)
C	10200 (1430; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.169

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.2582
<b>1</b>	D vs DT	0.08003

Kruskal-Wallis p value for the three-way comparison is 0.143

	Comparison	P value	Direction
<b>3</b>	V vs D	0.2113	V > D
<b>1</b>	D vs DT	0.1132	D > DT

## gastrocnemius cathepsin activity (RU)

Treatment	Average (SD; n)
V	8400 (290; n = 5)
D	7020 (239; n = 5)
T	8690 (261; n = 5)
C	7400 (400; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.0114

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.2113
<b>1</b>	D vs DT	0.009339

Kruskal-Wallis p value for the three-way comparison is 0.0437

	Comparison	P value	Direction
<b>3</b>	V vs D	0.179	V > D
<b>1</b>	D vs DT	0.006664	D < DT

## quadriceps cathepsin activity (RU)

Treatment	Average (SD; n)
V	39700 (1380; n = 5)
D	33900 (1140; n = 5)
T	42500 (1930; n = 5)
C	40400 (1440; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.0255

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.01421
<b>1</b>	D vs DT	0.01855

Kruskal-Wallis p value for the three-way comparison is 0.0265

	Comparison	P value	Direction
<b>3</b>	V vs D	0.009812	V > D
<b>1</b>	D vs DT	0.009812	D < DT

## triceps cathepsin activity (RU)

Treatment	Average (SD; n)
V	35500 (1650; n = 5)
D	26000 (1120; n = 5)
T	34400 (1290; n = 5)
C	31100 (1070; n = 5)

Kruskal-Wallis p value for the four-way comparison is 0.00565

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.04359
<b>1</b>	D vs DT	0.0008061

Kruskal-Wallis p value for the three-way comparison is 0.00592

	Comparison	P value	Direction
<b>3</b>	V vs D	0.02812	V > D
<b>1</b>	D vs DT	0.0007314	D < DT

## triceps calpain activity (RU)

Treatment	Average (SD; n)
V	11600 (1310; n = 5)
D	5430 (1000; n = 5)
T	13500 (870; n = 4)
C	3370 (606; n = 3)

Kruskal-Wallis p value for the four-way comparison is 0.00451

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.2576
<b>1</b>	D vs DT	0.02254

Kruskal-Wallis p value for the three-way comparison is 0.00964

	Comparison	P value	Direction
<b>3</b>	V vs D	0.1994	V > D
<b>1</b>	D vs DT	0.0115	D > DT

## gastrocnemius Ct(Becn1)- Ct(Gapdh)

Treatment	Average (SD; n)
V	5.74 (0.323; n = 4)
D	6.6 (0.454; n = 4)
T	5.64 (0.336; n = 4)
C	6.06 (0.443; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.314

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.1672
<b>1</b>	D vs DT	0.03735

Kruskal-Wallis p value for the three-way comparison is 0.174

	Comparison	P value	Direction
<b>3</b>	V vs D	0.1404	V < D
<b>1</b>	D vs DT	0.03122	D > DT

## gastrocnemius Ct(Bnip3)- Ct(Gapdh)

Treatment	Average (SD; n)
V	0.89 (0.297; n = 4)
D	1.05 (0.474; n = 4)
T	0.586 (0.516; n = 4)
C	0.575 (0.533; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.872

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.2762
<b>1</b>	D vs DT	0.5

Kruskal-Wallis p value for the three-way comparison is 0.874

	Comparison	P value	Direction
<b>3</b>	V vs D	0.312	V < D
<b>1</b>	D vs DT	0.4609	D > DT

### gastrocnemius Ct(Ctst)- Ct(Gapdh)

Treatment	Average (SD; n)
V	5.11 (0.246; n = 4)
D	5.16 (0.373; n = 4)
T	4.29 (0.282; n = 4)
C	4.88 (0.441; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.303

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.252
<b>1</b>	D vs DT	0.441

Kruskal-Wallis p value for the three-way comparison is 0.694

	Comparison	P value	Direction
<b>3</b>	V vs D	0.2462	V < D
<b>1</b>	D vs DT	0.4609	D > DT

### gastrocnemius Ct(Ddit4)- Ct(Gapdh)

Treatment	Average (SD; n)
V	6.88 (0.614; n = 4)
D	3.98 (0.608; n = 4)
T	6.82 (0.202; n = 4)
C	3.97 (0.256; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.01

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.441
<b>1</b>	D vs DT	0.01295

Kruskal-Wallis p value for the three-way comparison is 0.0244



	Comparison	P value	Direction
<b>3</b>	V vs D	0.4223	V > D
<b>1</b>	D vs DT	0.01206	D > DT

### gastrocnemius Ct(Fbxo32)- Ct(Gapdh)

Treatment	Average (SD; n)
V	2.29 (0.525; n = 4)
D	0.992 (0.389; n = 4)
T	2.88 (0.27; n = 4)
C	0.675 (0.184; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.0168

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.3016
<b>1</b>	D vs DT	0.06874

Kruskal-Wallis p value for the three-way comparison is 0.069

	Comparison	P value	Direction
<b>3</b>	V vs D	0.2462	V > D
<b>1</b>	D vs DT	0.05833	D > DT

### gastrocnemius Ct(Foxo1)- Ct(Gapdh)

Treatment	Average (SD; n)
V	8 (0.4; n = 4)
D	6.61 (0.537; n = 4)
T	8.28 (0.167; n = 4)
C	6.5 (0.332; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.021

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.4704
<b>1</b>	D vs DT	0.03735

Kruskal-Wallis p value for the three-way comparison is 0.087

	Comparison	P value	Direction
<b>3</b>	V vs D	0.4609	V > D
<b>1</b>	D vs DT	0.03122	D > DT

### gastrocnemius Ct(Foxo3a)- Ct(Gapdh)

Treatment	Average (SD; n)
V	8.83 (0.529; n = 4)
D	8.54 (0.625; n = 4)
T	9.79 (0.215; n = 4)
C	8.4 (0.302; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.201

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.441
<b>1</b>	D vs DT	0.328

Kruskal-Wallis p value for the three-way comparison is 0.874

	Comparison	P value	Direction
<b>3</b>	V vs D	0.4609	V > D
<b>1</b>	D vs DT	0.312	D > DT

### gastrocnemius Ct(Foxo4)- Ct(Gapdh)

Treatment	Average (SD; n)
V	5.44 (0.327; n = 4)
D	6.59 (0.551; n = 4)
T	6.34 (0.234; n = 4)
C	5.88 (0.35; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.306

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.2289
<b>1</b>	D vs DT	0.05116

Kruskal-Wallis p value for the three-way comparison is 0.292

	Comparison	P value	Direction
<b>3</b>	V vs D	0.2164	V < D
<b>1</b>	D vs DT	0.05833	D > DT

## gastrocnemius Ct(Igf1)- Ct(Gapdh)

Treatment	Average (SD; n)
V	7.27 (0.418; n = 4)
D	9.38 (0.533; n = 4)
T	6.96 (0.389; n = 4)
C	8.23 (0.396; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.0296

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.1672
<b>1</b>	D vs DT	0.005787

Kruskal-Wallis p value for the three-way comparison is 0.0488

	Comparison	P value	Direction
<b>3</b>	V vs D	0.1404	V < D
<b>1</b>	D vs DT	0.007114	D > DT

## gastrocnemius Ct(Igf1r)- Ct(Gapdh)

Treatment	Average (SD; n)
V	7.86 (0.439; n = 4)
D	8.84 (0.448; n = 4)
T	8.66 (0.235; n = 4)
C	8.42 (0.255; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.235

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.1672
<b>1</b>	D vs DT	0.02675

Kruskal-Wallis p value for the three-way comparison is 0.146

	Comparison	P value	Direction
<b>3</b>	V vs D	0.1634	V < D
<b>1</b>	D vs DT	0.02493	D > DT

## gastrocnemius Ct(Klf15)- Ct(Gapdh)

Treatment	Average (SD; n)
V	6.88 (0.405; n = 4)
D	6.9 (0.376; n = 4)
T	7.86 (0.188; n = 4)
C	7.09 (0.251; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.203

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.2762
<b>1</b>	D vs DT	0.3552

Kruskal-Wallis p value for the three-way comparison is 0.735

	Comparison	P value	Direction
<b>3</b>	V vs D	0.2164	V < D
<b>1</b>	D vs DT	0.3474	D < DT

## gastrocnemius Ct(Map1lc3b)- Ct(Gapdh)

Treatment	Average (SD; n)
V	3.19 (0.362; n = 4)
D	4 (0.486; n = 4)
T	3.34 (0.204; n = 4)
C	3.89 (0.424; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.239

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.3832
<b>1</b>	D vs DT	0.03169

Kruskal-Wallis p value for the three-way comparison is 0.167

	Comparison	P value	Direction
<b>3</b>	V vs D	0.3843	V < D
<b>1</b>	D vs DT	0.03878	D > DT

## gastrocnemius Ct(Nr3c1)- Ct(Gapdh)

Treatment	Average (SD; n)
V	3.67 (0.242; n = 4)
D	5.87 (0.483; n = 4)
T	5.06 (0.257; n = 4)
C	5.73 (0.57; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.0308

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.3552
<b>1</b>	D vs DT	0.003001

Kruskal-Wallis p value for the three-way comparison is 0.0373

	Comparison	P value	Direction
<b>3</b>	V vs D	0.3843	V < D
<b>1</b>	D vs DT	0.009301	D > DT

## gastrocnemius Ct(Odc)- Ct(Gapdh)

Treatment	Average (SD; n)
V	0.854 (0.189; n = 4)
D	0.165 (0.395; n = 4)
T	0.453 (0.478; n = 4)
C	-1.37 (0.617; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.0667

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.04382
<b>1</b>	D vs DT	0.207

Kruskal-Wallis p value for the three-way comparison is 0.0435

	Comparison	P value	Direction
<b>3</b>	V vs D	0.04776	V > D
<b>1</b>	D vs DT	0.2164	D > DT

### gastrocnemius Ct(Stk11)- Ct(Gapdh)

Treatment	Average (SD; n)
V	4.57 (0.333; n = 4)
D	5.39 (0.485; n = 4)
T	5.5 (0.23; n = 4)
C	5.15 (0.443; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.292

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.3016
<b>1</b>	D vs DT	0.05116

Kruskal-Wallis p value for the three-way comparison is 0.368

	Comparison	P value	Direction
<b>3</b>	V vs D	0.3474	V < D
<b>1</b>	D vs DT	0.08491	D > DT

### gastrocnemius Ct(Trim63)- Ct(Gapdh)

Treatment	Average (SD; n)
V	1.2 (0.306; n = 4)
D	0.274 (0.256; n = 4)
T	1.16 (0.4; n = 4)
C	0.154 (0.371; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.119

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.4704
<b>1</b>	D vs DT	0.03735

Kruskal-Wallis p value for the three-way comparison is 0.125

	Comparison	P value	Direction
<b>3</b>	V vs D	0.5	V > D
<b>1</b>	D vs DT	0.03878	D > DT

### quadriceps Ct(Becn1)- Ct(Gapdh)

Treatment	Average (SD; n)
V	7.79 (0.248; n = 4)
D	3.62 (1.11; n = 4)
T	7.58 (0.0748; n = 4)
C	3.19 (0.343; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.00835

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.3832
<b>1</b>	D vs DT	0.002387

Kruskal-Wallis p value for the three-way comparison is 0.0231

	Comparison	P value	Direction
<b>3</b>	V vs D	0.3474	V > D
<b>1</b>	D vs DT	0.005394	D > DT

### quadriceps Ct(Bnip3)- Ct(Gapdh)

Treatment	Average (SD; n)
V	3.56 (0.302; n = 4)
D	0.444 (0.611; n = 4)
T	3.38 (0.198; n = 4)
C	2.21 (0.0899; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.0068

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.1864
<b>1</b>	D vs DT	0.001487

Kruskal-Wallis p value for the three-way comparison is 0.0125

	Comparison	P value	Direction
<b>3</b>	V vs D	0.1197	V > D
<b>1</b>	D vs DT	0.001632	D < DT

### quadriceps Ct(Ctstl)- Ct(Gapdh)

Treatment	Average (SD; n)
V	4.7 (0.232; n = 4)
D	1.32 (0.561; n = 4)
T	4.13 (0.212; n = 4)
C	0.761 (0.418; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.00723

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.328
<b>1</b>	D vs DT	0.005787

Kruskal-Wallis p value for the three-way comparison is 0.021

	Comparison	P value	Direction
<b>3</b>	V vs D	0.2781	V > D
<b>1</b>	D vs DT	0.01974	D > DT

### quadriceps Ct(Ddit4)- Ct(Gapdh)

Treatment	Average (SD; n)
V	8.62 (0.248; n = 4)
D	0.32 (1.33; n = 4)
T	7.74 (0.549; n = 4)
C	0.712 (0.236; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.00673

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.2762
<b>1</b>	D vs DT	0.001165

Kruskal-Wallis p value for the three-way comparison is 0.0183



	Comparison	P value	Direction
<b>3</b>	V vs D	0.2164	V > D
<b>1</b>	D vs DT	0.00302	D < DT

### quadriceps Ct(Fbxo32)- Ct(Gapdh)

Treatment	Average (SD; n)
V	5.14 (0.333; n = 4)
D	-0.51 (0.967; n = 4)
T	5.87 (0.238; n = 4)
C	-0.0944 (0.155; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.00673

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.2762
<b>1</b>	D vs DT	0.01066

Kruskal-Wallis p value for the three-way comparison is 0.0183

	Comparison	P value	Direction
<b>3</b>	V vs D	0.2164	V > D
<b>1</b>	D vs DT	0.00302	D < DT

### quadriceps Ct(Foxo1)- Ct(Gapdh)

Treatment	Average (SD; n)
V	9.34 (0.283; n = 4)
D	3.02 (1.25; n = 4)
T	8.76 (0.408; n = 4)
C	3.2 (0.229; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.00738

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.2762
<b>1</b>	D vs DT	0.001487

Kruskal-Wallis p value for the three-way comparison is 0.0183

	Comparison	P value	Direction
<b>3</b>	V vs D	0.2164	V > D
<b>1</b>	D vs DT	0.00302	D < DT

### quadriceps Ct(Foxo3a)- Ct(Gapdh)

Treatment	Average (SD; n)
V	10.7 (0.389; n = 4)
D	4.37 (1.37; n = 4)
T	10.3 (0.254; n = 4)
C	4 (0.252; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.00835

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.2762
<b>1</b>	D vs DT	0.002387

Kruskal-Wallis p value for the three-way comparison is 0.0183

	Comparison	P value	Direction
<b>3</b>	V vs D	0.2164	V > D
<b>1</b>	D vs DT	0.00302	D > DT

### quadriceps Ct(Foxo4)- Ct(Gapdh)

Treatment	Average (SD; n)
V	7.36 (0.252; n = 4)
D	3.48 (1.17; n = 4)
T	7.42 (0.289; n = 4)
C	2.81 (0.208; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.0183

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.441
<b>1</b>	D vs DT	0.01564

Kruskal-Wallis p value for the three-way comparison is 0.0373

	Comparison	P value	Direction
<b>3</b>	V vs D	0.3843	V > D
<b>1</b>	D vs DT	0.009301	D > DT

## quadriceps Ct(Igf1)- Ct(Gapdh)

Treatment	Average (SD; n)
V	9.61 (0.467; n = 4)
D	8.79 (0.561; n = 4)
T	8.52 (0.048; n = 4)
C	8.2 (0.232; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.0795

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.3552
<b>1</b>	D vs DT	0.02248

Kruskal-Wallis p value for the three-way comparison is 0.069

	Comparison	P value	Direction
<b>3</b>	V vs D	0.2462	V > D
<b>1</b>	D vs DT	0.05833	D > DT

## quadriceps Ct(Igf1r)- Ct(Gapdh)

Treatment	Average (SD; n)
V	9.15 (0.326; n = 4)
D	5.26 (1.24; n = 4)
T	9.19 (0.413; n = 4)
C	4.82 (0.329; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.0327

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.3832
<b>1</b>	D vs DT	0.01564

Kruskal-Wallis p value for the three-way comparison is 0.0498

	Comparison	P value	Direction
<b>3</b>	V vs D	0.2781	V > D
<b>1</b>	D vs DT	0.009301	D > DT

### quadriceps Ct(Klf15)- Ct(Gapdh)

Treatment	Average (SD; n)
V	7.4 (0.254; n = 4)
D	3.6 (0.808; n = 4)
T	8.02 (0.343; n = 4)
C	3.96 (0.153; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.00673

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.2762
<b>1</b>	D vs DT	0.01066

Kruskal-Wallis p value for the three-way comparison is 0.0183

	Comparison	P value	Direction
<b>3</b>	V vs D	0.2164	V > D
<b>1</b>	D vs DT	0.00302	D < DT

### quadriceps Ct(Map1lc3b)- Ct(Gapdh)

Treatment	Average (SD; n)
V	3.16 (0.273; n = 4)
D	-0.376 (0.677; n = 4)
T	2.84 (0.199; n = 4)
C	-0.85 (0.352; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.00861

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.441
<b>1</b>	D vs DT	0.004673

Kruskal-Wallis p value for the three-way comparison is 0.0244

	Comparison	P value	Direction
<b>3</b>	V vs D	0.4223	V > D
<b>1</b>	D vs DT	0.01206	D > DT

### quadriceps Ct(Nr3c1)- Ct(Gapdh)

Treatment	Average (SD; n)
V	3.92 (0.335; n = 4)
D	2.03 (0.62; n = 4)
T	3.55 (0.107; n = 4)
C	2.04 (0.186; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.0496

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.441
<b>1</b>	D vs DT	0.01066

Kruskal-Wallis p value for the three-way comparison is 0.0498

	Comparison	P value	Direction
<b>3</b>	V vs D	0.2781	V > D
<b>1</b>	D vs DT	0.009301	D < DT

### quadriceps Ct(Odc)- Ct(Gapdh)

Treatment	Average (SD; n)
V	3.18 (0.168; n = 4)
D	-0.829 (0.712; n = 4)
T	2.43 (0.11; n = 4)
C	-2.57 (0.725; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.00411

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.2289
<b>1</b>	D vs DT	0.004673

Kruskal-Wallis p value for the three-way comparison is 0.0154

	Comparison	P value	Direction
<b>3</b>	V vs D	0.1634	V > D
<b>1</b>	D vs DT	0.03122	D > DT

## quadriceps Ct(Stk11)- Ct(Gapdh)

Treatment	Average (SD; n)
V	7.29 (0.302; n = 4)
D	3.9 (0.948; n = 4)
T	7.25 (0.384; n = 4)
C	3.29 (0.243; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.0184

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.5
<b>1</b>	D vs DT	0.008742

Kruskal-Wallis p value for the three-way comparison is 0.0388

	Comparison	P value	Direction
<b>3</b>	V vs D	0.4609	V > D
<b>1</b>	D vs DT	0.01206	D > DT

## quadriceps Ct(Trim63)- Ct(Gapdh)

Treatment	Average (SD; n)
V	6.25 (0.347; n = 4)
D	0.307 (0.91; n = 4)
T	6.08 (0.382; n = 4)
C	0.795 (0.187; n = 4)

Kruskal-Wallis p value for the four-way comparison is 0.00835

	contrastsfour	dunns.P
<b>5</b>	V vs D	0.2762
<b>1</b>	D vs DT	0.002387

Kruskal-Wallis p value for the three-way comparison is 0.0183

	Comparison	P value	Direction
<b>3</b>	V vs D	0.2164	$V > D$
<b>1</b>	D vs DT	0.00302	$D < DT$