

# Three Body Equation

Let  $\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3 \in \mathbb{R}^3$  be the positions of the three bodies, with masses  $m_1, m_2, m_3$  where  $m_1 \geq m_2 \geq m_3$ .

## Three Body Equation

Let  $\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3 \in \mathbb{R}^3$  be the positions of the three bodies, with masses  $m_1, m_2, m_3$  where  $m_1 \geq m_2 \geq m_3$ .

$$\ddot{\mathbf{r}}_1 = G \left( \frac{m_2}{|\mathbf{r}_{12}|^3} \mathbf{r}_{12} + \frac{m_3}{|\mathbf{r}_{13}|^3} \mathbf{r}_{13} \right)$$

$\mathbf{r}_{ij} = \mathbf{r}_j - \mathbf{r}_i$  is the vector from body  $i$  to body  $j$ .

## Three Body Equation

Let  $\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3 \in \mathbb{R}^3$  be the positions of the of the three bodies, with masses  $m_1, m_2, m_3$  where  $m_1 \geq m_2 \geq m_3$ .

$$\begin{aligned}\ddot{\mathbf{r}}_1 &= G \left( \frac{m_2}{|\mathbf{r}_{12}|^3} \mathbf{r}_{12} + \frac{m_3}{|\mathbf{r}_{13}|^3} \mathbf{r}_{13} \right) \\ \ddot{\mathbf{r}}_2 &= G \left( \frac{m_1}{|\mathbf{r}_{21}|^3} \mathbf{r}_{21} + \frac{m_3}{|\mathbf{r}_{23}|^3} \mathbf{r}_{23} \right)\end{aligned}$$

$\mathbf{r}_{ij} = \mathbf{r}_j - \mathbf{r}_i$  is the vector from body  $i$  to body  $j$ .

## Three Body Equation

Let  $\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3 \in \mathbb{R}^3$  be the positions of the of the three bodies, with masses  $m_1, m_2, m_3$  where  $m_1 \geq m_2 \geq m_3$ .

$$\ddot{\mathbf{r}}_1 = G \left( \frac{m_2}{|\mathbf{r}_{12}|^3} \mathbf{r}_{12} + \frac{m_3}{|\mathbf{r}_{13}|^3} \mathbf{r}_{13} \right)$$

$$\ddot{\mathbf{r}}_2 = G \left( \frac{m_1}{|\mathbf{r}_{21}|^3} \mathbf{r}_{21} + \frac{m_3}{|\mathbf{r}_{23}|^3} \mathbf{r}_{23} \right)$$

$$\ddot{\mathbf{r}}_3 = G \left( \frac{m_1}{|\mathbf{r}_{31}|^3} \mathbf{r}_{31} + \frac{m_2}{|\mathbf{r}_{32}|^3} \mathbf{r}_{32} \right)$$

$\mathbf{r}_{ij} = \mathbf{r}_j - \mathbf{r}_i$  is the vector from body  $i$  to body  $j$ .

# Restriction

## Restriction

Assume that  $m_3 \ll m_2$ .

## Restriction

Assume that  $m_3 \ll m_2$ .

$$\ddot{\mathbf{r}}_1 = G \frac{m_2}{|\mathbf{r}_{12}|^3} \mathbf{r}_{12}$$

$$\ddot{\mathbf{r}}_2 = G \frac{m_1}{|\mathbf{r}_{21}|^3} \mathbf{r}_{21}$$

## Restriction

Assume that  $m_3 \ll m_2$ .

$$\ddot{\mathbf{r}}_1 = G \frac{m_2}{|\mathbf{r}_{12}|^3} \mathbf{r}_{12}$$

$$\ddot{\mathbf{r}}_2 = G \frac{m_1}{|\mathbf{r}_{21}|^3} \mathbf{r}_{21}$$

$$\ddot{\mathbf{r}}_3 = G \left( \frac{m_1}{|\mathbf{r}_{31}|^3} \mathbf{r}_{31} + \frac{m_2}{|\mathbf{r}_{32}|^3} \mathbf{r}_{32} \right)$$



# Simplifying

# Simplifying

- XY-Plane be the plane where  $m_1$  and  $m_2$  orbit.

# Simplifying

- XY-Plane be the plane where  $m_1$  and  $m_2$  orbit.
- Let the origin be the center of mass at  $t = 0$

## Simplifying

- XY-Plane be the plane where  $m_1$  and  $m_2$  orbit.
- Let the origin be the center of mass at  $t = 0$

- $\mathbf{r}_1 = \begin{bmatrix} -x_1 \\ 0 \\ 0 \end{bmatrix}$

## Simplifying

- XY-Plane be the plane where  $m_1$  and  $m_2$  orbit.
- Let the origin be the center of mass at  $t = 0$

- $\mathbf{r}_1 = \begin{bmatrix} -x_1 \\ 0 \\ 0 \end{bmatrix}, \mathbf{r}_2 = \begin{bmatrix} x_2 \\ 0 \\ 0 \end{bmatrix}$  for  $x_1, x_2 \geq 0$ .

# Changing Frame

# Changing Frame

Define  $R_\omega : \mathbb{R}^3 \times \mathbb{R} \rightarrow \mathbb{R}^3$

## Changing Frame

Define  $R_\omega : \mathbb{R}^3 \times \mathbb{R} \rightarrow \mathbb{R}^3$

$$R_\omega(t) = \begin{pmatrix} \cos(\omega t) & \sin(\omega t) & 0 \\ -\sin(\omega t) & \cos(\omega t) & 0 \\ 0 & 0 & 1 \end{pmatrix}$$



## Changing Frame

Define  $R_\omega : \mathbb{R}^3 \times \mathbb{R} \rightarrow \mathbb{R}^3$

$$R_\omega(t) = \begin{pmatrix} \cos(\omega t) & \sin(\omega t) & 0 \\ -\sin(\omega t) & \cos(\omega t) & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$R_\omega^{-1}(t) = \begin{pmatrix} \cos(\omega t) & -\sin(\omega t) & 0 \\ \sin(\omega t) & \cos(\omega t) & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

## Changing Frame 2

## Changing Frame 2

- $\mathbf{p} = \begin{bmatrix} x \\ y \\ z \end{bmatrix} = R_{\omega}(t) \begin{bmatrix} r_3^1 \\ r_3^2 \\ r_3^3 \end{bmatrix} = R_{\omega}(t) \mathbf{r}_3$

## Changing Frame 2

- $\mathbf{p} = \begin{bmatrix} x \\ y \\ z \end{bmatrix} = R_\omega(t) \begin{bmatrix} r_3^1 \\ r_3^2 \\ r_3^3 \end{bmatrix} = R_\omega(t) \mathbf{r}_3$
- $\dot{\mathbf{r}}_3 = \dot{R}_\omega^{-1}(t) \mathbf{p} + R_\omega^{-1}(t) \dot{\mathbf{p}}$

## Changing Frame 2

- $\mathbf{p} = \begin{bmatrix} x \\ y \\ z \end{bmatrix} = R_\omega(t) \begin{bmatrix} r_3^1 \\ r_3^2 \\ r_3^3 \end{bmatrix} = R_\omega(t) \mathbf{r}_3$
- $\dot{\mathbf{r}}_3 = \dot{R}_\omega^{-1}(t) \mathbf{p} + R_\omega^{-1}(t) \dot{\mathbf{p}}$
- $\ddot{\mathbf{r}}_3 = \ddot{R}_\omega^{-1}(t) \mathbf{p} + 2\dot{R}_\omega^{-1}(t) \dot{\mathbf{p}} + R_\omega^{-1}(t) \ddot{\mathbf{p}}$

## Changing Frame 2

- $\mathbf{p} = \begin{bmatrix} x \\ y \\ z \end{bmatrix} = R_\omega(t) \begin{bmatrix} r_3^1 \\ r_3^2 \\ r_3^3 \end{bmatrix} = R_\omega(t) \mathbf{r}_3$
- $\dot{\mathbf{r}}_3 = \dot{R}_\omega^{-1}(t) \mathbf{p} + R_\omega^{-1}(t) \dot{\mathbf{p}}$
- $\ddot{\mathbf{r}}_3 = \ddot{R}_\omega^{-1}(t) \mathbf{p} + 2\dot{R}_\omega^{-1}(t) \dot{\mathbf{p}} + R_\omega^{-1}(t) \ddot{\mathbf{p}}$
- $|\mathbf{r}_{ij}| = |\mathbf{r}_j - \mathbf{r}_i| = |\mathbf{p}_j - \mathbf{p}_i| = |\mathbf{p}_{ij}|$

# New Equation

## New Equation

$$\ddot{\mathbf{r}}_3 = G \left( \frac{m_1}{|\mathbf{r}_{31}|^3} \mathbf{r}_{31} + \frac{m_2}{|\mathbf{r}_{32}|^3} \mathbf{r}_{32} \right)$$



## New Equation

$$\ddot{\mathbf{r}}_3 = G \left( \frac{m_1}{|\mathbf{r}_{31}|^3} \mathbf{r}_{31} + \frac{m_2}{|\mathbf{r}_{32}|^3} \mathbf{r}_{32} \right)$$

$$\ddot{R}_\omega^{-1}(t)\mathbf{p} + 2\dot{R}_\omega^{-1}(t)\dot{\mathbf{p}} + R_\omega^{-1}(t)\ddot{\mathbf{p}}$$

## New Equation

$$\ddot{\mathbf{r}}_3 = G \left( \frac{m_1}{|\mathbf{r}_{31}|^3} \mathbf{r}_{31} + \frac{m_2}{|\mathbf{r}_{32}|^3} \mathbf{r}_{32} \right)$$

$$\ddot{R}_\omega^{-1}(t) \mathbf{p} + 2\dot{R}_\omega^{-1}(t) \dot{\mathbf{p}} + R_\omega^{-1}(t) \ddot{\mathbf{p}} = GR_\omega^{-1}(t) \left( \frac{m_1}{|\mathbf{p}_{31}|^3} \mathbf{p}_{31} + \frac{m_2}{|\mathbf{p}_{32}|^3} \mathbf{p}_{32} \right)$$

## New Equation

$$\ddot{\mathbf{r}}_3 = G \left( \frac{m_1}{|\mathbf{r}_{31}|^3} \mathbf{r}_{31} + \frac{m_2}{|\mathbf{r}_{32}|^3} \mathbf{r}_{32} \right)$$

$$\ddot{R}_\omega^{-1}(t) \mathbf{p} + 2\dot{R}_\omega^{-1}(t) \dot{\mathbf{p}} + R_\omega^{-1}(t) \ddot{\mathbf{p}} = GR_\omega^{-1}(t) \left( \frac{m_1}{|\mathbf{p}_{31}|^3} \mathbf{p}_{31} + \frac{m_2}{|\mathbf{p}_{32}|^3} \mathbf{p}_{32} \right)$$

$$\ddot{\mathbf{p}} = -R_\omega(t) \ddot{R}_\omega^{-1}(t) \mathbf{p} - 2R_\omega(t) \dot{R}_\omega^{-1}(t) \dot{\mathbf{p}} + G \left( \frac{m_1}{|\mathbf{p}_{31}|^3} \mathbf{p}_{31} + \frac{m_2}{|\mathbf{p}_{32}|^3} \mathbf{p}_{32} \right)$$

## New Equation 2

- $R_\omega(t)\ddot{R}_\omega^{-1}(t) = -\omega^2 \mathbf{I}_{3 \times 3}$

## New Equation 2

- $R_\omega(t)\ddot{R}_\omega^{-1}(t) = -\omega^2 \mathbf{I}_{3 \times 3}$
- $R_\omega(t)\dot{R}_\omega^{-1}(t) = \begin{pmatrix} 0 & -\omega & 0 \\ \omega & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$

## New Equation 2

- $R_\omega(t)\ddot{R}_\omega^{-1}(t) = -\omega^2 \mathbf{I}_{3 \times 3}$
- $R_\omega(t)\dot{R}_\omega^{-1}(t) = \begin{pmatrix} 0 & -\omega & 0 \\ \omega & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$
- $\mathbf{p}_{31} = \mathbf{p}_1 - \mathbf{p}_3 = \begin{bmatrix} -x_1 \\ 0 \\ 0 \end{bmatrix} - \begin{bmatrix} x \\ y \\ z \end{bmatrix} = -\begin{bmatrix} x_1 + x \\ y \\ z \end{bmatrix}$

## New Equation 2

- $R_\omega(t)\ddot{R}_\omega^{-1}(t) = -\omega^2 \mathbf{I}_{3 \times 3}$
- $R_\omega(t)\dot{R}_\omega^{-1}(t) = \begin{pmatrix} 0 & -\omega & 0 \\ \omega & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$
- $\mathbf{p}_{31} = \mathbf{p}_1 - \mathbf{p}_3 = \begin{bmatrix} -x_1 \\ 0 \\ 0 \end{bmatrix} - \begin{bmatrix} x \\ y \\ z \end{bmatrix} = -\begin{bmatrix} x_1 + x \\ y \\ z \end{bmatrix}$
- $\mathbf{p}_{32} = \mathbf{p}_2 - \mathbf{p}_3 = \begin{bmatrix} x_2 \\ 0 \\ 0 \end{bmatrix} - \begin{bmatrix} x \\ y \\ z \end{bmatrix} = -\begin{bmatrix} x - x_2 \\ y \\ z \end{bmatrix}$

## New Equation 3



## New Equation 3

$$\ddot{\mathbf{p}} = -\omega^2 \mathbf{I}_{3 \times 3} \mathbf{p} + 2 \begin{pmatrix} 0 & -\omega & 0 \\ \omega & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \dot{\mathbf{p}} + G \left( \frac{m_1}{|\mathbf{p}_{31}|^3} \mathbf{p}_{31} + \frac{m_2}{|\mathbf{p}_{32}|^3} \mathbf{p}_{32} \right)$$

## New Equation 3

$$\ddot{\mathbf{p}} = -\omega^2 \mathbf{I}_{3 \times 3} \mathbf{p} + 2 \begin{pmatrix} 0 & -\omega & 0 \\ \omega & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \dot{\mathbf{p}} + G \left( \frac{m_1}{|\mathbf{p}_{31}|^3} \mathbf{p}_{31} + \frac{m_2}{|\mathbf{p}_{32}|^3} \mathbf{p}_{32} \right)$$

$$\begin{bmatrix} \ddot{x} \\ \ddot{y} \\ \ddot{z} \end{bmatrix} = \begin{bmatrix} \omega^2 x + 2\omega \dot{y} - G \left( \frac{m_1}{|\mathbf{p}_{31}|^3} (x + x_1) + \frac{m_2}{|\mathbf{p}_{32}|^3} (x - x_2) \right) \\ \omega^2 y - 2\omega \dot{x} - y G \left( \frac{m_1}{|\mathbf{p}_{31}|^3} + \frac{m_2}{|\mathbf{p}_{32}|^3} \right) \\ -z G \left( \frac{m_1}{|\mathbf{p}_{31}|^3} + \frac{m_2}{|\mathbf{p}_{32}|^3} \right) \end{bmatrix}$$

$$|\mathbf{p}_{31}| = \sqrt{(x + x_1)^2 + y^2 + z^2}, \quad |\mathbf{p}_{32}| = \sqrt{(x - x_2)^2 + y^2 + z^2}$$

# Removing Dimensions

## Removing Dimensions

- $\omega = \sqrt{\frac{G(m_1+m_2)}{|r_{12}|^3}}$

## Removing Dimensions

- $\omega = \sqrt{\frac{G(m_1+m_2)}{|r_{12}|^3}}$
- $-m_1x_1 + m_2x_2 = 0$

## Removing Dimensions

- $\omega = \sqrt{\frac{G(m_1+m_2)}{|r_{12}|^3}}$
- $-m_1x_1 + m_2x_2 = 0$
- $|r_{12}| = \sqrt{(x_2 - (-x_1))^2 + 0^2 + 0^2} = x_2 + x_1$

## Removing Dimensions

- $\omega = \sqrt{\frac{G(m_1+m_2)}{|\mathbf{r}_{12}|^3}}$
- $-m_1x_1 + m_2x_2 = 0$
- $|\mathbf{r}_{12}| = \sqrt{(x_2 - (-x_1))^2 + 0^2 + 0^2} = x_2 + x_1$
- $x_1 = \frac{m_2}{m_1+m_2}|\mathbf{r}_{12}|$

## Removing Dimensions

- $\omega = \sqrt{\frac{G(m_1+m_2)}{|\mathbf{r}_{12}|^3}}$
- $-m_1x_1 + m_2x_2 = 0$
- $|\mathbf{r}_{12}| = \sqrt{(x_2 - (-x_1))^2 + 0^2 + 0^2} = x_2 + x_1$
- $x_1 = \frac{m_2}{m_1+m_2}|\mathbf{r}_{12}| = \mu|\mathbf{r}_{12}|$



## Removing Dimensions

- $\omega = \sqrt{\frac{G(m_1+m_2)}{|\mathbf{r}_{12}|^3}}$
- $-m_1x_1 + m_2x_2 = 0$
- $|\mathbf{r}_{12}| = \sqrt{(x_2 - (-x_1))^2 + 0^2 + 0^2} = x_2 + x_1$
- $x_1 = \frac{m_2}{m_1+m_2}|\mathbf{r}_{12}| = \mu|\mathbf{r}_{12}|$
- $x_2 = (1 - \mu)|\mathbf{r}_{12}|$

## Removing Dimensions

- $\omega = \sqrt{\frac{G(m_1+m_2)}{|\mathbf{r}_{12}|^3}}$
- $-m_1 x_1 + m_2 x_2 = 0$
- $|\mathbf{r}_{12}| = \sqrt{(x_2 - (-x_1))^2 + 0^2 + 0^2} = x_2 + x_1$
- $x_1 = \frac{m_2}{m_1+m_2} |\mathbf{r}_{12}| = \mu |\mathbf{r}_{12}|$
- $x_2 = (1 - \mu) |\mathbf{r}_{12}|$
- $|\mathbf{r}_{12}| := 1 du$

## Removing Dimensions

- $\omega = \sqrt{\frac{G(m_1+m_2)}{|\mathbf{r}_{12}|^3}}$
- $-m_1 x_1 + m_2 x_2 = 0$
- $|\mathbf{r}_{12}| = \sqrt{(x_2 - (-x_1))^2 + 0^2 + 0^2} = x_2 + x_1$
- $x_1 = \frac{m_2}{m_1+m_2} |\mathbf{r}_{12}| = \mu |\mathbf{r}_{12}|$
- $x_2 = (1 - \mu) |\mathbf{r}_{12}|$
- $|\mathbf{r}_{12}| := 1 du$
- $m_1 + m_2 := 1 mu$

## Removing Dimensions

- $\omega = \sqrt{\frac{G(m_1+m_2)}{|\mathbf{r}_{12}|^3}}$
- $-m_1 x_1 + m_2 x_2 = 0$
- $|\mathbf{r}_{12}| = \sqrt{(x_2 - (-x_1))^2 + 0^2 + 0^2} = x_2 + x_1$
- $x_1 = \frac{m_2}{m_1+m_2} |\mathbf{r}_{12}| = \mu |\mathbf{r}_{12}|$
- $x_2 = (1 - \mu) |\mathbf{r}_{12}|$
- $|\mathbf{r}_{12}| := 1 du$
- $m_1 + m_2 := 1 mu$
- $tu$  such that the frequency is  $2\pi$

## Removing Dimensions

- $\omega = \sqrt{\frac{G(m_1+m_2)}{|\mathbf{r}_{12}|^3}}$
- $-m_1x_1 + m_2x_2 = 0$
- $|\mathbf{r}_{12}| = \sqrt{(x_2 - (-x_1))^2 + 0^2 + 0^2} = x_2 + x_1$
- $x_1 = \frac{m_2}{m_1+m_2}|\mathbf{r}_{12}| = \mu|\mathbf{r}_{12}|$
- $x_2 = (1 - \mu)|\mathbf{r}_{12}|$
- $|\mathbf{r}_{12}| := 1du$
- $m_1 + m_2 := 1mu$
- $tu$  such that the frequency is  $2\pi$
- $\omega := 1rad/tu$

# Consequences Removing Dimensions

## Consequences Removing Dimensions

- $G = \left( \omega \frac{(1du)^3}{1mu} \right)^2$

## Consequences Removing Dimensions

- $G = \left( \omega \frac{(1du)^3}{1mu} \right)^2 = 1 \left( \frac{rad \times (du)^3}{tu \times mu} \right)^2$



## Consequences Removing Dimensions

- $G = \left( \omega \frac{(1du)^3}{1mu} \right)^2 = 1 \left( \frac{rad \times (du)^3}{tu \times mu} \right)^2$
- $\mu = m_2$

## Consequences Removing Dimensions

- $G = \left( \omega \frac{(1du)^3}{1mu} \right)^2 = 1 \left( \frac{rad \times (du)^3}{tu \times mu} \right)^2$
- $\mu = m_2 \leq \frac{1}{2}$

## Consequences Removing Dimensions

- $G = \left( \omega \frac{(1du)^3}{1mu} \right)^2 = 1 \left( \frac{rad \times (du)^3}{tu \times mu} \right)^2$
- $\mu = m_2 \leq \frac{1}{2}$
- $m_1 = 1 - \mu$

## Consequences Removing Dimensions

- $G = \left( \omega \frac{(1du)^3}{1mu} \right)^2 = 1 \left( \frac{rad \times (du)^3}{tu \times mu} \right)^2$
- $\mu = m_2 \leq \frac{1}{2}$
- $m_1 = 1 - \mu$
- $x_1 = \mu$

## Consequences Removing Dimensions

- $G = \left( \omega \frac{(1du)^3}{1mu} \right)^2 = 1 \left( \frac{rad \times (du)^3}{tu \times mu} \right)^2$
- $\mu = m_2 \leq \frac{1}{2}$
- $m_1 = 1 - \mu$
- $x_1 = \mu$
- $x_2 = 1 - \mu$

## Consequences Removing Dimensions

- $G = \left( \omega \frac{(1du)^3}{1mu} \right)^2 = 1 \left( \frac{rad \times (du)^3}{tu \times mu} \right)^2$
- $\mu = m_2 \leq \frac{1}{2}$
- $m_1 = 1 - \mu$
- $x_1 = \mu$
- $x_2 = 1 - \mu$

$$\begin{bmatrix} \ddot{x} \\ \ddot{y} \\ \ddot{z} \end{bmatrix} = \begin{bmatrix} x + 2\dot{y} - \frac{1-\mu}{|r_{31}|^3}(x + \mu) - \frac{\mu}{|r_{32}|^3}[x - (1 - \mu)] \\ y - 2\dot{x} - y \left( \frac{1-\mu}{|r_{31}|^3} + \frac{\mu}{|r_{32}|^3} \right) \\ -z \left( \frac{1-\mu}{|r_{31}|^3} + \frac{\mu}{|r_{32}|^3} \right) \end{bmatrix}$$

$$|r_{31}| = \sqrt{(x + \mu)^2 + y^2 + z^2}, \quad |r_{32}| = \sqrt{(x - (1 - \mu))^2 + y^2 + z^2}$$

# Clean Up

## Clean Up

- $U(x, y, z) := \frac{1}{2}(x^2 + y^2) + \frac{1-\mu}{|\mathbf{r}_{31}|} + \frac{\mu}{|\mathbf{r}_{32}|}$



## Clean Up

- $U(x, y, z) := \frac{1}{2}(x^2 + y^2) + \frac{1-\mu}{|\mathbf{r}_{31}|} + \frac{\mu}{|\mathbf{r}_{32}|}$

$$\begin{bmatrix} \ddot{x} \\ \ddot{y} \\ \ddot{z} \end{bmatrix} = \begin{bmatrix} 2\dot{y} + \frac{\partial}{\partial x} U \\ -2\dot{x} + \frac{\partial}{\partial y} U \\ \frac{\partial}{\partial z} U \end{bmatrix}$$

## Clean Up

- $U(x, y, z) := \frac{1}{2}(x^2 + y^2) + \frac{1-\mu}{|\mathbf{r}_{31}|} + \frac{\mu}{|\mathbf{r}_{32}|}$

$$\begin{bmatrix} \ddot{x} \\ \ddot{y} \\ \ddot{z} \end{bmatrix} = \begin{bmatrix} 2\dot{y} + \frac{\partial}{\partial x} U \\ -2\dot{x} + \frac{\partial}{\partial y} U \\ \frac{\partial}{\partial z} U \end{bmatrix}$$

- $\mathbf{q} = \dot{\mathbf{p}} = (\dot{x}, \dot{y}, \dot{z}) = (u, v, w)$

## Clean Up

- $U(x, y, z) := \frac{1}{2}(x^2 + y^2) + \frac{1-\mu}{|\mathbf{r}_{31}|} + \frac{\mu}{|\mathbf{r}_{32}|}$

$$\begin{bmatrix} \ddot{x} \\ \ddot{y} \\ \ddot{z} \end{bmatrix} = \begin{bmatrix} 2\dot{y} + \frac{\partial}{\partial x} U \\ -2\dot{x} + \frac{\partial}{\partial y} U \\ \frac{\partial}{\partial z} U \end{bmatrix}$$

- $\mathbf{q} = \dot{\mathbf{p}} = (\dot{x}, \dot{y}, \dot{z}) = (u, v, w)$

$$\begin{bmatrix} \dot{\mathbf{p}} \\ \dot{\mathbf{q}} \end{bmatrix} = \begin{bmatrix} \dot{x} \\ \dot{y} \\ \dot{z} \\ \dot{u} \\ \dot{v} \\ \dot{w} \end{bmatrix} = f(\mathbf{p}, \mathbf{q})$$

# Energy

## Energy

$$\ddot{x}\dot{x} + \ddot{y}\dot{y} + \ddot{z}\dot{z} = \dot{x}\left(2\dot{y} + \frac{\partial}{\partial x}U\right) + \dot{y}\left(-2\dot{x} + \frac{\partial}{\partial y}U\right) + \dot{z}\left(\frac{\partial}{\partial z}U\right)$$

## Energy

$$\begin{aligned}\ddot{x}\dot{x} + \ddot{y}\dot{y} + \ddot{z}\dot{z} &= \dot{x}\left(2\dot{y} + \frac{\partial}{\partial x}U\right) + \dot{y}\left(-2\dot{x} + \frac{\partial}{\partial y}U\right) + \dot{z}\left(\frac{\partial}{\partial z}U\right) \\ &= \dot{x}U_x + \dot{y}U_y + \dot{z}U_z\end{aligned}$$

## Energy

$$\begin{aligned}\ddot{x}\dot{x} + \ddot{y}\dot{y} + \ddot{z}\dot{z} &= \dot{x}\left(2\dot{y} + \frac{\partial}{\partial x}U\right) + \dot{y}\left(-2\dot{x} + \frac{\partial}{\partial y}U\right) + \dot{z}\left(\frac{\partial}{\partial z}U\right) \\ &= \dot{x}U_x + \dot{y}U_y + \dot{z}U_z \\ &= \dot{U}\end{aligned}$$

## Energy

$$\begin{aligned}\ddot{x}\dot{x} + \ddot{y}\dot{y} + \ddot{z}\dot{z} &= \dot{x}\left(2\dot{y} + \frac{\partial}{\partial x}U\right) + \dot{y}\left(-2\dot{x} + \frac{\partial}{\partial y}U\right) + \dot{z}\left(\frac{\partial}{\partial z}U\right) \\ &= \dot{x}U_x + \dot{y}U_y + \dot{z}U_z \\ &= \dot{U}\end{aligned}$$

- $\frac{1}{2}\frac{d}{dt}\left(\dot{x}^2 + \dot{y}^2 + \dot{z}^2\right) = \ddot{x}\dot{x} + \ddot{y}\dot{y} + \ddot{z}\dot{z}$



## Energy

$$\begin{aligned}\ddot{x}\dot{x} + \ddot{y}\dot{y} + \ddot{z}\dot{z} &= \dot{x}\left(2\dot{y} + \frac{\partial}{\partial x}U\right) + \dot{y}\left(-2\dot{x} + \frac{\partial}{\partial y}U\right) + \dot{z}\left(\frac{\partial}{\partial z}U\right) \\ &= \dot{x}U_x + \dot{y}U_y + \dot{z}U_z \\ &= \dot{U}\end{aligned}$$

- $\frac{1}{2}\frac{d}{dt}\left(\dot{x}^2 + \dot{y}^2 + \dot{z}^2\right) = \ddot{x}\dot{x} + \ddot{y}\dot{y} + \ddot{z}\dot{z}$
- $\frac{1}{2}\left(\dot{x}^2 + \dot{y}^2 + \dot{z}^2\right) = U(x, y, z) - \frac{c}{2}$

## Energy

$$\begin{aligned}\ddot{x}\dot{x} + \ddot{y}\dot{y} + \ddot{z}\dot{z} &= \dot{x}\left(2\dot{y} + \frac{\partial}{\partial x}U\right) + \dot{y}\left(-2\dot{x} + \frac{\partial}{\partial y}U\right) + \dot{z}\left(\frac{\partial}{\partial z}U\right) \\ &= \dot{x}U_x + \dot{y}U_y + \dot{z}U_z \\ &= \dot{U}\end{aligned}$$

- $\frac{1}{2}\frac{d}{dt}\left(\dot{x}^2 + \dot{y}^2 + \dot{z}^2\right) = \ddot{x}\dot{x} + \ddot{y}\dot{y} + \ddot{z}\dot{z}$
- $\frac{1}{2}\left(\dot{x}^2 + \dot{y}^2 + \dot{z}^2\right) = U(x, y, z) - \frac{C}{2}$
- $x^2 + y^2 + 2\frac{1-\mu}{|r_{31}|} + 2\frac{\mu}{|r_{32}|} - \left(\dot{x}^2 + \dot{y}^2 + \dot{z}^2\right) = C$