Metadata for Whiteadder Moor (WHI), East Lothian, Scotland, UK

Description of data set	Magnetic time-series from the BGS SWIGS differential magnetometer method (DMM) systems. Funded by NERC, grant number: NE/P017231/1 "Space Weather Impact on Ground-based Systems (SWIGS)". These data consist of measurements of the Earth's natural magnetic field at the remote site (WHIR) and the natural magnetic field plus the field created by GIC at the underline site (WHIU). The database will include .xyz files with the DMM data and one document with metadata. See Hübert et al. (2020) for further details.
Location (in lat/long): underline station	55.882456 N, -2.58699 E
Location (in lat/long): remote station	55.884003 N, -2.590714 E
Distance between DMM sites	289 m
Description of site (i.e. open moorland),	Open moorland
Equipment used	Sensys 3DFM fluxgate magnetometer and Earth Data digitisers (EDR 209, 24bit)
Installation date	09/08/2018
Removal date	30/04/2019
Number of days collected	265
Data sampling rate	1 Hz
Notes of known data gaps	NA
Qualitative estimate of site quality	Very good
Orientation angle of HV wires relative to true north	58.7° (i.e. lines run approximately ENE-WSW)
Alignment of magnetometers	Magnetic north as of 09/08/2018
Voltage level of the lines (Torness – Smeaton)	400 kV
Height above the ground of each wire (m)	Lower: 12.21 m, Middle: 21.33 m, Upper: 30.19 m
Underline magnetometer	Beneath northern circuit (18m to southern circuit)
File Type	ASCII text (.xyz)
File format	YYYY-MM-DD HH:MM:SS X Y Z Five columns with 86400 rows (plus one header row). Magnetic values in units of nT (10^-9 T). Empty values denoted by: 999999.00
Citation	Hübert, J., Beggan, C. D., Richardson, G. S., Martyn, T., & Thomson, A. W. P. (2020). Differential magnetometer measurements of geomagnetically induced currents in a complex high voltage network. Space Weather, 18, e2019SW002421. doi: 10.1029/2019SW002421