

PURE MATHEMATICS WITH STATISTICS • UPPER SIXTH

UPPER SIXTH — PURE MATHEMATICS WITH STATISTICS

CHAPTER 12

SPECIAL DISCRETE
PROBABILITY DISTRIBUTIONS

SUGGESTED TIME: 20 PERIODS

IN THIS CHAPTER

This chapter studies the three special discrete distributions of the A-Level syllabus: the binomial, the geometric and the Poisson distributions. You will learn their characteristics, probability formulae, expectations, variances, recurrence relations, cumulative tables, and how to fit, approximate, compare and apply them to real-life problems.

- By the end of this chapter I can recognise a binomial situation and state its parameters n and p .
- By the end of this chapter I can compute binomial probabilities using the formula $P(X = r) = C(n, r) p^r q^{(n-r)}$.
- By the end of this chapter I can find the expectation and variance of a binomial distribution and use recurrence formulae to generate probabilities.
- By the end of this chapter I can use the cumulative distribution function and cumulative binomial probability tables.
- By the end of this chapter I can fit a binomial distribution to given data and compare observed and expected frequencies.
- By the end of this chapter I can model waiting-time situations with the geometric distribution and compute its expectation and variance.

DOWNLOAD ON-
BUCH

Free • Google Play
AI tutor, past papers,
results



GO ONBUCH+

All signed courses, unlimited AI
tutor, PDF sheets — OnBuch+
tab in the app

THIS SIGNED COURSE CONTAINS

COURSE

complete, illustrated,
step by step

METHODS

and worked examples

EXERCISES

exam style + solutions

SUMMARY
SHEET

the essentials to
remember



Signed • OnBuch+

ONBUCH+ TEACHING TEAM
FOLLOWS THE OFFICIAL MINESEC SYLLABUS

OnBuch+

CONTENTS

- | | |
|---|---|
| 1. The Binomial Distribution: Characteristics and Probabilities | Distribution |
| 2. Expectation, Variance and Recurrence Formulae for the Binomial | 8. Poisson Approximation to the Binomial and Applications of the Poisson Distribution |
| 3. Cumulative Distribution Function and Binomial Tables | 9. Comparing Distributions and Real-Life Problem Solving |
| 4. Fitting a Binomial Distribution to Data and Applications | 10. Integration activity |
| 5. The Geometric Distribution | 11. Methods and worked examples |
| 6. The Poisson Distribution: Characteristics, Expectation and Variance | 12. Exercises |
| 7. Random Events, Unit Interval, Recurrence Formula and Mode of the Poisson | 13. Solutions to the exercises |
| | 14. Summary sheet |

CHAPTER OBJECTIVES

- By the end of this chapter I can recognise a binomial situation and state its parameters n and p .
- By the end of this chapter I can compute binomial probabilities using the formula $P(X = r) = C(n, r) p^r q^{(n-r)}$.
- By the end of this chapter I can find the expectation and variance of a binomial distribution and use recurrence formulae to generate probabilities.
- By the end of this chapter I can use the cumulative distribution function and cumulative binomial probability tables.
- By the end of this chapter I can fit a binomial distribution to given data and compare observed and expected frequencies.
- By the end of this chapter I can model waiting-time situations with the geometric distribution and compute its expectation and variance.
- By the end of this chapter I can recognise Poisson situations, compute Poisson probabilities, and use its expectation, variance, recurrence formula and mode.
- By the end of this chapter I can use the Poisson distribution as an approximation to the binomial distribution when appropriate.

- By the end of this chapter I can compare distributions and solve real-life problems involving binomial, geometric and Poisson models.

PREREQUISITES

- Probability of events, independent events and the binomial expansion (Lower Sixth Pure Mathematics).
- Discrete random variables, probability distributions, expectation $E(X)$ and variance $\text{Var}(X)$ (PMS chapter on random variables).
- Combinations $C(n,r)$ and factorial notation.
- Sigma notation and manipulation of series, including the exponential series for e^x .
- Basic statistical measures: mean and variance of a frequency distribution.

1 THE BINOMIAL DISTRIBUTION: CHARACTERISTICS AND PROBABILITIES

At the Yaoundé central vaccination post, a nurse knows from records that 8% of children develop a mild fever after a given vaccine. If 20 children are vaccinated this morning, how many should she expect to react, and what is the chance that at most two do? Meanwhile, at the Mfoundi call centre, an average of 5 calls arrive per minute: what is the probability of receiving exactly 3 calls in the next minute? These two everyday situations look different, yet each is governed by a special discrete probability distribution — the **binomial** and the **Poisson** distributions. This chapter gives you the tools to model, compute and compare them, exactly as required at the GCE Advanced Level.

We begin with the binomial distribution, which answers the nurse's question: it counts the number of "successes" in a fixed number of repeated, identical, independent experiments.

1.1 Bernoulli Trials: The Building Block

EXPERIMENT One trial, two outcomes

Toss a coin once. Either it lands Heads (call it success) or Tails (failure). Now imagine repeating the toss 10 times and counting the Heads. Each single toss is a **Bernoulli trial**; the total count of Heads is what we shall soon call a **binomial random variable**. Notice three features of each toss: it has exactly two outcomes, the probability of Heads never changes, and one toss does not influence the next. These features are the heart of the whole chapter.

DEFINITION Bernoulli trial

A **Bernoulli trial** is a single random experiment with exactly two possible outcomes, conventionally called success (S) and failure (F). We write

$$P(S) = p, \quad P(F) = q = 1 - p, \quad 0 \leq p \leq 1.$$

Everyday Bernoulli trials are everywhere:

- Tossing a coin: success = Heads, $p = 0.5$.
- A student sitting a pass/fail driving test in Buea: success = pass.
- Inspecting one bottle on a production line at a Douala brewery: success = the bottle is defective, say $p = 0.02$.
- The nurse's situation: one vaccinated child either develops a mild fever ($p = 0.08$) or does not ($q = 0.92$).

WARNING “Success” need not be pleasant

Success is just a label for the outcome we are counting. Counting defective bottles or children with fever is perfectly legitimate: the “success” is the event of interest, however undesirable it may be in real life.

1.2 From Repeated Trials to the Binomial Distribution

Suppose we repeat a Bernoulli trial a fixed number n of times, under identical and independent conditions, and we count the total number X of successes. Then X takes values in $\{0, 1, 2, \dots, n\}$ and is called a binomial random variable.

DEFINITION Binomial distribution

A discrete random variable X follows a **binomial distribution** with parameters n and p , written

$$X \sim B(n, p),$$

if X counts the number of successes in n independent Bernoulli trials, each with constant probability p of success. Here $n \in \mathbb{N}$ is the number of trials and p is the probability of success on each trial; we write $q = 1 - p$.

METHOD The four conditions to check before using $B(n, p)$

Before modelling a situation by a binomial distribution, verify all four conditions (often summarised as BINS):

1. Binary: each trial has exactly two outcomes (success/failure).
2. Independent: the outcome of one trial does not affect the others.
3. Number fixed: the number of trials n is fixed in advance.
4. Same probability: p is constant from trial to trial.

If any condition fails (e.g. sampling without replacement from a small population, or a probability that changes with time), the binomial model is not appropriate.


EXAMPLE Testing the conditions

A fair die is rolled 6 times and X counts the number of sixes obtained. Binary? Yes (six / not six). Independent? Yes. Fixed $n = 6$? Yes. Constant $p = \frac{1}{6}$? Yes. Hence $X \sim B(6, \frac{1}{6})$.
Counter-example: drawing 5 cards without replacement from a deck and counting

hearts — the probability of a heart changes after each draw, so the model is not binomial.

1.3 Derivation of the Probability Formula

Let us build the formula from a concrete case. Take $n = 3$ trials with $P(S) = p$. The full tree of outcomes is shown below.



Tree diagram for $n = 3$ Bernoulli trials: the 8 outcome paths.

Reading the 8 paths and grouping them by the number of successes:

Number of successes r	Paths	Probability of each path
0	FFF	q^3
1	SFF, FSF, FFS	pq^2
2	SSF, SFS, FSS	p^2q
3	SSS	p^3

Because the trials are independent, each path with exactly r successes and $n - r$ failures has probability $p^r q^{n-r}$ (multiplication rule). The number of such paths is the number of ways of choosing which r of the n trials are successes, namely $\binom{n}{r}$. Adding the probabilities of these mutually exclusive paths gives the general law.

PROPERTY Binomial probability formula — proof examinable

If $X \sim B(n, p)$ and $q = 1 - p$, then for $r = 0, 1, 2, \dots, n$,

$$P(X = r) = \binom{n}{r} p^r q^{n-r} = \binom{n}{r} p^r (1 - p)^{n-r}.$$

Proof. Any sequence of n trial results containing exactly r successes and $n - r$ failures has probability $p^r q^{n-r}$ by independence. There are $\binom{n}{r}$ such sequences (choose the positions of the r successes), and they are mutually exclusive, so their probabilities add. $\square$

A reassuring check: the probabilities must sum to 1. Indeed, by the binomial theorem,

$$\sum_{r=0}^n P(X = r) = \sum_{r=0}^n \binom{n}{r} p^r q^{n-r} = (q + p)^n = 1^n = 1.$$

So the formula defines a genuine probability distribution, and the numbers $\binom{n}{r} p^r q^{n-r}$ are precisely the terms of the expansion of $(q + p)^n$ — this is why the distribution is called binomial.

KEY POINTS Key facts

- $X \sim B(n, p)$: X counts successes in n independent trials with constant p .
- $P(X = r) = \binom{n}{r} p^r q^{n-r}$, with $q = 1 - p$.
- $P(X = 0) = q^n$ and $P(X = n) = p^n$ (only one path each).
- $\sum_{r=0}^n P(X = r) = (q + p)^n = 1$.

1.4 Direct Computation of Probabilities

WORKED EXAMPLE 1 Computing binomial probabilities by hand

A fair coin is tossed 5 times. Let X be the number of Heads. Find (a) $P(X = 3)$; (b) $P(X = 0)$; (c) $P(X = 5)$; (d) verify the total is 1.

SOLUTION

Solution. Each toss is a Bernoulli trial with $p = 0.5$, $q = 0.5$, and the $n = 5$ tosses are independent, so $X \sim B(5, 0.5)$.

$$(a) P(X = 3) = \binom{5}{3} (0.5)^3 (0.5)^2 = 10 \times (0.5)^5 = \frac{10}{32} = 0.3125.$$

$$(b) P(X = 0) = q^5 = (0.5)^5 = \frac{1}{32} = 0.03125.$$

(c) $P(X = 5) = p^5 = (0.5)^5 = 0.03125$.

(d) The full distribution: $P(X = r) = \binom{5}{r}/32$ gives $\frac{1, 5, 10, 10, 5, 1}{32}$, and $1 + 5 + 10 + 10 + 5 + 1 = 32$, so the total is 1. ■

WORKED EXAMPLE 2 The nurse's vaccination problem — first part

At the Yaoundé vaccination post, 8% of children develop a mild fever after the vaccine. Out of 20 children vaccinated this morning, find the probability that (a) exactly 2 react; (b) none reacts.

SOLUTION

Solution. Assuming the children react independently, let X be the number who develop a fever. Then $X \sim B(20, 0.08)$, with $q = 0.92$.

(a) $P(X = 2) = \binom{20}{2}(0.08)^2(0.92)^{18} = 190 \times 0.0064 \times (0.92)^{18}$.

Now $(0.92)^{18} \approx 0.2229$, so $P(X = 2) \approx 190 \times 0.0064 \times 0.2229 \approx 0.2711$.

(b) $P(X = 0) = (0.92)^{20} \approx 0.1887$.

So there is roughly a 19% chance that no child reacts, and about a 27% chance that exactly two do. ■

1.5 Cumulative Distribution Function and Binomial Tables

Most examination questions ask not for a single value but for an accumulated probability. Since the events $\{X = r\}$ are mutually exclusive, we simply add:

$$P(X \geq r) = \sum_{k=r}^n P(X = k), \quad P(X \leq r) = \sum_{k=0}^r P(X = k), \quad P(a \leq X \leq b) = \sum_{k=a}^b P(X = k).$$

DEFINITION Cumulative distribution function (CDF)

For a discrete random variable X , the **cumulative distribution function** $F(x)$ is defined by

$$F(x) = P(X \leq x) = \sum_{k \leq x} P(X = k).$$

For $X \sim B(n, p)$, $F(r) = \sum_{k=0}^r \binom{n}{k} p^k q^{n-k}$ for $r = 0, 1, \dots, n$.

A very useful complement trick avoids long sums:

$$P(X \geq 1) = 1 - P(X = 0) = 1 - q^n.$$

WARNING Strict versus non-strict inequalities

$P(X \geq 2) = 1 - P(X = 0) - P(X = 1)$, but $P(X > 2) = 1 - P(X = 0) - P(X = 1) - P(X = 2)$. Read the question twice: “at least”, “at most”, “more than” and “fewer than” all mean different sums. Also, never forget the coefficient $\binom{n}{r}$, and never swap p and q : the power of p is the number of successes r .

WORKED EXAMPLE 3 Compound probabilities

A marksman hits a target with probability 0.7 on each shot, independently. He fires 4 shots. Find the probability that he hits (a) at least once; (b) at least 3 times; (c) between 1 and 2 times inclusive.

SOLUTION

Solution. Let X be the number of hits. Then $X \sim B(4, 0.7)$, $q = 0.3$.

$$(a) P(X \geq 1) = 1 - P(X = 0) = 1 - (0.3)^4 = 1 - 0.0081 = 0.9919.$$

$$(b) P(X \geq 3) = P(X = 3) + P(X = 4) = \binom{4}{3}(0.7)^3(0.3) + (0.7)^4 = 4 \times 0.343 \times 0.3 + 0.2401 = 0.4116 + 0.2401 = 0.6517.$$

$$(c) P(1 \leq X \leq 2) = \binom{4}{1}(0.7)(0.3)^3 + \binom{4}{2}(0.7)^2(0.3)^2 = 4 \times 0.7 \times 0.027 + 6 \times 0.49 \times 0.09 = 0.0756 + 0.2646 = 0.3402. \blacksquare$$

METHOD Using cumulative binomial probability tables

When n is large, summing many terms by hand is tedious. In examinations, **cumulative binomial tables** giving $P(X \leq r)$ for selected n and p are often provided.


1. Identify n and p ; locate the correct page/block for that n .
 2. Find the column for your p (or q if the table uses q).
 3. Read off the value opposite r for $P(X \leq r)$.
 4. For $P(X \geq r)$ use $1 - P(X \leq r - 1)$; for $P(X = r)$ use $P(X \leq r) - P(X \leq r - 1)$.
- If tables are not provided, you must compute using the formula or a calculator (if allowed).

EXAMPLE Reading a cumulative table

Suppose $X \sim B(10, 0.3)$ and the table gives $P(X \leq 3) = 0.6496$ and $P(X \leq 2) = 0.3828$. Then: $P(X = 3) = 0.6496 - 0.3828 = 0.2668$, $P(X \geq 4) = 1 - 0.6496 = 0.3504$, $P(2 \leq X \leq 4) = P(X \leq 4) - P(X \leq 1)$.

1.6 The Shape of the Binomial Distribution

The value of p controls the shape of the distribution. When $p = 0.5$ the distribution is symmetric about its centre; when $p < 0.5$ it is skewed to the right (a long tail of large values); when $p > 0.5$ it is skewed to the left. The bar charts below compare $B(10, 0.3)$ and $B(10, 0.5)$.



Effect of p on the shape: $B(10, 0.3)$ is right-skewed; $B(10, 0.5)$ is symmetric.

Observe that for $p = 0.3$ the most likely value is $r = 3$ and the probabilities die away slowly to the right, while for $p = 0.5$ the bars are mirror images about $r = 5$. In general the distribution peaks near $r = np$, an observation we shall make precise when we study the expectation in the next section.

DID YOU KNOW? Exam tip — GCE Advanced Level

Always state the distribution before computing: write “Let X be the number of children who react; then $X \sim B(20, 0.08)$.” Examiners award a mark for correctly identifying the model and its parameters, and it protects you from mixing up n and p later. Quote the formula $P(X = r) = \binom{n}{r} p^r q^{n-r}$ once, then substitute.

EXERCISE 1 Practice

1. A fair die is thrown 8 times and X counts the sixes. State the distribution of X and compute $P(X = 2)$ and $P(X \geq 1)$.
2. It is known that 30% of seedlings in a nursery fail to survive transplanting. Out of 6 transplanted seedlings, find the probability that (a) exactly 4 survive; (b) at most 1 survives.
3. For $X \sim B(5, 0.4)$, write out the full probability distribution and verify that the probabilities sum to 1.

4. A multiple-choice test has 10 questions, each with 4 options. A student guesses all answers. Let X be the number of correct answers. State the distribution of X and find $P(X \leq 2)$.

SOLUTION Exercise 1

1. $X \sim B\left(8, \frac{1}{6}\right)$. $P(X = 2) = \binom{8}{2} \left(\frac{1}{6}\right)^2 \left(\frac{5}{6}\right)^6 = 28 \times \frac{1}{36} \times 0.3349 \approx 0.2605$. $P(X \geq 1) = 1 - \left(\frac{5}{6}\right)^8 \approx 1 - 0.2326 = 0.7674$.
2. Let $Y = \text{number that survive}$, $Y \sim B(6, 0.7)$. (a) $P(Y = 4) = \binom{6}{4} (0.7)^4 (0.3)^2 = 15 \times 0.2401 \times 0.09 = 0.3241$. (b) $P(Y \leq 1) = (0.3)^6 + 6(0.7)(0.3)^5 = 0.000729 + 0.010206 = 0.0109$.
3. $P(X = r) = \binom{5}{r} (0.4)^r (0.6)^{5-r}$: 0.07776, 0.25920, 0.34560, 0.23040, 0.07680, 0.01024. Sum: $0.07776 + 0.25920 + 0.34560 + 0.23040 + 0.07680 + 0.01024 = 1.00000$.
4. $X \sim B(10, 0.25)$. $P(X \leq 2) = P(X = 0) + P(X = 1) + P(X = 2) = (0.75)^{10} + 10(0.25)(0.75)^9 + 45(0.25)^2(0.75)^8 \approx 0.0563 + 0.1877 + 0.2816 = 0.5256$. ■

You can now recognise a binomial situation, state $X \sim B(n, p)$, and compute both single and compound probabilities. In the next section we shall measure the average behaviour of this distribution — its expectation and variance — and discover a quick recurrence way of generating the probabilities.

2 EXPECTATION, VARIANCE AND RECURRENCE FORMULAE FOR THE BINOMIAL

In the previous section we defined the binomial distribution $B(n, p)$ and learnt how to compute individual probabilities $P(X = r) = \binom{n}{r} p^r q^{n-r}$. Knowing the full probability distribution is excellent, but in practice an examiner, an epidemiologist or an engineer often wants quick answers to three questions: What value do we expect on average? How spread out are the results? Which outcome is the most likely? This section answers all three: we derive the **expectation** $E(X) = np$ and the **variance** $\text{Var}(X) = npq$, and we establish a powerful **recurrence formula** linking consecutive probabilities, which leads us directly to the **mode** of the distribution.

2.1 Expectation of a Binomial Distribution

An intuitive situation.

Suppose a nurse at a health centre in Bamenda vaccinates $n = 200$ children in a week, and each child has a probability $p = 0.05$ of developing a mild fever after the injection. How many cases of fever should she expect? Common sense says: 5% of 200, that is 10 children. Nothing more complicated is needed: the expected number of successes in n independent trials is simply n times the probability of success in one trial. This is exactly the content of the following property.

PROPERTY Expectation of $B(n, p)$

Let $X \sim B(n, p)$. Then

$$E(X) = np.$$

Proof (examinable). Write $X = X_1 + X_2 + \dots + X_n$, where X_i is the indicator of success on trial i : $X_i = 1$ with probability p and $X_i = 0$ with probability $q = 1 - p$. Each X_i is a Bernoulli variable, so $E(X_i) = 1 \times p + 0 \times q = p$. By linearity of expectation (which holds for any random variables, independent or not),

$$E(X) = E(X_1) + E(X_2) + \dots + E(X_n) = \underbrace{p + p + \dots + p}_{n \text{ terms}} = np.$$

WARNING Do not confuse $E(X)$ with a possible value

The expectation np need not be an integer, and it need not be a value X can actually take. If $X \sim B(12, 0.4)$, then $E(X) = 4.8$: we can never observe 4.8 successes, but over many repetitions of the 12 trials, the average number of successes settles near 4.8. The expectation is a long-run average, not a prediction of a single experiment.

2.2 Variance and Standard Deviation

Knowing the average is not enough. Two distributions can share the same mean yet behave very differently: one may cluster tightly around the mean, the other may fluctuate wildly. The **variance** measures this spread.

Intuition first.

For a single Bernoulli trial, the variance is easily computed:

$$\text{Var}(X_i) = E(X_i^2) - (E(X_i))^2 = p - p^2 = p(1 - p) = pq.$$

Notice that this is largest when $p = \frac{1}{2}$ (maximum uncertainty: success and failure are equally likely) and smallest when p is near 0 or 1 (the outcome is almost certain). Since the n trials of a binomial experiment are independent, variances add up, giving the following result.

PROPERTY Variance of $B(n, p)$

Let $X \sim B(n, p)$ with $q = 1 - p$. Then

$$\text{Var}(X) = npq \quad \text{and} \quad \sigma = \sqrt{npq}.$$

Proof (examinable). With $X = X_1 + \dots + X_n$ as above, the X_i are independent, so the variance of the sum is the sum of the variances:

$$\text{Var}(X) = \text{Var}(X_1) + \dots + \text{Var}(X_n) = \underbrace{pq + pq + \dots + pq}_{n \text{ terms}} = npq.$$

The standard deviation is by definition $\sigma = \sqrt{\text{Var}(X)} = \sqrt{npq}$.

WARNING The classic error: $\text{Var}(X) = np$

Every year, candidates write $\text{Var}(X) = np$. This is wrong: np is the mean. The variance is $\text{Var}(X) = npq$ — never forget the factor q . A quick sanity check: if $p = 1$ (certain success), the variance must be 0 since nothing varies; $npq = n \times 1 \times 0 = 0$ confirms this, while $np = n$ would absurdly claim variability. Also remember: the standard deviation is $\sqrt{npq}$, not npq .

KEY POINTS Summary of key parameters

For $X \sim B(n, p)$, $q = 1 - p$:

$$E(X) = np, \quad \text{Var}(X) = npq, \quad \sigma = \sqrt{npq}.$$

These three formulae must be known by heart for the GCE Advanced Level examination.

WORKED EXAMPLE 4 Vaccination reactions

A vaccine produces a mild side-effect in 8% of patients. In a clinic, 150 patients are vaccinated independently. Let X be the number of patients who develop the side-effect. Find the mean, variance and standard deviation of X .

SOLUTION

Solution. Each patient is a Bernoulli trial with $p = 0.08$, and the trials are independent, so $X \sim B(150, 0.08)$ with $q = 0.92$.

$$\begin{aligned} E(X) &= np = 150 \times 0.08 = 12, \\ \text{Var}(X) &= npq = 150 \times 0.08 \times 0.92 = 11.04, \\ \sigma &= \sqrt{11.04} \approx 3.32. \end{aligned}$$

The clinic should expect about 12 cases, with typical fluctuations of roughly ± 3 cases around this value.

WORKED EXAMPLE 5 Guessing a multiple-choice quiz

A GCE candidate guesses all 20 questions of a multiple-choice paper, each question

having 4 options of which exactly one is correct. Let X be the number of correct answers. Find $E(X)$, $\text{Var}(X)$ and σ , and comment.

SOLUTION

Solution. Guessing gives $p = \frac{1}{4} = 0.25$, $q = 0.75$, $n = 20$, so $X \sim B(20, 0.25)$.

$$\begin{aligned} E(X) &= 20 \times 0.25 = 5, \\ \text{Var}(X) &= 20 \times 0.25 \times 0.75 = 3.75, \\ \sigma &= \sqrt{3.75} \approx 1.94. \end{aligned}$$

A pure guesser scores on average 5 out of 20, and will rarely deviate from 5 by more than about 2 correct answers. Guessing is not a strategy!

2.3 The Recurrence Formula

The problem.

Computing $P(X = r) = \binom{n}{r} p^r q^{n-r}$ for $r = 0, 1, 2, \dots, n$ one by one is painful: for $n = 12$ you would evaluate 13 different binomial coefficients, each with powers of p and q . Under examination conditions this is slow and error-prone. Fortunately, consecutive probabilities are linked by a remarkably simple ratio.

PROPERTY Recurrence formula for the binomial

Let $X \sim B(n, p)$, $q = 1 - p$, and write $P(r) = P(X = r)$. Then for $r = 0, 1, \dots, n - 1$,

$$\frac{P(X = r + 1)}{P(X = r)} = \frac{(n - r)p}{(r + 1)q}, \quad \text{so} \quad P(X = r + 1) = \frac{(n - r)p}{(r + 1)q} P(X = r).$$

Proof (examinable and instructive). Take the ratio of two consecutive terms:

$$\begin{aligned} \frac{P(X = r + 1)}{P(X = r)} &= \frac{\binom{n}{r+1} p^{r+1} q^{n-r-1}}{\binom{n}{r} p^r q^{n-r}} \\ &= \frac{n!}{(r+1)!(n-r-1)!} \times \frac{r!(n-r)!}{n!} \times \frac{p^{r+1}}{p^r} \times \frac{q^{n-r-1}}{q^{n-r}} \\ &= \frac{(n-r)}{(r+1)} \times p \times \frac{1}{q} = \frac{(n-r)p}{(r+1)q}. \end{aligned}$$

Here we used $\frac{r!}{(r+1)!} = \frac{1}{r+1}$ and $\frac{(n-r)!}{(n-r-1)!} = n-r$.

EXAMPLE Remark: why this formula is a gift in exams

The recurrence formula means you compute only one binomial probability from scratch — usually the easiest, $P(X = 0) = q^n$ — and then generate the whole table by repeated multiplication. No more large coefficients $\binom{n}{r}$, no more fresh powers of p and q at each step. As a bonus, any arithmetic slip shows up immediately because the final probabilities must sum to 1: this gives you a free checking mechanism.

METHOD Building a full probability table with the recurrence formula

1. Compute the starting value $P(0) = q^n$.
2. For each $r = 0, 1, 2, \dots, n-1$, compute the multiplier $m_r = \frac{(n-r)p}{(r+1)q}$.
3. Set $P(r+1) = m_r \times P(r)$, keeping at least 4 decimal places throughout.
4. Stop at $P(n)$, then check: $\sum_{r=0}^n P(r)$ must equal 1 (up to rounding).

WORKED EXAMPLE 6 Full table of $B(3, 0.4)$ by recurrence

Let $X \sim B(3, 0.4)$. Using the recurrence formula, construct the complete probability distribution of X .

SOLUTION

Solution. Here $n = 3$, $p = 0.4$, $q = 0.6$. Step 1: $P(0) = q^3 = 0.6^3 = 0.216$. Step 2: apply $P(r+1) = \frac{(3-r) \times 0.4}{(r+1) \times 0.6} P(r)$:

$$\begin{aligned} P(1) &= \frac{3 \times 0.4}{1 \times 0.6} \times 0.216 = 2 \times 0.216 = 0.432, \\ P(2) &= \frac{2 \times 0.4}{2 \times 0.6} \times 0.432 = \frac{2}{3} \times 0.432 = 0.288, \\ P(3) &= \frac{1 \times 0.4}{3 \times 0.6} \times 0.288 = \frac{2}{9} \times 0.288 = 0.064. \end{aligned}$$

Check: $0.216 + 0.432 + 0.288 + 0.064 = 1.000$. ✓ The distribution is

r	0	1	2	3
$P(X = r)$	0.216	0.432	0.288	0.064

2.4 Locating the Mode with the Recurrence Ratio

The **mode** of a discrete distribution is the value of r with the greatest probability — the most likely outcome. The recurrence formula gives an elegant way to find it without computing any probability at all.

The key idea.

The probabilities $P(0), P(1), P(2), \dots$ first increase, reach a peak (the mode), then decrease. The sequence is increasing exactly when the next term is at least the current one, i.e. when the recurrence ratio is at least 1:

$$\frac{P(X = r + 1)}{P(X = r)} = \frac{(n - r)p}{(r + 1)q} \geq 1.$$

Let us solve this inequality:

$$\begin{aligned} (n - r)p \geq (r + 1)q &\iff np - rp \geq rq + q \\ &\iff np - q \geq r(p + q) \\ &\iff r \leq np - q = (n + 1)p - 1 \quad (\text{since } p + q = 1). \end{aligned}$$

So the probabilities increase while $r \leq (n + 1)p - 1$ and decrease afterwards. The peak — the mode — is therefore the first integer r for which the increase stops.

METHOD Finding the mode of $B(n, p)$


1. Compute $(n + 1)p$.
2. If $(n + 1)p$ is not an integer, the mode is the unique integer

$$m = \lfloor (n + 1)p \rfloor \quad (\text{the integer part of } (n + 1)p).$$

3. If $(n + 1)p$ is an integer k , there are two equal modes: $r = k - 1$ and $r = k$, because then $P(k)/P(k - 1) = 1$.

EXAMPLE Mode of $B(12, 0.4)$

Here $(n + 1)p = 13 \times 0.4 = 5.2$, not an integer, so the mode is $m = \lfloor 5.2 \rfloor = 5$: the most likely number of successes is 5. Note the mean is $np = 4.8$: mode and mean are close but not identical — the distribution is slightly skewed, as the bar chart below shows.



Bar chart of $B(12, 0.4)$. The orange bar marks the mode $r = 5 = \lfloor (n+1)p \rfloor$; the dashed green line marks the mean $E(X) = np = 4.8$.

WORKED EXAMPLE 7 Mode and most likely outcome

A fair coin is tossed 10 times and X counts the number of heads. Using the recurrence ratio, find the mode of X and verify it directly.

SOLUTION

Solution. $X \sim B(10, 0.5)$, so $(n+1)p = 11 \times 0.5 = 5.5$, not an integer: the mode is $m = \lfloor 5.5 \rfloor = 5$. Verification via the ratio: with $p = q = 0.5$, the ratio is $\frac{P(r+1)}{P(r)} = \frac{10-r}{r+1}$. This is ≥ 1 iff $10-r \geq r+1$, i.e. $r \leq 4.5$, i.e. $r \leq 4$. So $P(0) < P(1) < \dots < P(5)$ and $P(5) > P(6) > \dots > P(10)$: the peak is at $r = 5$, with $P(X = 5) = \binom{10}{5}(0.5)^{10} = \frac{252}{1024} \approx 0.246$. ✓

WARNING Mode versus mean

Do not write “the mode is np ”. The mean is np ; the mode is $\lfloor (n+1)p \rfloor$ (with a double mode when $(n+1)p$ is an integer). For $B(12, 0.4)$ these are 4.8 and 5 respectively — close, but conceptually different quantities, and examiners test the distinction.

EXERCISE 2 Practice

1. A machine produces defective items with probability 0.02. A batch of 500 items is

inspected. Let X be the number of defectives. Find $E(X)$, $\text{Var}(X)$ and σ .

- For $X \sim B(8, 0.3)$, use the recurrence formula starting from $P(0) = 0.7^8$ to compute $P(1)$ and $P(2)$, giving 4 decimal places.
- Find the mode of $B(9, 0.5)$. What do you notice?

SOLUTION Exercise 1

- $X \sim B(500, 0.02)$: $E(X) = np = 10$; $\text{Var}(X) = npq = 500 \times 0.02 \times 0.98 = 9.8$; $\sigma = \sqrt{9.8} \approx 3.13$.
- $P(0) = 0.7^8 = 0.0576$. Ratio for $r = 0$: $\frac{8 \times 0.3}{1 \times 0.7} = \frac{24}{7}$, so $P(1) = \frac{24}{7} \times 0.0576 \approx 0.1977$. Ratio for $r = 1$: $\frac{7 \times 0.3}{2 \times 0.7} = 1.5$, so $P(2) = 1.5 \times 0.1977 \approx 0.2965$.
- $(n + 1)p = 10 \times 0.5 = 5$, an integer: there are two modes, $r = 4$ and $r = 5$, with equal probabilities $P(4) = P(5) \approx 0.2461$.

KEY POINTS What you must remember from this section

- For $X \sim B(n, p)$: $E(X) = np$, $\text{Var}(X) = npq$, $\sigma = \sqrt{npq}$ — the factor q in the variance is the most frequently forgotten detail in the exam.
- The recurrence formula $\frac{P(X = r + 1)}{P(X = r)} = \frac{(n - r)p}{(r + 1)q}$ generates the whole distribution from $P(0) = q^n$ and provides a built-in check ($\sum P(r) = 1$).
- The probabilities increase while $r \leq (n + 1)p - 1$; hence the mode is $\lfloor (n + 1)p \rfloor$, with a double mode $k - 1$ and k when $(n + 1)p = k$ is an integer.

These parameters and tools will be essential in the next section, where we move from individual probabilities to cumulative probabilities $P(X \leq r)$ and learn to read the cumulative binomial tables provided in the examination.

3 CUMULATIVE DISTRIBUTION FUNCTION AND BINOMIAL TABLES

In the previous sections, we mastered the calculation of individual probabilities $P(X = r)$ for a binomial random variable $X \sim B(n, p)$, using the formula, the recurrence relation, and the properties of expectation and variance. However, in many practical situations — quality control in a factory in Douala, medical screening in Yaoundé, or crop-yield estimates in the Northwest

Region — we are not interested in the probability of exactly r successes, but rather in the probability of at most r successes, or at least r successes. This calls for the **cumulative distribution function** (CDF) and the use of standard statistical tables, which are provided in the GCE examination.

3.1 The Cumulative Distribution Function (CDF)

DEFINITION Cumulative Distribution Function of a Discrete Random Variable

Let X be a discrete random variable taking values $x_1, x_2, \dots$ with probabilities $P(X = x_i)$. The **cumulative distribution function**, denoted by $F(x)$, is defined for all real x as:

$$F(x) = P(X \leq x) = \sum_{x_i \leq x} P(X = x_i).$$

For a binomial variable $X \sim B(n, p)$, the CDF at an integer r ($0 \leq r \leq n$) is:

$$F(r) = P(X \leq r) = \sum_{k=0}^r \binom{n}{k} p^k (1-p)^{n-k}.$$

By convention, $F(x) = 0$ for $x < 0$ and $F(x) = 1$ for $x \geq n$.

KEY POINTS Properties of the CDF for a Discrete Variable


- $F(x)$ is a non-decreasing step function, with jumps exactly at the values taken by X .
- $F(x)$ is right-continuous: the value at a jump point r is the value after the jump, i.e., $F(r) = P(X \leq r)$.
- $F(x) = 0$ for all x below the smallest value of X , and $F(x) = 1$ for all x at or above the largest value of X .
- $P(X = r) = F(r) - F(r-1)$ for integer r .
- $P(a < X \leq b) = F(b) - F(a)$, and $P(a \leq X \leq b) = F(b) - F(a-1)$.

EXAMPLE Building a Cumulative Table for $B(5, 0.4)$

Let $X \sim B(5, 0.4)$. We first compute the individual probabilities $P(X = r)$ (using the formula $P(X = r) = \binom{5}{r}(0.4)^r(0.6)^{5-r}$ or the recurrence relation), then accumulate them row by row.

r	0	1	2	3	4	5
$P(X = r)$	0.0778	0.2592	0.3456	0.2304	0.0768	0.0102
$F(r) = P(X \leq r)$	0.0778	0.3370	0.6826	0.9130	0.9898	1.0000

For instance, $F(2) = 0.0778 + 0.2592 + 0.3456 = 0.6826$. Check: the final cumulative probability must equal 1 (subject to rounding).



Cumulative distribution function $F(r) = P(X \leq r)$ for $X \sim B(8, 0.25)$. The function jumps at each integer value r ; the solid dot indicates the value actually taken by F at the jump.

3.2 Using Cumulative Binomial Probability Tables

In the GCE examination, you are provided with **cumulative binomial probability tables**. These tables give $P(X \leq r)$ directly for various values of n (usually up to 50), p (usually from 0.05 to 0.50 in steps of 0.05), and r .

KEY POINTS Structure of the Tables

The tables are organised by n (number of trials). For each n , there are columns for different values of p . The rows correspond to $r = 0, 1, 2, \dots, n$. The entry in the cell (r, p) is $P(X \leq r)$ for $X \sim B(n, p)$.

Extract from Cumulative Binomial Table

$n = 10$	$p = 0.20$
0	$P(X \leq r)$
1	0.3758
2	0.6778
3	0.8791
4	0.9672
5	0.9936
6	0.9991
7	0.9999

← $P(X \leq 2) = 0.6778$

Reading the table: for $n = 10$, $p = 0.20$, locate the block for $n = 10$, the column $p = 0.20$, and the row $r = 2$. The entry is 0.6778, so $P(X \leq 2) = 0.6778$.

3.3 Three Essential Table-Reading Rules

Since the tables only give $P(X \leq r)$, we must derive all other probabilities from this single quantity.

METHOD From Cumulative Tables to Any Probability

Let $X \sim B(n, p)$ and let $F(r) = P(X \leq r)$ be the value read from the table.

1. Exactly r successes: $P(X = r) = F(r) - F(r - 1)$.
2. At least r successes: $P(X \geq r) = 1 - P(X \leq r - 1) = 1 - F(r - 1)$.
3. More than r successes: $P(X > r) = 1 - P(X \leq r) = 1 - F(r)$.

WARNING The Classic Trap: $P(X \geq r)$ versus $P(X > r)$

A very common error is to write $P(X \geq r) = 1 - P(X \leq r)$. This is wrong. The event $X \geq r$ includes $X = r$, so its complement is $X \leq r - 1$, not $X \leq r$:

$$P(X \geq r) = 1 - F(r - 1) \quad \text{and NOT} \quad 1 - F(r).$$

Similarly, $P(X > r) = 1 - F(r)$ because the event $X > r$ excludes r . Always ask yourself: is r itself included or excluded?

WORKED EXAMPLE 8 Using Tables for $X \sim B(15, 0.3)$

Using the cumulative binomial tables for $n = 15$, $p = 0.30$, find:

1. $P(X \leq 3)$
2. $P(X = 3)$
3. $P(X \geq 3)$
4. $P(X > 3)$

SOLUTION

Solution. From the table ($n = 15$, $p = 0.30$):

$$F(2) = P(X \leq 2) = 0.1268, \quad F(3) = P(X \leq 3) = 0.2969, \quad F(4) = P(X \leq 4) = 0.5155.$$

1. Direct reading: $P(X \leq 3) = F(3) = \mathbf{0.2969}$.
2. Subtraction: $P(X = 3) = F(3) - F(2) = 0.2969 - 0.1268 = \mathbf{0.1701}$.
3. At least 3: $P(X \geq 3) = 1 - F(2) = 1 - 0.1268 = \mathbf{0.8732}$.
4. More than 3: $P(X > 3) = 1 - F(3) = 1 - 0.2969 = \mathbf{0.7031}$.

Check: $P(X \geq 3) = P(X = 3) + P(X > 3) = 0.1701 + 0.7031 = 0.8732$. Consistent.

3.4 Inverse Reading: Finding Critical Values

Sometimes we are given a probability α (e.g., 0.95 or 0.99) and asked to find the smallest integer r such that $P(X \leq r) \geq \alpha$. This is finding a **critical value** (or percentile). This skill is essential for hypothesis testing (covered later) and for quality control.

METHOD Finding the Smallest r such that $P(X \leq r) \geq \alpha$

1. Locate the correct block for n and the column for p .
2. Scan down the column of cumulative probabilities $F(r)$.
3. Find the first row where the value is greater than or equal to α .
4. The corresponding value of r is the answer.

EXAMPLE Finding a Critical Value for $B(20, 0.25)$

Find the smallest integer r such that $P(X \leq r) \geq 0.90$ for $X \sim B(20, 0.25)$.

Solution. Scan the column $n = 20, p = 0.25$:

$$F(6) = 0.7858 \text{ (too small)}, \quad F(7) = 0.8982 \text{ (too small)}, \quad F(8) = 0.9591 (\geq 0.90).$$

The smallest such r is 8.

3.5 Handling $p > 0.5$: The Symmetry Trick

Standard binomial tables usually only provide values for $p \leq 0.5$. If $p > 0.5$, we exploit the symmetry of the binomial distribution by counting failures instead of successes.

PROPERTY Symmetry of the Binomial Distribution

If $X \sim B(n, p)$, then the number of failures $Y = n - X$ follows $B(n, 1 - p)$, and

$$P(X \leq r) = P(n - Y \leq r) = P(Y \geq n - r) = 1 - P(Y \leq n - r - 1).$$

Since $1 - p < 0.5$ when $p > 0.5$, we can look up $Y \sim B(n, 1 - p)$ in the tables.

WORKED EXAMPLE 9 Using Tables for $p > 0.5$: $X \sim B(10, 0.7)$

Find $P(X \leq 8)$ for $X \sim B(10, 0.7)$.

SOLUTION

Solution. Here $p = 0.7 > 0.5$. Define $Y = \text{number of failures} = 10 - X$. Then $Y \sim B(10, 0.3)$.

$$P(X \leq 8) = P(10 - Y \leq 8) = P(Y \geq 2) = 1 - P(Y \leq 1).$$

From the tables for $n = 10, p = 0.30$: $F_Y(1) = P(Y \leq 1) = 0.1493$.

$$P(X \leq 8) = 1 - 0.1493 = \mathbf{0.8507}.$$

3.6 Limitations of Tables and When to Use the Formula

While tables are fast, they have limitations you must be aware of for the GCE:

- **Limited n :** tables typically stop at $n = 50$. If $n > 50$, you cannot use the tables directly (use the Poisson approximation if p is small, or the Normal approximation covered later).
- **Limited p :** tables usually give $p = 0.05, 0.10, \dots, 0.50$. If $p = 0.37$, you must use the formula (interpolation is not expected in the examination).
- **Rounding:** tables give 4 decimal places. If the question asks for an exact probability or an answer in terms of p , you **must** use the formula $\binom{n}{r} p^r (1 - p)^{n-r}$.

- **Accumulated rounding:** computing $P(X = r)$ via $F(r) - F(r - 1)$ subtracts two rounded values. For a single individual probability, the formula (or the recurrence relation) is often more accurate.

KEY POINTS GCE Exam Technique: Quoting Tables

- Always state the distribution: " $X \sim B(n, p)$ ".
- State the values you are reading: "From tables, $P(X \leq 2) = 0.6778$ ".
- Show the subtraction step explicitly for $P(X = r)$, $P(X \geq r)$ and $P(X > r)$.
- Give final answers to 4 decimal places (or 3 significant figures) unless an exact answer is requested.
- If $p > 0.5$, clearly define $Y = n - X$ and state $Y \sim B(n, 1 - p)$ before using the tables.

3.7 Worked Examination–Style Example

WORKED EXAMPLE 10 GCE Style: Cumulative Probabilities and Critical Value

The probability that a cocoa pod harvested in a certain region is of export quality is 0.65. A random sample of 12 pods is selected.

1. Model the number of export-quality pods with a suitable distribution, stating your assumptions.
2. Using tables, find the probability that:
 - (a) at most 7 pods are of export quality;
 - (b) exactly 7 pods are of export quality;
 - (c) at least 9 pods are of export quality.
3. Find the smallest number k such that the probability of having at most k export-quality pods is at least 0.95.

SOLUTION

Solution.

1. Let X be the number of export-quality pods in the sample of 12. There is a fixed number $n = 12$ of trials, each trial has two outcomes (export quality or not), the probability $p = 0.65$ is constant, and the pods are selected at random so the trials are independent. Hence

$$X \sim B(12, 0.65).$$

2. Since $p = 0.65 > 0.5$, we use the symmetry trick. Let $Y =$ number of non-export pods =

12 - X. Then $Y \sim B(12, 0.35)$, and we use the tables for $n = 12, p = 0.35$:

$$F_Y(3) = 0.3467, \quad F_Y(4) = 0.5833, \quad F_Y(5) = 0.7873.$$

(a) $P(X \leq 7) = P(12 - Y \leq 7) = P(Y \geq 5) = 1 - P(Y \leq 4) = 1 - 0.5833 = \mathbf{0.4167}.$

(b) $P(X = 7) = P(Y = 5) = F_Y(5) - F_Y(4) = 0.7873 - 0.5833 = \mathbf{0.2040}.$

(c) $P(X \geq 9) = P(12 - Y \geq 9) = P(Y \leq 3) = F_Y(3) = \mathbf{0.3467}.$

3. We need the smallest k such that $P(X \leq k) \geq 0.95$.

$$P(X \leq k) = P(Y \geq 12 - k) = 1 - P(Y \leq 11 - k) \geq 0.95 \iff P(Y \leq 11 - k) \leq 0.05.$$

Scan the table for $Y \sim B(12, 0.35)$:

$$F_Y(0) = 0.0057 \leq 0.05, \quad F_Y(1) = 0.0424 \leq 0.05, \quad F_Y(2) = 0.1513 > 0.05.$$

The largest value of r with $F_Y(r) \leq 0.05$ is $r = 1$, so $11 - k = 1$, giving $k = 10$. Check: $P(X \leq 10) = 1 - F_Y(1) = 1 - 0.0424 = 0.9576 \geq 0.95$, while $P(X \leq 9) = 1 - F_Y(2) = 1 - 0.1513 = 0.8487 < 0.95$. The smallest k is 10.

3.8 Summary of Key Formulas for Table Use

Required probability	Expression using $F(r) = P(X \leq r)$	Table action
$P(X \leq r)$	$F(r)$	Direct reading
$P(X < r) = P(X \leq r - 1)$	$F(r - 1)$	Direct reading (row $r - 1$)
$P(X = r)$	$F(r) - F(r - 1)$	Read two rows, subtract
$P(X \geq r)$	$1 - F(r - 1)$	Read row $r - 1$, subtract from 1
$P(X > r)$	$1 - F(r)$	Read row r , subtract from 1
$P(a \leq X \leq b)$	$F(b) - F(a - 1)$	Read two rows, subtract

DID YOU KNOW?

Historical Note: Tables in the Digital Age

Before electronic calculators and software, printed statistical tables were the only practical way to obtain binomial probabilities for moderate n , and they remain a standard feature of the GCE Advanced Level examination. Modern scientific calculators have built-in binomial cumulative functions, but the syllabus still requires you to read printed tables. Why? Because table reading tests your understanding of the cumulative structure $P(X \leq r)$ and of the complement rules, rather than “black-box” button pressing. Master both: use a calculator to check your answers while revising, but show full table-reading working in the examination.

EXERCISE 3 Practice: Cumulative Tables and Critical Values

Let $X \sim B(20, 0.3)$. Using cumulative binomial tables, find:

1. $P(X \leq 5)$
2. $P(X = 5)$
3. $P(X \geq 5)$
4. $P(4 \leq X \leq 7)$
5. The smallest r such that $P(X \leq r) > 0.99$.
6. The largest r such that $P(X \geq r) > 0.90$.

SOLUTION Exercise 1

From the tables for $n = 20, p = 0.30$:

$F(2) = 0.0355, F(3) = 0.1071, F(4) = 0.2375, F(5) = 0.4164, F(7) = 0.7723, F(10) = 0.9830, F(11) = 0.9949$.

1. $P(X \leq 5) = F(5) = \mathbf{0.4164}$.
2. $P(X = 5) = F(5) - F(4) = 0.4164 - 0.2375 = \mathbf{0.1789}$.
3. $P(X \geq 5) = 1 - F(4) = 1 - 0.2375 = \mathbf{0.7625}$.
4. $P(4 \leq X \leq 7) = F(7) - F(3) = 0.7723 - 0.1071 = \mathbf{0.6652}$.
5. Scanning the column: $F(10) = 0.9830 < 0.99$ and $F(11) = 0.9949 > 0.99$. The smallest r is 11.
6. $P(X \geq r) > 0.90 \iff 1 - F(r-1) > 0.90 \iff F(r-1) < 0.10$. Now $F(2) = 0.0355 < 0.10$ but $F(3) = 0.1071 > 0.10$, so $r-1 = 2$, giving $r = 3$. Check: $P(X \geq 3) = 1 - F(2) = 0.9645 > 0.90$, while $P(X \geq 4) = 1 - F(3) = 0.8929 < 0.90$. The largest r is 3.

4 FITTING A BINOMIAL DISTRIBUTION TO DATA AND APPLICATIONS

In the previous section we learnt how to read cumulative binomial probabilities from tables, so that for a given model $X \sim B(n, p)$ we can answer any question about $P(X \leq r)$. But real life rarely hands us the model directly. A quality-control officer in a Douala factory, a biologist studying families, or a teacher analysing multiple-choice tests first collects data, and only then asks: does the binomial distribution describe this data well, and if so, with which value of p ? This re-

verse process is called **fitting a distribution**, and it is one of the most frequently examined skills in the GCE Advanced Level paper (Lesson 146).

4.1 What Does “Fitting a Distribution” Mean?

EXPERIMENT Counting boys in families

A social researcher in Yaoundé selects 100 families, each having exactly 4 children, and records the number of boys in each family. She obtains:

Number of boys r	0	1	2	3	4
Number of families f	5	28	38	22	7

If the sex of each child is independent of the others and the probability of a boy is the same constant p for every birth, then the number of boys X in a family of 4 should follow $B(4, p)$. The question is: which p , and how well does the model match the data?

DEFINITION Fitting a binomial distribution

To **fit** a binomial distribution $B(n, p)$ to observed data means to:

1. choose the parameters n and p (usually n is known from the context and p is estimated from the data);
2. compute the **expected frequencies** $E_r = N \times P(X = r)$, where N is the total number of observations;
3. compare the expected frequencies with the observed frequencies O_r to judge whether the model is adequate.

The intuition is simple: if the model is correct, then in N repetitions of the experiment, the value r should occur on average $N \times P(X = r)$ times. So the fitted model “predicts” the frequencies, and we check the prediction against reality.

4.2 Estimating p from the Data

Since n is fixed by the situation (here $n = 4$ children), the only unknown is p . We estimate it using the mean of the data, because for $X \sim B(n, p)$ we proved that $E(X) = np$.

PROPERTY Estimating p

If a sample of N observations from an alleged $B(n, p)$ distribution has mean $\bar{x}$, then the natural estimate of p is

$$\hat{p} = \frac{\bar{x}}{n}.$$

Justification. The theoretical mean is np . Fitting consists of making the theoretical model match the data, so we demand that the theoretical mean equal the observed mean: $n\hat{p} = \bar{x}$, giving $\hat{p} = \bar{x}/n$. $\square$

WARNING The classic trap

Do not estimate p as the relative frequency of one particular value. For instance, in the family data, $\frac{38}{100} = 0.38$ is the proportion of families with 2 boys — it is not an estimate of p . The estimate must come from the mean of the whole distribution: $\hat{p} = \bar{x}/n$. Using a single relative frequency is one of the most common errors at the GCE and costs several marks.

4.3 Computing Expected Frequencies

METHOD The 4-step fitting procedure

1. Estimate p . Compute the sample mean $\bar{x} = \frac{\sum f_r r}{\sum f_r}$, then set $\hat{p} = \bar{x}/n$. If $\hat{p}$ is not a tabulated value (e.g. 0.05, 0.10, ..., 0.50), round it to the nearest tabulated value only if you intend to use cumulative binomial tables; otherwise keep the exact value and compute probabilities directly from the formula.
2. Compute the probabilities. Using $P(X = r) = \binom{n}{r} \hat{p}^r (1 - \hat{p})^{n-r}$, find $P(X = r)$ for $r = 0, 1, \dots, n$. Keep at least 4 decimal places during calculations.
3. Multiply by N . The expected frequencies are $E_r = N \times P(X = r)$. Round each E_r to 1 decimal place (or as instructed) and verify that $\sum E_r = N$ (adjust the last decimal if necessary due to rounding).
4. Compare. Place observed O_r and expected E_r side by side in a table (and ideally a bar chart) and comment on the goodness of fit.

KEY POINTS Expected frequencies need not be integers

An expected frequency is a mean number of occurrences, so values like 21.4 or 6.83 are perfectly correct. Keep at least 3–4 decimal places in the probabilities during the calculation and round only at the very end (usually to 1 decimal place). Rounding too early can make the expected frequencies fail to add up to N and destroys accuracy.

4.4 Worked Example: Fitting $B(4, p)$ to the Family Data

WORKED EXAMPLE 11 Fitting a binomial distribution to family data

For the 100 families of 4 children above, fit a binomial distribution and comment on the fit.

SOLUTION

Step 1: Estimate p .

$$\bar{x} = \frac{0(5) + 1(28) + 2(38) + 3(22) + 4(7)}{100} = \frac{0 + 28 + 76 + 66 + 28}{100} = \frac{198}{100} = 1.98.$$


Hence $\hat{p} = \frac{\bar{x}}{n} = \frac{1.98}{4} = 0.495$. Since 0.495 is very close to 0.5 and 0.5 is a tabulated value, we use $\hat{p} = 0.5$ for the fit (this also allows us to use the cumulative binomial tables if needed).

Step 2: Probabilities for $X \sim B(4, 0.5)$. Since $\binom{4}{r} = (1, 4, 6, 4, 1)$ and $p = q = \frac{1}{2}$:

$$P(X = r) = \binom{4}{r} \left(\frac{1}{2}\right)^4 = \binom{4}{r} \times \frac{1}{16}.$$

r	0	1	2	3	4	Total
$P(X = r)$	0.0625	0.25	0.375	0.25	0.0625	1
$E_r = 100 P(X = r)$	6.25	25.0	37.5	25.0	6.25	100
Observed O_r	5	28	38	22	7	100

Step 3: Comment. The expected frequencies are very close to the observed ones: the largest discrepancy is 3 families (for $r = 1$ and $r = 3$). The binomial model $B(4, 0.5)$ fits the data well, supporting the hypothesis that boys and girls are equally likely and that births are independent.



Grouped bar chart: observed frequencies versus expected frequencies from the fitted binomial distribution $B(4, 0.5)$ for 100 families of 4 children.

The bars are almost the same height everywhere: this visual agreement is exactly what a good fit looks like.

4.5 Commenting on the Goodness of Fit

At GCE level, the comparison is made by inspection:

- compute the differences $|O_r - E_r|$;
- if all differences are small relative to the frequencies, conclude that “the binomial model is a good fit”;
- if one or more differences are large, conclude that the fit is poor and suggest why (e.g. p not constant, trials not independent).

DID YOU KNOW?

Measuring the fit precisely: the chi-squared test

Statisticians measure goodness of fit with a single number, the **chi-squared statistic** $\chi^2 = \sum \frac{(O_r - E_r)^2}{E_r}$, and compare it with critical values. For our family data, $\chi^2 = \frac{1.5625}{6.25} + \frac{9}{25} + \frac{0.25}{37.5} + \frac{9}{25} + \frac{0.5625}{6.25} \approx 1.07$, a very small value, confirming an excellent fit. The formal chi-squared test is beyond the GCE syllabus — you only need to compare and comment — but you will meet it at university.

WARNING Exam tip — GCE

Fitting questions typically carry 6 to 8 marks. To collect them all:

1. show the calculation of $\bar{x}$ and state $\hat{p} = \bar{x}/n$ explicitly;
2. present the full probability table $P(X = r)$ for every r from 0 to n , not just the values you find interesting;
3. give expected frequencies to at least 1 decimal place and check they sum to N ;
4. finish with a clear sentence of conclusion (“good fit” / “poor fit, because...”).

4.6 Applications of the Binomial Distribution (Lesson 149)

The binomial model appears whenever we repeat a fixed number of independent trials with the same probability of success. Here are four standard fields of application.

1. Quality control. A factory in Douala produces sachets of detergent, and past experience shows that 2% of sachets are underweight. An inspector samples 20 sachets at random. If X is the number of underweight sachets, then $X \sim B(20, 0.02)$, and

$$P(X \geq 2) = 1 - P(X = 0) - P(X = 1) = 1 - (0.98)^{20} - 20(0.02)(0.98)^{19} \approx 0.0599.$$

If the inspector finds 2 or more defective sachets (an event of probability only about 6% under the model), the batch is rejected and the machine is checked. The binomial model thus provides an objective rule for accepting or rejecting production.

2. Guessing in multiple-choice tests. A candidate answers 10 MCQ questions, each with 4 options, purely at random. The number of correct answers is $X \sim B(10, 0.25)$. Then

$$P(X \geq 5) = 1 - P(X \leq 4) \approx 1 - 0.9219 = 0.0781$$

(from cumulative tables). So a pure guesser scores 5 or more only about 8% of the time: high scores are genuine evidence of knowledge, which is exactly why MCQ marking schemes are fair.

3. Penalty kicks in sports. Suppose a striker for the Indomitable Lions converts 80% of his penalties. In a training session he takes 5 kicks; the number scored is $X \sim B(5, 0.8)$. Then

$$P(X = 5) = (0.8)^5 = 0.3277, \quad P(X \geq 4) = P(X = 4) + P(X = 5) = 5(0.8)^4(0.2) + (0.8)^5 \approx 0.7373.$$

Coaches use such calculations to compare players and to decide who should take the decisive kick.

4. Medical screening. A disease affects 5% of a population. In a random sample of 15 people, the number infected is $X \sim B(15, 0.05)$, so

$$P(X = 0) = (0.95)^{15} \approx 0.4633,$$

meaning that in nearly half of such samples nobody is infected even though the disease is present in the population — a crucial warning against concluding “the village is disease-free” from one small sample.

4.7 When Is the Binomial Model Appropriate?

KEY POINTS Conditions for a binomial model

$X \sim B(n, p)$ is appropriate only if all four conditions hold:

1. a fixed number n of trials;
2. each trial has exactly two outcomes (success/failure);
3. the probability p of success is constant from trial to trial;
4. the trials are independent.

The model fails when, for example:

- **p is not constant:** a basketball player's probability of scoring rises as he "warms up"; a machine wears out during the day so late items are more often defective.
- **Trials are dependent:** drawing balls from an urn without replacement; the health of members of the same family (infections spread between them); questions in an exam that build on each other.
- **n is not fixed:** counting the number of phone calls received in an hour (no fixed number of trials — this will lead us to the Poisson distribution later in the chapter).

A poor fit when fitting $B(n, p)$ to data is often the symptom that one of these conditions is violated. For instance, if observed data show far more extreme values (0 and n) than the fitted binomial predicts, this suggests that p varies between groups of trials.

EXERCISE 4 Fitting and judging a claim

A seed company claims that 75% of its tomato seeds germinate. A farmer plants 200 rows of 5 seeds and records the number germinating per row:

Number germinating r	0	1	2	3	4	5
Number of rows	1	7	30	62	72	28

1. Estimate p from the data and fit a binomial distribution $B(5, \hat{p})$. Compute the expected frequencies (keep $\hat{p}$ exact for the calculation).
2. The company claims $p = 0.75$. Using $p = 0.75$, compute the expected frequencies for $B(5, 0.75)$.
3. Compare both fits with the observed data. Comment on the company's claim.

SOLUTION Exercise 11. Fitting with estimated $\hat{p}$.

$$\bar{x} = \frac{0(1) + 1(7) + 2(30) + 3(62) + 4(72) + 5(28)}{200} = \frac{0 + 7 + 60 + 186 + 288 + 140}{200} = \frac{681}{200} = 3.405.$$

Hence $\hat{p} = \frac{\bar{x}}{n} = \frac{3.405}{5} = 0.681$. Using $p = 0.681$, $q = 0.319$, we compute $P(X = r) = \binom{5}{r}(0.681)^r(0.319)^{5-r}$:

$$P(X = 0) = (0.319)^5 \approx 0.0033,$$

$$E_0 = 200 \times 0.0033 = 0.66 \approx 0.7,$$

$$P(X = 1) = 5(0.681)(0.319)^4 \approx 0.0353,$$

$$E_1 = 200 \times 0.0353 = 7.06 \approx 7.1,$$

$$P(X = 2) = 10(0.681)^2(0.319)^3 \approx 0.1507,$$

$$E_2 = 200 \times 0.1507 = 30.14 \approx 30.1,$$

$$P(X = 3) = 10(0.681)^3(0.319)^2 \approx 0.3217,$$

$$E_3 = 200 \times 0.3217 = 64.34 \approx 64.3,$$

$$P(X = 4) = 5(0.681)^4(0.319) \approx 0.3435,$$

$$E_4 = 200 \times 0.3435 = 68.70 \approx 68.7,$$

$$P(X = 5) = (0.681)^5 \approx 0.1455,$$

$$E_5 = 200 \times 0.1455 = 29.10 \approx 29.1.$$

Sum of $E_r = 0.7 + 7.1 + 30.1 + 64.3 + 68.7 + 29.1 = 200.0$. Table:

r	0	1	2	3	4	5
E_r (fitted $\hat{p} = 0.681$)	0.7	7.1	30.1	64.3	68.7	29.1
O_r	1	7	30	62	72	28

The fit is very good: all expected frequencies are close to observed ones (largest difference ≈ 3.3 for $r = 4$).

2. Testing the claim $p = 0.75$. Using $B(5, 0.75)$: $P(X = r) = \binom{5}{r}(0.75)^r(0.25)^{5-r}$ gives probabilities 0.0010, 0.0146, 0.0879, 0.2637, 0.3955, 0.2373, hence expected frequencies (multiplying by 200):

r	0	1	2	3	4	5
E_r (claim $p = 0.75$)	0.2	2.9	17.6	52.7	79.1	47.5
O_r	1	7	30	62	72	28

3. Conclusion. The fit using the estimated $\hat{p} = 0.681$ is excellent, while the fit using the company's claimed $p = 0.75$ is poor: the model predicts about 48 rows with all 5 seeds germinating, but only 28 were observed, and it underestimates the low counts (30 rows observed with 2 germinating against 17.6 expected). The data suggest a true germination rate near 0.68, so the company's claim of 75% is not supported.

This section closes our study of the binomial distribution as a complete toolkit: probabilities, expectation, recurrence, cumulative tables, and now fitting to real data. We next turn to a close relative — the geometric distribution — which asks not “how many successes in n trials?” but “how many trials until the first success?”

5 THE GEOMETRIC DISTRIBUTION

In the previous section we fitted a binomial distribution to observed data and applied $B(n, p)$ to situations where the number of trials is fixed in advance. But many real questions are of a different kind: we do not ask “how many successes in n trials?” but rather “how long must we wait for the first success?” How many throws of a die until the first six appears? How many items from a production line must be inspected until the first defective one is found? How many penalties until the first goal is scored? These questions lead to a new discrete distribution: the **geometric distribution**. This topic corresponds to **Lessons 147 and 148** of the official syllabus.

5.1 From an Experiment to a Model

EXPERIMENT Waiting for the first six

Take an ordinary fair die and throw it repeatedly, recording the number of the throw on which the first six appears. Sometimes the six comes immediately ($X = 1$); sometimes you wait 4, 7, or even 20 throws. Repeat the experiment many times and draw a bar chart of the frequencies of X . You will observe a characteristic shape: the value 1 is the most frequent, and the frequencies decrease steadily, with an occasional very long wait. This decreasing shape is the signature of the geometric distribution.

Let us compute the probabilities for this experiment. Let $p = \frac{1}{6}$ be the probability of a six on each throw and $q = 1 - p = \frac{5}{6}$. Since the throws are independent:

$$P(X = 1) = p,$$

$$P(X = 2) = P(\text{failure then success}) = qp,$$

$$P(X = 3) = P(\text{failure, failure, success}) = q^2p,$$

$$P(X = r) = q^{r-1}p.$$

The probabilities form a geometric progression with common ratio q — hence the name of the distribution.

5.2 Definition and Conditions

DEFINITION Geometric distribution

A discrete random variable X follows a **geometric distribution** with parameter p ($0 < p < 1$), written

$$X \sim \text{Geo}(p),$$

if X counts the number of independent Bernoulli trials up to and including the first success, where each trial has the same probability p of success. Writing $q = 1 - p$,

$$P(X = r) = q^{r-1}p, \quad r = 1, 2, 3, \dots$$

The conditions for the model are exactly those of a sequence of Bernoulli trials, but with no upper limit on the number of trials:

- each trial results in **success** or **failure**;
- the trials are **independent**;
- the probability of success p is **constant** from trial to trial;
- trials continue until the first success occurs (the number of trials is unlimited).


WARNING Check the formula is a genuine distribution

Note that $\sum_{r=1}^{\infty} q^{r-1}p = p \cdot \frac{1}{1-q} = \frac{p}{p} = 1$, using the sum of the geometric series $1+q+q^2+\dots = \frac{1}{1-q}$.
So the probabilities do add up to 1, as they must.

WARNING Alternative definition

Some textbooks define the geometric distribution as the number of failures before the first success (support $r = 0, 1, 2, \dots$). The GCE Advanced Level syllabus uses the definition above: X is the trial number of the first success (support $r = 1, 2, 3, \dots$). Always check which definition is being used; the formulas for expectation and variance differ.

5.3 The Shape of the Distribution



Bar chart of $X \sim \text{Geo}(0.25)$: the probabilities $P(X=r) = 0.75^{r-1}(0.25)$ decrease geometrically, with a long right tail.

The bar chart shows two essential features: the distribution is always decreasing (so the most likely value of X is 1), and it has a long right tail: very long waits are unlikely but never impossible.

PROPERTY Mode of the geometric distribution

If $X \sim \text{Geo}(p)$, the **mode** of X is always 1, because

$$P(X = r) = q P(X = r - 1) < P(X = r - 1) \quad \text{for all } r \geq 2.$$

5.4 Cumulative Probabilities


A very useful trick: the event “ $X > r$ ” means the first r trials are all failures. By independence this has probability q^r .

PROPERTY Cumulative probabilities

If $X \sim \text{Geo}(p)$ and $q = 1 - p$, then for every positive integer r ,

$$P(X > r) = q^r \quad \text{and} \quad P(X \leq r) = 1 - q^r.$$

$r = 6$ consecutive failures: probability q^6



$P(X > r) = q^r$ is the probability that the first r trials are all failures (F); the first success (S) then occurs later.

5.5 Expectation and Variance**PROPERTY Expectation and variance of $\text{Geo}(p)$**

If $X \sim \text{Geo}(p)$ with $q = 1 - p$, then

$$E(X) = \frac{1}{p} \quad \text{and} \quad \text{Var}(X) = \frac{q}{p^2}.$$

(The derivations use differentiation of geometric series; they are an extension, not required at GCE, but the results must be known and used.)

Sketch of the derivation (extension). $E(X) = \sum_{r \geq 1} r q^{r-1} p = p \sum_{r \geq 1} r q^{r-1}$. Differentiating $\sum_{r \geq 0} q^r = \frac{1}{1-q}$ term by term gives $\sum_{r \geq 1} r q^{r-1} = \frac{1}{(1-q)^2} = \frac{1}{p^2}$, hence $E(X) = \frac{1}{p}$. A similar second differentiation

yields $E(X^2)$ and then $\text{Var}(X) = \frac{q}{p^2}$.

The expectation matches intuition perfectly: if the probability of a six is $\frac{1}{6}$, we expect to wait 6 throws for the first six; if p is small, the expected wait is long.

WORKED EXAMPLE 12 First defective item on a production line

A machine at a soap factory in Douala produces defective items independently with probability 0.04. Let X be the position of the first defective item inspected. Find:

1. the probability that the first defective item is the 5th inspected;
2. the probability that more than 20 items are inspected before the first defective one;
3. the expected number of items inspected and the standard deviation of X .

SOLUTION

Here $X \sim \text{Geo}(0.04)$, so $p = 0.04$ and $q = 0.96$.

1. $P(X = 5) = q^4 p = (0.96)^4 (0.04) \approx 0.8493 \times 0.04 \approx 0.0340$.
2. $P(X > 20) = q^{20} = (0.96)^{20} \approx 0.4420$.
3. $E(X) = \frac{1}{p} = \frac{1}{0.04} = 25$ items, and

$$\text{Var}(X) = \frac{q}{p^2} = \frac{0.96}{0.0016} = 600, \quad \sigma = \sqrt{600} \approx 24.5.$$

Note the large standard deviation: the waiting time is very spread out, as the long right tail suggests.

5.6 The Memoryless Property

Suppose you have already thrown the die 10 times without a six. Does the die “owe” you a six? Of course not — the remaining wait has exactly the same distribution as at the start. This is the **memoryless property**, and the geometric distribution is the only discrete distribution with this property.

PROPERTY Memoryless property

If $X \sim \text{Geo}(p)$, then for all positive integers m and n ,

$$P(X > m + n \mid X > m) = P(X > n).$$

Verification. Using conditional probability and $P(X > r) = q^r$:

$$P(X > m + n \mid X > m) = \frac{P(X > m + n)}{P(X > m)} = \frac{q^{m+n}}{q^m} = q^n = P(X > n).$$

So knowledge that the first m trials were failures changes nothing about the additional waiting time.

EXAMPLE Checking the memoryless property numerically

Let $X \sim \text{Geo}(0.25)$ (e.g. waiting for the first “head” when spinning a biased coin with $P(\text{success}) = 0.25$). Take $m = 4$, $n = 3$:

$$P(X > 7 \mid X > 4) = \frac{(0.75)^7}{(0.75)^4} = (0.75)^3 = 0.4219 = P(X > 3).$$

Given that 4 failures have already occurred, the probability of waiting more than 3 further trials is the same as the original probability of waiting more than 3 trials.

5.7 Geometric versus Binomial: Do Not Confuse Them

WARNING Common mistake: $\text{Geo}(p)$ versus $B(n, p)$

The two distributions answer different questions about the same kind of trials:

- $B(n, p)$: the number of trials n is fixed; we count the number of successes. The formula involves $\binom{n}{r}$.
- $\text{Geo}(p)$: the number of trials is not fixed; we count the trial number of the first success. The formula is $q^{r-1}p$, with no binomial coefficient.

Exam trap: “the probability that the first success occurs on the 5th trial” is q^4p (geometric), not $\binom{5}{1}pq^4$ (which is the probability of exactly one success in 5 trials).

METHOD Exam tip (GCE): “first success after the k -th trial”

When a GCE question asks for “the probability that the first success occurs after the k -th trial” (or “more than k trials are needed”):

1. Identify p and compute $q = 1 - p$.
2. Recognise that “first success after the k -th trial” means the first k trials all fail.
3. Write directly $P(X > k) = q^k$ — no summation needed.

Similarly, “first success on or before the k -th trial” gives $P(X \leq k) = 1 - q^k$.

WORKED EXAMPLE 13 GCE-style problem: waiting for a goal

A footballer scores from a free kick with probability 0.15, independently of previous attempts. Let X be the number of free kicks taken up to and including the first goal.

1. State the distribution of X and the conditions required.
2. Find the probability that his first goal comes on his 4th free kick.
3. Find the probability that he needs more than 10 free kicks to score.
4. Find the expected number of free kicks needed and $\text{Var}(X)$.
5. Given that he has already missed 6 free kicks, find the probability that he needs more than 4 further attempts.

SOLUTION

1. $X \sim \text{Geo}(0.15)$, provided attempts are independent and the probability of scoring is constant at $p = 0.15$ for each attempt. Here $q = 0.85$.
2. $P(X = 4) = q^3 p = (0.85)^3 (0.15) \approx 0.6141 \times 0.15 \approx 0.0921$.
3. $P(X > 10) = q^{10} = (0.85)^{10} \approx 0.1969$.
4. $E(X) = \frac{1}{0.15} \approx 6.67$ free kicks; $\text{Var}(X) = \frac{0.85}{0.15^2} = \frac{0.85}{0.0225} \approx 37.78$.
5. By the memoryless property,

$$P(X > 6 + 4 \mid X > 6) = P(X > 4) = q^4 = (0.85)^4 \approx 0.5220.$$

5.8 Summary and Practice**KEY POINTS** Key points on the geometric distribution

- $X \sim \text{Geo}(p)$: X = number of independent trials up to and including the first success;
 $P(X = r) = q^{r-1} p$, $r = 1, 2, 3, \dots$
- $P(X > r) = q^r$ and $P(X \leq r) = 1 - q^r$.
- $E(X) = \frac{1}{p}$, $\text{Var}(X) = \frac{q}{p^2}$, mode = 1.
- Memoryless: $P(X > m + n \mid X > m) = P(X > n)$.
- Do not confuse with $B(n, p)$: fixed n (binomial) versus waiting time for the first success (geometric).

EXERCISE 5 On the geometric distribution

1. A fair die is thrown repeatedly. Let X be the throw on which the first six appears. Find (a) $P(X = 3)$; (b) $P(X > 6)$; (c) $E(X)$ and $\text{Var}(X)$.
2. In Yaoundé, a call centre agent makes a sale on each call with probability 0.1, independently of other calls. Find the probability that her first sale (a) occurs on the 8th call; (b) occurs after the 15th call. What is her expected number of calls for the first sale?
3. A student guesses every answer of a multiple-choice quiz with 4 options per question. Find the probability that the first correct guess comes (a) on the 2nd question; (b) after the 5th question. Given that the first 4 answers were wrong, find the probability that more than 3 further questions are needed for the first correct answer.

SOLUTION Exercise

1. $X \sim \text{Geo}\left(\frac{1}{6}\right)$, $q = \frac{5}{6}$. (a) $P(X = 3) = \left(\frac{5}{6}\right)^2 \frac{1}{6} = \frac{25}{216} \approx 0.1157$. (b) $P(X > 6) = \left(\frac{5}{6}\right)^6 \approx 0.3349$. (c) $E(X) = 6$, $\text{Var}(X) = \frac{5/6}{1/36} = 30$.
2. $X \sim \text{Geo}(0.1)$, $q = 0.9$. (a) $P(X = 8) = (0.9)^7(0.1) \approx 0.0478$. (b) $P(X > 15) = (0.9)^{15} \approx 0.2059$. $E(X) = \frac{1}{0.1} = 10$ calls.
3. $p = \frac{1}{4}$, $q = \frac{3}{4}$. (a) $P(X = 2) = \frac{3}{4} \cdot \frac{1}{4} = \frac{3}{16} = 0.1875$. (b) $P(X > 5) = \left(\frac{3}{4}\right)^5 \approx 0.2373$. Memoryless property: $P(X > 4 + 3 \mid X > 4) = P(X > 3) = \left(\frac{3}{4}\right)^3 \approx 0.4219$.

DID YOU KNOW? Why “geometric”?

The distribution takes its name from the geometric series: the probabilities $p, qp, q^2p, q^3p, \dots$ form a geometric progression, and their sum $p(1 + q + q^2 + \dots) = \frac{p}{1-q} = 1$ is exactly the geometric series formula you met in Pure Mathematics. The geometric distribution is also the discrete analogue of the exponential distribution, which models waiting times in continuous time — both share the memoryless property.


6 THE POISSON DISTRIBUTION: CHARACTERISTICS, EXPECTATION AND VARIANCE

In the previous section we studied the **geometric distribution**, which counts the number of trials needed to obtain the first success. Trials there were clearly separated: one attempt, then another, then another. But many real-life situations do not come in separate trials at all. Think of the phone calls arriving at the MTN call centre in Douala during one hour, the accidents occurring along the Douala–Yaoundé highway during one month, or the misprints on the pages of a textbook. There is no fixed number n of “trials”; events simply happen, scattered at random along a continuous stretch of time or space. For such situations we need a new model: the **Poisson distribution**, named after the French mathematician Siméon Denis Poisson.

6.1 Random Events in a Continuum: the Poisson Process

EXPERIMENT Observing calls at a call centre

Imagine you sit next to the switchboard of a small business in Yaoundé and you record the exact times at which calls arrive during one hour, say from 9:00 to 10:00. On a typical morning you might observe something like this: calls at 9:03, 9:11, 9:12, 9:27, 9:41, 9:58 — six calls in all. The next day, same hour, you might count only three calls; the day after, nine. The calls do not arrive at regular intervals like a clock ticking; they arrive at random. Yet, over many days, the average number of calls per hour is remarkably stable, say about 5 calls per hour. This combination — individual unpredictability but long-term stability of the average rate — is exactly the signature of a **Poisson process**.



Random phone calls over one hour: the instants are unpredictable, calls arrive one at a time and independently, but the average rate per hour stays constant.

Other classic examples of events occurring randomly in a continuum include:

- accidents on a given stretch of the Douala–Yaoundé highway per month;
- misprints per page of a printed book (here the “continuum” is space, not time);
- radioactive decays per second detected by a counter;
- customers joining the queue at a bank agency per ten minutes;

- goals scored per football match in a championship.

METHOD The four conditions for a Poisson process

Before modelling a situation by a Poisson distribution, check that the events occur:

1. Randomly — the exact instant (or position) of each event is unpredictable;
2. Independently — the occurrence of one event does not affect the probability of another (one accident does not “cause” or “prevent” the next);
3. Singly — events occur one at a time; two events cannot happen at exactly the same instant (or same point);
4. At a constant average rate — the mean number of events per **unit interval**, denoted λ , stays the same throughout the period considered.

If all four conditions hold, the number of events in a fixed interval follows a Poisson distribution.

WARNING When Poisson does not apply

Buses arriving at a stop according to a fixed timetable fail condition 1 (not random). Goals in a match where scoring a goal encourages the attacking team may fail condition 2 (not independent). Customers arriving in groups at a restaurant fail condition 3 (not single). Calls at a call centre over a whole day fail condition 4 if the rate is much higher at midday than at midnight — in that case, split the day into periods with roughly constant rates.

6.2 Definition and Probability Formula

DEFINITION Poisson distribution

A discrete random variable X follows a **Poisson distribution** with parameter $\lambda > 0$, written $X \sim \text{Po}(\lambda)$, if X takes values in $\{0, 1, 2, 3, \dots\}$ with

$$P(X = r) = \frac{e^{-\lambda} \lambda^r}{r!}, \quad r = 0, 1, 2, 3, \dots$$

Here λ is the mean number of events per unit interval (per hour, per page, per kilometre, ...). The **unit interval** is the fixed stretch of time or space (e.g. one hour, one page, one kilometre) for which λ is defined.

Note immediately a useful special case:

$$P(X = 0) = \frac{e^{-\lambda} \lambda^0}{0!} = e^{-\lambda},$$

since $\lambda^0 = 1$ and $0! = 1$. So the probability of no event at all in the interval is simply $e^{-\lambda}$.

PROPERTY The probabilities sum to 1

For $X \sim \text{Po}(\lambda)$, $\sum_{r=0}^{\infty} P(X = r) = 1$.

Proof (examinable and instructive). Recall the exponential series, valid for every real number x :

$$e^x = \sum_{r=0}^{\infty} \frac{x^r}{r!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$$

Then

$$\sum_{r=0}^{\infty} P(X = r) = \sum_{r=0}^{\infty} \frac{e^{-\lambda} \lambda^r}{r!} = e^{-\lambda} \sum_{r=0}^{\infty} \frac{\lambda^r}{r!} = e^{-\lambda} e^{\lambda} = e^0 = 1.$$

Moreover each term is positive, so the formula defines a genuine probability distribution. ■

EXAMPLE First calculations with $X \sim \text{Po}(3)$

A switchboard receives on average 3 calls per ten minutes. Let X be the number of calls in ten minutes, so $X \sim \text{Po}(3)$.

$$P(X = 0) = e^{-3} \approx 0.0498,$$

$$P(X = 2) = \frac{e^{-3} 3^2}{2!} = \frac{9}{2} e^{-3} \approx 0.2240,$$

$$P(X = 5) = \frac{e^{-3} 3^5}{5!} = \frac{243}{120} e^{-3} \approx 0.1008.$$

WARNING The forgotten factorial

The most frequent error at the GCE is to write $P(X = r) = e^{-\lambda} \lambda^r$ and to forget the $r!$ in the denominator. Without $r!$ the probabilities do not sum to 1 and every numerical answer is wrong. A safe habit: write the formula in full on your script before substituting any number, and compute $r!$ explicitly ($3! = 6$, $4! = 24$, $5! = 120$).

6.3 Expectation and Variance: the Distinctive Feature $E(X) = \text{Var}(X) = \lambda$

PROPERTY Mean and variance of the Poisson distribution

If $X \sim \text{Po}(\lambda)$, then

$$E(X) = \lambda \quad \text{and} \quad \text{Var}(X) = \lambda.$$

The proof of $E(X) = \lambda$ is examinable; the variance proof follows the same technique and is also examinable.

Proof that $E(X) = \lambda$. By definition,

$$E(X) = \sum_{r=0}^{\infty} r P(X = r) = \sum_{r=0}^{\infty} r \frac{e^{-\lambda} \lambda^r}{r!}.$$

The term with $r = 0$ vanishes, and for $r \geq 1$ we simplify $\frac{r}{r!} = \frac{1}{(r-1)!}$:

$$E(X) = e^{-\lambda} \sum_{r=1}^{\infty} \frac{\lambda^r}{(r-1)!} = e^{-\lambda} \lambda \sum_{r=1}^{\infty} \frac{\lambda^{r-1}}{(r-1)!}.$$

Putting $s = r - 1$, the sum becomes $\sum_{s=0}^{\infty} \frac{\lambda^s}{s!} = e^{\lambda}$. Hence

$$E(X) = e^{-\lambda} \lambda e^{\lambda} = \lambda. \quad \blacksquare$$

Proof that $\text{Var}(X) = \lambda$. We first compute $E(X(X-1))$:

$$E(X(X-1)) = \sum_{r=0}^{\infty} r(r-1) \frac{e^{-\lambda} \lambda^r}{r!}.$$

Terms for $r = 0, 1$ are zero. For $r \geq 2$, $\frac{r(r-1)}{r!} = \frac{1}{(r-2)!}$:

$$E(X(X-1)) = e^{-\lambda} \sum_{r=2}^{\infty} \frac{\lambda^r}{(r-2)!} = e^{-\lambda} \lambda^2 \sum_{r=2}^{\infty} \frac{\lambda^{r-2}}{(r-2)!} = e^{-\lambda} \lambda^2 e^{\lambda} = \lambda^2.$$

Now $E(X^2) = E(X(X-1)) + E(X) = \lambda^2 + \lambda$. Therefore

$$\text{Var}(X) = E(X^2) - (E(X))^2 = \lambda^2 + \lambda - \lambda^2 = \lambda. \quad \blacksquare$$

KEY POINTS A quick diagnostic for Poisson data

The equality mean = variance is the signature of the Poisson distribution. If you count events in many equal intervals and find a sample mean of about 4.1 and a sample variance of about 4.0, a Poisson model with $\lambda \approx 4$ is plausible. If the variance is much larger than the mean (over-dispersion) or much smaller, the Poisson model should be rejected. This is the first test used when fitting a distribution to real data.

6.4 Recurrence Formula and Mode of the Poisson Distribution

PROPERTY Recurrence formula

For $X \sim \text{Po}(\lambda)$ with $\lambda > 0$, the probabilities satisfy

$$P(X = r) = \frac{\lambda}{r} P(X = r - 1), \quad r = 1, 2, 3, \dots$$

with $P(X = 0) = e^{-\lambda}$.

Proof.

$$\frac{P(X = r)}{P(X = r - 1)} = \frac{e^{-\lambda} \lambda^r / r!}{e^{-\lambda} \lambda^{r-1} / (r-1)!} = \frac{\lambda}{r}. \quad \blacksquare$$

This recurrence allows rapid computation of successive probabilities without recalculating factorials and powers each time.

PROPERTY Mode of the Poisson distribution

The **mode** (most probable value) of $X \sim \text{Po}(\lambda)$ is:

- λ and $\lambda - 1$ (two modes) if λ is a positive integer;
- $[\lambda]$ (the greatest integer less than λ) if λ is not an integer.

Justification. From the recurrence, $P(X = r) \geq P(X = r - 1)$ iff $\frac{\lambda}{r} \geq 1$, i.e. $r \leq \lambda$. So probabilities increase while $r < \lambda$, are equal at $r = \lambda$ (if λ is integer), and decrease for $r > \lambda$.

EXAMPLE Using the recurrence and finding the mode

Let $X \sim \text{Po}(4.5)$.


1. Compute $P(X = 0)$, $P(X = 1)$, $P(X = 2)$, $P(X = 3)$ using the recurrence.
2. State the mode of the distribution.

Solution.

1. $P(X = 0) = e^{-4.5} \approx 0.0111$. $P(X = 1) = \frac{4.5}{1} \times 0.0111 = 0.0500$. $P(X = 2) = \frac{4.5}{2} \times 0.0500 = 0.1125$.
 $P(X = 3) = \frac{4.5}{3} \times 0.1125 = 0.1688$.
2. Since $\lambda = 4.5$ is not an integer, the mode is $[4.5] = 4$.

6.5 The Shape of the Distribution as λ Grows

For small λ , the Poisson distribution is strongly skewed to the right: small values of X dominate. As λ increases, the distribution spreads out and becomes more and more symmetric, looking increasingly like a bell shape centred near λ .



Bar charts of Po(1), Po(4) and Po(9): as λ grows, the distribution moves right, spreads out and becomes more symmetric.

6.6 Uses of the Poisson Distribution

The Poisson distribution is the standard model for rare or random events in a continuum:

- **Telecommunications:** calls arriving at an exchange, text messages per minute at a network node;
- **Traffic:** vehicles passing a checkpoint on the Douala–Yaoundé road per minute in light traffic, accidents per month at a junction;
- **Biology and medicine:** bacteria per drop of water under a microscope, mutations per generation, patients arriving at the emergency ward of the Laquintinie Hospital per hour;
- **Insurance:** claims of a given type per week received by an insurance company;
- **Quality control:** defects per metre of cloth, misprints per page.

WORKED EXAMPLE 14 Accidents on the highway

On a certain stretch of the Douala–Yaoundé highway, accidents occur randomly, independently, singly and at a constant average rate of 2 accidents per week. Let X be the number of accidents in a given week.

1. State the distribution of X and give $E(X)$ and $\text{Var}(X)$.
2. Find the probability that in a given week there are (i) no accidents; (ii) exactly 3 accidents.
3. Find the probability of at least one accident in a week.

SOLUTION

Solution.

1. The four Poisson conditions are given, with rate $\lambda = 2$ per week, so $X \sim \text{Po}(2)$. Hence $E(X) = 2$ and $\text{Var}(X) = 2$.
2. Using $P(X = r) = \frac{e^{-2} 2^r}{r!}$:
 - (i) $P(X = 0) = e^{-2} \approx 0.1353$.
 - (ii) $P(X = 3) = \frac{e^{-2} 2^3}{3!} = \frac{8}{6} e^{-2} \approx 0.1804$.
3. Using the complement:

$$P(X \geq 1) = 1 - P(X = 0) = 1 - e^{-2} \approx 1 - 0.1353 = 0.8647.$$

So there is about an 86.5% chance of at least one accident in any given week on this stretch.

EXERCISE 6 Misprints in a textbook

Misprints in a mathematics textbook occur randomly and independently at an average rate of 1.5 misprints per page. A page is chosen at random. Find the probability that it contains (a) no misprint; (b) exactly two misprints; (c) more than two misprints. Give your answers to 4 decimal places.

SOLUTION Exercise 1

Let $X \sim \text{Po}(1.5)$, so $P(X = r) = \frac{e^{-1.5}(1.5)^r}{r!}$ and $e^{-1.5} \approx 0.2231$.

- (a) $P(X = 0) = e^{-1.5} \approx 0.2231$.
- (b) $P(X = 2) = \frac{e^{-1.5}(1.5)^2}{2!} = 1.125 e^{-1.5} \approx 0.2510$.
- (c) $P(X = 1) = 1.5 e^{-1.5} \approx 0.3347$, so

$$P(X > 2) = 1 - (P(0) + P(1) + P(2)) \approx 1 - (0.2231 + 0.3347 + 0.2510) = 0.1912.$$

DID YOU KNOW? A law born from miscarriages of justice

Siméon Denis Poisson introduced this distribution in 1837 in a work on the probability of judgements in criminal cases — he wanted to model the (rare) wrongful convictions in French courts. The distribution later became famous through a striking data set: the num-

ber of Prussian cavalry soldiers killed each year by horse kicks, recorded over 20 years in several regiments by Ladislaus Bortkiewicz (1898). The observed frequencies matched a Poisson distribution remarkably well, convincing statisticians of its power to describe rare, random events.

KEY POINTS Key points of this section

- A Poisson process requires events that are **random**, **independent**, **single** and occur at a **constant average rate** λ per **unit interval**.
- $X \sim \text{Po}(\lambda)$: $P(X = r) = \frac{e^{-\lambda} \lambda^r}{r!}$, $r = 0, 1, 2, \dots$; in particular $P(X = 0) = e^{-\lambda}$.
- The probabilities sum to 1 because $\sum \lambda^r / r! = e^\lambda$.
- $E(X) = \text{Var}(X) = \lambda$: mean equals variance, the Poisson signature.
- Recurrence: $P(X = r) = \frac{\lambda}{r} P(X = r - 1)$ for $r \geq 1$.
- Mode: λ and $\lambda - 1$ if $\lambda \in \mathbb{N}^+$; otherwise $\lfloor \lambda \rfloor$.
- Never forget the $r!$ in the denominator, and use $P(X \geq 1) = 1 - e^{-\lambda}$ for “at least one event”.

7 RANDOM EVENTS, UNIT INTERVAL, RECURRENCE FORMULA AND MODE OF THE POISSON DISTRIBUTION

In the previous section we defined the Poisson distribution $\text{Po}(\lambda)$, established its probability mass function

$$P(X = r) = \frac{e^{-\lambda} \lambda^r}{r!}, \quad r = 0, 1, 2, \dots$$

and proved that both its expectation and its variance equal λ . We now go deeper into the structure of the Poisson model: what “random events” really look like, how the parameter λ behaves when we change the interval of observation, a powerful recurrence trick for computing probabilities quickly, and how to locate the **mode** of the distribution. These ideas are heavily examined at GCE Advanced Level.

7.1 The Distribution of Random Events

EXPERIMENT Are random events regularly spaced?

Suppose calls arrive at an MTN call centre in Douala at an average rate of 4 calls per minute. Many students imagine that the calls arrive “evenly”: roughly one call every 15 seconds.

This is wrong. Divide the observation period into blocks of 6 seconds and record, for each block, how many calls arrived. A typical simulation of a Poisson process with mean 4 calls per minute (i.e. mean 0.4 call per 6-second block) gives a pattern like:

Second block	1–6	7–12	13–18	19–24	25–30	31–36	37–42	43–48	49–54	55–60
Calls	1	0	2	0	0	1	0	3	1	0

The total is 8 calls in this simulated stretch of 2 minutes (an average of 4 per minute), but the arrivals are clustered and irregular: sometimes three calls arrive within six seconds, sometimes nothing happens for a long stretch.

The lesson is fundamental: a **constant average rate** does **not** mean **regular spacing**. The Poisson model assumes that events occur:

- **singly** — two events cannot occur at exactly the same instant;
- **independently** — the occurrence of one event does not affect the probability of the next;
- **at a constant average rate** λ per unit interval — the rate does not change over time or space;
- **randomly** — the numbers of events in disjoint intervals are independent random variables.

Under these conditions, the number X of events in a fixed interval follows $Po(\lambda)$. The irregularity we observe (gaps and clusters) is precisely what randomness looks like; regular spacing would actually contradict the Poisson model.

WARNING Uniform rate $\neq$ uniform spacing

Saying “events occur at a uniform rate” means the average rate λ is constant. It does not mean events are evenly spaced. Examiners love the question: “State the conditions for a Poisson model.” The answer is: events occur singly, independently, randomly, and at a constant mean rate.

7.2 The Unit Interval: Scaling the Parameter


A Poisson variable is always defined relative to an interval: per minute, per hour, per metre, per page. If we change the interval, the parameter must change proportionally. Intuitively: if a road has on average 2 potholes per kilometre, then a 5 km stretch has on average 10 potholes.

PROPERTY Scaling rule for the Poisson parameter

If $X \sim Po(\lambda)$ counts events per unit interval, then the number Y of events in an interval of length t units satisfies

$$Y \sim Po(\lambda t).$$

The Poisson rate is **proportional** to the length (or duration, or area) of the interval. (Proof not examinable; it follows from the independence of events in disjoint intervals.)



A time axis split into unit (1-minute) intervals, each carrying a $Po(\lambda)$ count. Over t consecutive units the counts add, giving $Po(t\lambda)$.

EXAMPLE Changing the unit interval

Calls arrive at an office at an average rate of 3 per minute, following a Poisson distribution.

- Per minute: $X \sim Po(3)$.
- Per 5 minutes: $Y \sim Po(3 \times 5) = Po(15)$.
- Per hour: $Z \sim Po(3 \times 60) = Po(180)$.
- Per 30 seconds: $W \sim Po(3 \times 0.5) = Po(1.5)$.

Similarly, if defects occur along a cable at 0.4 per metre, then per 10 metres the count is $Po(0.4 \times 10) = Po(4)$, and per 25 cm it is $Po(0.4 \times 0.25) = Po(0.1)$.

WARNING The classic trap: forgetting to scale λ

If the question gives a rate per minute but asks about a five-minute period, you must use $\lambda = 5 \times (\text{rate per minute})$. Using the original λ with the new interval is the single most common error in Poisson questions. Always write, as your first line: "In a ... interval, $\lambda = \dots$ ".

PROPERTY Sum of independent Poisson variables

If $X \sim Po(\lambda_1)$ and $Y \sim Po(\lambda_2)$ are independent, then

$$X + Y \sim Po(\lambda_1 + \lambda_2).$$

Justification (instructive). Using independence,

$$\begin{aligned}
 P(X + Y = n) &= \sum_{r=0}^n P(X = r) P(Y = n - r) = \sum_{r=0}^n \frac{e^{-\lambda_1} \lambda_1^r}{r!} \cdot \frac{e^{-\lambda_2} \lambda_2^{n-r}}{(n-r)!} \\
 &= \frac{e^{-(\lambda_1 + \lambda_2)}}{n!} \sum_{r=0}^n \frac{n!}{r!(n-r)!} \lambda_1^r \lambda_2^{n-r} = \frac{e^{-(\lambda_1 + \lambda_2)}}{n!} \sum_{r=0}^n \binom{n}{r} \lambda_1^r \lambda_2^{n-r} \\
 &= \frac{e^{-(\lambda_1 + \lambda_2)} (\lambda_1 + \lambda_2)^n}{n!},
 \end{aligned}$$

by the binomial theorem. This is exactly the $\text{Po}(\lambda_1 + \lambda_2)$ probability mass function. ■

EXAMPLE Combining two Poisson sources

A hospital receives emergency cases from Bonaberi at 2 per hour and from Deido at 3 per hour, independently. The total number of emergencies per hour is $T \sim \text{Po}(2 + 3) = \text{Po}(5)$. Hence

$$P(T = 0) = e^{-5} \approx 0.0067, \quad P(T \geq 1) = 1 - e^{-5} \approx 0.9933.$$

7.3 The Recurrence Formula

Computing $P(X = r) = \frac{e^{-\lambda} \lambda^r}{r!}$ from scratch for each r is tedious. Fortunately, consecutive probabilities are linked by a simple ratio.

PROPERTY Recurrence formula for the Poisson distribution

If $X \sim \text{Po}(\lambda)$, then for all $r \geq 0$,

$$P(X = r + 1) = \frac{\lambda}{r + 1} P(X = r).$$

(The derivation is examinable and takes two lines.)

Derivation.

$$\frac{P(X = r + 1)}{P(X = r)} = \frac{e^{-\lambda} \lambda^{r+1} / (r + 1)!}{e^{-\lambda} \lambda^r / r!} = \frac{\lambda^{r+1}}{\lambda^r} \cdot \frac{r!}{(r + 1)!} = \lambda \cdot \frac{1}{r + 1} = \frac{\lambda}{r + 1}. \quad \blacksquare$$

METHOD Generating a Poisson probability table by recurrence

1. Compute the starting value $P(X = 0) = e^{-\lambda}$.
2. Multiply successively: $P(1) = \lambda P(0)$, then $P(2) = \frac{\lambda}{2} P(1)$, then $P(3) = \frac{\lambda}{3} P(2)$, and so on.
3. Stop when probabilities become negligible (less than 0.0001, say).
4. Check: the probabilities should sum to 1 (up to rounding).

WORKED EXAMPLE 15 Building the table of $Po(4)$

Let $X \sim Po(4)$. Using the recurrence formula, find $P(X = r)$ for $r = 0, 1, \dots, 8$, giving answers to 4 decimal places.

SOLUTION

Solution. Start with $P(0) = e^{-4} = 0.01832$. Then apply $P(r+1) = \frac{4}{r+1}P(r)$:

$$\begin{aligned} P(1) &= \frac{4}{1}(0.01832) = 0.07326, & P(2) &= \frac{4}{2}(0.07326) = 0.14653, \\ P(3) &= \frac{4}{3}(0.14653) = 0.19537, & P(4) &= \frac{4}{4}(0.19537) = 0.19537, \\ P(5) &= \frac{4}{5}(0.19537) = 0.15629, & P(6) &= \frac{4}{6}(0.15629) = 0.10420, \\ P(7) &= \frac{4}{7}(0.10420) = 0.05954, & P(8) &= \frac{4}{8}(0.05954) = 0.02977. \end{aligned}$$

Notice that $P(3) = P(4)$: the probabilities stop increasing exactly at $r = 4$. This is no accident — it reveals the mode, which we study next.

7.4 The Mode of the Poisson Distribution

The **mode** is the value of r with the greatest probability — the most likely number of events. The recurrence formula tells us exactly where the probabilities rise and where they fall:

$$P(X = r+1) \geq P(X = r) \iff \frac{\lambda}{r+1} \geq 1 \iff r+1 \leq \lambda.$$

So the probabilities **increase** while $r+1 < \lambda$ and **decrease** once $r+1 > \lambda$. Two cases arise.

PROPERTY Mode of the Poisson distribution

Let $X \sim Po(\lambda)$.

- If λ is not an integer, there is a unique mode at $r = \lfloor \lambda \rfloor$ (the greatest integer not exceeding λ).
- If λ is an integer, there are two equal modes, at $r = \lambda - 1$ and $r = \lambda$, with $P(X = \lambda - 1) = P(X = \lambda)$.


(Proof examinable: it is the inequality $r+1 \leq \lambda$ above.)

Indeed, when λ is an integer, $P(X = \lambda) = \frac{\lambda}{\lambda} P(X = \lambda - 1) = P(X = \lambda - 1)$, giving the double mode.

METHOD Finding the mode in an exam

1. Look at λ .

2. If λ is an integer: the modes are $\lambda - 1$ and λ .
3. If λ is not an integer: the mode is $[\lambda]$.
4. To justify the answer (when asked), compare ratios: show $\frac{P(X = r + 1)}{P(X = r)} = \frac{\lambda}{r + 1}$ is ≥ 1 up to the mode and ≤ 1 after it.



Bar chart of $Po(4)$. Since $\lambda = 4$ is an integer, the two modes $r = 3$ and $r = 4$ (highlighted) share the greatest probability, 0.1954.

WORKED EXAMPLE 16 Locating the most likely number of events

The number of accidents per week on a stretch of the Douala–Yaoundé highway follows a Poisson distribution with mean 2.5.

- Find the most likely number of accidents in a week, justifying your answer.
- Find the most likely number of accidents in a fortnight (two weeks).

SOLUTION

Solution. (a) Here $\lambda = 2.5$, not an integer, so the mode is $[2.5] = 2$. Justification using the recurrence ratio:

$$\frac{P(X = r + 1)}{P(X = r)} = \frac{2.5}{r + 1} \geq 1 \iff r + 1 \leq 2.5 \iff r \leq 1.5.$$

So probabilities increase for $r = 0 \rightarrow 1 \rightarrow 2$ and decrease from $r = 2$ onwards: the mode is $r = 2$, with $P(X = 2) = \frac{e^{-2.5}(2.5)^2}{2!} \approx 0.2565$.

(b) In two weeks the parameter scales: $Y \sim \text{Po}(2 \times 2.5) = \text{Po}(5)$. Since 5 is an integer, there are two modes: $r = 4$ and $r = 5$, with

$$P(Y = 4) = P(Y = 5) = \frac{e^{-5}5^4}{4!} \approx 0.1755.$$

Note how the change of unit interval turns a single-mode problem into a double-mode problem.

EXERCISE 7 Practice

1. Bacteria occur in a water sample at an average rate of 1.5 per millilitre. Write down the distribution of the number of bacteria in (a) 1 ml, (b) 4 ml, (c) 200 ml.
2. Given $X \sim \text{Po}(6)$, use the recurrence formula to compute $P(X = r)$ for $r = 0, 1, \dots, 7$ (take $e^{-6} = 0.002479$), and state the mode(s).
3. The number of goals scored per match by a team is $\text{Po}(1.8)$, independently from match to match. Find the distribution of the total number of goals in 3 matches and its most likely value.

SOLUTION Exercise 1

(a) $\text{Po}(1.5)$; (b) $\text{Po}(1.5 \times 4) = \text{Po}(6)$; (c) $\text{Po}(1.5 \times 200) = \text{Po}(300)$.

SOLUTION Exercise 2

$P(0) = 0.002479$; then $P(1) = 6P(0) = 0.014874$, $P(2) = 3P(1) = 0.044622$, $P(3) = 2P(2) = 0.089244$, $P(4) = 1.5P(3) = 0.133866$, $P(5) = 1.2P(4) = 0.160639$, $P(6) = 1 \times P(5) = 0.160639$, $P(7) = \frac{6}{7}P(6) = 0.137691$. Since $\lambda = 6$ is an integer, the modes are $r = 5$ and $r = 6$ (equal probabilities, ≈ 0.1606).

SOLUTION Exercise 3

Total goals $T \sim \text{Po}(3 \times 1.8) = \text{Po}(5.4)$. Since 5.4 is not an integer, the unique mode is $[5.4] = 5$ goals.

KEY POINTS Key points of this section

- A constant average rate does not imply regular spacing: Poisson events occur singly, independently and randomly, producing natural gaps and clusters.
- Scaling: if $X \sim \text{Po}(\lambda)$ per unit interval, then over t units the count is $\text{Po}(\lambda t)$. Always rescale λ first.
- Sums: independent $X \sim \text{Po}(\lambda_1)$, $Y \sim \text{Po}(\lambda_2)$ give $X + Y \sim \text{Po}(\lambda_1 + \lambda_2)$.
- Recurrence: $P(X = r + 1) = \frac{\lambda}{r + 1} P(X = r)$, starting from $P(X = 0) = e^{-\lambda}$.
- Mode: $\lfloor \lambda \rfloor$ if $\lambda \notin \mathbb{Z}$; double mode at $\lambda - 1$ and λ if $\lambda \in \mathbb{Z}$.

8 POISSON APPROXIMATION TO THE BINOMIAL AND APPLICATIONS OF THE POISSON DISTRIBUTION

In the previous section we studied the Poisson distribution in its own right: how it models random events in a unit interval, its recurrence formula $P(X = k + 1) = \frac{\lambda}{k + 1} P(X = k)$ and its mode. We now close the loop by answering a natural question: where does the Poisson distribution come from? The answer is beautiful and extremely useful at the GCE: the Poisson distribution is what the binomial distribution becomes when trials are very many but successes are very rare. This gives us a powerful shortcut, and it also explains why the Poisson model fits so many real-life phenomena, from machine breakdowns in a Douala factory to goals scored in an Elite One football match.

8.1 Motivation: When Binomial Computations Become Heavy

Suppose a machine produces $n = 200$ items per day, each defective independently with probability $p = 0.02$, and we want the probability of exactly 5 defectives. The exact binomial answer is

$$P(X = 5) = \binom{200}{5} (0.02)^5 (0.98)^{195}.$$

The binomial coefficient $\binom{200}{5}$ has 13 digits, and $(0.98)^{195}$ must be handled with logarithms or a calculator. For $n = 1000$ the situation is worse. Yet notice something: the mean $np = 200 \times 0.02 = 4$ is a small, comfortable number. Although 200 items are produced, only about 4 defectives are expected. The action happens entirely in the small values $k = 0, 1, 2, \dots, 12$; the probabilities for large k are negligible.

This suggests an idea: when n is large and p is small, perhaps the complicated binomial formula can be replaced by a simpler law that depends only on the mean $\lambda = np$. That simpler law is precisely the Poisson distribution.

PROPERTY Poisson approximation to the binomial

If $X \sim B(n, p)$ where n is **large** and p is **small**, so that $\lambda = np$ is moderate, then

$$B(n, p) \approx \text{Po}(\lambda), \quad \text{i.e.} \quad P(X = k) \approx e^{-\lambda} \frac{\lambda^k}{k!}, \quad \lambda = np.$$

The standard working conditions for the GCE Advanced Level examination are $n \geq 50$ and $p \leq 0.1$ (or $q = 1 - p \leq 0.1$ if we count failures instead of successes).

Why the approximation works (matching the means). A rigorous limit proof is beyond the syllabus, but the key idea is instructive and easy to grasp. Fix $\lambda = np$ and write $p = \lambda/n$. Then

$$\begin{aligned} P(X = k) &= \binom{n}{k} p^k (1-p)^{n-k} \\ &= \frac{n(n-1)\cdots(n-k+1)}{k!} \left(\frac{\lambda}{n}\right)^k \left(1 - \frac{\lambda}{n}\right)^{n-k} \\ &= \frac{\lambda^k}{k!} \cdot \underbrace{\frac{n(n-1)\cdots(n-k+1)}{n^k}}_{\rightarrow 1 \text{ as } n \rightarrow \infty} \cdot \underbrace{\left(1 - \frac{\lambda}{n}\right)^n}_{\rightarrow e^{-\lambda}} \cdot \underbrace{\left(1 - \frac{\lambda}{n}\right)^{-k}}_{\rightarrow 1}. \end{aligned}$$

As $n \rightarrow \infty$ with λ fixed, the three bracketed factors tend to 1, $e^{-\lambda}$ and 1 respectively, leaving exactly the Poisson formula $e^{-\lambda} \lambda^k / k!$. Notice also that the two distributions agree on the mean: $E(X) = np = \lambda$ in both cases, and the binomial variance $np(1-p) = \lambda(1-p) \approx \lambda$ when p is small, matching the Poisson variance. The approximation is therefore consistent in mean and in variance.

METHOD Approximation checklist

To approximate $X \sim B(n, p)$ by a Poisson distribution:

1. Check $n \geq 50$ (large number of trials).
2. Check $p \leq 0.1$ (success is rare). Equivalently, if failures are rare ($q = 1 - p \leq 0.1$), work with q and count failures.
3. Compute $\lambda = np$ and state $X \approx Y$ where $Y \sim \text{Po}(np)$.
4. Compute Poisson probabilities using $P(Y = k) = e^{-\lambda} \lambda^k / k!$, the recurrence formula $P(Y = k+1) = \frac{\lambda}{k+1} P(Y = k)$, or cumulative tables.

WARNING When NOT to use the Poisson approximation

A very common mistake is to approximate $B(n, p)$ by a Poisson law when p is close to 0.5. For example $B(100, 0.5)$ is not rare-event territory: its mean is 50 and its spread is large, and $\text{Po}(50)$ fits it poorly (the binomial is nearly symmetric, the Poisson is right-skewed).

For large n with moderate p , the appropriate approximation is the normal distribution (met later in the course), not the Poisson. Reserve the Poisson approximation for **rare** events: p small, $\lambda = np$ moderate.


8.2 Comparing Exact and Approximate Values

EXAMPLE B(100, 0.04) versus Po(4)

Let $X \sim B(100, 0.04)$: $n = 100 \geq 50$, $p = 0.04 \leq 0.1$, so the approximation with $\lambda = np = 4$ is legitimate. The table compares exact binomial probabilities (computed from the binomial formula, rounded to 4 decimal places) with the Poisson approximations $e^{-4}4^k/k!$.

k	$B(100, 0.04)$ exact	Po(4) approx.	Absolute error
0	0.0169	0.0183	0.0014
1	0.0703	0.0733	0.0030
2	0.1450	0.1465	0.0015
3	0.1973	0.1954	0.0019
4	0.1994	0.1954	0.0040
5	0.1595	0.1563	0.0032
6	0.1053	0.1042	0.0011
7	0.0589	0.0595	0.0006
8	0.0285	0.0298	0.0013
9	0.0122	0.0132	0.0010
10	0.0046	0.0053	0.0007

The agreement is excellent: every error is below 0.005, and cumulative probabilities agree to two decimal places. The figure below shows the two distributions superimposed.



Bars: exact binomial probabilities for $B(100, 0.04)$. Dots: Poisson approximation $Po(4)$. The two are almost indistinguishable.

WORKED EXAMPLE 17 Quality control in a soap factory

A factory in Bonabéri produces 400 tablets of soap per batch. Each tablet is defective with probability 0.005, independently of the others. Using a suitable approximation, find the probability that a batch contains (a) exactly 3 defectives; (b) at least one defective.

SOLUTION

Solution. Let X be the number of defectives, $X \sim B(400, 0.005)$. Since $n = 400 \geq 50$ and $p = 0.005 \leq 0.1$, we approximate by $Y \sim Po(\lambda)$ with

$$\lambda = np = 400 \times 0.005 = 2.$$

$$(a) P(X = 3) \approx P(Y = 3) = e^{-2} \frac{2^3}{3!} = 0.1353 \times \frac{8}{6} \approx 0.1804.$$

(b) Using the “at least one” trick:

$$P(X \geq 1) \approx P(Y \geq 1) = 1 - P(Y = 0) = 1 - e^{-2} \approx 1 - 0.1353 = 0.8647.$$

So a batch contains at least one defective about 86% of the time.

8.3 Applications of the Poisson Distribution

Because rare-event binomial situations are everywhere, the Poisson distribution is one of the most applied laws in statistics. Typical modelling situations (all with n large, p small, events independent, in a fixed interval of time, length, area or volume):

- **Rare diseases:** number of cases of a rare illness per year in a region of Cameroon, with $\lambda =$

average annual number of cases.

- **Machine breakdowns:** number of breakdowns of a generator per month in a factory; maintenance teams use $P(X \geq k)$ to plan spare parts.
- **Typing and printing errors:** number of misprints per page of a textbook; a proofreader expects λ errors per page and computes $P(X = 0) = e^{-\lambda}$, the probability of a clean page.
- **Customer arrivals:** number of customers joining the queue at an MTN service point per 10 minutes, or calls arriving at a call centre per minute.
- **Goals in football:** the number of goals scored by a team in a match is well modelled by a Poisson law with λ equal to the team's average goals per match.

DID YOU KNOW? A law born from data

The Poisson distribution was introduced by Siméon Denis Poisson in the 1830s, and became famous when Ladislaus Bortkiewicz showed in 1898 that it fitted the number of soldiers killed by horse kicks per year in Prussian army corps: many corps (large n), tiny probability of such a death (small p), and a small average λ . It remains the standard model for rare events in insurance, epidemiology and telecommunications.

8.4 Multi-Step Problems: Scaling, Cumulative Probabilities and “At Least One”

GCE questions often combine three ideas in one problem: the **scaling rule** (if $X \sim \text{Po}(\lambda)$ counts events in one unit interval, then the count in t units is $\text{Po}(\lambda t)$), **cumulative probabilities** $P(X \leq k)$ obtained by summing, and the **at-least-one event** $P(X \geq 1) = 1 - e^{-\lambda}$.

WORKED EXAMPLE 18 Calls at a call centre

Calls arrive at a customer-care centre in Yaoundé at an average rate of 3 calls per 10 minutes, independently, following a Poisson process.

1. Find the probability of exactly 2 calls in a 10-minute period.
2. Find the probability of at least one call in a 5-minute period.
3. Find the probability of at most 2 calls in a 20-minute period.

SOLUTION

Solution.

1. Here $\lambda = 3$: $P(X = 2) = e^{-3} \frac{3^2}{2!} = 0.04979 \times 4.5 \approx 0.2240$.
2. Scale the interval: in 5 minutes, $\lambda = 3 \times \frac{5}{10} = 1.5$. Then

$$P(X \geq 1) = 1 - e^{-1.5} \approx 1 - 0.2231 = 0.7769.$$

3. In 20 minutes, $\lambda = 3 \times 2 = 6$. Then

$$P(X \leq 2) = e^{-6} \left(1 + 6 + \frac{6^2}{2!} \right) = e^{-6}(1 + 6 + 18) = 25e^{-6} \approx 0.0620.$$

WARNING Scale λ before computing

The most frequent error in such questions is forgetting to adjust λ when the interval changes. If the rate is given per 10 minutes and the question concerns 5 minutes, you must halve λ first. Writing $1 - e^{-3}$ instead of $1 - e^{-1.5}$ loses all the marks for that part.

8.5 Using the Recurrence Formula for Efficient Computation

When calculating several consecutive Poisson probabilities, the recurrence formula avoids repeated evaluation of $e^{-\lambda}$ and factorials.

EXAMPLE Using the recurrence formula

For the call centre with $\lambda = 3$ (per 10 minutes), compute $P(X = 0)$, $P(X = 1)$, $P(X = 2)$, $P(X = 3)$ using the recurrence $P(X = k + 1) = \frac{\lambda}{k+1} P(X = k)$.

Solution.

$$P(X = 0) = e^{-3} \approx 0.04979.$$

$$P(X = 1) = \frac{3}{1} \times 0.04979 = 0.14936.$$

$$P(X = 2) = \frac{3}{2} \times 0.14936 = 0.22404.$$

$$P(X = 3) = \frac{3}{3} \times 0.22404 = 0.22404.$$

Check: $P(X = 3) = e^{-3} \frac{27}{6} = 0.04979 \times 4.5 = 0.22404$. The recurrence is faster and reduces rounding errors.

8.6 Estimating λ from Data and Checking the Fit

In practice λ is unknown and must be **estimated** from data. Since $E(X) = \lambda$ for a Poisson distribution, the natural estimate is the sample mean:

$$\hat{\lambda} = \bar{x} = \frac{\sum f_k k}{\sum f_k}.$$

A quick diagnostic for whether a Poisson model is reasonable exploits the signature property of the law: **mean $\approx$ variance**. Compute the sample mean $\bar{x}$ and the sample variance s^2 ; if they are

close, a Poisson fit is plausible. If the variance is much larger than the mean (“overdispersion”), the Poisson model should be rejected.

EXAMPLE Goals scored per match

Over 40 matches, a team scores the following numbers of goals: 0 goals in 14 matches, 1 goal in 14 matches, 2 goals in 8 matches, 3 goals in 3 matches, 4 goals in 1 match. Then

$$\hat{\lambda} = \bar{x} = \frac{0(14) + 1(14) + 2(8) + 3(3) + 4(1)}{40} = \frac{43}{40} = 1.075.$$

The sample variance is

$$s^2 = \frac{\sum f_k k^2}{\sum f_k} - \bar{x}^2 = \frac{0 + 14 + 32 + 27 + 16}{40} - (1.075)^2 = \frac{89}{40} - 1.155625 = 2.225 - 1.155625 = 1.069375 \approx 1.07,$$

very close to the mean, so $\text{Po}(1.075)$ is a sensible model. For instance the fitted probability of scoring no goal is $e^{-1.075} \approx 0.341$, predicting about $40 \times 0.341 \approx 13.6$ goalless matches, against 14 observed: an excellent fit.

8.7 Comparing the Binomial, Poisson and Normal Approximations

Lesson 158 of the syllabus requires comparing distributions. The table below summarises when to use each approximation for a binomial $B(n, p)$.

Situation	Approximation	Conditions	Parameter(s)
n large, p small (rare successes)	Poisson	$n \geq 50, p \leq 0.1$	$\lambda = np$
n large, $q = 1 - p$ small (rare failures)	Poisson (on failures)	$n \geq 50, q \leq 0.1$	$\lambda = nq$
n large, p moderate (not near 0 or 1)	Normal	$np > 5, nq > 5$	$\mu = np, \sigma^2 = npq$

WARNING Continuity correction for Normal, not for Poisson

When using the Normal approximation to the binomial, a **continuity correction** is mandatory (e.g. $P(X \leq k) \approx P(Z \leq \frac{k+0.5-\mu}{\sigma})$). The Poisson approximation is discrete-to-discrete: no continuity correction is needed or allowed. Simply compute $P(Y = k)$ or $P(Y \leq k)$ for $Y \sim \text{Po}(np)$.

8.8 Real-Life Problem Solving: A Comprehensive Example

WORKED EXAMPLE 19 Inspection of electrical cables

A factory produces electrical cables. Experience shows that 0.8% of the cables have a microscopic defect. A quality controller selects a random sample of 500 cables.

1. State the exact distribution of the number of defective cables in the sample. Justify why a Poisson approximation is appropriate and give its parameter.
2. Use the Poisson approximation to find the probability that the sample contains (i) no defective cables, (ii) exactly two defective cables, (iii) at most one defective cable.
3. The factory produces 10 000 cables per day. Estimate the number of days in a 300-day working year on which a sample of 500 cables would contain more than 3 defectives.

SOLUTION

Solution.

1. Let X be the number of defective cables in the sample. Exact distribution: $X \sim B(500, 0.008)$. Since $n = 500 \geq 50$ and $p = 0.008 \leq 0.1$, the Poisson approximation is valid with $\lambda = np = 500 \times 0.008 = 4$. So $X \approx Y \sim \text{Po}(4)$.
2. Using $Y \sim \text{Po}(4)$:
 - (i) $P(Y = 0) = e^{-4} \approx 0.0183$.
 - (ii) $P(Y = 2) = e^{-4} \frac{4^2}{2!} = 0.0183 \times 8 = 0.1465$.
 - (iii) $P(Y \leq 1) = P(Y = 0) + P(Y = 1) = e^{-4}(1 + 4) = 5e^{-4} \approx 5 \times 0.0183 = 0.0916$.
3. For a daily sample of 500, $P(\text{more than 3 defectives}) = P(Y > 3) = 1 - P(Y \leq 3)$. $P(Y \leq 3) = e^{-4} \left(1 + 4 + \frac{16}{2} + \frac{64}{6}\right) = e^{-4}(1 + 4 + 8 + 10.6667) = 23.6667e^{-4} \approx 0.4335$. So $P(Y > 3) \approx 1 - 0.4335 = 0.5665$. Expected number of days in 300 days = $300 \times 0.5665 \approx 170$ days.

METHOD GCE exam technique: “use a suitable approximation”

When a question says “use a suitable approximation”:

1. Name the approximating distribution explicitly: “ $X \sim B(n, p) \approx \text{Po}(\lambda)$ ”.
2. Justify the conditions: state $n \geq 50$ and $p \leq 0.1$ (or $q \leq 0.1$).
3. Give the parameter: $\lambda = np$, with its numerical value.
4. Compute with the Poisson formula, recurrence, or tables, and round sensibly (usually 4 decimal places).

Examiners award method marks for the statement of the approximating distribution and its parameter, even before any computation.

EXERCISE 8 Insurance claims

An insurance company receives claims from 600 policyholders; each policyholder makes a claim in a given month with probability 0.004, independently.

1. Justify the use of a Poisson approximation and state its parameter.
2. Find the probability that exactly 2 claims are made in a month.
3. Find the probability that at least 3 claims are made in a month.

SOLUTION Exercise 1

1. $n = 600 \geq 50$, $p = 0.004 \leq 0.1$: the Poisson approximation is valid with $\lambda = np = 600 \times 0.004 = 2.4$, so $X \approx Y \sim \text{Po}(2.4)$.
2. $P(Y = 2) = e^{-2.4} \frac{2.4^2}{2!} = 0.0907 \times 2.88 \approx 0.2613$.
3. $P(Y \geq 3) = 1 - P(Y \leq 2) = 1 - e^{-2.4} \left(1 + 2.4 + \frac{2.4^2}{2} \right) = 1 - e^{-2.4}(1 + 2.4 + 2.88) = 1 - 0.0907 \times 6.28 \approx 1 - 0.5697 = 0.4303$.

KEY POINTS Key points

- If $n \geq 50$ and $p \leq 0.1$, then $B(n, p) \approx \text{Po}(np)$; set $\lambda = np$ and say so explicitly.
- The approximation matches the mean exactly ($np = \lambda$) and the variance approximately ($np(1 - p) \approx \lambda$).
- Never use the Poisson approximation when p is near 0.5; that is normal-approximation territory.
- “At least one event”: $P(X \geq 1) = 1 - e^{-\lambda}$. Change of interval: scale λ in proportion to the interval length.
- Estimate λ by the sample mean, and check the Poisson fit by verifying mean $\approx$ variance.
- Use the recurrence formula $P(X = k + 1) = \frac{\lambda}{k+1} P(X = k)$ for fast, accurate computation of several probabilities.
- No continuity correction for Poisson approximation (discrete to discrete).

9 COMPARING DISTRIBUTIONS AND REAL-LIFE PROBLEM SOLVING

We have now studied three fundamental discrete probability distributions: the **binomial**, the **geometric** and the **Poisson**. Each arises from a different random experiment, yet they share a common language of parameters, expectations and variances. In this final section we bring them together, develop a systematic method for choosing the right model in a worded problem, and practise the kind of multi-part questions that appear in GCE Paper 2.

9.1 Summary Comparison of the Three Distributions

Before solving problems, it is essential to have a clear mental picture of the differences and similarities. The table below summarises the key features. Note that for the Geometric distribution, we adopt the convention X = number of trials up to and including the first success (support $x = 1, 2, \dots$), which is the standard for the GCE Advanced Level syllabus.


KEY POINTS

Summary Table: Binomial vs Geometric vs Poisson

Feature	Binomial $B(n, p)$	Geometric $Geo(p)$	Poisson $Po(\lambda)$
Random experiment	Fixed number n of independent Bernoulli trials.	Repeated independent Bernoulli trials until the first success.	Counting random events occurring in a fixed interval of time/space.
Parameters	$n \in \mathbb{N}^*$ (number of trials), $p \in (0, 1)$ (success probability).	$p \in (0, 1)$ (success probability).	$\lambda > 0$ (average rate per unit interval).
Support (possible values)	$x = 0, 1, 2, \dots, n$	$x = 1, 2, 3, \dots$	$x = 0, 1, 2, \dots$
Probability mass function	$P(X = x) = \binom{n}{x} p^x (1 - p)^{n-x}$	$P(X = x) = (1 - p)^{x-1} p$	$P(X = x) = \frac{e^{-\lambda} \lambda^x}{x!}$
Expectation	$E(X) = np$	$E(X) = \frac{1}{p}$	$E(X) = \lambda$
Variance	$Var(X) = np(1 - p)$	$Var(X) = \frac{1 - p}{p^2}$	$Var(X) = \lambda$
Key condition	Fixed n , constant p , independence.	No fixed n , constant p , independence, memoryless property.	Events occur singly, independently, at constant average rate.
Shape	Symmetric if $p = 0.5$, skewed otherwise; bounded.	Strictly decreasing; unbounded right tail.	Skewed for small λ , approx. symmetric for large λ ; unbounded.

9.2 Decision Flowchart: Which Distribution to Use?

When faced with a worded problem, follow the logical steps in the flowchart below. It guides you from the description of the random experiment to the appropriate distribution.



9.3 Comparing Binomial and Poisson Distributions

The binomial and Poisson distributions are closely linked. The Poisson distribution arises as a limiting case of the binomial when $n \rightarrow \infty$, $p \rightarrow 0$ while $np = \lambda$ remains constant. This is the basis of the **Poisson approximation to the binomial**, which we studied in the previous section.

PROPERTY When to Use Poisson Approximation to Binomial

The approximation $B(n, p) \approx Po(\lambda)$ with $\lambda = np$ is appropriate when:

- n is large (typically $n \geq 50$),
- p is small (typically $p \leq 0.1$),
- $np = \lambda$ is moderate (typically $\lambda \leq 5$ or $\lambda \leq 10$).

The approximation works because the binomial's fixed upper bound n becomes irrelevant when p is tiny; the Poisson's infinite tail matches the binomial's effective range.

WARNING Common Mistake: Applying Poisson When There Is a Fixed Maximum

A frequent error is to model a situation with a fixed maximum number of occurrences using a Poisson distribution. For example, the number of defective items in a batch of 100 can-

not exceed 100. If $n = 100$ and $p = 0.02$, then $np = 2$. Although the Poisson approximation $Po(2)$ is numerically excellent, the correct model is $B(100, 0.02)$ because the experiment has a fixed number of trials. In GCE exams, you must state the exact distribution first, then mention the approximation if appropriate.

9.4 Modelling Strategy for Real-Life Problems

GCE Paper 2 questions often present a real-life scenario and ask you to "state the distribution", "calculate the probability that...", and "interpret your result". Use this four-step method.

METHOD 4-Step Modelling Strategy

1. Identify the random variable. Define X clearly in words (e.g., " X = number of faulty bulbs in a box of 20").
2. Check the conditions. Match the scenario to the assumptions of a distribution (fixed n ? waiting for first success? events in an interval?).
3. State the distribution and its parameters. Write, for example, " $X \sim B(20, 0.05)$ " or " $X \sim Po(3.2)$ ".
4. Compute and interpret. Perform the calculation (using formula, tables, or approximation) and finish with a sentence in context: "The probability that a box contains at most 1 faulty bulb is 0.736, so about 74% of boxes meet the quality standard."

9.5 Mixed Problems: Combining Binomial and Poisson

Many exam questions combine two distributions in a single problem. A classic pattern: a Poisson process generates a random number of "items", and each item is then subjected to a binomial trial (e.g., each item is inspected and may be defective).

WORKED EXAMPLE 20 Mixed Binomial-Poisson Problem

Statement: In a factory, the number of machines that break down in a week follows a Poisson distribution with mean 2.5. Each broken machine is repaired independently; the probability that a repair is completed within the same week is 0.6. Find the probability that in a given week, exactly 2 machines break down and both are repaired within the week.

SOLUTION

Detailed solution:

1. Define variables. Let N = number of breakdowns in a week. $N \sim \text{Po}(2.5)$. Given $N = n$, let R = number of those n machines repaired within the week. Then $R \mid N = n \sim B(n, 0.6)$.

2. We want $P(N = 2 \text{ and } R = 2)$. By the multiplication rule:

$$P(N = 2, R = 2) = P(N = 2) \times P(R = 2 \mid N = 2).$$

3. Compute each factor.

$$P(N = 2) = \frac{e^{-2.5}(2.5)^2}{2!} = \frac{6.25}{2}e^{-2.5} = 3.125e^{-2.5}.$$

Using $e^{-2.5} \approx 0.082085$:

$$P(N = 2) \approx 3.125 \times 0.082085 = 0.2565.$$

$$P(R = 2 \mid N = 2) = \binom{2}{2}(0.6)^2(0.4)^0 = 0.36.$$

4. Multiply.

$$P(N = 2, R = 2) \approx 0.2565 \times 0.36 = 0.09234.$$

5. Interpret. The probability that exactly two machines break down and both are repaired within the same week is approximately 0.0923 (or 9.23%).

9.6 Interpreting Results in Context


A numerical answer without interpretation is incomplete in GCE Pure Mathematics with Statistics. Always relate the probability or expectation back to the real situation.

KEY POINTS What "Interpret" Means in GCE Questions

- If you compute $P(X \leq k) = 0.85$, say: "There is an 85% chance that at most k events occur."
- If you find $E(X) = 4.2$, say: "On average, we expect 4.2 events per interval (e.g., per week)."
- If you compare observed and expected frequencies (goodness-of-fit), say: "The model fits the data well / does not fit the data well because..."
- If you use an approximation, say: "Using the Poisson approximation to the binomial, the probability is approximately 0.123."

9.7 Visual Comparison of Shapes

To reinforce intuition, the figure below plots the probability mass functions of $B(10, 0.5)$, $Geo(0.3)$ (counting trials until first success) and $Po(3)$ on aligned axes. Notice the bounded symmetric binomial, the strictly decreasing geometric, and the skewed Poisson with a mode at 3.



Comparison of probability mass functions: $B(10, 0.5)$ (blue dots), $Geo(0.3)$ (orange squares), $Po(3)$ (green triangles).

9.8 Chapter Synthesis and GCE Exam Technique

This chapter has equipped you with three powerful discrete distributions. Mastery comes from recognising the structure of the random experiment, not just memorising formulas. Here is a checklist for exam success.

METHOD GCE Exam Technique Checklist for Discrete Distributions

1. Read twice. Underline key phrases: "fixed number of trials", "until the first success", "average rate per hour", "randomly scattered".
2. Define X explicitly. " $X = \text{number of } \dots$ " — this earns a mark and focuses your thinking.
3. State the distribution with parameters. Write $X \sim B(n, p)$, $X \sim Geo(p)$ or $X \sim Po(\lambda)$. If using an approximation, write: " $X \sim B(100, 0.02) \approx Po(2)$ ".
4. Quote the formula. For binomial: $P(X = x) = \binom{n}{x} p^x (1 - p)^{n-x}$. For Poisson: $P(X = x) = e^{-\lambda} \lambda^x / x!$. For geometric: $P(X = x) = (1 - p)^{x-1} p$. This shows you know the theory.
5. Show substitution. Write the numbers into the formula before using a calculator or tables.

6. Use tables correctly. If using cumulative binomial tables, state clearly: "From tables with $n = 20, p = 0.3, P(X \leq 5) = 0.4164$." For Poisson tables, similarly.
7. Round appropriately. Probabilities to 3 or 4 significant figures (or as requested). Expectations/variances to 1 or 2 decimal places.
8. Conclude in context. Always end with a sentence interpreting the result. This is often a separate mark in the mark scheme.
9. Check variance vs mean. For Poisson, variance = mean. For binomial, variance < mean. For geometric, variance > mean. This can be a quick sanity check.

DID YOU KNOW?**Historical Note: The Poisson Distribution in Cameroon**

The Poisson distribution is named after Siméon Denis Poisson (1781–1840), a French mathematician who introduced it in 1837. In Cameroon, it is widely used in queueing theory for modelling customer arrivals at banks (e.g., Ecobank, Afriland First Bank) and in telecommunications for call arrivals at MTN or Orange switching centres. It also models the distribution of rare diseases in epidemiology (e.g., cases of cholera in a health district per month) and the number of defects in manufactured goods (e.g., bottles at the Cameroon Breweries plant in Douala).

WORKED EXAMPLE 21**Real-Life Problem: Choosing and Justifying the Model**

Statement: A call centre receives an average of 12 calls per hour. The centre has 5 operators. Each call lasts an average of 5 minutes. Assume calls arrive randomly and independently.

1. State the distribution of the number of calls received in a 30-minute period.
2. Find the probability that more than 8 calls are received in a 30-minute period.
3. The manager claims that the probability of all 5 operators being busy at a given instant is negligible. Comment on this claim using a suitable model.

SOLUTION

Detailed solution:

1. Identify: Calls arrive randomly, independently, at a constant average rate. This is a Poisson process. Rate per hour $\lambda = 12$. For 30 minutes (0.5 hour), $\lambda_{30} = 12 \times 0.5 = 6$. Let X = number of calls in 30 minutes. Then $X \sim Po(6)$.
2. $P(X > 8) = 1 - P(X \leq 8)$. Using Poisson cumulative tables with $\lambda = 6$: $P(X \leq 8) = 0.8472$ (from tables). So $P(X > 8) = 1 - 0.8472 = 0.1528$. Interpretation: There is about a 15.3%

chance that more than 8 calls arrive in a half-hour period.

3. The number of operators busy at an instant depends on the number of calls that have arrived in the last 5 minutes and are still ongoing. We approximate the number of simultaneous calls by the number of arrivals in a 5-minute interval. Average calls in 5 minutes = $12 \times (5/60) = 1$. Let Y = number of calls in a 5-minute interval. Then $Y \sim \text{Po}(1)$. $P(\text{all 5 busy}) \approx P(Y \geq 5) = 1 - P(Y \leq 4)$. From Poisson tables with $\lambda = 1$: $P(Y \leq 4) = 0.9963$. Thus $P(Y \geq 5) = 1 - 0.9963 = 0.0037$. This is very small (0.37%), so the manager's claim is reasonable under the Poisson assumption. However, real call centres often have bursty traffic (not perfectly Poisson), so the actual probability may be higher.

EXERCISE 9 Practice: Mixed and Comparison Problems

1. A factory produces bolts. The probability a bolt is defective is 0.02. Bolts are packed in boxes of 100.
 - (a) State the exact distribution of the number of defective bolts in a box.
 - (b) Use a Poisson approximation to find the probability that a box contains at most 3 defective bolts.
 - (c) A quality inspector checks 10 boxes. Find the probability that exactly 2 boxes contain more than 3 defectives. (Hint: two-stage binomial–Poisson).
2. In a large field, daisies are randomly distributed with an average density of 4 per square metre. A botanist places a circular hoop of radius 0.5 m at a random location.
 - (a) State the distribution of the number of daisies inside the hoop.
 - (b) Find the probability that the hoop contains exactly 2 daisies.
 - (c) The botanist repeats this in 20 independent locations. Find the probability that at least 15 hoops contain exactly 2 daisies.
3. A student takes a multiple-choice test with 20 questions, each with 4 options. She guesses all answers.
 - (a) State the distribution of the number of correct answers.
 - (b) Find the probability she passes (score ≥ 10).
 - (c) If she repeats the test until she passes, what is the distribution of the number of attempts needed? Find the expected number of attempts.

SOLUTION Exercise 1

- a) $X \sim B(100, 0.02)$.
- b) $\lambda = np = 2$. $X \approx Po(2)$. $P(X \leq 3) = e^{-2} \left(1 + 2 + \frac{2^2}{2!} + \frac{2^3}{3!}\right) = e^{-2} \left(1 + 2 + 2 + \frac{8}{6}\right) = e^{-2} (5 + 1.3333) = 6.3333 \times 0.13534 = 0.8571$. (Tables give 0.8571.)
- c) Let $p = P(\text{box has } > 3 \text{ defectives}) = 1 - 0.8571 = 0.1429$. Let $Y = \text{number of such boxes out of 10}$. $Y \sim B(10, 0.1429)$. $P(Y = 2) = \binom{10}{2} (0.1429)^2 (0.8571)^8 = 45 \times 0.02042 \times 0.291 \approx 0.268$.

SOLUTION Exercise 2

- a) Area of hoop $= \pi r^2 = \pi \times 0.25 \approx 0.7854 \text{ m}^2$. Mean daisies $\lambda = 4 \times 0.7854 = 3.1416$. $X \sim Po(3.1416)$.
- b) $P(X = 2) = e^{-3.1416} \frac{3.1416^2}{2!} \approx 0.216$.
- c) $p = 0.216$. $Y \sim B(20, 0.216)$. $P(Y \geq 15) = 1 - P(Y \leq 14)$. This is extremely small (use tables or normal approximation). $P(Y \geq 15) \approx 1.2 \times 10^{-7}$.

SOLUTION Exercise 3

- a) $X \sim B(20, 0.25)$.
- b) $P(X \geq 10) = 1 - P(X \leq 9)$. Using binomial tables with $n = 20, p = 0.25$: $P(X \leq 9) = 0.9861$, so $P(X \geq 10) = 0.0139$.
- c) Let $N = \text{number of attempts until first pass}$. $N \sim Geo(p)$ with $p = 0.0139$. $E(N) = 1/p = 1/0.0139 \approx 72 \text{ attempts}$.

WARNING Final Warning: "State the Distribution" Means Write the Symbol

In GCE exams, "State the distribution of X " requires the answer " $X \sim B(n, p)$ " or " $X \sim Po(\lambda)$ " with the correct numerical parameters. Writing only "binomial" or "Poisson" without parameters loses the mark. Always include the parameters.

With this section, we complete the chapter on Special Discrete Probability Distributions. You now possess the toolkit to model a wide variety of random phenomena — from quality control in a Douala factory to call arrivals at a Yaoundé call centre, from the number of attempts to pass a driving test in Buea to the distribution of malaria cases in a health district. The key is not

just computation, but modelling: translating a real situation into a mathematical framework, solving it, and translating the answer back. That is the essence of statistics at Advanced Level. Good luck in your revision and in the GCE examination!

10 INTEGRATION ACTIVITY

11 COMPARING DISTRIBUTIONS AND REAL-LIFE PROBLEM SOLVING

11.1 Integration Activity: The Biyem–Assi Health Centre Case Study

EXPERIMENT Context and Objectives

This integration activity consolidates your understanding of the **binomial** and **Poisson** distributions by applying them to a realistic health–centre scenario in Cameroon. You will:

- Identify the appropriate probability model for a given random phenomenon (binomial for a fixed number of independent trials with constant success probability; Poisson for events occurring randomly in time/space at a known average rate).
- Compute and interpret expectation, variance, and probabilities (both exact and using cumulative tables).
- Apply the **Poisson approximation to the binomial distribution** and critically assess its validity using the standard rule of thumb.
- Use the **scaling property** of the Poisson process to change the time interval.
- Determine the **mode** of a Poisson distribution via its recurrence formula.
- Communicate findings clearly in a concise report suitable for a non-specialist (the head nurse).

EXPERIMENT Situation: Biyem–Assi District Health Centre

The Biyem–Assi District Health Centre in the Centre Region of Cameroon serves a semi-urban population. The centre’s laboratory and reception desk generate two distinct streams of data that the head nurse wishes to understand better for staffing and resource planning.

Part A – Malaria Rapid Diagnostic Tests (RDTs). Historical records show that, on average, 6% of patients tested for malaria are positive. Each morning, the laboratory tests exactly 25 patients (a fixed number). The result for each patient (positive / negative) is assumed independent of the others, and the probability of a positive result remains constant at 0.06 throughout the morning session.

Part B – Incoming Telephone Calls at the Reception Desk. The reception desk receives

telephone calls randomly throughout the day. Over a long observation period, the average rate is 4 calls per 15-minute interval. Calls are independent of each other, and the rate is constant during working hours.

The head nurse has asked your group to produce a one-page statistical brief answering the following operational questions:

1. For the morning malaria testing session: what is the expected number of positive cases? What is the variability? What is the probability that at most 2 patients test positive? (Use the cumulative binomial probability table.)
2. The laboratory technician suggests using a Poisson approximation to simplify calculations. Compute the Poisson approximation for the probability in (1) and comment on whether the approximation is justified.
3. For the reception desk: what is the probability that no call arrives during a given 15-minute period?
4. What is the probability that more than 6 calls arrive in a 30-minute period? (Hint: scale the rate.)
5. What is the most likely number of calls per hour (i.e., the mode of the hourly distribution)? Use the recurrence formula for the Poisson distribution to justify your answer.


EXPERIMENT Tasks for Group Work

Work in groups of 3–4. Each group will submit a single report (maximum one A4 page) containing:

1. Model Justification: For each part (A and B), state the name of the distribution used, list the assumptions that make it appropriate, and define the parameter(s) clearly.
2. Calculations for Part A:
 - (a) Write the probability mass function of the binomial distribution for the number X of positive tests.
 - (b) Compute $E(X)$ and $Var(X)$.
 - (c) Using the cumulative binomial table (provided in the examination booklet), find $P(X \leq 2)$.
 - (d) Compute the Poisson approximation $\tilde{P}(X \leq 2)$ with $\lambda = np$. Show the steps.
 - (e) Compare the two probabilities and state whether the approximation is acceptable (use the rule of thumb: $n \geq 50$, $p \leq 0.1$, $np \leq 5$).
3. Calculations for Part B:
 - (a) Write the probability mass function of the Poisson distribution for the number Y of

calls in a 15-minute interval.

- (b) Find $P(Y = 0)$.
 - (c) For a 30-minute interval, define the new random variable and its parameter. Compute $P(\text{more than 6 calls})$.
 - (d) For a 60-minute (1-hour) interval, define the random variable Z and its parameter λ_Z . Use the recurrence formula $P(Z = k) = \frac{\lambda_Z}{k} P(Z = k - 1)$ to determine the mode (the value of k that maximises $P(Z = k)$).
4. Interpretation and Recommendations: In plain language, explain what the probabilities mean for the head nurse. Suggest one practical action for the laboratory and one for the reception desk based on your findings.



Comparison of the exact Binomial $B(25, 0.06)$ and the Poisson $Po(1.5)$ approximation for the number of positive malaria tests. The approximation is visually excellent.

WORKED EXAMPLE 22 Model Solution and Examiner's Commentary

1. Model Justification

- Part A (Malaria Tests): The number of positive tests X follows a binomial distribution $B(n = 25, p = 0.06)$. Assumptions verified:
 - Fixed number of trials: $n = 25$ patients tested each morning.
 - Two outcomes per trial: Positive (success) / Negative (failure).
 - Constant success probability: $p = 0.06$ (historical average).
 - Independence: The result for one patient does not affect another.
- Part B (Phone Calls): The number of calls in a given time interval follows a Poisson dis-

tribution. For a 15-minute interval, $Y \sim Po(\lambda = 4)$. Assumptions verified:

- Events occur randomly and independently in time.
- Constant average rate: 4 calls per 15 minutes.
- Two events cannot occur at exactly the same instant (negligible probability of simultaneous calls).

2. Calculations for Part A

1. Probability Mass Function (PMF):

$$P(X = k) = \binom{25}{k} (0.06)^k (0.94)^{25-k}, \quad k = 0, 1, \dots, 25.$$

2. Expectation and Variance:

$$\begin{aligned} E(X) &= np = 25 \times 0.06 = 1.5, \\ \text{Var}(X) &= np(1-p) = 25 \times 0.06 \times 0.94 = 1.41. \end{aligned}$$

The standard deviation is $\sigma_X = \sqrt{1.41} \approx 1.19$.

3. Cumulative Probability $P(X \leq 2)$ using Tables: From the cumulative binomial table for $n = 25$, $p = 0.06$ (or $p = 0.05$ and $p = 0.10$ with interpolation if necessary; here we assume the exact table for $p = 0.06$ is available):

$$P(X \leq 2) = 0.8106.$$

(Verification by direct calculation: $P(X = 0) = 0.94^{25} \approx 0.2129$, $P(X = 1) = 25 \times 0.06 \times 0.94^{24} \approx 0.3389$, $P(X = 2) = \binom{25}{2} 0.06^2 0.94^{23} \approx 0.2588$, Sum ≈ 0.8106 .)

4. Poisson Approximation: Take $\lambda = np = 1.5$. Then $X \approx \tilde{X} \sim Po(1.5)$.

$$\tilde{P}(X \leq 2) = e^{-1.5} \left(1 + 1.5 + \frac{1.5^2}{2!} \right) = e^{-1.5} (1 + 1.5 + 1.125) = e^{-1.5} \times 3.625.$$

Using $e^{-1.5} \approx 0.223130$, we get

$$\tilde{P}(X \leq 2) \approx 0.223130 \times 3.625 = 0.8088.$$

5. Comparison and Validity Assessment:

Measure	Exact Binomial	Poisson Approx.
$P(X \leq 2)$	0.8106	0.8088
Absolute Error	0.0018	
Relative Error	0.22%	

Rule of Thumb Check: $n = 25$ (not ≥ 50), $p = 0.06 \leq 0.1$, $np = 1.5 \leq 5$. Conclusion: Although $n < 50$, the approximation is excellent (error $< 0.3\%$) because p is small and np is moderate. It is acceptable for practical operational planning.

3. Calculations for Part B

1. PMF for 15-minute interval: $Y \sim Po(4)$.

$$P(Y = k) = \frac{e^{-4}4^k}{k!}, \quad k = 0, 1, 2, \dots$$

2. Probability of No Call in 15 Minutes:

$$P(Y = 0) = e^{-4} \approx \mathbf{0.0183} \quad (1.83\%).$$

3. 30-Minute Interval – Scaling the Rate: The Poisson rate scales linearly with the length of the interval. For 30 minutes (2 units of 15 minutes), $\lambda_{30} = 4 \times 2 = 8$. Let $W \sim Po(8)$ be the number of calls in 30 minutes.

$$P(W > 6) = 1 - P(W \leq 6).$$

Using the cumulative Poisson table for $\lambda = 8$: $P(W \leq 6) = 0.3134$. Hence

$$P(W > 6) = 1 - 0.3134 = \mathbf{0.6866} \quad (68.66\%).$$

4. 1-Hour (60-Minute) Interval and Mode: For 60 minutes (4 units of 15 minutes), $\lambda_{60} = 4 \times 4 = 16$. Let $Z \sim Po(16)$. The **recurrence formula** for Poisson probabilities is:

$$P(Z = k) = \frac{\lambda}{k} P(Z = k - 1), \quad k \geq 1.$$

Probabilities increase while $\frac{\lambda}{k} > 1 \iff k < \lambda$, and decrease when $k > \lambda$. Since $\lambda = 16$ is an integer, the maximum probability occurs at two adjacent values:

$$k = \lambda - 1 = 15 \quad \text{and} \quad k = \lambda = 16.$$

Both give the same probability. Thus the distribution has two modes: 15 and 16 calls per hour.

4. Interpretation and Recommendations for the Head Nurse

- Laboratory (Malaria Testing): On average, 1.5 patients test positive each morning. However, there is a $1 - 0.8106 = 0.1894 \approx 19\%$ chance that more than 2 patients are positive. The Poisson approximation (19.12%) confirms this risk.

WARNING Operational Implication

Preparing only 2 treatment kits each morning would leave the centre unprepared roughly 1 morning in 5.

Recommendation: Prepare at least 3 treatment kits every morning to cover the vast majority ($> 81\%$) of sessions. Keep a small buffer stock for the remaining 19% of days.

- **Reception Desk (Phone Calls):** The probability of a completely quiet 15-minute period is very low (1.8%). In any 30-minute window, there is a 69% chance of receiving more than 6 calls. The hourly mode of 15–16 calls indicates a consistently high and predictable workload. Recommendation: Schedule a second receptionist during peak hours (e.g., 8 am–12 pm) to manage the high call volume and keep patient waiting times acceptable. Consider an automated call-back system for non-urgent queries.

KEY POINTS Key Examination Takeaways for PMS12

- **Binomial $B(n, p)$:** Fixed n , independent trials, constant p . $E = np$, $Var = np(1 - p)$. Use tables for cumulative probabilities.
- **Poisson $Po(\lambda)$:** Random events in time/space, constant rate, independence. $E = Var = \lambda$. PMF: $P(X = k) = e^{-\lambda} \lambda^k / k!$.
- **Poisson Approximation to Binomial:** Valid when n is large (≥ 50), p is small (≤ 0.1), np is moderate (≤ 5). Use $\lambda = np$. Always check conditions and comment on accuracy.
- **Scaling Poisson Rate:** If rate is λ per unit time, then for t units the rate is $t\lambda$. The random variable for the new interval is $Po(t\lambda)$.
- **Mode of $Po(\lambda)$:** Use recurrence $P(k) = \frac{\lambda}{k} P(k-1)$. If λ not integer, mode = $\lfloor \lambda \rfloor$. If λ integer, modes are λ and $\lambda - 1$.
- **Contextual Justification:** Always explicitly link the assumptions of the distribution to the wording of the problem (fixed n vs. rate in time/space).
- **Communication:** Translate probabilities into plain language (e.g., “19% chance of shortage”) and propose concrete actions.

METHOD Choosing Between Binomial and Poisson Models

1. Identify the random variable: Is it a count of successes in a fixed number of trials? → Binomial. Is it a count of events occurring randomly in a continuous interval (time, area, length)? → Poisson.
2. Check for a fixed n : If the problem states “out of 25 patients” or “in a sample of 100”, n is fixed → Binomial. If it states “average 4 calls per 15 minutes” with no upper limit, n is not fixed → Poisson.
3. Check for constant p vs. constant rate λ : Binomial needs constant success probability per trial. Poisson needs constant average rate per unit interval.

4. Check independence: Required for both.
5. Approximation? If Binomial conditions hold but n is large and p small, Poisson approximation may simplify calculations. State the rule of thumb and verify.

WARNING Common Pitfalls in Examinations

- Confusing parameters: Writing $Po(np)$ for the approximation but forgetting to define $\lambda = np$ explicitly.
- Forgetting to scale λ : Using $\lambda = 4$ for a 30-minute or 1-hour interval without multiplying by the number of 15-minute units.
- Mode of Poisson: Stating the mode is λ (e.g., 16) without checking if λ is integer. If λ is integer, you must give both λ and $\lambda - 1$.
- Table usage: Writing $P(X < 2)$ instead of $P(X \leq 2)$ when reading cumulative tables. Remember tables give $P(X \leq x)$.
- Continuity correction: Do not apply continuity correction for Poisson approximation to Binomial (that is for Normal approximation).
- Units in final answer: Always include units (e.g., “1.5 patients”, “16 calls per hour”).

DID YOU KNOW? Historical Note: The Poisson Distribution in Cameroon

The Poisson distribution is named after Siméon Denis Poisson (1781–1840), a French mathematician. In Cameroon, it is widely used in queueing theory to model customer arrivals at banks (e.g., BICEC, Afriland First Bank), call centres (MTN, Orange), and health facilities like the one in this activity. The Erlang formula, derived from the Poisson process, is the mathematical backbone of modern telecommunications network dimensioning used by Camtel and mobile operators to determine the number of lines needed to keep call blocking probability low.

12 METHODS AND WORKED EXAMPLES

12.1 Methods and Worked Examples

METHOD Recognising and Computing Binomial Probabilities

1. Identify the binomial setting: a fixed number n of independent trials, each with two outcomes (Success/Failure), and constant success probability p .
2. Define the random variable: $X \sim B(n, p)$ where X = number of successes in the n trials.
3. Write the probability mass function: $\mathbb{P}(X = k) = \binom{n}{k} p^k (1 - p)^{n-k}$ for $k = 0, 1, \dots, n$.
4. Compute: substitute n, p and the required k ; evaluate $\binom{n}{k}$ and the powers.
5. Check: probabilities must sum to 1; each $\mathbb{P}(X = k) \in [0, 1]$.

WORKED EXAMPLE 23 Binomial Probability: Defective Items in a Batch

A factory produces light bulbs. The probability that a bulb is defective is 0.03. A quality control officer selects a random sample of 20 bulbs.

1. Find the probability that exactly 2 bulbs are defective.
2. Find the probability that at most 1 bulb is defective.
3. Find the probability that at least 3 bulbs are defective.

SOLUTION

Solution.

Let X be the number of defective bulbs in the sample of 20. The trials are independent, $n = 20$ is fixed and $p = 0.03$ is constant, so $X \sim B(20, 0.03)$.

1.

$$\mathbb{P}(X = 2) = \binom{20}{2} (0.03)^2 (0.97)^{18} = 190 \times 0.0009 \times (0.97)^{18}.$$

Since $(0.97)^{18} \approx 0.5780$,

$$\mathbb{P}(X = 2) \approx 190 \times 0.0009 \times 0.5780 \approx 0.0988.$$

2. $\mathbb{P}(X \leq 1) = \mathbb{P}(X = 0) + \mathbb{P}(X = 1)$.

$$\mathbb{P}(X = 0) = (0.97)^{20} \approx 0.5438,$$

$$\mathbb{P}(X = 1) = \binom{20}{1} (0.03)(0.97)^{19} = 20 \times 0.03 \times 0.5616 \approx 0.3370.$$

Hence $\mathbb{P}(X \leq 1) \approx 0.5438 + 0.3370 = 0.8808$.

3. Using the complement,

$$\mathbb{P}(X \geq 3) = 1 - \mathbb{P}(X \leq 2) = 1 - [\mathbb{P}(X \leq 1) + \mathbb{P}(X = 2)].$$

$\mathbb{P}(X \leq 2) \approx 0.8808 + 0.0988 = 0.9796$, so

$$\mathbb{P}(X \geq 3) \approx 1 - 0.9796 = 0.0204.$$

WARNING “At least”, “at most”, “more than”

For a discrete variable, $\mathbb{P}(X \geq k) = 1 - \mathbb{P}(X \leq k-1)$ and $\mathbb{P}(X > k) = 1 - \mathbb{P}(X \leq k)$. A very common GCE error is to subtract $\mathbb{P}(X \leq k)$ when “at least k ” is required, thus losing the term $\mathbb{P}(X = k)$.

METHOD Expectation, Variance and Recurrence Formula for the Binomial Distribution

1. Expectation: $\mathbb{E}(X) = np$.
2. Variance: $\text{Var}(X) = np(1-p) = npq$ where $q = 1-p$.
3. Standard deviation: $\sigma = \sqrt{npq}$.
4. Recurrence formula: $\mathbb{P}(X = k+1) = \frac{n-k}{k+1} \cdot \frac{p}{q} \cdot \mathbb{P}(X = k)$ for $k = 0, 1, \dots, n-1$. It is obtained by dividing the expression for $\mathbb{P}(X = k+1)$ by that for $\mathbb{P}(X = k)$ and simplifying the binomial coefficients.
5. Mode: the value(s) of k maximising $\mathbb{P}(X = k)$. If $(n+1)p$ is an integer, there are two modes, $(n+1)p$ and $(n+1)p - 1$; otherwise the mode is $[(n+1)p]$.

WORKED EXAMPLE 24 Binomial Expectation, Variance, Recurrence and Mode

The random variable $X \sim B(12, 0.4)$.

1. Find $\mathbb{E}(X)$ and $\text{Var}(X)$.
2. Given that $\mathbb{P}(X = 3) = 0.1419$ (to 4 d.p.), use the recurrence formula to find $\mathbb{P}(X = 4)$ and $\mathbb{P}(X = 5)$.
3. Determine the mode of the distribution.

SOLUTION

Solution.

$$1. \mathbb{E}(X) = np = 12 \times 0.4 = 4.8.$$

$$\text{Var}(X) = npq = 12 \times 0.4 \times 0.6 = 2.88, \quad \sigma = \sqrt{2.88} \approx 1.697.$$

$$2. \text{ Here } \frac{p}{q} = \frac{0.4}{0.6} = \frac{2}{3}, \text{ so the recurrence formula gives}$$

$$\mathbb{P}(X = k + 1) = \frac{12 - k}{k + 1} \times \frac{2}{3} \times \mathbb{P}(X = k).$$

For $k = 3$:

$$\mathbb{P}(X = 4) = \frac{9}{4} \times \frac{2}{3} \times 0.1419 = 1.5 \times 0.1419 = 0.2129.$$

For $k = 4$:

$$\mathbb{P}(X = 5) = \frac{8}{5} \times \frac{2}{3} \times 0.2129 = \frac{16}{15} \times 0.2129 \approx 0.2271.$$

$$3. (n + 1)p = 13 \times 0.4 = 5.2, \text{ which is not an integer, so the mode is } \lfloor 5.2 \rfloor = 5.$$

Check: $\mathbb{P}(X = 6) = \frac{7}{6} \times \frac{2}{3} \times 0.2271 \approx 0.1766 < \mathbb{P}(X = 5)$, and $\mathbb{P}(X = 4) = 0.2129 < \mathbb{P}(X = 5)$.
Since the probabilities increase up to the mode and decrease after it, $k = 5$ is indeed the mode.

METHOD Using Cumulative Binomial Tables and Complementary Probabilities

1. Identify the parameters n and p . If $p > 0.5$, use the symmetry: $X \sim B(n, p) \Rightarrow Y = n - X \sim B(n, 1 - p)$, and tables for $1 - p$.
2. Read cumulative probabilities: tables give $\mathbb{P}(X \leq x)$.
3. Convert the required probability:
 - $\mathbb{P}(X \leq k)$: read directly from the table;
 - $\mathbb{P}(X < k) = \mathbb{P}(X \leq k - 1)$;
 - $\mathbb{P}(X \geq k) = 1 - \mathbb{P}(X \leq k - 1)$;
 - $\mathbb{P}(X > k) = 1 - \mathbb{P}(X \leq k)$;
 - $\mathbb{P}(a \leq X \leq b) = \mathbb{P}(X \leq b) - \mathbb{P}(X \leq a - 1)$.
4. If $p > 0.5$: with $Y = n - X \sim B(n, 1 - p)$, use $\mathbb{P}(X = k) = \mathbb{P}(Y = n - k)$ and $\mathbb{P}(X \geq k) = \mathbb{P}(Y \leq n - k)$.

WORKED EXAMPLE 25 Cumulative Binomial Tables with $p > 0.5$

The probability that a student passes a test is 0.75. In a group of 15 students, find the probability that:

1. exactly 10 pass,
2. at least 12 pass,
3. between 8 and 11 inclusive pass.

Use cumulative tables for $B(15, 0.25)$, which give $\mathbb{P}(Y \leq 3) = 0.4613$, $\mathbb{P}(Y \leq 4) = 0.6865$, $\mathbb{P}(Y \leq 5) = 0.8516$ and $\mathbb{P}(Y \leq 7) = 0.9827$.

SOLUTION

Solution.

Let X = number of passes, so $X \sim B(15, 0.75)$. Since $p = 0.75 > 0.5$, set $Y = 15 - X \sim B(15, 0.25)$ (the number of failures).

1. $\mathbb{P}(X = 10) = \mathbb{P}(Y = 5) = \mathbb{P}(Y \leq 5) - \mathbb{P}(Y \leq 4) = 0.8516 - 0.6865 = 0.1651$.
2. $\mathbb{P}(X \geq 12) = \mathbb{P}(Y \leq 3) = 0.4613$.
3. $\mathbb{P}(8 \leq X \leq 11) = \mathbb{P}(X \leq 11) - \mathbb{P}(X \leq 7)$.

$$\mathbb{P}(X \leq 11) = \mathbb{P}(Y \geq 4) = 1 - \mathbb{P}(Y \leq 3) = 1 - 0.4613 = 0.5387,$$

$$\mathbb{P}(X \leq 7) = \mathbb{P}(Y \geq 8) = 1 - \mathbb{P}(Y \leq 7) = 1 - 0.9827 = 0.0173.$$

$$\text{Hence } \mathbb{P}(8 \leq X \leq 11) = 0.5387 - 0.0173 = 0.5214.$$

METHOD Fitting a Binomial Distribution to Observed Data

1. Calculate the sample mean: $\bar{x} = \frac{\sum f_i x_i}{\sum f_i}$.
2. Estimate p : for $B(n, p)$, $\mathbb{E}(X) = np$, so set $\hat{p} = \frac{\bar{x}}{n}$ (the value of n is known from the context).
3. Compute expected frequencies: $E_i = N \times \mathbb{P}(X = x_i)$, where $N = \sum f_i$ and $\mathbb{P}(X = x_i) = \binom{n}{x_i} \hat{p}^{x_i} (1 - \hat{p})^{n-x_i}$.
4. Pool adjacent classes if any expected frequency is less than 5 (a requirement for the χ^2 goodness-of-fit test).
5. Compare observed and expected frequencies to judge the fit.

WORKED EXAMPLE 26 Fitting a Binomial Distribution

The following data show the number of boys in 200 families, each with 5 children.

Number of boys (x)	0	1	2	3	4	5
Number of families (f)	8	32	70	60	25	5

Fit a binomial distribution and calculate the expected frequencies.

SOLUTION

Solution.

Here $n = 5$ is fixed and $N = 200$.

$$\bar{x} = \frac{0(8) + 1(32) + 2(70) + 3(60) + 4(25) + 5(5)}{200} = \frac{0 + 32 + 140 + 180 + 100 + 25}{200} = \frac{477}{200} = 2.385.$$

Estimate p : $\hat{p} = \frac{\bar{x}}{n} = \frac{2.385}{5} = 0.477$, so $\hat{q} = 1 - 0.477 = 0.523$.

Compute $\mathbb{P}(X = x) = \binom{5}{x}(0.477)^x(0.523)^{5-x}$, using $(0.523)^2 = 0.2735$, $(0.523)^3 = 0.1431$, $(0.523)^4 = 0.0748$, $(0.523)^5 = 0.0391$.

- $x = 0$: $(0.523)^5 = 0.0391$; $E_0 = 200 \times 0.0391 = 7.83$.
- $x = 1$: $5(0.477)(0.523)^4 = 5 \times 0.477 \times 0.0748 = 0.1784$; $E_1 = 35.69$.
- $x = 2$: $10(0.477)^2(0.523)^3 = 10 \times 0.2275 \times 0.1431 = 0.3255$; $E_2 = 65.10$.
- $x = 3$: $10(0.477)^3(0.523)^2 = 10 \times 0.1085 \times 0.2735 = 0.2968$; $E_3 = 59.35$.
- $x = 4$: $5(0.477)^4(0.523) = 5 \times 0.0518 \times 0.523 = 0.1354$; $E_4 = 27.08$.
- $x = 5$: $(0.477)^5 = 0.0247$; $E_5 = 200 \times 0.0247 = 4.94$.

The expected frequencies sum to $7.83 + 35.69 + 65.10 + 59.35 + 27.08 + 4.94 \approx 200$, as required.

x	0	1	2	3	4	5
Observed f	8	32	70	60	25	5
Expected E	7.8	35.7	65.1	59.4	27.1	4.9

The agreement is close, so a binomial model is reasonable. Note that $E_5 = 4.94 < 5$: for a χ^2 goodness-of-fit test, the last two classes ($x = 4$ and $x = 5$) should be pooled, giving observed 30 and expected 32.0.

METHOD Geometric Distribution: Probabilities, Expectation and Variance

1. Setting: independent Bernoulli trials with success probability p ; X = number of trials

up to and including the first success. Support: $x = 1, 2, 3, \dots$

2. PMF: $\mathbb{P}(X = x) = (1 - p)^{x-1}p = q^{x-1}p.$

3. Expectation: $\mathbb{E}(X) = \frac{1}{p}.$

4. Variance: $\text{Var}(X) = \frac{q}{p^2}.$

5. Tail probabilities: $\mathbb{P}(X > x) = q^x$ (the first x trials all fail); $\mathbb{P}(X \geq x) = q^{x-1}.$

6. Memoryless property: $\mathbb{P}(X > m + n \mid X > m) = \mathbb{P}(X > n)$: past failures do not change future probabilities.

WORKED EXAMPLE 27 Geometric Distribution: Drilling for Oil

An oil company drills wells independently; each well has probability 0.2 of striking oil.

1. Find the probability that the first strike occurs on the 4th well drilled.
2. Find the probability that the first strike occurs on or before the 3rd well.
3. Find the expected number of wells drilled until the first strike, and the variance.
4. Given that the first 5 wells are dry, find the probability that the first strike occurs on the 8th well.

SOLUTION

Solution.

$X \sim \text{Geo}(0.2)$, with $p = 0.2$ and $q = 0.8$.

1. $\mathbb{P}(X = 4) = q^3p = (0.8)^3 \times 0.2 = 0.512 \times 0.2 = 0.1024.$

2. $\mathbb{P}(X \leq 3) = 1 - \mathbb{P}(X > 3) = 1 - q^3 = 1 - 0.512 = 0.488.$

(Check: $\mathbb{P}(X = 1) + \mathbb{P}(X = 2) + \mathbb{P}(X = 3) = 0.2 + 0.16 + 0.128 = 0.488.$)

3. $\mathbb{E}(X) = \frac{1}{p} = \frac{1}{0.2} = 5$ wells; $\text{Var}(X) = \frac{q}{p^2} = \frac{0.8}{0.04} = 20.$

4. By the memoryless property,

$$\mathbb{P}(X = 8 \mid X > 5) = \mathbb{P}(X = 3) = q^2p = (0.8)^2 \times 0.2 = 0.128.$$

$$(\text{Directly: } \mathbb{P}(X = 8 \mid X > 5) = \frac{\mathbb{P}(X = 8)}{\mathbb{P}(X > 5)} = \frac{q^7p}{q^5} = q^2p.)$$

METHOD Poisson Distribution: Characteristics, Expectation, Variance, Recurrence and Mode

1. Setting: random events occurring independently, singly, and at a constant average rate λ per unit interval (of time, length, area, volume, ...).
2. PMF: $\mathbb{P}(X = x) = \frac{e^{-\lambda} \lambda^x}{x!}$, for $x = 0, 1, 2, \dots$
3. Expectation and variance: $\mathbb{E}(X) = \lambda$ and $\text{Var}(X) = \lambda$ (a hallmark of the Poisson distribution: mean equals variance).
4. Recurrence formula: $\mathbb{P}(X = x + 1) = \frac{\lambda}{x + 1} \mathbb{P}(X = x)$, obtained by dividing consecutive terms of the PMF.
5. Mode: if λ is an integer, there are two modes, λ and $\lambda - 1$; otherwise the mode is $\lfloor \lambda \rfloor$.
6. Sum of independent Poisson variables: if $X \sim \text{Po}(\lambda_1)$ and $Y \sim \text{Po}(\lambda_2)$ are independent, then $X + Y \sim \text{Po}(\lambda_1 + \lambda_2)$. In particular, changing the unit interval simply scales λ .

WORKED EXAMPLE 28 Poisson Distribution: Calls at a Call Centre

A call centre in Douala receives an average of 6 calls per hour. Assume a Poisson distribution.

1. Find the probability of exactly 4 calls in a given hour.
2. Find the probability of at most 2 calls in a 30-minute period.
3. Find the probability of more than 10 calls in a 2-hour period, given that $\mathbb{P}(Z \leq 10) = 0.3472$ for $Z \sim \text{Po}(12)$.
4. Determine the mode of the distribution for a 1-hour period.

SOLUTION

Solution.

1. For one hour, $X \sim \text{Po}(6)$.

$$\mathbb{P}(X = 4) = \frac{e^{-6} 6^4}{4!} = e^{-6} \times \frac{1296}{24} = 54 e^{-6} \approx 54 \times 0.002479 = 0.1339.$$

2. For 30 minutes the rate is $\lambda = 6 \times 0.5 = 3$, so $Y \sim \text{Po}(3)$.

$$\mathbb{P}(Y \leq 2) = e^{-3} \left(1 + 3 + \frac{3^2}{2!} \right) = 8.5 e^{-3} \approx 8.5 \times 0.049787 = 0.4232.$$

3. For 2 hours, $\lambda = 12$, so $Z \sim \text{Po}(12)$.

$$\mathbb{P}(Z > 10) = 1 - \mathbb{P}(Z \leq 10) = 1 - 0.3472 = 0.6528.$$

4. $\lambda = 6$ is an integer, so there are two modes: 5 and 6.

Check with the recurrence formula: $\mathbb{P}(X = 6) = \frac{6}{6} \mathbb{P}(X = 5) = \mathbb{P}(X = 5)$, so the two probabilities are equal; and $\mathbb{P}(X = 7) = \frac{6}{7} \mathbb{P}(X = 6) < \mathbb{P}(X = 6)$, confirming the maximum.

METHOD Poisson Approximation to the Binomial Distribution

1. Conditions: n large (typically $n \geq 50$), p small (typically $p \leq 0.1$), and $\lambda = np$ moderate (typically $\lambda \leq 5$).
2. Approximation: $X \sim B(n, p) \approx Y \sim \text{Po}(\lambda)$ with $\lambda = np$.
3. No continuity correction is needed (discrete approximated by discrete).
4. Compute: $\mathbb{P}(X = k) \approx \frac{e^{-\lambda} \lambda^k}{k!}$, and similarly for cumulative probabilities.
5. Decide: if n is small, use the exact binomial; if p is not small but $np > 5$ and $nq > 5$, a normal approximation is preferable.

WORKED EXAMPLE 29 Poisson Approximation to the Binomial

A machine produces 2% defective items. In a batch of 200 items, use a Poisson approximation to find the probability that:

1. there are exactly 3 defectives,
2. there are at most 2 defectives,
3. there are more than 5 defectives.

Comment on the suitability of the approximation.

SOLUTION

Solution.

$n = 200$, $p = 0.02$, so $\lambda = np = 4$. The conditions hold: $n = 200 \geq 50$, $p = 0.02 \leq 0.1$, $\lambda = 4 \leq 5$. Thus $X \sim B(200, 0.02) \approx Y \sim \text{Po}(4)$.

1. $\mathbb{P}(X = 3) \approx \mathbb{P}(Y = 3) = \frac{e^{-4} 4^3}{3!} = \frac{64}{6} e^{-4} \approx 10.6667 \times 0.018316 = 0.1954$.
(Exact binomial value: $\binom{200}{3}(0.02)^3(0.98)^{197} \approx 0.1964$. Very close.)
2. $\mathbb{P}(X \leq 2) \approx \mathbb{P}(Y \leq 2) = e^{-4} \left(1 + 4 + \frac{4^2}{2!}\right) = 13 e^{-4} \approx 13 \times 0.018316 = 0.2381$.
(Exact binomial value: ≈ 0.2351 .)

3. $\mathbb{P}(X > 5) \approx 1 - \mathbb{P}(Y \leq 5)$. Using the recurrence formula $\mathbb{P}(Y = x + 1) = \frac{4}{x+1} \mathbb{P}(Y = x)$:

$$\mathbb{P}(Y = 3) = 0.1954, \quad \mathbb{P}(Y = 4) = \frac{4}{4}(0.1954) = 0.1954, \quad \mathbb{P}(Y = 5) = \frac{4}{5}(0.1954) = 0.1563.$$

$$\mathbb{P}(Y \leq 5) \approx 0.2381 + 0.1954 + 0.1954 + 0.1563 = 0.7852,$$

$$\mathbb{P}(X > 5) \approx 1 - 0.7852 = 0.2148.$$

(Exact binomial value: ≈ 0.2131 .)

The approximation is excellent here because n is large, p is small and $\lambda = 4$ is moderate; the errors are of the order of 0.003 or less.

METHOD Comparing Distributions and Solving Real-Life Problems

1. Identify the random process:
 - fixed number n of trials, two outcomes, constant $p \rightarrow$ Binomial;
 - number of trials up to the first success $\rightarrow$ Geometric;
 - events occurring randomly, singly and at a constant average rate in a continuous interval $\rightarrow$ Poisson.
2. Write down the parameters: (n, p) for the binomial; p for the geometric; $\lambda = \text{rate} \times \text{interval}$ for the Poisson.
3. Choose the appropriate distribution and compute the required probability (exact, cumulative, recurrence or approximation).
4. Use approximations when appropriate: Poisson to binomial (n large, p small); normal to binomial ($np > 5$ and $nq > 5$, with continuity correction); normal to Poisson (λ large, with continuity correction).
5. Interpret the result in context, stating clearly what the probability means for the real situation.

WORKED EXAMPLE 30 Real-Life Problems: Choosing the Right Distribution

1. A typist makes on average 1.5 errors per page. Find the probability that a 3-page letter contains no errors.
2. A multiple-choice test has 50 questions, each with 4 options. A student guesses all the answers. Using a suitable approximation, find the probability that the student gets at least 15 correct.

3. A machine breaks down on average once every 10 days. Find the probability that it breaks down at least twice in a 30-day month.
4. A box contains 100 fuses, 5 of which are defective. A sample of 10 fuses is taken without replacement. Find the probability that exactly 1 is defective, naming the exact distribution and a suitable approximation.

SOLUTION

Solution.

1. Errors occur randomly and independently at a constant average rate over a continuous “interval” (pages): Poisson. For 3 pages, $\lambda = 1.5 \times 3 = 4.5$, so $X \sim \text{Po}(4.5)$.

$$\mathbb{P}(X = 0) = e^{-4.5} \approx 0.0111.$$

2. Fixed $n = 50$ trials, two outcomes, constant $p = \frac{1}{4} = 0.25$, independent guesses: binomial $X \sim B(50, 0.25)$.

Here $np = 12.5 > 5$ and $nq = 37.5 > 5$, but p is not small, so the Poisson approximation is unsuitable; use the normal approximation with continuity correction:

$$X \approx Y \sim N(12.5, 9.375), \quad \sigma = \sqrt{9.375} \approx 3.062.$$

$$\mathbb{P}(X \geq 15) \approx \mathbb{P}\left(Z > \frac{14.5 - 12.5}{3.062}\right) = \mathbb{P}(Z > 0.653) \approx 1 - 0.7431 = 0.2569.$$

3. Breakdowns occur randomly in time at a constant average rate: Poisson. Rate = $\frac{1}{10}$ per day, so for 30 days $\lambda = 3$ and $X \sim \text{Po}(3)$.

$$\mathbb{P}(X \geq 2) = 1 - \mathbb{P}(X = 0) - \mathbb{P}(X = 1) = 1 - e^{-3} - 3e^{-3} = 1 - 4e^{-3} \approx 1 - 0.1992 = 0.8008.$$

4. Sampling without replacement from a finite population: the exact model is the hypergeometric distribution:

$$\mathbb{P}(X = 1) = \frac{\binom{5}{1}\binom{95}{9}}{\binom{100}{10}} \approx 0.3394.$$

Since the sample fraction $\frac{n}{N} = 0.1$ is small, the binomial approximation $B(10, 0.05)$ is suitable:

$$\mathbb{P}(X = 1) \approx \binom{10}{1}(0.05)(0.95)^9 = 10 \times 0.05 \times 0.6302 = 0.3151.$$

(A Poisson approximation with $\lambda = np = 0.5$ gives $\mathbb{P}(Y = 1) = 0.5e^{-0.5} \approx 0.3033$, less accurate here because the binomial approximation is already good.)

WARNING Choosing the wrong model

Two frequent examination errors: (i) using the Poisson approximation when p is not small (e.g. $p = 0.25$) — the results are then badly wrong; (ii) forgetting to rescale λ when the interval changes (e.g. from 1 hour to 30 minutes, λ must be halved). Always state the distribution and its parameters before computing.

13 EXERCISES**13.1 I check my knowledge****EXERCISE 10 MCQ: Identifying a binomial experiment**

Which of the following scenarios can be modelled by a binomial distribution?

1. The number of heads obtained when a fair coin is tossed 10 times.
2. The number of throws of a fair die until a six appears.
3. The number of defective items in a random sample of 20 items taken from a large batch where 5% are defective, without replacement.
4. The number of customers arriving at a shop in a 10-minute interval, given an average of 3 customers per minute.

EXERCISE 11 True/False: Properties of the Poisson distribution

For each statement, state whether it is true or false and justify your answer.

1. If $X \sim \text{Po}(\lambda)$, then $E(X) = \text{Var}(X) = \lambda$.
2. The Poisson distribution is always symmetric.
3. The Poisson distribution can be used to approximate a binomial distribution when n is large and p is small.
4. If $X \sim \text{Po}(4)$, then the mode of X is 4.

EXERCISE 12 Definition: Geometric distribution

Define the geometric distribution with parameter p . Write down its probability mass function, expectation, and variance. Give a concrete example of a random experiment that follows this distribution.

EXERCISE 13 Direct calculation: Binomial probability

A fair die is thrown 10 times. Let X be the number of sixes obtained.

1. State the distribution of X with its parameters.

2. Find $E(X)$ and $\text{Var}(X)$.
3. Calculate $P(X = 2)$.
4. Calculate $P(X \geq 1)$.

13.2 I practise

EXERCISE 14 Binomial distribution: Recurrence formula and mode

Given $X \sim B(12, 0.35)$.

1. Use the recurrence formula $P(X = r + 1) = \frac{n-r}{r+1} \cdot \frac{p}{1-p} P(X = r)$ to compute $P(X = 0)$, $P(X = 1)$, and $P(X = 2)$.
2. Hence find the mode of X .
3. Verify your result using the formula for the mode of a binomial distribution.

EXERCISE 15 Cumulative binomial tables

Using cumulative binomial tables for $X \sim B(20, 0.15)$, find:

1. $P(X = 4)$.
2. $P(X < 3)$.
3. $P(X \geq 5)$.
4. The smallest integer k such that $P(X \leq k) \geq 0.95$.

EXERCISE 16 Fitting a binomial distribution to data

In an experiment, 200 samples of 5 seeds each were planted. The number of germinating seeds per sample was recorded as follows:

Number of germinating seeds (x)	0	1	2	3	4	5
Frequency (f)	12	48	60	52	24	4

1. Estimate the probability p that a single seed germinates.
2. Fit a binomial distribution $B(5, \hat{p})$ to the data and compute the expected frequencies.
3. Compare the observed and expected frequencies. Comment on the goodness of fit.

EXERCISE 17 Poisson distribution: Recurrence formula and mode

Let $X \sim \text{Po}(5)$.

1. Use the recurrence formula $P(X = r + 1) = \frac{\lambda}{r+1}P(X = r)$ to find the mode(s) of X .
2. Compute the probability(ies) at the mode(s).

Now let $Y \sim \text{Po}(4)$.

3. Find the mode(s) of Y and compute the corresponding probability(ies).
4. Comment on the difference between the modes of X and Y .

13.3 I prepare for the GCE**EXERCISE 18 Geometric distribution: Penalty kicks] [15 marks**

A footballer scores on 20% of his penalty kicks. He continues taking penalties until he scores his first goal.

1. Name the distribution that models the number of kicks needed to score the first goal, and state its parameter. [2]
2. Find the probability that his first goal occurs on the 4th kick. [3]
3. Find the probability that he needs more than 5 kicks to score his first goal. [3]
4. Find the expected number of kicks required to score the first goal. [2]
5. What is the probability that he scores at least one goal in his first 3 kicks? [3]
6. If he takes 10 penalties, what is the probability that he scores exactly 2 goals? Name the distribution used. [2]

EXERCISE 19 Poisson distribution: Call centre and rare disease] [20 marks

Part A: Call centre Calls arrive at an office at an average rate of 3 per minute. Assume that the number of calls in disjoint time intervals are independent and that the rate is constant.

1. State the distribution of the number of calls in a one-minute interval, giving its parameter. [2]
2. Find the probability of exactly 2 calls in a given minute. [2]
3. Find the probability of no call in a 30-second interval. [3]
4. Find the probability of more than 8 calls in a 2-minute interval. [3]

5. State two assumptions necessary for the Poisson model to be valid in this context. [2]

Part B: Rare disease A rare disease affects 0.5% of a population. In a village of 400 people, a health survey is conducted.

6. State an exact probability model for the number of cases in the village, and a suitable approximation. Justify the approximation. [3]

7. Using the approximation, find the probability of exactly 2 cases. [2]

8. Find the probability of at least one case. [2]

9. Find the expected number of cases and comment on the result. [1]

EXERCISE 20 Comparison of distributions and real-life problems] [20 marks

For each of the following situations, choose the most appropriate distribution from: Binomial, Geometric, Poisson. Justify your choice by stating the conditions that are satisfied. Then solve the probability question asked.

1. Exam guessing: A multiple-choice test has 20 questions, each with 4 options. A student guesses all answers. Find the probability that the student gets exactly 5 questions correct. [4]
2. First success in a game: A player wins a game with probability 0.3 each time they play, independently. Find the probability that the player's first win occurs on the 4th game. [4]
3. Accidents per week: Accidents occur at a factory at an average rate of 2 per week. Find the probability of exactly 3 accidents in a given week. [4]
4. Defective items in a sample: A batch of 100 items contains 5 defectives. A sample of 10 items is selected at random without replacement. Find the probability that the sample contains exactly 1 defective item. [4]
5. Emails per hour: Emails arrive at a server at an average rate of 12 per hour. Find the probability that more than 15 emails arrive in a given hour. [4]

14 SOLUTIONS TO THE EXERCISES

SOLUTION Exercise 1 — MCQ: Identifying a binomial experiment

1. Yes. Fixed number of trials ($n = 10$), two outcomes (head/tail), constant probability $p = 0.5$, independent tosses. $X \sim B(10, 0.5)$.
2. No. The number of trials is not fixed; we count trials until the first success. This is a **geometric** distribution with $p = \frac{1}{6}$.
3. Yes (as an approximation). Sampling is without replacement, so strictly the distribution is hypergeometric; however the batch is large, so $p = 0.05$ stays essentially constant and $X \approx B(20, 0.05)$ is an excellent model.
4. No. There is no fixed number of trials; events occur randomly in continuous time at an average rate. This is a **Poisson** distribution, $Po(30)$ for a 10-minute interval.

SOLUTION Exercise 2 — True/False: Properties of the Poisson distribution

1. True. For $X \sim Po(\lambda)$, both the expectation and the variance equal the parameter: $E(X) = \text{Var}(X) = \lambda$. This equality is a signature of the Poisson model.
2. False. The Poisson distribution is skewed to the right (positively skewed), especially for small λ . It only becomes approximately symmetric (bell-shaped) when λ is large.
3. True. When n is large, p is small and $\lambda = np$ is moderate, $B(n, p) \approx Po(np)$. The usual rule of thumb is $n \geq 50$ and $p \leq 0.1$.
4. True (with a precision). Since $\lambda = 4$ is an integer, the Poisson distribution has two modes: $\lambda - 1 = 3$ and $\lambda = 4$, with $P(X = 3) = P(X = 4)$. So 4 is indeed a mode, but it shares this status with 3.

SOLUTION Exercise 3 — Definition: Geometric distribution

Definition (Geometric distribution). A discrete random variable X follows a **geometric distribution** with parameter p ($0 < p < 1$), written $X \sim \text{Geo}(p)$, if X counts the number of independent trials needed to obtain the first success, where each trial has constant success probability p .

Probability mass function:

$$P(X = r) = (1 - p)^{r-1}p, \quad r = 1, 2, 3, \dots$$

Expectation and variance:

$$E(X) = \frac{1}{p}, \quad \text{Var}(X) = \frac{1-p}{p^2}.$$

Concrete example: a candidate at the GCE tries a multiple-choice question by guessing ($p = \frac{1}{4}$); the number of guesses needed before the first correct answer follows $\text{Geo}(0.25)$.

Another example: the number of throws of a fair die until the first six appears, $X \sim \text{Geo}(\frac{1}{6})$.

SOLUTION Exercise 4 — Direct calculation: Binomial probability

1. Each throw is an independent trial with success (a six) probability $p = \frac{1}{6}$, and there are $n = 10$ fixed trials:

$$X \sim B(10, \frac{1}{6}).$$

2.

$$E(X) = np = 10 \times \frac{1}{6} = \frac{5}{3} \approx 1.67, \quad \text{Var}(X) = np(1-p) = 10 \times \frac{1}{6} \times \frac{5}{6} = \frac{25}{18} \approx 1.39.$$

3.

$$P(X = 2) = \binom{10}{2} \left(\frac{1}{6}\right)^2 \left(\frac{5}{6}\right)^8 = 45 \times \frac{1}{36} \times 0.23257 \approx 0.2907.$$

4. Using the complement:

$$P(X \geq 1) = 1 - P(X = 0) = 1 - \left(\frac{5}{6}\right)^{10} = 1 - 0.16151 \approx 0.8385.$$

SOLUTION Exercise 5 — Binomial distribution: Recurrence formula and mode

1. With $n = 12$, $p = 0.35$, $\frac{p}{1-p} = \frac{0.35}{0.65} = \frac{7}{13} \approx 0.53846$.

$$P(X = 0) = 0.65^{12} \approx 0.005687.$$

$$P(X = 1) = \frac{12-0}{0+1} \times \frac{7}{13} \times P(X = 0) = 12 \times 0.53846 \times 0.005687 \approx 0.03675.$$

$$P(X = 2) = \frac{12-1}{1+1} \times \frac{7}{13} \times P(X = 1) = \frac{11}{2} \times 0.53846 \times 0.03675 \approx 0.10883.$$

2. The ratio $\frac{P(X = r+1)}{P(X = r)} = \frac{12-r}{r+1} \cdot \frac{7}{13}$ exceeds 1 while $7(12-r) > 13(r+1)$, i.e. $84 - 7r > 13r + 13$, i.e. $r < 3.55$. So probabilities increase up to $r = 4$ and decrease afterwards: the mode is 4.

3. Formula check: $(n+1)p = 13 \times 0.35 = 4.55$, not an integer, so the mode is $\lfloor 4.55 \rfloor = 4$. ✓

SOLUTION Exercise 6 — Cumulative binomial tables

For $X \sim B(20, 0.15)$ the cumulative tables give:

$$P(X \leq 2) = 0.4049, \quad P(X \leq 3) = 0.6477, \quad P(X \leq 4) = 0.8298,$$

$$P(X \leq 5) = 0.9327, \quad P(X \leq 6) = 0.9781.$$

- $P(X = 4) = P(X \leq 4) - P(X \leq 3) = 0.8298 - 0.6477 = 0.1821$.
- $P(X < 3) = P(X \leq 2) = 0.4049$.
- $P(X \geq 5) = 1 - P(X \leq 4) = 1 - 0.8298 = 0.1702$.
- $P(X \leq 5) = 0.9327 < 0.95$ but $P(X \leq 6) = 0.9781 \geq 0.95$, so the smallest such integer is $k = 6$.

SOLUTION Exercise 7 — Fitting a binomial distribution to data

1. Total number of germinating seeds:

$$\sum fx = 0(12) + 1(48) + 2(60) + 3(52) + 4(24) + 5(4) = 48 + 120 + 156 + 96 + 20 = 440.$$

Sample mean: $\bar{x} = \frac{440}{200} = 2.2$ germinating seeds per sample of 5. Since $E(X) = n\hat{p} = 5\hat{p}$, we get $\hat{p} = \frac{2.2}{5} = 0.44$.

2. Expected frequencies $E_r = 200 \binom{5}{r} (0.44)^r (0.56)^{5-r}$:

x	0	1	2	3	4	5
Observed f	12	48	60	52	24	4
Expected E	11.0	43.3	68.0	53.4	21.0	3.3

3. The expected frequencies follow the same pattern as the observed ones: both rise to a peak at $x = 2$ and then fall. The largest discrepancy is at $x = 2$ (60 observed vs 68.0 expected), but the differences are small relative to the frequencies, and $\sum E = 200$ as required. The binomial model $B(5, 0.44)$ fits the data well: germination behaves like 5 independent trials with constant probability $p \approx 0.44$.

SOLUTION Exercise 8 — Poisson distribution: Recurrence formula and modeCase $X \sim \text{Po}(5)$.

1. The ratio of consecutive probabilities is

$$\frac{P(X = r + 1)}{P(X = r)} = \frac{5}{r + 1}.$$

This is > 1 when $r + 1 < 5$, i.e. $r \leq 3$ (probabilities increase), and $= 1$ when $r + 1 = 5$, i.e. $P(X = 5) = P(X = 4)$. For $r \geq 5$ the probabilities decrease. Hence the modes are 4 and 5.

- 2.

$$P(X = 4) = P(X = 5) = e^{-5} \frac{5^4}{4!} = e^{-5} \times \frac{625}{24} \approx 0.0067379 \times 26.0417 \approx 0.1755.$$

Case $Y \sim \text{Po}(4)$.

3. Similarly
- $\frac{P(Y = r + 1)}{P(Y = r)} = \frac{4}{r + 1}$
- , which equals 1 when
- $r = 3$
- : the modes are 3 and 4, with

$$P(Y = 3) = P(Y = 4) = e^{-4} \frac{4^4}{4!} = e^{-4} \times \frac{256}{24} \approx 0.0183156 \times 10.6667 \approx 0.1954.$$

4. In both cases λ is an integer, so each distribution has two consecutive modes, $\lambda - 1$ and λ . The modal probability is larger for $\text{Po}(4)$ ($0.1954 > 0.1755$): a smaller λ concentrates the distribution more sharply near its centre, while a larger λ spreads the probability mass over more values.

SOLUTION Exercise 9 — Geometric distribution: Penalty kicksLet $p = 0.2$ (probability of scoring), $q = 0.8$.

1. The number of kicks until the first goal follows a **geometric distribution**: $X \sim \text{Geo}(0.2)$.
[2 marks: name 1, parameter 1]
2. $P(X = 4) = q^3 p = 0.8^3 \times 0.2 = 0.512 \times 0.2 = 0.1024$. [3 marks: formula 1, substitution 1, answer 1]
3. “More than 5 kicks” means the first 5 kicks all fail:

$$P(X > 5) = q^5 = 0.8^5 = 0.32768 \approx 0.3277.$$

[3 marks]

- 4.
- $E(X) = \frac{1}{p} = \frac{1}{0.2} = 5$
- kicks.

[2 marks]

5. “At least one goal in the first 3 kicks” is the complement of “no goal in 3 kicks”:

$$P(X \leq 3) = 1 - q^3 = 1 - 0.512 = 0.488.$$

[3 marks]

6. Now the number of trials is fixed at $n = 10$ and we count successes: $Y \sim B(10, 0.2)$.

$$P(Y = 2) = \binom{10}{2}(0.2)^2(0.8)^8 = 45 \times 0.04 \times 0.16777 \approx 0.3020.$$

[2 marks: binomial named 1, probability 1]

Examiner's expectations: the candidate must clearly distinguish the geometric setting (number of trials until first success) from the binomial setting (number of successes in a fixed number of trials). Writing $P(X > 5) = q^5$ directly (memorylessness) earns full credit; a term-by-term sum is accepted but slower.

SOLUTION Exercise 10 — Poisson distribution: Call centre and rare disease

Part A: Call centre

1. Calls occur randomly, independently, at a constant average rate of 3 per minute, so the number of calls in one minute is $X \sim \text{Po}(3)$. [2 marks]

2. $P(X = 2) = e^{-3} \frac{3^2}{2!} = 0.049787 \times 4.5 \approx 0.2240$. [2 marks]

3. In 30 seconds the mean is $\lambda = 3 \times 0.5 = 1.5$, so $N \sim \text{Po}(1.5)$:

$$P(N = 0) = e^{-1.5} \approx 0.2231.$$

[3 marks: new λ 1, model 1, answer 1]

4. In 2 minutes, $\lambda = 6$, so $M \sim \text{Po}(6)$. From cumulative Poisson tables, $P(M \leq 8) = 0.8472$:

$$P(M > 8) = 1 - 0.8472 = 0.1528.$$

[3 marks]

5. Any two of: calls arrive independently of each other; calls arrive at a constant average rate; calls occur singly (two calls cannot arrive at exactly the same instant); calls occur randomly in time. [2 marks]

Part B: Rare disease

6. Exact model: $Y \sim B(400, 0.005)$. Since $n = 400$ is large, $p = 0.005$ is small and $\lambda = np = 2$ is moderate, the suitable approximation is

$$Y \approx \text{Po}(2).$$

[3 marks: exact 1, approximation 1, justification 1]

7. $P(Y = 2) = e^{-2} \frac{2^2}{2!} = 2e^{-2} \approx 0.2707$. [2 marks]

8. $P(Y \geq 1) = 1 - P(Y = 0) = 1 - e^{-2} \approx 1 - 0.1353 = 0.8647$. [2 marks]

9. $E(Y) = \lambda = 2$ cases. In a village of 400 people one expects about 2 cases of the disease, which confirms that the disease is rare but likely to appear in a community of this size. [1 mark]

Examiner's expectations: marks are awarded for adjusting the rate to the interval (questions 3 and 4), for quoting the conditions n large, p small, np moderate when justifying the Poisson approximation, and for using the complement in question 8.

SOLUTION Exercise 11 — Comparison of distributions and real-life problems

1. Binomial. Fixed number of trials ($n = 20$), two outcomes per question, constant success probability $p = \frac{1}{4} = 0.25$, independent guesses: $X \sim B(20, 0.25)$.

$$P(X = 5) = \binom{20}{5}(0.25)^5(0.75)^{15} = 15504 \times 0.0009766 \times 0.013363 \approx 0.2023.$$

[4 marks]

2. Geometric. Independent plays, constant win probability $p = 0.3$, counting trials until the first win: $X \sim \text{Geo}(0.3)$.

$$P(X = 4) = (0.7)^3(0.3) = 0.343 \times 0.3 = 0.1029.$$

[4 marks]

3. Poisson. Accidents occur randomly, independently, at a constant average rate $\lambda = 2$ per week, with no fixed number of trials: $X \sim \text{Po}(2)$.

$$P(X = 3) = e^{-2} \frac{2^3}{3!} = e^{-2} \times \frac{8}{6} \approx 0.1353 \times 1.3333 \approx 0.1804.$$

[4 marks]

4. Binomial (approximation). Strictly, sampling without replacement gives a hypergeometric distribution; however the batch (100 items) is large compared with the sample (10), so the defective proportion $p = 0.05$ stays nearly constant and $X \approx B(10, 0.05)$:

$$P(X = 1) = \binom{10}{1}(0.05)(0.95)^9 = 0.5 \times 0.63025 \approx 0.3151.$$

(The exact hypergeometric value is about 0.34; the binomial approximation is the expected GCE answer when a choice among the three distributions is imposed.) [4 marks]


5. Poisson. Emails arrive randomly and independently at a constant average rate $\lambda = 12$ per hour: $X \sim \text{Po}(12)$. From cumulative Poisson tables, $P(X \leq 15) \approx 0.8444$:

$$P(X > 15) = 1 - P(X \leq 15) \approx 1 - 0.8444 = 0.1556.$$

[4 marks]

Examiner's expectations: for each situation the mark scheme rewards (i) the correct distribution by name, (ii) the parameters, (iii) a justification quoting the defining conditions (fixed n and constant p for binomial; "until the first success" for geometric; random independent events at constant rate for Poisson), and (iv) the correct numerical probability rounded to 4 decimal places. In question 4, mentioning that the sampling is without replacement and that the binomial is an approximation shows the deeper understanding that earns full credit.

15 SUMMARY SHEET



Mind map of Chapter PMS12: Special Discrete Probability Distributions

FICHE BILAN L'essentiel à connaître pour l'examen

Key Definitions

- **Binomial Distribution:** $X \sim B(n, p)$ with n independent Bernoulli trials, success probability p . PMF: $P(X = k) = \binom{n}{k} p^k (1 - p)^{n-k}$, $k = 0, \dots, n$.
- **Geometric Distribution:** $X \sim Geo(p)$ counting trials until first success (including success). PMF: $P(X = k) = p(1 - p)^{k-1}$, $k \geq 1$. Alternative: failures before first success $P(Y = k) = p(1 - p)^k$, $k \geq 0$.
- **Poisson Distribution:** $X \sim Po(\lambda)$ for rare events in a fixed interval. PMF: $P(X = k) = e^{-\lambda} \frac{\lambda^k}{k!}$, $k \geq 0$.

Laws, Formulas, Theorems

- **Expectation & Variance:**

$$B(n, p) : E(X) = np, \quad \text{Var}(X) = np(1 - p).$$

$$Geo(p) : E(X) = \frac{1}{p}, \quad \text{Var}(X) = \frac{1 - p}{p^2}.$$

$$Po(\lambda) : E(X) = \lambda, \quad \text{Var}(X) = \lambda.$$

- Recurrence Formulae:

$$B(n, p) : P(X = k + 1) = \frac{n - k}{k + 1} \frac{p}{1 - p} P(X = k).$$

$$Po(\lambda) : P(X = k + 1) = \frac{\lambda}{k + 1} P(X = k).$$

- Mode:

- Binomial: $[(n + 1)p]$ (or two modes if $(n + 1)p$ integer).
- Poisson: $[\lambda]$ (or λ and $\lambda - 1$ if λ integer).
- Poisson Approximation to Binomial: If n large, p small, $np = \lambda$ moderate, then $B(n, p) \approx Po(\lambda)$. Rule of thumb: $n \geq 50$, $p \leq 0.05$, $np \leq 5$.
- Cumulative Distribution Function: $F(k) = P(X \leq k) = \sum_{i=0}^k P(X = i)$. Use binomial tables for $n \leq 50$; for larger n use normal approximation (not in this chapter).

Methods

1. Identify the distribution: Check for fixed n and two outcomes $\rightarrow$ Binomial; waiting for first success $\rightarrow$ Geometric; rare events in continuous interval $\rightarrow$ Poisson.
2. Compute probabilities: Use PMF directly, recurrence to build successive terms, or cumulative tables.
3. Fit a binomial to data: Estimate p by $\hat{p} = \frac{\text{total successes}}{n \times \text{frequency}}$; compute expected frequencies $E_i = N \times P(X = i)$; check χ^2 goodness-of-fit (if required).
4. Approximate binomial by Poisson: When conditions hold, replace $B(n, p)$ with $Po(np)$ for easier calculation.
5. Solve real-life problems: Translate word problem into parameters, choose distribution, compute required probability or expectation, interpret answer in context.

Common Mistakes

- Confusing the two definitions of geometric distribution (trials vs failures). Always state which version you use.
- Forgetting that binomial requires independent trials with constant p .
- Using Poisson approximation when np is not moderate (e.g. $np > 5$ or p not small).
- Misreading cumulative tables: $P(X \leq k)$ vs $P(X < k)$.
- In fitting, not pooling expected frequencies < 5 before χ^2 test.
- Mixing up variance formulas: $np(1 - p)$ for binomial, λ for Poisson, $\frac{1-p}{p^2}$ for geometric.

GCE Tips

- Show all steps: Write the distribution name, parameters, formula, substitution, and final answer.

- Use recurrence to compute several probabilities quickly in exams; it saves time and reduces arithmetic errors.
- Tables: If $n \leq 50$ and p is in the table, use it; otherwise compute manually or approximate.
- Approximation justification: State clearly why Poisson approximation is valid (large n , small p , np moderate).
- Interpretation: Always conclude with a sentence answering the question in context (e.g. “The probability that at most 2 defects occur is 0.92”).
- Time management: Binomial and Poisson questions often carry 6–8 marks; allocate 8–10 minutes each.

**Signed • OnBuch+**ONBUCH+ TEACHING TEAM
FOLLOWS THE OFFICIAL MINESEC SYLLABUS**OnBuch+**