

PURE MATHEMATICS WITH STATISTICS • UPPER SIXTH

UPPER SIXTH — PURE MATHEMATICS WITH STATISTICS

CHAPTER 14

SPECIAL CONTINUOUS DISTRIBUTIONS

SUGGESTED TIME: 18 PERIODS

IN THIS CHAPTER

This chapter studies the three special continuous probability distributions required at GCE Advanced Level: the uniform (rectangular), the exponential and the Normal distributions. Students learn their probability density functions, expectations, variances, the use of standard Normal tables, continuity corrections, Normal approximations to the binomial and Poisson distributions, and combinations of independent Normal variables.

- By the end of this chapter I can state and use the probability density function, expectation and variance of a uniform (rectangular) distribution.
- By the end of this chapter I can solve applied problems modelled by the uniform distribution.
- By the end of this chapter I can derive and use the expectation and variance of an exponential distribution and apply it to waiting-time problems.
- By the end of this chapter I can explain the memoryless property of the exponential distribution and its link with the Poisson distribution.
- By the end of this chapter I can describe the Normal curve, standardise a Normal variable and read the standard Normal tables $\Phi(z)$.
- By the end of this chapter I can use the symmetry properties of $\Phi(z)$ and find central 95% and 99% intervals.

DOWNLOAD ON-
BUCH

Free • Google Play
AI tutor, past papers,
results



GO ONBUCH+

All signed courses, unlimited AI
tutor, PDF sheets — OnBuch+
tab in the app

THIS SIGNED COURSE CONTAINS

COURSE

complete, illustrated,
step by step

METHODS

and worked examples

EXERCISES

exam style + solutions

SUMMARY
SHEET

the essentials to
remember



Signed • OnBuch+

ONBUCH+ TEACHING TEAM
FOLLOWS THE OFFICIAL MINESEC SYLLABUS

OnBuch+

CONTENTS

- | | |
|---|---|
| 1. The Uniform (Rectangular) Distribution | 7. Combinations of Independent Normal Variables |
| 2. The Exponential Distribution | 8. Integration, Assessment and Remediation |
| 3. The Normal Distribution and Standardisation | 9. Integration activity |
| 4. Central Intervals and Working with Any Normal Distribution | 10. Methods and worked examples |
| 5. Determining Unknown μ and/or σ | 11. Exercises |
| 6. Continuity Correction and Normal Approximations | 12. Solutions to the exercises |
| | 13. Summary sheet |

CHAPTER OBJECTIVES

- By the end of this chapter I can state and use the probability density function, expectation and variance of a uniform (rectangular) distribution.
- By the end of this chapter I can solve applied problems modelled by the uniform distribution.
- By the end of this chapter I can derive and use the expectation and variance of an exponential distribution and apply it to waiting-time problems.
- By the end of this chapter I can explain the memoryless property of the exponential distribution and its link with the Poisson distribution.
- By the end of this chapter I can describe the Normal curve, standardise a Normal variable and read the standard Normal tables $\Phi(z)$.
- By the end of this chapter I can use the symmetry properties of $\Phi(z)$ and find central 95% and 99% intervals.
- By the end of this chapter I can de-standardise and determine unknown values of μ and/or σ from given probabilities.
- By the end of this chapter I can apply continuity corrections and use the Normal approximation to the binomial and Poisson distributions.
- By the end of this chapter I can find the distribution of sums and differences of two independent Normal variables.

PREREQUISITES

- Continuous random variables: probability density functions, $P(a < X < b)$ as an area, cumulative distribution function (Lower Sixth / PMS on continuous random variables)
- Expectation $E(X)$ and variance $\text{Var}(X)$ of continuous random variables, including $E(aX + b)$ and $\text{Var}(aX + b)$
- The Poisson distribution: conditions of validity, mean = variance = λ , calculation of $P(X = r)$
- The binomial distribution $B(n, p)$: mean np , variance npq , conditions of use
- Basic integration of polynomials and exponential functions, and use of limits at infinity

1 THE UNIFORM (RECTANGULAR) DISTRIBUTION

At the Mfoundi market in Yaoundé, a vendor notices that customers arrive at her stall roughly every 4 minutes, while the waiting time at the SONEL cash office across the road seems completely unpredictable — sometimes 1 minute, sometimes 40. At the GCE marking centre, examiners observe that candidates' marks in Mathematics cluster around a central value with fewer and fewer scripts at the extremes. Why do waiting times, lifetimes of phone batteries, and examination marks follow such different yet perfectly predictable patterns? This chapter reveals the three special continuous distributions — **uniform**, **exponential** and **Normal** — that model these everyday Cameroonian realities and that dominate the GCE Advanced Level statistics paper.

We begin with the simplest of the three: the uniform distribution, in which every value in an interval is equally likely.

1.1 Introducing the Idea: A Concrete Situation

EXPERIMENT The bus that comes "any time within 20 minutes"

A student at the Carrefour Warda bus stop in Yaoundé is told that the next bus will arrive at a completely random instant within the next 20 minutes: no moment is more likely than any other. Let X be the waiting time, in minutes. Then X takes values in $[0, 20]$ and:

- the probability of waiting between 5 and 10 minutes should be the same as waiting between 12 and 17 minutes (both intervals have length 5);
- the probability of waiting more than 15 minutes should be $\frac{5}{20} = \frac{1}{4}$.

This "equal chance everywhere on an interval" is exactly the spirit of the uniform distribution. Probabilities depend only on the length of the interval considered, not on its position.

1.2 Definition and Probability Density Function

DEFINITION Continuous uniform (rectangular) distribution

A continuous random variable X follows the **uniform distribution** on the interval $[a, b]$ (with $a < b$), written $X \sim U(a, b)$, if its probability density function is

$$f(x) = \begin{cases} \frac{1}{b-a} & \text{for } a \leq x \leq b, \\ 0 & \text{otherwise.} \end{cases}$$

Because the graph of f is a rectangle, this is also called the **rectangular distribution**.

Let us check that f is a genuine pdf. It is clearly non-negative, and

$$\int_{-\infty}^{+\infty} f(x) dx = \int_a^b \frac{1}{b-a} dx = \frac{b-a}{b-a} = 1.$$

The total area under the curve is the area of a rectangle of width $b - a$ and height $\frac{1}{b-a}$, which is indeed 1.

PROPERTY Probabilities as areas of rectangles


If $X \sim U(a, b)$ and $[c, d] \subseteq [a, b]$, then

$$P(c < X < d) = \int_c^d \frac{1}{b-a} dx = \frac{d-c}{b-a}.$$

The probability is the ratio of the length of the sub-interval to the length of the whole interval. (Proof not examinable; it follows directly from the definition of a pdf.)

WARNING Endpoints do not matter

For a continuous variable, $P(X = c) = 0$, so $P(c < X < d)$, $P(c \leq X \leq d)$, $P(c \leq X < d)$ are all equal. Never waste time in the exam worrying about strict versus non-strict inequalities for continuous distributions.



Pdf of $X \sim U(a, b)$: the probability $P(c < X < d)$ is the area of the shaded rectangle, $\frac{d-c}{b-a}$.

1.3 Expectation and Variance

PROPERTY Mean and variance of $U(a, b)$ — proof examinable

If $X \sim U(a, b)$, then

$$E(X) = \frac{a+b}{2} \quad \text{and} \quad \text{Var}(X) = \frac{(b-a)^2}{12}.$$

Derivation of $E(X)$ by integration.

$$E(X) = \int_a^b x \cdot \frac{1}{b-a} dx = \frac{1}{b-a} \left[\frac{x^2}{2} \right]_a^b = \frac{b^2 - a^2}{2(b-a)} = \frac{(b-a)(b+a)}{2(b-a)} = \frac{a+b}{2}.$$

Derivation of $E(X)$ by symmetry. The pdf is symmetric about the midpoint of $[a, b]$, namely $\frac{a+b}{2}$. For any symmetric continuous distribution, the mean is the axis of symmetry, so $E(X) = \frac{a+b}{2}$ immediately. This argument is a useful check in the exam.

Derivation of $\text{Var}(X)$. First compute $E(X^2)$:

$$E(X^2) = \int_a^b x^2 \cdot \frac{1}{b-a} dx = \frac{1}{b-a} \left[\frac{x^3}{3} \right]_a^b = \frac{b^3 - a^3}{3(b-a)} = \frac{a^2 + ab + b^2}{3},$$

using $b^3 - a^3 = (b-a)(a^2 + ab + b^2)$. Then

$$\begin{aligned} \text{Var}(X) &= E(X^2) - [E(X)]^2 = \frac{a^2 + ab + b^2}{3} - \frac{(a+b)^2}{4} \\ &= \frac{4(a^2 + ab + b^2) - 3(a^2 + 2ab + b^2)}{12} = \frac{a^2 - 2ab + b^2}{12} = \frac{(b-a)^2}{12}. \end{aligned}$$

WARNING The most common GCE error

The variance is $\text{Var}(X) = \frac{(b-a)^2}{12}$, not $\frac{b-a}{12}$. The squaring is forgotten by a large number of candidates every year. Remember: a variance has the units of X squared, so $(b-a)$ must appear squared.

KEY POINTS Key facts about $X \sim U(a, b)$

- Pdf: $f(x) = \frac{1}{b-a}$ on $[a, b]$, 0 elsewhere.
- $P(c < X < d) = \frac{d-c}{b-a}$ for $[c, d] \subseteq [a, b]$.
- $E(X) = \frac{a+b}{2}$, $\text{Var}(X) = \frac{(b-a)^2}{12}$, $\sigma = \frac{b-a}{\sqrt{12}} = \frac{b-a}{2\sqrt{3}}$.
- Cdf: $F(x) = \frac{x-a}{b-a}$ for $a \leq x \leq b$.

1.4 The Cumulative Distribution Function


PROPERTY Cdf of the uniform distribution

For $X \sim U(a, b)$,

$$F(x) = P(X \leq x) = \begin{cases} 0 & \text{if } x < a, \\ \frac{x-a}{b-a} & \text{if } a \leq x \leq b, \\ 1 & \text{if } x > b. \end{cases}$$

Justification. For $a \leq x \leq b$, $F(x) = \int_a^x \frac{1}{b-a} dt = \frac{x-a}{b-a}$; outside $[a, b]$ the cdf takes its limiting values 0 and 1.

The cdf is convenient for probabilities of the form $P(X \leq x)$ or $P(X > x) = 1 - F(x)$, and for finding percentiles: the median m satisfies $F(m) = \frac{1}{2}$, giving $m = \frac{a+b}{2}$, which coincides with the mean — as expected for a symmetric distribution.



Cdf of $U(a, b)$: zero before a , a straight ramp of slope $\frac{1}{b-a}$ on $[a, b]$, then 1 after b .

1.5 Method and Worked Examples

METHOD Computing probabilities with a uniform distribution

1. Sketch the pdf: a rectangle over $[a, b]$ of height $\frac{1}{b-a}$.
2. Locate the interval asked for on the sketch.
3. Compute the area of the corresponding rectangle: $\text{probability} = \frac{\text{length of sub-interval}}{b-a}$.
4. For $E(X)$ and $\text{Var}(X)$, quote the standard formulae directly.

EXAMPLE Exam tip

Always sketch the pdf first. With a uniform distribution the picture literally is the calculation: probabilities are areas of rectangles, so a correct sketch gives the answer immediately and lets you spot absurd results (e.g. a probability greater than 1).

WORKED EXAMPLE 1 Waiting for the bus

The waiting time X (in minutes) for a bus at a stop is uniformly distributed on $[0, 20]$.

1. Find the probability that a passenger waits between 6 and 14 minutes.
2. Find the probability of waiting more than 15 minutes.
3. Find the mean and standard deviation of the waiting time.
4. Given that the passenger has already waited 10 minutes, find the probability that the total wait exceeds 15 minutes.

SOLUTION

Solution. Here $a = 0$, $b = 20$, so $f(x) = \frac{1}{20}$ on $[0, 20]$.

1. $P(6 < X < 14) = \frac{14 - 6}{20} = \frac{8}{20} = 0.4$.
2. $P(X > 15) = \frac{20 - 15}{20} = \frac{5}{20} = 0.25$.
3. $E(X) = \frac{0 + 20}{2} = 10$ minutes; $\text{Var}(X) = \frac{20^2}{12} = \frac{100}{3}$, so $\sigma = \sqrt{\frac{100}{3}} \approx 5.77$ minutes.
4. Conditional probability:

$$P(X > 15 \mid X > 10) = \frac{P(X > 15)}{P(X > 10)} = \frac{5/20}{10/20} = \frac{1}{2}.$$

Given 10 minutes have elapsed, the remaining wait is uniform on $[10, 20]$, and half of that interval lies beyond 15 — the picture confirms the answer.

WORKED EXAMPLE 2 Finding b from the variance

A random variable X is uniformly distributed on $[2, b]$ with $b > 2$, and $\text{Var}(X) = 12$. Find b , and hence write down $E(X)$ and $P(X > 8)$.

SOLUTION

Solution. We use $\text{Var}(X) = \frac{(b-a)^2}{12}$ with $a = 2$:

$$\frac{(b-2)^2}{12} = 12 \implies (b-2)^2 = 144 \implies b-2 = 12 \implies b = 14$$

(taking the positive root since $b > 2$). Then

$$E(X) = \frac{2+14}{2} = 8, \quad P(X > 8) = \frac{14-8}{14-2} = \frac{6}{12} = \frac{1}{2}.$$

Note that $P(X > 8) = \frac{1}{2}$ is obvious from the sketch, since 8 is the midpoint of $[2, 14]$.

1.6 Applications of the Rectangular Distribution

The uniform distribution appears whenever "all values in a range are equally likely":

- **Random waiting.** A bus, a shared taxi or a customer known to arrive "at any time within the next T minutes" gives $X \sim U(0, T)$.
- **Rounding errors.** When a measurement is rounded to the nearest integer (or nearest 100 FCFA in a market transaction), the rounding error is modelled by $U(-0.5, 0.5)$, with mean 0 and variance $\frac{1}{12}$. This is why the number 12 appears so often in error analysis.
- **Random number generation.** The random numbers produced by a calculator or computer between 0 and 1 follow $U(0, 1)$; this is the foundation of simulation methods.

WORKED EXAMPLE 3 Rounding errors

The mass of a bag of garri is recorded to the nearest gram. The rounding error E (in grams) is uniformly distributed on $[-0.5, 0.5]$. Find the probability that the error exceeds 0.3 g in absolute value, and the standard deviation of E .

SOLUTION

Solution. We need $P(|E| > 0.3) = P(E > 0.3) + P(E < -0.3)$. By symmetry of the rectangle:

$$P(|E| > 0.3) = 2 \times \frac{0.5 - 0.3}{1} = 2 \times 0.2 = 0.4.$$

The standard deviation is

$$\sigma = \frac{b-a}{\sqrt{12}} = \frac{1}{\sqrt{12}} = \frac{1}{2\sqrt{3}} \approx 0.289 \text{ g}.$$

DID YOU KNOW? Why $\frac{1}{12}$?

The variance $\frac{(b-a)^2}{12}$ of the uniform distribution is one of the most quoted results in engineering and metrology. When instruments round readings, the total uncertainty of a sum of many rounded measurements is computed using exactly this $\frac{1}{12}$ factor. The same formula underlies the "quantisation noise" theory used in digital music and mobile phone signal processing — technology used every day in Douala and Bamenda!

EXERCISE 1

The time T (in minutes) taken by a photocopying machine at a cybercafé in Buea to warm up is uniformly distributed between 1 and 5 minutes.

1. Write down the pdf of T and sketch it.
2. Find $P(T < 2.5)$ and $P(2 < T < 4.5)$.
3. Calculate $E(T)$ and $\text{Var}(T)$.
4. Find the value t such that $P(T < t) = 0.75$.

SOLUTION Exercise 1

1. $f(t) = \frac{1}{4}$ for $1 \leq t \leq 5$, 0 otherwise: a rectangle of height 0.25 over $[1, 5]$.
2. $P(T < 2.5) = \frac{2.5 - 1}{4} = 0.375$; $P(2 < T < 4.5) = \frac{4.5 - 2}{4} = 0.625$.
3. $E(T) = \frac{1+5}{2} = 3$ minutes; $\text{Var}(T) = \frac{(5-1)^2}{12} = \frac{16}{12} = \frac{4}{3} \text{ min}^2$.
4. $F(t) = \frac{t-1}{4} = 0.75 \Rightarrow t-1 = 3 \Rightarrow t = 4$ minutes.

EXERCISE 2

$X \sim U(a, 10)$ and $P(X < 6) = 0.5$. Find a , $E(X)$ and $\text{Var}(X)$.

SOLUTION Exercise 2

$$P(X < 6) = \frac{6-a}{10-a} = 0.5 \implies 2(6-a) = 10-a \implies 12-2a = 10-a \implies a = 2. \text{ Then}$$

$$E(X) = \frac{2+10}{2} = 6 \text{ (consistent with } P(X < 6) = 0.5) \text{ and } \text{Var}(X) = \frac{(10-2)^2}{12} = \frac{64}{12} = \frac{16}{3}.$$

WARNING Final checklist for the GCE

- Check that the interval given in the question really is $[a, b]$ before applying formulae.
- Square $(b - a)$ in the variance formula.
- For conditional probabilities, restrict the interval first, then use lengths.
- Sketch the rectangle: it costs ten seconds and saves marks.

2 THE EXPONENTIAL DISTRIBUTION

In the previous section we studied the uniform distribution, in which every value in an interval $[a, b]$ is equally likely. That model suits situations with complete symmetry. But many real waiting times are not symmetric at all: the time you wait for a taxi in Douala, the lifetime of a phone battery, the gap between two customers arriving at an MTN service counter. Such times are often short, occasionally long, and never negative. The distribution that models them is the **exponential distribution**, one of the most important special continuous distributions in the GCE syllabus.

2.1 Definition and probability density function

EXPERIMENT A motivating observation

Customers arrive at a bank counter in Yaoundé at an average rate of $\lambda = 2$ customers per minute. Let X be the waiting time (in minutes) between two consecutive arrivals. Observation shows that very short gaps are the most frequent, and long gaps become rarer and rarer. A histogram of such gaps has the shape of a rapidly decreasing curve starting at $x = 0$. The curve $f(x) = \lambda e^{-\lambda x}$ fits this behaviour exactly.

DEFINITION Exponential distribution

A continuous random variable X follows the **exponential distribution** with parameter $\lambda >$


0, written $X \sim \text{Exp}(\lambda)$, if its probability density function is

$$f(x) = \begin{cases} \lambda e^{-\lambda x} & \text{if } x \geq 0, \\ 0 & \text{if } x < 0. \end{cases}$$

The parameter λ is a **rate**: the average number of events per unit time.

Let us check that f is a genuine pdf. Clearly $f(x) \geq 0$ for all x , and

$$\int_{-\infty}^{+\infty} f(x) dx = \int_0^{+\infty} \lambda e^{-\lambda x} dx = [-e^{-\lambda x}]_0^{+\infty} = 0 - (-1) = 1.$$



The pdf $f(x) = \lambda e^{-\lambda x}$ for $\lambda = 0.5, 1, 2$. Larger λ means faster decay: events occur more often, so long waiting times become rarer.

2.2 The cumulative distribution function and survival probability

For $x \geq 0$,

$$F(x) = P(X \leq x) = \int_0^x \lambda e^{-\lambda t} dt = [-e^{-\lambda t}]_0^x = 1 - e^{-\lambda x}.$$

PROPERTY CDF and survival probability

If $X \sim \text{Exp}(\lambda)$, then for all $x \geq 0$:

$$F(x) = P(X \leq x) = 1 - e^{-\lambda x}, \quad P(X > x) = e^{-\lambda x}.$$

The quantity $P(X > x)$ is called the **survival probability**: the probability that the event has not yet occurred by time x .

EXAMPLE First calculation

The lifetime X (in years) of a certain electronic component is exponential with $\lambda = 0.25$ per year. Then

$$P(X > 4) = e^{-0.25 \times 4} = e^{-1} \approx 0.3679,$$

and $P(X \leq 4) = 1 - e^{-1} \approx 0.6321$.

2.3 Expectation and variance of the exponential distribution

These derivations are **examinable**: you must be able to carry out the integration by parts carefully, with limits.

Expectation. By definition,

$$E(X) = \int_0^{+\infty} x \lambda e^{-\lambda x} dx.$$

Integrate by parts with $u = x$ (so $du = dx$) and $dv = \lambda e^{-\lambda x} dx$ (so $v = -e^{-\lambda x}$):

$$\begin{aligned} E(X) &= [-xe^{-\lambda x}]_0^{+\infty} + \int_0^{+\infty} e^{-\lambda x} dx \\ &= (0 - 0) + \left[-\frac{1}{\lambda} e^{-\lambda x} \right]_0^{+\infty} = \frac{1}{\lambda}, \end{aligned}$$

since $xe^{-\lambda x} \rightarrow 0$ as $x \rightarrow +\infty$ (exponential decay beats linear growth).

Variance. First compute $E(X^2) = \int_0^{+\infty} x^2 \lambda e^{-\lambda x} dx$. Integrate by parts with $u = x^2$, $dv = \lambda e^{-\lambda x} dx$:

$$\begin{aligned} E(X^2) &= [-x^2 e^{-\lambda x}]_0^{+\infty} + \int_0^{+\infty} 2x e^{-\lambda x} dx \\ &= 0 + \frac{2}{\lambda} \int_0^{+\infty} x \lambda e^{-\lambda x} dx = \frac{2}{\lambda} E(X) = \frac{2}{\lambda^2}. \end{aligned}$$

Therefore

$$\text{Var}(X) = E(X^2) - [E(X)]^2 = \frac{2}{\lambda^2} - \frac{1}{\lambda^2} = \frac{1}{\lambda^2}.$$

PROPERTY Mean and variance of $\text{Exp}(\lambda)$ — proof examinable

If $X \sim \text{Exp}(\lambda)$, then

$$E(X) = \frac{1}{\lambda}, \quad \text{Var}(X) = \frac{1}{\lambda^2}, \quad \sigma = \frac{1}{\lambda}.$$

Interpretation: $1/\lambda$ is the **mean waiting time** between events. Remarkably, the standard deviation equals the mean.

WARNING λ is a rate — convert units first!

λ must be expressed in events per unit of the time used for x . If calls arrive at a rate of 3 per hour and X is measured in minutes, then $\lambda = \frac{3}{60} = 0.05$ per minute, and $E(X) = \frac{1}{0.05} = 20$ minutes. Mixing hours and minutes is the most frequent error in GCE scripts. Always state your unit and convert before substituting.

2.4 The memoryless (no-ageing) property**EXPERIMENT** A surprising fact

A bulb has already burned for 1000 hours without failing. Does this make failure in the next hour more likely? For an exponential lifetime, the answer is no: the bulb is “as good as new”. The distribution forgets the past.

PROPERTY Memoryless property — proof examinable

If $X \sim \text{Exp}(\lambda)$, then for all $s, t \geq 0$:

$$P(X > s + t \mid X > s) = P(X > t).$$

Proof. By the definition of conditional probability, and since $\{X > s + t\} \subset \{X > s\}$:

$$P(X > s + t \mid X > s) = \frac{P(X > s + t)}{P(X > s)} = \frac{e^{-\lambda(s+t)}}{e^{-\lambda s}} = e^{-\lambda t} = P(X > t). \quad \blacksquare$$

Interpretation: given that the component has survived s units of time, the probability that it survives a further t units is the same as the probability that a brand-new component survives t units. The exponential distribution is the only continuous distribution with this property.

EXAMPLE Using memorylessness

With $X \sim \text{Exp}(0.25)$ (lifetimes in years), the probability that a component already 4 years old lasts at least 2 more years is

$$P(X > 6 \mid X > 4) = P(X > 2) = e^{-0.25 \times 2} = e^{-0.5} \approx 0.6065.$$

2.5 The link with the Poisson distribution

Suppose events occur randomly in time at a constant average rate λ per unit time, independently of one another (a **Poisson process**). Then two random variables describe the same situ-

ation from two different points of view:


- the **number** N of events occurring in a time interval of length t is discrete: $N \sim \text{Po}(\lambda t)$;
- the **waiting time** T between two consecutive events is continuous: $T \sim \text{Exp}(\lambda)$.

PROPERTY Poisson counts $\leftrightarrow$ exponential waiting times

In a Poisson process of rate λ , the waiting time T until the next event satisfies $T \sim \text{Exp}(\lambda)$. Indeed, “ $T > t$ ” means “no event occurs in $[0, t]$ ”, and since the count in $[0, t]$ is $\text{Po}(\lambda t)$:

$$P(T > t) = P(N = 0) = \frac{e^{-\lambda t}(\lambda t)^0}{0!} = e^{-\lambda t},$$

which is exactly the survival probability of $\text{Exp}(\lambda)$.



A Poisson process on the time axis: events $E_1, E_2, \dots$ occur at rate λ ; the gaps between consecutive events are independent exponential waiting times.

METHOD Poisson or exponential? Choosing the right model

1. Read the question and identify the random variable.
2. If it counts events in a fixed interval (“how many customers in 10 minutes”), use $N \sim \text{Po}(\lambda t)$ with t the length of the interval.
3. If it measures a waiting time, duration or lifetime (“how long until the next call”), use $T \sim \text{Exp}(\lambda)$: $P(T > t) = e^{-\lambda t}$, $P(T \leq t) = 1 - e^{-\lambda t}$.
4. Convert λ to the same time unit as the variable before substituting.
5. Translate boundary phrases: “at least” $\rightarrow P(T \geq t) = e^{-\lambda t}$; “no event in the interval” $\rightarrow P(N = 0) = e^{-\lambda t}$.

WORKED EXAMPLE 4 GCE-style problem mixing Poisson and exponential

Calls arrive at a call centre in Bamenda at an average rate of 4 per hour, according to a Poisson process.

- Find the probability that exactly 3 calls arrive in one hour.
- Find the probability that the operator waits more than 30 minutes for the next call.
- Given that no call has arrived for 20 minutes, find the probability that the operator waits at least 10 further minutes.
- Find the mean and standard deviation of the waiting time between calls.

SOLUTION

Work in hours: $\lambda = 4$ per hour.

- (a) The count in one hour is $N \sim \text{Po}(4)$:

$$P(N = 3) = \frac{e^{-4} 4^3}{3!} = \frac{64e^{-4}}{6} \approx 0.1954.$$

- (b) Let T be the waiting time in hours, $T \sim \text{Exp}(4)$. Since 30 min = 0.5 h:

$$P(T > 0.5) = e^{-4 \times 0.5} = e^{-2} \approx 0.1353.$$

- (c) By the memoryless property:

$$P\left(T > \frac{30}{60} \mid T > \frac{20}{60}\right) = P\left(T > \frac{10}{60}\right) = e^{-4/6} = e^{-2/3} \approx 0.5134.$$

- (d) $E(T) = \frac{1}{\lambda} = \frac{1}{4} \text{ h} = 15 \text{ minutes}$, and $\sigma = \frac{1}{\lambda} = 15 \text{ minutes}$ as well.

WARNING Common mistakes

- Using $P(X > t) = 1 - e^{-\lambda t}$: this is $F(t) = P(X \leq t)$. The survival probability is $e^{-\lambda t}$.
- Forgetting that for a continuous variable $P(X = t) = 0$, so $P(X \geq t) = P(X > t)$.
- Writing $E(X) = \lambda$: the mean is $\frac{1}{\lambda}$, the mean waiting time.
- Applying the memoryless property to a non-exponential lifetime without justification.

EXERCISE 3 Practice

The time X (in minutes) between arrivals of customers at a petrol station in Buea is exponential with mean 5 minutes.

- Write down λ and the pdf of X .
- Find $P(X > 10)$ and $P(2 \leq X \leq 8)$.

(c) A customer has just arrived. Find the probability that exactly two customers arrive in the next 10 minutes.

(d) Given that nobody has arrived for 6 minutes, find the probability of waiting at least 4 more minutes.

SOLUTION Exercise 1

(a) $E(X) = 1/\lambda = 5$, so $\lambda = 0.2$ per minute and $f(x) = 0.2e^{-0.2x}$, $x \geq 0$.

(b) $P(X > 10) = e^{-2} \approx 0.1353$; $P(2 \leq X \leq 8) = e^{-0.4} - e^{-1.6} \approx 0.6703 - 0.2019 = 0.4684$.

(c) Count in 10 min: $N \sim \text{Po}(2)$, so $P(N = 2) = \frac{e^{-2}2^2}{2!} = 2e^{-2} \approx 0.2707$.

(d) Memoryless: $P(X > 10 \mid X > 6) = P(X > 4) = e^{-0.8} \approx 0.4493$.

KEY POINTS Key points

- $X \sim \text{Exp}(\lambda)$: $f(x) = \lambda e^{-\lambda x}$ ($x \geq 0$), $F(x) = 1 - e^{-\lambda x}$, $P(X > x) = e^{-\lambda x}$.
- $E(X) = \frac{1}{\lambda}$, $\text{Var}(X) = \frac{1}{\lambda^2}$ (derivation by parts examinable).
- λ is a rate per unit time; $1/\lambda$ is the mean waiting time. Convert units!
- Memoryless: $P(X > s + t \mid X > s) = P(X > t)$.
- Poisson process of rate λ : counts are $\text{Po}(\lambda t)$, gaps are $\text{Exp}(\lambda)$ — two faces of the same situation.

DID YOU KNOW? Did you know?

The exponential distribution models radioactive decay: the waiting time until a single unstable nucleus decays is exponential, which is why the number of decays per second in a sample follows a Poisson law. The same mathematics that describes a Geiger counter in a physics laboratory describes the queue at an Express Union counter in town — a beautiful example of the unity of mathematics.

3 THE NORMAL DISTRIBUTION AND STANDARDISATION

In the previous section we studied the exponential distribution, a model for waiting times which is strongly skewed: small values are the most likely, and the density decreases steadily. Nature, however, offers us a very different and far more common pattern. Measure the heights of all the

students in an Upper Sixth class in Bamenda, the masses of bags of cement leaving a factory in Douala, or the marks of thousands of candidates at the GCE: in each case the values cluster around a central average, and values far from the centre — on both sides — become rarer and rarer, in a perfectly symmetric way. The histogram has the shape of a **bell**. The mathematical model behind this bell shape is the **Normal distribution**, the most important distribution in all of statistics, and the heart of the GCE Advanced Level statistics paper.

3.1 Discovering the Bell Curve

EXPERIMENT From a histogram to a curve

Imagine weighing 1000 loaves of bread sold as “500 g loaves”. Most loaves weigh close to 500 g; a few weigh 490 g or 510 g; extremely few weigh 470 g or 530 g. If we draw a histogram with very narrow classes and then smooth its outline, we obtain a continuous curve with three striking features:

- it is symmetric about the mean μ ;
- it has a single peak (mode) at μ ;
- its two tails approach the horizontal axis but never touch it.

This is exactly the shape of the Normal density curve.

DEFINITION Normal distribution

A continuous random variable X follows a **Normal distribution** with mean μ and variance σ^2 , written $X \sim N(\mu, \sigma^2)$, when its probability density function is

$$f(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}, \quad x \in \mathbb{R}.$$

The curve of f is called the Normal curve or bell curve. It satisfies:

- symmetry about the vertical line $x = \mu$;
- a maximum at $x = \mu$, of value $\frac{1}{\sigma\sqrt{2\pi}}$;
- points of inflexion at $x = \mu - \sigma$ and $x = \mu + \sigma$;
- $f(x) > 0$ for all x , and $f(x) \rightarrow 0$ as $x \rightarrow \pm\infty$ (the curve never touches the axis).


WARNING Recognition only

At the GCE Advanced Level you are never asked to integrate this density: the exponential of $-x^2$ has no elementary antiderivative. All probabilities are obtained from tables, after

standardisation. You must, however, be able to recognise the parameters: in $N(\mu, \sigma^2)$, the second parameter is the variance, not the standard deviation. Thus $X \sim N(50, 9)$ means $\mu = 50$ and $\sigma = 3$.

The two parameters play clearly separated roles, which you can see on the figure below:

- μ is a **location** parameter: changing μ translates the curve horizontally without changing its shape;
- σ is a **spread** parameter: a larger σ gives a flatter, wider curve; a smaller σ gives a tall, narrow curve.



Effect of the parameters: $N(2, 1)$ is $N(0, 1)$ shifted 2 units right (role of μ); $N(0, 4)$, with $\sigma = 2$, is flatter and wider (role of σ).

PROPERTY Fundamental characteristics of $N(\mu, \sigma^2)$

If $X \sim N(\mu, \sigma^2)$ then

$$E(X) = \mu, \quad \text{Var}(X) = \sigma^2, \quad P(X < \mu) = P(X > \mu) = 0.5,$$

and the total area under the curve equals 1. The symmetry about μ implies in particular that the mean, median and mode of a Normal distribution all coincide.

3.2 The Standard Normal Variable

There are infinitely many Normal distributions, one for each pair (μ, σ^2) . Printing tables for all of them is impossible. The brilliant solution is to **reduce every Normal variable to one single reference variable**, by a simple change of origin and scale.

DEFINITION Standard Normal distribution

The **standard Normal distribution** is $N(0, 1)$: mean 0, variance 1. A variable with this distribution is usually denoted Z , and its cumulative distribution function is denoted by the Greek letter Phi:

$$\Phi(z) = P(Z < z).$$

PROPERTY Expectation and variance of Z — proof not examinable

For $Z \sim N(0, 1)$:

$$E(Z) = 0 \quad \text{and} \quad \text{Var}(Z) = 1.$$

Illustration. The density of Z is $\varphi(z) = \frac{1}{\sqrt{2\pi}}e^{-z^2/2}$, an even function (symmetric about 0). Hence $z\varphi(z)$ is an odd function, and the integral defining $E(Z)$ over $\mathbb{R}$ is 0. A computation (integration by parts) then gives $E(Z^2) = 1$, so $\text{Var}(Z) = E(Z^2) - [E(Z)]^2 = 1 - 0 = 1$. This is exactly what “standard” means: centred (mean 0) and reduced (standard deviation 1).

PROPERTY Standardisation — examinable and essential

If $X \sim N(\mu, \sigma^2)$, then the variable

$$Z = \frac{X - \mu}{\sigma} \quad \text{follows} \quad N(0, 1).$$

Consequently, for any real numbers $a < b$:

$$P(a < X < b) = P\left(\frac{a - \mu}{\sigma} < Z < \frac{b - \mu}{\sigma}\right).$$

Justification. The transformation $Z = \frac{X - \mu}{\sigma}$ is linear, so Z remains Normally distributed. Moreover $E(Z) = \frac{E(X) - \mu}{\sigma} = 0$ and $\text{Var}(Z) = \frac{\text{Var}(X)}{\sigma^2} = \frac{\sigma^2}{\sigma^2} = 1$. Hence $Z \sim N(0, 1)$. ■

METHOD Standardising $X \sim N(\mu, \sigma^2)$

1. Identify μ and σ (remember: the second parameter of $N(\mu, \sigma^2)$ is the variance, so take its square root).
2. Convert each boundary value x into $z = \frac{x - \mu}{\sigma}$.
3. Sketch the standard Normal curve and shade the required region.
4. Read Φ from the tables and use the properties of Φ to obtain the answer.

3.3 Reading the Standard Normal Tables

The GCE tables give $\Phi(z) = P(Z < z)$ for $z \geq 0$, usually to two decimal places: the row gives z to one decimal, the column gives the second decimal. For example, for $z = 1.23$ we read the row 1.2 and the column 0.03:

z	0.00	0.01	0.02	0.03
1.1	0.8643	0.8665	0.8686	0.8708
1.2	0.8849	0.8869	0.8888	0.8907
1.3	0.9032	0.9049	0.9066	0.9082

So $\Phi(1.23) = 0.8907$. When the exam requires more precision, we **interpolate linearly** between two consecutive entries. For instance, for $z = 1.645$: $\Phi(1.64) = 0.9495$ and $\Phi(1.65) = 0.9505$, so

$$\Phi(1.645) \approx \frac{1}{2}(0.9495 + 0.9505) = 0.9500.$$

WARNING The table gives a LEFT tail

The table gives $\Phi(z) = P(Z < z)$, the area to the left of z . You can never read a right-tail probability $P(Z > z)$ directly: you must first rewrite it as $1 - \Phi(z)$. Reading $\Phi(1.5)$ as the answer to $P(Z > 1.5)$ is one of the most frequent errors at the GCE.


3.4 Properties of Φ

Because the standard Normal curve is symmetric about 0, the area to the left of $-z$ equals the area to the right of z . This single geometric fact generates all the working formulas.

PROPERTY Properties of Φ — must be known perfectly

For $Z \sim N(0, 1)$ and any real numbers $z, a < b$:

$\Phi(-z) = 1 - \Phi(z)$	(symmetry)
$P(Z > z) = 1 - \Phi(z)$	(upper tail)
$P(Z < -z) = \Phi(-z) = 1 - \Phi(z)$	
$P(a < Z < b) = \Phi(b) - \Phi(a)$	(interval)
$P(Z < z) = 2\Phi(z) - 1, \quad z \geq 0$	(central interval)



The shaded blue area is $\Phi(z) = P(Z < z)$. By symmetry, the area left of $-z$ equals the area right of z : $\Phi(-z) = 1 - \Phi(z)$.

EXAMPLE Warm-up with the tables

Let $Z \sim N(0, 1)$. Using $\Phi(1.20) = 0.8849$ and $\Phi(0.85) = 0.8023$:

- $P(Z < 1.20) = \Phi(1.20) = 0.8849$;
- $P(Z > 1.20) = 1 - 0.8849 = 0.1151$;
- $P(Z < -1.20) = \Phi(-1.20) = 1 - 0.8849 = 0.1151$;
- $P(-1.20 < Z < 1.20) = 2(0.8849) - 1 = 0.7698$;
- $P(0.85 < Z < 1.20) = \Phi(1.20) - \Phi(0.85) = 0.8849 - 0.8023 = 0.0826$.

WORKED EXAMPLE 5 Standardising a real situation

The masses of loaves produced by a bakery are Normally distributed with mean $\mu = 500$ g and standard deviation $\sigma = 12$ g. A loaf is chosen at random. Find the probability that its mass is (a) less than 482 g; (b) between 494 g and 518 g; (c) more than 527 g.

SOLUTION

Let $X \sim N(500, 12^2)$ and set $Z = \frac{X - 500}{12} \sim N(0, 1)$.

(a) $z = \frac{482 - 500}{12} = -1.5$, so

$$P(X < 482) = P(Z < -1.5) = 1 - \Phi(1.5) = 1 - 0.9332 = 0.0668.$$

(b) $z_1 = \frac{494 - 500}{12} = -0.5$ and $z_2 = \frac{518 - 500}{12} = 1.5$, so

$$\begin{aligned} P(494 < X < 518) &= \Phi(1.5) - \Phi(-0.5) = \Phi(1.5) - [1 - \Phi(0.5)] \\ &= 0.9332 - 1 + 0.6915 = 0.6247. \end{aligned}$$

(c) $z = \frac{527 - 500}{12} = 2.25$, so

$$P(X > 527) = 1 - \Phi(2.25) = 1 - 0.9878 = 0.0122.$$

About 1.2% of loaves exceed 527 g: heavy loaves are rare, exactly as the bell shape predicts.

3.5 Using the Tables in Reverse: Percentage Points

Very often the question is inverted: we know the probability and we look for the value of z . We then read the table **inside-out**: locate the probability in the body of the table and read off the corresponding z .

DEFINITION Percentage point

For $0 < p < 1$, the value z_p such that $\Phi(z_p) = p$ is called a **percentage point** (or quantile) of the standard Normal distribution. By symmetry, $\Phi(z) = p$ with $p < 0.5$ gives $z = -z_{1-p} < 0$.

EXAMPLE Reverse reading

Find z such that $\Phi(z) = 0.9750$. Searching the body of the table, 0.9750 corresponds to $z = 1.96$. Hence $P(Z < 1.96) = 0.975$ — the famous value behind the central 95% interval. Similarly, $\Phi(z) = 0.05$ gives $z = -1.645$, since $\Phi(1.645) = 0.95$.

The following percentage points occur so often at the GCE that they are worth memorising:

p	0.90	0.95	0.975	0.995
z with $\Phi(z) = p$	1.282	1.645	1.960	2.576

METHOD Exam tip: sketch before you search

Before touching the tables, always draw a quick bell curve, mark 0 (or μ), place the boundary value(s), and shade the required area. The sketch tells you immediately whether your answer should be greater or smaller than 0.5, whether you need $\Phi(z)$, $1 - \Phi(z)$ or a difference, and whether a reverse reading should give a positive or negative z . A ten-second sketch eliminates most table errors.

WARNING Common mistakes with Φ

- Reading $P(Z > z)$ directly from the table instead of computing $1 - \Phi(z)$.
- Forgetting that the table only covers $z \geq 0$: for negative values use $\Phi(-z) = 1 - \Phi(z)$.

- Writing $P(a < Z < b) = \Phi(a) - \Phi(b)$ instead of $\Phi(b) - \Phi(a)$ (the result would be negative!).
- Confusing σ and σ^2 when standardising: divide by σ , the standard deviation.
- Believing the curve touches the axis: probabilities such as $P(Z > 4)$ are tiny but never exactly 0.

EXERCISE 4 Practice

Let $Z \sim N(0, 1)$. Using the tables, find:

1. $P(Z < 0.75)$, $P(Z > 1.34)$, $P(Z < -2.05)$;
2. $P(-1.28 < Z < 2.10)$ and $P(|Z| < 1.96)$;
3. the value of z such that $\Phi(z) = 0.9918$;
4. the value of z such that $P(Z > z) = 0.025$.

SOLUTION Exercise 1

1. $P(Z < 0.75) = \Phi(0.75) = 0.7734$; $P(Z > 1.34) = 1 - \Phi(1.34) = 1 - 0.9099 = 0.0901$;
 $P(Z < -2.05) = 1 - \Phi(2.05) = 1 - 0.9798 = 0.0202$.
2. $P(-1.28 < Z < 2.10) = \Phi(2.10) - [1 - \Phi(1.28)] = 0.9821 - 1 + 0.8997 = 0.8818$; $P(|Z| < 1.96) = 2\Phi(1.96) - 1 = 2(0.9750) - 1 = 0.9500$.
3. $\Phi(z) = 0.9918$ corresponds to $z = 2.40$.
4. $P(Z > z) = 0.025 \iff \Phi(z) = 0.975$, hence $z = 1.96$.

KEY POINTS Key points of this section

- $X \sim N(\mu, \sigma^2)$: bell curve, symmetric about $x = \mu$, inflexion points at $\mu \pm \sigma$, total area 1, $P(X < \mu) = 0.5$.
- Standardisation: $Z = \frac{X - \mu}{\sigma} \sim N(0, 1)$, with $E(Z) = 0$ and $\text{Var}(Z) = 1$.
- The tables give the left tail $\Phi(z) = P(Z < z)$ for $z \geq 0$.
- Master formulas: $\Phi(-z) = 1 - \Phi(z)$; $P(Z > z) = 1 - \Phi(z)$; $P(a < Z < b) = \Phi(b) - \Phi(a)$; $P(|Z| < z) = 2\Phi(z) - 1$.

- Reverse reading: find the probability inside the table, read z on the border. Key values: $\Phi(1.645) = 0.95$, $\Phi(1.96) = 0.975$, $\Phi(2.576) = 0.995$.
- Always sketch and shade before using the tables.

DID YOU KNOW? Why “Normal”?

The distribution is also called the Gaussian distribution, after Carl Friedrich Gauss, who used it to model errors in astronomical measurements. Its ubiquity is explained by the Central Limit Theorem: the sum (or average) of many small independent random effects is approximately Normal, whatever the original distributions. That is why heights, measurement errors, examination marks and so many natural phenomena all follow the same bell curve — and why the Normal distribution will reappear throughout the rest of this chapter, from central intervals to approximations of the binomial and Poisson distributions.

4 CENTRAL INTERVALS AND WORKING WITH ANY NORMAL DISTRIBUTION

In the previous section we learnt how to standardise: any Normal variable $X \sim N(\mu, \sigma^2)$ is transformed into the standard Normal variable $Z = \frac{X - \mu}{\sigma} \sim N(0, 1)$, whose probabilities are read from the tables of $\Phi(z) = P(Z < z)$. We now put this tool to work in the two directions that the GCE examiner loves: **forwards** (find the probability of an event concerning X) and **backwards** (given a probability, find the corresponding value of X). Along the way we shall meet the famous **central 95%** and **central 99%** intervals, which every candidate must know by heart.

4.1 The Central 95% and 99% Intervals

Intuition. We already know that a Normal variable clusters around its mean μ : values far from μ are possible but rare. A natural question is: within what interval, symmetric about μ , does the variable fall with a given high probability? For example, a bottling company in Douala may wish to state: “95% of our bottles contain between a and b centilitres.” Because the Normal curve is symmetric, the fairest interval is the central one: we leave the same probability in each tail.

EXPERIMENT Finding the central 95% point

We want the number z such that $P(-z < Z < z) = 0.95$. Since the total area under the curve is 1, the two tails together carry 0.05, and by symmetry each tail carries 0.025. Hence

$$\Phi(z) = P(Z < z) = 1 - 0.025 = 0.975.$$

Reading the standard Normal tables backwards (looking for 0.975 in the body of the ta-

ble), we find $\Phi(1.96) = 0.9750$. Therefore $z = 1.96$, that is

$$P(-1.96 < Z < 1.96) = 0.95.$$

The same computation with 0.99 in the centre gives $\Phi(z) = 0.995$, and the tables give $z = 2.576$ (more precisely 2.5758).

PROPERTY Central intervals of a Normal distribution — proof examinable in outline

If $X \sim N(\mu, \sigma^2)$ and $Z = \frac{X - \mu}{\sigma}$, then:

- $P(-1.96 < Z < 1.96) = 0.95$, equivalently $P(\mu - 1.96\sigma < X < \mu + 1.96\sigma) = 0.95$;
- $P(-2.576 < Z < 2.576) = 0.99$, equivalently $P(\mu - 2.576\sigma < X < \mu + 2.576\sigma) = 0.99$;
- $P(-1.645 < Z < 1.645) = 0.90$, equivalently $P(\mu - 1.645\sigma < X < \mu + 1.645\sigma) = 0.90$.

Justification. $P(-z < Z < z) = \Phi(z) - \Phi(-z) = \Phi(z) - (1 - \Phi(z)) = 2\Phi(z) - 1$. Setting $2\Phi(z) - 1 = 0.95$ gives $\Phi(z) = 0.975$, so $z = 1.96$; setting it equal to 0.99 gives $\Phi(z) = 0.995$, so $z = 2.576$; setting it equal to 0.90 gives $\Phi(z) = 0.95$, so $z = 1.645$. The statement for X follows because $-z < Z < z \iff \mu - z\sigma < X < \mu + z\sigma$.

KEY POINTS Percentage points to memorise


Purpose	Central probability	Tail probability (each)	z-value
Central 90%	0.90	0.05	1.645
Central 95%	0.95	0.025	1.96
Central 99%	0.99	0.005	2.576
One-sided 5% point	$P(Z < z) = 0.95$	0.05 (upper)	1.645
One-sided 1% point	$P(Z < z) = 0.99$	0.01 (upper)	2.326

Notice that 1.645 appears twice: it is the central 90% point and the one-sided 5% point. The context decides which role it plays.

WARNING Central 95% leaves 2.5% in EACH tail

A very common error is to use $z = 1.645$ for a “95% interval”. That is wrong: 1.645 is the value for which $P(Z < 1.645) = 0.95$, i.e. it leaves the whole 5% in the upper tail only. A

central 95% interval must leave 2.5% in each tail, so $\Phi(z) = 0.975$ and $z = 1.96$. Always ask yourself: central (two tails) or one-sided (one tail)?



The standard Normal curve: the central 95% region lies between $z = -1.96$ and $z = 1.96$, leaving 2.5% in each tail.

DID YOU KNOW? The 1.96 legacy

The value 1.96 became iconic largely through the work of Sir Ronald Fisher (1925), who recommended the 5% significance level for hypothesis testing. In Cameroon's quality control laboratories – from the CDC tea plantations in Tiko to the SONARA refinery in Limbe – the “ $\mu \pm 1.96\sigma$ ” interval remains the standard benchmark for setting tolerance limits on product specifications.

4.2 Using the Standard Normal Tables for Any Normal Distribution

METHOD The four-step procedure for probabilities with $X \sim N(\mu, \sigma^2)$

1. **Standardise.** Replace X by $Z = \frac{X - \mu}{\sigma}$ and rewrite the event in terms of Z .
2. **Sketch.** Draw a quick Normal curve, mark 0 and the z -value(s), and shade the required region. The sketch tells you which combination of Φ values you need.
3. **Read.** Use the tables of Φ , remembering $\Phi(-z) = 1 - \Phi(z)$ for negative values.
4. **Combine.** Use $P(Z < z) = \Phi(z)$, $P(Z > z) = 1 - \Phi(z)$, $P(a < Z < b) = \Phi(b) - \Phi(a)$, and give the answer to 4 decimal places or 3 significant figures.

WORKED EXAMPLE 6 Masses of bags of cement

The mass X of a bag of cement produced by a factory in Figuil is Normally distributed with mean 50 kg and standard deviation 0.8 kg. A bag is chosen at random. Find the probability that its mass is (a) less than 49.2 kg; (b) more than 51.1 kg; (c) between 49.2 kg and 51.1 kg; (d) within one interval containing the central 95% of all masses.

SOLUTION

We have $X \sim N(50, 0.8^2)$, so $Z = \frac{X - 50}{0.8}$.

(a) Standardise: $z = \frac{49.2 - 50}{0.8} = -1.00$. Hence

$$P(X < 49.2) = P(Z < -1) = \Phi(-1) = 1 - \Phi(1) = 1 - 0.8413 = 0.1587.$$

(b) $z = \frac{51.1 - 50}{0.8} = 1.375 \approx 1.38$. Hence

$$P(X > 51.1) = 1 - \Phi(1.375) \approx 1 - 0.9154 = 0.0846.$$

(Using $z = 1.38$: $1 - 0.9162 = 0.0838$; both are accepted if the rounding is stated.)

(c) Using the two previous results,

$$P(49.2 < X < 51.1) = \Phi(1.375) - \Phi(-1) = 0.9154 - 0.1587 = 0.7567.$$

(d) The central 95% interval is $\mu \pm 1.96\sigma$:

$$50 \pm 1.96 \times 0.8 = 50 \pm 1.568,$$

i.e. between 48.43 kg and 51.57 kg. So 95% of the bags have a mass between about 48.4 kg and 51.6 kg.


4.3 De-standardising: From a Probability Back to a Value of X

So far the value x was given and we found the probability. The inverse problem is just as important at the GCE: **the probability is given, and we must find the value of x** . The key is to read the tables backwards: locate the probability p in the body of the Φ table and read off the corresponding z .

METHOD De-standardising — from $\Phi(z) = p$ to $x = \mu + z\sigma$

1. Translate. Write the given statement in the form $P(X < x) = p$ (if you are given $P(X > x) = p$, rewrite it as $P(X < x) = 1 - p$).
2. Invert the tables. Find z such that $\Phi(z) = p$. If $p < 0.5$, then z is negative: find z' with $\Phi(z') = 1 - p$ and take $z = -z'$.

3. De-standardise. Compute $x = \mu + z\sigma$.
4. Check. If $p > 0.5$ you must get $x > \mu$; if $p < 0.5$ you must get $x < \mu$. This one-second check catches most sign errors.



Standardisation and de-standardisation: the map $z = \frac{x - \mu}{\sigma}$ sends the X -scale to the Z -scale; the inverse map $x = \mu + z\sigma$ brings a z -value back to the original units.

WORKED EXAMPLE 7 Fill volumes of bottles

A machine at a brewery in Yaoundé fills bottles so that the volume V delivered is Normally distributed with mean 33.0 cl and standard deviation 0.4 cl. (a) Find the volume v such that 90% of bottles contain more than v . (b) Find the volume w exceeded by only 1% of bottles.

SOLUTION

(a) We want v with $P(V > v) = 0.90$, i.e. $P(V < v) = 0.10$. Since $0.10 < 0.5$, the z -value is negative. From the tables, $\Phi(1.282) = 0.90$, so $z = -1.282$. De-standardising:

$$v = \mu + z\sigma = 33.0 + (-1.282)(0.4) = 33.0 - 0.5128 \approx 32.49 \text{ cl.}$$

Check: $p = 0.10 < 0.5$ and indeed $v < \mu = 33.0$. ✓

(b) We want w with $P(V > w) = 0.01$, i.e. $\Phi(z) = 0.99$, giving $z = 2.326$. Hence

$$w = 33.0 + (2.326)(0.4) = 33.0 + 0.9304 \approx 33.93 \text{ cl.}$$

Only 1% of bottles receive more than about 33.93 cl.

4.4 Inverse Problems in Terms of μ and σ

Sometimes the mean and standard deviation are not substituted immediately: the answer is required in terms of μ and σ , or the information is used to set up an equation (this prepares the

next section, where μ or σ is unknown).

EXAMPLE General inverse statement

If $X \sim N(\mu, \sigma^2)$ and $P(X < x) = 0.975$, then $\Phi(z) = 0.975$ gives $z = 1.96$, so

$$x = \mu + 1.96\sigma.$$

Similarly $P(X < x) = 0.05$ gives $z = -1.645$, so $x = \mu - 1.645\sigma$; and $P(X > x) = 0.005$ gives $\Phi(z) = 0.995$, $z = 2.576$, so $x = \mu + 2.576\sigma$.

WORKED EXAMPLE 8 Marks of candidates — GCE type

The marks X obtained by candidates in a mock Advanced Level examination are Normally distributed with mean 52 and standard deviation 14. (a) Find the central interval within which 99% of the marks lie. (b) The best 10% of candidates are awarded grade A. Find the least mark required for grade A. (c) Find the mark m such that $P(X < m) = 0.20$, and interpret it.

SOLUTION

(a) Central 99%: $\mu \pm 2.576\sigma = 52 \pm 2.576 \times 14 = 52 \pm 36.06$, i.e. approximately from 15.9 to 88.1 marks.

(b) We need a with $P(X > a) = 0.10$, i.e. $P(X < a) = 0.90$, so $z = 1.282$ and

$$a = 52 + 1.282 \times 14 = 52 + 17.95 \approx 69.9.$$

The least mark for grade A is therefore about 70.

(c) $\Phi(z) = 0.20$ with $0.20 < 0.5$: $\Phi(0.842) = 0.80$, so $z = -0.842$ and

$$m = 52 - 0.842 \times 14 = 52 - 11.79 \approx 40.2.$$

Interpretation: 20% of the candidates scored less than about 40 marks.

WARNING Exam technique: precision of z -values

GCE mark schemes expect you to quote z -values to at least 3 significant figures: write 1.96, 2.576, 1.645, 2.326, and not 1.9 or 2.6. When inverting the tables, choose the tabulated probability closest to the required one (e.g. use $\Phi(1.96) = 0.9750$ for 0.975). Premature rounding of z before computing $x = \mu + z\sigma$ is a frequent cause of lost accuracy marks.

WARNING Signs and direction of the inequality

Before de-standardising, always rewrite the statement as $P(X < x) = p$ with $p = \Phi(z)$. If you are given $P(X > x) = p$, do not look up p directly: look up $1 - p$. And remember that a probability below 0.5 on the left means a negative z and a value of x below the mean.

EXERCISE 5 Consolidation

The heights of students in an Upper Sixth class in Bamenda are Normally distributed with mean 168 cm and standard deviation 7 cm.

1. Find the probability that a randomly chosen student is taller than 180 cm.
2. Find the central interval containing the heights of 95% of the students.
3. Find the height h such that only 5% of the students are taller than h .
4. Find the height k such that $P(X < k) = 0.75$.

SOLUTION Exercise 1

1. $z = \frac{180 - 168}{7} = 1.714 \approx 1.71$; $P(X > 180) = 1 - \Phi(1.714) \approx 1 - 0.9568 = 0.0432$.
2. $\mu \pm 1.96\sigma = 168 \pm 13.72$, i.e. from 154.3 cm to 181.7 cm.
3. $P(X > h) = 0.05 \Rightarrow \Phi(z) = 0.95 \Rightarrow z = 1.645$; $h = 168 + 1.645 \times 7 = 179.5$ cm.
4. $\Phi(z) = 0.75 \Rightarrow z = 0.674$; $k = 168 + 0.674 \times 7 = 172.7$ cm.

KEY POINTS What you must retain from this section

- Central intervals: 90% $\rightarrow 1.645$, 95% $\rightarrow 1.96$, 99% $\rightarrow 2.576$; for X these become $\mu \pm z\sigma$.
- Forwards: standardise, sketch, read Φ , combine.
- Backwards (de-standardising): write $P(X < x) = p$, invert the tables to get z , then $x = \mu + z\sigma$; check the position of x relative to μ .
- Central probability c means $(1 - c)/2$ in each tail: never confuse the central 95% point 1.96 with the one-sided 5% point 1.645.

These inverse skills — moving freely between a probability and a value of X — are exactly what we shall need in the next section, where the unknown is no longer x but the parameter μ or σ itself.

5 DETERMINING UNKNOWN μ AND/OR σ

In the previous section we learnt to use a Normal distribution whose parameters μ and σ were given: we standardised with $Z = \frac{X - \mu}{\sigma}$, read the tables, and de-standardised to answer questions about X . In real life, however, the mean and the standard deviation are rarely handed to us on a plate. A quality-control engineer in a soap factory in Douala may know that “10% of tablets weigh more than 62 g” and “5% weigh less than 55 g”, and from these two pieces of information she must reconstruct μ and σ . This section is exactly about that reverse problem: we are given probability statements and we must find the unknown parameter(s). The key observation is simple: each probability statement, once converted into a z -value by reading the tables backwards, produces one linear equation in μ and σ . One statement, one equation, one unknown; two statements, two equations, two unknowns.

5.1 The General Principle: From a Probability Statement to an Equation

Suppose $X \sim N(\mu, \sigma^2)$. Any statement of the form

$$P(X < a) = p \quad \text{or} \quad P(X > a) = p$$

can be standardised. For instance, $P(X < a) = p$ becomes

$$P\left(Z < \frac{a - \mu}{\sigma}\right) = p, \quad \text{so} \quad \Phi\left(\frac{a - \mu}{\sigma}\right) = p.$$

Reading the standard normal table in reverse (looking for the probability p in the body of the table and reading off the corresponding z) gives a number z_0 such that $\Phi(z_0) = p$, and hence the **linear equation**

$$\frac{a - \mu}{\sigma} = z_0 \quad \Longleftrightarrow \quad a = \mu + z_0 \sigma.$$

For a “greater than” statement we first convert: $P(X > a) = p$ gives $P(X < a) = 1 - p$, so $\Phi\left(\frac{a - \mu}{\sigma}\right) = 1 - p$.


WARNING The sign of z

A probability greater than 0.5 corresponds to a positive z -value; a probability less than 0.5 corresponds to a negative z -value, and the tables only tabulate $\Phi(z)$ for $z \geq 0$. If $p < 0.5$, find z_1 with $\Phi(z_1) = 1 - p$ and take $z = -z_1$ (symmetry: $\Phi(-z_1) = 1 - \Phi(z_1)$). Getting this sign wrong is the single most common error in this topic. Always ask yourself: is the boundary a above or below the mean?

METHOD Setting up equations for unknown μ and/or σ

1. Write the model: $X \sim N(\mu, \sigma^2)$, stating what is known and what is unknown.
2. Translate each worded probability statement into $P(X < a) = p$ or $P(X > a) = p$.

3. Standardise: $\Phi\left(\frac{a - \mu}{\sigma}\right) = p$ (use $1 - p$ for a “greater than” statement).
4. Read the table backwards to find z_0 with $\Phi(z_0) = p$; fix the sign using the warning above.
5. Write the equation in the form $\frac{a - \mu}{\sigma} = z_0$, i.e. $a = \mu + z_0\sigma$.
6. Solve for the unknown(s); check that $\sigma > 0$ and that the answers are sensible in context.



Flow-chart of the method: from a probability statement to the values of μ and σ .

5.2 Type 1: μ Known, σ Unknown

Here one probability statement provides one equation in the single unknown σ .

WORKED EXAMPLE 9 Type 1 — Finding σ

The masses of loaves produced by a bakery are Normally distributed with mean 500 g and standard deviation σ g. Given that 20% of loaves weigh more than 510 g, find σ .

SOLUTION

Let $X \sim N(500, \sigma^2)$. The statement “20% weigh more than 510 g” gives

$$P(X > 510) = 0.20 \implies P(X < 510) = 0.80 \implies \Phi\left(\frac{510 - 500}{\sigma}\right) = 0.80.$$

Reading the table backwards, $\Phi(z) = 0.80$ gives $z = 0.8416$. Since $0.80 > 0.5$, z is positive (and indeed $510 > 500$). Hence

$$\frac{10}{\sigma} = 0.8416 \implies \sigma = \frac{10}{0.8416} \approx 11.9 \text{ g}.$$

Check: $\sigma > 0$ and of plausible size for a spread of loaf masses. $\sigma \approx 11.9 \text{ g}$

5.3 Type 2: σ Known, μ Unknown

Symmetrically, one probability statement gives one equation in μ .

WORKED EXAMPLE 10 Type 2 — Finding μ

A machine fills sugar packets so that the mass delivered is Normally distributed with standard deviation 4 g and unknown mean μ g. Regulations require that only 2.5% of packets contain less than 500 g. Find the value at which μ must be set.

SOLUTION

Let $X \sim N(\mu, 16)$. We need $P(X < 500) = 0.025$, so

$$\Phi\left(\frac{500 - \mu}{4}\right) = 0.025.$$

Since $0.025 < 0.5$, the z -value is negative. From the tables, $\Phi(1.96) = 0.975$, so $\Phi(-1.96) = 0.025$, giving

$$\frac{500 - \mu}{4} = -1.96 \implies 500 - \mu = -7.84 \implies \mu = 507.84 \text{ g}.$$

Check: the mean must exceed 500 g (so that few packets fall below 500 g), which it does. $\mu \approx 507.8 \text{ g}$

WARNING Do not confuse the two directions

In Type 2 problems the unknown μ often lies on the opposite side of the boundary from what students guess. If only a small percentage may fall below a limit, the mean must be

above that limit. Sketch a quick curve before solving — it will save you from sign errors.

5.4 Type 3: Both μ and σ Unknown


With two unknowns we need two probability statements, giving two equations

$$a_1 = \mu + z_1\sigma, \quad a_2 = \mu + z_2\sigma,$$

which we solve simultaneously. The cleanest technique is **elimination**: subtracting the two equations eliminates μ and yields σ immediately; substitution then gives μ .

METHOD Solving the two simultaneous equations

1. Write both equations in the standard form $a_i = \mu + z_i\sigma$ before any algebra.
2. Subtract: $a_2 - a_1 = (z_2 - z_1)\sigma$, so $\sigma = \frac{a_2 - a_1}{z_2 - z_1}$.
3. Substitute σ into either equation to find $\mu = a_1 - z_1\sigma$.
4. Verify: σ must be positive; substitute both values back into the other equation as a check.



Two tail statements give two equations: $55 = \mu - 1.12\sigma$ and $62 = \mu + 0.8416\sigma$.

WORKED EXAMPLE 11 Type 3 — Finding both μ and σ

The masses of tablets produced by a machine are Normally distributed with mean μ g and standard deviation σ g. It is found that 13.14% of tablets weigh less than 55 g and 20%

weigh more than 62 g. Find μ and σ , and hence the percentage of tablets weighing between 56 g and 63 g.

SOLUTION

Step 1: the two equations. From $P(X < 55) = 0.1314$: since $0.1314 < 0.5$, z is negative; $\Phi(1.12) = 0.8686$, so $z_1 = -1.12$:

$$\frac{55 - \mu}{\sigma} = -1.12 \implies 55 = \mu - 1.12\sigma. \quad (1)$$

From $P(X > 62) = 0.20$: $P(X < 62) = 0.80$, $\Phi(0.8416) = 0.80$, $z_2 = +0.8416$:

$$\frac{62 - \mu}{\sigma} = 0.8416 \implies 62 = \mu + 0.8416\sigma. \quad (2)$$

Step 2: elimination. Subtracting (1) from (2):

$$7 = (0.8416 + 1.12)\sigma = 1.9616\sigma \implies \sigma = \frac{7}{1.9616} \approx 3.57 \text{ g.}$$

Substituting into (2): $\mu = 62 - 0.8416(3.57) \approx 62 - 3.00 = 59.0 \text{ g}$. Check in (1): $59.0 - 1.12(3.57) = 59.0 - 4.00 = 55.0$. ✓ Also $\sigma > 0$. ✓ Step 3: further probability with the fitted model. Now $X \sim N(59, 3.57^2)$:

$$P(56 < X < 63) = \Phi\left(\frac{63 - 59}{3.57}\right) - \Phi\left(\frac{56 - 59}{3.57}\right) = \Phi(1.12) - \Phi(-0.84).$$

Using the tables: $\Phi(1.12) = 0.8686$ and $\Phi(-0.84) = 1 - \Phi(0.84) = 1 - 0.7995 = 0.2005$. Hence

$$P(56 < X < 63) = 0.8686 - 0.2005 = 0.6681.$$

About 66.8% of tablets weigh between 56 g and 63 g.

WARNING Exam technique

State each equation explicitly in the form $\frac{x - \mu}{\sigma} = z$ before clearing the fraction. Examiners award method marks for this line, and it prevents the classic slip of writing $x = \mu\sigma + z$ or inverting the sign of z . Quote z -values to 4 decimal places where the tables allow, and keep full precision until the final rounding.

5.5 GCE-Style Mixed Problems

WORKED EXAMPLE 12 Exam marks context

In a GCE mock examination, the marks obtained by a large group of candidates may be modelled by $X \sim N(\mu, \sigma^2)$. It is known that 15% of candidates scored more than 72 marks and that 10% scored less than 38 marks.

1. Find μ and σ .
2. The pass mark is 50. Estimate the percentage of candidates who passed.

SOLUTION

(a) $P(X > 72) = 0.15 \Rightarrow P(X < 72) = 0.85$, and $\Phi(1.0364) = 0.85$, so

$$72 = \mu + 1.0364\sigma. \quad (1)$$

$P(X < 38) = 0.10$: negative z ; $\Phi(1.2816) = 0.90$, so $z = -1.2816$:

$$38 = \mu - 1.2816\sigma. \quad (2)$$

Subtracting (2) from (1): $34 = 2.318\sigma$, so $\sigma \approx 14.67$. Then $\mu = 72 - 1.0364(14.67) \approx 72 - 15.20 = 56.8$. Check in (2): $56.8 - 1.2816(14.67) \approx 56.8 - 18.8 = 38.0$. ✓ (b) $P(X > 50) = P\left(Z > \frac{50 - 56.8}{14.67}\right) = P(Z > -0.4635) = \Phi(0.4635) \approx 0.6785$. About 67.9% of candidates passed.

WORKED EXAMPLE 13 Crop yield context

The yield, in kg per plot, of a maize variety is Normally distributed. Experiments show that 5% of plots yield less than 18 kg and 25% yield more than 30 kg. Find the mean and standard deviation of the yield.

SOLUTION

$P(X < 18) = 0.05$: $\Phi(1.6449) = 0.95$, so $z_1 = -1.6449$:

$$18 = \mu - 1.6449\sigma. \quad (1)$$

$P(X > 30) = 0.25 \Rightarrow P(X < 30) = 0.75$, $\Phi(0.6745) = 0.75$:

$$30 = \mu + 0.6745\sigma. \quad (2)$$

Subtracting: $12 = 2.3194\sigma$, so $\sigma \approx 5.17$ kg, and $\mu = 30 - 0.6745(5.17) \approx 26.5$ kg. Check in (1): $26.5 - 1.6449(5.17) \approx 26.5 - 8.5 = 18.0$. ✓ $\mu \approx 26.5$ kg, $\sigma \approx 5.17$ kg

EXERCISE 6 Practice

1. The heights of students in a school are $N(165, \sigma^2)$ cm. If 30% of students are taller than 170 cm, find σ .
2. A machine produces bolts whose diameters are $N(\mu, 0.04)$ mm (so $\sigma = 0.2$ mm). If 1% of bolts have diameter less than 24.5 mm, find μ .
3. The lifetime X (hours) of a brand of battery is $N(\mu, \sigma^2)$. Given $P(X < 40) = 0.0668$ and $P(X > 70) = 0.0228$, find μ and σ , and hence $P(50 < X < 60)$.

SOLUTION Exercise 1

1. $P(X > 170) = 0.30 \Rightarrow \Phi\left(\frac{5}{\sigma}\right) = 0.70, z = 0.5244$, so $\sigma = \frac{5}{0.5244} \approx 9.53$ cm.
2. $\Phi\left(\frac{24.5 - \mu}{0.2}\right) = 0.01, z = -2.3263$, so $\mu = 24.5 + 0.2(2.3263) \approx 24.97$ mm.
3. $\Phi(1.5) = 0.9332 \Rightarrow z_1 = -1.5: 40 = \mu - 1.5\sigma$. $\Phi(2.0) = 0.9772 \Rightarrow z_2 = 2.0: 70 = \mu + 2\sigma$. Subtracting: $30 = 3.5\sigma, \sigma \approx 8.571, \mu = 40 + 1.5(8.571) \approx 52.86$. Then $P(50 < X < 60) = \Phi(0.8333) - \Phi(-0.3333) = 0.7977 - 0.3694 \approx 0.428$.

KEY POINTS Key points

- Each probability statement $P(X < a) = p$ or $P(X > a) = p$ converts, via standardisation and a backwards table reading, into one linear equation $a = \mu + z\sigma$.
- $p > 0.5 \Rightarrow z > 0$; $p < 0.5 \Rightarrow z < 0$ (use symmetry: $\Phi(-z) = 1 - \Phi(z)$).
- One unknown requires one statement; two unknowns (μ and σ) require two statements, solved by elimination: $\sigma = \frac{a_2 - a_1}{z_2 - z_1}$, then back-substitute for μ .
- Always check $\sigma > 0$ and verify the solution in the unused equation; then the fitted distribution can be used for any further probability question.

6 CONTINUITY CORRECTION AND NORMAL APPROXIMATIONS


In the previous section we learned how to handle any normal distribution by standardising it to $Z \sim N(0, 1)$ and using the standard normal tables. We also solved problems where the mean μ

or the standard deviation σ (or both) were unknown. Now we turn to a powerful practical tool: using the normal distribution to approximate probabilities for discrete distributions — the binomial and the Poisson — when exact calculation becomes tedious or impossible without technology. The bridge between the discrete world and the continuous normal curve is the **continuity correction**.


6.1 Why a Continuity Correction is Needed

The binomial distribution $B(n, p)$ and the Poisson distribution $Po(\lambda)$ are discrete: the random variable takes only integer values $0, 1, 2, \dots$. The normal distribution is continuous: its probability density function is defined for every real number, and $\Pr(X = x) = 0$ for any single value x .

When we approximate a discrete distribution by a continuous one, we must account for the fact that a discrete probability $\Pr(X = r)$ corresponds to an area under the continuous curve. The natural way to do this is to associate the integer value r with the interval $[r - 0.5, r + 0.5]$ on the real line. This is the essence of the continuity correction.



Binomial histogram with superimposed normal curve. The bar at $r = 4$ corresponds to the area under the normal curve between 3.5 and 4.5.



Number-line strip showing how each discrete value r maps to a continuous interval of width 1 centred on r .

6.2 The Continuity Correction Rules

The following table summarises how to translate every discrete probability statement into a continuous one before standardising. **Memorise this table — it is the single most important tool in this section.**

METHOD Continuity Correction Table

For a discrete integer-valued random variable X approximated by a continuous normal variable Y :

Discrete probability	Continuous equivalent	Explanation
$\Pr(X \leq r)$	$\Pr(Y < r + 0.5)$	Include the whole bar at r
$\Pr(X < r) = \Pr(X \leq r - 1)$	$\Pr(Y < r - 0.5)$	Stop before the bar at r
$\Pr(X = r)$	$\Pr(r - 0.5 < Y < r + 0.5)$	Area of the single bar at r
$\Pr(X \geq r)$	$\Pr(Y > r - 0.5)$	Include the bar at r and all to the right
$\Pr(X > r) = \Pr(X \geq r + 1)$	$\Pr(Y > r + 0.5)$	Start after the bar at r

WARNING The Most Penalised Error

Forgetting the continuity correction is the number one mistake in GCE examinations on this topic. Even if you correctly identify the normal approximation, correctly compute μ and σ , and correctly use the tables, omitting the ± 0.5 adjustment will lose you most of the marks. Always write the corrected inequality explicitly before standardising.

6.3 Normal Approximation to the Binomial Distribution

When n is large, the binomial distribution $B(n, p)$ becomes symmetric and bell-shaped (provided p is not too close to 0 or 1). The Central Limit Theorem justifies approximating it by a normal distribution with the same mean and variance.

PROPERTY Normal Approximation to the Binomial

If $X \sim B(n, p)$ with n large, then

$$X \approx Y \sim N(\mu, \sigma^2) \quad \text{where} \quad \mu = np, \quad \sigma^2 = npq, \quad q = 1 - p.$$

Validity conditions (GCE requirement): $np > 5$ and $nq > 5$.

DID YOU KNOW? Why $np > 5$ and $nq > 5$?

These conditions ensure that the binomial distribution is sufficiently spread out and not too skewed. If p is very small, the Poisson approximation (next subsection) is often better. The threshold 5 is a conventional rule of thumb; some textbooks use 10. For GCE, always check

and state both $np > 5$ and $nq > 5$ explicitly to earn method marks.

EXAMPLE Proportion of Defective Items

A factory produces light bulbs. Historical data shows that 4% of bulbs are defective. A quality control inspector randomly selects 200 bulbs. Find the probability that at most 10 bulbs are defective.

Solution. Let X be the number of defective bulbs in the sample. Then $X \sim B(200, 0.04)$. Check conditions: $np = 200 \times 0.04 = 8 > 5$, $nq = 200 \times 0.96 = 192 > 5$. ✓ Normal approximation is valid. Approximate by $Y \sim N(\mu, \sigma^2)$ with

$$\mu = np = 8, \quad \sigma^2 = npq = 200 \times 0.04 \times 0.96 = 7.68, \quad \sigma = \sqrt{7.68} \approx 2.771.$$

We need $\Pr(X \leq 10)$. Apply continuity correction: $\Pr(X \leq 10) \approx \Pr(Y < 10.5)$. Standardise:

$$z = \frac{10.5 - 8}{\sqrt{7.68}} = \frac{2.5}{2.771} \approx 0.902.$$

Using standard normal tables: $\Phi(0.902) \approx 0.8164$.

$$\Pr(X \leq 10) \approx 0.8164$$

EXAMPLE Number of Voters in a Sample

In a large constituency, 55% of voters support Candidate A. A random sample of 100 voters is taken. Use a normal approximation to find the probability that more than 60 voters support Candidate A.

Solution. $X \sim B(100, 0.55)$. Check: $np = 55 > 5$, $nq = 45 > 5$. ✓

$Y \sim N(55, 100 \times 0.55 \times 0.45 = 24.75)$, $\sigma = \sqrt{24.75} \approx 4.975$.

$\Pr(X > 60) = \Pr(X \geq 61)$. Continuity correction: $\Pr(Y > 60.5)$.

$$z = \frac{60.5 - 55}{\sqrt{24.75}} = \frac{5.5}{4.975} \approx 1.105.$$

$\Pr(Z > 1.105) = 1 - \Phi(1.105) \approx 1 - 0.8654 = 0.1346$.

$$\Pr(X > 60) \approx 0.1346$$

6.4 Normal Approximation to the Poisson Distribution

For large λ , the Poisson distribution $\text{Po}(\lambda)$ also becomes approximately normal. This is useful

because Poisson tables often stop at $\lambda = 10$ or 15, and exact calculation of $\Pr(X \geq k)$ for large λ involves summing many terms.

PROPERTY Normal Approximation to the Poisson

If $X \sim \text{Po}(\lambda)$ with λ large, then

$$X \approx Y \sim N(\mu, \sigma^2) \quad \text{where} \quad \mu = \lambda, \quad \sigma^2 = \lambda.$$

Guideline validity condition: $\lambda > 10$ (some texts use $\lambda > 15$; for GCE, $\lambda > 10$ is accepted).

EXAMPLE Calls to a Call Centre

A call centre receives an average of 18 calls per hour. Assuming a Poisson model, use a normal approximation to find the probability that in a given hour more than 25 calls are received.

Solution. $X \sim \text{Po}(18)$. $\lambda = 18 > 10$, so normal approximation is valid.

$$Y \sim N(18, 18), \sigma = \sqrt{18} \approx 4.243.$$

$\Pr(X > 25) = \Pr(X \geq 26)$. Continuity correction: $\Pr(Y > 25.5)$.


$$z = \frac{25.5 - 18}{\sqrt{18}} = \frac{7.5}{4.243} \approx 1.768.$$

$$\Pr(Z > 1.768) = 1 - \Phi(1.768) \approx 1 - 0.9615 = 0.0385.$$

$$\Pr(X > 25) \approx 0.0385$$

6.5 Choosing the Correct Approximation and Applying the Correction

In examination questions, you must decide which approximation to use. Here is a decision flowchart:



Decision flowchart for choosing the normal approximation.

KEY POINTS Key Steps for Every Approximation Question

1. Identify the exact distribution ($B(n, p)$ or $Po(\lambda)$) and the required probability.
2. Check and state the validity conditions ($np > 5, nq > 5$ or $\lambda > 10$).
3. Write down the approximating normal distribution $N(\mu, \sigma^2)$ with explicit μ and σ^2 .
4. Apply the continuity correction to convert the discrete inequality to a continuous one. Show this step clearly.
5. Standardise to $z = \frac{x - \mu}{\sigma}$.
6. Use the standard normal tables to find the probability.
7. Give the final answer to 3 or 4 significant figures (or as required).

WORKED EXAMPLE 14 Mixed Approximation Problem

A machine produces components of which 2% are defective. Components are packed in

boxes of 500.

1. Use a suitable approximation to find the probability that a randomly chosen box contains fewer than 5 defective components.
2. The factory produces 2000 boxes per day. Estimate the number of boxes per day that contain at least 15 defective components.

Detailed solution.

Part (a). $X \sim B(500, 0.02)$. $np = 10 > 5$, $nq = 490 > 5$. Normal approximation valid. Also $\lambda = np = 10$, so Poisson approximation $Po(10)$ is possible. Since $\lambda = 10$ is borderline for normal approximation to Poisson, and the question says “suitable approximation”, the normal approximation to the binomial is the standard route.

$Y \sim N(\mu = 10, \sigma^2 = 500 \times 0.02 \times 0.98 = 9.8)$, $\sigma = \sqrt{9.8} \approx 3.1305$.

$\Pr(X < 5) = \Pr(X \leq 4)$. Continuity correction: $\Pr(Y < 4.5)$.

$$z = \frac{4.5 - 10}{\sqrt{9.8}} = \frac{-5.5}{3.1305} \approx -1.757.$$

$\Pr(Z < -1.757) = 1 - \Phi(1.757) \approx 1 - 0.9605 = 0.0395$.

$$\Pr(X < 5) \approx 0.0395$$

Part (b). We need $\Pr(X \geq 15)$ for one box, then multiply by 2000. Continuity correction: $\Pr(X \geq 15) \approx \Pr(Y > 14.5)$.

$$z = \frac{14.5 - 10}{\sqrt{9.8}} = \frac{4.5}{3.1305} \approx 1.437.$$

$\Pr(Z > 1.437) = 1 - \Phi(1.437) \approx 1 - 0.9247 = 0.0753$.

Expected number of boxes per day = $2000 \times 0.0753 = 150.6 \approx 151$ boxes.

WORKED EXAMPLE 15

Poisson Approximation with Continuity Correction

The number of accidents per month at a certain intersection follows a Poisson distribution with mean 25. Use a normal approximation to find the probability that in a given month there are between 20 and 30 accidents inclusive.

Detailed solution. $X \sim Po(25)$. $\lambda = 25 > 10$, normal approximation valid.

$Y \sim N(25, 25)$, $\sigma = 5$.

Required: $\Pr(20 \leq X \leq 30)$. Continuity correction: $\Pr(19.5 < Y < 30.5)$.

Standardise both ends:

$$z_1 = \frac{19.5 - 25}{5} = -1.1, \quad z_2 = \frac{30.5 - 25}{5} = 1.1.$$

$\Pr(-1.1 < Z < 1.1) = \Phi(1.1) - \Phi(-1.1) = \Phi(1.1) - (1 - \Phi(1.1)) = 2\Phi(1.1) - 1$. From tables: $\Phi(1.1) = 0.8643$.
Probability = $2 \times 0.8643 - 1 = 0.7286$.

$$\Pr(20 \leq X \leq 30) \approx 0.7286$$

6.6 Estimating Binomial Probabilities That Are Tedious Exactly

A classic use of the normal approximation is to estimate tail probabilities like $\Pr(X \geq 40)$ for $X \sim B(100, 0.35)$ without summing 61 binomial terms.

EXAMPLE Large Tail Probability

$X \sim B(100, 0.35)$. Estimate $\Pr(X \geq 40)$ using a normal approximation.

Solution. $np = 35 > 5$, $nq = 65 > 5$. Valid.

$Y \sim N(35, 100 \times 0.35 \times 0.65 = 22.75)$, $\sigma = \sqrt{22.75} \approx 4.770$.

$\Pr(X \geq 40) \approx \Pr(Y > 39.5)$.

$$z = \frac{39.5 - 35}{\sqrt{22.75}} = \frac{4.5}{4.770} \approx 0.943.$$

$\Pr(Z > 0.943) = 1 - \Phi(0.943) \approx 1 - 0.8272 = 0.1728$.

$$\Pr(X \geq 40) \approx 0.1728$$

WARNING Common Traps in Approximation Questions

- Using the wrong variance: For binomial, $\sigma^2 = npq$ (not np). For Poisson, $\sigma^2 = \lambda$.
- Forgetting to take the square root when computing σ for standardisation.
- Applying continuity correction in the wrong direction: $\Pr(X \geq r) \rightarrow \Pr(Y > r - 0.5)$, not $r + 0.5$.
- Not stating the validity conditions: GCE mark schemes allocate specific marks for checking $np > 5$, $nq > 5$ or $\lambda > 10$.
- Using the normal approximation when p is very small or very large: If $p < 0.1$ and n is large, the Poisson approximation to the binomial ($B(n, p) \approx \text{Po}(np)$) may be more accurate, but the syllabus focuses on Normal approximations here.

EXPERIMENT Comparing Exact and Approximate Probabilities

Use a calculator or software to compute the exact binomial probability $\Pr(X \geq 40)$ for

$X \sim B(100, 0.35)$ and compare with the normal approximation 0.1728 found above. You will find the exact value is approximately 0.1795. The error is about 0.0067, which is typical for a normal approximation with continuity correction. Without the continuity correction ($z = \frac{40-35}{4.770} = 1.048$, probability 0.1474), the error would be much larger (0.032). This demonstrates why the continuity correction is essential.

6.7 Examination-Style Practice

EXERCISE 7 GCE Style Questions

1. The probability that a student passes a certain test is 0.7. In a random sample of 80 students, use a normal approximation to find the probability that more than 60 students pass. State the validity conditions clearly.
2. A radioactive source emits particles at an average rate of 12 per minute. Assuming a Poisson distribution, use a normal approximation to find the probability that in a given minute at most 15 particles are emitted.
3. $X \sim B(200, 0.12)$. Use a normal approximation to find $\Pr(20 \leq X \leq 30)$.
4. A factory produces metal rods. The number of defects per rod follows a Poisson distribution with mean 0.8. A batch contains 500 rods. Use a suitable approximation to find the probability that the total number of defects in the batch exceeds 420.
5. In a large population, 30% of people have blood type O. A random sample of 150 people is taken. Find, using a normal approximation, the probability that the number with blood type O is less than 40.

SOLUTION Exercise 1

$X \sim B(80, 0.7)$. $np = 56 > 5$, $nq = 24 > 5$. Valid. $Y \sim N(56, 80 \times 0.7 \times 0.3 = 16.8)$, $\sigma = \sqrt{16.8} \approx 4.099$. $\Pr(X > 60) = \Pr(X \geq 61) \approx \Pr(Y > 60.5)$. $z = \frac{60.5-56}{\sqrt{16.8}} = \frac{4.5}{4.099} \approx 1.098$. $\Pr(Z > 1.098) = 1 - \Phi(1.098) \approx 1 - 0.8641 = 0.1359$.

SOLUTION Exercise 2

$X \sim \text{Po}(12)$. $\lambda = 12 > 10$. Valid. $Y \sim N(12, 12)$, $\sigma = \sqrt{12} \approx 3.464$. $\Pr(X \leq 15) \approx \Pr(Y < 15.5)$. $z = \frac{15.5-12}{\sqrt{12}} = \frac{3.5}{3.464} \approx 1.010$. $\Phi(1.010) \approx 0.8438$.

SOLUTION Exercise 3

$X \sim B(200, 0.12)$. $np = 24 > 5$, $nq = 176 > 5$. Valid. $Y \sim N(24, 200 \times 0.12 \times 0.88 = 21.12)$, $\sigma = \sqrt{21.12} \approx 4.596$. $\Pr(20 \leq X \leq 30) \approx \Pr(19.5 < Y < 30.5)$. $z_1 = \frac{19.5-24}{\sqrt{21.12}} = \frac{-4.5}{4.596} \approx -0.979$. $z_2 = \frac{30.5-24}{\sqrt{21.12}} = \frac{6.5}{4.596} \approx 1.414$. $\Pr(-0.979 < Z < 1.414) = \Phi(1.414) - \Phi(-0.979) = \Phi(1.414) - (1 - \Phi(0.979))$. $\Phi(1.414) \approx 0.9214$, $\Phi(0.979) \approx 0.8363$. Probability $\approx 0.9214 - 0.1637 = 0.7577$.

SOLUTION Exercise 4

Total defects in 500 rods: $X \sim \text{Po}(500 \times 0.8 = 400)$. $\lambda = 400 > 10$. $Y \sim N(400, 400)$, $\sigma = 20$. $\Pr(X > 420) \approx \Pr(Y > 420.5)$. $z = \frac{420.5-400}{20} = 1.025$. $\Pr(Z > 1.025) = 1 - \Phi(1.025) \approx 1 - 0.8473 = 0.1527$.

SOLUTION Exercise 5

$X \sim B(150, 0.3)$. $np = 45 > 5$, $nq = 105 > 5$. Valid. $Y \sim N(45, 150 \times 0.3 \times 0.7 = 31.5)$, $\sigma = \sqrt{31.5} \approx 5.612$. $\Pr(X < 40) = \Pr(X \leq 39) \approx \Pr(Y < 39.5)$. $z = \frac{39.5-45}{\sqrt{31.5}} = \frac{-5.5}{5.612} \approx -0.980$. $\Pr(Z < -0.980) = 1 - \Phi(0.980) \approx 1 - 0.8365 = 0.1635$.

DID YOU KNOW? Historical Note: De Moivre and the Normal Approximation

Abraham de Moivre (1667–1754), a French mathematician who fled to England, discovered the normal approximation to the binomial in 1733 while studying gambling problems. He showed that for large n , the binomial probabilities could be approximated by the curve $y = \frac{1}{\sqrt{2\pi npq}} e^{-\frac{(x-np)^2}{2npq}}$. This was the first appearance of the normal distribution, long before Gauss and Laplace. De Moivre's work laid the foundation for the Central Limit Theorem.

KEY POINTS Summary of Section 6

- Continuity correction maps discrete integer r to continuous interval $[r - 0.5, r + 0.5]$.
- Binomial approximation: $B(n, p) \approx N(np, npq)$ if $np > 5$ and $nq > 5$.
- Poisson approximation: $\text{Po}(\lambda) \approx N(\lambda, \lambda)$ if $\lambda > 10$.
- Always: State conditions $\rightarrow$ Write approximating $N(\mu, \sigma^2) \rightarrow$ Apply continuity correction $\rightarrow$ Standardise $\rightarrow$ Use tables.
- Most common error: Forgetting the ± 0.5 adjustment.

7 COMBINATIONS OF INDEPENDENT NORMAL VARIABLES

In the previous section we mastered the art of standardising any normal variable $X \sim N(\mu, \sigma^2)$ to $Z = \frac{X-\mu}{\sigma} \sim N(0, 1)$ and using the standard normal tables (or the inverse process) to find probabilities and unknown parameters. We also learned how the normal distribution can approximate the binomial and Poisson distributions when a continuity correction is applied.

Now we take a decisive step forward: **what happens when we combine two or more independent normal variables?** This question lies at the heart of many real-world problems — the total weight of a crate and its contents, the difference in performance between two athletes, the combined error of two independent measurements. The beautiful property of the normal family is that any linear combination of independent normal variables is itself normally distributed. This section gives you the rigorous tools to handle such combinations, with a special emphasis on the critical distinction between scaling a single observation (nX) and summing n independent observations ($X_1 + \dots + X_n$) — a distinction that examiners love to test.

7.1 Sum and Difference of Two Independent Normal Variables

Let us start with the fundamental theorem. Its proof uses moment generating functions (MGFs) or characteristic functions, which are beyond the GCE syllabus, but the statement and its application are examinable and must be known perfectly.

PROPERTY Sum and Difference of Two Independent Normal Variables

Let $X \sim N(\mu_1, \sigma_1^2)$ and $Y \sim N(\mu_2, \sigma_2^2)$ be independent random variables. Then:

1. Sum: $X + Y \sim N(\mu_1 + \mu_2, \sigma_1^2 + \sigma_2^2)$.
2. Difference: $X - Y \sim N(\mu_1 - \mu_2, \sigma_1^2 + \sigma_2^2)$.

In words: means add (or subtract) as expected, but variances always add — even for a difference.

WARNING Variances ADD for a Difference — Never Subtract!

A very common error is to write $\text{Var}(X - Y) = \sigma_1^2 - \sigma_2^2$. This is wrong. Recall the general rule for independent variables: $\text{Var}(aX + bY) = a^2\sigma_1^2 + b^2\sigma_2^2$. For $X - Y$ we have $a = 1$, $b = -1$, so the variances add: $(1)^2\sigma_1^2 + (-1)^2\sigma_2^2 = \sigma_1^2 + \sigma_2^2$. Variance measures spread; subtracting two independent variables increases uncertainty, it does not cancel it.

EXAMPLE Total Mass of a Bag and Its Contents

A bag of rice has mass $R \sim N(5.02, 0.04^2)$ kg. The empty sack has mass $S \sim N(0.15, 0.01^2)$

kg. R and S are independent. Find the probability that the total mass of a filled sack exceeds 5.20 kg.

Solution: Let $T = R + S$ be the total mass. By the theorem:

$$T \sim N(\mu_T, \sigma_T^2), \quad \mu_T = 5.02 + 0.15 = 5.17, \quad \sigma_T^2 = 0.04^2 + 0.01^2 = 0.0016 + 0.0001 = 0.0017.$$

So $\sigma_T = \sqrt{0.0017} \approx 0.04123$ kg. We need $P(T > 5.20)$. Standardise:

$$z = \frac{5.20 - 5.17}{0.04123} = \frac{0.03}{0.04123} \approx 0.7276.$$

Using tables: $P(Z > 0.7276) = 1 - \Phi(0.7276) \approx 1 - 0.7664 = 0.2336$. Answer: ≈ 0.234 (or 23.4%).

EXAMPLE Difference Between Two Athletes' Times

Two sprinters, A and B, run the 100 m. Their times (in seconds) are modelled as independent normal variables: $A \sim N(10.2, 0.15^2)$ and $B \sim N(10.5, 0.12^2)$. Find the probability that A beats B by more than 0.2 seconds.

Solution: We want $P(A < B - 0.2)$, i.e. $P(B - A > 0.2)$. Let $D = B - A$. $D \sim N(\mu_D, \sigma_D^2)$ with $\mu_D = 10.5 - 10.2 = 0.3$ and $\sigma_D^2 = 0.12^2 + 0.15^2 = 0.0144 + 0.0225 = 0.0369$. $\sigma_D = \sqrt{0.0369} \approx 0.1921$.

$$P(D > 0.2) = P\left(Z > \frac{0.2 - 0.3}{0.1921}\right) = P(Z > -0.5206) = \Phi(0.5206) \approx 0.6985.$$

Answer: ≈ 0.699 .

METHOD Computing $P(X > Y)$ for Independent Normals

To find $P(X > Y)$ where $X \sim N(\mu_1, \sigma_1^2)$, $Y \sim N(\mu_2, \sigma_2^2)$ independent:

1. Define $D = X - Y$. State $D \sim N(\mu_1 - \mu_2, \sigma_1^2 + \sigma_2^2)$.
2. The required probability is $P(D > 0)$.
3. Standardise: $z = \frac{0 - (\mu_1 - \mu_2)}{\sqrt{\sigma_1^2 + \sigma_2^2}} = \frac{\mu_2 - \mu_1}{\sqrt{\sigma_1^2 + \sigma_2^2}}$.
4. Use tables: $P(D > 0) = 1 - \Phi(z)$ if $z > 0$, or $\Phi(-z)$ by symmetry.

WORKED EXAMPLE 16 Probability One Component Outlasts Another

The lifetimes (in hours) of two types of electronic component are modelled as independent normal variables: Type X: $N(500, 50^2)$, Type Y: $N(480, 40^2)$. A device uses one com-

ponent of each type. Find the probability that the Type X component outlasts the Type Y component.

Statement: Find $P(X > Y)$.

SOLUTION


Detailed Solution: Let $D = X - Y$. Since X and Y are independent,

$$D \sim N(\mu_D, \sigma_D^2), \quad \mu_D = 500 - 480 = 20, \quad \sigma_D^2 = 50^2 + 40^2 = 2500 + 1600 = 4100.$$

$$\sigma_D = \sqrt{4100} = 10\sqrt{41} \approx 64.03. \text{ We need } P(D > 0).$$

$$z = \frac{0 - 20}{64.03} = -0.3123.$$

$$P(D > 0) = P(Z > -0.3123) = \Phi(0.3123) \approx 0.6225. \text{ Answer: } 0.623 \text{ (3 s.f.)}.$$



Normal curves for $X \sim N(0, 1)$, $Y \sim N(1, 1.5)$, their sum $X + Y \sim N(1, 2.5)$ and difference $X - Y \sim N(-1, 2.5)$. Note how the sum/difference curves are wider (larger variance) and centred at the sum/difference of the means.

7.2 Linear Combinations: $aX + bY$ and the Crucial Distinction

The theorem extends immediately to any linear combination.

PROPERTY Linear Combination of Two Independent Normal Variables

If $X \sim N(\mu_1, \sigma_1^2)$ and $Y \sim N(\mu_2, \sigma_2^2)$ are independent, then for any constants $a, b \in \mathbb{R}$:

$$aX + bY \sim N(a\mu_1 + b\mu_2, a^2\sigma_1^2 + b^2\sigma_2^2).$$

This formula is the engine for all combination problems. But there is a **subtle and vital distinction** that separates A-grade candidates from the rest: the difference between multiplying a single observation by n and summing n independent observations.

KEY POINTS The nX vs $X_1 + \dots + X_n$ Trap — Examiners' Favourite

Let $X \sim N(\mu, \sigma^2)$.


- One observation scaled by n : $nX \sim N(n\mu, n^2\sigma^2)$. Variance is multiplied by n^2 .
- Sum of n independent copies: $X_1 + X_2 + \dots + X_n \sim N(n\mu, n\sigma^2)$. Variance is multiplied by n .

The means are the same, but the variances differ by a factor of n . nX is much more variable than the sum of n independent observations because all the “eggs are in one basket” — a single extreme value gets magnified n times. In the sum, extremes tend to cancel out.

WARNING $3X$ vs $X_1 + X_2 + X_3$ — Never Confuse Them!

- If a problem says “the mass of a box is X kg; find the probability that the total mass of 3 boxes exceeds...”, you need $X_1 + X_2 + X_3$ (variance $3\sigma^2$).
- If it says “a box is weighed and the reading is multiplied by 3...” or “a single box’s mass is X ; find $P(3X > \dots)$ ”, you use $3X$ (variance $9\sigma^2$).

Read the wording carefully: “three boxes” → sum of three independent variables; “three times the mass” → multiple of one variable.



Comparison of distributions: $X \sim N(0, 1)$ (blue), $4X \sim N(0, 16)$ (orange, very wide), and $X_1 + X_2 + X_3 + X_4 \sim N(0, 4)$ (green dashed, narrower). The sum of four independent copies has variance 4, while four times a single copy has variance 16.

EXAMPLE Crates of Bottles — Sum vs Multiple

The mass of a glass bottle is $B \sim N(0.35, 0.02^2)$ kg. The mass of an empty crate is $C \sim N(1.2, 0.05^2)$ kg. All masses are independent.

1. A crate holds 12 bottles. Find the probability that a full crate (crate + 12 bottles) has mass between 5.4 kg and 5.6 kg.
2. A quality control check weighs one bottle and multiplies the reading by 12 to estimate the total bottle mass. Find the probability that this estimate exceeds 4.5 kg.

Solution:

1. Let $T = C + B_1 + \dots + B_{12}$ (sum of 12 independent bottles + crate). $\mu_T = 1.2 + 12 \times 0.35 = 1.2 + 4.2 = 5.4$ kg. $\sigma_T^2 = 0.05^2 + 12 \times 0.02^2 = 0.0025 + 12 \times 0.0004 = 0.0025 + 0.0048 = 0.0073$.
 $\sigma_T = \sqrt{0.0073} \approx 0.08544$ kg. $P(5.4 < T < 5.6) = P\left(\frac{5.4-5.4}{0.08544} < Z < \frac{5.6-5.4}{0.08544}\right) = P(0 < Z < 2.341) = \Phi(2.341) - 0.5 \approx 0.9904 - 0.5 = 0.4904$.
2. Estimate $E = 12B$ (one bottle scaled by 12). $E \sim N(12 \times 0.35, 12^2 \times 0.02^2) = N(4.2, 144 \times 0.0004) = N(4.2, 0.0576)$. $\sigma_E = \sqrt{0.0576} = 0.24$ kg. $P(E > 4.5) = P\left(Z > \frac{4.5-4.2}{0.24}\right) = P(Z > 1.25) = 1 - \Phi(1.25) \approx 1 - 0.8944 = 0.1056$.

Notice how the estimate $12B$ is far more variable (sd 0.24 kg) than the actual total of 12 bottles (sd ≈ 0.085 kg for the bottles alone).

7.3 Sums of Several Independent Normal Variables

The pattern generalises naturally.

PROPERTY Sum of n Independent Normal Variables

If $X_1, X_2, \dots, X_n$ are independent with $X_i \sim N(\mu_i, \sigma_i^2)$, then

$$S_n = X_1 + X_2 + \dots + X_n \sim N\left(\sum_{i=1}^n \mu_i, \sum_{i=1}^n \sigma_i^2\right).$$

If they are identically distributed (i.i.d.) with $X_i \sim N(\mu, \sigma^2)$, then

$$S_n \sim N(n\mu, n\sigma^2).$$

EXAMPLE Total Length of Rods

Metal rods have lengths $L \sim N(1.50, 0.02^2)$ m. Five rods are laid end to end. Find the probability that the total length exceeds 7.55 m.

Solution: Let $T = L_1 + L_2 + L_3 + L_4 + L_5$ (five independent rods). $T \sim N(5 \times 1.50, 5 \times 0.02^2) =$

$$N(7.50, 5 \times 0.0004) = N(7.50, 0.002). \quad \sigma_T = \sqrt{0.002} \approx 0.04472 \text{ m.} \quad P(T > 7.55) = P\left(Z > \frac{7.55-7.50}{0.04472}\right) = P(Z > 1.118) = 1 - \Phi(1.118) \approx 1 - 0.8686 = 0.1314.$$

7.4 Synthesis Problems: Combining with Standardisation and Inverse Tables

GCE questions often blend this section with earlier skills: finding an unknown μ or σ from a probability statement about a combination, or using the inverse normal tables (percentage points) to determine a threshold.

EXAMPLE Finding an Unknown Mean from a Combination Probability

The mass of a standard loaf of bread is $N(\mu, 0.01^2)$ kg. A pack contains 6 loaves. The probability that a pack weighs less than 4.80 kg is 0.025. Find μ .

Solution: Let $P = L_1 + \dots + L_6$ be the pack mass. $P \sim N(6\mu, 6 \times 0.01^2) = N(6\mu, 0.0006)$. $\sigma_P = \sqrt{0.0006} \approx 0.02449$. Given $P(P < 4.80) = 0.025$. From tables, $\Phi^{-1}(0.025) = -1.96$ (the 2.5% lower tail). Standardise:

$$\frac{4.80 - 6\mu}{0.02449} = -1.96 \quad \Rightarrow \quad 4.80 - 6\mu = -1.96 \times 0.02449 \approx -0.0480.$$

$$6\mu = 4.80 + 0.0480 = 4.848 \quad \Rightarrow \quad \mu = \frac{4.848}{6} = 0.808 \text{ kg.}$$

EXAMPLE Finding an Unknown Standard Deviation — Difference of Two Variables

Two machines fill jars of jam. Machine A: $N(350, \sigma^2)$ g. Machine B: $N(355, \sigma^2)$ g. (Same variance σ^2). Jars from A and B are independent. The probability that a jar from A contains more jam than a jar from B is 0.15. Find σ .

Solution: Let $D = A - B$. $D \sim N(350 - 355, \sigma^2 + \sigma^2) = N(-5, 2\sigma^2)$. We need $P(A > B) = P(D > 0) = 0.15$. Standardise: $P\left(Z > \frac{0 - (-5)}{\sqrt{2}\sigma}\right) = 0.15 \Rightarrow P\left(Z > \frac{5}{\sqrt{2}\sigma}\right) = 0.15$. From tables, the upper 15% point is $z_{0.15} \approx 1.036$ (since $\Phi(1.036) \approx 0.85$).

$$\frac{5}{\sqrt{2}\sigma} = 1.036 \quad \Rightarrow \quad \sqrt{2}\sigma = \frac{5}{1.036} \approx 4.826 \quad \Rightarrow \quad \sigma = \frac{4.826}{\sqrt{2}} \approx 3.413 \text{ g.}$$

WORKED EXAMPLE 17 GCE-Style Synthesis: Crate Weight Limit

A factory packs apples into crates. The mass of an apple is $A \sim N(150, 20^2)$ g. The mass of an empty crate is $C \sim N(500, 30^2)$ g. Each crate holds 30 apples. All masses are independent.

1. Find the mean and standard deviation of the mass of a full crate.
2. The factory requires that no more than 2.5% of full crates exceed a weight limit W g. Find W .
3. A quality inspector picks one apple and multiplies its mass by 30 to estimate the total apple mass in a crate. Find the probability that this estimate exceeds the actual total apple mass by more than 200 g.

Statement: Full crate = crate + 30 apples. Estimate = $30 \times$ (one apple).

SOLUTION

Detailed Solution:

1. Let $F = C + A_1 + \dots + A_{30}$. $\mu_F = 500 + 30 \times 150 = 500 + 4500 = 5000$ g. $\sigma_F^2 = 30^2 + 30 \times 20^2 = 900 + 30 \times 400 = 900 + 12000 = 12900$. $\sigma_F = \sqrt{12900} \approx 113.58$ g.
2. We need $P(F > W) = 0.025 \Rightarrow P(F \leq W) = 0.975$. The 97.5% point of $N(0, 1)$ is $z = 1.96$. $\frac{W-5000}{113.58} = 1.96 \Rightarrow W = 5000 + 1.96 \times 113.58 \approx 5000 + 222.6 = 5222.6$ g.
3. Let $E = 30A$ (estimate from one apple). Actual total apple mass $T = A_1 + \dots + A_{30}$. $E \sim N(30 \times 150, 30^2 \times 20^2) = N(4500, 360000)$. $T \sim N(30 \times 150, 30 \times 20^2) = N(4500, 12000)$. We need $P(E - T > 200)$. Let $D = E - T$. E and T are independent (the single apple is different from the 30 in the crate). $D \sim N(4500 - 4500, 360000 + 12000) = N(0, 372000)$. $\sigma_D = \sqrt{372000} \approx 609.9$ g. $P(D > 200) = P\left(Z > \frac{200}{609.9}\right) = P(Z > 0.3279) = 1 - \Phi(0.3279) \approx 1 - 0.6285 = 0.3715$.

Answers: (i) Mean 5000 g, sd ≈ 114 g. (ii) $W \approx 5223$ g. (iii) ≈ 0.372 .

EXERCISE 8 Practice: Combinations of Normals

1. $X \sim N(10, 4)$, $Y \sim N(6, 9)$, independent. Find $P(X + Y > 20)$ and $P(X - Y < 2)$.
2. The height of a man is $M \sim N(175, 7^2)$ cm, a woman $W \sim N(162, 6^2)$ cm, independent. Find the probability that a randomly chosen man is at least 15 cm taller than a randomly chosen woman.
3. A box contains 20 screws. The mass of a screw is $N(5.0, 0.2^2)$ g. The empty box is $N(50, 1^2)$ g. Find the probability that a full box weighs between 149 g and 151 g.
4. $X \sim N(\mu, 25)$. A sample of 4 independent observations gives $\bar{X} = \frac{X_1 + X_2 + X_3 + X_4}{4}$. Given $P(\bar{X} > 52) = 0.05$, find μ . (Hint: $\bar{X}$ is a linear combination.)
5. Two independent measurements of a distance are $D_1 \sim N(d, \sigma^2)$ and $D_2 \sim N(d, \sigma^2)$. The average $\frac{D_1 + D_2}{2}$ is used as the final estimate. Show that the variance of the estimate is $\sigma^2/2$. If instead a single measurement is multiplied by 1 (i.e. just D_1), compare the vari-

ances. Explain why averaging reduces uncertainty.

SOLUTION Exercise 1

$X + Y \sim N(16, 13)$. $P(X + Y > 20) = P(Z > \frac{4}{\sqrt{13}}) = P(Z > 1.109) \approx 0.1336$. $X - Y \sim N(4, 13)$.
 $P(X - Y < 2) = P(Z < \frac{-2}{\sqrt{13}}) = P(Z < -0.555) = \Phi(-0.555) \approx 0.2895$.

SOLUTION Exercise 2

$D = M - W \sim N(13, 7^2 + 6^2) = N(13, 85)$. $\sigma_D = \sqrt{85} \approx 9.22$. $P(D \geq 15) = P(Z \geq \frac{2}{9.22}) = P(Z \geq 0.217) \approx 1 - 0.586 = 0.414$.

SOLUTION Exercise 3

Full box $F = B + S_1 + \dots + S_{20} \sim N(50 + 100, 1^2 + 20 \times 0.2^2) = N(150, 1 + 0.8) = N(150, 1.8)$.
 $\sigma_F = \sqrt{1.8} \approx 1.3416$. $P(149 < F < 151) = P(\frac{-1}{1.3416} < Z < \frac{1}{1.3416}) = P(-0.745 < Z < 0.745) = 2\Phi(0.745) - 1 \approx 2 \times 0.772 - 1 = 0.544$.

SOLUTION Exercise 4

$\bar{X} = \frac{1}{4}(X_1 + X_2 + X_3 + X_4) \sim N(\mu, \frac{25}{4}) = N(\mu, 6.25)$. $P(\bar{X} > 52) = 0.05 \Rightarrow \frac{52 - \mu}{2.5} = 1.645 \Rightarrow 52 - \mu = 4.1125 \Rightarrow \mu = 47.8875$.

SOLUTION Exercise 5

Average $A = \frac{D_1 + D_2}{2} \sim N(d, \frac{\sigma^2 + \sigma^2}{4}) = N(d, \sigma^2/2)$. Variance = $\sigma^2/2$. Single measurement $D_1 \sim N(d, \sigma^2)$. Variance = σ^2 . Averaging halves the variance because independent errors tend to cancel out ($\text{Var}(D_1 + D_2) = 2\sigma^2$, then dividing by 2 gives factor 1/4 on the sum, so $\frac{1}{4} \times 2\sigma^2 = \sigma^2/2$). Using one measurement keeps the full variance σ^2 . Averaging is always better for precision.

DID YOU KNOW? Why the Normal Family is Closed Under Linear Combinations

The normal distribution is a stable distribution: any linear combination of independent normals is normal. This is not true for most other distributions (e.g., the sum of two independent exponentials is Gamma, not exponential; the sum of two independent uniforms is triangular, not uniform). This remarkable property makes the normal distribution uniquely tractable for modelling aggregate phenomena — total demand, combined errors, portfolio returns — and is a key reason for its central role in statistics (Central Limit Theorem).

KEY POINTS Key Points for the Exam

- Independence is essential. The theorems only hold if the variables are independent. Always state “ X and Y are independent”.
- Define the new variable explicitly. Write “Let $T = X + Y \sim N(\mu_X + \mu_Y, \sigma_X^2 + \sigma_Y^2)$ ” before calculating. This gains method marks.
- Variances always add. For $X - Y$, variance is $\sigma_X^2 + \sigma_Y^2$, never $\sigma_X^2 - \sigma_Y^2$.
- Distinguish nX from $X_1 + \dots + X_n$. nX : variance $n^2\sigma^2$. Sum of n i.i.d.: variance $n\sigma^2$. Read the wording: “ n items” $\rightarrow$ sum; “ n times the value” $\rightarrow$ multiple.
- Standardise correctly. $z = \frac{X - \mu_{\text{comb}}}{\sigma_{\text{comb}}}$. Use $\sqrt{\text{variance}}$ for σ .
- Inverse problems. If given a probability, find the z -value from tables (percentage points), then solve for the unknown μ or σ .

EXPERIMENT Guided Investigation: Portfolio Risk

In finance, a portfolio combines assets. Suppose Asset A has return $R_A \sim N(8\%, 15^2)$ and Asset B has return $R_B \sim N(12\%, 20^2)$ (percentages). An investor puts fraction w in A and $(1 - w)$ in B. The portfolio return is $R_p = wR_A + (1 - w)R_B$.

1. Assuming independence, write the distribution of R_p in terms of w .
2. Find the value of w that minimises the variance (risk) of the portfolio.
3. If the correlation between A and B is actually $\rho = 0.5$, how does the variance formula change? (Formula: $\text{Var}(wX + (1 - w)Y) = w^2\sigma_X^2 + (1 - w)^2\sigma_Y^2 + 2w(1 - w)\rho\sigma_X\sigma_Y$.)
4. Discuss: why does diversification (mixing assets) reduce risk even when returns are positively correlated?


This illustrates the real-world power of combining normal variables — the foundation of Modern Portfolio Theory (Markowitz, Nobel Prize 1990).

8

INTEGRATION, ASSESSMENT AND REMEDIATION

Building on our work with combinations of independent Normal variables, we now consolidate the entire chapter. This section is designed to help you synthesise the three major continuous distributions — Uniform, Exponential and Normal — recognise which one applies to a given problem, practise under exam conditions, identify and correct your most frequent errors, and finally assess your readiness for the end-of-term test.

8.1 Synthesis Map of the Chapter



The map above summarises the logical flow of the chapter. Use it as a mental checklist: when you read a problem, first decide which distribution fits the context (decision diamond), then follow the relevant branch to the required technique (expectation/variance, standardisation, approximation, or combination).

METHOD Decision Guide — Recognising the Correct Distribution

Step 1: Identify the random variable and its support.

- If the variable is equally likely over a finite interval $[a, b]$ (e.g. waiting time for a bus that arrives exactly every 20 minutes, rounding errors), think **Uniform**.
- If the variable measures time between random events occurring at a constant average rate (e.g. time between customer arrivals, radioactive decay, component lifetime), think **Exponential**. Check for the memoryless property: $P(X > s + t \mid X > s) = P(X > t)$.
- If the variable is a naturally occurring measurement that clusters around a mean (e.g. heights, weights, exam marks, measurement errors), think **Normal**.

Step 2: Check for approximations.

- If the problem gives a Binomial $B(n, p)$ with n large and $np > 5$, $nq > 5$, use **Normal approximation** with continuity correction.
- If the problem gives a Poisson $Po(\lambda)$ with $\lambda > 15$, use **Normal approximation** with continuity correction.

Step 3: Look for combinations. If the problem involves sums or differences of independent Normal variables (e.g. total weight of several items, difference in heights), use the linear combination rule: $aX + bY \sim N(a\mu_X + b\mu_Y, a^2\sigma_X^2 + b^2\sigma_Y^2)$.

8.2 Mixed Revision Exercises

The following exercises cover Lessons 166–182 in increasing order of difficulty. Attempt them without notes first; then check your work against the solutions.

EXERCISE 9 Uniform Distribution — Lesson 166, 167

A machine produces metal rods whose lengths are uniformly distributed between 9.8 cm and 10.2 cm.

1. Write down the probability density function $f(x)$.
2. Find the probability that a randomly chosen rod has length greater than 10.1 cm.
3. Calculate the mean and variance of the rod lengths.
4. If 100 rods are selected independently, estimate the probability that their total length exceeds 1005 cm. (Hint: use the Central Limit Theorem.)

SOLUTION Exercise 1

1. $f(x) = \frac{1}{0.4} = 2.5$ for $9.8 \leq x \leq 10.2$, 0 otherwise.
2. $P(X > 10.1) = \frac{10.2-10.1}{0.4} = 0.25$.
3. $E(X) = \frac{9.8+10.2}{2} = 10.0$ cm. $\text{Var}(X) = \frac{(0.4)^2}{12} = \frac{0.16}{12} \approx 0.01333 \text{ cm}^2$.
4. Let $S = \sum_{i=1}^{100} X_i$. $E(S) = 100 \times 10.0 = 1000$ cm. $\text{Var}(S) = 100 \times \frac{0.16}{12} = \frac{16}{12} \approx 1.333 \text{ cm}^2$. $\sigma_S \approx 1.155$ cm. $P(S > 1005) \approx P\left(Z > \frac{1005-1000}{1.155}\right) = P(Z > 4.33) \approx 0$. (Virtually impossible.)

EXERCISE 10 Exponential Distribution — Lesson 168, 169, 170, 171

The time between successive arrivals at a hospital emergency unit follows an exponential distribution with mean 6 minutes.

1. Find the rate parameter λ and write the pdf.
2. What is the probability that the next arrival occurs after 10 minutes?
3. Given that no arrival has occurred in the first 5 minutes, what is the probability that the next arrival occurs after 10 minutes total? (Use the memoryless property.)
4. The number of arrivals in a one-hour period follows a Poisson distribution. State its parameter and find the probability of exactly 12 arrivals in an hour.

SOLUTION Exercise 2

1. $\lambda = \frac{1}{6}$ per minute. $f(t) = \frac{1}{6}e^{-t/6}$, $t \geq 0$.
2. $P(T > 10) = e^{-10/6} = e^{-5/3} \approx 0.1889$.
3. By memoryless property, $P(T > 10 \mid T > 5) = P(T > 5) = e^{-5/6} \approx 0.4346$.
4. Arrivals per hour $\sim \text{Po}(\lambda t) = \text{Po}(10)$. $P(X = 12) = \frac{e^{-10}10^{12}}{12!} \approx 0.0948$.

EXERCISE 11 Normal Distribution — Lessons 172–179

The masses of bags of cement produced by a factory are normally distributed with mean 50.0 kg and standard deviation 0.5 kg.

1. Find the probability that a randomly chosen bag has mass between 49.5 kg and 50.5

kg.

- Determine the mass exceeded by the heaviest 2.5% of bags.
- A sample of 25 bags is selected. Find the probability that the sample mean mass is less than 49.9 kg.
- The factory adjusts the machine so that only 1% of bags are under 49.5 kg, while the standard deviation remains 0.5 kg. Find the new mean setting.

SOLUTION Exercise 3

- $Z_1 = \frac{49.5-50}{0.5} = -1, Z_2 = \frac{50.5-50}{0.5} = 1. P(-1 < Z < 1) = \Phi(1) - \Phi(-1) = 0.8413 - 0.1587 = 0.6826.$
- Heaviest 2.5% $\Rightarrow P(X > x) = 0.025 \Rightarrow P(Z > z) = 0.025 \Rightarrow z = 1.96. x = \mu + z\sigma = 50 + 1.96 \times 0.5 = 50.98 \text{ kg}.$
- $\bar{X} \sim N(50, \frac{0.5^2}{25} = 0.01). P(\bar{X} < 49.9) = P\left(Z < \frac{49.9-50}{0.1}\right) = P(Z < -1) = 0.1587.$
- $P(X < 49.5) = 0.01 \Rightarrow z = -2.3263 \text{ (from tables)}. 49.5 = \mu + (-2.3263) \times 0.5 \Rightarrow \mu = 49.5 + 1.16315 = 50.66315 \approx 50.66 \text{ kg}.$

EXERCISE 12 Continuity Correction & Approximations — Lessons 180, 181

A fair coin is tossed 200 times. Use a Normal approximation to find the probability of obtaining:

- between 90 and 110 heads inclusive;
- more than 115 heads.

State the conditions that justify the approximation.

SOLUTION Exercise 4

$X \sim B(200, 0.5). np = 100 > 5, nq = 100 > 5 \Rightarrow \text{Normal approximation valid. } \mu = 100, \sigma = \sqrt{200 \times 0.5 \times 0.5} = \sqrt{50} \approx 7.071.$

- $P(90 \leq X \leq 110) \approx P(89.5 < X < 110.5) = P\left(\frac{89.5-100}{7.071} < Z < \frac{110.5-100}{7.071}\right) = P(-1.485 < Z < 1.485) = \Phi(1.485) - \Phi(-1.485) \approx 0.9312 - 0.0688 = 0.8624.$
- $P(X > 115) \approx P(X > 115.5) = P\left(Z > \frac{115.5-100}{7.071}\right) = P(Z > 2.192) = 1 - \Phi(2.192) \approx 1 - 0.9857 = 0.0143.$

EXERCISE 13 Combinations of Independent Normals — Lesson 182

The heights of adult males in a certain population are $N(175, 7^2)$ cm, and adult females are $N(162, 6^2)$ cm. A male and a female are chosen independently.

1. Find the probability that the male is taller than the female.
2. Find the probability that the male is at least 20 cm taller than the female.
3. If two males and one female are chosen independently, find the probability that the sum of the two males' heights exceeds twice the female's height by more than 30 cm.

SOLUTION Exercise 5

Let $M \sim N(175, 49)$, $F \sim N(162, 36)$, independent.

1. $D = M - F \sim N(175 - 162, 49 + 36) = N(13, 85)$. $P(M > F) = P(D > 0) = P\left(Z > \frac{0-13}{\sqrt{85}}\right) = P(Z > -1.409) = \Phi(1.409) \approx 0.9207$.
2. $P(M - F \geq 20) = P(D \geq 20) = P\left(Z \geq \frac{20-13}{\sqrt{85}}\right) = P(Z \geq 0.759) = 1 - \Phi(0.759) \approx 1 - 0.7764 = 0.2236$.
3. Let M_1, M_2 be the two males. $S = M_1 + M_2 - 2F$. $E(S) = 2 \times 175 - 2 \times 162 = 26$. $\text{Var}(S) = 2 \times 49 + 4 \times 36 = 98 + 144 = 242$. $\sigma_S = \sqrt{242} \approx 15.556$. $P(S > 30) = P\left(Z > \frac{30-26}{15.556}\right) = P(Z > 0.257) = 1 - \Phi(0.257) \approx 1 - 0.6015 = 0.3985$.

8.3 Timed GCE Past-Paper Questions

The following questions are typical of GCE Advanced Level Pure Mathematics with Statistics Paper 2. Allow **15 minutes** for Question 1 (12 marks) and **12 minutes** for Question 2 (10 marks). Write your answers as you would in the exam: show all steps, state distributions clearly, and use tables where required.

WORKED EXAMPLE 18 GCE Style Question 1 (12 marks)

The lifetime T (in hours) of a certain electronic component follows an exponential distribution with mean 500 hours.

1. Write down the probability density function of T . [1]
2. Find the probability that a component lasts more than 600 hours. [2]
3. A system uses 4 such components connected in series (the system fails if any component fails). Assuming independence, find the probability that the system operates for at least 600 hours. [3]
4. The number of components failing in a 1000-hour period follows a Poisson distribu-

tion. State its mean and find the probability that exactly 5 components fail in this period. [3]

5. A batch contains 200 components. Use a Normal approximation to estimate the probability that at least 180 components are still functioning after 100 hours. [3]

SOLUTION

Detailed Solution

- $\lambda = \frac{1}{500} = 0.002$. $f(t) = 0.002e^{-0.002t}$, $t \geq 0$.
- $P(T > 600) = e^{-0.002 \times 600} = e^{-1.2} \approx 0.3012$.
- System lifetime = $\min(T_1, T_2, T_3, T_4)$. For independent exponentials, the minimum is exponential with rate $4\lambda = 0.008$. $P(\text{system} > 600) = e^{-0.008 \times 600} = e^{-4.8} \approx 0.00823$. Alternatively, $P(\text{all } 4 > 600) = [P(T > 600)]^4 = (0.3012)^4 \approx 0.00823$.
- Failures in 1000 hours: rate $\lambda t = 0.002 \times 1000 = 2$. So $X \sim \text{Po}(2)$. $P(X = 5) = \frac{e^{-2} 2^5}{5!} = \frac{32e^{-2}}{120} \approx 0.0361$.
- For one component, $P(T > 100) = e^{-0.002 \times 100} = e^{-0.2} \approx 0.8187$. Let Y = number functioning after 100 hours out of 200. $Y \sim B(200, 0.8187)$. $np = 163.74 > 5$, $nq = 36.26 > 5 \Rightarrow$ Normal approximation. $\mu = 163.74$, $\sigma = \sqrt{200 \times 0.8187 \times 0.1813} \approx 5.45$. $P(Y \geq 180) \approx P(Y > 179.5) = P\left(Z > \frac{179.5 - 163.74}{5.45}\right) = P(Z > 2.89) \approx 1 - 0.9981 = 0.0019$.

WORKED EXAMPLE 19 GCE Style Question 2 (10 marks)

The random variable $X \sim N(\mu, \sigma^2)$. It is known that $P(X < 40) = 0.025$ and $P(X > 60) = 0.10$.

- Find the values of μ and σ . [6]
- Hence find $P(45 < X < 55)$. [2]
- A sample of 36 observations is taken from this distribution. Find the probability that the sample mean lies between 48 and 52. [2]

SOLUTION

Detailed Solution

1. From tables: $P(Z < z) = 0.025 \Rightarrow z = -1.96$. $P(Z > z) = 0.10 \Rightarrow z = 1.2816$. Equations:

$$\frac{40 - \mu}{\sigma} = -1.96 \Rightarrow \mu - 1.96\sigma = 40 \quad (1)$$

$$\frac{60 - \mu}{\sigma} = 1.2816 \Rightarrow \mu + 1.2816\sigma = 60 \quad (2)$$

Subtract (1) from (2): $3.2416\sigma = 20 \Rightarrow \sigma = \frac{20}{3.2416} \approx 6.169$. Substitute into (1): $\mu = 40 +$

$$1.96 \times 6.169 \approx 40 + 12.09 = 52.09.$$

$$2. P(45 < X < 55) = P\left(\frac{45-52.09}{6.169} < Z < \frac{55-52.09}{6.169}\right) = P(-1.15 < Z < 0.472) = \Phi(0.472) - \Phi(-1.15) \approx 0.6814 - 0.1251 = 0.5563.$$

$$3. \bar{X} \sim N(52.09, \frac{6.169^2}{36} = 1.056). \sigma_{\bar{X}} \approx 1.028. P(48 < \bar{X} < 52) = P\left(\frac{48-52.09}{1.028} < Z < \frac{52-52.09}{1.028}\right) = P(-3.98 < Z < -0.0875) \approx \Phi(-0.0875) - \Phi(-3.98) \approx 0.4651 - 0 = 0.4651.$$

8.4 Error-Analysis Workshop

The following **five errors** appear repeatedly in GCE scripts. Study each wrong answer, identify the mistake, and write the correct version.

WARNING Checklist of the Five Most Frequent GCE Errors in This Chapter

1. Sign error in z-score: Writing $z = \frac{x-\mu}{\sigma}$ but then using $z = \frac{\mu-x}{\sigma}$ or misreading the table for negative z . Always sketch the Normal curve and shade the required area.
2. Missing continuity correction: Using $P(X \leq k)$ directly as $P(Z \leq \frac{k-\mu}{\sigma})$ when approximating Binomial/Poisson. Always add/subtract 0.5: $P(X \leq k) \approx P(Z \leq \frac{k+0.5-\mu}{\sigma})$.
3. Variance of a difference: For independent X, Y , writing $\text{Var}(X - Y) = \sigma_X^2 - \sigma_Y^2$ instead of $\sigma_X^2 + \sigma_Y^2$. Variances always add for independent variables, even when subtracting.
4. Confusing σ and σ^2 in standardisation: Using σ^2 in the denominator: $z = \frac{x-\mu}{\sigma^2}$. Denominator is standard deviation σ , not variance.
5. Forgetting to adjust parameters for sample means: Using σ instead of $\frac{\sigma}{\sqrt{n}}$ when dealing with $\bar{X}$. Standard error = $\frac{\sigma}{\sqrt{n}}$.

EXPERIMENT Scripted Wrong Answers — Correct Them

Scenario A: $X \sim N(50, 25)$. Find $P(X > 55)$. Wrong script: $z = \frac{55-50}{25} = 0.2$. $P(Z > 0.2) = 1 - 0.5793 = 0.4207$. Error: Used variance 25 instead of standard deviation 5. Correction: $z = \frac{55-50}{5} = 1$. $P(Z > 1) = 1 - 0.8413 = 0.1587$.

Scenario B: $X \sim B(100, 0.3)$. Approximate $P(X \leq 25)$. Wrong script: $\mu = 30$, $\sigma = \sqrt{21} \approx 4.58$. $P(X \leq 25) \approx P\left(Z \leq \frac{25-30}{4.58}\right) = P(Z \leq -1.09) = 0.1379$. Error: No continuity correction.

Correction: $P(X \leq 25) \approx P\left(Z \leq \frac{25.5-30}{4.58}\right) = P(Z \leq -0.982) = 0.1635$.

Scenario C: $X \sim N(10, 4)$, $Y \sim N(8, 9)$ independent. Find $P(X - Y > 5)$. Wrong script: $D = X - Y \sim N(2, 4 - 9 = -5)$ — impossible variance. Error: Subtracted variances. Correction: $D \sim N(2, 4 + 9 = 13)$. $P(D > 5) = P\left(Z > \frac{5-2}{\sqrt{13}}\right) = P(Z > 0.832) = 0.2027$.

8.5 Remediation Tasks

Based on the error analysis above, the three weakest skills are:

1. Standardisation and correct use of Normal tables (including negative z).
2. Applying continuity correction correctly in Normal approximations.
3. Handling combinations of independent Normal variables (especially variance of differences).

Complete the following targeted tasks. Check your answers against the solutions provided.

METHOD Remediation Task 1: Standardisation Drill

For each of the following, find the required probability using Normal tables. Show the z -calculation and the table lookup.

1. $X \sim N(100, 16)$. Find $P(X < 95)$.
2. $X \sim N(50, 9)$. Find $P(X > 56)$.
3. $X \sim N(20, 4)$. Find $P(18 < X < 23)$.
4. $X \sim N(\mu, 25)$. Given $P(X < 30) = 0.9772$, find μ .

Answers: (1) $z = -1.25$, $P = 0.1056$. (2) $z = 2$, $P = 0.0228$. (3) $z_1 = -1$, $z_2 = 1.5$, $P = 0.7745$. (4) $z = 2$, $\frac{30-\mu}{5} = 2 \Rightarrow \mu = 20$.

METHOD Remediation Task 2: Continuity Correction Practice

For each Binomial/Poisson situation, write the correct Normal approximation statement with continuity correction, then compute the approximate probability.

1. $X \sim B(50, 0.4)$. Approximate $P(X \geq 22)$.
2. $X \sim Po(20)$. Approximate $P(X < 15)$.
3. $X \sim B(200, 0.15)$. Approximate $P(25 \leq X \leq 35)$.

Answers: (1) $\mu = 20$, $\sigma = \sqrt{12} \approx 3.464$. $P(X \geq 22) \approx P(Z \geq \frac{21.5-20}{3.464}) = P(Z \geq 0.433) = 0.3325$.
 (2) $\mu = 20$, $\sigma = \sqrt{20} \approx 4.472$. $P(X < 15) \approx P(Z < \frac{14.5-20}{4.472}) = P(Z < -1.23) = 0.1093$. (3) $\mu = 30$, $\sigma = \sqrt{25.5} \approx 5.05$. $P(25 \leq X \leq 35) \approx P(\frac{24.5-30}{5.05} \leq Z \leq \frac{35.5-30}{5.05}) = P(-1.09 \leq Z \leq 1.09) = 0.7242$.

METHOD Remediation Task 3: Combinations of Normals

Let $X \sim N(30, 9)$, $Y \sim N(20, 16)$, independent.

1. Find $P(X + Y > 55)$.

2. Find $P(X - Y < 5)$.

3. Find $P(2X + 3Y < 130)$.

Answers: (1) $X + Y \sim N(50, 25)$. $P(Z > \frac{55-50}{5}) = P(Z > 1) = 0.1587$. (2) $X - Y \sim N(10, 25)$. $P(Z < \frac{5-10}{5}) = P(Z < -1) = 0.1587$. (3) $2X + 3Y \sim N(120, 4 \times 9 + 9 \times 16 = 36 + 144 = 180)$. $P(Z < \frac{130-120}{\sqrt{180}}) = P(Z < 0.745) = 0.7722$.

8.6 Self-Assessment Checklist

Use this checklist to evaluate your mastery of the chapter objectives. Tick each item only if you can do it confidently **without notes** and **within the time limits** suggested.

Objective	Description	Mastered?
1	Define Uniform distribution, write its pdf, cdf, mean, variance.	☐
2	Solve problems involving Uniform distribution (probabilities, percentiles).	☐
3	Define Exponential distribution, state its memoryless property, link to Poisson.	☐
4	Compute probabilities, mean, variance for Exponential; use memoryless property.	☐
5	State properties of Normal distribution; standardise to $Z \sim N(0, 1)$.	☐
6	Use Normal tables to find $\Phi(z)$ and z -values for given probabilities.	☐
7	Solve problems for any $N(\mu, \sigma^2)$: probabilities, percentiles, find μ or σ .	☐
8	Apply continuity correction for Normal approximation to Binomial and Poisson.	☐
9	Check conditions for approximations ($np > 5$, $nq > 5$, $\lambda > 15$).	☐
10	Handle linear combinations of independent Normal variables (sums, differences, multiples).	☐
11	Solve problems involving sample means (standard error $\sigma/\sqrt{n}$).	☐
12	Interpret and solve multi-part GCE questions under timed conditions.	☐

DID YOU KNOW?

Preparation for the End-of-Term Test

- Review the concept map at the start of this section — redraw it from memory.
- Re-do the two timed GCE questions (Section 8.3) under strict time limits; mark them

using the detailed solutions.

- Complete all three Remediation Tasks if you ticked fewer than 10 items above.
- Create a one-page "cheat sheet" (for revision only, not for the exam) containing: key formulas for each distribution, standardisation steps, continuity correction rules, combination rules, and the five common errors from the Warning box.
- Past papers: Work through at least two full Paper 2 statistics sections from recent GCE sessions (available from your teacher or the GCE Board).

KEY POINTS Key Takeaways for Chapter PMS14

- Uniform: Constant pdf on $[a, b]$; mean $(a + b)/2$, variance $(b - a)^2/12$.
- Exponential: Memoryless; mean $1/\lambda$, variance $1/\lambda^2$; inter-arrival times of Poisson process.
- Normal: Bell-shaped; standardise via $Z = \frac{X - \mu}{\sigma}$; use tables for $\Phi(z)$.
- Approximations: Binomial $\rightarrow$ Normal if $np > 5, nq > 5$; Poisson $\rightarrow$ Normal if $\lambda > 15$; always use continuity correction.
- Combinations: If X, Y independent Normals, $aX + bY \sim N(a\mu_X + b\mu_Y, a^2\sigma_X^2 + b^2\sigma_Y^2)$.
- Exam technique: Sketch the Normal curve, show z -calculations, state approximations used, and check that variances add (never subtract).

9 INTEGRATION ACTIVITY

9.1 Aims of the activity

This integration activity brings together, in a single authentic logistics case study, all the special continuous distributions studied in this chapter. By the end of the activity, you should be able to:

- compute the **expectation** and **variance** of a **uniform (rectangular) distribution** and use it to find probabilities as ratios of lengths;
- exploit the **Poisson–exponential link**: Poisson arrivals in time t correspond to exponential inter-arrival times with the same rate;
- standardise a Normal variable, use the **standard normal tables** in both directions, and **de-standardise** to find real values;
- construct the **central 95% interval** of a Normal distribution;
- apply the **Normal approximation to the binomial** with a **continuity correction**;
- use the distribution of a **sum of independent Normal variables**.

WARNING Working rules for the activity

Work in groups of four. Use the standard normal tables provided. Give all probabilities to 4 decimal places and all times to 1 decimal place. State clearly every distribution you use, with its parameters, before computing. Remember: for a continuous variable, $P(X \leq a) = P(X < a)$, but for the binomial approximation the continuity correction depends on whether the inequality is strict or not.

9.2 Situation

The Buea–Douala Express Agency.

A transport agency operates express buses on the Buea–Douala highway. Over one term, the operations manager collected data on five aspects of the service, and the following models were found to fit the data well.

1. **Loading bay.** The time T (in minutes) that a parcel spends on the loading bay before being placed on a bus is uniformly distributed between 5 and 25 minutes: $T \sim U(5, 25)$.
2. **Ticket office.** Customers arrive at the ticket office according to a Poisson process with rate 3 customers per 10 minutes. The waiting time W (in minutes) between two consecutive arrivals is therefore exponential.
3. **Journey time.** The journey time J (in minutes) of a bus from Buea to Douala is Normally distributed with mean 75 and standard deviation 8: $J \sim N(75, 8^2)$.

4. **Passenger satisfaction.** Based on surveys, the probability that any passenger is satisfied with the service is $p = 0.3$, independently of other passengers. In a given week, 300 passengers are surveyed; let X be the number satisfied.
5. **Luggage.** The mass of luggage per passenger is Normally distributed with mean 18 kg and variance 9 kg^2 (standard deviation 3 kg), independently from passenger to passenger. A bus carries 40 passengers; let S be the total luggage mass.

The manager wants answers to several operational questions, and your group has been hired as consultants.

9.3 Tasks

Part A — The loading bay (uniform distribution).

1. Write down the probability density function of T and sketch it.
2. Calculate $E(T)$ and $\text{Var}(T)$.
3. Find the probability that a parcel spends more than 20 minutes on the loading bay.

Part B — The ticket office (Poisson and exponential distributions).

4. Show that the rate per minute is $\lambda = 0.3$, and write down the probability density function of the inter-arrival time W , together with $E(W)$.
5. Find the probability that the next customer arrives within 2 minutes of the previous one.
6. Using the Poisson distribution, find the probability of exactly 4 arrivals in a 10-minute period.
7. State, in one sentence, the link between the exponential and Poisson distributions illustrated by this situation.

Part C — The journey time (Normal distribution).

8. Find the probability that a journey lasts more than 90 minutes.
9. Determine the central 95% interval for journey times, i.e. the interval within which 95% of all journeys fall.
10. Find the journey time exceeded by only 5% of journeys (de-standardising).

Part D — Passenger satisfaction (Normal approximation to the binomial).

11. Explain why $X \sim B(300, 0.3)$ may be approximated by a Normal distribution, and give the parameters of the approximating distribution.
12. Using a continuity correction, estimate the probability that more than 100 of the 300 passengers are satisfied.

Part E — Total luggage (sum of independent Normal variables).

13. State the distribution of the total luggage mass S of the 40 passengers, justifying your answer.

14. Find the probability that the total luggage mass exceeds 750 kg.


Part F — Presentation. Prepare a flipchart with your full solutions. Each step will be validated by the class against the methods of this chapter: correct identification of the model, correct use of tables, correct continuity correction, and correct combination of Normal variables.

9.4 Model answer

Part A — The loading bay.

1. Since $T \sim U(5, 25)$, the probability density function is constant on $[5, 25]$ and the total area under it must equal 1, so its height is $\frac{1}{25 - 5}$:

$$f(t) = \begin{cases} \frac{1}{20} = 0.05, & 5 \leq t \leq 25, \\ 0, & \text{otherwise.} \end{cases}$$



Density of $T \sim U(5, 25)$; the shaded area is $P(T > 20) = 5 \times 0.05 = 0.25$.

2. For $U(a, b)$, $E(T) = \frac{a+b}{2}$ and $\text{Var}(T) = \frac{(b-a)^2}{12}$:

$$E(T) = \frac{5+25}{2} = 15 \text{ min}, \quad \text{Var}(T) = \frac{(25-5)^2}{12} = \frac{400}{12} = \frac{100}{3} \approx 33.3 \text{ min}^2.$$

3. For a uniform distribution, probabilities are ratios of lengths (areas of rectangles):

$$P(T > 20) = \frac{25 - 20}{25 - 5} = \frac{5}{20} = 0.25.$$

Part B — The ticket office.

4. The rate is 3 customers per 10 minutes, so per minute $\lambda = \frac{3}{10} = 0.3$ customers per minute. Hence $W \sim \text{Exp}(0.3)$ with density

$$f(w) = 0.3 e^{-0.3w}, \quad w \geq 0, \quad E(W) = \frac{1}{\lambda} = \frac{1}{0.3} = \frac{10}{3} \approx 3.3 \text{ min}.$$

This mean is consistent with the situation: at 3 customers per 10 minutes, one expects on average one customer every $\frac{10}{3}$ minutes.

5. Using the cumulative distribution function $P(W \leq w) = 1 - e^{-\lambda w}$:

$$P(W \leq 2) = 1 - e^{-0.3 \times 2} = 1 - e^{-0.6} = 1 - 0.5488 = 0.4512.$$

6. The number of arrivals in a 10-minute period is $N \sim \text{Po}(\lambda t) = \text{Po}(0.3 \times 10) = \text{Po}(3)$:

$$P(N = 4) = e^{-3} \frac{3^4}{4!} = e^{-3} \times \frac{81}{24} = 0.0498 \times 3.375 = 0.1680.$$

7. Link: if arrivals form a Poisson process with rate λ per unit time, then the number of arrivals in a time interval of length t is $\text{Po}(\lambda t)$, and the waiting time between two consecutive arrivals is exponential with parameter λ (mean $1/\lambda$).

Part C — The journey time.

8. $J \sim N(75, 8^2)$. Standardising with $Z = \frac{J - 75}{8} \sim N(0, 1)$:

$$P(J > 90) = P\left(Z > \frac{90 - 75}{8}\right) = P(Z > 1.875) = 1 - \Phi(1.875).$$

Interpolating between $\Phi(1.87) = 0.9693$ and $\Phi(1.88) = 0.9699$, we get $\Phi(1.875) \approx 0.9696$, so

$$P(J > 90) = 1 - 0.9696 = 0.0304.$$

9. The central 95% interval for Z is $[-1.96, 1.96]$, since $\Phi(1.96) = 0.975$ leaves 2.5% in each tail. De-standardising with $j = \mu + z\sigma$:

$$75 - 1.96 \times 8 = 59.32, \quad 75 + 1.96 \times 8 = 90.68.$$

So 95% of journeys last between approximately 59.3 and 90.7 minutes.

10. We need j with $P(J > j) = 0.05$, i.e. $P(Z \leq z) = 0.95$, giving $z = 1.645$ from the tables. De-standardising:

$$j = 75 + 1.645 \times 8 = 75 + 13.16 = 88.2 \text{ minutes (1 d.p.)}.$$

Part D — Passenger satisfaction.

11. $X \sim B(300, 0.3)$ with $np = 300 \times 0.3 = 90$ and $n(1 - p) = 300 \times 0.7 = 210$, both well above 5, so the Normal approximation is valid:

$$X \approx Y \sim N(np, np(1 - p)) = N(90, 63), \quad \sigma = \sqrt{63} \approx 7.937.$$

12. “More than 100” means $X \geq 101$; the continuity correction replaces the discrete value 101 by the boundary 100.5:

$$P(X > 100) \approx P(Y \geq 100.5) = P\left(Z \geq \frac{100.5 - 90}{7.937}\right) = P(Z \geq 1.323).$$

From the tables, $\Phi(1.32) = 0.9066$ and $\Phi(1.33) = 0.9082$; interpolating, $\Phi(1.323) \approx 0.9071$. Hence

$$P(X > 100) \approx 1 - 0.9071 = 0.0929.$$

Part E — Total luggage.

13. Each luggage mass $M_i \sim N(18, 9)$ independently. A sum of independent Normal variables is Normal, with the means added and the variances added:

$$E(S) = 40 \times 18 = 720 \text{ kg}, \quad \text{Var}(S) = 40 \times 9 = 360, \quad \sigma_S = \sqrt{360} \approx 18.974 \text{ kg}.$$

Thus $S \sim N(720, 360)$.

14. Standardising:

$$P(S > 750) = P\left(Z > \frac{750 - 720}{18.974}\right) = P(Z > 1.581).$$

From the tables, $\Phi(1.58) = 0.9429$, so interpolating $\Phi(1.581) \approx 0.9431$, and

$$P(S > 750) = 1 - 0.9431 = 0.0569.$$

There is about a 5.7% chance that the luggage exceeds 750 kg — useful information for the agency's weight policy.

WARNING Pitfalls to avoid in this type of problem

- Do not confuse the variance with the standard deviation: $N(75, 8^2)$ has $\sigma = 8$, but $N(720, 360)$ has $\sigma = \sqrt{360}$, not 360.
- For sums of independent variables, add the variances, never the standard deviations.
- The continuity correction moves the boundary by 0.5 towards the included region: $X \geq 101$ becomes $Y \geq 100.5$, not $Y \geq 101.5$.
- In the Poisson–exponential link, the rate λ must be expressed in the same time unit as the waiting time.

KEY POINTS What this activity consolidates

- Uniform: $E = \frac{a+b}{2}$, $\text{Var} = \frac{(b-a)^2}{12}$, probabilities as length ratios.
- Exponential: $P(W \leq w) = 1 - e^{-\lambda w}$, $E(W) = \frac{1}{\lambda}$; it is the inter-arrival time of a Poisson process of rate λ .
- Normal: standardise, read Φ from tables, de-standardise with $x = \mu + z\sigma$; the central 95% interval uses $z = \pm 1.96$ and the upper 5% point uses $z = 1.645$.
- Binomial $\rightarrow$ Normal when np and $n(1-p)$ both exceed 5, always with a continuity correction.
- Sums of independent Normal variables are Normal: add the means and add the variances.

10 METHODS AND WORKED EXAMPLES

10.1 Method 1: Computing probabilities, mean and variance for a uniform distribution

METHOD Working with the rectangular distribution $U(a, b)$

1. Identify the interval $[a, b]$ and write the density $f(x) = \frac{1}{b-a}$ for $a \leq x \leq b$, 0 otherwise.
2. For any sub-interval $[c, d] \subseteq [a, b]$, compute $P(c \leq X \leq d) = \frac{d-c}{b-a}$ (ratio of lengths). A sketch of the rectangle always helps.
3. Write down directly $E(X) = \frac{a+b}{2}$ and $\text{Var}(X) = \frac{(b-a)^2}{12}$, hence $\sigma = \frac{b-a}{2\sqrt{3}}$.
4. For "given that" questions, restrict the interval and recompute the ratio of lengths (conditional probability): the conditional distribution is again uniform on the restricted interval.

Reflex: probabilities for a uniform variable are areas of rectangles, i.e. simple ratios of lengths — never integrate unless the question demands a proof.

WORKED EXAMPLE 20 Waiting for a bus in Bamenda

A bus arrives at a park in Bamenda at a random time between 8:00 and 8:20 each morning. The waiting time X (in minutes after 8:00) of a passenger who arrives at 8:00 is modelled by $X \sim U(0, 20)$.

1. Write down the probability density function of X .
2. Find the probability that the passenger waits more than 12 minutes.
3. Find $E(X)$ and $\text{Var}(X)$.
4. Given that the passenger has already waited 5 minutes, find the probability that he waits at least 10 minutes in total.

SOLUTION

Solution.

1. Here $a = 0$, $b = 20$, so

$$f(x) = \begin{cases} \frac{1}{20} & 0 \leq x \leq 20, \\ 0 & \text{otherwise.} \end{cases}$$

2. The favourable interval is $[12, 20]$, of length 8:

$$P(X > 12) = \frac{20 - 12}{20} = \frac{8}{20} = 0.4.$$

3. By the standard formulae:

$$E(X) = \frac{0 + 20}{2} = 10 \text{ minutes}, \quad \text{Var}(X) = \frac{(20 - 0)^2}{12} = \frac{400}{12} = \frac{100}{3} \approx 33.33 \text{ min}^2.$$

Hence $\sigma = \sqrt{100/3} \approx 5.77$ minutes.

4. We need $P(X \geq 10 \mid X \geq 5)$. Since $X \mid X \geq 5$ is uniform on $[5, 20]$ (length 15):

$$P(X \geq 10 \mid X \geq 5) = \frac{20 - 10}{20 - 5} = \frac{10}{15} = \frac{2}{3} \approx 0.667.$$

10.2 Method 2: Using the exponential distribution

METHOD Probabilities with $X \sim \text{Exp}(\lambda)$

1. Write the density $f(x) = \lambda e^{-\lambda x}$, $x \geq 0$, and the cumulative distribution function $F(x) = 1 - e^{-\lambda x}$.
2. Use the survival formula $P(X > t) = e^{-\lambda t}$ and $P(X \leq t) = 1 - e^{-\lambda t}$.
3. For an interval: $P(a < X < b) = e^{-\lambda a} - e^{-\lambda b}$.
4. Quote $E(X) = \frac{1}{\lambda}$, $\text{Var}(X) = \frac{1}{\lambda^2}$.
5. Use the **memoryless property**: $P(X > s + t \mid X > s) = P(X > t) = e^{-\lambda t}$.
6. If λ is unknown, find it from a given probability (e.g. solve $e^{-\lambda t_0} = p_0$ by taking natural logarithms, $\lambda = -\frac{\ln p_0}{t_0}$) or from a given mean ($\lambda = 1/E(X)$).

Reflex: check units — if X is in minutes, λ is "per minute".

WORKED EXAMPLE 21 Lifetime of a phone battery component

The lifetime T , in hours, of an electronic component follows an exponential distribution with mean 500 hours.

1. Write down the value of λ and the probability density function of T .

2. Find the probability that a component lasts more than 300 hours.
3. Find the probability that a component lasts between 200 and 600 hours.
4. Given that a component has already worked for 300 hours, find the probability that it works for at least 200 more hours. Comment.

SOLUTION

Solution.

1. $E(T) = \frac{1}{\lambda} = 500 \Rightarrow \lambda = \frac{1}{500} = 0.002$ per hour. Hence

$$f(t) = 0.002 e^{-0.002t}, \quad t \geq 0.$$

2. Using the survival formula:

$$P(T > 300) = e^{-0.002 \times 300} = e^{-0.6} \approx 0.5488.$$

- 3.

$$P(200 < T < 600) = e^{-0.002(200)} - e^{-0.002(600)} = e^{-0.4} - e^{-1.2} \approx 0.6703 - 0.3012 = 0.3691.$$

4. By the memoryless property:

$$P(T > 500 \mid T > 300) = P(T > 200) = e^{-0.002 \times 200} = e^{-0.4} \approx 0.6703.$$

Comment: the component "forgets" its age — the probability of surviving 200 extra hours is the same whether it is new or already 300 hours old. This is the defining feature of the exponential model.

10.3 Method 3: Linking the exponential and Poisson distributions

METHOD From Poisson events to exponential waiting times

1. If events occur randomly in time at rate λ per unit time, the number of events in an interval of length t is $N \sim \text{Po}(\lambda t)$.
2. The waiting time T between two consecutive events is $T \sim \text{Exp}(\lambda)$.
3. Key link: $\{T > t\}$ is exactly the event $\{N = 0 \text{ in } [0, t]\}$, so

$$P(T > t) = P(N = 0) = e^{-\lambda t}.$$

4. To answer questions, decide whether the variable asked about is a count (use Poisson) or a time (use exponential).

WORKED EXAMPLE 22 Calls at a call centre in Douala

Calls arrive at a customer-care centre at an average rate of 4 calls per hour, according to a Poisson process.

1. Find the probability that exactly 3 calls arrive in one hour.
2. Find the probability that no call arrives in a 30-minute period.
3. Let T be the waiting time, in hours, between two consecutive calls. State the distribution of T and find $P(T > 0.5)$.
4. Find the mean and variance of T .

SOLUTION

Solution.

1. $N \sim \text{Po}(4)$: $P(N = 3) = \frac{e^{-4}4^3}{3!} = \frac{64e^{-4}}{6} \approx 0.1954$.
2. In 30 minutes, the mean number of calls is $\lambda t = 4 \times 0.5 = 2$, so $N \sim \text{Po}(2)$:

$$P(N = 0) = e^{-2} \approx 0.1353.$$

3. $T \sim \text{Exp}(4)$ (rate 4 per hour). Then

$$P(T > 0.5) = e^{-4 \times 0.5} = e^{-2} \approx 0.1353.$$

Note this equals the answer to (b): "waiting more than 30 min for the next call" is the same event as "no call in the next 30 min" — the exponential–Poisson link in action.

4. $E(T) = \frac{1}{\lambda} = \frac{1}{4}$ hour = 15 minutes; $\text{Var}(T) = \frac{1}{\lambda^2} = \frac{1}{16} = 0.0625$ hour².

10.4 Method 4: Standardising and using the normal tables**METHOD** Probabilities for $X \sim N(\mu, \sigma^2)$

1. Standardise: $Z = \frac{X - \mu}{\sigma}$, where $Z \sim N(0, 1)$.
2. Rewrite the required probability in terms of Z and express it using $\Phi(z) = P(Z \leq z)$, with the properties

$$\Phi(-z) = 1 - \Phi(z), \quad P(Z > z) = 1 - \Phi(z), \quad P(a < Z < b) = \Phi(b) - \Phi(a).$$

3. Read Φ from the standard normal tables (4 decimal places).

4. For a "reverse" question (probability given, value wanted), read the table backwards: find the probability in the body of the table and read off z , then de-standardise with $x = \mu + \sigma z$.

Reflex: always draw a quick sketch of the bell curve and shade the required region before touching the tables.

KEY POINTS Special values of Φ to know

For the **central intervals**: $P(-1.96 < Z < 1.96) = 0.95$ and $P(-2.576 < Z < 2.576) = 0.99$. Thus the central 95% of any normal distribution lies in $\mu \pm 1.96\sigma$, and the central 99% in $\mu \pm 2.576\sigma$.

WORKED EXAMPLE 23 Masses of bags of rice

The mass X (in kg) of bags of rice sold in a market is modelled by $X \sim N(50, 1.44)$.

- Find $P(X < 51.2)$.
- Find $P(X > 48.5)$.
- Find $P(48.5 < X < 51.2)$.
- Find the mass m such that 90% of bags are heavier than m .

SOLUTION

Solution. Here $\mu = 50$, $\sigma = \sqrt{1.44} = 1.2$.

1. $z = \frac{51.2 - 50}{1.2} = 1.00$, so $P(X < 51.2) = \Phi(1.00) = 0.8413$.

2. $z = \frac{48.5 - 50}{1.2} = -1.25$, so

$$P(X > 48.5) = P(Z > -1.25) = \Phi(1.25) = 0.8944.$$

3.

$$P(48.5 < X < 51.2) = \Phi(1.00) - \Phi(-1.25) = 0.8413 - (1 - 0.8944) = 0.7357.$$

4. We need $P(X > m) = 0.90$, i.e. $P(X < m) = 0.10$. From the tables, $\Phi(z) = 0.10$ gives $z = -1.2816$. De-standardising:

$$m = \mu + \sigma z = 50 + 1.2(-1.2816) \approx 48.46 \text{ kg}.$$

10.5 Method 5: Finding unknown μ and/or σ

METHOD Determining parameters from given probabilities

1. Translate each given probability statement into an equation of the form $\Phi\left(\frac{x - \mu}{\sigma}\right) = p$.
2. Read the corresponding z -value from the tables (take care with signs: if $p < 0.5$, then z is negative).
3. Obtain linear equations in μ and σ : $x = \mu + z\sigma$.
4. With two statements, solve the simultaneous equations for μ and σ ; with one statement and one known parameter, solve directly for the other.

Reflex: write each equation as $x - \mu = z\sigma$ before substituting numbers — this avoids sign errors.

WORKED EXAMPLE 24 Filling bottles of palm oil

A machine fills bottles so that the volume dispensed is normally distributed. It is known that 10% of bottles contain less than 498 ml and 5% contain more than 506 ml.

1. Find the mean μ and standard deviation σ of the volume dispensed.
2. The firm wants only 1% of bottles to contain less than 500 ml. Keeping σ unchanged, to what value should the mean be adjusted?

SOLUTION

Solution.

1. From the tables: $\Phi(z) = 0.10 \Rightarrow z = -1.2816$; $\Phi(z) = 0.95 \Rightarrow z = 1.6449$. The two statements give:

$$498 = \mu - 1.2816\sigma \quad (1) \quad 506 = \mu + 1.6449\sigma \quad (2)$$

Subtracting (1) from (2): $8 = 2.9265\sigma \Rightarrow \sigma \approx 2.734$ ml. Then from (1): $\mu = 498 + 1.2816(2.734) \approx 501.50$ ml. So $\mu \approx 501.5$ ml and $\sigma \approx 2.73$ ml.

2. We require $P(X < 500) = 0.01$, i.e. $\Phi\left(\frac{500 - \mu}{2.734}\right) = 0.01$, so $\frac{500 - \mu}{2.734} = -2.3263$.

$$\mu = 500 + 2.3263 \times 2.734 \approx 506.36 \text{ ml.}$$

The mean should be set to about 506.4 ml.

10.6 Method 6: Continuity correction and normal approximations

METHOD Approximating binomial and Poisson distributions

1. Check validity. Binomial $B(n, p)$: use $N(np, npq)$ when n is large and $np > 5, nq > 5$. Poisson $Po(\lambda)$: use $N(\lambda, \lambda)$ when λ is large (say $\lambda > 10$).
2. Apply the continuity correction (the discrete variable X takes integer values):
 - $P(X \leq k) \rightarrow P(Y < k + 0.5); \quad P(X < k) \rightarrow P(Y < k - 0.5);$
 - $P(X \geq k) \rightarrow P(Y > k - 0.5); \quad P(X > k) \rightarrow P(Y > k + 0.5);$
 - $P(X = k) \rightarrow P(k - 0.5 < Y < k + 0.5).$
3. Standardise with $\mu = np$ (or λ) and $\sigma = \sqrt{npq}$ (or $\sqrt{\lambda}$), then use the normal tables.

WARNING Forgetting the continuity correction

Omitting the ± 0.5 is the most common error at the GCE. Without it, $P(X = k)$ would be 0 for a continuous variable! Always ask: "which integers are included?" and place the boundary halfway between included and excluded values.

WORKED EXAMPLE 25 Defective items in a factory

A machine produces items of which 8% are defective. A random sample of 200 items is taken. Using a suitable approximation, find the probability that

1. fewer than 12 items are defective;
2. exactly 16 items are defective.

SOLUTION

Solution. Let $X \sim B(200, 0.08)$. Then $np = 16 > 5$ and $nq = 184 > 5$, so the normal approximation is valid:

$$X \approx Y \sim N(16, 200 \times 0.08 \times 0.92) = N(16, 14.72), \quad \sigma = \sqrt{14.72} \approx 3.837.$$

1. "Fewer than 12" means $X \leq 11$, corrected to $Y < 11.5$:

$$Z = \frac{11.5 - 16}{3.837} = -1.173, \quad P(Y < 11.5) = \Phi(-1.173) = 1 - \Phi(1.173) \approx 1 - 0.8796 = 0.1204.$$

2. $P(X = 16)$ becomes $P(15.5 < Y < 16.5)$:

$$z_1 = \frac{15.5 - 16}{3.837} = -0.130, \quad z_2 = \frac{16.5 - 16}{3.837} = 0.130.$$

$$P = \Phi(0.130) - \Phi(-0.130) = 2\Phi(0.130) - 1 \approx 2(0.5517) - 1 = 0.1034.$$

10.7 Method 7: Combinations of independent normal variables

METHOD Sums and differences of independent normal variables

1. If $X \sim N(\mu_1, \sigma_1^2)$ and $Y \sim N(\mu_2, \sigma_2^2)$ are **independent**, then

$$aX + bY \sim N(a\mu_1 + b\mu_2, a^2\sigma_1^2 + b^2\sigma_2^2).$$

In particular $X - Y \sim N(\mu_1 - \mu_2, \sigma_1^2 + \sigma_2^2)$ — variances add even for a difference.

2. For a sum of n independent copies of X : $S_n \sim N(n\mu, n\sigma^2)$.
 3. Compute the new mean and variance, then standardise and use the tables as usual.

WARNING Variance of a difference

$\text{Var}(X - Y) = \sigma_1^2 + \sigma_2^2$, not $\sigma_1^2 - \sigma_2^2$. Subtracting variances is a classic GCE trap.

WORKED EXAMPLE 26 Heavier boy than girl?

At a school in Yaoundé, the masses of boys are $B \sim N(62, 25)$ kg and the masses of girls are $G \sim N(54, 16)$ kg, independently. A boy and a girl are chosen at random.

- Find the probability that the boy is heavier than the girl.
- Find the probability that their combined mass exceeds 125 kg.

SOLUTION

Solution.

1. Let $D = B - G$. Then

$$D \sim N(62 - 54, 25 + 16) = N(8, 41), \quad \sigma_D = \sqrt{41} \approx 6.403.$$

We want $P(D > 0)$:

$$Z = \frac{0 - 8}{6.403} = -1.249, \quad P(D > 0) = P(Z > -1.249) = \Phi(1.249) \approx 0.8942.$$

2. Let $S = B + G$. Then

$$S \sim N(62 + 54, 25 + 16) = N(116, 41).$$

$$Z = \frac{125 - 116}{6.403} = 1.406, \quad P(S > 125) = 1 - \Phi(1.406) \approx 1 - 0.9201 = 0.0799.$$

So the boy is heavier with probability about 0.894, and the combined mass exceeds 125 kg with probability about 0.080.

11 EXERCISES

11.1 I check my knowledge

EXERCISE 14 Uniform distribution basics

Let $X \sim U(2, 10)$.

1. Write down the probability density function $f(x)$.
2. Find $E(X)$ and $\text{Var}(X)$.
3. Calculate $\mathbb{P}(X > 6)$.
4. Determine the value of k such that $\mathbb{P}(X < k) = 0.75$.

EXERCISE 15 Exponential distribution: true/false with justification

The lifetime T (in months) of a phone battery follows an exponential distribution with mean 30. State whether each of the following is true or false, and justify your answer.

1. $\mathbb{P}(T > 36) = e^{-1.2}$.
2. Given that the battery has already lasted 24 months, the probability it lasts at least another 12 months is $\mathbb{P}(T > 12)$.
3. The median lifetime is $30 \ln 2$ months.
4. The variance of T is 900.

EXERCISE 16 Poisson–Exponential link

Customers arrive at an MTN service point according to a Poisson process with an average rate of 2 per minute. Let X be the number of arrivals in a 2-minute interval and Y be the time (in minutes) between two successive arrivals.

1. Write down the distribution of X and its parameter.
2. Write down the distribution of Y and its parameter.
3. Find $\mathbb{P}(X = 5)$.
4. Find $\mathbb{P}(Y > 1)$.

EXERCISE 17 Normal distribution: standardisation

The marks in a mock GCE examination are normally distributed with mean 52 and variance 144. Let X be the mark of a randomly chosen candidate.

1. Find the z -score corresponding to $X = 40$.
2. Use standard normal tables to find $\mathbb{P}(X < 40)$.
3. Find $\mathbb{P}(45 < X < 65)$.
4. Determine the mark exceeded by the top 10% of candidates.

11.2 I practise**EXERCISE 18** Finding parameters of a Normal distribution

A random variable $X \sim N(\mu, \sigma^2)$ satisfies $\mathbb{P}(X < 30) = 0.1587$ and $\mathbb{P}(X > 55) = 0.0228$. Determine the values of μ and σ .

EXERCISE 19 Quality control: adjusting the standard deviation

A machine fills sugar packets. The masses (in grams) are normally distributed with mean 505 and standard deviation 4. Packets with mass below 500 g are rejected.

1. Find the percentage of packets rejected.
2. The manufacturer wants to reduce the rejection rate to 1% by improving the machine (keeping $\mu = 505$). Find the required new standard deviation σ .

EXERCISE 20 Normal approximation to the Binomial with continuity correction

In a large batch of seeds, each seed germinates independently with probability 0.85. A random sample of 200 seeds is taken.

1. State the exact distribution of the number G of seeds that germinate.
2. Explain why a Normal approximation is appropriate.
3. Using a Normal approximation with continuity correction, estimate $\mathbb{P}(G < 160)$.

EXERCISE 21 Normal approximation to the Poisson distribution

The number of accidents per week on a certain highway follows a Poisson distribution with mean 12.

1. State the conditions under which a Normal approximation to the Poisson distribution is valid. Are they satisfied here?
2. Using a Normal approximation with continuity correction, estimate the probability that there are at most 8 accidents in a given week.

11.3 I prepare for the GCE**EXERCISE 22** Combining independent Normal variables: filled sacks] [12 marks

The mass of a coffee bag (in kg) is normally distributed with mean 50.2 and variance 0.16. The mass of an empty sack (in kg) is normally distributed with mean 0.8 and variance 0.04. The masses of bags and sacks are independent. A filled sack consists of one bag placed inside one sack.

1. Find the mean and variance of the mass of a filled sack. [3]
2. Find the probability that a randomly chosen filled sack weighs more than 51.5 kg. [4]
3. A quality control check selects 4 filled sacks at random. Find the probability that exactly 2 of them weigh more than 51.5 kg. [5]

EXERCISE 23 Two athletes: difference of Normal variables] [10 marks

Two athletes, A and B, run the 100 m. Their times (in seconds) are independent and normally distributed as follows:

$$T_A \sim N(11.2, 0.09), \quad T_B \sim N(11.5, 0.16).$$

In a given race, the athlete with the smaller time wins.

1. Find the distribution of $D = T_A - T_B$. [3]
2. Find the probability that athlete A beats athlete B. [4]
3. In a series of 5 independent races, find the probability that A wins at least 3 races. [3]

EXERCISE 24 Determining unknown parameters and using approximations] [14 marks

A factory produces metal rods. The length L (in cm) of a rod is normally distributed with mean μ and standard deviation σ . It is known that 2.5% of rods are shorter than 98.0 cm and 5% are longer than 102.0 cm.

1. Find the values of μ and σ . [6]
2. Rods are packed in boxes of 50. A box is accepted if at least 48 rods have lengths between 98.0 cm and 102.0 cm. Using a suitable approximation, estimate the probability that a randomly chosen box is accepted. [8]

12 SOLUTIONS TO THE EXERCISES

13 INTEGRATION EXERCISES AND CORRECTIONS: SPECIAL CONTINUOUS DISTRIBUTIONS

SOLUTION Exercise 1 — Uniform distribution basics

We have $X \sim U(2, 10)$, so $a = 2$ and $b = 10$, with $b - a = 8$.

1. The probability density function is

$$f(x) = \begin{cases} \frac{1}{b-a} = \frac{1}{8} & \text{if } 2 \leq x \leq 10, \\ 0 & \text{otherwise.} \end{cases}$$

2. Using the standard results $\mathbb{E}(X) = \frac{a+b}{2}$ and $\text{Var}(X) = \frac{(b-a)^2}{12}$:

$$\mathbb{E}(X) = \frac{2+10}{2} = 6, \quad \text{Var}(X) = \frac{(10-2)^2}{12} = \frac{64}{12} = \frac{16}{3} \approx 5.33.$$

3. For a uniform distribution, probabilities are ratios of lengths:

$$\mathbb{P}(X > 6) = \frac{10-6}{10-2} = \frac{4}{8} = \frac{1}{2}.$$

(This is also immediate from symmetry, since 6 is the mean.)

4. We require $\mathbb{P}(X < k) = \frac{k-2}{8} = 0.75$, so

$$k - 2 = 8 \times 0.75 = 6 \implies k = 8.$$

SOLUTION Exercise 2 — Exponential distribution: true/false with justification

The mean is 30 months, so the rate parameter is $\lambda = \frac{1}{30}$ and the cumulative distribution function is $\mathbb{P}(T \leq t) = 1 - e^{-t/30}$, hence $\mathbb{P}(T > t) = e^{-t/30}$.

1. True. $\mathbb{P}(T > 36) = e^{-36/30} = e^{-1.2}$.
2. True. By the **memoryless property** of the exponential distribution,

$$\mathbb{P}(T > 24 + 12 \mid T > 24) = \frac{\mathbb{P}(T > 36)}{\mathbb{P}(T > 24)} = \frac{e^{-36/30}}{e^{-24/30}} = e^{-12/30} = \mathbb{P}(T > 12).$$

3. True. The median m satisfies $\mathbb{P}(T > m) = \frac{1}{2}$, i.e. $e^{-m/30} = \frac{1}{2}$, so $\frac{m}{30} = \ln 2$ and $m = 30 \ln 2 \approx 20.8$ months.

4. True. For an exponential distribution, $\text{Var}(T) = \frac{1}{\lambda^2} = 30^2 = 900$ months².

All four statements are true.

SOLUTION Exercise 3 — Poisson–Exponential link

1. The number of arrivals in a 2-minute interval follows a Poisson distribution with mean $2 \times 2 = 4$:

$$X \sim \text{Po}(4).$$

2. The waiting time between two successive arrivals in a Poisson process of rate $\lambda = 2$ per minute is exponential with the same rate:

$$Y \sim \text{Exp}(2), \quad \mathbb{E}(Y) = \frac{1}{2} \text{ minute.}$$

$$3. \mathbb{P}(X = 5) = \frac{e^{-4} 4^5}{5!} = \frac{e^{-4} \times 1024}{120} \approx 0.1563.$$

$$4. \mathbb{P}(Y > 1) = e^{-\lambda \times 1} = e^{-2} \approx 0.1353.$$

Note the link: $\mathbb{P}(Y > 1)$ (no arrival during 1 minute) equals $\mathbb{P}(N = 0)$ where $N \sim \text{Po}(2)$, and indeed $\mathbb{P}(N = 0) = e^{-2}$.

SOLUTION Exercise 4 — Normal distribution: standardisation

Here $X \sim N(52, 144)$, so $\mu = 52$ and $\sigma = \sqrt{144} = 12$.

$$1. z = \frac{40 - 52}{12} = \frac{-12}{12} = -1.$$

$$2. \mathbb{P}(X < 40) = \mathbb{P}(Z < -1) = 1 - \Phi(1) = 1 - 0.8413 = 0.1587.$$

3. Standardising both bounds:

$$z_1 = \frac{45 - 52}{12} = -0.5833 \dots, \quad z_2 = \frac{65 - 52}{12} = 1.0833 \dots$$

Hence

$$\begin{aligned} \mathbb{P}(45 < X < 65) &= \Phi(1.0833 \dots) - \Phi(-0.5833 \dots) \\ &= \Phi(1.0833 \dots) - [1 - \Phi(0.5833 \dots)] \\ &\approx 0.8607 - (1 - 0.7202) = 0.8607 - 0.2798 = 0.5809. \end{aligned}$$

(Using $\Phi(1.08) = 0.8599$, $\Phi(0.58) = 0.7190$ gives ≈ 0.579 ; answers around 0.58 are accepted.)

4. We need the mark m with $\mathbb{P}(X > m) = 0.10$, i.e. $\mathbb{P}(X < m) = 0.90$. From the tables, $\Phi(z) = 0.90$ gives $z = 1.2816$. De-standardising:

$$m = \mu + z\sigma = 52 + 1.2816 \times 12 = 52 + 15.3792 \approx 67.4.$$

The top 10% of candidates score above approximately 67.4 marks.

SOLUTION Exercise 5 — Finding parameters of a Normal distribution

From $\mathbb{P}(X < 30) = 0.1587$: since $\Phi(1) = 0.8413$, we have $1 - \Phi(1) = 0.1587$, so the standardised value is -1 :

$$\frac{30 - \mu}{\sigma} = -1 \implies \mu - \sigma = 30. \quad (1)$$

From $\mathbb{P}(X > 55) = 0.0228$: since $\Phi(2) = 0.9772$, we have $1 - \Phi(2) = 0.0228$, so the standardised value is 2 :

$$\frac{55 - \mu}{\sigma} = 2 \implies \mu + 2\sigma = 55. \quad (2)$$

Subtracting (1) from (2): $3\sigma = 25$, so $\sigma = \frac{25}{3} \approx 8.33$. Then from (1):

$$\mu = 30 + \sigma = 30 + \frac{25}{3} = \frac{115}{3} \approx 38.33.$$

$$\boxed{\mu = \frac{115}{3} \approx 38.3, \quad \sigma = \frac{25}{3} \approx 8.33.}$$

Check: $\frac{30 - 38.33}{8.33} = -1$ ✓ and $\frac{55 - 38.33}{8.33} = 2$ ✓.

SOLUTION Exercise 6 — Quality control: adjusting the standard deviation

Let $M \sim N(505, 4^2)$.

1. Standardising the rejection limit 500:

$$z = \frac{500 - 505}{4} = -1.25.$$

$$\mathbb{P}(M < 500) = \mathbb{P}(Z < -1.25) = 1 - \Phi(1.25) = 1 - 0.8944 = 0.1056.$$

So about 10.56% of packets are rejected.

2. We now require $\mathbb{P}(M < 500) = 0.01$ with $\mu = 505$ and unknown σ . From the tables,

$\Phi(z) = 0.99$ gives $z = 2.3263$, so the standardised value of 500 must be -2.3263 :

$$\frac{500 - 505}{\sigma} = -2.3263 \implies \sigma = \frac{5}{2.3263} \approx 2.15 \text{ g.}$$

The machine must be improved so that the standard deviation is reduced from 4 g to about 2.15 g.

SOLUTION Exercise 7 — Normal approximation to the Binomial with continuity correction

- $G \sim B(200, 0.85)$.
- The conditions for a Normal approximation are $np > 5$ and $n(1 - p) > 5$ (some texts use $npq > 5$). Here

$$np = 200 \times 0.85 = 170 > 5, \quad n(1 - p) = 200 \times 0.15 = 30 > 5,$$

so the approximation $G \sim N(np, npq)$ is appropriate, with

$$\mu = 170, \quad \sigma^2 = 200 \times 0.85 \times 0.15 = 25.5, \quad \sigma = \sqrt{25.5} \approx 5.05.$$

- Applying the **continuity correction**, $\mathbb{P}(G < 160) = \mathbb{P}(G \leq 159)$ becomes $\mathbb{P}(Y < 159.5)$ where $Y \sim N(170, 25.5)$:

$$z = \frac{159.5 - 170}{5.05} = \frac{-10.5}{5.05} \approx -2.08.$$

$$\mathbb{P}(G < 160) \approx \mathbb{P}(Z < -2.08) = 1 - \Phi(2.08) = 1 - 0.9812 = 0.0188.$$

So $\mathbb{P}(G < 160) \approx 0.019$ (about 1.9%).

SOLUTION Exercise 8 — Normal approximation to the Poisson distribution

Let $X \sim \text{Po}(12)$.

- A Poisson distribution $\text{Po}(\lambda)$ can be approximated by $N(\lambda, \lambda)$ when λ is sufficiently large — the usual rule of thumb is $\lambda > 10$ (some texts say $\lambda \geq 15$). Here $\lambda = 12 > 10$, so the approximation is acceptable (though only moderately accurate).
- With the approximation $X \approx N(12, 12)$, we have $\sigma = \sqrt{12} \approx 3.464$. "At most 8 accidents" means $\mathbb{P}(X \leq 8)$; with the continuity correction this becomes $\mathbb{P}(Y < 8.5)$:

$$z = \frac{8.5 - 12}{\sqrt{12}} = \frac{-3.5}{3.464} \approx -1.01.$$

$$\mathbb{P}(X \leq 8) \approx \mathbb{P}(Z < -1.01) = 1 - \Phi(1.01) = 1 - 0.8438 = 0.1562.$$

So the probability of at most 8 accidents in a week is approximately 0.156.

SOLUTION Exercise 9 — Combining independent Normal variables: filled sacks

Let $B \sim N(50.2, 0.16)$ (bag) and $S \sim N(0.8, 0.04)$ (sack), independent, and $F = B + S$.

(a) For independent Normal variables, means add and variances add:

$$\mathbb{E}(F) = 50.2 + 0.8 = 51.0 \text{ kg}, \quad \text{Var}(F) = 0.16 + 0.04 = 0.20 \text{ kg}^2,$$

so $F \sim N(51.0, 0.20)$, with $\sigma_F = \sqrt{0.20} \approx 0.4472$.

Mark scheme: M1 for stating that means add; M1 for variances add (independence quoted); A1 for both values correct. [3 marks]

(b)

$$z = \frac{51.5 - 51.0}{\sqrt{0.20}} = \frac{0.5}{0.4472} \approx 1.12.$$

$$\mathbb{P}(F > 51.5) = 1 - \Phi(1.12) = 1 - 0.8686 = 0.1314.$$

Mark scheme: M1 standardising with their σ_F ; A1 for $z = 1.12$; M1 for using $1 - \Phi$; A1 for 0.1314 (accept 0.131–0.132). [4 marks]

(c) Let Y be the number of the 4 sacks weighing more than 51.5 kg. Then $Y \sim B(4, p)$ with $p = 0.1314$:

$$\begin{aligned} \mathbb{P}(Y = 2) &= \binom{4}{2} p^2 (1-p)^2 = 6 \times (0.1314)^2 \times (0.8686)^2 \\ &= 6 \times 0.017266 \times 0.754466 \approx 0.0781. \end{aligned}$$

Mark scheme: B1 for identifying the binomial model $B(4, 0.1314)$; M1 for the binomial formula with $n = 4, k = 2$; A1 for $\binom{4}{2} = 6$ used; M1 correct substitution; A1 for ≈ 0.078 . [5 marks]
Examiner's expectations: candidates must state the distribution of F explicitly, quote independence when adding variances, and carry their answer from (b) into (c) (error carried forward applies).

SOLUTION Exercise 10 — Two athletes: difference of Normal variables

(a) For independent Normal variables, $D = T_A - T_B$ is Normal with

$$\mathbb{E}(D) = 11.2 - 11.5 = -0.3, \quad \text{Var}(D) = 0.09 + 0.16 = 0.25,$$

(variances add even for a difference), so

$$D \sim N(-0.3, 0.25), \quad \sigma_D = 0.5.$$

Mark scheme: M1 for subtracting means; M1 for adding variances; A1 for $N(-0.3, 0.25)$. [3 marks]

(b) A beats B when $T_A < T_B$, i.e. $D < 0$:

$$z = \frac{0 - (-0.3)}{0.5} = 0.6.$$

$$\mathbb{P}(D < 0) = \Phi(0.6) = 0.7257.$$

Athlete A wins with probability about 0.726.

Mark scheme: B1 for translating "A wins" into $D < 0$; M1 standardising; A1 for $z = 0.6$; A1 for 0.7257. [4 marks]

(c) Let W be the number of races won by A out of 5: $W \sim B(5, 0.7257)$.

$$\begin{aligned}\mathbb{P}(W \geq 3) &= \mathbb{P}(W = 3) + \mathbb{P}(W = 4) + \mathbb{P}(W = 5) \\ &= \binom{5}{3}p^3q^2 + \binom{5}{4}p^4q + p^5, \quad p = 0.7257, q = 0.2743 \\ &= 10(0.3822)(0.07524) + 5(0.2773)(0.2743) + 0.2012 \\ &\approx 0.2876 + 0.3803 + 0.2012 = 0.8691.\end{aligned}$$

So $\mathbb{P}(W \geq 3) \approx 0.869$.

Mark scheme: B1 for the binomial model; M1 for summing the three correct terms; A1 for ≈ 0.869 . [3 marks]

Common trap: subtracting the variances in (a) — the variance of a difference of independent variables is always the sum of the variances.

SOLUTION Exercise 11 — Determining unknown parameters and using approximations

(a) We have $L \sim N(\mu, \sigma^2)$.

$\mathbb{P}(L < 98.0) = 0.025$: from the tables, $\Phi(z) = 0.975$ gives $z = 1.96$, so the standardised value of 98.0 is -1.96 :

$$\frac{98.0 - \mu}{\sigma} = -1.96 \implies \mu - 1.96\sigma = 98.0. \quad (1)$$

$\mathbb{P}(L > 102.0) = 0.05$: from the tables, $\Phi(z) = 0.95$ gives $z = 1.6449$:

$$\frac{102.0 - \mu}{\sigma} = 1.6449 \implies \mu + 1.6449\sigma = 102.0. \quad (2)$$

Subtracting (1) from (2):

$$(1.6449 + 1.96)\sigma = 4.0 \implies 3.6049\sigma = 4.0 \implies \sigma \approx 1.110 \text{ cm.}$$

Substituting in (1):

$$\mu = 98.0 + 1.96 \times 1.110 = 98.0 + 2.175 \approx 100.17 \text{ cm.}$$

$$\boxed{\mu \approx 100.2 \text{ cm,} \quad \sigma \approx 1.11 \text{ cm.}}$$

Mark scheme: B1 for $z = \pm 1.96$; B1 for $z = 1.6449$ (accept 1.645); M1 for setting up both standardised equations; M1 for solving the simultaneous system; A1 for σ ; A1 for μ . [6 marks]

(b) First find the probability p that one rod has length between 98.0 and 102.0 cm:

$$p = 1 - 0.025 - 0.05 = 0.925.$$

Let X be the number of rods (out of 50) with acceptable length: $X \sim B(50, 0.925)$. The box is accepted if $X \geq 48$.

Since $np = 46.25 > 5$ and $n(1-p) = 3.75 < 5$, the direct Normal approximation to X is poor. Instead, consider the number of defective rods $Y = 50 - X \sim B(50, 0.075)$. Here $np = 3.75$ is small, p is small and n large, so a Poisson approximation is suitable:

$$Y \approx \text{Po}(3.75).$$

The box is accepted when $X \geq 48$, i.e. $Y \leq 2$:


$$\begin{aligned} \mathbb{P}(Y \leq 2) &\approx e^{-3.75} \left(1 + 3.75 + \frac{3.75^2}{2} \right) \\ &= e^{-3.75} (1 + 3.75 + 7.03125) \\ &= 0.02352 \times 11.78125 \approx 0.2771. \end{aligned}$$

So the probability that a box is accepted is approximately 0.277.

Mark scheme: B1 for $p = 0.925$; B1 for identifying $X \sim B(50, 0.925)$; M1 for checking the Normal-approximation conditions and rejecting it ($nq < 5$); M1 for switching to defective rods $Y \sim B(50, 0.075)$; B1 for the Poisson approximation $\text{Po}(3.75)$ with justification; M1 for translating acceptance into $Y \leq 2$; M1 for the Poisson cumulative sum; A1 for ≈ 0.277 . [8 marks]

Examiner's expectations: the key insight is recognising that the Normal approximation fails here ($nq = 3.75 < 5$) and choosing the Poisson approximation to the binomial instead. Candidates who apply the Normal approximation blindly (even with continuity correction) score only the first two marks. Note also that no continuity correction is needed for the Poisson calculation since it remains a discrete distribution.

14 SUMMARY SHEET



Mind map of Special Continuous Distributions

FICHE BILAN L'essentiel à connaître pour l'examen

Key Definitions

- **Uniform (Rectangular) Distribution:** Continuous random variable X on $[a, b]$ with pdf $f(x) = \frac{1}{b-a}$ for $a \leq x \leq b$, zero elsewhere. $E(X) = \frac{a+b}{2}$, $\text{Var}(X) = \frac{(b-a)^2}{12}$.
- **Exponential Distribution:** Continuous random variable $X \geq 0$ with pdf $f(x) = \lambda e^{-\lambda x}$ ($x \geq 0$), $\lambda > 0$. $E(X) = \frac{1}{\lambda}$, $\text{Var}(X) = \frac{1}{\lambda^2}$. Memoryless property: $P(X > s+t \mid X > s) = P(X > t)$.
- **Normal Distribution:** $X \sim N(\mu, \sigma^2)$ with pdf $f(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$. Standard normal $Z \sim N(0, 1)$ with cdf $\Phi(z)$.
- **Standardisation:** $Z = \frac{X-\mu}{\sigma}$ transforms any $N(\mu, \sigma^2)$ to $N(0, 1)$.
- **Continuity Correction:** When approximating discrete (Binomial/Poisson) by Normal, add/subtract 0.5 to the discrete bound.

Laws, Formulas & Theorems to Know

- **Uniform:** $P(c \leq X \leq d) = \frac{d-c}{b-a}$ for $a \leq c < d \leq b$.
- **Exponential:** $P(X > x) = e^{-\lambda x}$, $P(X \leq x) = 1 - e^{-\lambda x}$.
- **Link Exponential–Poisson:** If events occur as Poisson process with rate λ , inter-arrival times are Exponential(λ).

- Normal: $\Phi(-z) = 1 - \Phi(z)$; $\Phi(0) = 0.5$; $\Phi(1.96) \approx 0.975$, $\Phi(2.576) \approx 0.995$.
- Central intervals: $P(\mu - 1.96\sigma < X < \mu + 1.96\sigma) = 0.95$; $P(\mu - 2.576\sigma < X < \mu + 2.576\sigma) = 0.99$.
- Linear combination: If $X_i \sim N(\mu_i, \sigma_i^2)$ independent, then $\sum a_i X_i \sim N(\sum a_i \mu_i, \sum a_i^2 \sigma_i^2)$.
- Normal approximation to Binomial: $X \sim B(n, p) \approx N(np, npq)$ if $np > 5$ and $nq > 5$.
- Normal approximation to Poisson: $X \sim Po(\lambda) \approx N(\lambda, \lambda)$ if $\lambda > 10$ (or > 15).

Methods

1. Uniform: Identify a, b ; compute probabilities as lengths; use $E(X)$, $\text{Var}(X)$ formulas.
2. Exponential: Identify λ (rate) or mean $1/\lambda$; use cdf $1 - e^{-\lambda x}$ or survival $e^{-\lambda x}$; apply memoryless property when needed.
3. Normal (standard): Use tables to find $\Phi(z)$ or z given probability; exploit symmetry $\Phi(-z) = 1 - \Phi(z)$.
4. Any Normal: Standardise $z = \frac{x-\mu}{\sigma}$; use tables; de-standardise $x = \mu + z\sigma$.
5. Finding μ and/or σ : Set up equations from given probabilities (e.g. $P(X < x_1) = p_1$, $P(X > x_2) = p_2$); standardise to get z -values; solve simultaneous equations.
6. Continuity correction: For $P(X \leq k)$ use $P(X < k + 0.5)$; for $P(X \geq k)$ use $P(X > k - 0.5)$; for $P(X = k)$ use $P(k - 0.5 < X < k + 0.5)$.
7. Independent Normals: For X, Y independent normals, $X \pm Y$ is normal with mean $\mu_X \pm \mu_Y$ and variance $\sigma_X^2 + \sigma_Y^2$.

Common Mistakes

- Forgetting that the uniform pdf is zero outside $[a, b]$; integrating over wrong limits.
- Confusing rate λ and mean $1/\lambda$ in exponential; using λ as mean.
- Misreading normal tables: using $\Phi(z)$ as tail probability instead of cumulative.
- Forgetting symmetry: $\Phi(-z) = 1 - \Phi(z)$.
- Omitting continuity correction when approximating discrete distributions.
- Using normal approximation when conditions ($np > 5$, $nq > 5$ or $\lambda > 10$) are not met.
- Incorrectly adding variances for independent normals: variances add, standard deviations do not.
- Mixing up z -values for central 95% (1.96) and 99% (2.576).

GCE Tips

- Always define the distribution and its parameters clearly at the start of a solution.
- Show the standardisation step explicitly: $z = \frac{x-\mu}{\sigma}$.

- When using tables, state the value you look up (e.g. $\Phi(1.645) = 0.95$).
- For inverse problems (find μ, σ), write the equations in terms of z-scores before solving.
- In hypothesis testing context (though not in this chapter), remember the critical values 1.645, 1.96, 2.576.
- Check that probabilities sum to 1 or lie in $[0, 1]$ as a sanity check.
- For exponential, the memoryless property is a favourite exam question; recognise wording like “given that it has lasted s units, probability it lasts t more”.
- When approximating binomial/poisson, state the conditions and apply continuity correction; lose marks if omitted.
- For combinations of independent normals, write the resulting mean and variance clearly.

**Signed • OnBuch+**ONBUCH+ TEACHING TEAM
FOLLOWS THE OFFICIAL MINESEC SYLLABUS**OnBuch+**