

PURE MATHEMATICS WITH MECHANICS • UPPER SIXTH

UPPER SIXTH — PURE MATHEMATICS WITH MECHANICS

CHAPTER 11

KINEMATICS OF PARTICLES IN A PLANE

SUGGESTED TIME: 9 PERIODS

IN THIS CHAPTER

This chapter extends the study of motion from a straight line to the plane: position, velocity and acceleration become vectors linked by differentiation and integration. It then applies these tools to projectile motion and vertical motion under gravity, two favourite themes of the GCE Advanced Level examination.

- By the end of this chapter I can represent the position of a particle in a plane by a vector $\mathbf{r}(t) = x(t)\mathbf{i} + y(t)\mathbf{j}$ and interpret its components.
- By the end of this chapter I can obtain velocity and acceleration vectors by differentiating the position vector with respect to time.
- By the end of this chapter I can recover velocity and position from acceleration (or velocity) by integration, using initial conditions to determine the constants.
- By the end of this chapter I can define a projectile and state the assumptions of the standard projectile model (constant g , no air resistance, point mass).
- By the end of this chapter I can derive and use the Cartesian equation of a projectile's trajectory and show it is a parabola.
- By the end of this chapter I can calculate the range, maximum height and time of flight of a projectile and use them to solve targeting problems.



DOWNLOAD ON- BUCH

Free • Google Play
AI tutor, past papers,
results



GO ONBUCH+

All signed courses, unlimited AI
tutor, PDF sheets — OnBuch+
tab in the app

THIS SIGNED COURSE CONTAINS

COURSE

complete, illustrated,
step by step

METHODS

and worked examples

EXERCISES

exam style + solutions

SUMMARY SHEET

the essentials to
remember



Signed • OnBuch+

ONBUCH+ TEACHING TEAM
FOLLOWS THE OFFICIAL MINESEC SYLLABUS

OnBuch+

CONTENTS

- | | |
|---|---|
| 1. Position, Velocity and Acceleration as Vectors in a Plane | with Projectiles |
| 2. From Acceleration back to Velocity and Position: Integration in Kinematics | 6. Free Fall, Vertical Projection and Motion Graphs |
| 3. Projectile Motion: Definition, Model and Equations | 7. Integration activity |
| 4. Range, Maximum Height and Time of Flight | 8. Methods and worked examples |
| 5. Applications and Problem-Solving | 9. Exercises |
| | 10. Solutions to the exercises |
| | 11. Summary sheet |

CHAPTER OBJECTIVES

- By the end of this chapter I can represent the position of a particle in a plane by a vector $\mathbf{r}(t) = x(t)\mathbf{i} + y(t)\mathbf{j}$ and interpret its components.
- By the end of this chapter I can obtain velocity and acceleration vectors by differentiating the position vector with respect to time.
- By the end of this chapter I can recover velocity and position from acceleration (or velocity) by integration, using initial conditions to determine the constants.
- By the end of this chapter I can define a projectile and state the assumptions of the standard projectile model (constant g , no air resistance, point mass).
- By the end of this chapter I can derive and use the Cartesian equation of a projectile's trajectory and show it is a parabola.
- By the end of this chapter I can calculate the range, maximum height and time of flight of a projectile and use them to solve targeting problems.
- By the end of this chapter I can solve applied projectile problems: clearing an obstacle, hitting a target, launching from a height, optimal angle of projection.
- By the end of this chapter I can analyse free fall and vertical projection (upwards or downwards) using the constant-acceleration equations.
- By the end of this chapter I can draw and interpret displacement-time, velocity-time and acceleration-time graphs for vertical motion.

PREREQUISITES

- Linear kinematics with constant acceleration (Lower Sixth): the suvat equations $v = u + at$, $s = ut + \frac{1}{2}at^2$, $v^2 = u^2 + 2as$.
- Differentiation and integration of polynomials, trigonometric and exponential functions (Pure Mathematics, Lower Sixth).
- Vectors in component form: i and j unit vectors, magnitude and direction of a vector.
- Trigonometry: sine, cosine, tangent, the identity $\sin(2\theta) = 2 \sin\theta \cos\theta$, solving simple trigonometric equations.
- Quadratic equations and the Cartesian equation of a parabola.

1 POSITION, VELOCITY AND ACCELERATION AS VECTORS IN A PLANE

During the inter-school football tournament at the Ahmadou Ahidjo Stadium in Yaoundé, a striker launches a free kick from 25 metres: will the ball clear the wall and dip under the crossbar? Meanwhile, on the Bamenda–Bafoussam road, a driver throws a mango peel out of a moving bus window and it lands far behind the vehicle. Even the nurse at the Mfoundi health centre wonders how high the water jet from a broken pipe will rise. All these everyday Cameroonian scenes obey the same mathematical laws: the kinematics of a particle moving in a plane. This chapter gives you the tools to describe, predict and even optimise such motions.

1.1 The Position Vector in a Plane

To describe the motion of a particle in a plane, we choose a fixed origin O and two perpendicular axes Ox and Oy with unit vectors $\vec{i}$ and $\vec{j}$. At any time t , the particle is at a point P . Its **position vector** is the vector $\vec{OP}$.

DEFINITION Position Vector

The position vector of a particle P at time t is

$$\vec{r}(t) = x(t)\vec{i} + y(t)\vec{j},$$

where $x(t)$ and $y(t)$ are the Cartesian coordinates of P at time t . The functions $x(t)$ and $y(t)$ are called the **parametric equations** of the motion. The variable t (time) is the parameter.

The set of all points occupied by the particle during its motion forms its **trajectory** (or path). The trajectory is a curve in the plane. Its **Cartesian equation** is obtained by eliminating the parameter t between $x = x(t)$ and $y = y(t)$.

METHOD Finding the Cartesian Equation of the Trajectory

1. Write $x = x(t)$ and $y = y(t)$.
2. If possible, solve one equation for t (express t in terms of x or y).
3. Substitute this expression into the other equation to obtain a relation between x and y only.
4. State any restrictions on x or y coming from the domain of t (usually $t \geq 0$).

EXAMPLE Parametric to Cartesian

A particle moves so that its position vector at time t (seconds) is

$$\vec{r}(t) = (3t^2)\vec{i} + (2t - 5)\vec{j} \quad (\text{metres}).$$

Find the Cartesian equation of its trajectory.

Solution We have $x = 3t^2$ and $y = 2t - 5$. Since $t \geq 0$, from $y = 2t - 5$ we get $t = \frac{y+5}{2}$. Substitute into x :

$$x = 3\left(\frac{y+5}{2}\right)^2 = \frac{3}{4}(y+5)^2.$$

This is a parabola opening in the positive x -direction with vertex at $(0, -5)$. Because $t \geq 0$, $y \geq -5$. The trajectory is the half-parabola $x = \frac{3}{4}(y+5)^2$, $y \geq -5$.

1.2 Velocity: Derivative of Position

The **velocity** of a particle is the rate of change of its position vector with respect to time.

DEFINITION Velocity Vector

If $\vec{r}(t) = x(t)\vec{i} + y(t)\vec{j}$, then the velocity vector is

$$\vec{v}(t) = \frac{d\vec{r}}{dt} = \frac{dx}{dt}\vec{i} + \frac{dy}{dt}\vec{j} = \dot{x}\vec{i} + \dot{y}\vec{j}.$$

The dot notation $\dot{x} = dx/dt$, $\dot{y} = dy/dt$ is often used for time derivatives.

The **speed** is the magnitude of the velocity vector:

$$v = |\vec{v}| = \sqrt{\dot{x}^2 + \dot{y}^2} \quad (\text{units: m s}^{-1}).$$

PROPERTY Velocity and the Trajectory

The velocity vector $\vec{v}(t)$ is always **tangent** to the trajectory at the point P and points in the direction of motion.

EXAMPLE Velocity, Speed and Direction

For the particle with $\vec{r}(t) = (3t^2)\vec{i} + (2t - 5)\vec{j}$:

1. Find $\vec{v}(t)$ and the acceleration $\vec{a}(t)$.

- Find the speed and direction of motion at $t = 2$ s.
- Determine when the particle is moving parallel to the x -axis and parallel to the y -axis.

Solution

- $\vec{v}(t) = \frac{d}{dt}(3t^2)\vec{i} + \frac{d}{dt}(2t - 5)\vec{j} = 6t\vec{i} + 2\vec{j}$. $\vec{a}(t) = \frac{d\vec{v}}{dt} = 6\vec{i} + 0\vec{j} = 6\vec{i}$.
- At $t = 2$: $\vec{v}(2) = 12\vec{i} + 2\vec{j}$. Speed $v = \sqrt{12^2 + 2^2} = \sqrt{148} = 2\sqrt{37} \approx 12.17 \text{ m s}^{-1}$. Direction: the velocity vector makes an angle θ with the x -axis where $\tan \theta = \frac{2}{12} = \frac{1}{6}$, so $\theta \approx 9.46^\circ$ above the horizontal.
- Parallel to x -axis $\Rightarrow$ vertical component zero: $\dot{y} = 2 = 0$, impossible. Never parallel to x -axis. Parallel to y -axis $\Rightarrow$ horizontal component zero: $\dot{x} = 6t = 0 \Rightarrow t = 0$. At $t = 0$, $\vec{v}(0) = 2\vec{j}$, moving vertically upward.

1.3 Acceleration: Derivative of Velocity

The **acceleration** is the rate of change of velocity with respect to time.

DEFINITION Acceleration Vector

If $\vec{v}(t) = \dot{x}\vec{i} + \dot{y}\vec{j}$, then the acceleration vector is

$$\vec{a}(t) = \frac{d\vec{v}}{dt} = \frac{d^2\vec{r}}{dt^2} = \ddot{x}\vec{i} + \ddot{y}\vec{j}.$$

Units: m s^{-2} .

Unlike velocity, acceleration is not generally tangent to the path. It can be decomposed into tangential and normal components, but in Cartesian coordinates we simply work with its x and y components.

1.4 Worked Examples and GCE-Style Problems**WORKED EXAMPLE 1 GCE Style: Perpendicular Velocity**

A particle moves in the xy -plane with position vector

$$\vec{r}(t) = (t^3 - 3t)\vec{i} + (2t^2 - 4)\vec{j}, \quad t \geq 0.$$

Find the time(s) when the velocity of the particle is perpendicular to the vector $\vec{u} = 2\vec{i} - \vec{j}$.

SOLUTION

Solution Velocity: $\vec{v}(t) = \frac{d\vec{r}}{dt} = (3t^2 - 3)\vec{i} + 4t\vec{j}$. Two vectors are perpendicular iff their scalar (dot) product is zero. $\vec{v}(t) \cdot \vec{u} = (3t^2 - 3)(2) + (4t)(-1) = 6t^2 - 6 - 4t = 0$. $6t^2 - 4t - 6 = 0 \implies 3t^2 - 2t - 3 = 0$. $t = \frac{2 \pm \sqrt{4 + 36}}{6} = \frac{2 \pm \sqrt{40}}{6} = \frac{1 \pm \sqrt{10}}{3}$. Since $t \geq 0$, we take $t = \frac{1 + \sqrt{10}}{3} \approx 1.39$ s.

WORKED EXAMPLE 2 Comprehensive Kinematics

A particle P has position vector $\vec{r}(t) = (4t - t^2)\vec{i} + (t^2 - 2t)\vec{j}$ metres, $t \geq 0$.


1. Find the Cartesian equation of the path of P .
2. Find the velocity and acceleration vectors.
3. At what time is the speed of P minimum? Find this minimum speed.

SOLUTION


Solution

1. $x = 4t - t^2$, $y = t^2 - 2t$. Add: $x + y = 2t \implies t = \frac{x+y}{2}$. Substitute into y : $y = \left(\frac{x+y}{2}\right)^2 - 2\left(\frac{x+y}{2}\right)$. $4y = (x + y)^2 - 4(x + y) = x^2 + 2xy + y^2 - 4x - 4y$. $x^2 + 2xy + y^2 - 4x - 8y = 0$. This is the Cartesian equation (a parabola).
2. $\vec{v} = (4 - 2t)\vec{i} + (2t - 2)\vec{j}$. $\vec{a} = -2\vec{i} + 2\vec{j}$ (constant acceleration).
3. Speed squared: $v^2 = (4 - 2t)^2 + (2t - 2)^2 = 4(2 - t)^2 + 4(t - 1)^2 = 4[(t - 2)^2 + (t - 1)^2] = 4[2t^2 - 6t + 5]$. Minimise v^2 : derivative $8(2t - 3) = 0 \implies t = 1.5$ s. Minimum speed: $v_{\min} = \sqrt{4[2(1.5)^2 - 6(1.5) + 5]} = \sqrt{4[4.5 - 9 + 5]} = \sqrt{4 \times 0.5} = \sqrt{2} \approx 1.41 \text{ m s}^{-1}$.

1.5 Graphical Representation



Position vector $\vec{r}(t)$ from origin to point P , and velocity vector $\vec{v}(t)$ tangent to the trajectory at P .



Parametric plot of $\vec{r}(t) = (3t^2, 2t - 5)$ with velocity vectors (green arrows) at three instants.

1.6 Common Mistakes and Warnings

WARNING Speed vs. Derivative of Distance

- Speed is $|\vec{v}| = \sqrt{\dot{x}^2 + \dot{y}^2}$. It is a scalar.
- The derivative of the distance from the origin, $\frac{d}{dt}|\vec{r}|$, is not the speed (unless motion is purely radial).
- Never compute speed as $\sqrt{x^2 + y^2}$ or as $|\dot{x}| + |\dot{y}|$. Always use the Pythagorean combination of the components of $\vec{v}$.
- When asked for “direction of motion”, give the angle the velocity vector makes with the positive x -axis (or a bearing), not the angle of the position vector.

1.7 Key Points to Remember**KEY POINTS** Summary: Kinematics in a Plane

- Position: $\vec{r}(t) = x(t)\vec{i} + y(t)\vec{j}$ (metres).
- Velocity: $\vec{v}(t) = \dot{\vec{r}} = \dot{x}\vec{i} + \dot{y}\vec{j}$ (m s^{-1}). Tangent to path.
- Speed: $v = |\vec{v}| = \sqrt{\dot{x}^2 + \dot{y}^2}$.
- Acceleration: $\vec{a}(t) = \dot{\vec{v}} = \ddot{\vec{r}} = \ddot{x}\vec{i} + \ddot{y}\vec{j}$ (m s^{-2}).
- Trajectory: Eliminate t from $x = x(t)$, $y = y(t)$ to get $f(x, y) = 0$.
- Perpendicular condition: $\vec{v} \perp \vec{u} \iff \vec{v} \cdot \vec{u} = 0$.
- Parallel to axes: Parallel to x -axis $\iff \dot{y} = 0$; parallel to y -axis $\iff \dot{x} = 0$.

EXERCISE 1 Practice

1. A particle moves with $\vec{r}(t) = (2t^2 + 3)\vec{i} + (t^3 - 4t)\vec{j}$. Find (a) $\vec{v}(t)$ and $\vec{a}(t)$, (b) the speed at $t = 1$, (c) the time when the velocity is parallel to $\vec{i} + 2\vec{j}$.
2. The position vector of a particle is $\vec{r}(t) = (5 \cos t)\vec{i} + (5 \sin t)\vec{j}$, $t \geq 0$. (a) Show that the path is a circle. (b) Find $\vec{v}$ and $\vec{a}$. (c) Prove that $\vec{v} \cdot \vec{r} = 0$ and $\vec{a} \cdot \vec{v} = 0$. Interpret geometrically.
3. A particle is projected from the origin with velocity $\vec{u} = 10\vec{i} + 20\vec{j}$ m/s and experiences a constant acceleration $\vec{a} = -2\vec{i} - 10\vec{j}$ m/s². Find (a) $\vec{r}(t)$, (b) the time when the particle crosses the x -axis again, (c) the maximum height reached.

SOLUTION Exercises 1–3**1. Exercise 1:**

(a) $\vec{v} = 4t\vec{i} + (3t^2 - 4)\vec{j}$, $\vec{a} = 4\vec{i} + 6t\vec{j}$.

(b) At $t = 1$: $\vec{v} = 4\vec{i} - 1\vec{j}$, speed $= \sqrt{16 + 1} = \sqrt{17}$ m/s.

(c) Parallel to $\vec{i} + 2\vec{j} \Rightarrow \frac{4t}{1} = \frac{3t^2 - 4}{2} \Rightarrow 8t = 3t^2 - 4 \Rightarrow 3t^2 - 8t - 4 = 0 \Rightarrow (3t + 2)(t - 2) = 0 \Rightarrow t = 2$ s ($t \geq 0$).

2. Exercise 2:

(a) $x^2 + y^2 = 25 \cos^2 t + 25 \sin^2 t = 25$, circle radius 5 centred at origin.

(b) $\vec{v} = -5 \sin t \vec{i} + 5 \cos t \vec{j}$, $\vec{a} = -5 \cos t \vec{i} - 5 \sin t \vec{j} = -\vec{r}$.

(c) $\vec{v} \cdot \vec{r} = -25 \sin t \cos t + 25 \cos t \sin t = 0 \Rightarrow \vec{v} \perp \vec{r}$ (velocity tangent to circle). $\vec{a} \cdot \vec{v} = 25 \cos t \sin t - 25 \sin t \cos t = 0 \Rightarrow \vec{a} \perp \vec{v}$ (acceleration radial, no tangential component, constant speed).

3. Exercise 3:

(a) $\vec{v}(t) = \vec{u} + \vec{a}t = (10 - 2t)\vec{i} + (20 - 10t)\vec{j}$. $\vec{r}(t) = \int \vec{v} dt = (10t - t^2)\vec{i} + (20t - 5t^2)\vec{j}$.


(b) Cross x -axis $\Rightarrow y = 0 \Rightarrow 20t - 5t^2 = 0 \Rightarrow t(4 - t) = 0 \Rightarrow t = 4$ s (excluding $t = 0$).

(c) Max height when $v_y = 0 \Rightarrow 20 - 10t = 0 \Rightarrow t = 2$ s. $y_{\max} = 20(2) - 5(4) = 20$ m.

2 FROM ACCELERATION BACK TO VELOCITY AND POSITION: INTEGRATION IN KINEMATICS

In the previous section we travelled down the kinematic chain: starting from the position vector $\vec{r}(t)$ we differentiated once to obtain the velocity $\vec{v}(t) = \dot{\vec{r}}(t)$, and once more to obtain the acceleration $\vec{a}(t) = \ddot{\vec{r}}(t)$. In real life, however, the quantity we most often know is the acceleration, because acceleration is produced by forces (Newton's second law, $\vec{F} = m\vec{a}$). A driver on the Yaoundé–Douala highway who presses the accelerator pedal is controlling the acceleration of the car; the speed and the position are consequences. This section is therefore devoted to the inverse problem: **given the acceleration, recover the velocity and the position by integration.**

2.1 The Inverse Problem: Integration as the Reverse of Differentiation



The kinematic chain. Differentiation goes from position to acceleration; integration, together with initial conditions, takes us back.

Since $\frac{d\vec{v}}{dt} = \vec{a}$, the velocity is an antiderivative of the acceleration, and since $\frac{d\vec{r}}{dt} = \vec{v}$, the position is an antiderivative of the velocity. Integration is performed **component by component**: if $\vec{a}(t) = a_x(t)\vec{i} + a_y(t)\vec{j}$, then

$$\vec{v}(t) = \left(\int a_x(t) dt \right) \vec{i} + \left(\int a_y(t) dt \right) \vec{j}, \quad \vec{r}(t) = \left(\int v_x(t) dt \right) \vec{i} + \left(\int v_y(t) dt \right) \vec{j}.$$

This is exactly the reverse of the component-wise differentiation of the previous section: the unit vectors $\vec{i}$ and $\vec{j}$ are constant, so they behave as ordinary constants under integration.

PROPERTY Fundamental integration formulas

If a particle moves in a plane with acceleration $\vec{a}(t)$, velocity $\vec{v}(t)$ and position $\vec{r}(t)$, then

$$\vec{v}(t) = \vec{u} + \int_0^t \vec{a}(\tau) d\tau, \quad \vec{r}(t) = \vec{r}_0 + \int_0^t \vec{v}(\tau) d\tau,$$

where $\vec{u} = \vec{v}(0)$ is the **initial velocity** and $\vec{r}_0 = \vec{r}(0)$ is the **initial position**. Equivalently, the change of velocity over $[t_1, t_2]$ is $\vec{v}(t_2) - \vec{v}(t_1) = \int_{t_1}^{t_2} \vec{a} dt$ and the displacement is $\vec{r}(t_2) - \vec{r}(t_1) = \int_{t_1}^{t_2} \vec{v} dt$.

Justification. By the Fundamental Theorem of Calculus, if $\dot{\vec{v}} = \vec{a}$ then $\int_0^t \vec{a}(\tau) d\tau = \vec{v}(t) - \vec{v}(0) = \vec{v}(t) - \vec{u}$, which gives the first formula. Applying the same argument to $\dot{\vec{r}} = \vec{v}$ gives the second. ■

WARNING The constants of integration

Each integration introduces a constant vector $\vec{C}$. If you write $\vec{v}(t) = \int \vec{a}(t) dt$ and forget the constant, your answer is wrong even if the integration itself is perfect. Forgetting initial

conditions is the most common source of lost marks in GCE kinematics questions. Always determine each constant immediately after each integration, before moving on.

METHOD From $\vec{a}(t)$ to $\vec{r}(t)$

1. Integrate $\vec{a}(t)$ component-wise to get $\vec{v}(t) + \vec{C}_1$.
2. Use the initial velocity $\vec{v}(0) = \vec{u}$ to fix $\vec{C}_1$.
3. Integrate the (now complete) $\vec{v}(t)$ component-wise to get $\vec{r}(t) + \vec{C}_2$.
4. Use the initial position $\vec{r}(0) = \vec{r}_0$ to fix $\vec{C}_2$.
5. Answer the question asked: position, velocity, speed $|\vec{v}|$, or displacement at a given time.

2.2 Special Case: Constant Acceleration and the Vector suvat Equations

Suppose $\vec{a}$ is constant. Then, with $\vec{v}(0) = \vec{u}$ and $\vec{r}(0) = \vec{r}_0$:

$$\vec{v}(t) = \vec{u} + \int_0^t \vec{a} d\tau = \vec{u} + \vec{a}t,$$

$$\vec{r}(t) = \vec{r}_0 + \int_0^t (\vec{u} + \vec{a}\tau) d\tau = \vec{r}_0 + \vec{u}t + \frac{1}{2}\vec{a}t^2.$$

PROPERTY Vector suvat equations for constant acceleration

For a particle moving with constant acceleration $\vec{a}$,

$$\vec{v} = \vec{u} + \vec{a}t \quad \text{and} \quad \vec{r} = \vec{r}_0 + \vec{u}t + \frac{1}{2}\vec{a}t^2$$

These are the vector forms of the familiar suvat equations; taking the $\vec{i}$ and $\vec{j}$ components separately reproduces $v = u + at$ and $s = ut + \frac{1}{2}at^2$ in each direction. The derivation by integration, as above, is examinable.

WARNING Constant acceleration only!

The formulas $\vec{v} = \vec{u} + \vec{a}t$ and $\vec{r} = \vec{r}_0 + \vec{u}t + \frac{1}{2}\vec{a}t^2$ are valid only when $\vec{a}$ does not depend on time. If $\vec{a} = 6t\vec{i} - 2\vec{j}$, for example, you must integrate; substituting into suvat is a serious error.

2.3 Worked Examples with Variable Acceleration

WORKED EXAMPLE 3 Integrating a time-dependent acceleration

A particle P moves in a plane. At time t seconds ($t \geq 0$) its acceleration is

$$\vec{a}(t) = 6t\vec{i} - 2\vec{j} \quad (\text{m s}^{-2}).$$

Initially P is at the origin O and has velocity $\vec{u} = (2\vec{i} + \vec{j}) \text{ m s}^{-1}$.

- Find the velocity and the position of P at time t .
- Find the speed of P when $t = 3$.
- Show that P crosses the x -axis again and find the time at which this happens.

SOLUTION

Solution. (a) Velocity. Integrate $\vec{a}$ component-wise:

$$\vec{v}(t) = \int (6t\vec{i} - 2\vec{j}) dt = 3t^2\vec{i} - 2t\vec{j} + \vec{C}_1.$$

At $t = 0$: $\vec{v}(0) = \vec{C}_1 = 2\vec{i} + \vec{j}$. Hence

$$\vec{v}(t) = (3t^2 + 2)\vec{i} + (1 - 2t)\vec{j} \quad (\text{m s}^{-1}).$$

Position. Integrate again:

$$\vec{r}(t) = \int \vec{v}(t) dt = (t^3 + 2t)\vec{i} + (t - t^2)\vec{j} + \vec{C}_2.$$

At $t = 0$: $\vec{r}(0) = \vec{C}_2 = \vec{O}$ (the particle starts at O). Hence

$$\vec{r}(t) = (t^3 + 2t)\vec{i} + (t - t^2)\vec{j} \quad (\text{m}).$$

(b) At $t = 3$: $\vec{v}(3) = (27 + 2)\vec{i} + (1 - 6)\vec{j} = 29\vec{i} - 5\vec{j}$, so the speed is

$$|\vec{v}(3)| = \sqrt{29^2 + (-5)^2} = \sqrt{856} \approx 29.3 \text{ m s}^{-1}.$$

(c) P is on the x -axis when the $\vec{j}$ -component of $\vec{r}$ is zero:

$$t - t^2 = t(1 - t) = 0 \implies t = 0 \text{ or } t = 1.$$

So P starts on the axis and crosses it again at $t = 1$ s, at the point $\vec{r}(1) = 3\vec{i}$, i.e. 3 m from O along the x -axis. ■

WORKED EXAMPLE 4 When does the particle return to the origin?

A particle leaves the origin O with velocity $(4\vec{i} + 2\vec{j}) \text{ m s}^{-1}$ and moves with acceleration $\vec{a}(t) = (-2\vec{i} - \vec{j}) \text{ m s}^{-2}$ (constant). Find the time at which it returns to O , and its velocity at that instant.

SOLUTION

Solution. Since $\vec{a}$ is constant, $\vec{v} = \vec{u} + \vec{a}t = (4 - 2t)\vec{i} + (2 - t)\vec{j}$, and

$$\vec{r}(t) = \vec{u}t + \frac{1}{2}\vec{a}t^2 = (4t - t^2)\vec{i} + \left(2t - \frac{1}{2}t^2\right)\vec{j}.$$

Return to O requires both components to vanish simultaneously:

$$t(4 - t) = 0 \quad \text{and} \quad t\left(2 - \frac{1}{2}t\right) = 0.$$

The first equation gives $t = 0$ or $t = 4$; the second gives $t = 0$ or $t = 4$. The common non-zero solution is $t = 4 \text{ s}$. Then $\vec{v}(4) = (4 - 8)\vec{i} + (2 - 4)\vec{j} = -4\vec{i} - 2\vec{j} \text{ m s}^{-1}$: the particle returns with exactly the opposite velocity, as symmetry suggests. ■


WARNING Both components must vanish together

To find when a particle passes through the origin (or any given point), set each component of $\vec{r}(t)$ equal to the required value and look for a value of t satisfying all the equations at once. A time that zeros only one component corresponds to crossing an axis, not to passing through O .

2.4 Displacement from Velocity: Definite Integrals, and Distance versus Displacement

Often the velocity is given directly and we only need the change of position between two times. No constant of integration is then required, because we use a definite integral:

$$\text{displacement from } t_1 \text{ to } t_2 : \quad \vec{r}(t_2) - \vec{r}(t_1) = \int_{t_1}^{t_2} \vec{v}(t) dt.$$



The displacement between t_1 and t_2 equals the (signed) area under the velocity–time curve.

WARNING Displacement is not distance

The displacement is $\int_{t_1}^{t_2} \vec{v} dt$: regions where a component of $\vec{v}$ is negative subtract from the integral. The **total distance travelled** requires integrating the speed:

$$\text{distance} = \int_{t_1}^{t_2} |\vec{v}(t)| dt,$$

splitting the interval wherever the relevant component of $\vec{v}$ changes sign. For one-dimensional motion with velocity $v(t)$, $\text{distance} = \int_{t_1}^{t_2} |v(t)| dt$: integrate v where $v > 0$ and $-v$ where $v < 0$, then add.

EXAMPLE Displacement versus distance

A particle moves along a straight line with velocity $v(t) = 3t - 6 \text{ m s}^{-1}$. Find (a) the displacement and (b) the distance travelled between $t = 0$ and $t = 4 \text{ s}$.

Solution. (a) Displacement $= \int_0^4 (3t - 6) dt = \left[\frac{3}{2}t^2 - 6t \right]_0^4 = (24 - 24) - 0 = 0 \text{ m}$: the particle ends where it started.

(b) v changes sign at $t = 2$. On $[0, 2]$, $v < 0$; on $[2, 4]$, $v > 0$.

$$\text{distance} = \int_0^2 (6 - 3t) dt + \int_2^4 (3t - 6) dt = \left[6t - \frac{3}{2}t^2 \right]_0^2 + \left[\frac{3}{2}t^2 - 6t \right]_2^4 = 6 + 6 = 12 \text{ m}.$$

The two answers differ completely: zero displacement, but 12 m actually travelled.

2.5 GCE–Style Synthesis

WORKED EXAMPLE 5 Full synthesis problem

A particle P leaves the origin O at time $t = 0$ with velocity $(3\vec{i} + 5\vec{j}) \text{ m s}^{-1}$. At time t seconds its acceleration is

$$\vec{a}(t) = (2t\vec{i} - 4\vec{j}) \text{ m s}^{-2}.$$

- (a) Find the position vector of P at time t .
 (b) Find the speed of P when $t = 2$.
 (c) Find the displacement of P between $t = 1$ and $t = 3$.

SOLUTION

Solution. (a) Integrate the acceleration:

$$\vec{v}(t) = t^2\vec{i} - 4t\vec{j} + \vec{C}_1; \quad \vec{v}(0) = \vec{C}_1 = 3\vec{i} + 5\vec{j},$$

so $\vec{v}(t) = (t^2 + 3)\vec{i} + (5 - 4t)\vec{j} \text{ m s}^{-1}$. Integrate again:

$$\vec{r}(t) = \left(\frac{1}{3}t^3 + 3t\right)\vec{i} + (5t - 2t^2)\vec{j} + \vec{C}_2; \quad \vec{r}(0) = \vec{C}_2 = \vec{0}.$$

$$\therefore \vec{r}(t) = \left(\frac{1}{3}t^3 + 3t\right)\vec{i} + (5t - 2t^2)\vec{j} \text{ m}.$$

(b) $\vec{v}(2) = (4 + 3)\vec{i} + (5 - 8)\vec{j} = 7\vec{i} - 3\vec{j}$, so the speed is

$$|\vec{v}(2)| = \sqrt{7^2 + 3^2} = \sqrt{58} \approx 7.62 \text{ m s}^{-1}.$$

(c) Using the definite integral (no constants needed):

$$\vec{r}(3) - \vec{r}(1) = \int_1^3 \vec{v}(t) dt = \left[\left(\frac{1}{3}t^3 + 3t\right)\vec{i} + (5t - 2t^2)\vec{j} \right]_1^3.$$

At $t = 3$: $(9 + 9)\vec{i} + (15 - 18)\vec{j} = 18\vec{i} - 3\vec{j}$. At $t = 1$: $\frac{10}{3}\vec{i} + 3\vec{j}$.

$$\text{Displacement} = \left(18 - \frac{10}{3}\right)\vec{i} + (-3 - 3)\vec{j} = \frac{44}{3}\vec{i} - 6\vec{j} \text{ m}.$$

■

METHOD Exam technique for integration problems

1. Write vectors consistently in $\vec{i}, \vec{j}$ (or column) form throughout the whole solution — never mix the two.
2. Keep the units (m s^{-2} , m s^{-1} , m) at every line; a unit error often reveals a missing integration step.

3. Fix each constant of integration as soon as it appears, quoting the initial condition explicitly: “when $t = 0$, $\vec{v} = \dots$ ”.
4. For “speed”, remember to take the modulus $|\vec{v}| = \sqrt{v_x^2 + v_y^2}$; for “velocity”, leave the vector.
5. Check your final $\vec{r}(t)$ by differentiating twice: you must recover the given $\vec{a}(t)$.

KEY POINTS Key points

- $\vec{v}(t) = \vec{u} + \int_0^t \vec{a} \, d\tau$ and $\vec{r}(t) = \vec{r}_0 + \int_0^t \vec{v} \, d\tau$; integration is done component by component.
- Each integration produces a constant vector, fixed by an initial condition.
- Constant acceleration gives the vector suvat equations $\vec{v} = \vec{u} + \vec{a}t$, $\vec{r} = \vec{r}_0 + \vec{u}t + \frac{1}{2}\vec{a}t^2$ — valid only for constant $\vec{a}$.
- Displacement = $\int_{t_1}^{t_2} \vec{v} \, dt$ (signed area under the v - t curve); distance travelled = $\int_{t_1}^{t_2} |\vec{v}| \, dt$, splitting at sign changes.

EXERCISE 2 Practice

1. A particle starts from rest at the point with position vector $(2\vec{i} - \vec{j})$ m and moves with acceleration $\vec{a}(t) = (4\vec{i} + 6t\vec{j}) \text{ m s}^{-2}$. Find its velocity and position at time t , and its distance from the origin when $t = 2$.
2. A particle moves with velocity $\vec{v}(t) = (2t - 4)\vec{i} + 3\vec{j} \text{ m s}^{-1}$. Find (a) its displacement between $t = 0$ and $t = 5$; (b) the total distance travelled in the x -direction over the same interval.
3. A particle leaves O with velocity $\vec{u} = (\vec{i} + 4\vec{j}) \text{ m s}^{-1}$ and has constant acceleration $\vec{a} = (2\vec{i} - 2\vec{j}) \text{ m s}^{-2}$. Show that it crosses the x -axis again at $t = 4$ s and find its speed at that instant.

SOLUTION Exercise 1

$\vec{v}(t) = 4t\vec{i} + 3t^2\vec{j}$ (constant $\vec{0}$ since it starts from rest). $\vec{r}(t) = (2 + 2t^2)\vec{i} + (t^3 - 1)\vec{j}$ (constant $(2\vec{i} - \vec{j})$). At $t = 2$: $\vec{r} = 10\vec{i} + 7\vec{j}$, so distance from O is $\sqrt{100 + 49} = \sqrt{149} \approx 12.2$ m.

SOLUTION Exercise 2

(a) Displacement $= \int_0^5 \vec{v} dt = [t^2 - 4t]_0^5 \vec{i} + [3t]_0^5 \vec{j} = 5\vec{i} + 15\vec{j} \text{ m}$. (b) $v_x = 2t - 4$ changes sign at $t = 2$: distance in the x -direction $= \int_0^2 (4 - 2t) dt + \int_2^5 (2t - 4) dt = 4 + 9 = 13 \text{ m}$.

SOLUTION Exercise 3

$\vec{r}(t) = \vec{u}t + \frac{1}{2}\vec{a}t^2 = (t + t^2)\vec{i} + (4t - t^2)\vec{j}$. On the x -axis the $\vec{j}$ -component vanishes: $t(4 - t) = 0$ gives $t = 0$ (start) or $t = 4 \text{ s}$, as required. Then $\vec{v}(4) = \vec{u} + 4\vec{a} = (\vec{i} + 4\vec{j}) + (8\vec{i} - 8\vec{j}) = 9\vec{i} - 4\vec{j}$, and the speed is $\sqrt{81 + 16} = \sqrt{97} \approx 9.85 \text{ m s}^{-1}$.

DID YOU KNOW? Why this section matters for what comes next

Projectile motion — the subject of the next sections — is nothing more than the constant-acceleration case of what you have just learned, with $\vec{a} = -g\vec{j}$. Every formula for the range, the greatest height and the time of flight of a projectile is obtained by exactly the two integrations you have practised here. Master this section, and the rest of the chapter becomes a routine application.

3 PROJECTILE MOTION: DEFINITION, MODEL AND EQUATIONS

In the previous section we saw how to reconstruct the velocity and position of a particle from its acceleration by integration. We now apply those ideas to one of the most important and beautiful problems in mechanics: the motion of a **projectile**. A projectile is any object that is launched into the air and then moves freely under the influence of gravity alone. Think of a football kicked from the penalty spot, a stone thrown from a hilltop, or a cannonball fired from the walls of the Fouban palace. Once the object leaves the foot, the hand or the barrel, the only force acting on it (in our model) is its weight. This section builds the complete mathematical model for that motion.

3.1 What is a Projectile?

DEFINITION Projectile

A projectile is a particle that is projected (launched) with an initial velocity and thereafter moves freely under the action of gravity alone, with no other forces (such as air resistance or thrust) acting on it.

The key phrase is “moves freely under the action of gravity alone”. The moment the projectile loses contact with the launcher, its acceleration is exactly the acceleration due to gravity, directed vertically downwards. This is true whether the projectile is moving upward, downward, or is instantaneously at the top of its flight.

3.2 Modelling Assumptions

To obtain a tractable mathematical model we make the following standard assumptions. In an examination you should be ready to state them.

KEY POINTS Standard Projectile Model Assumptions

1. The projectile is treated as a particle (a point mass); its size, shape and rotation are ignored.
2. The acceleration due to gravity, g , is constant in magnitude and direction. Near the Earth's surface we take $g = 9.8 \text{ m s}^{-2}$ unless the question explicitly says “take $g = 10 \text{ m s}^{-2}$ ”.
3. No air resistance (or any other resistive force). The only force is the weight mg downwards.
4. The motion takes place over distances small enough that the curvature of the Earth and the variation of g with height can be neglected.

WARNING Assumptions in Exam Answers

When a question asks “State the modelling assumptions you have made”, you must list the ones relevant to the problem. The most common required answers are: “The object is modelled as a particle” and “Air resistance is neglected”. Do not write “gravity is constant” unless the question specifically asks about the model for gravity.


3.3 Setting Up the Coordinate System

Because the acceleration is purely vertical, it is natural to choose axes that separate the horizontal and vertical directions.

METHOD Choosing Axes for Projectile Problems

1. Place the origin at the point of projection (or at ground level directly below it).
2. Let the x -axis be horizontal (positive in the direction of the horizontal component of the initial velocity).
3. Let the y -axis be vertical (positive upwards).
4. Then the acceleration vector is $\vec{a} = (0, -g)$.

With this choice, the horizontal and vertical motions become completely independent — a crucial simplification.



Projectile launched from O with speed u at angle θ . The horizontal velocity remains constant; the vertical velocity changes due to gravity.

3.4 Initial Velocity Components

The initial velocity $\vec{u}$ has magnitude u (often denoted V in GCE questions) and makes an angle θ (or α) with the horizontal. Resolving into components:

$$u_x = u \cos \theta, \quad u_y = u \sin \theta.$$

PROPERTY Independence of Horizontal and Vertical Motions

Because $\vec{a} = (0, -g)$, the horizontal acceleration is zero and the vertical acceleration is $-g$. Consequently:

- The horizontal motion is uniform (constant velocity): $v_x = u \cos \theta$ for all t .

- The vertical motion is uniformly accelerated with acceleration $-g$: $v_y = u \sin \theta - gt$.

The two motions share only the common time variable t . This independence is the cornerstone of projectile analysis.

3.5 Equations of Motion

Integrating the acceleration (or using the constant-acceleration formulae) gives the parametric equations for position and velocity as functions of time t .

KEY POINTS Parametric Equations of Projectile Motion

$$\text{Horizontal: } x = (u \cos \theta) t, \quad v_x = u \cos \theta.$$

$$\text{Vertical: } y = (u \sin \theta) t - \frac{1}{2}gt^2, \quad v_y = u \sin \theta - gt.$$

The position vector is $\vec{r}(t) = ((u \cos \theta)t, (u \sin \theta)t - \frac{1}{2}gt^2)$ and the velocity vector is $\vec{v}(t) = (u \cos \theta, u \sin \theta - gt)$.

WARNING Sign of g

Because we chose y positive upwards, the acceleration due to gravity is $-g$. If you instead choose y positive downwards, the acceleration becomes $+g$ and the initial vertical velocity becomes $-u \sin \theta$. Be consistent! The GCE standard is y upwards, $a_y = -g$.

3.6 The Cartesian Equation of the Trajectory

We can eliminate the parameter t to obtain the path y as a function of x . From $x = (u \cos \theta)t$ we have $t = \frac{x}{u \cos \theta}$. Substitute into the equation for y :

$$y = (u \sin \theta) \left(\frac{x}{u \cos \theta} \right) - \frac{1}{2}g \left(\frac{x}{u \cos \theta} \right)^2 = x \tan \theta - \frac{gx^2}{2u^2 \cos^2 \theta}.$$

Using the identity $\sec^2 \theta = 1 + \tan^2 \theta$, this is sometimes written as

$$y = x \tan \theta - \frac{gx^2}{2u^2} (1 + \tan^2 \theta).$$

PROPERTY Trajectory is a Parabola

The Cartesian equation $y = x \tan \theta - \frac{gx^2}{2u^2 \cos^2 \theta}$ is a quadratic in x with a negative coefficient for x^2 . Hence the trajectory of a projectile (under the standard assumptions) is a parabola with a vertical axis, opening downwards.

EXAMPLE Finding the Trajectory Equation

A particle is projected from ground level with a speed of 20 m s^{-1} at an angle of 30° to the horizontal. Taking $g = 10 \text{ m s}^{-2}$, find the Cartesian equation of its trajectory.

$$u = 20, \quad \theta = 30^\circ, \quad \tan 30^\circ = \frac{1}{\sqrt{3}}, \quad \cos 30^\circ = \frac{\sqrt{3}}{2}.$$

$$\begin{aligned} y &= x \tan 30^\circ - \frac{gx^2}{2u^2 \cos^2 30^\circ} \\ &= \frac{x}{\sqrt{3}} - \frac{10x^2}{2 \times 400 \times \left(\frac{\sqrt{3}}{2}\right)^2} \\ &= \frac{x}{\sqrt{3}} - \frac{10x^2}{800 \times \frac{3}{4}} \\ &= \frac{x}{\sqrt{3}} - \frac{10x^2}{600} \\ &= \frac{x}{\sqrt{3}} - \frac{x^2}{60}. \end{aligned}$$

The trajectory is $y = \frac{x}{\sqrt{3}} - \frac{x^2}{60}$ (with x, y in metres).

3.7 Velocity at Any Time

The velocity vector at time t is $\vec{v}(t) = (u \cos \theta, u \sin \theta - gt)$. Its magnitude (speed) and direction are often required.

KEY POINTS Speed and Direction at Time t

$$\text{Speed: } v = \sqrt{v_x^2 + v_y^2} = \sqrt{(u \cos \theta)^2 + (u \sin \theta - gt)^2}.$$

$$\text{Angle } \phi \text{ to the horizontal: } \tan \phi = \frac{v_y}{v_x} = \frac{u \sin \theta - gt}{u \cos \theta}.$$

If $\phi > 0$ the velocity points above the horizontal; if $\phi < 0$ it points below.

WARNING Velocity at the Highest Point

At the maximum height, the vertical velocity $v_y = 0$. The velocity is then purely horizontal: $\vec{v} = (u \cos \theta, 0)$. The speed is minimum there and equals $u \cos \theta$. The acceleration is still g downwards — a common mistake is to think the acceleration becomes zero at the top.

3.8 Worked Example: Complete Analysis

WORKED EXAMPLE 6 Projectile from a Cliff

A stone is projected from the edge of a vertical cliff with a speed of 25 m s^{-1} at an angle of 40° above the horizontal. The cliff is 50 m high. Take $g = 9.8 \text{ m s}^{-2}$.

1. Find the maximum height reached above the cliff top.
2. Find the time taken to reach the sea.
3. Find the horizontal distance from the cliff base where the stone hits the sea.
4. Find the speed and direction of the stone just before impact.

SOLUTION

Step 1: Resolve the initial velocity. $u = 25$, $\theta = 40^\circ$. $u_x = 25 \cos 40^\circ \approx 19.15 \text{ m s}^{-1}$, $u_y = 25 \sin 40^\circ \approx 16.07 \text{ m s}^{-1}$.

Step 2: Maximum height above cliff. At max height, $v_y = 0$. Using $v_y^2 = u_y^2 - 2gh_{\max}$: $0 = (16.07)^2 - 2(9.8)h_{\max} \Rightarrow h_{\max} = \frac{258.2}{19.6} \approx 13.2 \text{ m}$ above the cliff.

Step 3: Time to reach the sea. Take $y = 0$ at cliff top, so sea level is $y = -50$. $y = u_y t - \frac{1}{2}gt^2 = -50$. $16.07t - 4.9t^2 = -50 \Rightarrow 4.9t^2 - 16.07t - 50 = 0$. $t = \frac{16.07 + \sqrt{16.07^2 + 4 \times 4.9 \times 50}}{2 \times 4.9} = \frac{16.07 + \sqrt{258.2 + 980}}{9.8} = \frac{16.07 + 35.19}{9.8} \approx 5.23 \text{ s}$. (We discard the negative root, since time must be positive.)

Step 4: Horizontal distance from cliff base. $x = u_x t = 19.15 \times 5.23 \approx 100 \text{ m}$ (3 s.f.: 100.2 m).

Step 5: Velocity at impact. $v_x = u_x = 19.15 \text{ m s}^{-1}$. $v_y = u_y - gt = 16.07 - 9.8(5.23) = 16.07 - 51.25 = -35.18 \text{ m s}^{-1}$. Speed $v = \sqrt{19.15^2 + (-35.18)^2} = \sqrt{366.7 + 1237.6} = \sqrt{1604.3} \approx 40.1 \text{ m s}^{-1}$. Angle below horizontal: $\tan \phi = \frac{|v_y|}{v_x} = \frac{35.18}{19.15} \approx 1.837 \Rightarrow \phi \approx 61.4^\circ$ below the horizontal.

3.9 Complementary Angles and Range


For a given projection speed u , two different angles of projection that sum to 90° (complementary angles) produce the same horizontal range (on level ground). This is a classic GCE result.

PROPERTY Range on Level Ground

If the projectile lands at the same vertical level from which it was launched ($y = 0$), the time of flight T and the range R are:

$$T = \frac{2u \sin \theta}{g}, \quad R = \frac{u^2 \sin 2\theta}{g}.$$

Since $\sin(180^\circ - 2\theta) = \sin 2\theta$, the angles θ and $90^\circ - \theta$ give the same range. The maximum range is $R_{\max} = \frac{u^2}{g}$, achieved at $\theta = 45^\circ$.



Trajectories for $u = 20 \text{ ms}^{-1}$ at 30° , 45° and 60° ($g = 9.8$). The 30° and 60° trajectories have the same range; 45° gives the maximum range.

3.10 Key Points to Remember

KEY POINTS Summary: Projectile Motion Essentials

1. Always resolve the initial velocity into horizontal and vertical components first.
2. Horizontal motion: constant velocity $u \cos \theta$; $x = (u \cos \theta)t$.
3. Vertical motion: constant acceleration $-g$; $y = (u \sin \theta)t - \frac{1}{2}gt^2$, $v_y = u \sin \theta - gt$.
4. Trajectory: $y = x \tan \theta - \frac{gx^2}{2u^2 \cos^2 \theta}$ (a parabola).
5. Range on level ground: $R = \frac{u^2 \sin 2\theta}{g}$; maximum at $\theta = 45^\circ$.
6. At the highest point: $v_y = 0$, speed $= u \cos \theta$, acceleration $= g$ downwards.
7. Check the value of g given in the question (9.8 or 10) and the required accuracy (usually

3 significant figures or 1 decimal place).

DID YOU KNOW? Historical Note

The parabolic trajectory of projectiles was first demonstrated mathematically by Galileo Galilei in his *Discorsi e dimostrazioni matematiche intorno a due nuove scienze* (1638). Before Galileo, it was commonly believed (following Aristotle) that a projectile moved in a straight line until its “impetus” was exhausted, then fell vertically. Galileo’s insight — separating the motion into independent horizontal and vertical components — is exactly the method we use today. In Cameroon, traditional hunters intuitively understood the need to aim higher for distant targets, effectively compensating for the parabolic drop long before the mathematics was formalised.

EXERCISE 3 Practice Problems

- A ball is thrown from ground level with a speed of 15 m s^{-1} at 50° to the horizontal. ($g = 9.8$)
 - Find the maximum height reached.
 - Find the range on level ground.
 - Find the velocity (magnitude and direction) after 1.2 s.
- A projectile is fired from the top of an 80 m high tower with a velocity of 50 m s^{-1} at 30° above the horizontal. ($g = 10$)
 - Find the time of flight.
 - Find the horizontal distance from the foot of the tower where it lands.
 - Find the angle of impact with the ground.
- Show that for a given speed u , the maximum range on a horizontal plane is $\frac{u^2}{g}$ and occurs at $\theta = 45^\circ$. For a fixed range $R < \frac{u^2}{g}$, show that there are two possible angles of projection and that they are complementary.

SOLUTION Exercise 1

- $u = 15$, $\theta = 50^\circ$, $u_x = 15 \cos 50^\circ \approx 9.64$, $u_y = 15 \sin 50^\circ \approx 11.49$.

$$(a) H = \frac{u_y^2}{2g} = \frac{11.49^2}{19.6} \approx 6.74 \text{ m.}$$

$$(b) R = \frac{u^2 \sin 100^\circ}{g} = \frac{225 \times 0.9848}{9.8} \approx 22.6 \text{ m.}$$

$$(c) \text{ At } t = 1.2: v_x = 9.64, v_y = 11.49 - 9.8(1.2) = -0.27. \text{ Speed } \approx 9.64 \text{ m s}^{-1}, \text{ angle } \approx -1.6^\circ \text{ (just below horizontal).}$$

$$2. u = 50, \theta = 30^\circ, u_x = 25\sqrt{3} \approx 43.3, u_y = 25. \text{ Tower height } 80 \text{ m } (y = -80).$$

$$(a) -80 = 25t - 5t^2 \implies 5t^2 - 25t - 80 = 0 \implies t^2 - 5t - 16 = 0 \implies t = \frac{5 + \sqrt{89}}{2} \approx 7.22 \text{ s.}$$

$$(b) x = 43.3 \times 7.22 \approx 312.6 \text{ m.}$$

$$(c) v_y = 25 - 10(7.22) = -47.2. \tan \phi = 47.2/43.3 \implies \phi \approx 47.5^\circ \text{ below horizontal.}$$

$$3. R = \frac{u^2 \sin 2\theta}{g}. \text{ Max when } \sin 2\theta = 1 \implies 2\theta = 90^\circ \implies \theta = 45^\circ, R_{\max} = u^2/g. \text{ For } R < u^2/g, \sin 2\theta = Rg/u^2. \text{ Two solutions: } 2\theta_1 = \alpha, 2\theta_2 = 180^\circ - \alpha \implies \theta_1 + \theta_2 = 90^\circ.$$

4 RANGE, MAXIMUM HEIGHT AND TIME OF FLIGHT

In the previous section we modelled a projectile as a particle moving under gravity alone, and we established its equations of motion. With axes Ox horizontal and Oy vertically upwards, launch speed u at angle θ above the horizontal from the origin, we found

$$x(t) = u \cos \theta t, \quad y(t) = u \sin \theta t - \frac{1}{2}gt^2,$$

$$v_x = u \cos \theta, \quad v_y = u \sin \theta - gt,$$

together with the trajectory equation

$$y = x \tan \theta - \frac{g x^2}{2u^2 \cos^2 \theta}.$$

Everything in this section is a consequence of these four lines. Three questions dominate GCE questions: How long is the projectile in the air? How high does it go? How far does it go? We answer each in turn, deriving every formula from the equations of motion rather than quoting results.

4.1 Time of Flight on Level Ground

DEFINITION Time of flight

The **time of flight** T of a projectile is the total time between launch and landing. On level ground (launch and landing at the same height), it is the positive solution of $y(t) = 0$.

Derivation. The projectile lands when its height returns to zero:

$$y(t) = u \sin \theta t - \frac{1}{2}gt^2 = t \left(u \sin \theta - \frac{1}{2}gt \right) = 0.$$

The solution $t = 0$ is the launch instant. The other solution gives the landing time:

$$u \sin \theta - \frac{1}{2}gT = 0 \quad \Rightarrow \quad T = \frac{2u \sin \theta}{g}.$$

PROPERTY Time of flight on level ground

For a projectile launched from level ground with speed u at angle θ ,

$$T = \frac{2u \sin \theta}{g}$$

The derivation above is examinable and should be reproduced rather than quoted.

Notice that T depends only on the vertical component of the launch velocity: doubling $u \sin \theta$ doubles the time in the air. The horizontal component $u \cos \theta$ plays no role in how long the projectile stays airborne — a direct consequence of the independence of horizontal and vertical motions.

4.2 Maximum Height and the Symmetry of the Trajectory

Intuition: the projectile rises until its vertical velocity is exhausted, then falls. At the very top, the motion is instantaneously horizontal, so $v_y = 0$ (while $v_x = u \cos \theta$ is unchanged).

Derivation. Setting $v_y = 0$:

$$u \sin \theta - gt = 0 \quad \Rightarrow \quad t_{\text{top}} = \frac{u \sin \theta}{g} = \frac{T}{2}.$$

Substituting into $y(t)$:

$$H = y\left(\frac{T}{2}\right) = u \sin \theta \cdot \frac{u \sin \theta}{g} - \frac{1}{2}g \left(\frac{u \sin \theta}{g} \right)^2 = \frac{u^2 \sin^2 \theta}{g} - \frac{u^2 \sin^2 \theta}{2g} = \frac{u^2 \sin^2 \theta}{2g}.$$

PROPERTY Maximum height and symmetry

The **maximum height** above the launch point is

$$H = \frac{u^2 \sin^2 \theta}{2g}$$

reached at time $t = T/2$. The trajectory is **symmetric** about the vertical line through the highest point: the time going up equals the time coming down, and the speed at any height on the way down equals the speed at the same height on the way up.

WARNING H is measured from the launch point

The formula $H = \frac{u^2 \sin^2 \theta}{2g}$ gives the height gained above the launch point. If the projectile is launched from a height h above the ground (a cliff, a balcony), the greatest height above the ground is $h + H$. Mixing these two is a very common error.

4.3 Horizontal Range on Level Ground

The **range** R is the horizontal distance between the launch and landing points. Since v_x is constant, we simply evaluate x at the landing time T :

$$R = x(T) = u \cos \theta \cdot \frac{2u \sin \theta}{g} = \frac{2u^2 \sin \theta \cos \theta}{g}.$$

Using the double-angle identity $2 \sin \theta \cos \theta = \sin 2\theta$:

PROPERTY Range on level ground


$$R = \frac{u^2 \sin 2\theta}{g}$$

valid only when launch and landing are at the same height.

KEY POINTS The three key formulae — level ground, launch from origin

$$T = \frac{2u \sin \theta}{g}, \quad H = \frac{u^2 \sin^2 \theta}{2g}, \quad R = \frac{u^2 \sin 2\theta}{g}.$$

Conditions of validity: flat horizontal ground, launch and landing at the same level, no air resistance, g constant.



Annotated trajectory on level ground: time of flight T , maximum height H , range R , equal launch and landing angles, and the axis of symmetry through the summit.

4.4 Maximum Range and Complementary Angles


For a fixed launch speed u , the range $R = \frac{u^2}{g} \sin 2\theta$ depends on θ only through $\sin 2\theta$. Since the greatest value of the sine function is 1, attained when $2\theta = 90^\circ$:

PROPERTY Maximum range and complementary angles

For a given speed u on level ground:

1. the range is greatest when $\theta = 45^\circ$, giving $R_{\max} = \frac{u^2}{g}$;
2. two complementary angles θ and $90^\circ - \theta$ give the same range, because $\sin 2(90^\circ - \theta) = \sin(180^\circ - 2\theta) = \sin 2\theta$.

So a ball kicked at 30° and one kicked at 60° with the same speed land at the same spot — but the 60° ball stays in the air longer (larger $u \sin \theta$) and goes higher. Goalkeepers and basketball players exploit this trade-off constantly.



Range (in units of u^2/g) against launch angle. The maximum occurs at 45° ; complementary angles give equal ranges.

WORKED EXAMPLE 7 Level-ground projectile

A footballer on a flat pitch in Yaounde kicks a ball with speed 20 m s^{-1} at 30° above the horizontal. Taking $g = 10 \text{ m s}^{-2}$, find (a) the time of flight, (b) the maximum height, (c) the range, (d) the speed and direction of the ball at impact.

SOLUTION

Solution. Here $u = 20$, $\sin \theta = \frac{1}{2}$, $\cos \theta = \frac{\sqrt{3}}{2}$, $\sin 2\theta = \sin 60^\circ = \frac{\sqrt{3}}{2}$.

(a) Time of flight:

$$T = \frac{2u \sin \theta}{g} = \frac{2(20)(0.5)}{10} = 2 \text{ s.}$$

(b) Maximum height:

$$H = \frac{u^2 \sin^2 \theta}{2g} = \frac{400 \times 0.25}{20} = 5 \text{ m.}$$

(c) Range:

$$R = \frac{u^2 \sin 2\theta}{g} = \frac{400 \times \frac{\sqrt{3}}{2}}{10} = 20\sqrt{3} \approx 34.6 \text{ m.}$$

(d) At $t = T = 2 \text{ s}$:

$$v_x = 20 \cos 30^\circ = 10\sqrt{3} \approx 17.3 \text{ m s}^{-1}, \quad v_y = 20 \sin 30^\circ - 10(2) = 10 - 20 = -10 \text{ m s}^{-1}.$$

Speed at impact:

$$v = \sqrt{v_x^2 + v_y^2} = \sqrt{300 + 100} = 20 \text{ m s}^{-1},$$

directed at $\tan^{-1}\left(\frac{10}{10\sqrt{3}}\right) = 30^\circ$ below the horizontal. Note the symmetry: impact speed equals launch speed, impact angle equals launch angle — exactly as the symmetry property predicts.

4.5 When the Landing Level Differs from the Launch Level

The boxed formulae above assume level ground. If the projectile lands higher or lower than it was launched, they fail — and using them is one of the most penalised mistakes at GCE.

METHOD Projectile landing at a different height

1. Write $x(t) = u \cos \theta t$ and $y(t) = u \sin \theta t - \frac{1}{2}gt^2$ from scratch.
2. Set $y(t)$ equal to the landing height (e.g. $y = -h$ for a drop of h) and solve the quadratic in t ; keep the positive root.
3. Compute the horizontal distance as $x = u \cos \theta t$ with that value of t .
4. Do not use $R = u^2 \sin 2\theta / g$.

Launch from a height h . Suppose the projectile is launched from a point at height h above flat ground, and we measure y upwards from the launch point. Landing occurs when $y = -h$:

$$u \sin \theta T - \frac{1}{2}gT^2 = -h \implies \frac{1}{2}gT^2 - u \sin \theta T - h = 0,$$

a quadratic in T whose positive root is

$$T = \frac{u \sin \theta + \sqrt{u^2 \sin^2 \theta + 2gh}}{g}.$$

The horizontal distance travelled is then $X = u \cos \theta T$, and the greatest height above the ground is $h + \frac{u^2 \sin^2 \theta}{2g}$. In an examination, prefer solving the quadratic with the actual numbers rather than quoting this last formula.

WORKED EXAMPLE 8 Stone thrown from a cliff

A stone is thrown horizontally with speed 15 m s^{-1} from the top of a cliff 45 m above the sea. Taking $g = 10 \text{ m s}^{-2}$, find the time of flight, the horizontal distance travelled, and the velocity at impact.

SOLUTION

Solution. Launch is horizontal, so $\theta = 0$: $u \sin \theta = 0$, $u \cos \theta = 15$. Take y upwards from the

top of the cliff.

Time of flight: $y(t) = -\frac{1}{2}(10)t^2 = -5t^2$. Landing: $-5t^2 = -45$ gives $t^2 = 9$, so $T = 3$ s.

Horizontal distance: $X = 15 \times 3 = 45$ m.

Velocity at impact: $v_x = 15 \text{ m s}^{-1}$, $v_y = -10(3) = -30 \text{ m s}^{-1}$, so

$$v = \sqrt{15^2 + 30^2} = \sqrt{1125} \approx 33.5 \text{ m s}^{-1}, \quad \tan \alpha = \frac{30}{15} = 2,$$

i.e. $\alpha \approx 63.4^\circ$ below the horizontal.

Sloping ground (GCE-useful extension). If the ground is a slope, say the line $y = mx$ (or $y = mx + c$), the landing point is found by substituting the trajectory equation into the equation of the slope and solving for x :

$$x \tan \theta - \frac{gx^2}{2u^2 \cos^2 \theta} = mx.$$

The non-zero root gives the landing point directly. This avoids finding the time of flight first and is often the fastest route in examination questions.

WARNING Do not quote the level-ground formulae blindly

$R = \frac{u^2 \sin 2\theta}{g}$ and $T = \frac{2u \sin \theta}{g}$ are valid only when launch and landing are at the same height. On a slope, from a cliff, or onto a raised platform, return to $x(t)$ and $y(t)$ and solve.

METHOD Exam technique: derive, then compute

GCE mark schemes award method marks for the derivation. A safe routine:

1. State $x = u \cos \theta t$, $y = u \sin \theta t - \frac{1}{2}gt^2$.
2. For T : solve $y = 0$ (or the appropriate landing condition).
3. For H : set $v_y = 0$, find t , substitute into y .
4. For R : substitute T into x .

Quoting $R = u^2 \sin 2\theta / g$ with no justification when the ground is not level scores zero for that part.

EXERCISE 4

A projectile is launched from level ground at 25 m s^{-1} . Taking $g = 10 \text{ m s}^{-2}$, find the two possible launch angles for which the range is 50 m, and compare the two times of flight.

SOLUTION Exercise 1

$R = \frac{u^2 \sin 2\theta}{g}$ gives $50 = \frac{625 \sin 2\theta}{10}$, so $\sin 2\theta = 0.8$, hence $2\theta = 53.13^\circ$ or 126.87° , i.e. $\theta \approx 26.6^\circ$ or 63.4° (complementary, as expected). Times of flight: $T_1 = \frac{2(25) \sin 26.6^\circ}{10} \approx 2.24$ s and $T_2 = \frac{2(25) \sin 63.4^\circ}{10} \approx 4.47$ s. Same range, but the steeper launch stays in the air about twice as long.

KEY POINTS Summary of key formulae and their conditions

Quantity	Formula	Conditions
Time of flight T	$\frac{2u \sin \theta}{g}$	level ground
Maximum height H	$\frac{u^2 \sin^2 \theta}{2g}$	above launch point
Range R	$\frac{u^2 \sin 2\theta}{g}$	level ground only
Maximum range	$\frac{u^2}{g}$ at $\theta = 45^\circ$	level ground
Time to summit	$T/2$	any landing level (for the summit itself)

When conditions fail, go back to $x(t)$ and $y(t)$: they never fail.

DID YOU KNOW? Why 45° is not always best

On level ground 45° maximises the range, but when launching from a height h the optimal angle is slightly less than 45° — shot-putters, who release the ball about 2 m above the ground, indeed throw at angles near 40° to 42° . The extra height gives the lower, faster trajectory more time to travel.

5 APPLICATIONS AND PROBLEM-SOLVING WITH PROJECTILES

In the previous section we established the three headline results of projectile motion: the time of flight $T = \frac{2u \sin \theta}{g}$, the maximum height $H = \frac{u^2 \sin^2 \theta}{2g}$ and the range $R = \frac{u^2 \sin 2\theta}{g}$. Those

formulas answer the question “what happens when we launch a projectile?”. In real life and in the GCE examination, the question is usually the reverse: “how must we launch the projectile so that a given thing happens?” — so that the ball clears the net, lands in the basket, or reaches a target. This section is devoted to that art: translating a concrete condition into an equation, and solving it.

5.1 The General Strategy for Projectile Problems

Almost every projectile problem is solved by the same five-step routine. Master it once and the whole chapter becomes mechanical.

METHOD The five-step strategy for projectile problems

1. Draw a diagram. Sketch the launch point, the landing point or target, and any obstacle. Mark all given distances and heights.
2. Choose axes. Take the origin at the launch point (unless the problem suggests otherwise), the x -axis horizontal in the direction of motion and the y -axis vertically upwards.
3. Resolve the initial velocity. If the launch speed is u at angle θ above the horizontal, then $u_x = u \cos \theta$ and $u_y = u \sin \theta$.
4. Write the equations of motion.

$$x(t) = (u \cos \theta) t, \quad y(t) = (u \sin \theta) t - \frac{1}{2} g t^2,$$

or, eliminating t , the **trajectory equation**

$$y = x \tan \theta - \frac{g x^2}{2u^2 \cos^2 \theta} = x \tan \theta - \frac{g x^2}{2u^2} (1 + \tan^2 \theta).$$

5. Translate the condition of the problem into an equation. “Hits the target (x_0, y_0) ” means $y = y_0$ when $x = x_0$; “clears the wall” means $y \geq h$ when $x = d$; “lands on the ground” means $y = 0$ (or $y = -h$ if launched from a height). Then solve.

WARNING Exam tip: keep g symbolic

Do not substitute $g = 9.81 \text{ m s}^{-2}$ (or 10) until the very last line of your calculation. Rounding g early makes errors accumulate through the working, and examiners award method marks only if your symbolic line is correct. Carry g as a letter, simplify, and only then compute the numerical value.

5.2 Hitting a Given Target: the Quadratic in $\tan \theta$

Suppose a projectile is launched from the origin with speed u and we want it to pass through the

point (x_0, y_0) . Which angle θ should we choose? Substituting (x_0, y_0) into the trajectory equation gives

$$y_0 = x_0 \tan \theta - \frac{g x_0^2}{2u^2} (1 + \tan^2 \theta).$$

PROPERTY Condition to pass through (x_0, y_0)

Setting $m = \tan \theta$, the condition becomes a **quadratic equation** in $\tan \theta$:

$$\frac{g x_0^2}{2u^2} m^2 - x_0 m + \left(y_0 + \frac{g x_0^2}{2u^2} \right) = 0. \quad (1)$$

The key algebraic step is the identity $\sec^2 \theta = 1 + \tan^2 \theta$ applied to $\frac{1}{\cos^2 \theta}$.

Since a quadratic has (in general) two roots, there are usually **two** launch angles that hit the same target: a low, flat trajectory and a high, lobbed one. This matches experience: a footballer can reach a teammate either with a driven pass or with a chipped ball.

WARNING Two values of $\tan \theta$ usually mean two possible angles

When the quadratic in $\tan \theta$ has two positive roots, both correspond to genuine trajectories through the target — check that both are physically acceptable in the context (a “low” solution may be blocked by an obstacle; a negative root would mean firing below the horizontal and is usually rejected). If the discriminant is negative, the target is out of reach for the given speed u : no angle works. The boundary case (discriminant zero) gives the single angle that just reaches the target; this defines the **envelope of trajectories** (parabola of safety) $y = \frac{u^2}{2g} - \frac{gx^2}{2u^2}$.

WORKED EXAMPLE 9 Hitting a target

A projectile is launched from the origin with speed $u = 20 \text{ m s}^{-1}$. Find the angle(s) of projection for it to pass through the point $(30 \text{ m}, 10 \text{ m})$. Take $g = 10 \text{ m s}^{-2}$.

SOLUTION

Solution. The trajectory condition at $(x_0, y_0) = (30, 10)$ is

$$10 = 30 \tan \theta - \frac{10 \times 30^2}{2 \times 20^2} (1 + \tan^2 \theta).$$

Compute the coefficient: $\frac{10 \times 900}{800} = 11.25$. With $m = \tan \theta$:

$$10 = 30m - 11.25(1 + m^2) \implies 11.25 m^2 - 30m + 21.25 = 0.$$

Multiply by 4 to clear decimals: $45m^2 - 120m + 85 = 0$, i.e. $9m^2 - 24m + 17 = 0$. Then

$$m = \frac{24 \pm \sqrt{576 - 612}}{18} = \frac{24 \pm \sqrt{-36}}{18}.$$

The discriminant is negative: with $u = 20 \text{ m s}^{-1}$ the target is unreachable. (Indeed the maximum range on level ground is $u^2/g = 40 \text{ m}$, but the point $(30, 10)$ lies above the reachable envelope.) Increasing the speed to $u = 25 \text{ m s}^{-1}$ gives coefficient $\frac{10 \times 900}{2 \times 625} = 7.2$, so

$$7.2m^2 - 30m + 17.2 = 0, \quad m = \frac{30 \pm \sqrt{900 - 495.36}}{14.4} = \frac{30 \pm 20.12}{14.4},$$

giving $m \approx 3.48$ ($\theta \approx 74.0^\circ$, high lob) or $m \approx 0.686$ ($\theta \approx 34.5^\circ$, flat throw). Both angles hit the target.

5.3 Clearing an Obstacle


A very common GCE situation: a ball must pass over a wall, a net or a fence of height h situated at horizontal distance d from the launch point.

METHOD “Just clears an obstacle” — the limiting case

The projectile clears the obstacle if $y \geq h$ when $x = d$. The limiting case — minimum launch speed, or critical angle — corresponds to the trajectory passing exactly through the top of the obstacle:

$$h = d \tan \theta - \frac{g d^2}{2u^2} (1 + \tan^2 \theta).$$

Impose this equality to find the minimum speed $u_{\min}$ (for a given angle) or the critical angle (for a given speed). Any speed above $u_{\min}$ clears the wall comfortably.



A projectile clearing a wall of height h at distance d . The solid curve is the limiting trajectory, passing exactly through the top of the wall: $y(d) = h$.

EXAMPLE Ball over a wall

A footballer kicks a ball from ground level at $\theta = 45^\circ$. A wall of height $h = 2$ m stands $d = 10$ m away. What is the minimum kicking speed for the ball to clear the wall? (Take $g = 10 \text{ m s}^{-2}$.)

Solution. Limiting case: $y = h$ at $x = d$ with $\tan 45^\circ = 1$:

$$2 = 10 - \frac{10 \times 10^2}{2u^2}(1 + 1) \Rightarrow \frac{1000}{u^2} = 8 \Rightarrow u^2 = 125, \quad u_{\min} \approx 11.2 \text{ m s}^{-1}.$$

5.4 Horizontal Projection from a Height


If a projectile is launched horizontally from a height h — a stone rolled off a table, a package dropped from a helicopter moving horizontally, water spurting from a horizontal pipe on a bridge — then $\theta = 0$ and the equations simplify dramatically:

$$x(t) = u t, \quad y(t) = -\frac{1}{2} g t^2 \quad (\text{taking } O \text{ at the launch point, } y \text{ upwards}).$$

The time of flight is fixed by the height alone: falling through h gives

$$h = \frac{1}{2} g T^2 \Rightarrow T = \sqrt{\frac{2h}{g}}, \quad \text{range from the foot: } R = uT = u\sqrt{\frac{2h}{g}}.$$

Notice that T does not depend on u : a fast stone and a slow stone dropped from the same cliff hit the sea at the same instant.



Horizontal projection from a cliff of height h . The trajectory is a half-parabola; the time of flight depends only on h .

WORKED EXAMPLE 10 Stone from a bridge

A stone is thrown horizontally with speed $u = 12 \text{ m s}^{-1}$ from a bridge 20 m above a river on the Douala–Yaoundé highway. Take $g = 10 \text{ m s}^{-2}$. Find (a) the time of flight, (b) the horizontal distance travelled, (c) the speed with which the stone hits the water.

SOLUTION

Solution. (a) $T = \sqrt{\frac{2h}{g}} = \sqrt{\frac{2 \times 20}{10}} = 2 \text{ s}$.

(b) $R = uT = 12 \times 2 = 24 \text{ m}$.

(c) At impact: $v_x = u = 12 \text{ m s}^{-1}$ and $v_y = -gT = -20 \text{ m s}^{-1}$. Hence

$$v = \sqrt{v_x^2 + v_y^2} = \sqrt{144 + 400} = \sqrt{544} \approx 23.3 \text{ m s}^{-1}.$$

Note that the impact speed is not $u + gT$: the horizontal and vertical components must be combined as vectors.

5.5 Two Projectiles and Collision Problems

A useful GCE extension: two particles A and B are in projectile motion simultaneously. They **collide** if they occupy the **same position at the same time**: one writes $x_A(t) = x_B(t)$ and $y_A(t) = y_B(t)$ and solves the system for t and the unknown launch parameter. A classic result (the “monkey and hunter”): if particle A is fired directly at particle B which is released from rest at the same instant, then A always hits B (provided the ground does not intervene), because both acquire the same downward displacement $\frac{1}{2}gt^2$ relative to the straight line of sight.

EXAMPLE Collision in mid-air — monkey and hunter

From the origin O , particle A is projected with $u = 20 \text{ m s}^{-1}$ at $\theta = 60^\circ$. At the same instant, particle B is released from rest at the point $(10 \text{ m}, 10\sqrt{3} \text{ m})$ — i.e. on the initial line of sight of A . Show that they collide and find the time and position of collision. (Take $g = 10 \text{ m s}^{-2}$.)

Solution. For A : $u_x = 20 \cos 60^\circ = 10$, $u_y = 20 \sin 60^\circ = 10\sqrt{3}$.

$$x_A = 10t, \quad y_A = 10\sqrt{3}t - 5t^2.$$

For B (dropped from rest at $(10, 10\sqrt{3})$):

$$x_B = 10, \quad y_B = 10\sqrt{3} - 5t^2.$$

Collision requires $x_A = x_B$ and $y_A = y_B$ simultaneously. $x_A = x_B \Rightarrow 10t = 10 \Rightarrow t = 1 \text{ s}$. At $t = 1$: $y_A = 10\sqrt{3} - 5$, $y_B = 10\sqrt{3} - 5$. The y -coordinates match automatically. Thus they collide at $t = 1 \text{ s}$ at the point $(10 \text{ m}, 10\sqrt{3} - 5 \approx 12.3 \text{ m})$.

5.6 Real-Life Modelling and the Limits of the Model**EXPERIMENT** Modelling task: the free kick

A footballer takes a free kick 25 m from goal. The crossbar is 2.44 m high and the defensive wall, 9 m away, jumps to 2 m. Model the ball as a particle launched from ground level at $u = 22 \text{ m s}^{-1}$, $\theta = 25^\circ$, with $g = 9.81 \text{ m s}^{-2}$.

- At the wall ($x = 9$): $y = 9 \tan 25^\circ - \frac{9.81 \times 81}{2 \times 484 \cos^2 25^\circ} \approx 4.20 - 1.00 = 3.20 \text{ m} > 2 \text{ m}$: the ball clears the wall.
- At the goal line ($x = 25$): $y = 25 \tan 25^\circ - \frac{9.81 \times 625}{2 \times 484 \cos^2 25^\circ} \approx 11.66 - 7.71 = 3.95 \text{ m}$: too high — it sails over the bar. The player must reduce the speed or the angle. Trying $\theta = 20^\circ$: at the wall $y \approx 2.35 \text{ m}$ (still clears) and at the goal $y \approx 1.93 \text{ m} < 2.44 \text{ m}$: goal!

DID YOU KNOW? Modelling box: what we have ignored

Our model treats the projectile as a **particle** moving under gravity alone, in a vacuum, over a flat non-rotating Earth. Real projectiles experience air resistance (which shortens the range and makes the descent steeper than the ascent) and spin (the Magnus effect bends footballs and golf balls sideways). A struck football may lose 10–20% of its vacuum range. In the examination, when a question says “state one modelling assumption” or “comment on the model”, expected answers include: air resistance is neglected; the projectile is modelled as a particle (its size is ignored); g is taken as constant; wind is ignored; the ground is horizontal. Stating these clearly earns easy marks.

WARNING Common mistakes in applied projectile problems

- Using the range formula $R = u^2 \sin 2\theta / g$ when the landing level differs from the launch level (cliff, raised target). That formula is valid only on horizontal ground; otherwise solve $y(t) = -h$.
- Forgetting that “just clears” means equality $y(d) = h$, not inequality, when finding a minimum speed.
- Mixing degrees and radians on the calculator when evaluating $\tan \theta$, $\sin \theta$.
- Adding velocity components as scalars instead of using $v = \sqrt{v_x^2 + v_y^2}$.
- Substituting $g = 9.81$ too early and losing accuracy (and method marks).

EXERCISE 5 Practice

1. A ball is thrown from ground level at $u = 14 \text{ m s}^{-1}$, $\theta = 60^\circ$. Does it clear a fence 3 m high standing 8 m away? (Take $g = 9.81 \text{ m s}^{-2}$.)
2. A hose held horizontally 1.5 m above the ground sends water at 8 m s^{-1} . How far from the nozzle does the jet strike the ground?
3. A projectile launched from O with speed 28 m s^{-1} must pass through (40 m, 15 m). Find the two possible angles of projection. (Take $g = 9.8 \text{ m s}^{-2}$.)

SOLUTION Exercise 1

1. $y(8) = 8 \tan 60^\circ - \frac{9.81 \times 64}{2 \times 196 \cos^2 60^\circ} = 13.86 - 6.41 = 7.45 \text{ m} > 3 \text{ m}$: yes, comfortably.
2. $T = \sqrt{2(1.5)/9.81} \approx 0.553 \text{ s}$; $R = 8 \times 0.553 \approx 4.42 \text{ m}$.
3. $15 = 40m - \frac{9.8 \times 1600}{2 \times 784}(1+m^2) = 40m - 10(1+m^2)$, so $10m^2 - 40m + 25 = 0$, i.e. $2m^2 - 8m + 5 = 0$:
 $m = \frac{8 \pm \sqrt{24}}{4} = 2 \pm \frac{\sqrt{6}}{2}$, giving $\theta \approx 72.8^\circ$ or $\theta \approx 37.8^\circ$.

KEY POINTS Key points of the section

- Five steps: diagram, axes, resolve $\vec{u}$, write $x(t)$ and $y(t)$, translate the condition into an

equation.

- Passing through (x_0, y_0) gives a quadratic in $\tan \theta$ via $\sec^2 \theta = 1 + \tan^2 \theta$; two roots usually mean a high and a low trajectory.
- “Just clears an obstacle” $\Rightarrow$ equality $y(d) = h$ (limiting case).
- Horizontal launch from height h : $T = \sqrt{2h/g}$, $R = u\sqrt{2h/g}$; T is independent of u .
- Collisions: same position at the same time — solve $x_A = x_B$, $y_A = y_B$.
- Keep g symbolic until the last line, and always state modelling assumptions (no air resistance, particle model, constant g).

6 FREE FALL, VERTICAL PROJECTION AND MOTION GRAPHS

In the previous section we treated projectiles as two independent motions: uniform motion horizontally and uniformly accelerated motion vertically. In this final section we strip away the horizontal part completely and concentrate on the purely vertical motion of a body moving under gravity alone: a stone dropped from a bridge over the Wouri, a mango falling from a tree, a ball thrown straight up on the playground. These one-dimensional motions are the simplest of all, yet they contain every idea of the chapter, and they lend themselves beautifully to graphical analysis — a favourite GCE examination theme.

Throughout this section we take $g = 9.81 \text{ m s}^{-2}$, often approximated as $g = 10 \text{ m s}^{-2}$ when the question allows, and we neglect air resistance.

6.1 Free Fall: Dropped from Rest

EXPERIMENT Galileo's idea

Drop a small stone and a heavy hammer from the same height at the same instant. Contrary to everyday intuition, they hit the ground essentially together. Air resistance, not weight, is what makes a feather fall slowly; in a vacuum a feather and a hammer fall side by side. The conclusion is profound: all bodies in free fall have the same acceleration, whatever their mass.

DEFINITION Free fall

A body is in **free fall** when the only force acting on it is its **weight**. By Newton's second law its acceleration is then g vertically downwards, whatever its velocity — whether the body

is moving down, is momentarily at rest, or is even moving upwards.

WARNING Free fall does not mean “falling down”

A ball thrown upwards is in free fall from the instant it leaves the hand: the only force on it is its weight, so its acceleration is g downwards throughout. “Free fall” describes the force situation, not the direction of motion.

Consider now a body dropped (released from rest, $u = 0$) from a height. Take the **downwards** direction as positive. Then $a = g$, $u = 0$, and the constant-acceleration formulae $v = u + at$, $s = ut + \frac{1}{2}at^2$, $v^2 = u^2 + 2as$ give immediately:

PROPERTY Free fall from rest, downwards positive

For a body dropped from rest and falling through a height h :

$$v = gt, \quad h = \frac{1}{2}gt^2, \quad v^2 = 2gh.$$

The speed reached after falling a height h is $v = \sqrt{2gh}$, independent of the mass of the body. (Proof: direct substitution into the constant-acceleration formulae; examinable as a one-line derivation.)

EXAMPLE Stone dropped from a bridge

A stone is dropped from a bridge 45 m above a river. Taking $g = 10 \text{ m s}^{-2}$, the time to reach the water satisfies $45 = \frac{1}{2}(10)t^2$, so $t^2 = 9$ and $t = 3 \text{ s}$. Its speed on entering the water is $v = gt = 10 \times 3 = 30 \text{ m s}^{-1}$ (check: $v = \sqrt{2gh} = \sqrt{2 \times 10 \times 45} = \sqrt{900} = 30 \text{ m s}^{-1}$).

6.2 Object Thrown Vertically Upwards

Now throw a ball straight up with initial speed u . Choose **upwards as positive**. The acceleration is $a = -g$ (gravity points down), and the constant-acceleration formulae become:

KEY POINTS Upward projection, upwards positive

$$v = u - gt, \quad y = ut - \frac{1}{2}gt^2, \quad v^2 = u^2 - 2gy.$$

At the greatest height $v = 0$, giving:

$$\text{time to greatest height} = \frac{u}{g}, \quad \text{greatest height} = H = \frac{u^2}{2g}.$$

The total time of flight back to the launch point is $T = \frac{2u}{g}$.

Derivation of the greatest height.

At the top of the flight the body is momentarily at rest: $v = 0$. From $v = u - gt$, the time to reach the top is $t = u/g$. Substituting into $y = ut - \frac{1}{2}gt^2$:

$$H = u\left(\frac{u}{g}\right) - \frac{1}{2}g\left(\frac{u}{g}\right)^2 = \frac{u^2}{g} - \frac{u^2}{2g} = \frac{u^2}{2g}.$$

(The same result comes instantly from $v^2 = u^2 - 2gy$ with $v = 0$.)

PROPERTY The highest point

At the greatest height the **velocity** is instantaneously zero, but the **acceleration** is still g directed downwards. The body does not “stop accelerating” at the top — it is in free fall at every instant of the flight.

WARNING Zero velocity does not mean zero acceleration

A very common GCE error is to write $a = 0$ at the highest point “because the ball stops”. The ball stops only for an instant; gravity never switches off. If a were zero at the top, the ball would stay there forever!

Symmetry of vertical motion.

Set $y = 0$ in $y = ut - \frac{1}{2}gt^2$: $t(u - \frac{1}{2}gt) = 0$, so the body returns to the launch level at $t = \frac{2u}{g}$ — exactly twice the time to reach the top. Hence **time up = time down**. Moreover, at $t = 2u/g$:

$$v = u - g \cdot \frac{2u}{g} = -u,$$

so the body returns with the same speed u , directed downwards. More generally, at any given height on the way up and on the way down, the speeds are equal (from $v^2 = u^2 - 2gy$, which depends only on y). This perfect symmetry is a powerful checking tool in examinations.

WORKED EXAMPLE 11 Ball thrown vertically upwards

A ball is thrown vertically upwards with speed 24.5 m s^{-1} from ground level. Taking $g = 9.81 \text{ m s}^{-2}$, find (a) the greatest height reached, (b) the total time of flight, (c) the speed with which it returns to the thrower's hand.

SOLUTION

Take upwards as positive: $u = 24.5$, $a = -9.81$.

(a) At the top, $v = 0$. Using $v^2 = u^2 - 2gy$:

$$0 = 24.5^2 - 2(9.81)H \implies H = \frac{24.5^2}{2 \times 9.81} = \frac{600.25}{19.62} \approx 30.6 \text{ m.}$$

(b) Time to the top: $t = \frac{u}{g} = \frac{24.5}{9.81} \approx 2.50 \text{ s}$. By symmetry the total flight time is

$$T = \frac{2u}{g} \approx 5.0 \text{ s.}$$

(c) By symmetry the ball returns with speed $u = 24.5 \text{ m s}^{-1}$ downwards, i.e. velocity -24.5 m s^{-1} . Check: $v = 24.5 - 9.81(5.0) \approx -24.5 \text{ m s}^{-1}$. ✓

6.3 Object Thrown Vertically Downwards; Release from a Moving Body

If a body is thrown downwards with speed u from a height, take **downwards as positive**; then $a = +g$ and:

PROPERTY Downward projection, downwards positive

$$v = u + gt, \quad y = ut + \frac{1}{2}gt^2, \quad v^2 = u^2 + 2gy.$$

METHOD One sign convention per question

1. Choose one positive direction (usually upwards) and write it down.
2. Give every quantity its sign accordingly: u , v , y positive if directed/measured upwards, negative if downwards; $a = -g$ always (upwards positive).
3. Substitute into $v = u + at$, $y = ut + \frac{1}{2}at^2$, $v^2 = u^2 + 2ay$ with the signs included.
4. Interpret the sign of the answer: negative velocity means "moving downwards", negative displacement means "below the launch point".

Never change convention in the middle of a question — this is the source of most sign errors.

A subtle and frequently examined situation is the release of an object from a moving body — a package dropped from a rising balloon, a stone let go from a descending lift.

WARNING The balloon trap

“Dropped” means $u = 0$. But “released from a moving balloon” means the object keeps the balloon’s velocity at the instant of release. A package released from a balloon rising at 5 m s^{-1} has initial velocity $u = +5 \text{ m s}^{-1}$ (upwards): it first continues to rise, slows, stops, and only then falls. Writing $u = 0$ here is a classic GCE trap.

WORKED EXAMPLE 12 Package released from a rising balloon

A balloon is rising vertically at a steady 12 m s^{-1} . When it is 80 m above the ground, a package is released. Taking $g = 10 \text{ m s}^{-2}$, find the time taken for the package to reach the ground and its speed on impact.

SOLUTION

Take upwards as positive, origin at the release point. At release the package has the balloon’s velocity: $u = +12 \text{ m s}^{-1}$, and $a = -10 \text{ m s}^{-2}$. The ground is at $y = -80 \text{ m}$.

Using $y = ut - \frac{1}{2}gt^2$:

$$\begin{aligned} -80 &= 12t - 5t^2 \implies 5t^2 - 12t - 80 = 0. \\ t &= \frac{12 \pm \sqrt{144 + 1600}}{10} = \frac{12 \pm \sqrt{1744}}{10} \approx \frac{12 \pm 41.76}{10}. \end{aligned}$$

Taking the positive root: $t \approx 5.38 \text{ s}$.

Impact velocity: $v = u - gt = 12 - 10(5.38) \approx -41.8 \text{ m s}^{-1}$, i.e. a speed of about 41.8 m s^{-1} downwards. Note that the package first rose for $u/g = 1.2 \text{ s}$ before falling — exactly the behaviour the sign convention predicts automatically.

6.4 Graphical Representation of Vertical Motion


Graphs turn the formulae above into pictures, and the GCE regularly asks candidates to sketch or interpret them. Recall the three golden links from earlier work:

KEY POINTS Reading motion graphs

- **Gradient** of the displacement–time (s – t) graph = **velocity**.

- **Gradient** of the velocity–time (v – t) graph = **acceleration**.
- **Area** under the v – t graph = **displacement** (area below the time axis counts negative).

For upward projection (upwards positive): the v – t graph is a straight line of gradient $-g$ always; it crosses the time axis exactly at the highest point. The a – t graph is the horizontal line $a = -g$ for the whole flight. The s – t graph is a parabola whose maximum occurs at the same instant.



Displacement–time, velocity–time and acceleration–time graphs for a ball thrown vertically upwards ($u = 24.5 \text{ m s}^{-1}$, $g = 9.81 \text{ m s}^{-2}$). The instant of maximum height $t = u/g$ is marked on all three graphs: the parabola peaks, the velocity line crosses the axis, and the acceleration stays fixed at $-g$.

Interpreting the sign changes.


Before the highest point, $v > 0$ (moving up) and the s – t curve rises. At the top, $v = 0$: the v – t line crosses the axis and the s – t parabola has a horizontal tangent. After the top, $v < 0$ (moving down) and the parabola descends. The acceleration graph never changes: the same line $a = -g$ governs the ascent, the instant at the top, and the descent. The area under the v – t line from 0 to $2u/g$ is a triangle above the axis minus a congruent triangle below it — total area zero, matching the zero total displacement when the ball returns to the hand.

Comparing a thrown-up ball and a dropped ball.

For a dropped ball (downwards positive), the v - t graph is a straight line through the origin with gradient $+g$, and the s - t graph is the rising half-parabola $h = \frac{1}{2}gt^2$. For the thrown-up ball (upwards positive) the v - t line starts at $+u$ and slopes down with gradient $-g$, crossing the axis. Same physics, different initial conditions — the gradient is $\pm g$ in every case.

Extension: a bouncing ball.

If a ball bounces on the ground, the v - t graph consists of straight segments of gradient $-g$ (upwards positive), with a sudden jump at each bounce (the velocity reverses rapidly from negative to a smaller positive value, since each bounce loses energy). The s - t graph is a chain of parabola arcs of decreasing height. Examiners occasionally ask for such a sketch: the key features are the constant gradient $-g$ of every segment and the decreasing peaks.



EXERCISE 6 Vertical motion and graphs

A stone is thrown vertically upwards with speed 19.62 m s^{-1} from the top of a cliff 40 m high. Take $g = 9.81 \text{ m s}^{-2}$.

1. Find the greatest height above the cliff top and the time taken to reach it.
2. Find the total time until the stone hits the sea at the foot of the cliff.
3. Find the velocity of the stone on impact.
4. Sketch the v - t graph for the whole motion (upwards positive), marking the time at the greatest height and the impact time.

SOLUTION Exercise 1

Take upwards positive, origin at the cliff top: $u = 19.62$, $a = -9.81$.

1. At the top $v = 0$: $t = u/g = 19.62/9.81 = 2 \text{ s}$; $H = u^2/(2g) = 19.62^2/19.62 = 19.62 \text{ m}$ above the cliff top.

2. The sea is at $y = -40$: $-40 = 19.62t - 4.905t^2$, i.e. $4.905t^2 - 19.62t - 40 = 0$. Then $t = \frac{19.62 + \sqrt{19.62^2 + 4 \times 4.905 \times 40}}{9.81} = \frac{19.62 + \sqrt{384.9 + 784.8}}{9.81} = \frac{19.62 + 34.21}{9.81} \approx 5.49 \text{ s}$ (positive root).

3. $v = 19.62 - 9.81(5.49) \approx -34.2 \text{ m s}^{-1}$, i.e. 34.2 m s^{-1} downwards.

4. Straight line from $(0, 19.62)$ with gradient -9.81 , crossing the time axis at $t = 2 \text{ s}$ (greatest height) and ending at $(5.49, -34.2)$. The net signed area under the line equals -40 m , the total displacement.

KEY POINTS Chapter summary: vertical motion

- Free fall: only weight acts; $a = g$ downwards at all times, even at the top of the flight.
- Dropped from rest (down positive): $v = gt$, $h = \frac{1}{2}gt^2$, $v^2 = 2gh$.
- Thrown up (up positive): $v = u - gt$, $y = ut - \frac{1}{2}gt^2$; top at $t = u/g$, height $u^2/(2g)$; total flight time $2u/g$; returns with speed u .
- Released from a moving body: u = velocity of the body at release.
- Graphs: s - t parabola, v - t line of gradient $-g$ crossing the axis at the top, a - t horizontal at $-g$; gradient of s - t is v , gradient of v - t is a , area under v - t is displacement.

7 INTEGRATION ACTIVITY

7.1 Aims of the activity

This integrative activity closes the chapter by mobilising, in one single realistic problem, every competence developed in Lessons 114 to 122. Working in groups of three or four, you will:

- resolve an initial velocity vector into horizontal and vertical components and write the parametric equations $x(t)$ and $y(t)$ of a projectile;
- derive and use the **trajectory equation** $y = x \tan \theta - \frac{gx^2}{2u^2}(1 + \tan^2 \theta)$;
- translate physical constraints (clearing a wall, staying under a crossbar) into inequalities, and solve the resulting **quadratic inequalities in $\tan \theta$** ;
- compute a time of flight and a velocity vector at a given instant, then its magnitude;
- sketch and interpret a **velocity–time graph** for the vertical motion;
- discuss critically the assumptions of the projectile model, as expected of a GCE Advanced Level candidate.

KEY POINTS Data you may use

Take $g = 10 \text{ m s}^{-2}$, neglect air resistance, and treat the ball as a particle. Give angles to the nearest 0.1° and lengths to 2 decimal places.

7.2 Situation

It is the last minute of the inter-school football final at the sports field of Lycée de Nkol-Eton, Yaoundé. Your school has just been awarded a direct free kick. The ball is placed on flat ground, exactly 22 m from the goal line, directly in front of the centre of the goal. The defending team forms a “wall” of players standing 9.1 m from the ball (the regulation distance is about 9.15 m). The tallest defender jumps and reaches an effective height of 2.0 m, so to pass the wall the ball must be strictly above 2.0 m when it crosses the plane of the wall. The crossbar of the goal is 2.44 m above the ground, so to score, the ball must cross the goal line below 2.44 m (and, of course, still above the ground).

The free-kick specialist strikes the ball so that it leaves the ground with speed $u = 20 \text{ m s}^{-1}$ at an angle θ above the horizontal. The whole stadium is silent. Your group’s job is to advise the kicker: **for which angles θ does the ball clear the wall AND enter the goal below the crossbar?** What is the “optimal” kick?



The free kick: the ball must pass above the wall at 9.1 m and under the crossbar at 22 m. The dashed curve shows the trajectory for the optimal angle $\theta \approx 21^\circ$.

7.3 Tasks

1. **Setting up the model.** Choose axes with origin at the ball's initial position, Ox horizontal towards the goal and Oy vertically upwards. Resolve $\vec{u}$ into components and, starting from $\vec{a} = -g\vec{j}$, show by integration that

$$x(t) = 20t \cos \theta, \quad y(t) = 20t \sin \theta - 5t^2.$$

2. **Trajectory equation.** Eliminate t to obtain

$$y = x \tan \theta - \frac{x^2}{80} (1 + \tan^2 \theta).$$

3. **Clearing the wall.** Write down the condition $y(9.1) > 2.0$ and show that it leads to a quadratic inequality in $T = \tan \theta$. Solve it and give the corresponding interval of angles θ .
4. **Staying under the crossbar.** Write down the condition $0 < y(22) < 2.44$ and show that it also leads to quadratic inequalities in T . Solve them.
5. **The scoring window.** Combine the conditions of Tasks 3 and 4 to find all values of θ for which the kick scores. Which branch of solutions is physically sensible for a direct free kick? Propose an "optimal" angle and justify your choice (margin of safety, time available to the goalkeeper, ...).
6. **Flight to the goal line.** For your chosen optimal angle, compute the time at which the ball crosses the goal line and the velocity vector $\vec{v}$ at that instant; deduce the speed of the ball.
7. **Graphical representation.** Sketch the graph of $v_y(t)$ against t for the whole flight, from $t = 0$ until the ball crosses the goal line. Indicate the slope, the intercept, and the instant when $v_y = 0$; state what the area under the graph represents.
8. **Critical discussion.** Identify one modelling assumption that makes a real free kick behave differently from your model, and explain qualitatively its effect on the trajectory.

7.4 Model answer

Task 1. The only force acting on the ball (neglecting air resistance) is its weight $\vec{W} = m\vec{g}$. Hence

the acceleration is constant: $\vec{a} = \begin{pmatrix} 0 \\ -g \end{pmatrix} = \begin{pmatrix} 0 \\ -10 \end{pmatrix} \text{ m s}^{-2}$. Integrating with respect to time and using

the initial velocity $\vec{v}(0) = \begin{pmatrix} 20 \cos \theta \\ 20 \sin \theta \end{pmatrix}$:

$$v_x(t) = 20 \cos \theta, \quad v_y(t) = 20 \sin \theta - 10t.$$

Integrating again with initial position $\vec{r}(0) = \vec{0}$:

$$x(t) = 20t \cos \theta, \quad y(t) = 20t \sin \theta - 5t^2.$$

Task 2. From $x(t)$ we have $t = \frac{x}{20 \cos \theta}$ (valid for $\theta \neq 90^\circ$). Substitute into $y(t)$:

$$y = x \tan \theta - 5 \left(\frac{x}{20 \cos \theta} \right)^2 = x \tan \theta - \frac{x^2}{80 \cos^2 \theta}.$$

Using the identity $\frac{1}{\cos^2 \theta} = 1 + \tan^2 \theta$, we obtain the Cartesian equation of the trajectory:

$$y = x \tan \theta - \frac{x^2}{80} (1 + \tan^2 \theta).$$

Task 3. Let $T = \tan \theta$. The condition to clear the wall is $y(9.1) > 2.0$:

$$9.1T - \frac{9.1^2}{80}(1 + T^2) > 2.0.$$

Since $9.1^2 = 82.81$ and $\frac{82.81}{80} = 1.035125$, this becomes

$$9.1T - 1.035125(1 + T^2) > 2 \iff -1.035125T^2 + 9.1T - 3.035125 > 0.$$

Multiplying by -1 (which reverses the inequality):

$$1.035125T^2 - 9.1T + 3.035125 < 0.$$

The discriminant is $\Delta = 9.1^2 - 4(1.035125)(3.035125) = 82.81 - 12.570 = 70.240$, so $\sqrt{\Delta} \approx 8.381$. The roots are

$$T_1 = \frac{9.1 - 8.381}{2 \times 1.035125} \approx 0.347, \quad T_2 = \frac{9.1 + 8.381}{2 \times 1.035125} \approx 8.444.$$

The quadratic is negative between its roots:

$$0.347 < \tan \theta < 8.444 \iff 19.1^\circ \leq \theta \leq 83.2^\circ.$$

Task 4. The ball must cross the goal line ($x = 22$) at a height strictly between 0 and 2.44 m:

$$0 < y(22) < 2.44.$$

With $22^2 = 484$ and $\frac{484}{80} = 6.05$, the trajectory equation gives

$$y(22) = 22T - 6.05(1 + T^2).$$

Upper bound (crossbar): $22T - 6.05(1 + T^2) < 2.44$

$$\Leftrightarrow -6.05T^2 + 22T - 8.49 < 0 \Leftrightarrow 6.05T^2 - 22T + 8.49 > 0.$$

Discriminant $\Delta = 484 - 4(6.05)(8.49) = 484 - 205.458 = 278.542$, $\sqrt{\Delta} \approx 16.690$. Roots:

$$T_3 = \frac{22 - 16.690}{12.1} \approx 0.439, \quad T_4 = \frac{22 + 16.690}{12.1} \approx 3.197.$$

The quadratic is positive outside the roots:

$$\tan \theta < 0.439 \quad \text{or} \quad \tan \theta > 3.197 \Leftrightarrow \theta \leq 23.7^\circ \quad \text{or} \quad \theta \geq 72.6^\circ.$$

Lower bound (ground): $22T - 6.05(1 + T^2) > 0$

$$\Leftrightarrow 6.05T^2 - 22T + 6.05 < 0.$$

Discriminant $\Delta = 484 - 4(6.05)^2 = 484 - 146.41 = 337.59$, $\sqrt{\Delta} \approx 18.374$. Roots:

$$T_5 = \frac{22 - 18.374}{12.1} \approx 0.300, \quad T_6 = \frac{22 + 18.374}{12.1} \approx 3.337.$$

The quadratic is negative between its roots:

$$0.300 < \tan \theta < 3.337 \Leftrightarrow 16.7^\circ \leq \theta \leq 73.3^\circ.$$

Combining the upper and lower bounds for the crossbar condition:

$$(0.300 < \tan \theta < 0.439) \quad \text{or} \quad (3.197 < \tan \theta < 3.337).$$

Task 5. Intersect the wall condition ($0.347 < \tan \theta < 8.444$) with the crossbar condition above:

- Low branch: $\max(0.347, 0.300) < \tan \theta < \min(8.444, 0.439) \Rightarrow 0.347 < \tan \theta < 0.439$.
- High branch: $\max(0.347, 3.197) < \tan \theta < \min(8.444, 3.337) \Rightarrow 3.197 < \tan \theta < 3.337$.

In degrees:

$$\theta \in [19.1^\circ, 23.7^\circ] \quad \text{or} \quad \theta \in [72.6^\circ, 73.3^\circ].$$

The high branch corresponds to a very steep kick: the ball would be in the air for nearly 4 s, giving the goalkeeper ample time to react. It is not a practical direct free kick. The **physically sensible branch** is the low one: $\theta \in [19.1^\circ, 23.7^\circ]$.

An **optimal angle** is the midpoint of this window, maximising the vertical clearance margins both above the wall and below the crossbar:

$$\theta_{\text{opt}} \approx \frac{19.1^\circ + 23.7^\circ}{2} = 21.4^\circ \approx 21^\circ \text{ (to nearest degree).}$$

We adopt $\theta = 21^\circ$ for the remaining computations.

Task 6. For $\theta = 21^\circ$: $\cos 21^\circ \approx 0.9336$, $\sin 21^\circ \approx 0.3584$. Time to reach the goal line ($x = 22$):

$$t = \frac{22}{20 \cos 21^\circ} = \frac{22}{18.672} \approx 1.178 \text{ s.}$$

Velocity components at that instant:

$$v_x = 20 \cos 21^\circ \approx 18.67 \text{ m s}^{-1},$$

$$v_y = 20 \sin 21^\circ - 10t \approx 7.168 - 11.78 = -4.61 \text{ m s}^{-1}.$$

The negative sign confirms the ball is descending (dipping under the crossbar). The speed is


$$v = \sqrt{v_x^2 + v_y^2} = \sqrt{18.67^2 + 4.61^2} = \sqrt{348.6 + 21.3} \approx 19.2 \text{ m s}^{-1}.$$

Check height at goal line: $y(1.178) = 20(0.3584)(1.178) - 5(1.178)^2 \approx 8.44 - 6.94 = 1.50 \text{ m}$, safely below the bar.

Task 7. The vertical velocity is $v_y(t) = 7.17 - 10t$ (in m s^{-1}). This is a straight line with:

- slope $-g = -10 \text{ m s}^{-2}$,
- vertical intercept $v_y(0) = 7.17 \text{ m s}^{-1}$,
- zero at $t = 0.717 \text{ s}$ (instant of maximum height),
- value at goal line $t = 1.178 \text{ s}$: $v_y = -4.61 \text{ m s}^{-1}$.

The signed area between the graph and the t -axis represents the vertical displacement Δy .



Vertical component of velocity during the flight of the optimal free kick ($\theta = 21^\circ$).

Task 8. Possible answers (any one well argued):

- **Air resistance (drag):** reduces the horizontal speed and the maximum height, causing the ball to fall shorter and steeper than predicted. The real scoring window is narrower, and the optimal angle would be slightly lower.
- **The Magnus effect (spin):** a spinning ball experiences a force perpendicular to its velocity. Topspin makes the ball dip more sharply, helping it clear the wall and drop under the crossbar — this is exactly what elite free-kick takers exploit. Backspin would make it float, risking hitting the crossbar or going over.
- **Finite size of the ball / initial height:** the ball is struck slightly above the ground (on the boot), so $y(0) > 0$. This effectively raises the whole trajectory, slightly widening the window.

- **Wind:** a headwind increases drag and lift (if spinning), a tailwind does the opposite; cross-wind curves the trajectory laterally.

WARNING Marking points to watch

- Do not forget the factor $(1 + \tan^2 \theta)$ when replacing $1/\cos^2 \theta$ in the trajectory equation.
- A quadratic inequality in $\tan \theta$ is solved by studying the sign of the quadratic expression (using a sign table or sketch), not just by stating the roots.
- Remember that a velocity is a vector — the speed in Task 6 requires both components via Pythagoras.
- Always check the physical feasibility: the ball must be above ground ($y > 0$) at the goal line, not just below the crossbar.
- In the v_y-t graph, the slope is the acceleration ($-g$), the intercept is the initial vertical velocity, and the area under the curve gives the vertical displacement.

8 METHODS AND WORKED EXAMPLES

9 KINEMATICS OF PARTICLES IN A PLANE

9.1 Introduction and Syllabus Overview

In this chapter we extend the one-dimensional kinematics of Lower Sixth to motion in a plane. We describe the position of a particle by a vector $\vec{r}(t)$, obtain velocity and acceleration by differentiation, and recover them by integration when acceleration is known. The central application is **projectile motion** — motion under uniform gravity with no air resistance — where the horizontal and vertical motions are independent and linked only by the time t . We shall derive the standard formulas for range, maximum height and time of flight, solve problems where the launch and landing heights differ, analyse vertical projection (free fall and upward throws), and interpret motion graphs. This covers Lessons 114–122 of the official MINESEC/GCE Board progression.

DID YOU KNOW? Why vectors in kinematics?

In the Anglophone GCE Advanced Level, vectors $\vec{i}, \vec{j}$ are the standard language for planar motion. They allow us to treat the two perpendicular components independently while keeping the physical unity of the motion. This approach is also the gateway to the study of relative motion, circular motion and simple harmonic motion in later chapters.

9.2 Position, Velocity and Acceleration as Vectors in a Plane

DEFINITION Position vector

Let O be a fixed origin in the plane. The **position vector** of a particle P at time t is $\vec{r}(t) = \vec{OP}$. In Cartesian coordinates with constant unit vectors $\vec{i}$ (horizontal) and $\vec{j}$ (vertical), we write

$$\vec{r}(t) = x(t)\vec{i} + y(t)\vec{j},$$

where $x(t)$ and $y(t)$ are the scalar coordinates of P .

PROPERTY Velocity and acceleration by differentiation

If $\vec{r}(t) = x(t)\vec{i} + y(t)\vec{j}$ is differentiable, the **velocity** and **acceleration** vectors are

$$\vec{v}(t) = \frac{d\vec{r}}{dt} = \dot{x}(t)\vec{i} + \dot{y}(t)\vec{j}, \quad \vec{a}(t) = \frac{d\vec{v}}{dt} = \ddot{x}(t)\vec{i} + \ddot{y}(t)\vec{j}.$$

The **speed** is the magnitude $|\vec{v}| = \sqrt{\dot{x}^2 + \dot{y}^2}$; the magnitude of acceleration is $|\vec{a}| = \sqrt{\ddot{x}^2 + \ddot{y}^2}$.

METHOD Differentiating a position vector

1. Write $\vec{r}(t) = x(t)\vec{i} + y(t)\vec{j}$; identify $x(t)$ and $y(t)$.
2. Differentiate each component with respect to t to obtain $\vec{v}(t) = \dot{x}\vec{i} + \dot{y}\vec{j}$.
3. Differentiate again to obtain $\vec{a}(t) = \ddot{x}\vec{i} + \ddot{y}\vec{j}$.
4. For speed or magnitude of acceleration, compute the square root of the sum of squares of the components.
5. To find when the motion is parallel to an axis, set the other component of $\vec{v}$ to zero and solve for t .

Reflex: $\vec{i}$ and $\vec{j}$ are constant vectors, so only the scalar components are differentiated.

WORKED EXAMPLE 13 Velocity and acceleration from $\vec{r}(t)$

A particle P moves in the plane so that its position vector at time t seconds is

$$\vec{r}(t) = (2t^3 - 3t)\vec{i} + (t^2 + 4t)\vec{j} \text{ metres.}$$

- (a) Find the velocity of P at $t = 2$.
- (b) Find the acceleration of P and show that it is never zero.
- (c) Find the speed of P when $t = 1$.

SOLUTION

(a) Differentiate each component:

$$\vec{v}(t) = \frac{d\vec{r}}{dt} = (6t^2 - 3)\vec{i} + (2t + 4)\vec{j}.$$

At $t = 2$: $\vec{v}(2) = (6 \times 4 - 3)\vec{i} + (2 \times 2 + 4)\vec{j} = 21\vec{i} + 8\vec{j} \text{ m s}^{-1}$.

(b) Differentiate again:

$$\vec{a}(t) = \frac{d\vec{v}}{dt} = 12t\vec{i} + 2\vec{j}.$$

The $\vec{j}$ -component is the constant $2 \neq 0$, so $\vec{a}(t)$ can never be the zero vector. Hence the acceleration is never zero.

(c) At $t = 1$: $\vec{v}(1) = (6 - 3)\vec{i} + (2 + 4)\vec{j} = 3\vec{i} + 6\vec{j}$.

$$|\vec{v}(1)| = \sqrt{3^2 + 6^2} = \sqrt{45} = 3\sqrt{5} \approx 6.71 \text{ m s}^{-1}.$$

9.3 From Acceleration back to Velocity and Position: Integration in Kinematics

When the acceleration (or velocity) is given as a function of time, we recover the velocity (or position) by integration. Each integration introduces a constant vector which is determined by the initial conditions.

METHOD Integrating with initial conditions

Given $\vec{a}(t)$ (or $\vec{v}(t)$) and initial values $\vec{v}(t_0)$, $\vec{r}(t_0)$:

1. Integrate each component of $\vec{a}$ with respect to t to obtain $\vec{v}$; a constant of integration appears in each component (write them as $c_1\vec{i} + c_2\vec{j}$ or a constant vector $\vec{c}_1$).
2. Use the given velocity at a known time (often $t = 0$) to fix the constants.
3. Integrate $\vec{v}$ component by component to obtain $\vec{r}$; use the initial position to fix the new constants.
4. Write the final answers as single vectors in $\vec{i}, \vec{j}$ form.

Key idea: $\vec{v} = \int \vec{a} dt + \vec{c}_1$, $\vec{r} = \int \vec{v} dt + \vec{c}_2$. Never forget the constant vectors: they carry the physical initial conditions.

WORKED EXAMPLE 14 From acceleration back to position

A particle moves in a plane with acceleration $\vec{a}(t) = (6t)\vec{i} - 2\vec{j} \text{ m s}^{-2}$. Initially ($t = 0$) the particle is at the point with position vector $3\vec{j}$ metres and is moving with velocity $4\vec{i} \text{ m s}^{-1}$.

- (a) Find the velocity of the particle at time t .
- (b) Find the position vector of the particle at time t .

SOLUTION

(a) Integrate the acceleration component by component:

$$\vec{v}(t) = \int \vec{a} dt = (3t^2 + c_1)\vec{i} + (-2t + c_2)\vec{j}.$$

At $t = 0$, $\vec{v} = 4\vec{i}$, so $c_1 = 4$ and $c_2 = 0$. Hence

$$\vec{v}(t) = (3t^2 + 4)\vec{i} - 2t\vec{j} \text{ m s}^{-1}.$$

(b) Integrate the velocity:

$$\vec{r}(t) = \int \vec{v} dt = (t^3 + 4t + k_1)\vec{i} + (-t^2 + k_2)\vec{j}.$$

At $t = 0$, $\vec{r} = 3\vec{j}$, so $k_1 = 0$ and $k_2 = 3$. Therefore

$$\vec{r}(t) = (t^3 + 4t)\vec{i} + (3 - t^2)\vec{j} \text{ metres.}$$

Check: differentiating $\vec{r}$ twice returns $6t\vec{i} - 2\vec{j} = \vec{a}$. ✓

9.4 Projectile Motion: Definition, Model and Equations

DEFINITION Projectile

A **projectile** is a particle projected in a vertical plane and moving under the action of gravity alone. The standard model assumes:

- The acceleration due to gravity $\vec{g}$ is constant in magnitude ($g \approx 9.8 \text{ m s}^{-2}$ or 10 m s^{-2} as instructed) and direction (vertically downwards).
- Air resistance is negligible.
- The Earth's curvature and rotation are ignored over the range of motion.

PROPERTY Equations of motion of a projectile launched from the origin

A projectile is launched from the origin O with initial speed u at an angle α above the horizontal. Take $\vec{i}$ horizontal (positive in the direction of projection) and $\vec{j}$ vertically upward. Then $\vec{a} = -g\vec{j}$. Resolving the initial velocity: $u_x = u \cos \alpha$, $u_y = u \sin \alpha$.

Horizontal motion (no acceleration): $x(t) = u \cos \alpha \cdot t$, $v_x = u \cos \alpha$ (constant).

Vertical motion (constant acceleration $-g$): $y(t) = u \sin \alpha \cdot t - \frac{1}{2}gt^2$, $v_y = u \sin \alpha - gt$.

The **trajectory equation** (Cartesian equation) is obtained by eliminating t :

$$y = x \tan \alpha - \frac{gx^2}{2u^2 \cos^2 \alpha} = x \tan \alpha - \frac{gx^2}{2u^2} (1 + \tan^2 \alpha).$$

This is a parabola with vertical axis, opening downwards.

KEY POINTS Independence of motions

The horizontal and vertical motions are independent; only the time t links them. The horizontal velocity remains constant throughout the flight; the vertical velocity changes exactly as in one-dimensional motion under constant acceleration $-g$.

WORKED EXAMPLE 15 Trajectory of a stone

A stone is thrown from the top of a cliff with speed 20 m s^{-1} at an angle of 30° above the horizontal. Take $g = 10 \text{ m s}^{-2}$ and the point of projection as origin.

- (a) Write down $x(t)$ and $y(t)$.
 (b) Find the Cartesian equation of the trajectory.
 (c) Find the position and the velocity of the stone at $t = 1 \text{ s}$.

SOLUTION

(a) $u \cos 30^\circ = 20 \times \frac{\sqrt{3}}{2} = 10\sqrt{3}$, $u \sin 30^\circ = 20 \times \frac{1}{2} = 10$. Hence

$$x(t) = 10\sqrt{3}t, \quad y(t) = 10t - 5t^2.$$

(b) From $x = 10\sqrt{3}t$, $t = \frac{x}{10\sqrt{3}}$. Substituting:

$$y = 10 \cdot \frac{x}{10\sqrt{3}} - 5 \left(\frac{x}{10\sqrt{3}} \right)^2 = \frac{x}{\sqrt{3}} - \frac{x^2}{60}.$$

This is a parabola opening downwards, as expected.

(c) At $t = 1$: $x = 10\sqrt{3} \approx 17.3 \text{ m}$, $y = 10 - 5 = 5 \text{ m}$. So the stone is at $(17.3; 5) \text{ m}$. Velocity: $v_x = 10\sqrt{3} \text{ m s}^{-1}$ (constant), $v_y = 10 - 10 \times 1 = 0$. The vertical velocity is zero: at $t = 1 \text{ s}$ the stone is at the top of its path, moving horizontally with speed $10\sqrt{3} \approx 17.3 \text{ m s}^{-1}$.

9.5 Range, Maximum Height and Time of Flight

The following formulas apply only when the projectile is launched from and lands on the same horizontal level (level ground).

PROPERTY Standard projectile formulas (level ground)

For a projectile launched from level ground at speed u , angle α ($0 < \alpha < 90^\circ$):

1. **Time of flight:** $T = \frac{2u \sin \alpha}{g}$ (solve $y = 0$, $t \neq 0$).

2. **Maximum height:** $H = \frac{u^2 \sin^2 \alpha}{2g}$ (set $v_y = 0$ and use $v_y^2 = u_y^2 - 2gH$).
3. **Range:** $R = \frac{u^2 \sin 2\alpha}{g}$ (compute $x(T) = u \cos \alpha \cdot T$).
4. **Maximum range** for a given u : $R_{\max} = \frac{u^2}{g}$ at $\alpha = 45^\circ$.
5. **Complementary angles** α and $90^\circ - \alpha$ give the same range because $\sin(2(90^\circ - \alpha)) = \sin(180^\circ - 2\alpha) = \sin 2\alpha$.

METHOD Using the standard projectile formulas

1. Verify that launch and landing are at the same height. If not, go back to $x(t)$, $y(t)$.
2. Identify u , α , g .
3. Substitute into T , H , R formulas.
4. For complementary angle problems, state $90^\circ - \alpha$ and note the different T and H .

Warning: these formulas require launch and landing at the same height. Otherwise return to the equations of motion.

WORKED EXAMPLE 16 A footballer's kick

A footballer kicks a ball with speed 25 m s^{-1} at 40° above the horizontal from level ground. Take $g = 10 \text{ m s}^{-2}$, $\sin 40^\circ \approx 0.643$, $\cos 40^\circ \approx 0.766$, $\sin 80^\circ \approx 0.985$.

- (a) Find the time of flight.
- (b) Find the range.
- (c) Find the greatest height reached.
- (d) At what other angle, with the same speed, would the ball have the same range?

SOLUTION

$$(a) T = \frac{2u \sin \alpha}{g} = \frac{2 \times 25 \times 0.643}{10} = 3.22 \text{ s.}$$

$$(b) R = \frac{u^2 \sin 2\alpha}{g} = \frac{625 \times 0.985}{10} \approx 61.6 \text{ m.}$$

$$(c) H = \frac{u^2 \sin^2 \alpha}{2g} = \frac{625 \times (0.643)^2}{20} = \frac{625 \times 0.4134}{20} \approx 12.9 \text{ m.}$$

(d) The complementary angle $90^\circ - 40^\circ = 50^\circ$ gives the same range, since $\sin(2 \times 50^\circ) = \sin 100^\circ = \sin 80^\circ$. The ball would then go higher ($H \approx 18.3$ m) but stay longer in the air ($T \approx 3.83$ s).

9.6 Applications and Problem-Solving with Projectiles

When the launch and landing heights differ, or a specific target must be hit, the standard formulas do not apply. We must work from the parametric equations or the trajectory equation.

METHOD Non-level problems: work from the equations

1. Choose axes and write $x(t) = u \cos \alpha t$ and $y(t) = u \sin \alpha t - \frac{1}{2}gt^2$ with the correct signs for the initial position (e.g. if launched from a height h above the landing level, set $y(0) = h$ or $y(0) = 0$ and landing at $y = -h$).
2. Impose the condition of the problem (e.g. $y = -h$ at landing, or the trajectory passes through (x_0, y_0)).
3. Solve the resulting equation — often a quadratic in t or in $\tan \alpha$; keep the physically meaningful root ($t > 0$, α acute usually).
4. For "passes through (x_0, y_0) " problems, use the trajectory equation $y_0 = x_0 \tan \alpha - \frac{gx_0^2}{2u^2}(1 + \tan^2 \alpha)$ to get a quadratic in $\tan \alpha$.

Reflex: a quadratic in $\tan \alpha$ with two positive roots means the target can be hit by two possible throws (a low one and a high one).

WORKED EXAMPLE 17 Ball thrown from a balcony

From a balcony 15 m above the ground, a student throws a ball with speed 20 m s^{-1} at 30° above the horizontal. Take $g = 10 \text{ m s}^{-2}$.

- (a) Find the time at which the ball hits the ground.
- (b) Find the horizontal distance travelled before landing.
- (c) Find the speed of the ball just before impact.

SOLUTION

Take the point of projection as origin, y upwards. Then $u_x = 20 \cos 30^\circ = 10\sqrt{3}$, $u_y = 20 \sin 30^\circ = 10$, and the ground is at $y = -15$.

(a) Solve $y(t) = -15$:

$$10t - 5t^2 = -15 \implies 5t^2 - 10t - 15 = 0 \implies t^2 - 2t - 3 = 0.$$

Factorising: $(t - 3)(t + 1) = 0$, so $t = 3$ s (the root $t = -1$ is rejected).

(b) $x(3) = 10\sqrt{3} \times 3 = 30\sqrt{3} \approx 52.0$ m.

(c) At $t = 3$: $v_x = 10\sqrt{3} \text{ m s}^{-1}$, $v_y = 10 - 10 \times 3 = -20 \text{ m s}^{-1}$.

$$|\vec{v}| = \sqrt{(10\sqrt{3})^2 + (-20)^2} = \sqrt{300 + 400} = \sqrt{700} = 10\sqrt{7} \approx 26.5 \text{ m s}^{-1}.$$

Note that the impact speed exceeds the launch speed: the ball has gained kinetic energy by falling 15 m.

WORKED EXAMPLE 18 Hitting a target on a wall

A particle is projected from ground level with speed 25 m s^{-1} towards a vertical wall 30 m away. The particle just passes over the top of the wall which is 10 m high. Find the two possible angles of projection. Take $g = 10 \text{ m s}^{-2}$.

SOLUTION

The trajectory must pass through (30, 10). Using $y = x \tan \alpha - \frac{gx^2}{2u^2}(1 + \tan^2 \alpha)$:

$$10 = 30 \tan \alpha - \frac{10 \times 30^2}{2 \times 25^2}(1 + \tan^2 \alpha) = 30 \tan \alpha - \frac{9000}{1250}(1 + \tan^2 \alpha) = 30 \tan \alpha - 7.2(1 + \tan^2 \alpha).$$

Rearrange: $7.2 \tan^2 \alpha - 30 \tan \alpha + 17.2 = 0$. Multiply by 5: $36 \tan^2 \alpha - 150 \tan \alpha + 86 = 0$, divide by 2: $18 \tan^2 \alpha - 75 \tan \alpha + 43 = 0$.

$$\tan \alpha = \frac{75 \pm \sqrt{5625 - 3096}}{36} = \frac{75 \pm \sqrt{2529}}{36} \approx \frac{75 \pm 50.29}{36}.$$

$\tan \alpha_1 \approx 3.48 \implies \alpha_1 \approx 74^\circ$ (high trajectory), $\tan \alpha_2 \approx 0.686 \implies \alpha_2 \approx 34.5^\circ$ (low trajectory). Both are valid.

9.7 Free Fall, Vertical Projection and Motion Graphs

When the motion is purely vertical, we have one-dimensional motion with constant acceleration $\pm g$. The vector notation reduces to a sign convention.

METHOD One-dimensional vertical motion

All motion is on a vertical line; choose an origin and a positive direction, then use the

constant-acceleration formulas with $a = \pm g$:

$$v = u + at, \quad s = ut + \frac{1}{2}at^2, \quad v^2 = u^2 + 2as.$$

1. Dropped object (free fall): $u = 0$; taking downwards positive, $v = gt$, $h = \frac{1}{2}gt^2$.
2. Thrown upwards: take upwards positive, $a = -g$. At the highest point $v = 0$.
3. Time to the top: $t_{\text{up}} = \frac{u}{g}$; greatest height: $H = \frac{u^2}{2g}$.
4. Returning to the launch level, the object arrives with speed u (same speed, opposite direction) after total time $\frac{2u}{g}$.

Warning: fix the sign convention before substituting, and keep it throughout the whole solution.

WORKED EXAMPLE 19 Stone thrown upwards from a well

A stone is thrown vertically upwards from the top of a well with speed 14 m s^{-1} . It falls into the well and reaches the water surface 20 m below the point of projection. Take $g = 9.8 \text{ m s}^{-2}$.

- (a) Find the greatest height reached above the point of projection.
- (b) Find the total time of flight until the stone hits the water.
- (c) Find the speed of impact with the water.

SOLUTION

Take upwards as positive, origin at the point of projection; $u = 14$, $a = -9.8$.

(a) At the top, $v = 0$: $v^2 = u^2 + 2as$ gives $0 = 14^2 - 2 \times 9.8 \times H$, so

$$H = \frac{196}{19.6} = 10 \text{ m}.$$

(b) The water is at $s = -20 \text{ m}$. Using $s = ut + \frac{1}{2}at^2$:

$$-20 = 14t - 4.9t^2 \implies 4.9t^2 - 14t - 20 = 0.$$

By the quadratic formula:

$$t = \frac{14 \pm \sqrt{196 + 392}}{9.8} = \frac{14 \pm \sqrt{588}}{9.8} \approx \frac{14 \pm 24.25}{9.8}.$$

Keeping the positive root: $t \approx \frac{38.25}{9.8} \approx 3.90 \text{ s}$.

(c) $v = u + at = 14 - 9.8 \times 3.90 \approx -24.2 \text{ m s}^{-1}$. The impact speed is about 24.2 m s^{-1} downwards. Check with $v^2 = u^2 + 2as = 196 - 19.6 \times (-20) = 588$, so $|v| = \sqrt{588} \approx 24.2 \text{ m s}^{-1}$. ✓

9.8 Graphical Representation of Vertical Motion

For an object thrown vertically (acceleration $-g$, constant), the graphs are powerful tools for visualising and solving problems.

METHOD Graphs of vertical motion

For an object thrown vertically upwards with initial speed u from a reference level:

1. Velocity–time graph: a straight line of gradient $-g$, starting at $v = u$. The area between the line and the time axis gives the displacement (areas below the axis count negatively).
2. Displacement–time graph: a parabola $s = ut - \frac{1}{2}gt^2$; its gradient at any point is the velocity; the vertex corresponds to the highest point ($v = 0$).
3. Speed–time graph: the velocity graph reflected above the axis where $v < 0$ — it is V-shaped for an up–down flight. Speed is never negative.
4. Read off: time at the top (where $v = 0$), landing time (where s returns to its initial value), impact velocity (final value of v).

Reflex: gradient of s – t graph = velocity; gradient of v – t graph = acceleration; area under v – t graph = displacement.


WORKED EXAMPLE 20 Graphs for a ball thrown upwards

A ball is thrown vertically upwards with speed 20 m s^{-1} from ground level and is caught again at the same level. Take $g = 10 \text{ m s}^{-2}$.

- (a) Sketch the velocity–time graph for the whole flight.
- (b) Use the graph to find the total time of flight and the greatest height.
- (c) Describe the displacement–time graph.

SOLUTION

- (a) $v = 20 - 10t$: a straight line from $(0, 20)$ with gradient -10 , crossing the axis at $t = 2$ and reaching $v = -20$ at $t = 4$.



Velocity-time graph of the ball: gradient $-g = -10 \text{ m s}^{-2}$.

(b) The ball returns when the net area is zero. The two triangles have equal area $A = \frac{1}{2} \times 2 \times 20 = 20$, one positive, one negative, so the displacement is zero at $t = 4 \text{ s}$: the total flight time is 4 s. The greatest height is the positive area up to $t = 2$: $H = 20 \text{ m}$. (Check: $H = \frac{u^2}{2g} = \frac{400}{20} = 20 \text{ m}$. ✓)

(c) The displacement-time graph is the parabola $s = 20t - 5t^2$, starting at the origin, rising to a maximum of 20 m at $t = 2$ (where its gradient is zero, matching $v = 0$), and returning to $s = 0$ at $t = 4 \text{ s}$. Its gradient decreases steadily, reflecting the constant acceleration -10 m s^{-2} .

WARNING Frequent errors to avoid

- Mixing sign conventions inside one calculation: decide the positive direction first and write it down.
- Using $R = \frac{u^2 \sin 2\alpha}{g}$ when the landing point is not at the launch height — return to $x(t)$, $y(t)$.
- Forgetting that v_x is constant: at the top of a projectile path the velocity is not zero, only v_y is.
- Confusing speed and velocity on graphs: speed is $|\vec{v}|$ and is never negative.
- Dropping the constants of integration when working back from acceleration: they are fixed by the initial conditions.
- Using $g = 9.8$ and $g = 10$ in the same problem; follow the instruction in the question.

9.9 Consolidation Exercises

EXERCISE 7 Exercise 1: Vector kinematics

The position vector of a particle P at time t seconds is $\vec{r}(t) = (t^3 - 6t)\vec{i} + (4t^2 - 8t)\vec{j}$ metres.

1. Find the velocity and acceleration vectors at time t .
2. Find the time when P is moving parallel to the $\vec{j}$ -axis.
3. Find the speed of P when $t = 2$.

EXERCISE 8 Exercise 2: Integration from acceleration

A particle moves in the plane with acceleration $\vec{a}(t) = (4t - 2)\vec{i} + 6\vec{j} \text{ m s}^{-2}$. At $t = 0$, its velocity is $3\vec{i} - 2\vec{j} \text{ m s}^{-1}$ and its position vector is $5\vec{i} + 4\vec{j}$ metres. Find expressions for $\vec{v}(t)$ and $\vec{r}(t)$.

EXERCISE 9 Exercise 3: Projectile on level ground

A projectile is fired from ground level with speed 30 m s^{-1} at an angle α where $\sin \alpha = 0.6$ and $\cos \alpha = 0.8$. Take $g = 10 \text{ m s}^{-2}$.

1. Find the time of flight, the range and the maximum height.
2. Find the velocity of the projectile 2 s after projection.

EXERCISE 10 Exercise 4: Projectile from a height

A stone is thrown horizontally with speed 15 m s^{-1} from the top of a vertical cliff 80 m high. Take $g = 10 \text{ m s}^{-2}$.

1. Find the time taken to reach the sea.
2. Find the horizontal distance from the foot of the cliff where the stone hits the sea.
3. Find the speed and direction of motion at impact.

EXERCISE 11 Exercise 5: Vertical motion graphs

A ball is projected vertically upwards with speed 25 m s^{-1} from a point 5 m above the ground. Take $g = 10 \text{ m s}^{-2}$.

1. Sketch the velocity–time graph until the ball hits the ground.

2. Use the graph to find the total time of flight and the maximum height above the ground.
3. Sketch the corresponding displacement–time graph (displacement measured from the ground).

SOLUTION Exercise 1

1. $\vec{v}(t) = (3t^2 - 6)\vec{i} + (8t - 8)\vec{j}$; $\vec{a}(t) = 6t\vec{i} + 8\vec{j}$.
2. Parallel to $\vec{j}$ -axis $\Rightarrow v_x = 0 \Rightarrow 3t^2 - 6 = 0 \Rightarrow t = \sqrt{2}$ s ($t > 0$).
3. At $t = 2$: $\vec{v} = (6)\vec{i} + (8)\vec{j}$; speed $= \sqrt{36 + 64} = 10$ m s⁻¹.

SOLUTION Exercise 2

$$\vec{v}(t) = \int \vec{a} dt = (2t^2 - 2t + 3)\vec{i} + (6t - 2)\vec{j}. \quad \vec{r}(t) = \int \vec{v} dt = \left(\frac{2}{3}t^3 - t^2 + 3t + 5\right)\vec{i} + (3t^2 - 2t + 4)\vec{j}.$$

SOLUTION Exercise 3

$$u = 30, \sin \alpha = 0.6, \cos \alpha = 0.8, g = 10. \quad T = \frac{2 \times 30 \times 0.6}{10} = 3.6 \text{ s}. \quad R = \frac{30^2 \times 2 \times 0.6 \times 0.8}{10} = 86.4 \text{ m}.$$

$$H = \frac{30^2 \times 0.6^2}{20} = 16.2 \text{ m}. \quad \text{At } t = 2: v_x = 24 \text{ m s}^{-1}, v_y = 18 - 20 = -2 \text{ m s}^{-1}; \vec{v} = 24\vec{i} - 2\vec{j} \text{ m s}^{-1}.$$

SOLUTION Exercise 4

Horizontal throw $\Rightarrow \alpha = 0, u_x = 15, u_y = 0$. Origin at cliff top, y up, ground at $y = -80$.
 $y(t) = -5t^2 = -80 \Rightarrow t = 4$ s. $x(4) = 15 \times 4 = 60$ m. At $t = 4$: $v_x = 15, v_y = -40$; speed
 $= \sqrt{225 + 1600} = \sqrt{1825} \approx 42.7$ m s⁻¹; angle below horizontal $\theta = \arctan(40/15) \approx 69.4^\circ$.

SOLUTION Exercise 5

Upwards positive, origin at ground. Initial $s = 5, u = 25, a = -10$. $v = 25 - 10t$. Hits ground when $s = 0$: $5 + 25t - 5t^2 = 0 \Rightarrow t^2 - 5t - 1 = 0 \Rightarrow t = \frac{5 + \sqrt{29}}{2} \approx 5.19$ s. Max height: $v = 0 \Rightarrow t = 2.5$ s, $s_{\max} = 5 + 25(2.5) - 5(2.5)^2 = 36.25$ m. v - t graph: line from $(0, 25)$ to $(5.19, -26.9)$, crossing t -axis at 2.5. s - t graph: parabola from $(0, 5)$ to $(5.19, 0)$, vertex at $(2.5, 36.25)$.

10 EXERCISES**10.1 I check my knowledge****EXERCISE 12 Multiple choice questions**

For each question, choose the single correct answer, giving a brief justification.

1. A particle has position vector $\vec{r}(t) = (3t^2)\vec{i} + (4t - t^2)\vec{j}$ metres. Its velocity at time t is:
(a) $(6t)\vec{i} + (4 - 2t)\vec{j}$
(b) $(3t^2)\vec{i} + (4 - 2t)\vec{j}$
(c) $(6t)\vec{i} + (4t - 2t)\vec{j}$
(d) $(6)\vec{i} + (-2)\vec{j}$
2. A projectile is launched from level ground. At the highest point of its trajectory:
(a) its velocity and acceleration are both zero;
(b) its velocity is zero but its acceleration is g downwards;
(c) the horizontal component of its velocity is zero;
(d) the vertical component of its velocity is zero and its acceleration is g downwards.
3. For a projectile launched at speed u at angle θ on level ground, the range is:
(a) $\frac{u^2 \sin \theta}{g}$
(b) $\frac{u^2 \sin 2\theta}{g}$
(c) $\frac{u^2 \sin^2 \theta}{2g}$
(d) $\frac{2u \sin \theta}{g}$
4. A stone is dropped from rest from a height of 20 m. Taking $g = 10 \text{ m s}^{-2}$, the time taken to reach the ground is:
(a) 1 s
(b) $\sqrt{2}$ s
(c) 2 s
(d) 4 s

EXERCISE 13 True or false?

State whether each assertion is true or false, justifying your answer carefully.

1. If the acceleration of a particle is constant, then its velocity is constant.
2. Two projectiles launched from the same point on level ground with the same speed at complementary angles θ and $90^\circ - \theta$ have the same range.
3. A ball thrown vertically upwards has zero acceleration at the top of its flight.
4. The horizontal component of the velocity of a projectile remains constant throughout the flight (air resistance neglected).

EXERCISE 14 Key definitions

1. Define the **velocity** and the **acceleration** of a particle whose position vector is $\vec{r}(t)$.
2. State the two components of the acceleration of a projectile in the standard model (no air resistance), and justify them.
3. Define the **range**, the **time of flight** and the **maximum height** of a projectile.
4. What is meant by **free fall**? State the value of the acceleration of a body in free fall near the Earth's surface.

EXERCISE 15 Direct calculations

Take $g = 10 \text{ m s}^{-2}$ throughout.

1. A particle has velocity $\vec{v}(t) = (4t)\vec{i} + (3 - 2t)\vec{j} \text{ m s}^{-1}$. Find its acceleration and its speed at $t = 2$.
2. A stone is dropped from rest from a bridge and hits the water 3 s later. Find the height of the bridge above the water and the speed of impact.
3. A projectile is launched at 20 m s^{-1} at 30° above the horizontal. Find the horizontal and vertical components of its initial velocity.
4. A ball is thrown vertically upwards at 15 m s^{-1} . Find the greatest height reached and the time taken to reach it.

10.2 I practise

EXERCISE 16 Differentiating a position vector

A particle moves in a plane with position vector

$$\vec{r}(t) = (2t^2 - 3t)\vec{i} + (t^3 - 6t)\vec{j} \quad \text{metres.}$$

1. Find its velocity $\vec{v}(t)$ and acceleration $\vec{a}(t)$.
2. Find its velocity and acceleration at $t = 2$, and its speed at that instant.
3. Find the times when the particle moves parallel to the y -axis.
4. Find the Cartesian equation of its path.

EXERCISE 17 Integrating an acceleration

A particle starts from the origin O with velocity $(4\vec{i} + 3\vec{j}) \text{ m s}^{-1}$ and moves with acceleration

$$\vec{a}(t) = (2t\vec{i} - 6\vec{j}) \text{ m s}^{-2}.$$

1. Find its velocity vector at time t .
2. Find its position vector at time t .
3. Find its distance from O when $t = 3$.

EXERCISE 18 Standard projectile on level ground

A stone is projected at 25 m s^{-1} at 40° above the horizontal from level ground. Taking $g = 10 \text{ m s}^{-2}$, find:

1. the time of flight;
2. the maximum height reached;
3. the range on the horizontal plane through the point of projection.

(Give your answers to 3 significant figures; you may use $\sin 40^\circ \approx 0.643$ and $\cos 40^\circ \approx 0.766$.)

EXERCISE 19 Horizontal projection from a cliff

A ball is thrown horizontally at 12 m s^{-1} from the top of a cliff 45 m high, and falls into the sea. Taking $g = 10 \text{ m s}^{-2}$, find:

1. the time of flight;
2. the horizontal distance from the foot of the cliff to the point of impact;
3. the velocity (magnitude and direction) of the ball on hitting the water.

10.3 I prepare for the GCE

EXERCISE 20 Clearing the wall and scoring (GCE style)

(13 marks)

A footballer kicks a ball from ground level with speed 16 m s^{-1} at an angle θ above the horizontal. Take $g = 10 \text{ m s}^{-2}$ and neglect air resistance.

1. Write down the horizontal and vertical components of the displacement of the ball at time t , and hence show that the Cartesian equation of the trajectory is

$$y = x \tan \theta - \frac{5x^2}{128} (1 + \tan^2 \theta). \text{ (4 marks)}$$

2. A vertical wall 2 m high stands 9 m from the kicker. Show that for the ball to clear the wall, $\tan \theta$ must satisfy a certain inequality. (3 marks)
3. The crossbar of the goal, 2.4 m high, is 18 m away. Show that for the ball to pass under the crossbar and above the goal line at that point, $\tan \theta$ satisfies a quadratic equation at the limiting case, and find the possible values of θ for which the ball both clears the wall and enters the goal. (6 marks)

EXERCISE 21 Projectile from a rooftop (GCE style)

(12 marks)

A projectile is launched from the top of a building 20 m high with speed 15 m s^{-1} at 30° above the horizontal, and lands in the horizontal school field below. Take $g = 10 \text{ m s}^{-2}$.

1. Write down the equations giving the horizontal and vertical displacements of the projectile at time t after launch, measured from the foot of the building. (3 marks)
2. Show that the time of flight T satisfies $5T^2 - 7.5T - 20 = 0$, and hence find T . (4 marks)
3. Find the horizontal distance travelled by the projectile before landing. (2 marks)
4. Find the speed of the projectile and the angle its velocity makes with the horizontal at the instant it lands. (3 marks)

EXERCISE 22 Vertical motion and its graphs (GCE style)**(14 marks)**

Take $g = 10 \text{ m s}^{-2}$ throughout this question.

1. A mango falls from rest from a tree and takes 1.5 s to reach the ground. From what height did it fall, and with what speed does it hit the ground? (3 marks)
2. A stone is thrown vertically upwards at 20 m s^{-1} from the edge of a well and falls to the bottom of the well, 25 m below the point of projection. Find the total time of flight and the speed of the stone at the bottom. (4 marks)
3. An object is released from a balloon which is rising vertically at 8 m s^{-1} when the balloon is 100 m above the ground. Explain why the object initially moves upwards, and find the time taken for the object to reach the ground and its speed on impact. (4 marks)
4. A ball is thrown vertically upwards at 14 m s^{-1} and returns to the thrower's hand. Sketch, on separate axes, the displacement–time, velocity–time and acceleration–time graphs from launch until the ball returns to the thrower's hand, marking all key values. (3 marks)

11 SOLUTIONS TO THE EXERCISES

12 SOLUTIONS DÉTAILLÉES DES EXERCICES DU CHAPITRE PMM11

SOLUTION Exercice 1 – Questions à choix multiple

- Réponse (a). La vitesse est la dérivée de la position : $\vec{v}(t) = \frac{d\vec{r}}{dt} = (6t)\vec{i} + (4 - 2t)\vec{j}$. Les options (b) et (c) conservent des termes non dérivés ; (d) est l'accélération, pas la vitesse.
- Réponse (d). Au point le plus haut, la composante verticale de la vitesse est instantanément zéro (le projectile cesse de monter et commence à descendre), mais la composante horizontale $u \cos \theta$ est inchangée, et l'accélération est $\vec{g}$ vers le bas pendant tout le vol. Donc (a), (b) et (c) sont toutes fausses.
- Réponse (b). La portée sur un sol horizontal est $R = \frac{u^2 \sin 2\theta}{g}$. L'option (c) est la hauteur maximale ; l'option (d) est le temps de vol.
- Réponse (c). Pour un corps lâché sans vitesse initiale, $h = \frac{1}{2}gt^2$, donc $20 = \frac{1}{2}(10)t^2$, d'où $t^2 = 4$ et $t = 2$ s.

SOLUTION Exercice 2 – Vrai ou faux ?

- Faux. Une accélération constante signifie $\vec{v}(t) = \vec{u} + \vec{a}t$: la vitesse change uniformément. Une accélération constante implique une vitesse constante seulement si $\vec{a} = \vec{0}$. Contre-exemple : un projectile a une accélération constante $\vec{g}$ mais sa vitesse change continuellement.
- Vrai. $R = \frac{u^2 \sin 2\theta}{g}$, et $\sin 2(90^\circ - \theta) = \sin(180^\circ - 2\theta) = \sin 2\theta$. Donc des angles complémentaires donnent la même portée (mais des hauteurs maximales et des temps de vol différents).
- Faux. Au sommet du vol, la vitesse est zéro, mais l'accélération est $g \approx 9.81 \text{ m s}^{-2}$ vers le bas pendant tout le vol – la gravitation ne s'arrête jamais. Si l'accélération était nulle au sommet, le ballon y resterait.
- Vrai. En négligeant la résistance de l'air, la seule force est le poids, qui est vertical.

Donc l'accélération horizontale est nulle, donc $v_x = u \cos \theta$ est constante pendant tout le vol.

SOLUTION Exercice 3 – Définitions clés

1. La **vitesse** est le taux de variation de la position : $\vec{v}(t) = \frac{d\vec{r}}{dt}$. L'**accélération** est le taux de variation de la vitesse : $\vec{a}(t) = \frac{d\vec{v}}{dt} = \frac{d^2\vec{r}}{dt^2}$.
2. Dans le modèle standard du projectile, $\vec{a} = -g\vec{j}$, c'est-à-dire composante horizontale $a_x = 0$ (aucune force horizontale, donc aucune accélération horizontale) et composante verticale $a_y = -g$ (le poids est la seule force, agissant verticalement vers le bas, et d'après la deuxième loi de Newton $m\vec{a} = m\vec{g}$).
3. Le **temps de vol** est le temps total entre le lancement et l'atterrissage. La **portée** est la distance horizontale entre le point de lancement et le point d'atterrissage. La **hauteur maximale** est le plus grand déplacement vertical atteint au-dessus du point de lancement.
4. La **chute libre** est le mouvement d'un corps soumis à son poids seul (résistance de l'air négligée). Près de la surface de la Terre, l'accélération de la chute libre est $g \approx 9.81 \text{ m s}^{-2}$ (souvent prise comme 10 m s^{-2} dans les exercices), dirigée verticalement vers le bas.

SOLUTION Exercice 4 – Calculs directs

1. Accélération : $\vec{a} = \frac{d\vec{v}}{dt} = 4\vec{i} - 2\vec{j} \text{ m s}^{-2}$ (constante). à $t = 2$: $\vec{v}(2) = 8\vec{i} - 1\vec{j}$, donc la vitesse (module) est $|\vec{v}(2)| = \sqrt{8^2 + (-1)^2} = \sqrt{65} \approx 8.06 \text{ m s}^{-1}$.
2. Lâché sans vitesse initiale : $h = \frac{1}{2}gt^2 = \frac{1}{2}(10)(3^2) = 45 \text{ m}$. Vitesse à l'impact : $v = gt = 10 \times 3 = 30 \text{ m s}^{-1}$.
3. $u_x = u \cos \theta = 20 \cos 30^\circ = 20 \times \frac{\sqrt{3}}{2} = 10\sqrt{3} \approx 17.3 \text{ m s}^{-1}$; $u_y = u \sin \theta = 20 \sin 30^\circ = 10 \text{ m s}^{-1}$.
4. Sens vers le haut positif : $v = u - gt$. à la plus grande hauteur $v = 0$, donc $t = \frac{u}{g} = \frac{15}{10} = 1.5 \text{ s}$. Hauteur : $h = ut - \frac{1}{2}gt^2 = 15(1.5) - 5(1.5)^2 = 22.5 - 11.25 = 11.25 \text{ m}$ (équivalentment $h = \frac{u^2}{2g} = \frac{225}{20}$).

SOLUTION Exercice 5 – Dérivation d'un vecteur position

1. En dérivant composante par composante :

$$\vec{v}(t) = \frac{d\vec{r}}{dt} = (4t - 3)\vec{i} + (3t^2 - 6)\vec{j}, \quad \vec{a}(t) = \frac{d\vec{v}}{dt} = 4\vec{i} + 6t\vec{j}.$$

2. à
- $t = 2$
- :
- $\vec{v}(2) = 5\vec{i} + 6\vec{j} \text{ m s}^{-1}$
- et
- $\vec{a}(2) = 4\vec{i} + 12\vec{j} \text{ m s}^{-2}$
- . Vitesse :
- $|\vec{v}(2)| = \sqrt{25 + 36} = \sqrt{61} \approx 7.81 \text{ m s}^{-1}$
- .

3. Le mouvement est parallèle à l'axe
- y
- lorsque la composante horizontale de la vitesse est nulle (et la verticale non nulle) :
- $4t - 3 = 0 \Rightarrow t = \frac{3}{4} \text{ s}$
- . (Vérification :
- $3t^2 - 6 = \frac{27}{16} - 6 \neq 0$
- , la particule bouge bien.)

4. On a
- $x = 2t^2 - 3t$
- et
- $y = t^3 - 6t = t(t^2 - 6)$
- . Le paramètre
- t
- ne peut pas être éliminé par une algèbre élémentaire sous une forme close simple car
- x
- est quadratique et
- y
- cubique en
- t
- ; la trajectoire est donnée sous forme paramétrique par

$$x = 2t^2 - 3t, \quad y = t^3 - 6t,$$

et l'élimination de t (par exemple en résolvant $2t^2 - 3t - x = 0$, $t = \frac{3 \pm \sqrt{9 + 8x}}{4}$, et en substituant dans y) donne la relation cartésienne

$$y = \left(\frac{3 \pm \sqrt{9 + 8x}}{4} \right) \left(\left(\frac{3 \pm \sqrt{9 + 8x}}{4} \right)^2 - 6 \right),$$

qu'il est préférable de laisser sous forme paramétrique. (Au niveau GCE, écrire les équations paramétriques et la procédure de substitution rapporte tous les points.)

SOLUTION Exercice 6 – Intégration d'une accélération

- 1.
- $\vec{v}(t) = \int \vec{a} dt = (t^2 + c_1)\vec{i} + (-6t + c_2)\vec{j}$
- . à
- $t = 0$
- ,
- $\vec{v} = 4\vec{i} + 3\vec{j}$
- , donc
- $c_1 = 4$
- ,
- $c_2 = 3$
- :

$$\vec{v}(t) = (t^2 + 4)\vec{i} + (3 - 6t)\vec{j} \text{ m s}^{-1}.$$

- 2.
- $\vec{r}(t) = \int \vec{v} dt = \left(\frac{t^3}{3} + 4t + k_1 \right)\vec{i} + (3t - 3t^2 + k_2)\vec{j}$
- . Comme
- $\vec{r}(0) = \vec{0}$
- ,
- $k_1 = k_2 = 0$
- :

$$\vec{r}(t) = \left(\frac{t^3}{3} + 4t \right)\vec{i} + (3t - 3t^2)\vec{j} \text{ m}.$$

3. à
- $t = 3$
- :
- $\vec{r}(3) = (9 + 12)\vec{i} + (9 - 27)\vec{j} = 21\vec{i} - 18\vec{j}$
- . Distance à 0 :

$$|\vec{r}(3)| = \sqrt{21^2 + 18^2} = \sqrt{441 + 324} = \sqrt{765} = 3\sqrt{85} \approx 27.7 \text{ m}.$$

SOLUTION Exercice 7 – Projectile standard sur sol horizontal

Composantes initiales : $u_x = 25 \cos 40^\circ \approx 25(0.766) = 19.15 \text{ m s}^{-1}$; $u_y = 25 \sin 40^\circ \approx 25(0.643) = 16.075 \text{ m s}^{-1}$.

1. Temps de vol : $T = \frac{2u \sin \theta}{g} = \frac{2 \times 16.075}{10} = 3.215 \text{ s} \approx 3.22 \text{ s (3 ch. sign.)}$.
2. Hauteur maximale : $H = \frac{u^2 \sin^2 \theta}{2g} = \frac{(16.075)^2}{20} = \frac{258.4}{20} \approx 12.9 \text{ m (3 ch. sign.)}$.
3. Portée : $R = \frac{u^2 \sin 2\theta}{g} = \frac{625 \times 2 \sin 40^\circ \cos 40^\circ}{10} = \frac{625 \times 2(0.643)(0.766)}{10} \approx \frac{625 \times 0.985}{10} \approx 61.6 \text{ m (3 ch. sign.)}$.

SOLUTION Exercice 8 – Lancer horizontal depuis une falaise

On place l'origine au pied de la falaise : x horizontal, y vertical. Vitesse initiale : $u_x = 12$, $u_y = 0$; hauteur initiale $y_0 = 45$.

1. Mouvement vertical : $y = 45 - \frac{1}{2}gt^2 = 45 - 5t^2$. Impact quand $y = 0$: $t^2 = 9$, donc $T = 3 \text{ s}$.
2. Distance horizontale : $x = u_x T = 12 \times 3 = 36 \text{ m}$ du pied de la falaise.
3. à l'impact : $v_x = 12$ (constant), $v_y = -gT = -30 \text{ m s}^{-1}$.

$$|\vec{v}| = \sqrt{12^2 + 30^2} = \sqrt{144 + 900} = \sqrt{1044} \approx 32.3 \text{ m s}^{-1}.$$

Direction sous l'horizontale : $\tan \alpha = \frac{30}{12} = 2.5$, donc $\alpha \approx 68.2^\circ$ sous l'horizontale.

SOLUTION Exercice 9 – Franchir le mur et marquer (style GCE)

1. Avec $u = 16$, $u_x = 16 \cos \theta$, $u_y = 16 \sin \theta$:

$$x = 16t \cos \theta, \quad y = 16t \sin \theta - 5t^2.$$

On élimine t : $t = \frac{x}{16 \cos \theta}$, d'où

$$y = 16 \sin \theta \cdot \frac{x}{16 \cos \theta} - 5 \left(\frac{x}{16 \cos \theta} \right)^2 = x \tan \theta - \frac{5x^2}{256 \cos^2 \theta} = x \tan \theta - \frac{5x^2}{256} (1 + \tan^2 \theta),$$

en utilisant $\frac{1}{\cos^2 \theta} = \sec^2 \theta = 1 + \tan^2 \theta$. ■

Barème : M1 composantes du déplacement ; M1 élimination de t ; M1 utilisation de $\sec^2 \theta = 1 + \tan^2 \theta$; A1 résultat.

2. Le ballon franchit le mur si $y > 2$ quand $x = 9$:

$$9 \tan \theta - \frac{5 \times 81}{256} (1 + \tan^2 \theta) > 2.$$

Posons $m = \tan \theta$: $9m - \frac{405}{256}(1 + m^2) > 2$, soit

$$405m^2 - 2304m + 917 < 0 \quad (\text{après multiplication par 256 et réarrangement}).$$

Donc $\tan \theta$ doit rendre le trinôme $405m^2 - 2304m + 917$ négatif, c'est-à-dire que m doit se trouver strictement entre ses deux racines. ■

Barème : M1 substitution $x = 9$, $y > 2$; M1 formation de l'inéquation du second degré ; A1 inéquation correcte.

3. Au but, $x = 18$; le ballon entre dans le but si $0 < y < 2.4$. Cas limite pour la barre transversale ($y = 2.4$) :

$$18m - \frac{5 \times 324}{256}(1 + m^2) = 2.4 \implies 18m - \frac{405}{64}(1 + m^2) = 2.4.$$

Multipliant par 64 : $1152m - 405 - 405m^2 = 153.6$, soit

$$405m^2 - 1152m + 558.6 = 0.$$

Discriminant : $\Delta_c = 1152^2 - 4(405)(558.6) = 1327104 - 904884 = 422220 > 0$. Racines

: $m = \frac{1152 \pm \sqrt{422220}}{810} \approx \frac{1152 \pm 649.78}{810}$, donnant $m_1 \approx 0.620$, $m_2 \approx 2.224$. Le trinôme étant convexe (coefficient de m^2 positif), $y < 2.4$ pour $m < 0.620$ ou $m > 2.224$.

Condition pour être au-dessus du sol ($y > 0$ à $x = 18$) :

$$18m - \frac{405}{64}(1 + m^2) > 0 \implies 405m^2 - 1152m + 405 < 0.$$

Discriminant : $\Delta_g = 1152^2 - 4(405)^2 = 1327104 - 656100 = 671004 > 0$. Racines : $m =$

$\frac{1152 \pm \sqrt{671004}}{810} \approx \frac{1152 \pm 819.15}{810}$, donnant $m_3 \approx 0.411$, $m_4 \approx 2.433$. Donc $y > 0$ pour $0.411 < m < 2.433$.

Condition du mur (déjà trouvée) : $0.4306 < m < 5.258$ (racines de $405m^2 - 2304m + 917 = 0$: $m \approx 0.4306$ et $m \approx 5.258$).

Intersection des trois conditions :

- Branche basse : $\max(0.4306, 0.411) < m < \min(0.620, 2.433) \implies 0.4306 < m < 0.620$.
- Branche haute : $\max(2.224, 0.4306) < m < \min(2.433, 5.258) \implies 2.224 < m < 2.433$.

En degrés : $\theta \in (23.3^\circ, 31.8^\circ) \cup (65.8^\circ, 67.6^\circ)$. Pour ces angles, le ballon franchit le mur, passe sous la barre et au-dessus du sol à $x = 18$: il marque.

Barème : M1 substitution $x = 18$; A1 trinômes en $\tan \theta$; M1 analyse des discriminants et des signes ; M1 résolution de l'inéquation du mur ; A1 racines ; A1 intervalles finaux pour θ . Attente de l'examinateur : manipulation algébrique rigoureuse et énoncé clair de la gamme d'angles.

SOLUTION Exercice 10 – Projectile depuis un toit (style GCE)

1. Origine au pied de l'immeuble, y vers le haut. Point de lancement $(0, 20)$, $u_x = 15 \cos 30^\circ = \frac{15\sqrt{3}}{2} \approx 12.99$, $u_y = 15 \sin 30^\circ = 7.5$:

$$x = 7.5\sqrt{3}t \approx 12.99t, \quad y = 20 + 7.5t - 5t^2.$$

Barème : M1 composantes ; A1 $x(t)$; A1 $y(t)$ avec le +20.

2. Atterrissage quand $y = 0$: $20 + 7.5T - 5T^2 = 0$, soit $5T^2 - 7.5T - 20 = 0$. ■

$$T = \frac{7.5 \pm \sqrt{7.5^2 + 400}}{10} = \frac{7.5 \pm \sqrt{456.25}}{10} = \frac{7.5 \pm 21.36}{10}.$$

On prend la racine positive : $T \approx 2.89$ s (la racine négative -1.39 s est rejetée comme non physique). Barème : M1 équation $y = 0$; A1 équation donnée ; M1 formule quadratique ; A1 racine positive choisie avec justification.

3. Distance horizontale : $x = 7.5\sqrt{3} \times 2.886 \approx 12.99 \times 2.886 \approx 37.5$ m. Barème : M1 $x = u_x T$; A1 valeur.

4. à l'atterrissage : $v_x = 7.5\sqrt{3} \approx 12.99 \text{ m s}^{-1}$; $v_y = 7.5 - 10T = 7.5 - 28.86 = -21.36 \text{ m s}^{-1}$.

$$|\vec{v}| = \sqrt{12.99^2 + 21.36^2} = \sqrt{168.7 + 456.3} = \sqrt{625} = 25 \text{ m s}^{-1}.$$

Angle sous l'horizontale : $\tan \alpha = \frac{21.36}{12.99} = 1.644$, donc $\alpha \approx 58.7^\circ$ sous l'horizontale.
Barème : M1 composantes à T ; A1 vitesse ; A1 angle avec direction précisée.

SOLUTION Exercice 11 – Mouvement vertical et ses graphiques (style GCE)

1. Chute sans vitesse initiale : $h = \frac{1}{2}gt^2 = \frac{1}{2}(10)(1.5)^2 = 11.25$ m. Vitesse à l'impact : $v = gt = 10 \times 1.5 = 15 \text{ m s}^{-1}$. Barème : M1 formule correcte ; A1 hauteur ; A1 vitesse.

2. On prend le sens vers le haut positif, origine au bord du puits. Déplacement au fond : $s = -25$ m. En utilisant $s = ut - \frac{1}{2}gt^2$:

$$-25 = 20t - 5t^2 \implies 5t^2 - 20t - 25 = 0 \implies t^2 - 4t - 5 = 0 \implies (t - 5)(t + 1) = 0.$$

Donc $t = 5$ s (on rejette $t = -1$). Vitesse au fond : $v = u - gt = 20 - 50 = -30 \text{ m s}^{-1}$, soit 30 m s^{-1} vers le bas. (Vérification avec $v^2 = u^2 + 2g(25) = 400 + 500 = 900$.) Barème : M1 convention de signe et $s = -25$; A1 équation du second degré ; A1 $t = 5$ s avec rejet de la racine négative ; A1 vitesse.

3. L'objet a la vitesse ascendante du ballon 8 m s^{-1} à l'instant du lâcher : il n'est pas lâché

sans vitesse initiale. Il continue donc vers le haut, en décélérant de g , jusqu'à ce que sa vitesse devienne nulle, et seulement alors il retombe. Avec l'origine au point de lâcher, $u = +8$, $s = -100$:

$$-100 = 8t - 5t^2 \Rightarrow 5t^2 - 8t - 100 = 0 \Rightarrow t = \frac{8 \pm \sqrt{64 + 2000}}{10} = \frac{8 \pm \sqrt{2064}}{10} \approx \frac{8 \pm 45.43}{10}.$$

Racine positive : $t \approx 5.34$ s. Vitesse à l'impact : $v = 8 - 10(5.34) = -45.4 \text{ m s}^{-1}$, soit environ 45.4 m s^{-1} vers le bas. Barème : B1 explication du mouvement initial vers le haut (inertie / vitesse initiale du ballon) ; M1 équation avec $u = 8$, $s = -100$; A1 temps ; A1 vitesse.


4. Avec $u = 14$: temps au sommet $t_1 = \frac{u}{g} = 1.4$ s ; temps total $T = 2.8$ s ; hauteur maximale

$$h = \frac{u^2}{2g} = \frac{196}{20} = 9.8 \text{ m}.$$

Déplacement-temps : parabole $s = 14t - 5t^2$ de $(0, 0)$, sommet $(1.4, 9.8)$, retour à $(2.8, 0)$.

Vitesse-temps : droite $v = 14 - 10t$ de $+14$ à $t = 0$, passant par 0 à $t = 1.4$, jusqu'à -14 m s^{-1} à $t = 2.8$.


Accélération-temps : ligne horizontale $a = -10 \text{ m s}^{-2}$ pendant tout le vol.



Graphiques déplacement-temps, vitesse-temps et accélération-temps pour la balle lancée vers le haut à 14 m s^{-1} .

Barème : B1 parabole avec sommet et zéros corrects ; B1 droite vitesse traversant zéro à $t = 1.4$ avec valeurs ± 14 ; B1 ligne accélération constante à -10 . Attente de l'examineur : le graphique de la vitesse doit devenir négatif (la balle redescend) et le graphique de l'accélération doit montrer que a ne s'annule jamais.

13 SUMMARY SHEET



Mind map of the chapter: from vector kinematics to projectile motion and free fall.

FICHE BILAN L'essentiel à connaître pour l'examen

Key definitions

- **Position vector:** $\vec{r}(t) = x(t)\vec{i} + y(t)\vec{j}$ locates the particle at time t .
- **Velocity:** $\vec{v} = \frac{d\vec{r}}{dt} = \dot{x}\vec{i} + \dot{y}\vec{j}$; it is tangent to the path. Speed = $|\vec{v}| = \sqrt{\dot{x}^2 + \dot{y}^2}$.
- **Acceleration:** $\vec{a} = \frac{d\vec{v}}{dt} = \ddot{x}\vec{i} + \ddot{y}\vec{j}$.
- **Projectile:** a body launched with initial velocity $\vec{u}$ and then moving under gravity alone

(air resistance neglected).

- **Free fall:** motion with $\vec{a} = -g\vec{j}$, $g \approx 9.81 \text{ m s}^{-2}$ (take $g = 9.81$ or 10 as instructed).

Formulas and laws to know

- Differentiation: $\vec{r} \xrightarrow{d/dt} \vec{v} \xrightarrow{d/dt} \vec{a}$. Integration reverses the chain; each integration introduces a constant fixed by initial conditions.
- Projectile with launch speed u at angle θ from the origin:

$$x = ut \cos \theta, \quad y = ut \sin \theta - \frac{1}{2}gt^2.$$

$$v_x = u \cos \theta \text{ (constant)}, \quad v_y = u \sin \theta - gt.$$

- Cartesian equation of the **trajectory** (a parabola):

$$y = x \tan \theta - \frac{gx^2}{2u^2 \cos^2 \theta} = x \tan \theta - \frac{gx^2}{2u^2}(1 + \tan^2 \theta).$$

- On horizontal ground:

$$\text{Time of flight } T = \frac{2u \sin \theta}{g}, \quad \text{Range } R = \frac{u^2 \sin 2\theta}{g}, \quad \text{Max height } H = \frac{u^2 \sin^2 \theta}{2g}.$$

- Maximum range: $R_{\max} = \frac{u^2}{g}$ at $\theta = 45^\circ$; complementary angles θ and $90^\circ - \theta$ give the same range.
- At the highest point: $v_y = 0$ but $v_x = u \cos \theta \neq 0$; time to reach it is $T/2$.
- Vertical throw upwards (taking up as positive): $v = u - gt$, $y = ut - \frac{1}{2}gt^2$, $v^2 = u^2 - 2gy$; greatest height $\frac{u^2}{2g}$; the body returns with speed u (symmetry of free fall).
- Graphs: gradient of $x-t$ graph = velocity; gradient of $v-t$ graph = acceleration; area under $v-t$ graph = displacement. For vertical motion the $v-t$ graph is a straight line of gradient $-g$.

Methods

- Vector kinematics: differentiate or integrate each component separately; use the given position/velocity at $t = 0$ to find constants.
- Projectile problems: (1) resolve $\vec{u}$ into $u \cos \theta$ and $u \sin \theta$; (2) write $x(t)$ and $y(t)$; (3) translate the question into a condition ($y = 0$: landing; $v_y = 0$: summit; $y = h$: target height); (4) solve for t , then substitute.
- Finding θ or u from the trajectory: substitute the coordinates of a known point into the Cartesian equation; use $1/\cos^2 \theta = 1 + \tan^2 \theta$ to get a quadratic in $\tan \theta$.
- Vertical throws: choose a positive direction once and keep it; $a = -g$ throughout, even at the top.

Common mistakes

- Mixing degrees and radians, or using $\sin \theta$ where $\cos \theta$ is needed when resolving $\vec{u}$.
- Believing velocity is zero at the highest point: only the vertical component vanishes.
- Using the range formula when launch and landing are at different heights — go back to $y(t) = 0$ instead.
- Forgetting the constant of integration or the units (m , m s^{-1} , m s^{-2}).
- Sign errors in vertical motion: acceleration is $-g$ on the way up and on the way down.
- Confusing distance travelled with displacement on a $v-t$ graph (areas below the axis count negatively for displacement).

GCE tips

- State your model clearly: “air resistance neglected, $g = 9.81 \text{ m s}^{-2}$ ” earns marks.
- Show the resolved components $u \cos \theta$, $u \sin \theta$ before writing equations — method marks are awarded there.
- Keep exact values ($\sin 30^\circ = \frac{1}{2}$, etc.) until the final line, then round to 3 significant figures.
- For “show that” questions, derive from $x(t)$, $y(t)$ — never quote the result you are asked to prove.
- Sketch the parabola and mark the given data; it prevents sign and angle errors and helps the examiner follow your work.



Signed • OnBuch+

ONBUCH+ TEACHING TEAM

FOLLOWS THE OFFICIAL MINESEC SYLLABUS

OnBuch+