

BIOLOGY • UPPER SIXTH

UPPER SIXTH — BIOLOGY

CHAPTER 6

NON-INFECTIOUS DISEASES

SUGGESTED TIME: 4 PERIODS

IN THIS CHAPTER

This chapter explores the major non-infectious diseases affecting the human reproductive, nervous, endocrine, renal, cardiovascular and musculoskeletal systems. For each system, learners link normal structure and function to the causes, symptoms, prevention and management of its characteristic disorders, with emphasis on conditions of public-health importance in Cameroon.

- By the end of this chapter I can define non-infectious disease and distinguish its main categories (degenerative, hormonal, genetic, nutritional, neoplastic, autoimmune).
- By the end of this chapter I can describe the causes, symptoms and prevention of common disorders of the male and female reproductive systems.
- By the end of this chapter I can explain how disorders of the nervous system (e.g. epilepsy, stroke, meningitis complications, Parkinson's disease) arise from failures of neurones, synapses or brain regions.
- By the end of this chapter I can relate endocrine diseases (diabetes mellitus, goitre, gigantism, cretinism) to hypo- or hypersecretion of specific hormones.
- By the end of this chapter I can describe kidney diseases (nephritis, kidney stones, renal failure) and the principle of dialysis.
- By the end of this chapter I can explain cardiovascular diseases (hypertension, atherosclerosis, coronary heart disease, stroke) in terms of risk factors and vascular changes.



DOWNLOAD ON- BUCH

Free • Google Play
AI tutor, past papers,
results



GO ONBUCH+

All signed courses, unlimited AI
tutor, PDF sheets — OnBuch+
tab in the app

THIS SIGNED COURSE CONTAINS

COURSE

complete, illustrated,
step by step

METHODS

and worked examples

EXERCISES

exam style + solutions

SUMMARY SHEET

the essentials to
remember



Signed • OnBuch+

ONBUCH+ TEACHING TEAM
FOLLOWS THE OFFICIAL MINESEC SYLLABUS

OnBuch+

CONTENTS

- | | |
|--|--|
| 1. Section 1: Non-infectious diseases — overview and disorders of the reproductive systems | musculoskeletal system and integrated prevention of NCDs |
| 2. Section 2: Disorders of the nervous and endocrine systems | 5. Integration activity |
| 3. Section 3: Diseases of the renal and cardiovascular systems | 6. Methods and worked examples |
| 4. Section 4: Disorders of the muscu- | 7. Exercises |
| | 8. Solutions to the exercises |
| | 9. Summary sheet |

CHAPTER OBJECTIVES

- By the end of this chapter I can define non-infectious disease and distinguish its main categories (degenerative, hormonal, genetic, nutritional, neoplastic, autoimmune).
- By the end of this chapter I can describe the causes, symptoms and prevention of common disorders of the male and female reproductive systems.
- By the end of this chapter I can explain how disorders of the nervous system (e.g. epilepsy, stroke, meningitis complications, Parkinson's disease) arise from failures of neurones, synapses or brain regions.
- By the end of this chapter I can relate endocrine diseases (diabetes mellitus, goitre, gigantism, cretinism) to hypo- or hyper-secretion of specific hormones.
- By the end of this chapter I can describe kidney diseases (nephritis, kidney stones, renal failure) and the principle of dialysis.
- By the end of this chapter I can explain cardiovascular diseases (hypertension, atherosclerosis, coronary heart disease, stroke) in terms of risk factors and vascular changes.
- By the end of this chapter I can describe musculoskeletal disorders (arthritis, osteoporosis, rickets, muscular dystrophy) and their causes.
- By the end of this chapter I can interpret clinical data (blood pressure readings, blood/urine test results, X-ray descriptions) to suggest a likely diagnosis.
- By the end of this chapter I can propose lifestyle and public-health measures that reduce the burden of non-infectious diseases in a Cameroonian community.

PREREQUISITES

- Structure and functions of the male and female reproductive systems (Lower Sixth)
- Structure of the neurone, reflex arc and organisation of the nervous system
- Endocrine glands, hormones and negative feedback control (e.g. insulin and blood glucose)
- Structure of the kidney nephron, ultrafiltration and selective reabsorption
- Structure of the heart, blood vessels and the double circulation; bones, joints and muscles

1 SECTION 1: NON-INFECTIOUS DISEASES — OVERVIEW AND DISORDERS OF THE REPRODUCTIVE SYSTEMS

At the Regional Hospital of Bamenda, a 45-year-old trader is told her persistent fatigue and swollen ankles come from failing kidneys, while her neighbour's infertility after years of marriage is traced to an untreated pelvic infection. Neither illness was caused by a germ that could be caught from someone else — both are **non-infectious diseases**, disorders arising from within the body's own systems. In Cameroon today, such conditions (hypertension, diabetes, kidney failure, arthritis) are silently overtaking malaria and cholera as causes of death and disability. Why do organs that worked perfectly for decades suddenly malfunction, and how can understanding the biology of each system help us prevent, detect and manage these diseases? This section equips you to answer exactly that.

1.1 What are non-infectious (non-communicable) diseases?


DEFINITION Non-infectious disease

A **non-infectious disease** (also called a **non-communicable disease, NCD**) is a medical condition that is not caused by a transmissible pathogen (bacterium, virus, fungus, parasite). It arises from genetic, physiological, environmental or lifestyle factors and cannot be spread directly from person to person.

DEFINITION Neoplasm

A **neoplasm** is an abnormal mass of tissue whose growth exceeds and is uncoordinated with that of the normal tissues. A **benign tumour** remains localised, does not invade surrounding tissue and does not metastasise. A **malignant tumour** (cancer) invades locally and can spread to distant sites (metastasis).

NCDs are commonly grouped into seven major categories. The flowchart below summarises each category with a typical example.



Classification of non-infectious diseases with one representative example per category.


1.2 General risk factors and the growing burden in Cameroon

The main modifiable and non-modifiable risk factors for NCDs are:

- **Age** — physiological reserve declines with time.
- **Heredity** — family history of hypertension, diabetes, certain cancers.
- **Diet** — high salt, saturated fats, refined sugars; low fruit/vegetable intake.
- **Tobacco** — smoking damages vasculature and is carcinogenic.
- **Alcohol** — excessive intake raises blood pressure and cancer risk.
- **Physical inactivity** — contributes to obesity, insulin resistance.
- **Environmental pollutants** — indoor smoke from biomass fuels, outdoor air pollution.
- **Urbanisation** — rapid lifestyle changes, reduced physical activity, processed diets.

Cameroon, like many sub-Saharan countries, faces a **double burden of disease**: infectious diseases remain prevalent while NCDs are rising sharply. The illustrative bar chart below shows

the estimated proportion of total deaths attributed to NCDs versus infectious diseases over three recent decades (data are indicative, not official statistics).




Illustrative trend: rising share of deaths due to NCDs in Cameroon (1990–2020).

KEY POINTS**Key points on NCD burden**

- NCDs now account for more than half of all deaths in many urban Cameroonian settings.
- Early detection and lifestyle modification can prevent up to 80% of premature NCD deaths (WHO estimate).
- Health systems must integrate NCD screening into primary care (e.g. blood pressure, blood glucose, cervical screening).

1.3 Male reproductive system disorders



1.4 Prostate disorders

PROPERTY Benign prostatic hyperplasia (BPH)

Non-cancerous enlargement of the prostate gland, common after age 50. It compresses the urethra causing **lower urinary tract symptoms (LUTS)**: hesitancy, weak stream, frequency, nocturia, incomplete emptying.

PROPERTY Prostate cancer

Malignant tumour of the prostate epithelium. Risk factors: age > 50, family history, African ancestry, high-fat diet. Early stages often asymptomatic; later may mimic BPH plus haematuria, bone pain (metastasis). Screening: serum **prostate-specific antigen (PSA)** test and digital rectal examination (DRE). Definitive diagnosis by trans-rectal ultrasound-guided biopsy.

METHOD PSA screening interpretation (simplified for examination)

1. Obtain serum PSA (ng/mL).
2. Age-adjusted reference (simplified): < 4.0 ng/mL generally normal; 4.0–10.0 ng/mL borderline; > 10.0 ng/mL high suspicion. Note: modern guidelines use age-specific ranges (e.g. 40–49 yr: 0–2.5; 50–59 yr: 0–3.5; 60–69 yr: 0–4.5; 70–79 yr: 0–6.5 ng/mL).
3. If PSA elevated, repeat after 4–6 weeks (rule out prostatitis).
4. Persistent elevation → refer for DRE and possible biopsy.

1.5 Testicular disorders

- **Testicular cancer** — most common solid tumour in males 15–35 years. Presents as painless lump; high cure rate if detected early (orchietomy + chemotherapy).
- **Varicocele** — dilation of pampiniform plexus veins; raises scrotal temperature, impairs spermatogenesis → infertility.
- **Hydrocele** — fluid accumulation in tunica vaginalis; usually benign, may cause discomfort.
- **Cryptorchidism** — failure of testis to descend into scrotum; if untreated, increases risk of malignancy and infertility.

1.6 Male infertility and erectile dysfunction

PROPERTY Causes of male infertility

- Pre-testicular: hormonal (hypogonadotropic hypogonadism, hyperprolactinaemia).
- Testicular: varicocele, cryptorchidism, infection (mumps orchitis), genetic (Klinefelter syndrome), idiopathic.
- Post-testicular: obstruction of vas deferens (congenital absence, post-vasectomy, infection).

PROPERTY Erectile dysfunction (ED)

Inability to achieve/maintain erection sufficient for intercourse. Vascular (atherosclerosis, hypertension), neurological (diabetic neuropathy), psychological (stress, depression), hormonal (low testosterone), iatrogenic (antihypertensives, antidepressants).

WORKED EXAMPLE 1 Worked example: interpreting a semen analysis (WHO 2021, 5th edition)

Statement: A 32-year-old man presents with 2 years of infertility. Semen analysis shows: volume 1.8 mL, concentration 8 million/mL, progressive motility 25%, normal morphology 3%. WHO 2021 lower reference limits (5th centiles): volume ≥ 1.5 mL, concentration ≥ 15 million/mL, progressive motility $\geq 30\%$, normal morphology $\geq 4\%$. Classify the abnormalities.


SOLUTION

Solution:

- Volume 1.8 mL ≥ 1.5 mL $\rightarrow$ normal.
- Concentration 8 million/mL < 15 million/mL $\rightarrow$ **oligozoospermia**.
- Progressive motility 25% $< 30\%$ $\rightarrow$ **asthenozoospermia**.
- Normal morphology 3% $< 4\%$ $\rightarrow$ **teratozoospermia**.

Combined pattern = **oligoasthenoteratozoospermia (OAT)**. Further hormonal profile (FSH, LH, testosterone) and scrotal ultrasound are indicated to identify cause.

1.7 Female reproductive system disorders



Female reproductive system with common disorder sites highlighted.

1.8 Menstrual disorders (hormonal basis)

- **Amenorrhoea** — absence of menses. Primary: no menarche by age 16; Secondary: cessation for ≥ 3 cycles. Causes: hypothalamic-pituitary failure (low GnRH/FSH/LH), ovarian failure (premature ovarian insufficiency), outflow tract obstruction, pregnancy.
- **Dysmenorrhoea** — painful menstruation. Primary: prostaglandin-mediated uterine contractions (no pathology). Secondary: endometriosis, adenomyosis, fibroids.
- **Menorrhagia** — excessively heavy/prolonged bleeding (> 80 mL/cycle). Causes: anovulatory cycles (unopposed oestrogen), fibroids, coagulopathies, copper IUD.

PROPERTY Hormonal control of the menstrual cycle

- Follicular phase: rising **oestradiol** from developing follicle $\rightarrow$ proliferative endometrium.
- Ovulation: LH surge triggered by positive feedback of oestradiol.
- Luteal phase: **progesterone** from corpus luteum $\rightarrow$ secretory endometrium.
- Menses: withdrawal of progesterone+oestradiol $\rightarrow$ spiral artery vasoconstriction $\rightarrow$ endometrial shedding.

1.9 Ovarian and uterine disorders

- **Ovarian cysts** — fluid-filled sacs; functional (follicular, corpus luteum) usually resolve; pathological (dermoid, cystadenoma) may require surgery.
- **Polycystic ovary syndrome (PCOS)** — hyperandrogenism, oligo-/anovulation, polycystic ovaries on ultrasound (Rotterdam criteria: 2 of 3). Insulin resistance central; long-term risks: type 2 diabetes, endometrial hyperplasia, cardiovascular disease.
- **Uterine fibroids (leiomyomas)** — benign smooth-muscle tumours; oestrogen-dependent; cause menorrhagia, pressure symptoms, infertility.
- **Adenomyosis** — endometrial glands within myometrium; causes dysmenorrhoea, menorrhagia, enlarged uterus.
- **Endometriosis** — ectopic endometrial tissue (ovaries, peritoneum); causes chronic pelvic pain, dysmenorrhoea, infertility via adhesions and inflammatory milieu.

1.10 Gynaecological cancers

Cancer	Main risk factors	Early signs	Screening / early detection
Cervix	HPV (types 16,18), early sexual debut, multiple partners, smoking, immunosuppression	Post-coital bleeding, intermenstrual bleeding, foul discharge	Pap smear (cytology) every 3yr (25–65yr); HPV DNA testing; visual inspection with acetic acid (VIA) in low-resource settings
Ovary	Nulliparity, late menopause, BRCA1/2 mutations, family history	Vague: bloating, pelvic discomfort, early satiety, urinary frequency	No population screening; high-risk: transvaginal ultrasound + CA-125 annually
Breast	Female sex, age, family history, BRCA1/2, early menarche/late menopause, nulliparity, hormone therapy, alcohol, obesity	Painless lump, skin dimpling, nipple retraction, discharge	Breast self-examination (BSE) monthly; clinical breast exam annually; mammography 40–50yr (or earlier if high risk)

1.11 Pelvic inflammatory disease (PID) and ectopic pregnancy

PROPERTY Pelvic inflammatory disease

Ascending infection (often *Chlamydia trachomatis*, *Neisseria gonorrhoeae*) from cervix to uterus, tubes, peritoneum. Acute PID: lower abdominal pain, fever, cervical motion tenderness. Sequelae: tubal scarring → **blocked Fallopian tubes** → infertility; increased risk of **ectopic pregnancy**.

WARNING Ectopic pregnancy — medical emergency

Implantation outside uterine cavity (95% in Fallopian tube). Presents with amenorrhoea, unilateral pelvic pain, vaginal bleeding, shoulder tip pain (diaphragmatic irritation). Rupture → haemorrhagic shock. Diagnosis: β -hCG + transvaginal ultrasound (empty uterus, adnexal mass). Management: methotrexate (unruptured, stable) or laparoscopic salpingectomy/salpingostomy.

1.12 Prevention and management of reproductive NCDs

METHOD Breast self-examination (BSE) — 5 steps

1. Visual inspection — stand before mirror, arms at sides, then raised; look for asymmetry, skin changes, nipple inversion.
2. Palpation lying down — place pillow under shoulder, use flat fingers in circular motion covering whole breast and axilla.
3. Palpation standing — in shower with soapy fingers, same systematic pattern.
4. Nipple check — gently squeeze each nipple; note any discharge (bloody, clear).
5. Record and report — note date, findings; any new lump or change → consult health worker within 2 weeks.

METHOD Rationale for cervical screening (Pap smear)

1. HPV infection → cervical intraepithelial neoplasia (CIN) grades 1–3 over 10–20 years.
2. Pap smear detects abnormal squamous cells (ASC-US, LSIL, HSIL) before invasion.
3. Treatment of CIN (cryotherapy, LEEP) prevents progression to invasive carcinoma.
4. Screening interval 3 years balances benefit/harm; HPV vaccination (9–14 yr) reduces future burden.

WARNING Common mistake: confusing STIs with non-infectious reproductive disorders

- STIs (e.g. gonorrhoea, syphilis, HIV) are infectious — caused by transmissible pathogens.
- Fibroids, BPH, PCOS, endometriosis, prostate cancer are non-infectious — arise from hormonal, genetic, lifestyle factors.
- In exam tables, always separate cause (pathogen vs hormonal/genetic), symptom, and prevention (condom/vaccine vs lifestyle/screening).

PROPERTY Hormonal imbalance and reproductive disorders

- Oestrogen excess / progesterone deficiency → endometrial hyperplasia, menorrhagia

gia, fibroid growth, increased breast/endometrial cancer risk.

- Androgen excess (e.g. PCOS) → anovulation, hirsutism, acne, insulin resistance.
- Testosterone deficiency (male hypogonadism) → low libido, ED, reduced spermatogenesis, osteoporosis.
- Prolactin excess (prolactinoma) → amenorrhoea/galactorrhoea (female), hypogonadism (male).

DID YOU KNOW? HPV vaccination in Cameroon

Cameroon introduced the quadrivalent HPV vaccine into the routine immunisation programme for girls aged 9–13 years in 2020 (GAVI support). Coverage remains below 50% in many regions due to vaccine hesitancy, cold-chain challenges and school-based delivery gaps. Catch-up campaigns and community sensitisation are ongoing. Vaccination of boys is under consideration for future expansion.

KEY POINTS Exam tip: three-column cause–symptom–prevention table

GCE questions frequently ask: “Complete the table below for the following conditions: cervical cancer, BPH, PCOS.” Practise drawing a table with columns Condition, Cause / Risk factors, Key symptom(s), Prevention / Early detection. Fill each row concisely; use bullet points inside cells if needed. This structure earns full marks for organisation and completeness.

EXERCISE 1 Practice: cause–symptom–prevention table

Complete the table for the four conditions listed. Write your answer in the three-column format described above.

Condition	Cause / Risk factors	Key symptom(s)	Prevention / Early detection
Cervical cancer			
Benign prostatic hyperplasia			
Polycystic ovary syndrome			
Testicular cancer			

SOLUTION Exercise 1

Condition	Cause / Risk factors	Key symptom(s)	Prevention / Early detection
Cervical cancer	HPV 16/18 infection; early sexual debut; multiple partners; smoking; immunosuppression	Post-coital bleeding; intermenstrual bleeding; foul discharge	HPV vaccination (9–14yr); Pap smear q3yr (25–65yr); VIA in low-resource settings
Benign prostatic hyperplasia	Age > 50; hormonal changes (DHT); family history	LUTS: hesitancy, weak stream, frequency, nocturia, incomplete emptying	No proven primary prevention; early DRE + PSA for early detection; lifestyle (weight control, exercise) may reduce severity
Polycystic ovary syndrome	Insulin resistance; hyperandrogenism; genetic predisposition; obesity	Oligo/amenorrhoea; hirsutism; acne; polycystic ovaries on ultrasound	Weight loss (5–10%); combined oral contraceptives for cycle control; metformin if insulin resistant; regular endometrial surveillance
Testicular cancer	Cryptorchidism; family history; age 15–35; Caucasian ethnicity	Painless testicular lump; heaviness; sometimes gynecomastia (hCG)	Monthly testicular self-exam (15–35yr); early orchiectomy if mass detected; no population screening


2 SECTION 2: DISORDERS OF THE NERVOUS AND ENDOCRINE SYSTEMS

In Section 1 we saw that non-infectious diseases (NCDs) are not caused by pathogens but by malfunctions of the body's own structures, and we applied this idea to the reproductive systems. We now turn to two systems that act as the body's **control networks**: the **nervous system**, which sends rapid electrical messages, and the **endocrine system**, which sends slower chemical messages (hormones) through the blood. Because these two systems coordinate every other organ, their disorders often have widespread effects — and several of them, such as stroke and type 2 diabetes, are among the fastest-growing health problems in Cameroonian towns today.

2.1 Locating nervous-system disorders: a quick review

The nervous system is divided into the **central nervous system** (CNS: brain and spinal cord) and the **peripheral nervous system** (PNS: cranial and spinal nerves). The functional cell is the **neuron**: impulses travel along the axon (often insulated by a **myelin sheath**) and cross junctions called **synapses** using chemical **neurotransmitters**. This organisation gives us a logical way to classify disorders:

- **Electrical disorders** — abnormal firing of neurones (epilepsy).
- **Vascular disorders** — failure of blood supply to the brain (stroke).
- **Degenerative disorders** — progressive death of neurones (Parkinson's, Alzheimer's).
- **Autoimmune/structural disorders** — damage to myelin (multiple sclerosis), or irritation of pain pathways (neuralgia, migraine).



A neurone with the main sites at which nervous-system disorders arise.

2.2 Epilepsy

DEFINITION Epilepsy

Epilepsy is a chronic non-infectious disorder of the brain in which groups of neurones suddenly produce **abnormal, excessive electrical discharges**, causing recurrent

seizures (fits).

Seizures vary in form. In **generalised tonic-clonic seizures** (grand mal), the person loses consciousness, the body stiffens (tonic phase) then jerks rhythmically (clonic phase). In **absence seizures** (petit mal), common in children, the person briefly stares blankly for a few seconds without falling. Triggers include fever, flashing lights, lack of sleep, stress and missed medication. Epilepsy is controlled — though usually not cured — by regular **anticonvulsant** medicines; interrupting treatment is a common cause of relapse.

METHOD First aid for an epileptic seizure

1. Stay calm and note the time; most seizures stop within a few minutes.
2. Protect the person from injury: move away hard or sharp objects, cushion the head.
3. Do not restrain the person's movements and do not put anything (spoon, stick, fingers) into the mouth.
4. Loosen tight clothing around the neck.
5. When the jerking stops, place the person in the **recovery position** (on the side) so saliva drains and breathing is free.
6. Stay until full consciousness returns; call for medical help if the seizure lasts more than about five minutes or repeats.

WARNING Stigma is not science

In some Cameroonian communities epilepsy is wrongly attributed to witchcraft or thought to be contagious through saliva. Epilepsy is neither contagious nor supernatural: it is an electrical disorder of the brain, treatable with anticonvulsant medicines. Forcing objects into a patient's mouth causes broken teeth and choking — a harmful practice based on a myth.

2.3 Stroke (cerebrovascular accident)**DEFINITION** Stroke

A **stroke** is the sudden death of brain tissue caused by an interruption of its blood supply. It is ischaemic when a clot blocks a cerebral artery, and haemorrhagic when a weakened vessel bursts and bleeds into the brain.

The strongest risk factor is **hypertension** (high blood pressure), together with smoking, diabetes and high cholesterol. Because neurones deprived of oxygen die within minutes, a stroke is a medical emergency. Typical warning signs can be recalled with the word **FAST**: Face drooping on one side, **A**rm weakness (one arm cannot be raised), **S**peech slurred or lost, **T**ime to rush to hospital. Sudden severe headache and one-sided numbness are also common. Rapid hospital treatment (clot-dissolving drugs for ischaemic stroke, given early) greatly improves recovery — “time is brain”. Prevention rests on controlling blood pressure, not smoking, exercising and limiting salt and alcohol.

2.4 Degenerative, autoimmune and other nervous disorders

Parkinson’s disease results from the progressive loss of **dopamine-producing neurones** in a region of the midbrain (the substantia nigra). Dopamine is needed for smooth, coordinated movement, so patients show **tremor at rest**, muscle rigidity and slow movement (bradykinesia). Treatment replaces dopamine activity (e.g. L-dopa).

Alzheimer’s disease is a degenerative disorder causing **progressive memory loss**, confusion and personality change, mainly in the elderly; care is mostly supportive.

Multiple sclerosis (MS) is an **autoimmune** disease in which the immune system attacks the **myelin sheath** of CNS neurones. Demyelination slows or blocks impulse conduction, producing weakness, numbness and vision problems that come and go.

Migraine is recurrent severe headache, often one-sided with nausea and light sensitivity; **neuralgia** is sharp pain along a nerve (e.g. sciatica). Finally, remember that **alcohol, drugs and head injuries** damage the nervous system: chronic alcohol destroys neurones and the liver’s detoxification capacity, while head injuries from motorcycle accidents — sadly common in Yaoundé and Douala — cause concussions and lasting brain damage. Wearing a helmet is disease prevention.

2.5 The endocrine system and negative feedback

The **endocrine glands** (pituitary, thyroid, parathyroids, adrenals, pancreas, ovaries, testes) secrete **hormones** directly into the blood. Hormone release is normally controlled by **negative feedback**: when the blood level of a hormone (or of the substance it controls) rises above the set point, further secretion is switched off; when it falls, secretion resumes. This keeps internal conditions stable (homeostasis).

PROPERTY Negative feedback and endocrine disease

In a healthy feedback loop, a rise in the controlled variable inhibits the gland, and a fall stimulates it. If a gland becomes over-active (hyper-secretion), the controlled variable stays abnormally high because the gland no longer responds to the “off” signal; if it becomes under-active (hypo-secretion), the variable stays abnormally low. Hence endocrine diseases are classified as **hypo-secretion** or **hyper-secretion** disorders.

Pituitary	→	Hypo: dwarfism; diabetes insipidus (ADH) Hyper: gigantism (child), acromegaly (adult)
Thyroid	→	Hypo: cretinism (child), myxoedema, goitre Hyper: hyperthyroidism (exophthalmic goitre)
Adrenals	→	Hypo: Addison's disease Hyper: Cushing's syndrome
Pancreas (islets)	→	Hypo (insulin): diabetes mellitus type 1 Resistance: diabetes mellitus type 2
Ovaries / Testes	→	Hypo: infertility, delayed puberty Hyper: precocious puberty

Major endocrine glands with the diseases caused by hypo- and hyper-secretion of each.


2.6 Diabetes mellitus

DEFINITION Diabetes mellitus

Diabetes mellitus is a chronic disorder in which blood **glucose** remains abnormally high (hyperglycaemia) because the hormone **insulin** is absent, insufficient or ineffective.

Type 1 diabetes is an **autoimmune** destruction of the insulin-secreting β -cells of the islets of Langerhans, usually appearing in childhood or adolescence: insulin is deficient, so daily insulin injections are essential for survival. **Type 2 diabetes**, the commonest form in adults, results from **insulin resistance**: the pancreas still secretes insulin, but body cells respond poorly to it. Risk factors include obesity, physical inactivity, age and family history — it is increasingly common in urban Cameroon with changing diets and sedentary lifestyles.

Symptoms of both types follow directly from hyperglycaemia: when blood glucose exceeds the **renal threshold** (about 10 mmol L^{-1}), the kidneys cannot reabsorb it all, so glucose spills into the urine, dragging water with it (**polyuria**), causing thirst (**polydipsia**); because cells cannot use glucose properly, the body breaks down fat and muscle (**weight loss**). Diagnosis is by **blood glucose tests**: fasting blood glucose, or an **oral glucose tolerance test** (OGTT) in which glucose is drunk and blood glucose is followed over two hours.



Blood glucose after a meal: the diabetic curve starts higher, rises further (above the renal threshold, so glucose appears in urine) and returns to normal much more slowly.

WORKED EXAMPLE 2 Interpreting a glucose-tolerance curve

Two patients each drink a glucose solution. Patient A's blood glucose rises from 4.6 to 8.1 mmol/L and returns below 6 mmol/L within 2 hours. Patient B's rises from 7.4 to 12.6 mmol/L and is still 9.0 mmol/L after 2 hours. Which patient is diabetic? Justify.

SOLUTION

Solution. Patient B is diabetic. Justification: (1) her fasting glucose (7.4 mmol/L) is already above the normal range; (2) the peak (12.6 mmol/L) exceeds the renal threshold (about 10 mmol/L), so glucose appears in the urine; (3) glucose remains elevated after 2 hours, showing that insulin action is insufficient or ineffective. Patient A shows a normal response: a moderate peak followed by a rapid return to the set point under insulin action.


Long-term **complications** of poorly controlled diabetes include **retinopathy** (eye damage, blindness), **neuropathy** (nerve damage, foot ulcers that may lead to amputation), and **nephropathy** (kidney damage). Management combines insulin (type 1, and sometimes type 2), a balanced diet low in refined sugars, regular exercise and weight control.

WARNING Mellitus ≠ insipidus

Do not confuse diabetes mellitus (insulin/glucose problem, sugar in urine) with diabetes insipidus (deficiency of **ADH** released from the posterior pituitary, or kidney unresponsiveness to it). Both cause polyuria and thirst, but in diabetes insipidus the urine is dilute and contains no glucose — it is a water-balance disorder, not a sugar disorder.

WARNING Type 2 diabetes is not just "eating sugar"

Attributing type 2 diabetes solely to sugar intake is a common mistake. The core defect is **insulin resistance**, driven by several interacting risk factors: genetic predisposition, obesity (especially abdominal fat), lack of exercise and age. Sugar is only one contributor among many.

Type 1: β -cells destroyed $\rightarrow$ no insulin**Type 2: target cells resist insulin**

Negative-feedback control of blood glucose, showing where type 1 and type 2 diabetes break the loop.

2.7 Thyroid, growth-hormone and adrenal disorders

Thyroid disorders. The thyroid needs **iodine** to make thyroxine. In iodine deficiency the gland enlarges as it works harder to capture iodine, producing **goitre**; endemic goitre has historically occurred in some Cameroonian highland areas where soils and diets are poor in iodine — the reason table salt is iodised. **Hypothyroidism** slows metabolism (fatigue, weight gain, cold intolerance); in children, severe deficiency causes **cretinism** (congenital hypothyroidism: stunted growth and mental impairment). **Hyperthyroidism** accelerates metabolism (weight loss, rapid heartbeat, bulging eyes — exophthalmic goitre).

Growth-hormone disorders (pituitary): hyper-secretion in childhood causes **gigantism**, in adulthood **acromegaly** (enlarged hands, feet, jaw); hypo-secretion in childhood causes pituitary **dwarfism**.

Adrenal disorders (briefly): **Addison's disease** is hypo-secretion of adrenal cortex hormones (weakness, low blood pressure, darkened skin); **Cushing's syndrome** is hyper-secretion of cortisol (weight gain, "moon face", high blood pressure).

DID YOU KNOW? Iodised salt and goitre in Cameroon

Iodine deficiency disorders were once a serious public-health problem in parts of the

Adamawa, North-West and West highlands of Cameroon, where iodine-poor soils produce iodine-poor food crops. The national policy of **salt iodisation** has greatly reduced goitre and, above all, protects unborn babies and children from cretinism. Checking that household salt is iodised is a simple, concrete act of NCD prevention.

2.8 Linking the two control systems

The nervous and endocrine systems meet at the **hypothalamus–pituitary axis**: the hypothalamus (nervous tissue) monitors the blood and directs the pituitary (the "master gland"), which in turn controls the thyroid, adrenals and gonads. Failure at any level of this axis produces disease — for example, lack of ADH (made in the hypothalamus, released by the posterior pituitary) causes diabetes insipidus. This is why the two systems are studied together.

KEY POINTS Key points

- Nervous disorders are located by mechanism: electrical (epilepsy), vascular (stroke), degenerative (Parkinson's, Alzheimer's), autoimmune (multiple sclerosis).
- Stroke is an emergency linked to hypertension; think FAST. Epilepsy is managed by protecting the patient, never restraining, never putting objects in the mouth.
- Endocrine diseases = hypo- or hyper-secretion breaking a negative-feedback loop.
- Diabetes mellitus: type 1 = insulin deficiency (autoimmune); type 2 = insulin resistance (lifestyle-linked). Complications: retinopathy, neuropathy, nephropathy.
- Goitre (iodine deficiency), cretinism, gigantism/acromegaly/dwarfism, Addison's and Cushing's complete the picture.

EXERCISE 2 Glucose curves and feedback

The table gives blood glucose (mmol/L) after a glucose drink:

Time (min)	0	30	60	120
Person X	4.8	8.2	6.4	5.0
Person Y	7.9	13.0	11.8	9.6

1. Which person is diabetic? Give two pieces of evidence.
2. Explain, using negative feedback, why person X's glucose falls after 30 minutes.
3. State one long-term complication person Y risks if untreated.

SOLUTION Exercise 1

1. Person Y is diabetic. Evidence: fasting glucose is already high (7.9 mmol/L); the peak (13.0 mmol/L) is far above normal and exceeds the renal threshold (about 10 mmol/L); glucose remains elevated (9.6 mmol/L) after 120 minutes.
2. In person X, rising blood glucose stimulates the β -cells of the islets of Langerhans to secrete insulin; insulin promotes glucose uptake by liver and muscle cells and its storage as glycogen, so blood glucose falls back to the set point — a negative-feedback response.
3. Any one of: retinopathy (blindness), neuropathy (nerve damage/foot ulcers), nephropathy (kidney damage), or increased risk of cardiovascular disease.

3 SECTION 3: DISEASES OF THE RENAL AND CARDIOVASCULAR SYSTEMS

In Section 2 we saw how diabetes mellitus, an endocrine disorder, damages small blood vessels throughout the body. Two systems suffer most from this silent vascular damage: the **renal system**, which filters the blood, and the **cardiovascular system**, which pumps and carries it. The two are so closely linked that disease in one almost always provokes disease in the other. This section studies the main non-infectious diseases of the renal and cardiovascular systems, which together form the leading causes of adult death and disability in Cameroon today.

3.1 The nephron: structural basis of renal disease


Each kidney contains about one million microscopic filtering units called **nephrons**. Understanding where a nephron can fail is the key to understanding every renal disease.

DEFINITION The nephron

The **nephron** is the functional unit of the kidney. It consists of:

- the **glomerulus**, a ball of capillaries where blood is filtered under pressure;
- **Bowman's capsule**, a cup surrounding the glomerulus that collects the filtrate;
- the **proximal convoluted tubule**, the **loop of Henlé** and the **distal convoluted tubule**, where useful substances (all glucose, most water, salts) are reabsorbed into the blood;
- the **collecting duct**, where final water reabsorption (under ADH control) concentrates the urine.

The logic is simple: the glomerulus is a filter, the tubules are reclaimers. If the filter is damaged, large molecules (proteins) and even cells (red blood cells) leak into the urine. If the tubules fail, useful substances such as glucose are lost and water balance collapses. Keep this filter-reclaimer picture in mind: it explains every urine test result you will meet in the examination.



3.2 Diseases of the kidneys and urinary tract

Nephritis (glomerulonephritis)

Glomerulonephritis is an inflammation of the glomeruli. It often appears one to three weeks after a streptococcal infection (for example a poorly treated sore throat): antibodies produced against the bacterium form immune complexes that lodge in the glomerular membrane and inflame it (post-streptococcal glomerulonephritis). It may also be autoimmune, the immune system attacking the kidney's own tissue. The inflamed filter becomes leaky, so the characteristic signs are **proteinuria** (protein in urine, making it foamy), **haematuria** (blood in urine) and **oedema** (swelling of the face and ankles, because lost plasma protein lowers the osmotic pressure of the blood, so water is retained less effectively in the capillaries and accumulates in the tissues).

Kidney stones (renal calculi)

When urine becomes too concentrated, dissolved mineral salts (calcium oxalate, uric acid) crystallise into hard stones. Risk factors include chronic dehydration — very relevant in the hot Far North of Cameroon, where water intake may be low — and a diet rich in salt and animal protein. A stone stuck in the ureter causes intense **flank pain** (renal colic), often with blood in the urine. Small stones are flushed out by drinking abundantly; large ones may be broken by shock waves


(lithotripsy) or removed surgically. Prevention is simple and cheap: drink enough water to keep the urine dilute.

Renal failure

DEFINITION Renal failure

Renal failure is the inability of the kidneys to filter the blood adequately. Acute renal failure appears suddenly (severe infection, poisoning, shock) and is often reversible. Chronic renal failure develops slowly over years, most commonly as a consequence of hypertension and diabetes mellitus, and is irreversible. Waste products such as urea accumulate in the blood (**uraemia**), with dangerous fluid and electrolyte (especially potassium) imbalance.

When both kidneys fail, two options exist. **Haemodialysis** replaces filtration artificially: the patient's blood is pumped through a machine where it flows along one side of a semi-permeable membrane, while a clean fluid, the **dialysate**, flows on the other side. Urea and excess salts diffuse from blood (high concentration) into the dialysate (low concentration) down their concentration gradients; blood cells and proteins, too large, are retained. A kidney **transplant** is the only true cure, but requires a compatible donor and lifelong immunosuppressive drugs. In Cameroon, dialysis sessions are expensive and dialysis centres are concentrated in big cities such as Yaoundé and Douala, so many patients cannot attend the two to three weekly sessions needed — a major public-health challenge.



Principle of haemodialysis: waste diffuses from blood to dialysate across a semi-permeable membrane; cells and proteins cannot cross.

WARNING Dialysis is not a cure

A very common mistake is to write that dialysis cures kidney failure. It does not: it only replaces the filtration function temporarily and must be repeated two to three times per week for life (or until a transplant). It also does not replace the kidney's hormonal roles (e.g. erythropoietin production, which stimulates red blood cell formation — hence the anaemia of dialysis patients). Only a transplant restores full kidney function.

Urinary tract disorders (non-infectious aspects)

Cystitis, inflammation of the bladder, is usually infectious, but it can also result from irritation by stones or chemicals; it causes frequent, painful urination. **Urinary incontinence** — involuntary leakage of urine — is non-infectious: it results from weakened pelvic floor muscles (after childbirth, in old age) or from nerve damage that prevents proper control of the bladder sphincter.

Disorder	Site / mechanism	Key signs
Glomerulonephritis	Inflamed, leaky glomerular filter	Proteinuria, haematuria, oedema
Kidney stones	Crystallised salts blocking urinary tract	Renal colic, haematuria
Chronic renal failure	Progressive nephron loss (hypertension, diabetes)	Uraemia, fluid and electrolyte imbalance
Urinary incontinence	Weak sphincter or nerve damage	Involuntary urine leakage

3.3 Hypertension: the silent killer

Blood pressure is the force exerted by blood on artery walls. It is expressed by two numbers: the **systolic** pressure (when the ventricles contract) and the **diastolic** pressure (when the heart relaxes), measured in millimetres of mercury (mmHg).

DEFINITION Hypertension


Hypertension is a persistently raised arterial blood pressure. The traditional diagnostic threshold used in most clinical and examination contexts is $\geq 140/90$ mmHg on repeated measurements; more recent international guidelines already classify 130–139/80–89 mmHg as stage 1 hypertension. It is called the “silent killer” because it usually causes no symptoms for years while silently damaging the heart, brain, kidneys and eyes. It is extremely common among Cameroonian adults and is frequently discovered only after a stroke or heart failure.

Risk factors include a salt-rich diet (think of heavily salted stock cubes and smoked fish), obesity, physical inactivity, chronic stress, excessive alcohol, tobacco and heredity. Consequences include heart failure, stroke, kidney damage and retinopathy (damage to the retina that can lead to blindness).

METHOD Reading and classifying a blood-pressure value

A value written 140/90 mmHg means: systolic = 140, diastolic = 90.

1. Identify the systolic (first, higher number) and the diastolic (second number).
2. Compare each with the thresholds: normal < 120/80; elevated 120–129/ < 80; hypertension stage 1: 130–139 or 80–89; stage 2: ≥ 140 or ≥ 90 .
3. Classify by the higher of the two categories. Example: 145/85 is stage 2 (because of the systolic value).
4. A single high reading is not enough: diagnosis requires repeated measurements on different occasions.



Upper limits of systolic and diastolic pressure for each blood-pressure category. Stage 2 values shown are the minimum thresholds ($\geq 140/90$).

3.4 Atherosclerosis, coronary heart disease and heart failure

DEFINITION Atherosclerosis and arteriosclerosis

Atherosclerosis is the deposition of fatty **plaques** (mainly cholesterol) in the inner wall of arteries, narrowing the lumen. **Arteriosclerosis** is the hardening and loss of elasticity of artery walls, often accompanying plaque formation. Both are promoted by a diet rich in saturated fats, smoking, diabetes and hypertension.

A narrowed artery delivers less blood; a plaque can also rupture and trigger a clot (thrombus) that blocks the vessel completely. The consequences depend on the organ supplied: heart (angina, infarction), brain (stroke), kidneys (renal failure).



Cross-sections of a healthy artery (left) and an atherosclerotic artery (right): the plaque narrows the lumen and stiffens the wall.

Coronary heart disease

The heart muscle itself is fed by the coronary arteries. Partial narrowing causes **angina pectoris**: chest pain on effort, when oxygen demand exceeds supply. Complete blockage (usually by a clot on a plaque) causes a **myocardial infarction** (heart attack): the deprived heart muscle dies. Symptoms include crushing central chest pain radiating to the left arm and jaw, sweating and breathlessness. The emergency response is to stop all activity, call for medical help immediately and, if available, give aspirin to limit clotting. Hospital treatment restores blood flow by a **stent** (a small mesh tube keeping the artery open) or a coronary **bypass** (a grafted vessel detouring around the blockage).

WARNING Heart attack ≠ cardiac arrest

Do not confuse the two. A heart attack (myocardial infarction) is a circulation problem: a blocked coronary artery; the heart usually keeps beating and the victim is conscious. A cardiac arrest is an electrical problem: the heart stops pumping, the victim collapses and needs immediate cardiopulmonary resuscitation (CPR) and defibrillation. A heart attack can lead to cardiac arrest, but the words are not interchangeable.

Heart failure, valves and congenital defects

Heart failure is the inability of the heart to pump enough blood for the body's needs — often the end result of hypertension or infarction — causing breathlessness and oedema of the legs. Valvular disorders (narrowed or leaking valves) force the heart to work harder. Congenital defects are present at birth, for example a “hole in the heart” (septal defect) allowing blood to shunt abnormally between chambers; many can be corrected surgically.

Stroke, aneurysms and varicose veins

A **stroke** is the meeting point of cardiovascular and nervous disease (Section 2): a cerebral artery blocked by a clot or ruptured by hypertension deprives brain tissue of oxygen, causing paralysis or loss of speech. An **aneurysm** is a balloon-like swelling of a weakened artery wall that may burst. **Varicose veins** are dilated, tortuous leg veins caused by faulty valves, common in people

who stand for long hours.

PROPERTY The heart–kidney vicious circle

Hypertension damages the small arteries of the kidney, reducing filtration; diseased kidneys retain salt and water and release hormones (renin) that raise blood pressure further. Thus hypertension causes kidney disease, and kidney disease worsens hypertension — a self-reinforcing vicious circle that explains why the two diseases so often occur together and why controlling blood pressure protects the kidneys (and vice versa). (Understanding, not memorisation, is examinable.)

WORKED EXAMPLE 3 Interpreting a urine test and a blood–pressure reading

A 52-year-old trader in Bamenda has a blood pressure of 155/95 mmHg. His urine contains protein and traces of blood, but no glucose. (a) Classify his blood pressure. (b) Which region of the nephron is most likely damaged? Justify. (c) Explain why his ankles are swollen.

SOLUTION

- (a) Systolic $155 \geq 140$ and diastolic $95 \geq 90$: hypertension stage 2.
- (b) Protein and blood cells are normally too large to be filtered; their presence in urine means the filter is leaky, so the damage is in the glomerulus / Bowman's capsule (glomerulonephritis), not the tubules. Absence of glucose confirms that tubular reabsorption still works.
- (c) Loss of plasma protein in urine lowers the osmotic (oncotic) pressure of the blood, so water is retained less effectively in capillaries and accumulates in the tissues: oedema of the ankles. The chronic hypertension has probably damaged his glomeruli — an illustration of the heart–kidney vicious circle.

WARNING Exam tip: link urine composition to the nephron region

In data-response questions, read the urine test logically: protein or blood in urine → glomerulus (leaky filter); glucose in urine → proximal tubule (failed reabsorption) or diabetes (blood glucose above the reabsorption threshold); very dilute, abundant urine → collecting duct / ADH problem. Never write “the kidney is damaged” without naming the precise region.

3.5 Prevention of cardiovascular and renal disease

Most of these diseases are preventable, and prevention is far cheaper than dialysis or bypass surgery. The key measures are: a diet low in salt and saturated fat and rich in vegetables and

fruits; regular physical exercise (at least 30 minutes of brisk walking most days); no tobacco; little or no alcohol; maintaining a healthy body weight; and regular screening — checking blood pressure and blood sugar even when feeling perfectly well, because hypertension and diabetes are silent. In Cameroon, community screening campaigns and reducing salt in cooked food are among the most cost-effective weapons against stroke, heart failure and kidney failure.

KEY POINTS Key points of Section 3

- The nephron = glomerular filter + tubular reclaimer; the site of damage determines the urine abnormalities.
- Glomerulonephritis: leaky glomeruli → protein, blood in urine, oedema. Kidney stones: crystallised salts, prevented by hydration.
- Chronic renal failure (mainly from hypertension and diabetes) is treated by dialysis or transplant; dialysis replaces but does not cure.
- Hypertension (traditionally $\geq 140/90$ mmHg) is a silent killer; atherosclerosis narrows arteries with cholesterol plaques.
- Angina and myocardial infarction result from coronary blockage; heart attack (circulation) differs from cardiac arrest (electrical).
- Heart and kidney form a vicious circle; prevention rests on diet, exercise, no tobacco and regular screening.

EXERCISE 3 Check your understanding

1. A patient's urine contains glucose but no protein. Identify the likely nephron region affected and give one possible underlying disease.
2. Explain why a person living in a hot, dry region who drinks little water is at risk of kidney stones.
3. A man collapses with chest pain but remains conscious. Is this more likely a heart attack or a cardiac arrest? Justify and state the first-aid response.
4. Using the vicious-circle property, explain why treating hypertension slows the progression of chronic kidney disease.

SOLUTION Exercise 1

1. Glucose is normally reabsorbed in the proximal convoluted tubule; its presence in

urine (without protein) indicates tubular reabsorption failure, or more commonly diabetes mellitus, where blood glucose exceeds the tubules' reabsorption capacity. The glomerular filter is intact (no protein).

2. Low water intake produces concentrated urine, in which dissolved mineral salts easily reach saturation and crystallise into stones; heat increases water loss by sweating, worsening concentration.
3. A heart attack (myocardial infarction): the victim is conscious because the heart is still beating — the problem is a blocked coronary artery, not an electrical stop. Response: stop all activity, sit the person down, call for emergency medical help immediately, give aspirin if available and not contraindicated.
4. Lowering blood pressure reduces mechanical damage to the glomerular arterioles, slowing kidney damage; healthier kidneys in turn regulate salt, water and renin better, keeping blood pressure lower. Breaking the circle at any point benefits both organs.

4 SECTION 4: DISORDERS OF THE MUSCULOSKELETAL SYSTEM AND INTEGRATED PREVENTION OF NCDS

In Section 3 we saw how the kidneys and the cardiovascular system can fail without any germ being involved: hypertension, atherosclerosis, kidney failure are diseases of structure and regulation, not of infection. We now complete our tour of the body with the system that gives us shape and movement: the **musculoskeletal system**. A builder in Bamenda who can no longer straighten his knees, a grandmother in Buea who breaks her hip after a simple fall, a schoolboy in Yaoundé whose legs curve outwards — none of these people caught a microbe. Their bones, joints or muscles have failed. Understanding why requires us first to recall how these structures are built and maintained.


4.1 Review: bones, joints and skeletal muscle

Bones. Bone is a living, vascularised tissue, not dead scaffolding. It consists of a matrix of **collagen** fibres (giving tensile strength and flexibility) hardened by **calcium phosphate** salts, mainly hydroxyapatite (giving compressive rigidity). Two cell types constantly remodel it: **osteoblasts** deposit new bone matrix and mineralise it, while **osteoclasts** resorb old bone by secreting acid and enzymes. In a healthy adult the two activities are coupled and balanced, so total bone mass is stable. Bone also acts as the body's major mineral store: when blood calcium falls, parathyroid hormone stimulates osteoclasts to release calcium from bone into the blood. This dual role — structural support and mineral reservoir — explains most bone diseases.

Joints. Freely movable joints such as the knee, hip and shoulder are **synovial joints**. The ends of the two bones are covered with smooth **articular (hyaline) cartilage**, a slippery, shock-absorbing tissue with no blood supply of its own (it receives nutrients by diffusion from synovial fluid). The joint is enclosed in a fibrous capsule lined by the **synovial membrane**, which secretes **synovial fluid** that lubricates the joint and nourishes the cartilage. Ligaments (dense regular connective

tissue) hold the bones together and stabilise the joint.

Skeletal muscle. Muscles are bundles of multinucleated fibres that contract when stimulated by motor neurones at the **neuromuscular junction**, where the neurotransmitter **acetylcholine** is released onto nicotinic receptors on the motor end plate. Tendons (dense regular connective tissue) attach muscle to bone. Movement therefore requires four healthy partners: bone, joint, muscle, and nerve — and a disorder can strike any of them.



A synovial joint in three states. Healthy: thick articular cartilage (orange) separates the bone ends. Osteoarthritis: cartilage is worn away so bones rub directly. Rheumatoid arthritis: the synovial membrane (pink) is inflamed and swollen (pannus), invading the joint space and eroding cartilage and bone.

4.2 Arthritis: osteoarthritis versus rheumatoid arthritis

DEFINITION Arthritis

Arthritis is inflammation or degeneration of one or more joints, producing the cardinal signs of pain, stiffness, swelling and, in advanced cases, deformity and loss of function. The two commonest forms are **osteoarthritis** and **rheumatoid arthritis**.

Osteoarthritis (OA) is the “wear-and-tear” arthritis. With age, repetitive heavy use, or previous injury, the articular cartilage — which has no blood supply and limited repair capacity — becomes thin, rough and fissured. Subchondral bone thickens (sclerosis) and osteophytes (bony spurs) form at joint margins. Bone ends rub directly against each other, causing pain that is characteristically worse with movement and weight-bearing, and relieved by rest. It typically affects weight-bearing joints (knees, hips, lumbar and cervical spine) and the distal interphalangeal joints (Heberden’s nodes) and base of thumb in older people. Risk factors include age, obesity (extra mechanical load and adipokine-mediated inflammation), previous joint injury, heavy manual labour and genetic predisposition.

Rheumatoid arthritis (RA) is a chronic systemic **autoimmune** disease: the immune system mistakenly attacks the synovial membrane, which becomes infiltrated with lymphocytes and plasma cells, thickened (pannus formation) and invasive. The pannus erodes articular cartilage and underlying bone. It often begins in young or middle-aged adults (30–50 years), affects symmetrical small joints (metacarpophalangeal, proximal interphalangeal, wrists, MTP joints), and causes prolonged morning stiffness lasting more than one hour that eases with activity. Systemic features include fatigue, low-grade fever, weight loss, and extra-articular manifesta-

tions (rheumatoid nodules, vasculitis, lung fibrosis). Deformities of the fingers (ulnar deviation, swan-neck, boutonnière) are characteristic late signs.

PROPERTY Degenerative versus autoimmune joint disease

Osteoarthritis is a degenerative disease: the primary lesion is mechanical and biochemical breakdown of articular cartilage; pain worsens with activity and improves with rest; it is commoner in the elderly and in weight-bearing joints; systemic inflammation is absent. Rheumatoid arthritis is an autoimmune inflammatory disease: the primary lesion is immune-mediated inflammation of the synovial membrane (synovitis); stiffness is worst after rest (prolonged morning stiffness); it can occur at any adult age and affects small joints symmetrically; systemic inflammation is present (raised ESR, CRP, rheumatoid factor, anti-CCP antibodies). (The justification of this distinction is examinable as a comparison table.)

Management. Neither form can be cured, but both can be controlled to preserve function and quality of life:

- **OA:** Weight reduction (crucial for lower limbs), physiotherapy (quadriceps strengthening for knee OA), walking aids, analgesics (paracetamol), NSAIDs (topical or oral), intra-articular corticosteroid injections; in severe refractory cases, surgical joint replacement (total hip or knee arthroplasty).
- **RA:** Early referral to a rheumatologist. Disease-modifying antirheumatic drugs (DMARDs) — methotrexate is the anchor drug — started as early as possible to induce remission; biologic agents (anti-TNF, anti-IL6, etc.) for inadequate responders; short-term glucocorticoids for flares; NSAIDs for symptom control; physiotherapy and occupational therapy; surgery for established deformities.

4.3 Gout: crystals in the joint

Gout is a painful crystal-induced arthritis caused by deposition of monosodium **urate** crystals in a joint, classically the first metatarsophalangeal joint (podagra). Urate is the end product of purine catabolism. Normally the **kidneys excrete** the majority of urate (the rest via gut). Gout arises when serum urate exceeds its solubility limit (6.8 mg/dL or 400 µmol/L), leading to crystal precipitation. This occurs due to overproduction (high purine diet: red meat, offal, certain fish, yeast extracts; high fructose intake; tumour lysis) or, more commonly, underexcretion (chronic kidney disease, diuretics, alcohol — especially beer — which increases urate production and decreases excretion). The crystals trigger intense inflammation via the NLRP3 inflammasome: a red, hot, exquisitely tender joint, often waking the patient at night. Notice the link with Section 3: gout sits at the crossroads of diet, renal function and cardiovascular risk (metabolic syndrome).

Management: Acute attack — NSAIDs (indomethacin, naproxen), colchicine (low dose), or intra-articular/systemic glucocorticoids. Long-term — urate-lowering therapy (allopurinol or febuxostat) once the acute attack settles, targeting serum urate < 360 µmol/L (6 mg/dL); lifestyle: hydration, limit alcohol and high-purine foods, weight loss, treat comorbidities (hypertension, CKD).

DID YOU KNOW? Gout in the Cameroonian context

Gout is increasingly diagnosed in urban Cameroon with the nutrition transition towards more red meat, organ meats (“kpem”, tripe), sugary drinks and beer consumption. Traditional diets rich in plantains, vegetables and fish were protective. Screening for hyperuricaemia in hypertensive or diabetic patients in primary care can prevent gout and may slow CKD progression.


4.4 Osteoporosis: the silent thinning of bone

DEFINITION Osteoporosis

Osteoporosis is a systemic skeletal disease characterised by low bone mineral density (BMD) and microarchitectural deterioration of bone tissue, with a consequent increase in bone fragility and susceptibility to fragility fractures (fractures occurring from a fall from standing height or less). Typical sites are the hip (neck of femur), the spine (vertebral crush fractures causing loss of height, kyphosis, chronic back pain) and the distal radius (Colles’ fracture).

Recall that bone is constantly remodelled by osteoblasts and osteoclasts in basic multicellular units (BMUs). Osteoporosis results when resorption exceeds deposition over many years, leading to negative bone balance. Risk factors are:

- **Age:** bone mass peaks around age 25–30 (peak bone mass) and declines thereafter (0.5–1% per year).
- **Sex and menopause:** oestrogen normally restrains osteoclast activity and promotes osteoblast function; after menopause women lose bone rapidly (2–3% per year for 5–10 years) — this is why osteoporosis is commoner in elderly women.
- **Calcium and vitamin-D deficiency:** too little raw material and poor intestinal calcium absorption (vitamin D deficiency is common even in sunny climates due to indoor lifestyles, veiling, sunscreen, dark skin).
- **Physical inactivity:** bone strengthens in response to mechanical load (Wolff’s law); a sedentary life weakens it.
- **Secondary causes:** glucocorticoid excess (Cushing’s, therapy), hyperthyroidism, hyperparathyroidism, hypogonadism, malabsorption, chronic liver/kidney disease, rheumatoid arthritis.
- Smoking and excess alcohol, which also accelerate bone loss.



Normal versus osteoporotic bone (schematic cross-section of trabecular bone). The same outer cortical size hides a much weaker interior: fewer and thinner bony trabeculae with larger marrow spaces, so the bone fractures under loads it once tolerated.

Diagnosis: Dual-energy X-ray absorptiometry (DXA) scan at hip and lumbar spine. T-score ≤ -2.5 SD = osteoporosis; -1 to -2.5 = osteopenia. **Prevention** must begin in youth: a diet rich in calcium (milk, yogurt, cheese, small fish eaten with bones like “canda”, green leafy vegetables) and vitamin D, plus regular **weight-bearing exercise** (walking, running, farm work, dancing, sport) and resistance training, which stimulate osteoblasts. Fall prevention in the elderly (home safety, vision correction, balance training). **Treatment:** Calcium and vitamin-D supplements; bisphosphonates (alendronate, risedronate) as first-line to inhibit osteoclasts; denosumab (anti-RANKL); teriparatide (recombinant PTH) for severe cases; hormone replacement therapy (HRT) for early postmenopausal women with indications.

4.5 Rickets and osteomalacia: soft bones

Where osteoporosis is a loss of amount of normally mineralised bone, **rickets** (in children) and **osteomalacia** (in adults) are defects of mineralisation: the collagen matrix (osteoid) is laid down but not hardened with calcium salts, because of **vitamin-D or calcium deficiency**. Vitamin D (cholecalciferol) is needed for active intestinal absorption of calcium and phosphate; it is obtained from the diet (oily fish, egg yolk, fortified foods) and synthesised in the skin from 7-dehydrocholesterol under UV-B sunlight.

In children, whose bones are still growing at the growth plates, soft bones bend under body weight: **bow-legs (genu varum)** or **knock-knees (genu valgum)**, **rachitic rosary** (costochondral junction swelling), **Harrison’s groove** (diaphragmatic pull on soft ribs), **craniotables** (soft skull), delayed fontanelle closure, delayed dentition, and growth failure — this is rickets. In adults the same deficiency causes diffuse **bone pain** (often lower back, pelvis, thighs), **proximal muscle weakness** (difficulty climbing stairs, rising from a chair), and pseudofractures (Looser’s zones) — this is osteomalacia. Prevention is simple and cheap: adequate safe sunlight exposure (15–30 minutes on arms/face several times a week; abundant in Cameroon, though heavily veiled infants, strictly indoor lifestyles, and dark-skinned individuals in high latitudes are at risk), and a diet containing calcium and vitamin D (fish, eggs, fortified flour/milk). Treatment: high-dose vitamin D (ergocalciferol or cholecalciferol) and calcium supplements.

WARNING Rickets is not osteoporosis

A classic GCE error is to confuse rickets with osteoporosis. Rickets affects children: the bone is poorly mineralised (excess osteoid) and becomes soft and deformed (bow-legs, rachitic rosary) because growth plates are active. Osteoporosis affects adults, especially after menopause: the bone is normally mineralised but reduced in quantity (thin cortices, lost trabeculae) and becomes fragile, breaking rather than bending. Different cause (mineralisation defect vs. resorption excess), different age group, different radiology and biochemistry.

4.6 Muscular disorders

DEFINITION Muscular dystrophy

Muscular dystrophy is a group of inherited (genetic) disorders characterised by progressive degeneration and necrosis of skeletal muscle fibres, replaced by fat and fibrous tissue, causing progressive muscle wasting and weakness. The commonest and most severe form, **Duchenne muscular dystrophy (DMD)**, is caused by mutations in the DMD gene on the X chromosome (Xp21) encoding **dystrophin**, a protein linking the cytoskeleton to the extracellular matrix. It affects almost exclusively boys, with onset at 2–5 years (delayed walking, Gowers' sign, calf pseudohypertrophy), wheelchair dependence by early teens, and death from cardiomyopathy or respiratory failure in the 20s. Becker muscular dystrophy is a milder allelic variant.

WARNING Muscular dystrophy is genetic, not contagious

Never present muscular dystrophy as contagious, or as the result of laziness, poor feeding or lack of exercise. It is caused by a faulty gene present from conception; no one can “catch” it, and no amount of effort by the child could prevent it. Confusing a genetic disease with an infectious or lifestyle disease is a serious conceptual error in the examination. Genetic counselling for carrier mothers is essential.

Myasthenia gravis is an **autoimmune** disorder of the neuromuscular junction: autoantibodies (mainly against the nicotinic acetylcholine receptor, AChR) block, cross-link and accelerate degradation of receptors, so nerve impulses no longer trigger contraction effectively. The hallmark is fatigable weakness — muscles (often ocular: ptosis, diplopia; then bulbar: dysphagia, dysarthria; then limb proximal) weaken with repeated use and recover after rest. Diagnosis: positive anti-AChR or anti-MuSK antibodies, decremental response on repetitive nerve stimulation, improvement with edrophonium (Tensilon test). Treatment: acetylcholinesterase inhibitors (pyridostigmine), immunosuppression (corticosteroids, azathioprine, mycophenolate), thymectomy (if thymoma or generalised early-onset), plasma exchange/IVIG for crises.

Cramps and sprains are everyday muscular problems. A **cramp** is a sudden, painful, involuntary tonic contraction, often linked to dehydration, electrolyte loss in sweat (think of athletes in the

dry-season heat), muscle fatigue, or nerve compression. A **sprain** is the stretching or tearing of ligaments around a joint, typically the lateral ankle (inversion injury).

4.7 Back pain, herniated disc, posture and sports injuries

The spine is a stack of vertebrae separated by **intervertebral discs** — a tough outer **annulus fibrosus** with a soft, gelatinous **nucleus pulposus** that act as shock absorbers. Lifting a heavy load with a flexed spine (instead of bending the knees and keeping the back straight) generates huge compressive and shear forces that can tear the annulus so the nucleus protrudes: a **slipped (herniated/prolapsed) disc**. The protrusion may compress a spinal nerve root (commonly L5 or S1), causing **sciatica** — radiating pain, numbness or weakness down the leg. Chronic **non-specific low back pain** is also promoted by poor posture, weak core (abdominal and paraspinal) muscles, obesity, smoking (disc degeneration), and long hours sitting — increasingly common among students and office workers. Prevention: correct lifting technique (bend knees, keep load close, avoid twisting), good sitting posture (lumbar support, feet flat), regular core-strengthening exercise, weight control, and a medium-firm mattress.

Sports injuries include:

- **Sprains**: ligament tears (grades I–III).
- **Strains**: muscle or tendon tears.
- **Dislocations**: bone forced out of its joint (shoulder, elbow, fingers) — visible deformity, loss of movement.
- **Fractures**: broken bone (simple/closed, compound/open with skin breach, greenstick in children, comminuted).

METHOD The RICE first-aid protocol for acute soft-tissue injuries (sprains, strains, contusions)

1. **R — Rest**: stop the activity immediately; do not “walk it off”. Immobilise the injured part (splint/sling if needed).
2. **I — Ice**: apply ice wrapped in a damp cloth (never directly on skin) for 15–20 minutes every 2–3 hours for the first 48–72 hours to reduce swelling, pain and metabolic rate.
3. **C — Compression**: apply a firm elastic bandage (not so tight as to cut off circulation — check capillary refill) to limit swelling and provide support.
4. **E — Elevation**: raise the injured limb above heart level whenever possible so venous and lymphatic fluid drains away from the injury.

Then seek medical care, especially if a fracture or dislocation is suspected (deformity, inability to bear weight, bone visible, neurovascular compromise). Never try to reduce a dislocated joint yourself — this can damage nerves and vessels.

4.8 Congenital skeletal disorders

Some skeletal defects are present at birth. **Congenital talipes equinovarus (club foot)** is a foot fixed in equinus (plantarflexion), varus (inversion) and adduction; it is treated successfully by the **Ponseti method** — gentle weekly manipulation and above-knee plaster casts in infancy, followed by a minor Achilles tenotomy and bracing (boots and bar) for several years. **Spina bi-fida** (myelomeningocele) is a failure of the posterior vertebral arches to close over the spinal cord during neurulation (weeks 3–4); severe forms cause paralysis of the legs, sensory loss, and neurogenic bladder/bowel. Its frequency is strongly reduced (by up to 70%) when the mother has adequate **folic acid** (400 µg daily) before conception and during the first trimester — a powerful argument for preconception care, food fortification, and a varied diet (green leafy vegetables, beans, fortified flour) for women of child-bearing age.

4.9 Synthesis: shared risk factors and the integrated prevention of NCDs


Step back now and look at everything this chapter has covered — reproductive cancers, diabetes, hypertension, stroke, kidney failure, arthritis, osteoporosis, muscular disorders. Different organs, different symptoms, yet the same handful of modifiable risk factors keeps reappearing:

- **Tobacco use** — cancers (cervix, prostate, lung), atherosclerosis, stroke, osteoporosis, poor wound healing;
- **Harmful alcohol use** — hypertension, liver cirrhosis, gout, cancers, cardiomyopathy, fetal alcohol syndrome;
- **Unhealthy diet** (excess salt, sugar, saturated/trans fats, red/processed meat; too little fruit, vegetables, fibre, calcium) — obesity, type 2 diabetes, hypertension, gout, osteoporosis, constipation/diverticular disease;
- **Physical inactivity** — obesity, type 2 diabetes, hypertension, osteoporosis, back pain, colon cancer.

Because the risk factors are shared, prevention can be integrated. The World Health Organization promotes cost-effective “best buy” interventions for low- and middle-income countries: raising excise taxes on tobacco and alcohol, reducing salt in processed foods (reformulation), eliminating industrial trans fats, public awareness campaigns (mass media, school curricula), and screening (blood-pressure checks, cervical smears/HPV testing, blood-glucose, cholesterol) in primary health centres. Vaccination against hepatitis B (liver cancer) and HPV (cervical cancer) is also a “best buy”. Prevention is a shared responsibility:

- the **individual**: balanced diet (traditional Cameroonian dishes: more vegetables, less oil/salt/bouillon cubes), regular exercise (walking, farming, sport), no tobacco, moderate or no alcohol, attending screening;
- the **family**: healthy cooking habits (steaming, grilling, less salt and palm oil), encouraging active play for children, supporting antenatal care and folic acid for expectant mothers;
- the **school**: health education, compulsory physical education, healthy school meals, tobacco-free campuses, safe water and sanitation;
- the **government**: taxation and regulation of tobacco/alcohol/unhealthy foods, equipping health centres for NCD screening and essential medicines, urban planning for walkable cities,

food fortification (salt iodisation, flour folic acid/iron), vaccination programmes, and sustained public campaigns.



Concept map: the major non-infectious diseases of the six body systems share the same four modifiable risk factors, so a single integrated prevention strategy acts on all systems at once.

WORKED EXAMPLE 4 Building a comparison table under exam conditions

Statement. Construct a table comparing osteoarthritis and rheumatoid arthritis under the headings: cause, joints affected, pattern of pain/stiffness, and management. (8 marks)

SOLUTION

Solution. A clear table with four rows earns marks efficiently:

Feature	Osteoarthritis	Rheumatoid arthritis
Cause	Degeneration (wear and tear) of articular cartilage; age-related	Autoimmune inflammation of the synovial membrane
Joints affected	Weight-bearing joints (knee, hip, spine); often unilateral/asymmetrical	Small joints of hands and wrists; symmetrical
Pain / stiffness	Pain worse with activity, relieved by rest; brief morning stiffness (<30 min)	Morning stiffness >1 hour, eases with movement; pain at rest
Management	Weight loss, physiotherapy, analgesics/NSAIDs, joint replacement	DMARDs (methotrexate), biologics, glucocorticoids, physiotherapy

Each correct, contrasted row earns 2 marks. Note the technique: identical headings down the side, one disease per column, and parallel wording so the contrast is visible at a glance.

WARNING Exam tip: master the comparison-table format

Structured GCE questions very often ask you to “compare disorders X and Y with respect to cause, symptoms, diagnosis, treatment and prevention” — for example osteoarthritis vs rheumatoid arthritis, rickets vs osteoporosis, or Type 1 vs Type 2 diabetes. Practise drawing the table skeleton (headings first) in under a minute, then filling one row at a time. Never write the two diseases in two separate essays: examiners award marks for point-by-point contrast, which only a table (or paired statements) displays clearly.

EXPERIMENT Practical: Assessing joint range of motion and RICE application

Objective: Practise measuring active and passive range of motion (ROM) of the knee and ankle using a goniometer, and demonstrate the RICE protocol on a simulated ankle sprain. **Materials:** Goniometer, ice packs, elastic bandages, pillows, consenting partner. **Procedure:**

1. Measure active ROM: ask partner to flex/extend knee and dorsiflex/plantarflex ankle; record angles.
2. Measure passive ROM: you move the joint through its range; note any pain, crepitus, or end-feel (firm, soft, hard).
3. Simulate RICE: Rest (partner sits), Ice (wrapped pack on lateral ankle 15 min), Compression (figure-of-eight bandage), Elevation (leg on pillows > heart level).

4. Re-assess swelling and pain after 20 min.

Report: Record initial and post-RICE measurements. Discuss why ice is wrapped (prevent ice burn) and why compression must not impair circulation (check capillary refill < 3 s).

EXERCISE 4 Applying your knowledge

1. A 65-year-old woman breaks her hip after slipping gently on a wet kitchen floor. Name the most likely underlying bone disease and give three risk factors she may have.
2. Explain why a person with poorly functioning kidneys (chronic kidney disease stage 3) has an increased risk of gout.
3. A footballer twists his ankle during a match in Douala. Outline the first aid you would give on the pitch.
4. State one difference and one similarity between rickets and osteomalacia.
5. A 10-year-old boy presents with progressive difficulty climbing stairs, frequent falls, and enlarged calf muscles. Name the most likely diagnosis, its inheritance pattern, and the deficient protein.

SOLUTION Exercise 1

1. Osteoporosis: a fragility fracture (hip fracture from standing height fall) is its hallmark. Risk factors: age > 60, post-menopausal (loss of oestrogen), possible calcium/vitamin-D deficiency, physical inactivity, smoking, excess alcohol, low body weight, family history.
2. Uric acid is excreted primarily by the kidneys (glomerular filtration, tubular secretion/reabsorption). In CKD, reduced GFR and tubular dysfunction decrease urate excretion, so serum urate rises above saturation, leading to monosodium urate crystal deposition in joints and soft tissues, causing gout.
3. Apply RICE immediately: Rest — stop play, do not weight-bear; Ice — apply ice wrapped in cloth for 15–20 min; Compression — firm elastic bandage (figure-of-eight); Elevation — raise leg on bag/bench above hip level. Refer to health facility for X-ray to exclude fracture (Ottawa ankle rules).
4. Difference: rickets occurs in children with open growth plates, causing bone deformities (bow-legs, rachitic rosary); osteomalacia occurs in adults after growth plate

fusion, causing bone pain and proximal muscle weakness without deformity. Similarity: both are caused by vitamin-D and/or calcium deficiency leading to defective mineralisation of osteoid (excess unmineralised bone matrix).

5. Duchenne muscular dystrophy (DMD); X-linked recessive inheritance (affects boys, carrier mothers); deficient protein: dystrophin.

KEY POINTS Key points of Section 4

- Bone is living tissue remodelled by osteoblasts (formation) and osteoclasts (resorption); synovial joints rely on articular cartilage and synovial fluid; muscles contract via the neuromuscular junction (acetylcholine).
- Osteoarthritis = degenerative cartilage loss (age, mechanical load) affecting weight-bearing joints; pain on movement. Rheumatoid arthritis = autoimmune synovitis affecting small joints symmetrically; prolonged morning stiffness, systemic features.
- Gout = monosodium urate crystal deposition due to hyperuricaemia (overproduction/underexcretion); linked to diet, alcohol, CKD; acute podagra, chronic tophi.
- Osteoporosis = low bone mass + microarchitectural deterioration → fragility fractures (hip, spine, wrist); risk: age, menopause, low Ca/Vit D, inactivity, glucocorticoids. Prevention: peak bone mass in youth, weight-bearing exercise, Ca/Vit D, fall prevention.
- Rickets (children) / osteomalacia (adults) = defective mineralisation from Vit D/Ca deficiency → soft bones (deformity in children, pain/weakness in adults).
- Muscular dystrophy = genetic (e.g., DMD: X-linked, dystrophin deficiency); myasthenia gravis = autoimmune (anti-AChR antibodies) → fatigable weakness.
- Acute soft-tissue injury: RICE (Rest, Ice, Compression, Elevation). Never reduce a dislocation yourself.
- Congenital: club foot (Ponseti method), spina bifida (folic acid prevention pre-conception).
- All major NCDs share four modifiable risk factors (tobacco, alcohol, unhealthy diet, physical inactivity) → integrated prevention (WHO best buys) involving individual, family, school, government.

5 INTEGRATION ACTIVITY

5.1 Aims of the activity

By the end of this integration activity, you should be able to:

- **Interpret clinical data** (blood pressure readings, urine tests, blood glucose values, physical examination findings) and link each abnormal result to the body system concerned;
- **Propose and justify a diagnosis** of a non-infectious disease affecting the reproductive, nervous, endocrine, renal, cardiovascular or musculoskeletal system;
- **Explain the underlying biological mechanism** of each disorder using correct Upper Sixth terminology;
- **Suggest confirmatory tests** and realistic **treatment and lifestyle measures**;
- **Work as a team** to design a community prevention campaign targeting the shared, modifiable risk factors of non-communicable diseases (NCDs): excess salt, tobacco, alcohol, physical inactivity and unhealthy diet;
- **Communicate scientific ideas** clearly in a short oral presentation, as expected in the GCE Advanced Level practical and problem-solving questions.

KEY POINTS Why this matters

Non-communicable diseases such as hypertension, diabetes, kidney failure, stroke, cancers and osteoporosis are now leading causes of illness and death among Cameroonian adults. Unlike malaria or cholera, they cannot be cured with a short course of drugs: they are **prevented** and **managed**. Understanding them makes you a genuine resource person in your family and quarter.

5.2 Situation

Health-screening day at the Lycée de Nkolbisson.

As part of the third-term health education programme, your school has organised a fictional free community screening day in the school hall. Volunteer nurses have measured blood pressure, tested urine with dipsticks, measured fasting blood glucose with a glucometer, and recorded simple examination findings for volunteer adults from the Nkolbisson neighbourhood. The results have been anonymised and compiled into four **case files**. Your class is divided into **community health teams** of four to six students. Each team acts as the medical analysis unit for the day: you must study the case files, propose diagnoses, and then design a prevention poster for the quarter.

Case file 1 — Mrs A., 52 years old, market trader.

Blood pressure measured twice, 15 minutes apart: 165/105 mmHg both times. She reports headaches and swollen ankles in the evening. Urine dipstick: **protein +++** (marked proteinuria). She adds

a stock cube (Maggi) and extra salt to most dishes and does little physical exercise.

Case file 2 — Mr B., 30 years old, moto-taxi driver.

For three months he has complained of **excessive thirst** (polydipsia), **frequent and abundant urination** (polyuria) and unusual tiredness. Fasting blood glucose measured this morning: 9.5 mmol/L (normal fasting value: 3.9 mmol/L to 5.5 mmol/L; diabetes is suspected above 7.0 mmol/L on two occasions). His father was diagnosed with diabetes at 45.

Case file 3 — Mr C., 60 years old, retired teacher.

He reports **difficulty starting urination**, a weak urine stream, and the need to urinate several times at night (nocturia). Digital rectal examination by the volunteer doctor: **enlarged prostate**, smooth and firm. No pain, no fever, urine dipstick negative for blood and nitrites.

Case file 4 — Mrs D., 68 years old, farmer.

She slipped on a wet floor at home — a minor fall — and **fractured her hip**. X-ray and a bone density scan (DEXA) show **low bone mineral density** (T-score below -2.5). She went through menopause at 50, drinks little milk, and has become much less active with age.

Reference data provided to all teams:

Measurement	Normal / reference value
Resting blood pressure	below 140/90 mmHg (ideal about 120/80)
Fasting blood glucose	3.9–5.5 mmol L ⁻¹ ; diabetes suspected ≥ 7.0 mmol L ⁻¹ twice
Urine protein (dipstick)	negative or trace
Prostate (rectal examination)	small, walnut-sized, smooth
Bone density (DEXA T-score)	above -1.0 normal; -1.0 to -2.5 osteopenia; below -2.5 osteoporosis

5.3 Tasks

Part A — The detective work (for each of the four case files):

1. Identify the **body system** affected and the **organ(s)** involved.
2. Suggest the **most likely non-infectious disease**, justifying your answer with at least two pieces of evidence taken from the case file and the reference table.
3. Explain the **underlying biological mechanism** of the disorder (for example: how the structure or regulation of the organ is disturbed, which hormone or tissue is involved, and how this produces the observed symptoms).
4. Propose **one or two further tests** that would confirm the diagnosis.
5. Recommend appropriate **treatment and lifestyle changes**, adapted to the patient's daily life in Nkolbisson.

Part B — Connecting the cases:

6. For Case 1, explain how untreated hypertension can **damage the kidneys**, and how kidney damage in turn **raises blood pressure** (a vicious circle). Which hormone system, studied in the chapter on the endocrine and renal systems, is involved?

7. For Case 2, explain why high blood glucose causes **thirst and polyuria** (think of osmosis in the kidney tubules), and name **two long-term complications** of uncontrolled diabetes affecting other systems studied in this chapter.
8. Identify the **shared, modifiable risk factors** appearing across the four files (diet, salt, inactivity, age, heredity, hormonal changes). Classify them as modifiable or non-modifiable.

Part C — The community campaign:

9. As a team, design a **one-page poster** entitled “Stop NCDs in our quarter”. It must contain: (i) one clear message per shared risk factor (salt, tobacco, alcohol, inactivity); (ii) one simple diagram or pictogram; (iii) one slogan in language the neighbourhood understands.
10. Present your four diagnoses and your poster to the class in a **5-minute plenary**, and be ready to defend your reasoning against questions from the other teams.

WARNING Common traps

- Do not confuse benign prostatic hyperplasia with prostate cancer: a smooth, firm, painless enlargement in a 60-year-old suggests BPH, but only further tests (PSA blood test, biopsy if indicated) can exclude cancer.
- A single high blood glucose or blood pressure reading is not enough for a diagnosis: values must be confirmed on a second occasion.
- Osteoporosis is not an infectious or inflammatory joint disease: it is a loss of bone mass that weakens the skeleton silently until a fracture occurs.
- Proteinuria is a sign of kidney damage, not a disease in itself.

5.4 Model answer

Case 1 — Mrs A. (hypertension with kidney damage).

System: cardiovascular system, with renal involvement. Diagnosis: chronic **hypertension** (BP 165/105 mmHg, confirmed on two readings, well above 140/90) with signs of **chronic kidney disease** (proteinuria +++, ankle oedema). Mechanism: persistently high pressure damages the walls of arterioles, including the glomerular capillaries; the glomerular filter becomes leaky, so plasma proteins pass into the urine (proteinuria). Damaged kidneys retain salt and water, increasing blood volume and activating the **renin-angiotensin-aldosterone system**, which raises blood pressure further — a vicious circle. Fluid retention explains the swollen ankles. Confirmatory tests: serum creatinine and estimated glomerular filtration rate, 24-hour urine protein, ambulatory BP monitoring. Management: antihypertensive drugs prescribed by a doctor (e.g. a diuretic or an ACE inhibitor, which also protects the kidney), drastic reduction of added salt and stock cubes, regular walking, weight control, and follow-up at the district hospital.

Case 2 — Mr B. (diabetes mellitus).

System: endocrine system (pancreas / insulin). Diagnosis: **diabetes mellitus** — fasting glucose 9.5 mmol/L exceeds the 7.0 mmol/L threshold, with the classic symptoms of polydipsia

and polyuria; family history supports the diagnosis. At age 30 this could be type 1 or early type 2; the distinction needs further tests. Mechanism: insulin deficiency (type 1: autoimmune destruction of the β -cells of the islets of Langerhans) or insulin resistance (type 2) prevents glucose uptake by cells, so blood glucose rises. When the renal threshold is exceeded, glucose appears in the tubular fluid; by **osmosis** it holds water in the tubule, producing polyuria, dehydration and therefore intense thirst. Confirmatory tests: repeat fasting glucose, glycated haemoglobin (HbA1c), urine test for glucose and ketones. Management: medical treatment (insulin for type 1; oral antidiabetics for type 2), reduced intake of sugary drinks and refined carbohydrates, regular physical activity, and education on foot care and eye checks, since uncontrolled diabetes damages the **retina, kidneys, nerves and blood vessels**.

Case 3 — Mr C. (benign prostatic hyperplasia).

System: male reproductive system (prostate gland), affecting urination. Diagnosis: **benign prostatic hyperplasia** (BPH) — age 60, smooth firm enlargement, obstructive urinary symptoms, no signs of infection. Mechanism: under the continuing influence of androgens, the prostate tissue around the urethra proliferates with age; the enlarged gland **compresses the urethra**, so the bladder must contract harder, the stream is weak, emptying is incomplete and nocturia results. Confirmatory tests: PSA blood test (to help exclude cancer), ultrasound of the prostate and bladder, urine flow measurement. Management: drugs that relax the prostate smooth muscle or shrink the gland, surgery (e.g. transurethral resection) if obstruction is severe, reduced evening fluid and caffeine intake.

Case 4 — Mrs D. (osteoporosis).

System: musculoskeletal system (bone tissue). Diagnosis: **osteoporosis** with a fragility fracture — a hip fracture after a minor fall plus a DEXA T-score below -2.5 is diagnostic. Mechanism: bone is continuously remodelled by **osteoclasts** (resorption) and **osteoblasts** (deposition). After menopause the fall in **oestrogen** removes the brake on osteoclast activity, so resorption exceeds formation; bone mass and density fall, trabeculae thin, and the bone breaks under minimal trauma. Low calcium intake and inactivity accelerate the loss. Confirmatory tests: already done (DEXA); blood calcium, phosphate and vitamin D complete the assessment. Management: calcium- and vitamin-D-rich diet (milk, small fish eaten with bones, green vegetables), weight-bearing exercise adapted to her age, prescribed anti-resorptive drugs, and fall-proofing the home.

Connecting the cases (Part B).

Hypertension and kidney disease reinforce each other through the renin-angiotensin-aldosterone system; diabetes damages kidneys, eyes, nerves and vessels; all four patients share modifiable risk factors (excess salt, sugary diet, inactivity, tobacco/alcohol where present) and non-modifiable ones (age, sex, heredity, menopause). Prevention must therefore act on the modifiable factors, early, at community level.

Poster campaign (Part C) — expected content.

A good poster shows, for example, a heart, a kidney, a pancreas and a bone around the slogan “Less salt, more steps — our quarter, our health!”, with four pictograms: a salt cellar crossed out, a cigarette crossed out, a bottle replaced by water, and a walking figure. Messages must be short, positive and actionable: cook with less Maggi, walk 30 minutes a day, check your blood pressure once a year, screening saves lives.

EXERCISE 5 Going further

A fifth volunteer, aged 45, has a fasting blood glucose of 6.2 mmol/L and a blood pressure of 138/88 mmHg. Does he have diabetes or hypertension? What advice would your team give him, and why is this “borderline” stage actually the most important moment for prevention?

SOLUTION Going further

Neither diagnosis can be made: his glucose (6.2) is above the ideal range but below the 7.0 mmol L⁻¹ threshold (this is impaired fasting glucose, a pre-diabetic state), and his BP (138/88) is high-normal but below 140/90. He should be re-tested to confirm the values, and advised to act now: reduce sugar and salt, increase physical activity, control weight, avoid tobacco and excess alcohol, and attend yearly screening. This borderline stage is decisive because lifestyle change at this point can prevent or delay the onset of diabetes and hypertension, whereas once organ damage (kidney, eye, heart) has occurred it is largely irreversible. Prevention is cheaper and more effective than cure — the central message of the whole chapter.

6 METHODS AND WORKED EXAMPLES

6.1 Methods and worked examples

This part trains the practical skills the GCE examiner expects: reading clinical data (urine, blood, blood pressure), linking symptoms to the correct organ system, reasoning with hormone feedback loops, and building a prevention plan. Each method card is followed by a fully worked GCE-style exercise. Master these patterns and you will be able to transfer them to any unfamiliar case the examiner sets.

METHOD Identifying the affected organ system from a set of symptoms

When a question gives a list of signs and symptoms and asks you to name the disorder or the system affected:

1. List each symptom and translate it into plain physiology (e.g. “excessive thirst” = water loss or high blood solute concentration).
2. Group the symptoms: do they point to one organ (kidney, pancreas, heart, joint) or to a whole system?
3. Match the group to a disorder you have studied, checking that every symptom is explained by your answer.
4. Justify in your answer by quoting the key diagnostic symptom(s), not by listing everything.

Reflex: the examiner awards marks for the link between symptom and organ function, not for the name of the disease alone.

WORKED EXAMPLE 5 Matching symptoms to a system

A 52-year-old trader in Bamenda complains of persistent lower back pain, swelling of the ankles, reduced urine volume, and fatigue. Her urine is frothy. (a) State the organ system most likely affected. (b) Explain how each symptom supports your answer. (c) Suggest one further test to confirm the diagnosis.

SOLUTION

(a) The **renal (urinary) system** is affected; the picture suggests a kidney disorder such as chronic nephritis or chronic kidney disease.

(b) Symptom-by-symptom justification:

- Reduced urine volume: damaged nephrons have a reduced filtration capacity, so less filtrate and therefore less urine is produced.

- Frothy urine: the glomerular filter is leaky and plasma proteins (albumin) pass into the urine (proteinuria); protein makes urine foam, like beaten egg white.
- Swollen ankles (oedema): loss of plasma protein lowers the osmotic (oncotic) pressure of the blood, so tissue fluid is not drawn back into the capillaries and accumulates in the tissues; gravity makes it collect at the ankles.
- Fatigue: failing kidneys produce less erythropoietin, reducing red blood cell formation (anaemia), and toxic wastes such as urea accumulate in the blood (uraemia); both cause tiredness.
- Lower back pain: the kidneys lie against the muscles of the lower back (loin region), so kidney inflammation is felt there.

(c) A urinalysis to confirm protein (and possibly blood) in the urine, or a blood test showing raised urea and creatinine levels, would confirm reduced kidney function.

Examiner's note: full marks require each symptom to be tied to a mechanism, as above — a bare list scores poorly.

METHOD Interpreting urinalysis data

To analyse a urine test table:

1. Recall the normal values: urine normally contains no glucose, no protein, no blood cells; it does contain water, urea and salts.
2. Compare each patient's result with the normal values and flag abnormalities.
3. Interpret each abnormality: glucose → diabetes mellitus (insulin problem; blood glucose above the kidney's reabsorption threshold); protein or blood → damaged glomeruli (nephritis); very large volumes of dilute urine → failure of water reabsorption (ADH defect, i.e. diabetes insipidus).
4. State the likely disorder and the defective structure (β -cells of the islets of Langerhans, glomerulus, tubule/collecting duct, or the hypothalamus–posterior pituitary ADH axis).

WORKED EXAMPLE 6 Analysing urine samples from three patients

The table shows the results of urine tests on three patients at a district hospital in Buea.

Substance	Patient A	Patient B	Patient C
Glucose	Present	Absent	Absent
Protein	Absent	Present	Absent
Blood cells	Absent	Present	Absent
Volume per day	Very large	Normal	Very large, very dilute

For each patient, state the likely disorder and the defective structure or hormone.

SOLUTION

Patient A — diabetes mellitus. Glucose in the urine with normal protein and no blood means the glomerular filter is intact. The blood glucose concentration has exceeded the kidney tubules' maximum reabsorption capacity (the renal threshold), so glucose spills into the urine. The defect is a failure of the β -cells of the islets of Langerhans to secrete enough insulin (or of body cells to respond to it). The large urine volume follows because unreabsorbed glucose holds water in the tubules by osmosis (osmotic diuresis) — explaining the classic thirst and copious urination.

Patient B — nephritis (glomerulonephritis). Protein and blood cells in urine indicate a damaged glomerular basement membrane: normally these molecules and cells are far too large to be filtered. Their presence shows the filter has become leaky, usually through inflammation, often following a bacterial infection.

Patient C — diabetes insipidus. A very large volume of very dilute urine, with no glucose or protein, points to a deficiency of antidiuretic hormone (ADH) from the posterior pituitary (or failure of the collecting ducts to respond to it). Without ADH the collecting duct walls remain impermeable to water, water is not reabsorbed, and huge volumes of dilute urine are excreted, causing intense thirst.

Key discriminator: glucose in urine → think pancreas/insulin; protein or blood in urine → think glomerulus; dilute urine in large volume with no solute abnormality → think ADH.

METHOD

Interpreting blood pressure readings and cardiovascular risk

1. Know the format: a reading is written systolic/diastolic in mmHg, e.g. 120/80 mmHg. Systolic = pressure during ventricular contraction; diastolic = pressure during ventricular relaxation.
2. Classify: roughly, normal is about 120/80 mmHg; persistent readings at or above 140/90 mmHg indicate **hypertension**.
3. Explain consequences: sustained high pressure damages artery walls, promotes atheroma (atherosclerosis), and overloads the heart → risk of stroke, heart failure, kidney damage.

4. Link to risk factors: high salt diet, obesity, smoking, stress, lack of exercise, age, family history.
5. Recommend: reduce salt and fat, exercise regularly, stop smoking, manage stress, take prescribed antihypertensive drugs.

WORKED EXAMPLE 7 Blood pressure data over one month

A 60-year-old man has his blood pressure measured weekly. Readings (mmHg): 155/98, 148/95, 160/102, 152/96. (a) State what the two numbers in each reading represent. (b) Comment on the data and name the condition. (c) Explain two possible long-term consequences if untreated. (d) Give three lifestyle changes you would advise.

SOLUTION

(a) The first (higher) number is the systolic pressure — the pressure in the arteries when the ventricles contract and eject blood. The second (lower) number is the diastolic pressure — the pressure when the ventricles relax and the heart refills.

(b) Every reading has systolic pressure above 140 mmHg and diastolic pressure above 90 mmHg. Because the elevation is persistent (four separate occasions), this is **hypertension** (high blood pressure), not a temporary rise due to stress or activity.

(c) Long-term consequences (any two, explained):

- Atherosclerosis, heart attack and stroke: high pressure damages the endothelium (inner lining) of arteries; cholesterol is deposited at the damage sites, forming atheroma plaques. A plaque or clot blocking a coronary artery causes a heart attack (myocardial infarction); blocking or bursting a brain artery causes a stroke.
- Heart failure: the left ventricle must pump against chronically high pressure; its muscle thickens, then weakens, and can no longer pump effectively.
- Kidney damage: high pressure damages the delicate glomerular capillaries, progressively destroying nephrons (a cause of chronic kidney disease).

(d) Lifestyle advice (any three): reduce salt intake (less salt in cooking, avoid stock cubes in excess); take regular exercise such as brisk walking; stop smoking and reduce alcohol; reduce fatty/fried foods and maintain a healthy body weight; learn to manage stress.


Examiner's note: "persistent" is the key word — one high reading is not a diagnosis.

METHOD Reasoning with hormone feedback loops (endocrine disorders)

1. Draw the loop for the hormone concerned: gland → hormone → target organ → effect → negative feedback on the gland (or on the pituitary/hypothalamus that controls it).

2. Locate the fault described in the question: is the gland over-secreting, under-secreting, or is the target tissue unresponsive?
3. Predict the consequences by following the loop: if the gland fails, the hormone falls and its effects are lost; if the gland is overactive, effects are exaggerated and feedback cannot switch it off.
4. Explain symptoms as direct results of too much or too little hormone action.

Key examples to master: insulin (diabetes mellitus), thyroxine (goitre, hyper-/hypothyroidism), ADH (diabetes insipidus).



Negative feedback control of thyroxine secretion. If iodine is lacking, thyroxine falls, the dashed inhibition is removed, and excess TSH makes the thyroid swell (goitre).

WORKED EXAMPLE 8 Goitre and the thyroxine feedback loop

In some highland communities, iodine in the diet is scarce. (a) Name the hormone whose production requires iodine and the gland that produces it. (b) Using the idea of negative feedback, explain why iodine deficiency causes the gland to swell (goitre). (c) State two symptoms you would expect from the resulting hormone deficiency.

SOLUTION

(a) The hormone is thyroxine, produced by the thyroid gland in the neck. Iodine is an essential component of the thyroxine molecule.

(b) Feedback explanation, step by step:

1. With little iodine available, the thyroid cannot make enough thyroxine, so the blood thyroxine level falls.
2. Normally, thyroxine exerts negative feedback on the pituitary, limiting its secretion of thyroid-stimulating hormone (TSH).

3. With thyroxine low, this feedback is removed, so the pituitary secretes excess TSH.
4. TSH continuously stimulates the thyroid; the gland grows larger as its cells multiply and enlarge in a vain attempt to produce more thyroxine — this swelling of the neck is a **goitre**.

(c) Symptoms of thyroxine deficiency (hypothyroidism): tiredness and a low metabolic rate (feeling cold, weight gain), and slowed physical and mental activity; in children, deficiency impairs growth and mental development (cretinism).

Prevention link: the use of iodised table salt — a cheap public-health measure promoted in Cameroon — prevents iodine-deficiency goitre.

METHOD Distinguishing disorders of the reproductive systems and advising on management

1. Classify the disorder by cause: hormonal (e.g. infertility from low sperm count or failure of ovulation), structural (e.g. blocked Fallopian tubes, enlarged prostate), or cancerous (e.g. cancer of the cervix, breast, prostate).
2. Link cause to effect: blocked tubes prevent fertilisation; low testosterone reduces sperm production; an enlarged prostate compresses the urethra, causing difficulty in urination.
3. State the key screening method where relevant: Pap smear (cervical cancer), breast self-examination and mammography (breast cancer), PSA blood test and rectal examination (prostate).
4. Advise early detection, medical consultation, and a healthy lifestyle; stress that many reproductive cancers are treatable if caught early.

WORKED EXAMPLE 9 Infertility and cancer screening

- (a) A couple has failed to conceive after three years of regular unprotected intercourse. Give two possible causes in the man and two in the woman, explaining each briefly.
- (b) Explain why regular screening is emphasised for cancers of the breast, cervix and prostate.

SOLUTION

- (a) Possible causes of infertility:
In the man:

- Low sperm count / abnormal sperm: poor sperm production in the seminiferous tubules (due to hormonal imbalance, undescended testes, excessive heat around the

scrotum, smoking or infection) means too few functional sperm reach the egg.

- Blocked vas deferens: sperm are produced but cannot pass into the semen, often as a result of infection.

In the woman:

- Failure of ovulation: hormonal imbalance (FSH/LH) means no ovum is released, so fertilisation cannot occur.
- Blocked Fallopian tubes: often a result of pelvic inflammatory disease following an untreated sexually transmitted infection; the ovum and sperm cannot meet, and scarred tubes also raise the risk of ectopic pregnancy.

(b) Why screening matters: cancers of the breast, cervix and prostate often develop silently, with few symptoms in the early stages, yet they are among the commonest cancers in adults. Screening (breast self-examination and mammography; Pap smear for the cervix; PSA test and examination for the prostate) detects abnormal cells before the cancer spreads. Early-stage cancers can be treated successfully by surgery, radiotherapy or chemotherapy; late-stage cancers that have metastasised (spread to other organs) are far harder to cure. Hence “early detection saves lives” is the central public-health message.

METHOD Analysing nervous system disorders from described deficits

1. Identify the level of the problem: brain (stroke, epilepsy, Alzheimer’s disease), spinal cord (paralysis below the injury), or peripheral nerves (neuritis).
2. Use the deficit to locate the lesion: paralysis of one side of the body after a burst or blocked brain artery → stroke affecting the opposite cerebral hemisphere; progressive loss of memory and reasoning in an elderly person → degenerative brain disease such as Alzheimer’s; sudden recurrent seizures → epilepsy (abnormal electrical discharge in the brain).
3. Explain the mechanism: neurones cannot regenerate in the CNS, so damage is usually permanent; severity depends on the region and extent damaged.
4. Conclude with management: rehabilitation/physiotherapy, anticonvulsant drugs for epilepsy, control of hypertension to prevent stroke.

WORKED EXAMPLE 10 A stroke case study

A 65-year-old hypertensive man suddenly loses movement and sensation on the right side of his body and his speech becomes slurred. (a) Name the likely event and the organ

affected. (b) Explain why the right side of the body is affected. (c) Explain why recovery is often incomplete. (d) State two measures that could have reduced his risk.

SOLUTION

(a) This is a **stroke** (cerebrovascular accident): a region of the brain has been deprived of blood, either because an artery was blocked by a clot or because a weakened artery burst (haemorrhage). His hypertension is the major underlying cause.

(b) The motor and sensory pathways cross over to the opposite side on their way between the brain and the body, so the left cerebral hemisphere controls and receives sensation from the right side of the body. Damage to the left hemisphere therefore paralyses the right side; the speech centres are also located in the left hemisphere in most people, explaining the slurred speech.

(c) Neurones of the central nervous system have virtually no power of regeneration. Once brain cells die from lack of oxygen, the lost functions can only be partly recovered as neighbouring areas are retrained through physiotherapy and speech therapy — hence recovery is usually slow and incomplete.

(d) Prevention: control blood pressure with medication and regular checks; reduce salt, stop smoking, exercise, and maintain a healthy weight — all of which reduce damage to cerebral arteries.

METHOD Interpreting joint and bone problems (musculoskeletal disorders)

1. Separate bone from joint problems: bones → osteoporosis (loss of bone mass, fractures), rickets/osteomalacia (poor mineralisation from vitamin D or calcium deficiency); joints → arthritis (inflammation: osteoarthritis from wear of cartilage, rheumatoid arthritis from autoimmune attack), sprains and dislocations from injury.
2. Match the patient profile: elderly woman with a hip fracture after a minor fall → osteoporosis; elderly person with painful, stiff knees worsening with use → osteoarthritis; child with bowed legs and a poor diet → rickets.
3. Explain the mechanism at tissue level (cartilage worn away so bones rub; demineralised bone becomes porous and weak).
4. Advise: calcium- and vitamin D-rich diet, weight-bearing exercise, weight control, physiotherapy, and anti-inflammatory treatment as prescribed.

WORKED EXAMPLE 11 Two elderly patients with joint pain

Patient X, aged 70, has painful, stiff knee joints that worsen after walking and improve with rest; X-ray shows narrowed joint space. Patient Y, aged 68, broke her hip after a

minor fall; bone scans show reduced bone density. (a) Name the most likely disorder in each case. (b) Explain the underlying cause of each. (c) Suggest one preventive measure for each condition.

SOLUTION

(a) Patient X: **osteoarthritis**. Patient Y: **osteoporosis**.

(b) Underlying causes:

- Patient X: the cartilage covering the ends of the bones in the knee joint has been worn away by years of use. Without this smooth, cushioning layer, the bone ends rub directly against each other, causing pain, stiffness and the reduced joint space seen on X-ray. Pain worsens with use and eases with rest — the classic pattern.
- Patient Y: bone is living tissue constantly being broken down and rebuilt. In old age — especially in women after the menopause, when protective oestrogen falls — breakdown exceeds rebuilding, so bone loses calcium salts and mass, becoming porous and fragile. Even a minor fall then causes a fracture, typically of the hip, wrist or spine.

(c) Prevention:

- Osteoarthritis: maintain a healthy body weight to reduce the load on the knees; avoid repetitive joint strain; gentle regular exercise keeps joints mobile.
- Osteoporosis: a diet rich in calcium and vitamin D (milk, small fish eaten with the bones, green vegetables, sunlight exposure) plus regular weight-bearing exercise throughout life to build and maintain bone mass.

Common trap: do not confuse osteoarthritis (a joint/cartilage disease) with osteoporosis (a bone-mass disease) — the GCE examiner tests this distinction directly.

METHOD Building an integrated NCD prevention plan

For an essay or structured question on preventing non-infectious diseases (NCDs):

1. Define NCDs: diseases not caused by pathogens and not transmissible (cardiovascular diseases, diabetes, cancers, chronic respiratory and renal diseases, musculoskeletal disorders).
2. Identify shared risk factors: tobacco, harmful alcohol use, unhealthy diet (excess salt, sugar, fat), physical inactivity, obesity, plus non-modifiable factors (age, sex, heredity).
3. Organise prevention at three levels: individual (lifestyle choices), community (health education, screening campaigns, smoke-free spaces), and health system (affordable screening, treatment, follow-up).
4. Give concrete examples: iodised salt against goitre, blood pressure checks in churches and markets, vaccination against hepatitis B and HPV to prevent liver and

cervical cancers, school sports against obesity.

5. Conclude that most NCDs are largely preventable because their main risk factors are behavioural.

WORKED EXAMPLE 12 Essay: reducing the burden of NCDs in your community

“Non-infectious diseases are an increasing cause of death among Cameroonian adults.” Discuss the statement by (a) naming four NCDs and the organ system each affects, (b) explaining three shared, modifiable risk factors, and (c) proposing a realistic prevention plan for your community.

SOLUTION

(a) Four NCDs and their systems (examples):

- Hypertension / stroke — cardiovascular and nervous systems.
- Diabetes mellitus — endocrine system (pancreatic insulin defect) with renal and cardiovascular complications.
- Cancer of the cervix or prostate — reproductive systems.
- Chronic kidney disease — renal system.

(b) Three shared modifiable risk factors, explained:

1. Tobacco smoking: damages artery walls (promoting atherosclerosis, hypertension, stroke) and introduces carcinogens (lung and other cancers).
2. Unhealthy diet: excess salt raises blood pressure; excess sugar and fat promote obesity and type 2 diabetes; fatty diets accelerate atheroma formation.
3. Physical inactivity: promotes obesity, raises blood pressure and blood glucose, and weakens the heart, bones and muscles — increasing risk across nearly all NCDs.

(c) Community prevention plan:

- Health education: talks in schools, churches, mosques and markets on salt reduction, a balanced diet (local vegetables, fruits, less fried food), and the dangers of tobacco and excessive alcohol.
- Screening campaigns: regular free blood-pressure and blood-sugar measurement days; Pap smear and breast examination drives for women; PSA testing for men over 50 — because early detection greatly improves survival.
- Environment and policy: encourage walking, community sports grounds, smoke-free public places, and promotion of iodised salt.
- Health system: ensure affordable drugs for hypertension and diabetes and proper follow-up of diagnosed patients, since NCDs require lifelong management.

Conclusion: NCDs share a small number of behavioural risk factors, so coordinated lifestyle change plus early screening can prevent a large proportion of cases — making prevention far cheaper than treatment.

WARNING Frequent errors to avoid in this chapter

- Confusing diabetes mellitus (glucose in urine, insulin defect) with diabetes insipidus (dilute urine, ADH defect).
- Writing that protein in urine indicates diabetes — it indicates glomerular damage (nephritis).
- Calling any single high blood pressure reading “hypertension” — it must be persistent.
- Mixing up osteoarthritis (worn joint cartilage) with osteoporosis (porous, fragile bone).
- Describing NCDs as “caught” from other people — by definition they are not transmissible.
- Naming a disease without linking each symptom to the failing organ’s function — this is where marks are lost.

7 EXERCISES

7.1 I check my knowledge

EXERCISE 6 Multiple choice questions

1. Which of the following is a non-infectious disease of the male reproductive system?
 - A Prostatitis caused by *Escherichia coli*
 - B Benign prostatic hyperplasia (BPH)
 - C Gonorrhoea
 - D Syphilis
2. A patient presents with polyuria, polydipsia and hyperglycaemia but no ketones in urine. The most likely diagnosis is:
 - A Diabetes insipidus
 - B Type 1 diabetes mellitus
 - C Type 2 diabetes mellitus
 - D Cushing's syndrome
3. The glomerular filtration rate (GFR) is best estimated by the clearance of:
 - A Urea
 - B Creatinine
 - C Glucose
 - D Sodium
4. Which joint is characteristically spared in rheumatoid arthritis but commonly affected in osteoarthritis?
 - A Metacarpophalangeal joints
 - B Proximal interphalangeal joints
 - C Distal interphalangeal joints
 - D Wrist joints
5. Atherosclerosis primarily affects which layer of the arterial wall?
 - A Tunica adventitia
 - B Tunica media
 - C Tunica intima

D External elastic lamina

EXERCISE 7 True/False with justification

State whether each statement is true or false and justify your answer briefly.

1. Hypertension is defined as a sustained systolic blood pressure ≥ 140 mmHg and/or diastolic ≥ 90 mmHg.
2. In type 1 diabetes mellitus, insulin resistance is the primary defect.
3. Glomerulonephritis always presents with nephrotic syndrome (massive proteinuria, hypoalbuminaemia, oedema, hyperlipidaemia).
4. Osteoarthritis is an autoimmune disease characterised by symmetrical small-joint involvement.
5. Benign prostatic hyperplasia (BPH) increases the risk of prostate cancer.

EXERCISE 8 Definitions and key terms

Define the following terms precisely:

1. **Non-communicable disease (NCD)**
2. **Glomerular filtration rate (GFR)**
3. **Insulin resistance**
4. **Atheroma (atherosclerotic plaque)**
5. **Rheumatoid factor**
6. **Benign prostatic hyperplasia (BPH)**
7. **Goitre**
8. **Myocardial infarction**

EXERCISE 9 Direct calculations

1. A 45-year-old man has a serum creatinine of $120 \mu\text{mol/L}$. Using the simplified MDRD

equation (for adults):

$$\text{eGFR (mL/min/1.73 m}^2\text{)} = 186 \times \left(\frac{\text{creatinine in } \mu\text{mol/L}}{88.4} \right)^{-1.154} \times \text{age}^{-0.203} \times (0.742 \text{ if female})$$

Calculate his estimated GFR. Is this consistent with chronic kidney disease (CKD) stage 3 (GFR 30–59 mL/min/1.73 m²)?

2. A woman's blood pressure readings on three separate occasions are: 148/92, 152/96, 146/94 mmHg. Classify her hypertension according to WHO/ISH guidelines (Grade 1, 2 or 3).
3. Calculate the Body Mass Index (BMI) of a patient weighing 92 kg and measuring 1.68 m. Classify the nutritional status according to WHO categories.
4. In a glucose tolerance test, a patient's 2-hour plasma glucose is 11.5 mmol/L. Fasting glucose is 6.8 mmol/L. Interpret these results using WHO criteria (fasting ≥ 7.0 mmol/L or 2-h ≥ 11.1 mmol/L for diabetes; fasting 6.1–6.9 mmol/L for impaired fasting glycaemia; 2-h 7.8–11.0 mmol/L for impaired glucose tolerance).

7.2 I practise


EXERCISE 10 Kidney disease: proteinuria, haematuria and oedema

A 12-year-old boy presents with puffy eyes, swollen ankles and cola-coloured urine two weeks after a sore throat. Urinalysis shows +++ protein and numerous red blood cells.

1. Name the specific part of the nephron most likely damaged.
2. Name the probable disease.
3. Explain the pathophysiological mechanism leading to oedema in this condition.
4. List two other clinical features you would expect to find.
5. Outline the principle of management.

EXERCISE 11 Glucose tolerance test interpretation

The graph below shows plasma glucose concentrations (mmol/L) over 2 hours after a 75 g oral glucose load for three individuals A, B and C.



Oral glucose tolerance test curves for three individuals.

1. Identify which curve corresponds to: (i) a healthy individual, (ii) type 2 diabetes mellitus, (iii) type 1 diabetes mellitus. Justify each choice using WHO diagnostic criteria (fasting and 2-hour values).
2. Explain why the curve for type 1 diabetes typically shows a higher peak and slower return to baseline than type 2.
3. State two other laboratory tests that help distinguish type 1 from type 2 diabetes.

EXERCISE 12 Comparison: osteoarthritis vs rheumatoid arthritis

Complete the table below contrasting osteoarthritis (OA) and rheumatoid arthritis (RA).

Feature	Osteoarthritis (OA)	Rheumatoid arthritis (RA)
Primary cause / aetiology		
Typical age of onset		
Pattern of joint involvement (symmetry, joints affected)		
Morning stiffness duration		
Systemic features (e.g. fever, weight loss, nodules)		
Radiographic findings (key differences)		
Autoantibodies		
First-line pharmacological management		

EXERCISE 13 Case study: endemic goitre

A 35-year-old woman living in a remote highland village in the Northwest Region of Cameroon presents with a diffuse, non-tender swelling in the anterior neck. She reports fatigue, cold intolerance and weight gain. The village diet consists mainly of cassava and maize; iodised salt is unavailable.

1. Name the condition and its primary cause.
2. Explain the hormonal feedback failure involved, mentioning TRH, TSH, T3 and T4.
3. Why does the thyroid enlarge (goitre) in this condition?
4. Propose a national prevention measure that has proven effective in Cameroon.
5. List two other consequences of iodine deficiency in pregnancy and early childhood.

7.3 I prepare for the GCE**EXERCISE 14 Essay: NCDs as a public health threat in Cameroon] [25 marks**

“Non-infectious diseases are now a greater public-health threat in Cameroon than infectious diseases.” Discuss this statement with reference to at least four body systems. In your answer, include:

- Epidemiological evidence (trends, risk factors).
- Major NCDs affecting each system, their pathophysiology and complications.
- Socio-economic impact on individuals, families and the health system.
- Prevention strategies at primary, secondary and tertiary levels, with Cameroonian examples.

EXERCISE 15 Short answers: Hypertension] [15 marks

1. Define hypertension according to current WHO/ISH guidelines. [2 marks]
2. State two consequences of untreated hypertension on the kidney and two on the brain. [4 marks]
3. Explain why hypertension is called a “silent killer”. [3 marks]
4. Describe the mechanism of action of one class of antihypertensive drugs (e.g. ACE inhibitors, calcium channel blockers, thiazide diuretics, beta-blockers). [3 marks]

5. A 55-year-old trader has BP 160/100 mmHg on two occasions. Outline the initial investigations you would request and lifestyle advice you would give. [3 marks]

EXERCISE 16 Practical: Blood pressure measurement and interpretation] [15 marks

1. Describe, step by step, the correct procedure for measuring a classmate's blood pressure using a manual sphygmomanometer and stethoscope. Include patient preparation, cuff placement, inflation, deflation and reading. [6 marks]
2. A reading of 150/95 mmHg is obtained. Interpret this value (classification, urgency). [3 marks]
3. List four common errors that can falsely elevate or lower the reading. [2 marks]
4. Advise the person on immediate next steps (lifestyle, follow-up, when to seek medical care). [4 marks]

8 SOLUTIONS TO THE EXERCISES

9 CORRECTIONS AND WORKED SOLUTIONS — NON-INFECTIOUS DISEASES (BIO06)

SOLUTION Exercise 1 — Multiple choice questions

1. B — Benign prostatic hyperplasia (BPH). BPH is a non-infectious, age-related proliferation of prostatic stromal and glandular tissue driven by dihydrotestosterone. Options A, C and D are all infectious: prostatitis caused by *E. coli* is bacterial, gonorrhoea is caused by *Neisseria gonorrhoeae* and syphilis by *Treponema pallidum*.
2. C — Type 2 diabetes mellitus. The triad polyuria–polydipsia–hyperglycaemia indicates diabetes mellitus. The absence of ketones points to type 2: residual insulin secretion is sufficient to suppress lipolysis and ketogenesis. Type 1 (B) typically presents with ketoacidosis. Diabetes insipidus (A) causes polyuria and polydipsia but not hyperglycaemia. Cushing's syndrome (D) may cause hyperglycaemia but is not the most likely diagnosis here.
3. B — Creatinine. Creatinine is freely filtered at the glomerulus and is neither significantly reabsorbed nor secreted (only minimal secretion), so its clearance approximates GFR. Urea is partly reabsorbed (underestimates GFR); glucose is fully reabsorbed in health; sodium is extensively reabsorbed.
4. C — Distal interphalangeal joints. The DIP joints (Heberden's nodes) are characteristically affected in osteoarthritis but spared in rheumatoid arthritis, which targets MCP, PIP and wrist joints symmetrically.
5. C — Tunica intima. Atherosclerosis begins with endothelial dysfunction and accumulation of LDL-derived lipids in the **tunica intima**, forming fatty streaks then fibrous plaques. The media and adventitia are affected only secondarily.

SOLUTION Exercise 2 — True/False with justification

1. True. WHO/ISH defines hypertension as sustained systolic BP ≥ 140 mmHg and/or diastolic BP ≥ 90 mmHg, confirmed on at least two separate occasions.
2. False. In type 1 diabetes the primary defect is autoimmune destruction of pancreatic β -cells causing absolute insulin deficiency. Insulin resistance is the hallmark of type 2 diabetes.
3. False. Glomerulonephritis has several presentations: nephritic syndrome (haematuria, mild proteinuria, hypertension, oliguria), nephrotic syndrome, or rapidly pro-

gressive renal failure. Nephrotic syndrome occurs only in some forms (e.g. membranous nephropathy, minimal change disease).

4. False. Osteoarthritis is a degenerative (“wear-and-tear”) disease of articular cartilage, typically asymmetrical and involving weight-bearing joints and DIP joints. Symmetrical small-joint autoimmune disease describes rheumatoid arthritis.
5. False. BPH arises from the transition zone of the prostate and is benign; it does not transform into or increase the risk of prostate cancer (which usually arises in the peripheral zone). The two conditions share age as a risk factor and may coexist, which explains the confusion.

SOLUTION Exercise 3 — Definitions and key terms

1. Non-communicable disease (NCD): a chronic disease that is not caused by an infectious agent and cannot be transmitted from person to person; it results from genetic, physiological, environmental and behavioural factors (e.g. hypertension, diabetes, cancers, chronic respiratory diseases).
2. Glomerular filtration rate (GFR): the volume of plasma filtered through the glomeruli into Bowman’s capsule per unit time (normally about 120 mL/min/1.73 m² in young adults); it is the best overall index of kidney function, estimated clinically from creatinine clearance or equations such as MDRD or CKD-EPI.
3. Insulin resistance: a reduced biological response of target tissues (muscle, liver, adipose tissue) to normal circulating insulin concentrations, so that glucose uptake and utilisation are impaired and compensatory hyperinsulinaemia occurs.
4. Atheroma: a focal, raised lesion of the arterial intima composed of a lipid core (cholesterol esters, foam cells) covered by a fibrous cap of smooth-muscle cells and connective tissue; it narrows the lumen and may rupture, triggering thrombosis.
5. Rheumatoid factor (RF): an autoantibody (usually IgM) directed against the Fc portion of IgG, found in about 70–80% of patients with rheumatoid arthritis (also in some infections and healthy elderly people, so it is not fully specific).
6. Benign prostatic hyperplasia: a non-malignant, age- and androgen-dependent nodular enlargement of the transition zone of the prostate, causing bladder-outlet obstruction with hesitancy, weak stream, nocturia and urinary retention.
7. Goitre: any enlargement of the thyroid gland; in endemic goitre it results from prolonged TSH stimulation when iodine deficiency limits T3/T4 synthesis.
8. Myocardial infarction: irreversible necrosis of cardiac myocytes caused by prolonged ischaemia, usually from occlusion of a coronary artery by thrombus on a rup-

tured atherosclerotic plaque; diagnosed by typical chest pain, ECG changes and raised cardiac biomarkers (troponin).

SOLUTION Exercise 4 — Direct calculations

1. eGFR (MDRD). The patient is male, so the 0.742 factor is omitted. Serum creatinine = 120 $\mu\text{mol/L}$ ($= 120/88.4 = 1.357 \text{ mg/dL}$). Age = 45 years.

$$\begin{aligned} \text{eGFR} &= 186 \times (\text{Scr})^{-1.154} \times (\text{Age})^{-0.203} \\ &= 186 \times 1.357^{-1.154} \times 45^{-0.203} \\ 1.357^{-1.154} &= e^{-1.154 \ln 1.357} = e^{-1.154 \times 0.3056} = e^{-0.3527} \approx 0.703 \\ 45^{-0.203} &= e^{-0.203 \ln 45} = e^{-0.203 \times 3.807} = e^{-0.7728} \approx 0.462 \\ \text{eGFR} &= 186 \times 0.703 \times 0.462 \approx 60.4 \text{ mL/min/1.73 m}^2 \end{aligned}$$

The eGFR $\approx 60 \text{ mL/min/1.73 m}^2$ lies just above the 30–59 band, so it is not consistent with CKD stage 3; it corresponds to stage 2 (60–89) if chronic kidney damage markers are present, and is borderline normal otherwise. (Accept values in the 58–62 range depending on rounding; the conclusion about stage 3 is unchanged if the value falls below 60 — the key skill is correct substitution and comparison with the reference band.)

2. Blood pressure classification. Mean values: systolic $\approx (148 + 152 + 146)/3 \approx 149 \text{ mmHg}$; diastolic $\approx (92 + 96 + 94)/3 = 94 \text{ mmHg}$. All readings fall in the ranges SBP 140–159 and/or DBP 90–99 mmHg $\Rightarrow$ Grade 1 (mild) hypertension. The diagnosis is valid because elevated values were recorded on three separate occasions.
3. BMI.

$$\text{BMI} = \frac{m}{h^2} = \frac{92}{1.68^2} = \frac{92}{2.8224} \approx 32.6 \text{ kg/m}^2$$

WHO classification: 30.0–34.9 $\Rightarrow$ obesity class I (moderate obesity).

4. Glucose tolerance test. Fasting glucose 6.8 mmol/L lies in the 6.1–6.9 band $\Rightarrow$ impaired fasting glycaemia; but the 2-hour value 11.5 mmol/L $\geq 11.1 \text{ mmol/L} \Rightarrow$ diabetes mellitus. When categories conflict, the more severe category prevails: the patient has diabetes mellitus (confirmed by a repeat test on another day if asymptomatic).

SOLUTION Exercise 5 — Kidney disease: proteinuria, haematuria and oedema

1. Structure damaged: the **glomerulus**, specifically the glomerular basement membrane and its filtration barrier (capillary endothelium, basement membrane, podocytes).

2. Probable disease: acute post-streptococcal glomerulonephritis (nephritic syndrome), occurring 1–3 weeks after a group A β -haemolytic streptococcal pharyngitis.
3. Mechanism of oedema: immune complexes deposit in the glomerular basement membrane $\Rightarrow$ inflammation increases permeability and reduces the filtration surface $\Rightarrow$ GFR falls, so Na^+ and water are retained (increased tubular reabsorption, activation of the renin–angiotensin–aldosterone system) $\Rightarrow$ plasma volume expands and hydrostatic pressure rises $\Rightarrow$ fluid shifts into the interstitium. Loss of protein in urine additionally lowers plasma oncotic pressure, worsening the oedema (periorbital in the morning, ankles later).
4. Other expected features (any two): hypertension; oliguria; mild fever/malaise; loin discomfort; elevated serum creatinine; low complement C3.
5. Principle of management: mainly supportive — rest, salt and water restriction, treat hypertension and fluid overload (diuretics), eradicate residual streptococcal infection with penicillin, monitor renal function; most children recover fully.

SOLUTION Exercise 6 — Glucose tolerance test interpretation

1. Identification:
 - Individual A — healthy: fasting 5.2 mmol/L (< 6.1) and 2-h 5.8 mmol/L (< 7.8): both normal; glucose returns to baseline within 2 h.
 - Individual B — type 1 diabetes: fasting 7.5 mmol/L (≥ 7.0) and 2-h 15.2 mmol/L (≥ 11.1): diabetic by both criteria, with a very high peak (18.5) and persistently marked hyperglycaemia — the pattern of absolute insulin deficiency.
 - Individual C — type 2 diabetes: fasting 6.0 mmol/L (normal/IFG borderline) but 2-h 11.5 mmol/L (≥ 11.1): diabetic on the 2-h criterion; the moderate peak and partial decline reflect residual but insufficient insulin secretion with insulin resistance.
2. Why type 1 peaks higher and falls slower: in type 1 there is no endogenous insulin, so ingested glucose cannot be taken up by muscle and adipose tissue and hepatic glucose output is not suppressed — glucose accumulates to a higher peak and is cleared only slowly (mainly by renal excretion once the tubular threshold ≈ 10 mmol/L is exceeded, causing glycosuria). In type 2, residual insulin secretion, though delayed and inadequate against insulin resistance, still promotes some peripheral uptake, so the peak is lower and the fall faster.
3. Two distinguishing tests: (i) autoantibodies — anti-GAD, anti-islet cell or anti-insulin antibodies (present in type 1); (ii) C-peptide / fasting insulin — low or absent in type 1, normal or high in type 2. (Also acceptable: presence of ketones/ketoacidosis, age and BMI at presentation.)

SOLUTION Exercise 7 — Comparison: osteoarthritis vs rheumatoid arthritis

Feature	Osteoarthritis (OA)	Rheumatoid arthritis (RA)
Primary cause / aetiology	Degenerative “wear-and-tear” loss of articular cartilage; age, obesity, joint overload	Autoimmune chronic synovitis (autoantibodies, immune complexes) destroying cartilage and bone
Typical age of onset	Usually > 50 years	Any age, peak 30–50 years; female predominance
Pattern of joint involvement	Asymmetrical; weight-bearing joints (hips, knees, spine) and DIP joints	Symmetrical small joints: MCP, PIP, wrists; spares DIP
Morning stiffness duration	Short (< 30 min)	Prolonged (> 1 h)
Systemic features	Absent	Present: fever, fatigue, weight loss, rheumatoid nodules, anaemia
Radiographic findings	Joint-space narrowing, osteophytes, subchondral sclerosis and cysts	Juxta-articular osteoporosis, marginal erosions, uniform space narrowing, deformities (ulnar deviation)
Autoantibodies	None	Rheumatoid factor, anti-CCP
First-line pharmacological management	Analgesics (paracetamol), NSAIDs, physiotherapy, weight loss	DMARDs (e.g. methotrexate) early; NSAIDs/corticosteroids for flares

SOLUTION Exercise 8 — Case study: endemic goitre

1. Condition: endemic (iodine-deficiency) goitre with hypothyroidism. Primary cause:

chronic dietary iodine deficiency, aggravated by goitrogens in cassava (cyanogenic glucosides that inhibit iodine uptake).

2. Feedback failure: iodine is required to synthesise the thyroid hormones T3 and T4. Iodine deficiency $\Rightarrow$ low T3/T4 $\Rightarrow$ loss of negative feedback on the hypothalamus and pituitary $\Rightarrow$ TRH and especially TSH secretion rise continuously. Chronically elevated TSH is the driver of the disease.
3. Why the gland enlarges: TSH is trophic: it stimulates thyroid follicular cells to grow and divide (hypertrophy and hyperplasia) and increases colloid accumulation in an attempt to trap more iodine and produce more hormone. Sustained stimulation therefore produces diffuse thyroid enlargement — the goitre.
4. National prevention measure: universal salt iodisation — Cameroon has implemented mandatory iodisation of table salt, which has markedly reduced endemic goitre; supplementation with iodised oil in high-risk areas is complementary.
5. Consequences in pregnancy/childhood (any two): miscarriage and stillbirth; congenital hypothyroidism with cretinism (severe, irreversible mental retardation and stunted growth); impaired cognitive development and reduced school performance; deaf-mutism in severe deficiency.

SOLUTION Exercise 9 — Essay: NCDs as a public health threat in Cameroon (25 marks)

Indicative mark scheme and examiner expectations.

Introduction (2 marks): define NCDs; present the epidemiological transition — while malaria, HIV and tuberculosis remain important, urbanisation, dietary change, tobacco/alcohol use, physical inactivity and ageing are driving a rapid rise of NCDs, creating a double burden of disease. A nuanced thesis (NCDs are a growing, comparable threat rather than simply “greater”) earns full credit.

Body — at least four systems ($4 \times 3 = 12$ marks; 1 mark for the disease + pathophysiology, 1 for complications, 1 for Cameroonian context):

- Cardiovascular: hypertension (very prevalent, often undiagnosed), stroke, myocardial infarction, heart failure; atherosclerosis mechanism; complications (renal failure, sudden death).
- Endocrine/metabolic: type 2 diabetes rising with obesity; complications — retinopathy, nephropathy, neuropathy, diabetic foot amputations.
- Renal: chronic kidney disease from hypertension, diabetes and glomerulonephritis; dialysis is costly and scarce.
- Musculoskeletal: osteoarthritis in an ageing population, road-traffic injuries; disability and loss of productivity.

- Reproductive/other: cancers (cervix, breast, prostate) — late presentation, limited radiotherapy.

Socio-economic impact (4 marks): catastrophic out-of-pocket spending, loss of breadwinners, caregiving burden, absenteeism, pressure on an under-resourced health system, treatment costs (insulin, dialysis, antihypertensives) pushing families into poverty.

Prevention (5 marks):

- Primary: health education, salt reduction, tobacco and alcohol control, promotion of physical activity, iodised salt, vaccination against hepatitis B and HPV (cancer prevention).
- Secondary: screening — BP and blood glucose campaigns in churches, markets and workplaces; cervical cancer screening; early detection programmes.
- Tertiary: affordable essential medicines, management of complications, rehabilitation, physiotherapy, palliative care.

Conclusion (2 marks): balanced judgement — NCDs now rival infectious diseases and require an integrated national response.

Examiner expectations: accurate terminology, correct pathophysiology, genuine Cameroonian examples (not invented statistics), logical structure, and coverage of all four requested elements. Vague lists without explanation score poorly.

SOLUTION Exercise 10 — Short answers: Hypertension (15 marks)

1. Definition [2]: a sustained elevation of arterial blood pressure ≥ 140 mmHg systolic and/or ≥ 90 mmHg diastolic, confirmed on at least two separate measurements/occasions (WHO/ISH). [1 mark for values, 1 for “sustained/confirmed”.]
2. Consequences [4]: Kidney (any 2): nephrosclerosis (hyaline arteriosclerosis), chronic kidney disease/proteinuria, end-stage renal failure. Brain (any 2): ischaemic or haemorrhagic stroke, hypertensive encephalopathy, vascular dementia, transient ischaemic attacks.
3. “Silent killer” [3]: hypertension usually causes no symptoms for years while progressively damaging arteries, heart, brain, kidneys and retina; the first manifestation may be a fatal event (stroke, myocardial infarction, sudden death); hence many Cameroonians are unaware of their condition until complications occur — only routine screening reveals it.
4. Mechanism of one drug class [3] — e.g. ACE inhibitors: they inhibit angiotensin-converting enzyme $\Rightarrow$ less angiotensin II $\Rightarrow$ reduced vasoconstriction and reduced aldosterone secretion $\Rightarrow$ decreased Na^+ /water retention and vasodilation $\Rightarrow$ lower blood pressure; they also reduce glomerular pressure, protecting the kidney. (Ac-

cept: thiazides — inhibit NaCl reabsorption in the distal convoluted tubule causing diuresis and reduced plasma volume; beta-blockers — block β_1 receptors reducing heart rate and contractility; calcium channel blockers — block L-type Ca^{2+} channels in vascular smooth muscle causing vasodilation.)


5. Investigations and advice [3]: Investigations: confirm BP, urinalysis (proteinuria), serum creatinine/electrolytes, fasting glucose and lipids, ECG, fundoscopy. Lifestyle: reduce salt (< 5 g/day), reduce alcohol, stop tobacco, weight reduction, regular physical activity, fruit/vegetable-rich diet, stress management, adherence to follow-up.

SOLUTION Exercise 11 — Practical: Blood pressure measurement (15 marks)

1. Procedure [6 marks — 1 per key step]:
 - (a) Preparation: the subject rests seated for 5 minutes, back supported, feet flat, no caffeine, tobacco or exercise in the previous 30 min; arm bare and supported at heart level.
 - (b) Cuff placement: choose the correct cuff size (bladder length encircling $\approx$ 80% of arm circumference); wrap snugly 2–3 cm above the elbow crease, with the artery marker over the brachial artery.
 - (c) Locate the pulse: palpate the brachial artery in the antecubital fossa; place the stethoscope diaphragm over it (not under the cuff).
 - (d) Inflation: close the valve and inflate rapidly to about 20–30 mmHg above the point where the radial pulse disappears (avoiding auscultatory gap errors).
 - (e) Deflation: open the valve and deflate slowly ($\approx$ 2–3 mmHg/s). The first clear tapping sound (Korotkoff I) = systolic pressure; the point where sounds disappear (Korotkoff V) = diastolic pressure.
 - (f) Reading and recording: read to the nearest 2 mmHg, record arm, position and time; repeat after 1–2 min and average; measure on both arms at first assessment.
2. Interpretation of 150/95 mmHg [3]: both values fall in Grade 1 (mild) hypertension (SBP 140–159, DBP 90–99). It is not an emergency; it requires confirmation on at least one or two further occasions (or ambulatory/home monitoring) before diagnosis and treatment decisions.
3. Four common errors [2]: cuff too small or too loose (falsely high); arm below heart level or unsupported (falsely high); deflation too fast (inaccurate readings); talking or recent caffeine/tobacco/exercise (falsely high); cuff over clothing; stethoscope pressed too hard (falsely low diastolic). [Any four, $\frac{1}{2}$ mark each.]

4. Next steps [4]: repeat measurements on separate days to confirm; begin lifestyle measures immediately (salt reduction, weight control, exercise, limit alcohol, stop smoking); check for target-organ damage (urinalysis, creatinine, glucose, ECG); seek medical care promptly for confirmed persistent hypertension, and urgently if BP $\geq$ 180/110 mmHg or symptoms such as severe headache, chest pain, visual disturbance or neurological signs occur.

10 SUMMARY SHEET



Mind map of chapter BIO06: the main families of non-infectious diseases, their examples and prevention.

KEY POINTS Key definitions

- **Non-infectious disease (NCD)**: a disease not caused by a pathogen and not transmissible from person to person; it results from genetic factors, lifestyle, ageing, hormonal imbalance or environmental factors.

- **Risk factor**: any behaviour or condition (smoking, harmful alcohol use, obesity, high salt intake, physical inactivity, heredity) that increases the probability of developing an NCD.
- **Benign tumour**: a localised, slow-growing cell mass that remains encapsulated and does not spread; **malignant tumour** (cancer): invades neighbouring tissues and forms **metastases** (secondary tumours) via the blood or lymph.
- **Infertility**: the inability of a couple to achieve pregnancy after about one year of regular unprotected intercourse.
- **Hypertension**: persistently raised arterial blood pressure, conventionally $\geq 140/90$ mmHg (systolic/diastolic) on repeated measurements.
- **Atherosclerosis**: deposition of fatty plaques (rich in cholesterol) in the inner wall of arteries, narrowing the lumen and reducing blood flow.
- **Renal failure**: inability of the kidneys to filter the blood and excrete nitrogenous wastes (urea); treated by **dialysis** or kidney transplant.

KEY POINTS Essential facts by system

- Reproductive: male — benign prostatic hyperplasia (enlarged prostate, difficulty urinating), prostate and testicular cancers, low sperm count; female — uterine fibroids, ovarian cysts, cervical cancer (associated with human papillomavirus, HPV; preventable by vaccination and screening), breast cancer (detected early by self-examination and mammography).
- Nervous: **stroke** (interrupted blood supply to part of the brain), **epilepsy** (recurrent seizures from abnormal electrical discharges in the brain), **Parkinson disease** (degeneration of dopamine-producing neurones; tremor, rigidity, slow movement), **Alzheimer disease** (progressive memory loss and dementia).
- Endocrine: **diabetes mellitus** — chronic hyperglycaemia; type 1 (autoimmune destruction of the β cells of the islets of Langerhans, no insulin, insulin-dependent, usually juvenile onset), type 2 (insulin resistance, associated with obesity and lifestyle, usually adult onset). **Goitre** and hypothyroidism from iodine deficiency; hyperthyroidism (excess thyroxine: weight loss, rapid heartbeat, exophthalmia).
- Renal: nephritis (inflammation of the nephrons), kidney stones (mineral crystals in the kidney or ureter), chronic renal failure; normal urine contains no glucose and no protein.
- Cardiovascular: hypertension, atherosclerosis, coronary thrombosis $\Rightarrow$ **myocardial infarction** (heart attack: death of heart muscle), heart failure, stroke.

- Musculoskeletal: **osteoarthritis** (wear of articular cartilage), **rheumatoid arthritis** (autoimmune inflammation of joints), **osteoporosis** (loss of bone mineral density; calcium/vitamin D deficiency; common in post-menopausal women), **rickets** (vitamin D deficiency in children; bowed legs), muscular dystrophy (inherited progressive muscle wasting), sprains and fractures.

METHOD Methods to master

1. Interpret urine/blood test results: glucose in urine $\Rightarrow$ suspect diabetes mellitus; protein in urine $\Rightarrow$ suspect kidney damage; high blood urea $\Rightarrow$ suspect renal failure.
2. Read a blood pressure value: systolic over diastolic (e.g. 120/80 mmHg); classify as normal, elevated or hypertensive.
3. Compare type 1 and type 2 diabetes in a table (cause, age of onset, treatment).
4. Draw a labelled flow diagram: risk factor $\rightarrow$ lesion $\rightarrow$ disease $\rightarrow$ complication $\rightarrow$ prevention.

WARNING Common mistakes

- Confusing infectious and non-infectious diseases: NCDs are never contagious.
- Saying diabetes is caused directly by “eating sugar”: it is a failure of insulin secretion (type 1) or of insulin action (type 2).
- Confusing **osteoporosis** (porous, fragile bones) with **osteoarthritis** (worn joint cartilage).
- Confusing **rickets** (children, vitamin D deficiency) with **osteomalacia** (the equivalent condition in adults).
- Believing hypertension always produces symptoms: it is a “silent killer” often detected only by measurement.
- Writing that all tumours are cancers: benign tumours do not invade tissues and do not metastasise.
- Listing meningitis as a nervous system NCD: meningitis is an infectious disease; only its after-effects (sequelae) are non-infectious.

KEY POINTS GCE examination tips

- Structured questions often ask: “State two causes, two symptoms and two preventive measures” — prepare such triplets for each disease.
- Use the Cameroonian context wisely: cervical and prostate cancer screening campaigns, iodised salt against goitre, hypertension and diabetes as rising NCDs, sickle cell disease as a genetic (non-infectious) disorder.
- Always link prevention to the four shared behavioural risk factors: tobacco use, harmful alcohol use, unhealthy diet and physical inactivity.
- In data questions, quote figures from the table or graph to justify your diagnosis (e.g. a fasting blood glucose value, a blood pressure reading).

**Signed • OnBuch+**ONBUCH+ TEACHING TEAM
FOLLOWS THE OFFICIAL MINESEC SYLLABUS**OnBuch+**