

FURTHER MATHEMATICS • UPPER SIXTH

UPPER SIXTH — FURTHER MATHEMATICS

CHAPTER 14

VECTOR PRODUCT AND ITS APPLICATIONS

SUGGESTED TIME: 3 PERIODS

IN THIS CHAPTER

This chapter introduces the vector (cross) product of two vectors in space, its algebraic and geometric properties, and its spectacular applications: areas, volumes via the scalar triple product, and the vector and Cartesian equations of a plane. It is a cornerstone chapter of the GCE Advanced Level Further Mathematics paper, where spatial geometry questions are guaranteed.

- By the end of this chapter I can define the vector product of two vectors and compute it both from the determinant formula and from the geometric definition.
- By the end of this chapter I can state and use the algebraic properties of the vector product (anticommutativity, distributivity, behaviour with scalar multiples).
- By the end of this chapter I can interpret the magnitude of a vector product as the area of a parallelogram or triangle.
- By the end of this chapter I can compute the scalar triple product and use it to find volumes of parallelepipeds and tetrahedra.
- By the end of this chapter I can use the scalar triple product to test whether three vectors are coplanar or four points are coplanar.
- By the end of this chapter I can determine a normal vector to a plane defined by three points or by two directions.

DOWNLOAD ON-
BUCH

Free • Google Play
AI tutor, past papers,
results



GO ONBUCH+

All signed courses, unlimited AI
tutor, PDF sheets — OnBuch+
tab in the app

THIS SIGNED COURSE CONTAINS

COURSE

complete, illustrated,
step by step

METHODS

and worked examples

EXERCISES

exam style + solutions

SUMMARY
SHEET

the essentials to
remember



Signed • OnBuch+

ONBUCH+ TEACHING TEAM
FOLLOWS THE OFFICIAL MINESEC SYLLABUS

OnBuch+

CONTENTS

- | | |
|---|--------------------------------|
| 1. Section 1: The Vector Product — Definition and Computation | tor and Cartesian Forms |
| 2. Section 2: Algebraic Properties and Geometric Applications (Areas) | 5. Integration activity |
| 3. Section 3: The Scalar Triple Product and Volumes | 6. Methods and worked examples |
| 4. Section 4: Equations of a Plane — Vector | 7. Exercises |
| | 8. Solutions to the exercises |
| | 9. Summary sheet |

CHAPTER OBJECTIVES

- By the end of this chapter I can define the vector product of two vectors and compute it both from the determinant formula and from the geometric definition.
- By the end of this chapter I can state and use the algebraic properties of the vector product (anticommutativity, distributivity, behaviour with scalar multiples).
- By the end of this chapter I can interpret the magnitude of a vector product as the area of a parallelogram or triangle.
- By the end of this chapter I can compute the scalar triple product and use it to find volumes of parallelepipeds and tetrahedra.
- By the end of this chapter I can use the scalar triple product to test whether three vectors are coplanar or four points are coplanar.
- By the end of this chapter I can determine a normal vector to a plane defined by three points or by two directions.
- By the end of this chapter I can write the vector equation and the Cartesian equation of a plane and pass from one form to the other.
- By the end of this chapter I can find the perpendicular distance from a point to a plane and the angle between two planes (GCE extension).
- By the end of this chapter I can solve GCE-style problems combining vector products, triple products and plane equations.

PREREQUISITES

- Vectors in space: components, magnitude, addition and scalar multiplication (Lower Sixth).
- The scalar (dot) product and its use for angles and perpendicularity (Lower Sixth).
- Position vectors, vector equation of a line in space (Lower Sixth).
- Determinants of 2×2 and 3×3 matrices (Further Pure, Lower Sixth).
- Basic trigonometry: sine of an angle, area of a triangle $(1/2)ab \sin C$.

1 SECTION 1: THE VECTOR PRODUCT — DEFINITION AND COMPUTATION

At a construction site in Bonabéri, a mason must check that a metal gate is perfectly rectangular: he measures the diagonals, but an engineer instead computes a cross product of two edge vectors to test perpendicularity and find the exact area in one step. Meanwhile, a carpenter in Bamenda wants the volume of a wedge-shaped roof beam, and a surveyor near Mount Cameroon needs the equation of an inclined plane to position a water tank. All three problems, seemingly different, are solved by one powerful tool: the **vector product** (also called the **cross product**). How can a single operation between two vectors give directions, areas, volumes and even the equations of planes in space? This section introduces the vector product, its geometric meaning, its algebraic properties and the practical determinant method for computation.

1.1 Motivation and Geometric Definition


In three-dimensional space, given two non-zero vectors $\vec{a}$ and $\vec{b}$, there are infinitely many vectors perpendicular to both. To single out one, we need both a magnitude and an orientation. The vector product provides exactly that: a vector perpendicular to the plane containing $\vec{a}$ and $\vec{b}$, whose length equals the area of the parallelogram spanned by $\vec{a}$ and $\vec{b}$, and whose direction follows the **right-hand rule**.

DEFINITION Vector Product — Geometric Definition

Let $\vec{a}$ and $\vec{b}$ be two vectors in $\mathbb{R}^3$ and let $\theta \in [0, \pi]$ be the angle between them. The vector product (or cross product) $\vec{a} \times \vec{b}$ is the vector defined by

$$\vec{a} \times \vec{b} = |\vec{a}| |\vec{b}| \sin \theta \vec{n},$$

where $\vec{n}$ is the unit vector perpendicular to both $\vec{a}$ and $\vec{b}$ such that $(\vec{a}, \vec{b}, \vec{n})$ forms a right-handed system (right-hand rule: curl the fingers of the right hand from $\vec{a}$ towards $\vec{b}$; the extended thumb points in the direction of $\vec{n}$).



Geometric definition of $\vec{a} \times \vec{b}$: magnitude $|\vec{a}| |\vec{b}| \sin \theta$, direction by right-hand rule.

Immediate consequences of the definition:

- If $\vec{a} = \vec{0}$ or $\vec{b} = \vec{0}$, then $\vec{a} \times \vec{b} = \vec{0}$.
- $\vec{a} \times \vec{a} = \vec{0}$ because $\theta = 0$ and $\sin 0 = 0$.
- For non-zero vectors, $\vec{a} \times \vec{b} = \vec{0}$ **if and only if** $\vec{a}$ and $\vec{b}$ are parallel (collinear), i.e. $\theta = 0$ or π .

PROPERTY Zero Cross Product

For non-zero vectors $\vec{a}, \vec{b} \in \mathbb{R}^3$,

$$\vec{a} \times \vec{b} = \vec{0} \iff \vec{a} \text{ and } \vec{b} \text{ are collinear.}$$

1.2 Algebraic Properties of the Vector Product

The vector product is **not commutative**; reversing the order changes the sign because the right-hand rule flips the orientation.

PROPERTY Anti-commutativity

$$\vec{a} \times \vec{b} = -(\vec{b} \times \vec{a}).$$


The standard orthonormal basis $\{\vec{i}, \vec{j}, \vec{k}\}$ obeys a cyclic rule that is easy to remember.

PROPERTY Cross Products of Basis Vectors

$$\vec{i} \times \vec{j} = \vec{k}, \quad \vec{j} \times \vec{k} = \vec{i}, \quad \vec{k} \times \vec{i} = \vec{j},$$

and the reverse products give the opposite vectors:

$$\vec{j} \times \vec{i} = -\vec{k}, \quad \vec{k} \times \vec{j} = -\vec{i}, \quad \vec{i} \times \vec{k} = -\vec{j}.$$



Cyclic rule for the basis vectors: $\vec{i} \rightarrow \vec{j} \rightarrow \vec{k} \rightarrow \vec{i}$.

The vector product is **distributive** over addition:

$$\vec{a} \times (\vec{b} + \vec{c}) = \vec{a} \times \vec{b} + \vec{a} \times \vec{c}, \quad (\vec{a} + \vec{b}) \times \vec{c} = \vec{a} \times \vec{c} + \vec{b} \times \vec{c}.$$

It is **not associative**: $(\vec{a} \times \vec{b}) \times \vec{c} \neq \vec{a} \times (\vec{b} \times \vec{c})$ in general.

1.3 Component Formula and Determinant Method

If $\vec{a} = (a_1, a_2, a_3)$ and $\vec{b} = (b_1, b_2, b_3)$ in the orthonormal basis, expanding $(\vec{a} \times \vec{b})$ using the basis rules yields the component formula.

DEFINITION Vector Product — Component Form

$$\vec{a} \times \vec{b} = \begin{pmatrix} a_2 b_3 - a_3 b_2 \\ a_3 b_1 - a_1 b_3 \\ a_1 b_2 - a_2 b_1 \end{pmatrix}.$$

This is conveniently remembered as the expansion of the symbolic 3×3 determinant

$$\vec{a} \times \vec{b} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}.$$

METHOD Computing $\vec{a} \times \vec{b}$ with the Determinant Scheme

1. Write the determinant with first row $\vec{i}, \vec{j}, \vec{k}$, second row components of $\vec{a}$, third row components of $\vec{b}$.

2. Expand along the first row (cofactor expansion):

$$\begin{aligned}\vec{a} \times \vec{b} &= \vec{i} \begin{vmatrix} a_2 & a_3 \\ b_2 & b_3 \end{vmatrix} - \vec{j} \begin{vmatrix} a_1 & a_3 \\ b_1 & b_3 \end{vmatrix} + \vec{k} \begin{vmatrix} a_1 & a_2 \\ b_1 & b_2 \end{vmatrix} \\ &= (a_2 b_3 - a_3 b_2) \vec{i} + (a_3 b_1 - a_1 b_3) \vec{j} + (a_1 b_2 - a_2 b_1) \vec{k}.\end{aligned}$$

3. Check perpendicularity (optional but recommended): compute $\vec{a} \cdot (\vec{a} \times \vec{b})$ and $\vec{b} \cdot (\vec{a} \times \vec{b})$; both must be 0.

1.4 Worked Examples

WORKED EXAMPLE 1 Computing a Cross Product

Let $\vec{a} = 2\vec{i} - \vec{j} + 3\vec{k}$ and $\vec{b} = \vec{i} + 4\vec{j} - 2\vec{k}$. Compute $\vec{a} \times \vec{b}$ and verify that it is perpendicular to both $\vec{a}$ and $\vec{b}$.

SOLUTION

Solution.

$$\vec{a} \times \vec{b} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 2 & -1 & 3 \\ 1 & 4 & -2 \end{vmatrix} = \vec{i} \begin{vmatrix} -1 & 3 \\ 4 & -2 \end{vmatrix} - \vec{j} \begin{vmatrix} 2 & 3 \\ 1 & -2 \end{vmatrix} + \vec{k} \begin{vmatrix} 2 & -1 \\ 1 & 4 \end{vmatrix}.$$

Calculate the 2×2 determinants:

$$\begin{vmatrix} -1 & 3 \\ 4 & -2 \end{vmatrix} = (-1)(-2) - 3 \cdot 4 = 2 - 12 = -10,$$

$$\begin{vmatrix} 2 & 3 \\ 1 & -2 \end{vmatrix} = 2(-2) - 3 \cdot 1 = -4 - 3 = -7,$$

$$\begin{vmatrix} 2 & -1 \\ 1 & 4 \end{vmatrix} = 2 \cdot 4 - (-1) \cdot 1 = 8 + 1 = 9.$$

Hence

$$\vec{a} \times \vec{b} = -10\vec{i} - (-7)\vec{j} + 9\vec{k} = -10\vec{i} + 7\vec{j} + 9\vec{k}.$$

Verification:

$$\vec{a} \cdot (\vec{a} \times \vec{b}) = (2)(-10) + (-1)(7) + (3)(9) = -20 - 7 + 27 = 0,$$

$$\vec{b} \cdot (\vec{a} \times \vec{b}) = (1)(-10) + (4)(7) + (-2)(9) = -10 + 28 - 18 = 0.$$

Both dot products are zero, confirming perpendicularity.

EXAMPLE Cross Product of Parallel Vectors

Let $\vec{u} = 3\vec{i} - 6\vec{j} + 9\vec{k}$ and $\vec{v} = -\vec{i} + 2\vec{j} - 3\vec{k}$. Notice $\vec{u} = -3\vec{v}$, so they are collinear. Compute $\vec{u} \times \vec{v}$.

Because $\vec{u}$ and $\vec{v}$ are parallel, the angle $\theta = 0$ (or π), $\sin \theta = 0$, and the cross product must be the zero vector. Using the determinant:

$$\vec{u} \times \vec{v} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 3 & -6 & 9 \\ -1 & 2 & -3 \end{vmatrix} = \vec{i}(18 - 18) - \vec{j}(-9 + 9) + \vec{k}(6 - 6) = \vec{0}.$$

This illustrates the property $\vec{a} \times \vec{b} = \vec{0} \iff \vec{a} \parallel \vec{b}$ (for non-zero vectors).

1.5 Geometric Interpretation: Magnitude and Angle

The magnitude of the cross product gives the area of the parallelogram spanned by $\vec{a}$ and $\vec{b}$ and allows us to find $\sin \theta$.

PROPERTY Magnitude of the Cross Product

$$|\vec{a} \times \vec{b}| = |\vec{a}| |\vec{b}| \sin \theta.$$

Consequently, for non-zero vectors,

$$\sin \theta = \frac{|\vec{a} \times \vec{b}|}{|\vec{a}| |\vec{b}|}.$$

WORKED EXAMPLE 2 Finding the Sine of the Angle

Given $\vec{p} = 2\vec{i} + \vec{j} - \vec{k}$ and $\vec{q} = \vec{i} - 2\vec{j} + 2\vec{k}$, find $\sin \theta$ where θ is the angle between $\vec{p}$ and $\vec{q}$.

SOLUTION

First compute $\vec{p} \times \vec{q}$:

$$\vec{p} \times \vec{q} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 2 & 1 & -1 \\ 1 & -2 & 2 \end{vmatrix} = \vec{i}(2 - 2) - \vec{j}(4 + 1) + \vec{k}(-4 - 1) = 0\vec{i} - 5\vec{j} - 5\vec{k}.$$

Magnitudes:

$$|\vec{p} \times \vec{q}| = \sqrt{0^2 + (-5)^2 + (-5)^2} = \sqrt{50} = 5\sqrt{2},$$

$$|\vec{p}| = \sqrt{2^2 + 1^2 + (-1)^2} = \sqrt{6}, \quad |\vec{q}| = \sqrt{1^2 + (-2)^2 + 2^2} = \sqrt{9} = 3.$$

Thus

$$\sin \theta = \frac{5\sqrt{2}}{\sqrt{6} \cdot 3} = \frac{5\sqrt{2}}{3\sqrt{6}} = \frac{5}{3\sqrt{3}} = \frac{5\sqrt{3}}{9}.$$

1.6 Checking Perpendicularity — A Quick Test

The cross product is the standard tool to obtain a vector perpendicular to two given vectors. After computing $\vec{a} \times \vec{b}$, a quick dot-product check guarantees no algebraic slip.

METHOD Verifying Perpendicularity

After obtaining $\vec{c} = \vec{a} \times \vec{b}$, compute

$$\vec{a} \cdot \vec{c} \quad \text{and} \quad \vec{b} \cdot \vec{c}.$$

If both are zero, $\vec{c}$ is perpendicular to both $\vec{a}$ and $\vec{b}$. If not, re-check the determinant expansion.

EXAMPLE Perpendicular Vector to a Plane

Find a vector perpendicular to the plane containing points $A(1, 2, -1)$, $B(3, 0, 2)$ and $C(-1, 1, 3)$.

Two vectors in the plane are $\vec{AB} = B - A = (2, -2, 3)$ and $\vec{AC} = C - A = (-2, -1, 4)$. Their cross product is perpendicular to the plane:

$$\vec{AB} \times \vec{AC} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 2 & -2 & 3 \\ -2 & -1 & 4 \end{vmatrix} = \vec{i}(-8 + 3) - \vec{j}(8 + 6) + \vec{k}(-2 - 4) = -5\vec{i} - 14\vec{j} - 6\vec{k}.$$

Check: $(2, -2, 3) \cdot (-5, -14, -6) = -10 + 28 - 18 = 0$; $(-2, -1, 4) \cdot (-5, -14, -6) = 10 + 14 - 24 = 0$. Hence $(-5, -14, -6)$ is a normal vector to the plane.

1.7 Common Mistakes and Warnings

WARNING Cross Product is Not Commutative

$\vec{a} \times \vec{b} = -(\vec{b} \times \vec{a})$. Swapping the two rows of the determinant changes the sign of every com-

ponent. Always keep the order as given in the problem.

WARNING Do Not Confuse Dot and Cross Products

- Dot product $\vec{a} \cdot \vec{b}$ is a scalar (number): $a_1b_1 + a_2b_2 + a_3b_3$.
- Cross product $\vec{a} \times \vec{b}$ is a vector: $(a_2b_3 - a_3b_2, a_3b_1 - a_1b_3, a_1b_2 - a_2b_1)$.

Writing $\vec{a} \cdot \vec{b} = \vec{c}$ or $\vec{a} \times \vec{b} = k$ (scalar) is a serious error.

WARNING Determinant Sign Errors

When expanding the 3×3 determinant, the $\vec{j}$ component comes with a minus sign:

$$-\vec{j} \begin{vmatrix} a_1 & a_3 \\ b_1 & b_3 \end{vmatrix}.$$

Forgetting this minus sign is the most frequent computational mistake.

1.8 Summary of Key Points

KEY POINTS Vector Product — Essentials

- Geometric definition: $\vec{a} \times \vec{b} = |\vec{a}||\vec{b}| \sin \theta \vec{n}$, $\vec{n}$ by right-hand rule.
- Component formula: $\vec{a} \times \vec{b} = (a_2b_3 - a_3b_2, a_3b_1 - a_1b_3, a_1b_2 - a_2b_1)$.

• Determinant mnemonic:
$$\begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}.$$

- Anti-commutative: $\vec{a} \times \vec{b} = -(\vec{b} \times \vec{a})$.
- Zero cross product $\iff$ collinear (for non-zero vectors).
- Magnitude: $|\vec{a} \times \vec{b}| = |\vec{a}||\vec{b}| \sin \theta = \text{area of parallelogram}$.
- Perpendicularity check: $\vec{a} \cdot (\vec{a} \times \vec{b}) = \vec{b} \cdot (\vec{a} \times \vec{b}) = 0$.

The vector product is now at our disposal. In the next sections we will see how it yields areas

of triangles and parallelograms, volumes of parallelepipeds and tetrahedra via the scalar triple product, and how it provides the normal vector needed to write the vector and Cartesian equations of a plane.

2 SECTION 2: ALGEBRAIC PROPERTIES AND GEOMETRIC APPLICATIONS (AREAS)

In Section 1 we learned how to compute the vector product $\vec{a} \times \vec{b}$ using a determinant and we saw its geometric definition: a vector perpendicular to both $\vec{a}$ and $\vec{b}$ with magnitude $|\vec{a}| |\vec{b}| \sin \theta$. Now we turn to the algebraic rules that govern this operation. Understanding these properties is essential for manipulating vector expressions in mechanics, geometry, and further mathematics. We will also discover the most immediate geometric application: calculating areas of parallelograms and triangles in three-dimensional space.

2.1 Algebraic Properties of the Vector Product

The vector product behaves differently from the scalar product and from ordinary multiplication of numbers. The three fundamental algebraic properties are anticommutativity, distributivity over addition, and compatibility with scalar multiplication.

PROPERTY Algebraic Properties of the Cross Product

Let $\vec{a}, \vec{b}, \vec{c} \in \mathbb{R}^3$ and $k \in \mathbb{R}$.

1. Anticommutativity: $\vec{b} \times \vec{a} = -(\vec{a} \times \vec{b})$.
2. Distributivity over addition: $\vec{a} \times (\vec{b} + \vec{c}) = \vec{a} \times \vec{b} + \vec{a} \times \vec{c}$.
3. Compatibility with scalar multiplication: $(k\vec{a}) \times \vec{b} = k(\vec{a} \times \vec{b}) = \vec{a} \times (k\vec{b})$.

EXAMPLE Proof of Anticommutativity from the Determinant

Recall the determinant definition:

$$\vec{a} \times \vec{b} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}.$$

Swapping the second and third rows (which correspond to the components of $\vec{a}$ and $\vec{b}$) changes the sign of the determinant. Hence

$$\vec{b} \times \vec{a} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ b_1 & b_2 & b_3 \\ a_1 & a_2 & a_3 \end{vmatrix} = - \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} = -(\vec{a} \times \vec{b}).$$

This also follows immediately from the geometric definition: $\vec{a} \times \vec{b}$ and $\vec{b} \times \vec{a}$ have the same magnitude $|\vec{a}||\vec{b}|\sin\theta$ but opposite directions (right-hand rule vs left-hand rule).

WARNING Non-Associativity – A Critical Trap

The vector product is not associative. In general,

$$(\vec{a} \times \vec{b}) \times \vec{c} \neq \vec{a} \times (\vec{b} \times \vec{c}).$$

Counter-example: Take the standard basis vectors $\vec{i}, \vec{j}, \vec{k}$.

$$\vec{i} \times (\vec{i} \times \vec{j}) = \vec{i} \times \vec{k} = -\vec{j},$$

but

$$(\vec{i} \times \vec{i}) \times \vec{j} = \vec{0} \times \vec{j} = \vec{0}.$$

Since $-\vec{j} \neq \vec{0}$, associativity fails. Never write $\vec{a} \times \vec{b} \times \vec{c}$ without parentheses; the order of operations matters.

WORKED EXAMPLE 3 Using Algebraic Properties to Simplify

Simplify the expression $(\vec{a} + 2\vec{b}) \times (3\vec{a} - \vec{b})$.

SOLUTION

Solution: We expand using distributivity and scalar compatibility, treating the cross product like a bilinear operation but respecting the order.

$$\begin{aligned}(\vec{a} + 2\vec{b}) \times (3\vec{a} - \vec{b}) &= \vec{a} \times (3\vec{a} - \vec{b}) + 2\vec{b} \times (3\vec{a} - \vec{b}) \\ &= 3(\vec{a} \times \vec{a}) - \vec{a} \times \vec{b} + 6(\vec{b} \times \vec{a}) - 2(\vec{b} \times \vec{b}).\end{aligned}$$

Since $\vec{v} \times \vec{v} = \vec{0}$ for any vector and $\vec{b} \times \vec{a} = -\vec{a} \times \vec{b}$,

$$\begin{aligned}&= 3\vec{0} - \vec{a} \times \vec{b} + 6(-\vec{a} \times \vec{b}) - 2\vec{0} \\ &= -7(\vec{a} \times \vec{b}).\end{aligned}$$

2.2 Geometric Interpretation: Area of a Parallelogram

The magnitude of the cross product has a beautiful and extremely useful geometric meaning.

PROPERTY Area of a Parallelogram


Let $\vec{a}$ and $\vec{b}$ be two non-zero, non-parallel vectors in $\mathbb{R}^3$. The area A of the parallelogram having $\vec{a}$ and $\vec{b}$ as adjacent sides is

$$A = |\vec{a} \times \vec{b}|.$$

EXAMPLE Proof

The area of a parallelogram is base $\times$ height. Take $|\vec{a}|$ as the base. The height is the perpendicular distance from the tip of $\vec{b}$ to the line along $\vec{a}$, which is $|\vec{b}| \sin \theta$, where θ is the angle between $\vec{a}$ and $\vec{b}$. Hence

$$\text{Area} = |\vec{a}| \cdot (|\vec{b}| \sin \theta) = |\vec{a}| |\vec{b}| \sin \theta = |\vec{a} \times \vec{b}|.$$



The parallelogram spanned by $\vec{a}$ and $\vec{b}$. The height is $h = |\vec{b}| \sin \theta$, so area $= |\vec{a}| h = |\vec{a} \times \vec{b}|$. The cross product vector is perpendicular to the plane.

WORKED EXAMPLE 4 Area of a Parallelogram from Vertices

Find the area of the parallelogram with vertices $A(1, 2, -1)$, $B(3, 1, 0)$, $C(4, 3, 2)$, $D(2, 4, 1)$.

SOLUTION

Solution: Two adjacent sides are $\vec{AB}$ and $\vec{AD}$.

$$\vec{AB} = \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix}, \quad \vec{AD} = \begin{pmatrix} 1 \\ 2 \\ 2 \end{pmatrix}.$$

$$\vec{AB} \times \vec{AD} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 2 & -1 & 1 \\ 1 & 2 & 2 \end{vmatrix} = \vec{i}(-2-2) - \vec{j}(4-1) + \vec{k}(4+1) = \begin{pmatrix} -4 \\ -3 \\ 5 \end{pmatrix}.$$

$$\text{Area} = |\vec{AB} \times \vec{AD}| = \sqrt{(-4)^2 + (-3)^2 + 5^2} = \sqrt{16 + 9 + 25} = \sqrt{50} = 5\sqrt{2} \text{ square units.}$$


2.3 Area of a Triangle in Space

A triangle is exactly half of a parallelogram. This gives a very efficient formula for the area of a triangle given the coordinates of its vertices in 3D.

METHOD Area of Triangle ABC in Space

Given three points A, B, C with position vectors $\vec{a}, \vec{b}, \vec{c}$:

1. Compute two side vectors, e.g. $\vec{AB} = \vec{b} - \vec{a}$ and $\vec{AC} = \vec{c} - \vec{a}$.
2. Compute the cross product $\vec{AB} \times \vec{AC}$.
3. The area is $\frac{1}{2}|\vec{AB} \times \vec{AC}|$.



Triangle ABC in space. The vector $\vec{AB} \times \vec{AC}$ is perpendicular to the plane of the triangle; its magnitude is twice the area of the triangle.

WORKED EXAMPLE 5 Area of a Triangle with Given Vertices

Find the area of triangle ABC where $A(2, -1, 3)$, $B(4, 0, 1)$, $C(1, 2, -1)$.

SOLUTION

Solution:

$$\vec{AB} = \begin{pmatrix} 2 \\ 1 \\ -2 \end{pmatrix}, \quad \vec{AC} = \begin{pmatrix} -1 \\ 3 \\ -4 \end{pmatrix}.$$

$$\vec{AB} \times \vec{AC} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 2 & 1 & -2 \\ -1 & 3 & -4 \end{vmatrix} = \vec{i}(-4 + 6) - \vec{j}(-8 - 2) + \vec{k}(6 + 1) = \begin{pmatrix} 2 \\ 10 \\ 7 \end{pmatrix}.$$

$$|\vec{AB} \times \vec{AC}| = \sqrt{2^2 + 10^2 + 7^2} = \sqrt{4 + 100 + 49} = \sqrt{153} = 3\sqrt{17}.$$

$$\text{Area} = \frac{1}{2} \times 3\sqrt{17} = \frac{3\sqrt{17}}{2} \text{ square units.}$$

2.4 Unit Vector Perpendicular to Two Given Vectors

Since $\vec{a} \times \vec{b}$ is perpendicular to both $\vec{a}$ and $\vec{b}$, normalising it gives a unit normal vector. This is fundamental for defining plane orientations, flux integrals, and surface normals.

METHOD Finding a Unit Vector Perpendicular to $\vec{a}$ and $\vec{b}$

1. Compute $\vec{n} = \vec{a} \times \vec{b}$.
2. Compute its magnitude $|\vec{n}|$.
3. The required unit vectors are $\pm \frac{\vec{n}}{|\vec{n}|}$ (two opposite directions).

If $\vec{a}$ and $\vec{b}$ are parallel, $\vec{a} \times \vec{b} = \vec{0}$ and there is no unique perpendicular direction (infinitely many).

WORKED EXAMPLE 6 Unit Normal Vector

Find a unit vector perpendicular to both $\vec{a} = 2\vec{i} - \vec{j} + 3\vec{k}$ and $\vec{b} = \vec{i} + 2\vec{j} - \vec{k}$.

SOLUTION

Solution:

$$\vec{a} \times \vec{b} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 2 & -1 & 3 \\ 1 & 2 & -1 \end{vmatrix} = \vec{i}(1 - 6) - \vec{j}(-2 - 3) + \vec{k}(4 + 1) = \begin{pmatrix} -5 \\ 5 \\ 5 \end{pmatrix}.$$

Magnitude: $|\vec{a} \times \vec{b}| = \sqrt{25 + 25 + 25} = \sqrt{75} = 5\sqrt{3}$. Unit vector: $\hat{n} = \frac{1}{5\sqrt{3}} \begin{pmatrix} -5 \\ 5 \\ 5 \end{pmatrix} = \frac{1}{\sqrt{3}} \begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix}$. The opposite vector $-\hat{n} = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 \\ -1 \\ -1 \end{pmatrix}$ is also a valid answer.

2.5 Distance from a Point to a Line in Space (GCE Extension)

Although not explicitly listed in the core syllabus, this application appears frequently in GCE Advanced Level papers as a "useful extension". It combines the cross product magnitude (area) with the parallelogram area formula.

PROPERTY Distance from Point P to Line ℓ

Let ℓ be a line with direction vector $\vec{u}$ passing through point A . The perpendicular distance d from a point P to ℓ is

$$d = \frac{|\vec{AP} \times \vec{u}|}{|\vec{u}|}.$$

EXAMPLE Derivation

The area of the parallelogram formed by $\vec{AP}$ and $\vec{u}$ is $|\vec{AP} \times \vec{u}|$. This area also equals base $\times$ height $= |\vec{u}| \cdot d$. Equating the two expressions gives the formula.

WORKED EXAMPLE 7 Distance from Point to Line

Find the perpendicular distance from the point $P(1, 2, 3)$ to the line $\ell : \frac{x-2}{1} = \frac{y+1}{2} = \frac{z}{-1}$.

SOLUTION

Solution: The line ℓ passes through $A(2, -1, 0)$ and has direction vector $\vec{u} = \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix}$.

$$\vec{AP} = \begin{pmatrix} 1 - 2 \\ 2 - (-1) \\ 3 - 0 \end{pmatrix} = \begin{pmatrix} -1 \\ 3 \\ 3 \end{pmatrix}.$$

$$\vec{AP} \times \vec{u} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ -1 & 3 & 3 \\ 1 & 2 & -1 \end{vmatrix} = \vec{i}(-3 - 6) - \vec{j}(1 - 3) + \vec{k}(-2 - 3) = \begin{pmatrix} -9 \\ 2 \\ -5 \end{pmatrix}.$$

$$|\vec{AP} \times \vec{u}| = \sqrt{81 + 4 + 25} = \sqrt{110}, \quad |\vec{u}| = \sqrt{1 + 4 + 1} = \sqrt{6}.$$

$$d = \frac{\sqrt{110}}{\sqrt{6}} = \sqrt{\frac{110}{6}} = \sqrt{\frac{55}{3}} = \frac{\sqrt{165}}{3} \text{ units.}$$

KEY POINTS Key Points of Section 2

- Anticommutativity: $\vec{b} \times \vec{a} = -(\vec{a} \times \vec{b})$. Order matters!
- Distributivity: $\vec{a} \times (\vec{b} + \vec{c}) = \vec{a} \times \vec{b} + \vec{a} \times \vec{c}$.
- Scalar multiplication: $(k\vec{a}) \times \vec{b} = k(\vec{a} \times \vec{b})$.
- Non-associativity: $(\vec{a} \times \vec{b}) \times \vec{c} \neq \vec{a} \times (\vec{b} \times \vec{c})$ in general.
- Area of parallelogram: $|\vec{a} \times \vec{b}|$.
- Area of triangle ABC: $\frac{1}{2} |\vec{AB} \times \vec{AC}|$.
- Unit normal to $\vec{a}, \vec{b}$: $\pm \frac{\vec{a} \times \vec{b}}{|\vec{a} \times \vec{b}|}$.
- Distance point-to-line: $d = \frac{|\vec{AP} \times \vec{u}|}{|\vec{u}|}$.

DID YOU KNOW? The Vector Product in Physics and Engineering

The cross product is ubiquitous in physics. Torque (moment of a force) is $\vec{\tau} = \vec{r} \times \vec{F}$. The magnetic force on a moving charge is $\vec{F} = q\vec{v} \times \vec{B}$. Angular momentum is $\vec{L} = \vec{r} \times \vec{p}$. In all these cases, the magnitude gives the "strength" of the effect (proportional to $\sin \theta$) and the direction gives the axis of rotation or the direction of the force, following the right-hand rule.

The area interpretation is used in computer graphics to compute surface normals for lighting and shading calculations (e.g., in video games or CAD software used in Douala's engineering firms).

EXERCISE 1 Practice Exercises

1. Given $\vec{a} = \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix}$, $\vec{b} = \begin{pmatrix} 3 \\ 0 \\ 1 \end{pmatrix}$, $\vec{c} = \begin{pmatrix} -1 \\ 1 \\ 2 \end{pmatrix}$, verify the distributive law $\vec{a} \times (\vec{b} + \vec{c}) = \vec{a} \times \vec{b} + \vec{a} \times \vec{c}$.
2. Show that $(\vec{i} + \vec{j}) \times (\vec{j} + \vec{k}) \times (\vec{k} + \vec{i})$ depends on the placement of parentheses by evaluating both $(\vec{u} \times \vec{v}) \times \vec{w}$ and $\vec{u} \times (\vec{v} \times \vec{w})$ where $\vec{u} = \vec{i} + \vec{j}$, $\vec{v} = \vec{j} + \vec{k}$, $\vec{w} = \vec{k} + \vec{i}$.
3. Find the area of the triangle with vertices $A(1, 1, 0)$, $B(2, -1, 2)$, $C(0, 3, 1)$.
4. Find a unit vector perpendicular to the plane containing points $P(1, 0, 2)$, $Q(2, 1, 0)$, $R(0, 2, 1)$.
5. The line ℓ passes through $A(1, 2, -1)$ and has direction vector $\vec{d} = 2\vec{i} - \vec{j} + 2\vec{k}$. Find the shortest distance from the origin O to ℓ .

SOLUTION Exercise 1

$$\begin{aligned} \vec{b} + \vec{c} &= \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix}, \vec{a} \times (\vec{b} + \vec{c}) = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 1 & 2 & -1 \\ 2 & 1 & 3 \end{vmatrix} = \vec{i}(6 + 1) - \vec{j}(3 + 2) + \vec{k}(1 - 4) = \begin{pmatrix} 7 \\ -5 \\ -3 \end{pmatrix}, \vec{a} \times \vec{b} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 1 & 2 & -1 \\ 3 & 0 & 1 \end{vmatrix} = \\ \vec{i}(2 - 0) - \vec{j}(1 + 3) + \vec{k}(0 - 6) &= \begin{pmatrix} 2 \\ -4 \\ -6 \end{pmatrix}, \vec{a} \times \vec{c} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 1 & 2 & -1 \\ -1 & 1 & 2 \end{vmatrix} = \vec{i}(4 + 1) - \vec{j}(2 - 1) + \vec{k}(1 + 2) = \begin{pmatrix} 5 \\ -1 \\ 3 \end{pmatrix}. \text{ Sum} \\ &= \begin{pmatrix} 7 \\ -5 \\ -3 \end{pmatrix}. \text{ Verified.} \end{aligned}$$

SOLUTION Exercise 2

$$\vec{u} \times \vec{v} = (\vec{i} + \vec{j}) \times (\vec{j} + \vec{k}) = \vec{i} \times \vec{j} + \vec{i} \times \vec{k} + \vec{j} \times \vec{j} + \vec{j} \times \vec{k} = \vec{k} - \vec{j} + \vec{0} + \vec{i} = \vec{i} - \vec{j} + \vec{k}. (\vec{u} \times \vec{v}) \times \vec{w} = (\vec{i} - \vec{j} + \vec{k}) \times (\vec{k} + \vec{i}) = \vec{i} \times \vec{k} + \vec{i} \times \vec{i} - \vec{j} \times \vec{k} -$$

$\vec{j} \times \vec{i} + \vec{k} \times \vec{k} + \vec{k} \times \vec{i} = -\vec{j} + \vec{0} - \vec{i} + \vec{k} + \vec{0} + \vec{j} = -\vec{i} + \vec{k}$. $\vec{v} \times \vec{w} = (\vec{j} + \vec{k}) \times (\vec{k} + \vec{i}) = \vec{j} \times \vec{k} + \vec{j} \times \vec{i} + \vec{k} \times \vec{k} + \vec{k} \times \vec{i} = \vec{i} - \vec{k} + \vec{0} + \vec{j} = \vec{i} + \vec{j} - \vec{k}$.
 $\vec{u} \times (\vec{v} \times \vec{w}) = (\vec{i} + \vec{j}) \times (\vec{i} + \vec{j} - \vec{k}) = \vec{i} \times \vec{i} + \vec{i} \times \vec{j} - \vec{i} \times \vec{k} + \vec{j} \times \vec{i} + \vec{j} \times \vec{j} - \vec{j} \times \vec{k} = \vec{0} + \vec{k} + \vec{j} - \vec{k} + \vec{0} - \vec{i} = -\vec{i} + \vec{j}$.
 Since $-\vec{i} + \vec{k} \neq -\vec{i} + \vec{j}$, the results differ. Non-associativity confirmed.

SOLUTION Exercise 3

$$\vec{AB} = \begin{pmatrix} 1 \\ -2 \\ 2 \end{pmatrix}, \vec{AC} = \begin{pmatrix} -1 \\ 2 \\ 1 \end{pmatrix}. \vec{AB} \times \vec{AC} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 1 & -2 & 2 \\ -1 & 2 & 1 \end{vmatrix} = \vec{i}(-2-4) - \vec{j}(1+2) + \vec{k}(2-2) = \begin{pmatrix} -6 \\ -3 \\ 0 \end{pmatrix}. |\vec{AB} \times \vec{AC}| = \sqrt{36+9} = \sqrt{45} = 3\sqrt{5}. \text{ Area} = \frac{1}{2} \times 3\sqrt{5} = \frac{3\sqrt{5}}{2} \text{ square units.}$$

SOLUTION Exercise 4

$$\vec{PQ} = \begin{pmatrix} 1 \\ 1 \\ -2 \end{pmatrix}, \vec{PR} = \begin{pmatrix} -1 \\ 2 \\ -1 \end{pmatrix}. \vec{PQ} \times \vec{PR} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 1 & 1 & -2 \\ -1 & 2 & -1 \end{vmatrix} = \vec{i}(-1+4) - \vec{j}(-1-2) + \vec{k}(2+1) = \begin{pmatrix} 3 \\ 3 \\ 3 \end{pmatrix}. |\vec{PQ} \times \vec{PR}| = \sqrt{27} = 3\sqrt{3}. \text{ Unit normal } \hat{n} = \pm \frac{1}{3\sqrt{3}} \begin{pmatrix} 3 \\ 3 \\ 3 \end{pmatrix} = \pm \frac{1}{\sqrt{3}} \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}.$$

SOLUTION Exercise 5

$$A(1, 2, -1), \vec{d} = \begin{pmatrix} 2 \\ -1 \\ 2 \end{pmatrix}, P = O(0, 0, 0). \vec{AO} = \begin{pmatrix} -1 \\ -2 \\ 1 \end{pmatrix}. \vec{AO} \times \vec{d} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ -1 & -2 & 1 \\ 2 & -1 & 2 \end{vmatrix} = \vec{i}(-4+1) - \vec{j}(-2-2) + \vec{k}(1+4) = \begin{pmatrix} -3 \\ 4 \\ 5 \end{pmatrix}. |\vec{AO} \times \vec{d}| = \sqrt{9+16+25} = \sqrt{50} = 5\sqrt{2}. |\vec{d}| = \sqrt{4+1+4} = 3. d = \frac{5\sqrt{2}}{3} \text{ units.}$$

3 SECTION 3: THE SCALAR TRIPLE PRODUCT AND VOLUMES

In Section 2 we saw that the vector product $\vec{b} \times \vec{c}$ encodes areas: its magnitude is the area of the parallelogram built on $\vec{b}$ and $\vec{c}$. A natural question now arises: can we measure volumes in space with vectors? Imagine a mason in Bamenda building a concrete block shaped like a parallelepiped, with three edges given by vectors $\vec{a}$, $\vec{b}$ and $\vec{c}$ meeting at one corner. The base is a parallelogram of area $\|\vec{b} \times \vec{c}\|$, and the volume is “base area $\times$ height”. The height is the component of $\vec{c}$ along the direction perpendicular to the base — and that perpendicular direction is precisely the direction of $\vec{b} \times \vec{c}$. Combining the two ideas (a vector product followed by a scalar product) gives a single number that measures volume: the **scalar triple product**. This section defines it, computes it as a determinant, proves its cyclic properties, and applies it to volumes of parallelepipeds and tetrahedra and to coplanarity tests — all classic GCE Advanced Level material.

3.1 Definition and Determinant Formula

DEFINITION Scalar triple product

Let $\vec{a}$, $\vec{b}$, $\vec{c}$ be three vectors in space. The **scalar triple product** of $\vec{a}$, $\vec{b}$, $\vec{c}$ (in this order) is the real number

$$[\vec{a}, \vec{b}, \vec{c}] = \vec{a} \cdot (\vec{b} \times \vec{c}).$$

It is also called the mixed product. Note carefully: the result is a scalar, not a vector.

The name “mixed” comes from the fact that we mix the two products we know: first a vector product $\vec{b} \times \vec{c}$ (a vector), then a scalar product of $\vec{a}$ with that vector (a number).

PROPERTY Determinant form of the scalar triple product

In an orthonormal basis $(\vec{i}, \vec{j}, \vec{k})$, if

$$\vec{a} = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}, \quad \vec{b} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix}, \quad \vec{c} = \begin{pmatrix} c_1 \\ c_2 \\ c_3 \end{pmatrix},$$

then

$$\vec{a} \cdot (\vec{b} \times \vec{c}) = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}.$$

(The same value is obtained if the components are placed in columns instead of rows.) This formula is examinable; its derivation is instructive and given below.

Derivation.

From Section 1, $\vec{b} \times \vec{c} = \begin{pmatrix} b_2c_3 - b_3c_2 \\ b_3c_1 - b_1c_3 \\ b_1c_2 - b_2c_1 \end{pmatrix}$. Taking the scalar product with $\vec{a}$:

$$\vec{a} \cdot (\vec{b} \times \vec{c}) = a_1(b_2c_3 - b_3c_2) + a_2(b_3c_1 - b_1c_3) + a_3(b_1c_2 - b_2c_1),$$

which is exactly the expansion of the 3×3 determinant along its first row:

$$\begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} = a_1 \begin{vmatrix} b_2 & b_3 \\ c_2 & c_3 \end{vmatrix} - a_2 \begin{vmatrix} b_1 & b_3 \\ c_1 & c_3 \end{vmatrix} + a_3 \begin{vmatrix} b_1 & b_2 \\ c_1 & c_2 \end{vmatrix}. \quad \blacksquare$$

EXAMPLE First computation

Let $\vec{a} = \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix}$, $\vec{b} = \begin{pmatrix} 3 \\ -1 \\ 1 \end{pmatrix}$, $\vec{c} = \begin{pmatrix} 0 \\ 2 \\ 4 \end{pmatrix}$. Then

$$[\vec{a}, \vec{b}, \vec{c}] = \begin{vmatrix} 1 & 2 & 0 \\ 3 & -1 & 1 \\ 0 & 2 & 4 \end{vmatrix} = 1 \begin{vmatrix} -1 & 1 \\ 2 & 4 \end{vmatrix} - 2 \begin{vmatrix} 3 & 1 \\ 0 & 4 \end{vmatrix} + 0 = 1(-4 - 2) - 2(12 - 0) = -6 - 24 = -30.$$

The negative sign will be explained geometrically below: it means $\vec{a}$ lies on the opposite side of the plane of $(\vec{b}, \vec{c})$ from the vector $\vec{b} \times \vec{c}$.

3.2 Algebraic Properties: Cyclic Invariance and Sign Changes**PROPERTY Cyclic invariance and swap rule**

For any vectors $\vec{a}, \vec{b}, \vec{c}$:

$$\vec{a} \cdot (\vec{b} \times \vec{c}) = \vec{b} \cdot (\vec{c} \times \vec{a}) = \vec{c} \cdot (\vec{a} \times \vec{b}) \quad (\text{cyclic permutations preserve the value});$$

$$\vec{a} \cdot (\vec{b} \times \vec{c}) = -\vec{a} \cdot (\vec{c} \times \vec{b}) = -\vec{b} \cdot (\vec{a} \times \vec{c}) \quad (\text{swapping any two vectors changes the sign}).$$

Consequently one may also write $\vec{a} \cdot (\vec{b} \times \vec{c}) = (\vec{a} \times \vec{b}) \cdot \vec{c}$.

Justification.

These follow at once from the determinant form: a cyclic permutation of the three rows of a determinant can be achieved by two row swaps, so the sign changes twice and the value is unchanged; swapping two adjacent vectors swaps two rows, which changes the sign. The last identity comes from $\vec{c} \cdot (\vec{a} \times \vec{b}) = (\vec{a} \times \vec{b}) \cdot \vec{c}$ by commutativity of the scalar product.

WARNING Order matters

The scalar triple product is not symmetric in its three arguments: $[\vec{a}, \vec{b}, \vec{c}] = -[\vec{a}, \vec{c}, \vec{b}]$. Only cyclic reorderings leave it unchanged. In an exam, keep the order given in the question.

3.3 Geometric Interpretation: Volume of the Parallelepiped**PROPERTY** Volume of the parallelepiped

The volume V of the parallelepiped built on three vectors $\vec{a}, \vec{b}, \vec{c}$ drawn from the same point is

$$V = |\vec{a} \cdot (\vec{b} \times \vec{c})|.$$

This theorem is examinable; the proof below is instructive and frequently asked as a “show that” question.

Proof.


Take the parallelogram built on $\vec{b}$ and $\vec{c}$ as the base. Its area is $S = \|\vec{b} \times \vec{c}\|$ (Section 2). The vector $\vec{n} = \vec{b} \times \vec{c}$ is perpendicular to the base. The height h of the parallelepiped is the absolute value of the component of $\vec{a}$ along $\vec{n}$:

$$h = \|\vec{a}\| \cos \theta,$$

where θ is the angle between $\vec{a}$ and $\vec{n}$. Hence

$$V = S \times h = \|\vec{n}\| \|\vec{a}\| \cos \theta = |\vec{a} \cdot \vec{n}| = |\vec{a} \cdot (\vec{b} \times \vec{c})|. \quad \blacksquare$$

The sign of the triple product also carries information: it is positive when $(\vec{a}, \vec{b}, \vec{c})$ forms a right-handed triad relative to $\vec{b} \times \vec{c}$, i.e. when $\vec{a}$ and $\vec{b} \times \vec{c}$ point to the same side of the base plane.



Parallelepiped built on $\vec{a}, \vec{b}, \vec{c}$: base area $\|\vec{b} \times \vec{c}\|$, height $h = \text{component of } \vec{a} \text{ along } \vec{b} \times \vec{c}$, so $V = |\vec{a} \cdot (\vec{b} \times \vec{c})|$.

EXAMPLE Volume of a parallelepiped

Using the vectors of the previous example, the parallelepiped built on $\vec{a}, \vec{b}, \vec{c}$ has volume

$$V = |-30| = 30 \text{ cubic units.}$$


3.4 Volume of a Tetrahedron

A tetrahedron is a pyramid with a triangular base. If we cut a parallelepiped built on $\vec{AB}, \vec{AC}, \vec{AD}$ by the planes through its diagonal structure, the tetrahedron $ABCD$ occupies exactly one sixth of it (the triangular base has half the area of the parallelogram, and a pyramid has one third of the volume of the prism on the same base with the same height: $\frac{1}{2} \times \frac{1}{3} = \frac{1}{6}$).

PROPERTY Volume of a tetrahedron

For four points A, B, C, D in space,

$$V_{ABCD} = \frac{1}{6} |\vec{AB} \cdot (\vec{AC} \times \vec{AD})|.$$



Tetrahedron $ABCD$: the three edges from A give $V = \frac{1}{6} |\vec{AB} \cdot (\vec{AC} \times \vec{AD})|$.

WARNING The factor is $\frac{1}{6}$, not $\frac{1}{3}$

A very common error is to write $V = \frac{1}{3} |\vec{AB} \cdot (\vec{AC} \times \vec{AD})|$. The factor $\frac{1}{3}$ alone would be correct

for a pyramid whose base is the parallelogram built on $\vec{AC}, \vec{AD}$. But the tetrahedron's base is the triangle ACD , whose area is half the parallelogram. Hence $V = \frac{1}{3} \times (\text{triangle area}) \times h = \frac{1}{3} \times \frac{1}{2} \|\vec{AC} \times \vec{AD}\| \times h = \frac{1}{6} |\vec{AB} \cdot (\vec{AC} \times \vec{AD})|$.

3.5 Coplanarity Tests

If three vectors $\vec{a}, \vec{b}, \vec{c}$ lie in one plane (are **coplanar**), then the parallelepiped they build is completely flat: its volume is zero. Conversely, if the triple product is zero and the vectors are non-zero, no volume is spanned, so they must be coplanar.

PROPERTY Coplanarity criteria

- Three vectors $\vec{a}, \vec{b}, \vec{c}$ are coplanar if and only if $\vec{a} \cdot (\vec{b} \times \vec{c}) = 0$.
- Four points A, B, C, D are coplanar if and only if

$$\vec{AB} \cdot (\vec{AC} \times \vec{AD}) = 0.$$

WARNING Zero triple product does not mean zero vectors

$[\vec{a}, \vec{b}, \vec{c}] = 0$ does not imply that any of the vectors is $\vec{0}$. It only means the three vectors are coplanar (or one of them is zero, or two are parallel — all special cases of coplanarity). Never conclude “ $\vec{a} = \vec{0}$ ” from a vanishing triple product.

METHOD Testing coplanarity of four points

1. Form the three edge vectors from one vertex: $\vec{AB}, \vec{AC}, \vec{AD}$.
2. Write the 3×3 determinant of their components (rows or columns).
3. Expand the determinant (e.g. along a row).
4. Conclude: the points are coplanar $\iff$ the determinant equals 0. If a parameter is involved, solve the resulting equation.

WORKED EXAMPLE 8 Finding a parameter for coplanarity (GCE style)

The points $A(1, 0, 2)$, $B(2, 1, 1)$, $C(0, 3, 1)$ and $D(1, 1, \lambda)$ are coplanar. Find the value(s) of λ .

SOLUTION

Solution. Form the edge vectors from A:

$$\vec{AB} = \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}, \quad \vec{AC} = \begin{pmatrix} -1 \\ 3 \\ -1 \end{pmatrix}, \quad \vec{AD} = \begin{pmatrix} 0 \\ 1 \\ \lambda - 2 \end{pmatrix}.$$

Coplanarity requires

$$\begin{vmatrix} 1 & 1 & -1 \\ -1 & 3 & -1 \\ 0 & 1 & \lambda - 2 \end{vmatrix} = 0.$$

Expanding along the first row:

$$\begin{aligned} & 1 \begin{vmatrix} 3 & -1 \\ 1 & \lambda - 2 \end{vmatrix} - 1 \begin{vmatrix} -1 & -1 \\ 0 & \lambda - 2 \end{vmatrix} + (-1) \begin{vmatrix} -1 & 3 \\ 0 & 1 \end{vmatrix} \\ &= 1(3(\lambda - 2) + 1) - 1(-(\lambda - 2) - 0) - 1(-1 - 0) \\ &= 3\lambda - 5 + \lambda - 2 + 1 = 4\lambda - 6. \end{aligned}$$

Setting $4\lambda - 6 = 0$ gives $\lambda = \frac{3}{2}$.

WORKED EXAMPLE 9 Height of a tetrahedron from volume and base area

A tetrahedron ABCD has $A(0, 0, 0)$, $B(2, 0, 0)$, $C(0, 3, 0)$, $D(1, 1, 4)$.

(a) Compute its volume.

(b) Taking triangle ABC as base, deduce the height from D.

SOLUTION

Solution. (a) $\vec{AB} = \begin{pmatrix} 2 \\ 0 \\ 0 \end{pmatrix}$, $\vec{AC} = \begin{pmatrix} 0 \\ 3 \\ 0 \end{pmatrix}$, $\vec{AD} = \begin{pmatrix} 1 \\ 1 \\ 4 \end{pmatrix}$.

$$\vec{AB} \cdot (\vec{AC} \times \vec{AD}) = \begin{vmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 1 & 1 & 4 \end{vmatrix} = 2 \begin{vmatrix} 3 & 0 \\ 1 & 4 \end{vmatrix} = 2 \times 12 = 24.$$

Hence $V = \frac{1}{6} |24| = 4$ cubic units.

(b) Base area: $S_{ABC} = \frac{1}{2} \|\vec{AB} \times \vec{AC}\| = \frac{1}{2} \left\| \begin{pmatrix} 0 \\ 0 \\ 6 \end{pmatrix} \right\| = 3$. Since $V = \frac{1}{3}Sh$, we get $h = \frac{3V}{S} = \frac{3 \times 4}{3} = 4$ units. (Check: the base lies in the plane $z = 0$ and D has height 4 — consistent.)

EXERCISE 2 Practice

1. Compute $\vec{a} \cdot (\vec{b} \times \vec{c})$ for $\vec{a} = (1, -1, 2)$, $\vec{b} = (2, 0, 1)$, $\vec{c} = (1, 1, -1)$, and state the volume of the parallelepiped they determine.
2. Show that the points $A(1, 1, 0)$, $B(2, 0, 1)$, $C(0, 2, 1)$, $D(3, -1, 2)$ are coplanar.
3. Find k so that $A(0, 0, 0)$, $B(1, 2, 0)$, $C(2, 1, 1)$, $D(1, 1, k)$ are coplanar.
4. A tetrahedron has vertices $A(1, 0, 0)$, $B(0, 2, 0)$, $C(0, 0, 3)$, $D(2, 2, 2)$. Find its volume and the height from D to the plane ABC given that the area of triangle ABC is $\frac{7}{2}$.

SOLUTION Exercise 1

1. $\begin{vmatrix} 1 & -1 & 2 \\ 2 & 0 & 1 \\ 1 & 1 & -1 \end{vmatrix} = 1(0 - 1) - (-1)(-2 - 1) + 2(2 - 0) = -1 - 3 + 4 = 0$. The volume is 0: the three vectors are coplanar.
2. $\vec{AB} = (1, -1, 1)$, $\vec{AC} = (-1, 1, 1)$, $\vec{AD} = (2, -2, 2) = 2\vec{AB}$. The determinant has two proportional rows, hence its value is 0. The points are coplanar.
3. $\begin{vmatrix} 1 & 2 & 0 \\ 2 & 1 & 1 \\ 1 & 1 & k \end{vmatrix} = 1(k - 1) - 2(2k - 1) + 0 = k - 1 - 4k + 2 = -3k + 1 = 0$, so $k = \frac{1}{3}$.
4. $\vec{AB} = (-1, 2, 0)$, $\vec{AC} = (-1, 0, 3)$, $\vec{AD} = (1, 2, 2)$. Triple product: $-1(0 \cdot 2 - 3 \cdot 2) - 2((-1)(2) - 3 \cdot 1) + 0 = -1(-6) - 2(-5) = 6 + 10 = 16$. Volume $V = \frac{16}{6} = \frac{8}{3}$. Height: $h = \frac{3V}{S} = \frac{8}{7/2} = \frac{16}{7}$.

KEY POINTS Key points of Section 3

- Scalar triple product: $[\vec{a}, \vec{b}, \vec{c}] = \vec{a} \cdot (\vec{b} \times \vec{c}) = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}$ — a scalar.
- Cyclic permutations leave it unchanged; swapping two vectors changes the sign.
- Volume of parallelepiped: $V = |[\vec{a}, \vec{b}, \vec{c}]|$; volume of tetrahedron: $V = \frac{1}{6} |\vec{AB} \cdot (\vec{AC} \times \vec{AD})|$.
- Coplanarity: A, B, C, D coplanar $\iff \vec{AB} \cdot (\vec{AC} \times \vec{AD}) = 0$.
- Traps: the tetrahedron factor is $\frac{1}{6}$, not $\frac{1}{3}$; a zero triple product means coplanarity, not zero vectors.

DID YOU KNOW? Why “triple”?

The scalar triple product appears in physics as the volume element of crystallography: the volume of the unit cell of a crystal lattice (for instance in the study of minerals from the Cameroonian bauxite deposits) is $|\vec{a} \cdot (\vec{b} \times \vec{c})|$ where $\vec{a}, \vec{b}, \vec{c}$ are the lattice vectors. The same formula also detects whether three forces applied to a rigid body are coplanar — a key step in statics problems.

4 SECTION 4: EQUATIONS OF A PLANE — VECTOR AND CARTESIAN FORMS

In Section 3 we saw how the scalar triple product $\vec{a} \cdot (\vec{b} \times \vec{c})$ gives the volume of a parallelepiped. That same cross product $\vec{b} \times \vec{c}$ produces a vector perpendicular to both $\vec{b}$ and $\vec{c}$ — a normal vector to the plane containing them. This simple observation is the key to writing down the equation of any plane in space. Whether we are given a point and a normal, three points, or a point and two directions, the cross product provides the normal we need to switch effortlessly between vector (parametric) and Cartesian forms. Mastering these conversions is a staple of GCE Advanced Level Further Mathematics; they appear in almost every paper, often as the first step in a longer problem on lines, planes, distances or angles.

4.1 Specifying a Plane in Space

A plane in three-dimensional space can be uniquely determined in three equivalent ways:

- A point A (with position vector $\vec{a}$) and a **normal vector** $\vec{n} \neq \vec{0}$ perpendicular to the plane.
- A point A and two **non-parallel direction vectors** $\vec{u}$ and $\vec{v}$ lying in the plane.
- Three **non-collinear points** A, B, C (the vectors $\vec{AB}$ and $\vec{AC}$ then serve as the two direction vectors).

The cross product $\vec{u} \times \vec{v}$ immediately gives a normal vector $\vec{n}$ when we have the two-direction description. This links the three descriptions together.

4.2 Vector (Parametric) Equation of a Plane

If a plane passes through point $A(\vec{a})$ and is parallel to the non-parallel vectors $\vec{u}$ and $\vec{v}$, then any point $R(\vec{r})$ in the plane can be reached by starting at A and moving some distance along $\vec{u}$ and some distance along $\vec{v}$.

DEFINITION Vector (Parametric) Equation of a Plane


The vector equation of a plane through $A(\vec{a})$ with direction vectors $\vec{u}$ and $\vec{v}$ ($\vec{u} \nparallel \vec{v}$) is

$$\vec{r} = \vec{a} + s\vec{u} + t\vec{v}, \quad s, t \in \mathbb{R}.$$

The parameters s and t are independent real numbers; varying them sweeps out the whole plane.

WARNING Two Parameters are Essential

A single parameter s in $\vec{r} = \vec{a} + s\vec{u}$ describes a line, not a plane. A plane requires two independent parameters because it is a two-dimensional surface. Forgetting the second parameter is a very common error in examinations.



Vector equation $\vec{r} = \vec{a} + s\vec{u} + t\vec{v}$: the plane is spanned by $\vec{u}$ and $\vec{v}$ from point A ; the normal $\vec{n} = \vec{u} \times \vec{v}$ is perpendicular to the plane.

4.3 Normal Form and Cartesian Equation

If we know a point $A(\vec{a})$ on the plane and a normal vector $\vec{n} = \begin{pmatrix} a \\ b \\ c \end{pmatrix}$, the condition that $\vec{r} = \begin{pmatrix} x \\ y \\ z \end{pmatrix}$ lies in the plane is simply that the vector $\vec{r} - \vec{a}$ is perpendicular to $\vec{n}$:

$$(\vec{r} - \vec{a}) \cdot \vec{n} = 0.$$

PROPERTY Normal Form and Cartesian Equation

Let $\vec{n} = \begin{pmatrix} a \\ b \\ c \end{pmatrix}$ be a normal vector to a plane Π , and let $A(x_0, y_0, z_0)$ be a point on Π . Then Π has equation

$$(\vec{r} - \vec{a}) \cdot \vec{n} = 0 \iff a(x - x_0) + b(y - y_0) + c(z - z_0) = 0.$$

Expanding gives the Cartesian equation

$$ax + by + cz = d, \quad \text{where } d = ax_0 + by_0 + cz_0 = \vec{n} \cdot \vec{a}.$$

WARNING Coefficients Give the Normal

In the Cartesian equation $ax + by + cz = d$, the coefficients (a, b, c) are the components of a normal vector to the plane. They are not direction vectors lying in the plane. A direction vector $\vec{v}$ in the plane satisfies $av_x + bv_y + cv_z = 0$.

4.4 Finding the Cartesian Equation from Three Points

When three non-collinear points A, B, C are given, we form two vectors in the plane, e.g. $\vec{AB}$ and $\vec{AC}$, and take their cross product to obtain a normal vector.

METHOD Cartesian Equation Through Three Points

1. Compute $\vec{AB} = \vec{b} - \vec{a}$ and $\vec{AC} = \vec{c} - \vec{a}$.
2. Find a normal vector $\vec{n} = \vec{AB} \times \vec{AC}$.
3. Use point A (or B or C) to find $d = \vec{n} \cdot \vec{a}$.

4. Write $ax + by + cz = d$ where $\vec{n} = \begin{pmatrix} a \\ b \\ c \end{pmatrix}$.

WORKED EXAMPLE 10 Plane Through Three Points

Find the Cartesian equation of the plane passing through $A(1, 2, -1)$, $B(3, 0, 2)$ and $C(-1, 1, 3)$.

SOLUTION

Solution.

$$\vec{AB} = \begin{pmatrix} 2 \\ -2 \\ 3 \end{pmatrix}, \quad \vec{AC} = \begin{pmatrix} -2 \\ -1 \\ 4 \end{pmatrix}.$$

$$\begin{aligned} \vec{n} = \vec{AB} \times \vec{AC} &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -2 & 3 \\ -2 & -1 & 4 \end{vmatrix} = \mathbf{i}((-2)(4) - 3(-1)) - \mathbf{j}((2)(4) - 3(-2)) + \mathbf{k}((2)(-1) - (-2)(-2)) \\ &= \mathbf{i}(-8 + 3) - \mathbf{j}(8 + 6) + \mathbf{k}(-2 - 4) = \begin{pmatrix} -5 \\ -14 \\ -6 \end{pmatrix}. \end{aligned}$$

We can multiply by -1 for a simpler normal: $\vec{n} = \begin{pmatrix} 5 \\ 14 \\ 6 \end{pmatrix}$. Using point $A(1, 2, -1)$:

$$d = 5(1) + 14(2) + 6(-1) = 5 + 28 - 6 = 27.$$

The Cartesian equation is

$$5x + 14y + 6z = 27.$$

(Check: B gives $5(3) + 14(0) + 6(2) = 15 + 12 = 27$; C gives $5(-1) + 14(1) + 6(3) = -5 + 14 + 18 = 27$.)

4.5 Converting Between Parametric and Cartesian Forms

The cross product is the bridge between the two forms.

METHOD Parametric $\rightarrow$ Cartesian

Given $\vec{r} = \vec{a} + s\vec{u} + t\vec{v}$:

1. Compute $\vec{n} = \vec{u} \times \vec{v}$.
2. Compute $d = \vec{n} \cdot \vec{a}$.
3. Write $\vec{n} \cdot \vec{r} = d$, i.e. $ax + by + cz = d$.

METHOD Cartesian $\rightarrow$ Parametric

Given $ax + by + cz = d$ with normal $\vec{n} = \begin{pmatrix} a \\ b \\ c \end{pmatrix}$:

1. Find one point A on the plane (e.g. set two variables to 0 and solve for the third, provided the corresponding coefficient is non-zero).
2. Find two independent direction vectors $\vec{u}, \vec{v}$ perpendicular to $\vec{n}$ (i.e. $\vec{n} \cdot \vec{u} = 0, \vec{n} \cdot \vec{v} = 0$).

A quick way: if $a \neq 0$, take $\vec{u} = \begin{pmatrix} -b \\ a \\ 0 \end{pmatrix}$ and $\vec{v} = \begin{pmatrix} -c \\ 0 \\ a \end{pmatrix}$; adjust if some coefficients are zero.

3. Write $\vec{r} = \vec{a} + s\vec{u} + t\vec{v}$.

WORKED EXAMPLE 11 Converting Both Ways

1. Convert $\vec{r} = \begin{pmatrix} 1 \\ -2 \\ 3 \end{pmatrix} + s \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix} + t \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}$ to Cartesian form.

2. Convert $2x - y + 3z = 6$ to parametric form.

SOLUTION

Solution.

1. $\vec{u} = \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix}, \vec{v} = \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}, \vec{n} = \vec{u} \times \vec{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & 1 & 0 \\ -1 & 0 & 1 \end{vmatrix} = \mathbf{i}(1 \cdot 1 - 0 \cdot 0) - \mathbf{j}(2 \cdot 1 - 0 \cdot (-1)) + \mathbf{k}(2 \cdot 0 - 1 \cdot (-1)) = \begin{pmatrix} 1 \\ -2 \\ 1 \end{pmatrix}.$
 $d = \vec{n} \cdot \vec{a} = 1(1) + (-2)(-2) + 1(3) = 1 + 4 + 3 = 8$. Cartesian equation: $x - 2y + z = 8$.

2. Normal $\vec{n} = \begin{pmatrix} 2 \\ -1 \\ 3 \end{pmatrix}$. Find a point: set $y = 0, z = 0 \Rightarrow 2x = 6 \Rightarrow x = 3$, so $A(3, 0, 0)$. Two

direction vectors perpendicular to $\vec{n}$: $\vec{u} = \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix}$ (since $2(1) + (-1)(2) + 3(0) = 0$), $\vec{v} = \begin{pmatrix} 0 \\ 3 \\ 1 \end{pmatrix}$

(since $2(0) + (-1)(3) + 3(1) = 0$). Parametric form: $\vec{r} = \begin{pmatrix} 3 \\ 0 \\ 0 \end{pmatrix} + s \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix} + t \begin{pmatrix} 0 \\ 3 \\ 1 \end{pmatrix}, s, t \in \mathbb{R}.$

4.6 Perpendicular Distance from a Point to a Plane (GCE Extension)

The shortest distance from a point $P(x_1, y_1, z_1)$ to the plane $\Pi : ax + by + cz = d$ is the length of the projection of $\vec{AP}$ onto the normal $\vec{n}$, where A is any point on Π .

PROPERTY Distance from Point to Plane

The perpendicular distance from $P(x_1, y_1, z_1)$ to the plane $ax + by + cz = d$ is

$$\text{dist}(P, \Pi) = \frac{|ax_1 + by_1 + cz_1 - d|}{\sqrt{a^2 + b^2 + c^2}}.$$

WORKED EXAMPLE 12 Distance from Point to Plane

Find the distance from $P(2, -1, 3)$ to the plane $3x - 4y + 12z = 13$.


SOLUTION

Substitute into the formula:

$$\text{Numerator} = |3(2) - 4(-1) + 12(3) - 13| = |6 + 4 + 36 - 13| = 33.$$

$$\text{Denominator} = \sqrt{3^2 + (-4)^2 + 12^2} = \sqrt{9 + 16 + 144} = \sqrt{169} = 13.$$

$$\text{Distance} = \frac{33}{13} \text{ units.}$$



Perpendicular distance from point P to plane $\Pi : ax + by + cz = d$. The distance is the length of the projection of $\vec{AP}$ onto the normal $\vec{n}$. The foot F is correctly computed as the intersection of the line through P parallel to $\vec{n}$ with the plane.

4.7 Angle Between Two Planes (GCE Extension)

The angle between two planes is defined as the angle between their normal vectors (or its supplement, whichever is acute).

PROPERTY Angle Between Two Planes

Let $\Pi_1 : a_1x + b_1y + c_1z = d_1$ and $\Pi_2 : a_2x + b_2y + c_2z = d_2$ have normals $\vec{n}_1 = \begin{pmatrix} a_1 \\ b_1 \\ c_1 \end{pmatrix}$, $\vec{n}_2 = \begin{pmatrix} a_2 \\ b_2 \\ c_2 \end{pmatrix}$. The

acute angle θ between the planes satisfies

$$\cos \theta = \frac{|\vec{n}_1 \cdot \vec{n}_2|}{|\vec{n}_1| |\vec{n}_2|} = \frac{|a_1a_2 + b_1b_2 + c_1c_2|}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}}.$$

- $\Pi_1 \parallel \Pi_2 \iff \vec{n}_1 \parallel \vec{n}_2 \iff (a_1, b_1, c_1) = k(a_2, b_2, c_2)$ for some $k \neq 0$.
- $\Pi_1 \perp \Pi_2 \iff \vec{n}_1 \perp \vec{n}_2 \iff a_1a_2 + b_1b_2 + c_1c_2 = 0$.

WORKED EXAMPLE 13 Angle Between Planes

Find the acute angle between the planes $2x - y + 2z = 5$ and $3x + 6y - 2z = 7$.

SOLUTION

$$\vec{n}_1 = \begin{pmatrix} 2 \\ -1 \\ 2 \end{pmatrix}, \vec{n}_2 = \begin{pmatrix} 3 \\ 6 \\ -2 \end{pmatrix}. \vec{n}_1 \cdot \vec{n}_2 = 2(3) + (-1)(6) + 2(-2) = 6 - 6 - 4 = -4. |\vec{n}_1| = \sqrt{4 + 1 + 4} = 3, \quad |\vec{n}_2| = \sqrt{9 + 36 + 4} = 7. \cos \theta = \frac{|-4|}{3 \cdot 7} = \frac{4}{21}. \theta = \cos^{-1}\left(\frac{4}{21}\right) \approx 79.0^\circ.$$

4.8 Summary and Examination Tips**KEY POINTS** Key Points for GCE Advanced Level

- Vector form: $\vec{r} = \vec{a} + s\vec{u} + t\vec{v}$ ($s, t \in \mathbb{R}, \vec{u} \nparallel \vec{v}$).
- Normal form: $(\vec{r} - \vec{a}) \cdot \vec{n} = 0$.
- Cartesian form: $ax + by + cz = d$ with $\vec{n} = \begin{pmatrix} a \\ b \\ c \end{pmatrix}$.
- Normal from two directions: $\vec{n} = \vec{u} \times \vec{v}$.
- Plane through three points: $\vec{n} = \vec{AB} \times \vec{AC}$, then $d = \vec{n} \cdot \vec{a}$.
- Distance point-plane: $\frac{|ax_1 + by_1 + cz_1 - d|}{\sqrt{a^2 + b^2 + c^2}}$.
- Angle between planes: $\cos \theta = \frac{|\vec{n}_1 \cdot \vec{n}_2|}{|\vec{n}_1||\vec{n}_2|}$.
- Parallel planes: normals are parallel. Perpendicular planes: normals are perpendicular.

WARNING Common Examination Pitfalls

- Forgetting that the parametric form needs two parameters.
- Confusing the normal vector (a, b, c) with a direction vector in the plane.

- When converting Cartesian $\rightarrow$ parametric, choosing direction vectors that are not perpendicular to the normal (check $\vec{n} \cdot \vec{u} = 0$).
- In the distance formula, using the wrong sign for d (remember d is the constant on the right-hand side of $ax + by + cz = d$).
- For the angle between planes, always take the absolute value of the dot product to get the acute angle.

DID YOU KNOW? Planes in Real Life

The roof of a traditional Cameroonian case (round hut) is a conical surface, but the flat ceiling of a modern classroom is a plane. The equation $ax + by + cz = d$ describes that ceiling: the normal vector (a, b, c) points straight up (or down), and d fixes the height. In computer graphics, every flat polygon on your screen is a plane segment defined by exactly these equations — the cross product of two edges gives the normal used for lighting calculations.

With this section we complete Chapter FMA14. You now have a full toolkit: vector products for areas, scalar triple products for volumes, and the equations of planes for describing flat surfaces in space. These ideas are the foundation of three-dimensional analytic geometry and appear repeatedly in mechanics, computer graphics, and higher mathematics. Practice converting between forms until it becomes automatic — it is the single most useful skill for the GCE paper.

5 INTEGRATION ACTIVITY

5.1 Aims of the activity

By the end of this group activity, you should be able to:

- compute a **vector product** $\vec{AB} \times \vec{AC}$ from coordinates and interpret it geometrically as a normal vector and as an area;
- derive the **Cartesian equation of a plane** from a point and a normal vector;
- use the **scalar triple product** to test whether four points are coplanar and to compute the **volume of a tetrahedron**;
- compute the **perpendicular distance from a point to a plane** and interpret it in a real safety context;
- work cooperatively, present a reasoned solution on the board, and compare the efficiency of different methods.

KEY POINTS Toolbox for this activity

You will need the following results from this chapter:

- For $\vec{u}(x, y, z)$ and $\vec{v}(x', y', z')$,

$$\vec{u} \times \vec{v} = \begin{pmatrix} yz' - zy' \\ zx' - xz' \\ xy' - yx' \end{pmatrix}.$$

- $\vec{u} \times \vec{v}$ is perpendicular to both $\vec{u}$ and $\vec{v}$, and $|\vec{u} \times \vec{v}|$ is the area of the parallelogram built on $\vec{u}$ and $\vec{v}$; the triangle built on them has area $\frac{1}{2}|\vec{u} \times \vec{v}|$.
- Plane through A with normal vector $\vec{n}(a, b, c)$: $ax + by + cz + d = 0$, where d is found by substituting the coordinates of A .
- $[\vec{u}, \vec{v}, \vec{w}] = (\vec{u} \times \vec{v}) \cdot \vec{w}$; four points A, B, C, D are coplanar if and only if $[\vec{AB}, \vec{AC}, \vec{AD}] = 0$; the volume of the tetrahedron $ABCD$ is $\frac{1}{6} |[\vec{AB}, \vec{AC}, \vec{AD}]|$.
- Distance from $P(x_0, y_0, z_0)$ to the plane $ax + by + cz + d = 0$:

$$d = \frac{|ax_0 + by_0 + cz_0 + d|}{\sqrt{a^2 + b^2 + c^2}}.$$

5.2 Situation

EXPERIMENT Designing a Roof Truss for a School Hall in Buea

A contractor in Buea is roofing the new assembly hall of a secondary school. Because of the heavy rains on the slopes of Mount Cameroon, the roof must be steep enough for water to run off, and the carpenter wants exact measurements before cutting the timber and ordering the zinc sheets.

The contractor sets up a 3D coordinate system $(O; \vec{i}, \vec{j}, \vec{k})$, with the unit 1 metre on each axis, fixed to the hall under construction. One rectangular face of the sloping roof is modelled by the plane through the three points


$$A(0, 0, 4), \quad B(6, 0, 4), \quad C(6, 8, 6),$$

where A and B are the two ends of the lower edge of the roof face (at height 4 m) and C is one end of the upper edge (at height 6 m). The triangular panel ABC is half of that rectangular face and will be covered with corrugated zinc sheets, sold by square metre.

A wooden beam is to be placed at the fourth corner $D(0, 8, 6)$, which the contractor believes completes the rectangle $ABDC$ of the roof face. Before nailing anything, the carpenter must check that A , B , C and D really lie in the same plane; if they do, the “attic space” $ABCD$ is flat and the architect must raise the point D to create storage volume.

Finally, a lamp hangs from the ceiling framework at the point $P(3, 4, 2)$. Safety regulations require a clearance of at least 2.5 m between the lamp and the roof plane directly above it. The contractor asks your group to verify whether the regulation is satisfied.

Your group must produce a complete written report, then present it on the board.



The roof face $ABDC$, the triangular panel ABC , a normal vector $\vec{n}$ and the lamp at P .

5.3 Tasks

Work in groups of three or four. Answer the questions in order; each one uses a different tool of the chapter.

1. **A normal vector to the roof.** Compute the vectors $\vec{AB}$ and $\vec{AC}$. Calculate the vector product $\vec{AB} \times \vec{AC}$. Explain in one sentence why this vector is normal to the roof plane.
2. **Cartesian equation of the roof plane.** Using point A and your normal vector, show that the roof plane has Cartesian equation $y - 4z + 16 = 0$. Verify that B and C also satisfy this equation.

- Area of the zinc panel.** Compute $|\vec{AB} \times \vec{AC}|$ and deduce the exact area of the triangular panel ABC , then an approximate value to 2 decimal places. If zinc sheets cost 4 500 FCFA per square metre and the whole rectangular face is twice the triangle, estimate the cost for this face (round up to the next square metre).
- Is the beam D in the right place?** Compute the scalar triple product $[\vec{AB}, \vec{AC}, \vec{AD}]$ where $D(0, 8, 6)$. What do you conclude about the four points? What is the volume of the “tetrahedron” $ABCD$? Interpret this bad news for the carpenter.
- Creating an attic.** The architect moves the fourth corner to $D'(0, 8, 8)$. Compute the new scalar triple product and the volume of the tetrahedron $ABCD'$. Is the attic now usable?
- Safety clearance of the lamp.** Compute the perpendicular distance from $P(3, 4, 2)$ to the roof plane. Is the regulation clearance of 2.5 m respected?
- Presentation.** Prepare a poster showing: the normal vector, the plane equation, the panel area, the coplanarity test, the attic volume and the clearance. Compare with another group: did anyone use $\vec{AC} \times \vec{AB}$ instead? What changes, what does not?

METHOD How to attack each question

- Normal vector:** form $\vec{AB}$ and $\vec{AC}$ by subtracting coordinates, then apply the cross product formula component by component.
- Plane equation:** simplify the normal vector if possible (divide by a common factor), write $a(x - x_A) + b(y - y_A) + c(z - z_A) = 0$, expand, and check the other given points.
- Area:** take the modulus of the cross product (parallelogram area), then halve it for the triangle.
- Coplanarity and volume:** dot the cross product with the third vector; zero means coplanar, otherwise take $\frac{1}{6}$ of the absolute value.
- Distance:** substitute the point into the left-hand side of the plane equation, take the absolute value, and divide by the norm of the normal vector.

WARNING Traps to avoid in this activity

- The vector product is anti-commutative: $\vec{AC} \times \vec{AB} = -(\vec{AB} \times \vec{AC})$. The sign of the normal vector changes, but the plane equation and the areas do not.
- Do not forget the factor $\frac{1}{2}$ for the triangle and $\frac{1}{6}$ for the tetrahedron.
- A zero scalar triple product does not mean you made an error: it means the points are coplanar.
- In the distance formula, use the normalised denominator $\sqrt{a^2 + b^2 + c^2}$; forgetting it

gives a wrong clearance.

- When simplifying a normal vector by dividing by a common factor, the plane equation changes form but describes the same plane; always re-check with a known point.

5.4 Model answer

1. **Normal vector.** From the coordinates,

$$\vec{AB} = (6 - 0, 0 - 0, 4 - 4) = (6, 0, 0), \quad \vec{AC} = (6 - 0, 8 - 0, 6 - 4) = (6, 8, 2).$$

Applying the cross product formula with $\vec{AB} = (6, 0, 0)$ and $\vec{AC} = (6, 8, 2)$:

$$\vec{AB} \times \vec{AC} = \begin{pmatrix} (0)(2) - (0)(8) \\ (0)(6) - (6)(2) \\ (6)(8) - (0)(6) \end{pmatrix} = \begin{pmatrix} 0 \\ -12 \\ 48 \end{pmatrix}.$$

By definition, the vector product is perpendicular to both $\vec{AB}$ and $\vec{AC}$, which lie in the roof plane; hence $\vec{n} = (0, -12, 48)$ is normal to the roof plane. We may simplify to $\vec{n} = (0, -1, 4)$ (dividing by 12), since any non-zero scalar multiple of a normal vector is still normal to the plane.

2. **Equation of the plane.** A point $M(x, y, z)$ belongs to the plane if and only if $\vec{AM} \cdot \vec{n} = 0$. With $\vec{AM} = (x - 0, y - 0, z - 4)$ and $\vec{n} = (0, -1, 4)$:

$$(0)(x - 0) + (-1)(y - 0) + (4)(z - 4) = 0 \iff -y + 4z - 16 = 0 \iff y - 4z + 16 = 0.$$

Check: $B(6, 0, 4)$: $0 - 16 + 16 = 0$ ✓; $C(6, 8, 6)$: $8 - 24 + 16 = 0$ ✓. Both points lie in the plane, as expected.

3. **Area of the panel.**

$$|\vec{AB} \times \vec{AC}| = \sqrt{0^2 + (-12)^2 + 48^2} = \sqrt{144 + 2304} = \sqrt{2448} = \sqrt{144 \times 17} = 12\sqrt{17}.$$

This is the area of the parallelogram built on $\vec{AB}$ and $\vec{AC}$, so

$$\text{Area}(ABC) = \frac{1}{2} |\vec{AB} \times \vec{AC}| = 6\sqrt{17} \approx 24.74 \text{ m}^2.$$

The full rectangular face has area $12\sqrt{17} \approx 49.48 \text{ m}^2$, so the contractor orders 50 m^2 of zinc, costing about $50 \times 4\,500 = 225\,000 \text{ FCFA}$.

4. **Coplanarity test for $D(0, 8, 6)$.** We have $\vec{AD} = (0 - 0, 8 - 0, 6 - 4) = (0, 8, 2)$, and

$$[\vec{AB}, \vec{AC}, \vec{AD}] = (\vec{AB} \times \vec{AC}) \cdot \vec{AD} = (0)(0) + (-12)(8) + (48)(2) = -96 + 96 = 0.$$

The scalar triple product is zero, so A, B, C, D are **coplanar**: indeed D satisfies the plane equation ($8 - 24 + 16 = 0$). The “tetrahedron” $ABCD$ is flat:

$$V = \frac{1}{6} |[\vec{AB}, \vec{AC}, \vec{AD}]| = 0 \text{ m}^3.$$

The beam at D lies exactly in the roof plane: the carpenter’s belief is correct, but there is no attic volume at all.

5. Attic with $D'(0, 8, 8)$. Now $\vec{AD}' = (0, 8, 4)$ and

$$[\vec{AB}, \vec{AC}, \vec{AD}'] = (0)(0) + (-12)(8) + (48)(4) = -96 + 192 = 96.$$

$$V_{ABCD'} = \frac{1}{6}|96| = 16 \text{ m}^3.$$

The volume is non-zero: raising the corner to height 8 m creates an attic of 16 m^3 , which is usable storage space.

6. Clearance of the lamp. With the plane $y - 4z + 16 = 0$ (so $a = 0, b = 1, c = -4, d = 16$) and $P(3, 4, 2)$:

$$d = \frac{|(0)(3) + (1)(4) + (-4)(2) + 16|}{\sqrt{0^2 + 1^2 + (-4)^2}} = \frac{|4 - 8 + 16|}{\sqrt{17}} = \frac{12}{\sqrt{17}} \approx 2.91 \text{ m}.$$

Since $2.91 > 2.5$, the safety clearance is respected.

7. Comparison of methods. Using $\vec{AC} \times \vec{AB} = (0, 12, -48)$ gives the opposite normal vector, hence the equation $-y + 4z - 16 = 0$, which is the same plane (both sides multiplied by -1); areas, volumes (in absolute value) and distances are unchanged.

EXERCISE 3 Going further: a second roof face

The opposite face of the roof passes through $A'(0, 0, 4)$, $E(0, 8, 6)$ and $F(-6, 8, 6)$.

1. Find a normal vector to this face and its Cartesian equation.
2. Compute the area of the triangular panel $A'EF$.
3. Show that the two roof planes $y - 4z + 16 = 0$ and the plane you found intersect along a line parallel to $\vec{i}$, and interpret this line on the building.

SOLUTION Going further: a second roof face

1. $\vec{A'E} = (0, 8, 2)$ and $\vec{A'F} = (-6, 8, 2)$, so

$$\vec{A'E} \times \vec{A'F} = \begin{pmatrix} (8)(2) - (2)(8) \\ (2)(-6) - (0)(2) \\ (0)(8) - (8)(-6) \end{pmatrix} = \begin{pmatrix} 0 \\ -12 \\ 48 \end{pmatrix},$$

the same normal direction $(0, -1, 4)$. The plane through $A'(0, 0, 4)$ is $-y + 4(z - 4) = 0$, i.e. $y - 4z + 16 = 0$: the two faces are in fact coplanar in this model (a single sloping face).

$$2. \text{Area}(A'EF) = \frac{1}{2}\sqrt{0^2 + (-12)^2 + 48^2} = \frac{1}{2}(12\sqrt{17}) = 6\sqrt{17} \approx 24.74 \text{ m}^2.$$

3. Both planes have normal vector $(0, -1, 4)$ and pass through $A = A'$, so they coincide; their “intersection” is the whole plane, which contains the direction $\vec{i} = (1, 0, 0)$ since $(0, -1, 4) \cdot (1, 0, 0) = 0$. On the building, $\vec{i}$ is the horizontal direction along the lower edge AB of the roof.

DID YOU KNOW?**Why this matters beyond the classroom**

Roofing errors are expensive in the South-West Region, where wind-driven rain quickly finds any badly joined panel. The same mathematics used here — cross products for normals and areas, triple products for volumes — is built into the CAD software used by modern architects in Douala and Yaoundé, and into the machines that cut roof panels to measure.


6 METHODS AND WORKED EXAMPLES

7 THE VECTOR PRODUCT — DEFINITION AND COMPUTATION

DEFINITION Vector Product (Cross Product)

Let $\vec{a}$ and $\vec{b}$ be two vectors in three-dimensional space. The **vector product** (or **cross product**) $\vec{a} \times \vec{b}$ is a vector defined by:

- **Magnitude:** $|\vec{a} \times \vec{b}| = |\vec{a}| |\vec{b}| \sin \theta$, where $\theta \in [0, \pi]$ is the angle between $\vec{a}$ and $\vec{b}$.
- **Direction:** Perpendicular to both $\vec{a}$ and $\vec{b}$, oriented according to the **right-hand rule**: if the fingers of the right hand curl from $\vec{a}$ towards $\vec{b}$ through the angle θ , the extended thumb points in the direction of $\vec{a} \times \vec{b}$.
- **Zero vector:** If $\vec{a} = \vec{0}$ or $\vec{b} = \vec{0}$ or $\vec{a} \parallel \vec{b}$ ($\theta = 0$ or π), then $\vec{a} \times \vec{b} = \vec{0}$.



Geometric definition of the vector product: magnitude $|\vec{a}| |\vec{b}| \sin \theta$, direction by right-hand rule.

PROPERTY Component Formula (Determinant Form)

If $\vec{a} = a_1\vec{i} + a_2\vec{j} + a_3\vec{k}$ and $\vec{b} = b_1\vec{i} + b_2\vec{j} + b_3\vec{k}$, then

$$\vec{a} \times \vec{b} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} = (a_2b_3 - a_3b_2)\vec{i} - (a_1b_3 - a_3b_1)\vec{j} + (a_1b_2 - a_2b_1)\vec{k}.$$

The result is a vector perpendicular to both $\vec{a}$ and $\vec{b}$.

METHOD Computing the Vector Product in Component Form

Goal: Find $\vec{a} \times \vec{b}$ when $\vec{a} = a_1\vec{i} + a_2\vec{j} + a_3\vec{k}$ and $\vec{b} = b_1\vec{i} + b_2\vec{j} + b_3\vec{k}$.

1. Write the vectors as ordered triples: $\vec{a} = (a_1, a_2, a_3)$, $\vec{b} = (b_1, b_2, b_3)$.
2. Set up the 3×3 determinant with unit vectors $\vec{i}, \vec{j}, \vec{k}$ in the first row.
3. Expand along the first row (Laplace expansion), taking care of the minus sign for the $\vec{j}$ -component (cofactor expansion).
4. Simplify each component.
5. Verification (Reflex): Compute $\vec{a} \cdot (\vec{a} \times \vec{b})$ and $\vec{b} \cdot (\vec{a} \times \vec{b})$; both must equal 0.

Key point: The $\vec{j}$ -component comes with a minus sign: $-(a_1b_3 - a_3b_1)$.

WORKED EXAMPLE 14 Vector Product of Two Given Vectors

Let $\vec{a} = 2\vec{i} - 3\vec{j} + \vec{k}$ and $\vec{b} = -\vec{i} + 4\vec{j} - 2\vec{k}$.

1. Compute $\vec{a} \times \vec{b}$.
2. Verify that the result is orthogonal to both $\vec{a}$ and $\vec{b}$.

SOLUTION

Solution.

1. Components: $\vec{a} = (2, -3, 1)$, $\vec{b} = (-1, 4, -2)$.

$$\vec{a} \times \vec{b} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 2 & -3 & 1 \\ -1 & 4 & -2 \end{vmatrix}$$

Expand along the first row:

$$\vec{i}\text{-component} = (-3)(-2) - (1)(4) = 6 - 4 = 2,$$

$$\vec{j}\text{-component} = -[(2)(-2) - (1)(-1)] = -[-4 + 1] = 3,$$

$$\vec{k}\text{-component} = (2)(4) - (-3)(-1) = 8 - 3 = 5.$$

$$\text{Hence } \vec{a} \times \vec{b} = 2\vec{i} + 3\vec{j} + 5\vec{k}.$$

2. Orthogonality check:

$$\vec{a} \cdot (\vec{a} \times \vec{b}) = (2)(2) + (-3)(3) + (1)(5) = 4 - 9 + 5 = 0,$$

$$\vec{b} \cdot (\vec{a} \times \vec{b}) = (-1)(2) + (4)(3) + (-2)(5) = -2 + 12 - 10 = 0.$$

Both dot products are zero, confirming orthogonality.

KEY POINTS Key Formulas for the Vector Product

- $\vec{a} \times \vec{b} = (a_2b_3 - a_3b_2, a_3b_1 - a_1b_3, a_1b_2 - a_2b_1).$
- $|\vec{a} \times \vec{b}| = |\vec{a}||\vec{b}| \sin \theta.$
- $\vec{a} \times \vec{b} = \vec{0} \iff \vec{a} \parallel \vec{b}$ (or one is zero).
- $\vec{a} \cdot (\vec{a} \times \vec{b}) = 0$ and $\vec{b} \cdot (\vec{a} \times \vec{b}) = 0$ (orthogonality).

WARNING Common Mistakes in Cross Product

- Sign error on j -component: Forgetting the minus sign from the cofactor expansion. Always write $-(a_1b_3 - a_3b_1).$
- Order matters: $\vec{a} \times \vec{b} = -(\vec{b} \times \vec{a})$ (anti-commutativity). Do not swap vectors carelessly.
- Not associative: $(\vec{a} \times \vec{b}) \times \vec{c} \neq \vec{a} \times (\vec{b} \times \vec{c})$ in general. Always use brackets.
- Confusing with dot product: Cross product yields a vector; dot product yields a scalar.

8 ALGEBRAIC PROPERTIES AND GEOMETRIC APPLICATIONS (AREAS)

PROPERTY Algebraic Properties of the Vector Product

For any vectors $\vec{a}, \vec{b}, \vec{c} \in \mathbb{R}^3$ and scalar $\lambda \in \mathbb{R}$:

1. Anti-commutativity: $\vec{a} \times \vec{b} = -(\vec{b} \times \vec{a}).$
2. Distributivity over addition: $\vec{a} \times (\vec{b} + \vec{c}) = \vec{a} \times \vec{b} + \vec{a} \times \vec{c}, (\vec{a} + \vec{b}) \times \vec{c} = \vec{a} \times \vec{c} + \vec{b} \times \vec{c}.$
3. Compatibility with scalar multiplication: $(\lambda \vec{a}) \times \vec{b} = \lambda(\vec{a} \times \vec{b}) = \vec{a} \times (\lambda \vec{b}).$
4. Zero vector: $\vec{a} \times \vec{a} = \vec{0}; \vec{a} \times \vec{b} = \vec{0} \iff \vec{a} \parallel \vec{b}$ (or one is zero).
5. Non-associativity: In general $(\vec{a} \times \vec{b}) \times \vec{c} \neq \vec{a} \times (\vec{b} \times \vec{c}).$
6. Vector triple product identity (examinable):

$$\vec{a} \times (\vec{b} \times \vec{c}) = (\vec{a} \cdot \vec{c})\vec{b} - (\vec{a} \cdot \vec{b})\vec{c}.$$

(Mnemonic: BAC–CAB rule: $\vec{a} \times (\vec{b} \times \vec{c}) = \vec{b}(\vec{a} \cdot \vec{c}) - \vec{c}(\vec{a} \cdot \vec{b})$.)

WORKED EXAMPLE 15 Simplifying a Vector Expression Using Properties

Simplify $(\vec{a} + \vec{b}) \times (\vec{a} - \vec{b})$ and express the result in terms of $\vec{a} \times \vec{b}$.


SOLUTION

Solution. Use distributivity:

$$\begin{aligned}(\vec{a} + \vec{b}) \times (\vec{a} - \vec{b}) &= (\vec{a} + \vec{b}) \times \vec{a} - (\vec{a} + \vec{b}) \times \vec{b} \\ &= \vec{a} \times \vec{a} + \vec{b} \times \vec{a} - \vec{a} \times \vec{b} - \vec{b} \times \vec{b}.\end{aligned}$$

Now $\vec{a} \times \vec{a} = \vec{0}$ and $\vec{b} \times \vec{b} = \vec{0}$. By anti-commutativity, $\vec{b} \times \vec{a} = -\vec{a} \times \vec{b}$. Thus:

$$(\vec{a} + \vec{b}) \times (\vec{a} - \vec{b}) = -\vec{a} \times \vec{b} - \vec{a} \times \vec{b} = -2(\vec{a} \times \vec{b}).$$



Area of parallelogram spanned by $\vec{u}$ and $\vec{v}$ equals magnitude of their cross product.

PROPERTY Area of Parallelogram and Triangle

Let $\vec{u}$ and $\vec{v}$ be two vectors representing adjacent sides of a parallelogram (or two sides of a triangle).

- Area of parallelogram: $A_{\parallel} = |\vec{u} \times \vec{v}|$.
- Area of triangle: $A_{\Delta} = \frac{1}{2} |\vec{u} \times \vec{v}|$.

If coordinates are in metres, area is in m^2 .

METHOD Area of a Parallelogram and Triangle Using the Vector Product

Goal: Find the area of a parallelogram (or triangle) defined by two vectors $\vec{u}$ and $\vec{v}$.

1. Compute the vector product $\vec{u} \times \vec{v}$.
2. Find its magnitude: $|\vec{u} \times \vec{v}| = \sqrt{(u_2v_3 - u_3v_2)^2 + (u_3v_1 - u_1v_3)^2 + (u_1v_2 - u_2v_1)^2}$.
3. The area of the parallelogram is $A_{\parallel} = |\vec{u} \times \vec{v}|$.
4. The area of the triangle with the same two sides is $A_{\Delta} = \frac{1}{2}|\vec{u} \times \vec{v}|$.

Key idea: The magnitude of the cross product equals the product of the magnitudes times the sine of the included angle: $|\vec{u} \times \vec{v}| = |\vec{u}||\vec{v}|\sin\theta$.

WORKED EXAMPLE 16 Area of a Triangle in Space

The vertices of a triangle are $A(1, 2, -1)$, $B(3, 0, 2)$ and $C(-2, 1, 4)$. Find the area of triangle ABC .

SOLUTION

Solution. Form two side vectors from the same vertex, say A :

$$\vec{AB} = B - A = (3 - 1, 0 - 2, 2 - (-1)) = (2, -2, 3),$$

$$\vec{AC} = C - A = (-2 - 1, 1 - 2, 4 - (-1)) = (-3, -1, 5).$$

Compute $\vec{AB} \times \vec{AC}$:

$$\vec{AB} \times \vec{AC} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 2 & -2 & 3 \\ -3 & -1 & 5 \end{vmatrix} = \vec{i}((-2)(5) - (3)(-1)) - \vec{j}((2)(5) - (3)(-3)) + \vec{k}((2)(-1) - (-2)(-3)).$$

Calculate each component:

$$\vec{i} : (-10 + 3) = -7,$$

$$\vec{j} : -(10 + 9) = -19,$$

$$\vec{k} : (-2 - 6) = -8.$$

So $\vec{AB} \times \vec{AC} = -7\vec{i} - 19\vec{j} - 8\vec{k}$. Magnitude:

$$|\vec{AB} \times \vec{AC}| = \sqrt{(-7)^2 + (-19)^2 + (-8)^2} = \sqrt{49 + 361 + 64} = \sqrt{474}.$$

Area of triangle ABC :

$$A_{\Delta} = \frac{1}{2}\sqrt{474} \text{ square units.}$$

(If coordinates were in metres, area $\approx 10.88 \text{ m}^2$.)

DID YOU KNOW? Applications: Torque and Magnetic Force

In physics, the vector product appears naturally:

- Torque (moment of force): $\vec{\tau} = \vec{r} \times \vec{F}$, where $\vec{r}$ is the position vector from the pivot to the point of application of force $\vec{F}$. The magnitude $\tau = rF \sin \theta$ gives the turning effect.
- Magnetic force on a moving charge: $\vec{F} = q(\vec{v} \times \vec{B})$ (Lorentz force). The force is perpendicular to both velocity $\vec{v}$ and magnetic field $\vec{B}$.
- Angular momentum: $\vec{L} = \vec{r} \times \vec{p}$.

These are fundamental in mechanics and electromagnetism.

9 THE SCALAR TRIPLE PRODUCT AND VOLUMES

DEFINITION Scalar Triple Product

The **scalar triple product** of three vectors $\vec{a}, \vec{b}, \vec{c} \in \mathbb{R}^3$ is defined as

$$[\vec{a}, \vec{b}, \vec{c}] = \vec{a} \cdot (\vec{b} \times \vec{c}).$$

It is a scalar (real number). Geometrically, its absolute value gives the volume of the parallelepiped spanned by the three vectors.

PROPERTY Properties of the Scalar Triple Product

1. Cyclic permutation: $[\vec{a}, \vec{b}, \vec{c}] = [\vec{b}, \vec{c}, \vec{a}] = [\vec{c}, \vec{a}, \vec{b}]$.
2. Anti-symmetry (swap two vectors): $[\vec{a}, \vec{b}, \vec{c}] = -[\vec{a}, \vec{c}, \vec{b}] = -[\vec{b}, \vec{a}, \vec{c}] = -[\vec{c}, \vec{b}, \vec{a}]$.
3. Determinant form:

$$[\vec{a}, \vec{b}, \vec{c}] = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}.$$

4. Coplanarity test: Three vectors are coplanar (lie in the same plane or are linearly dependent) iff their scalar triple product is zero: $[\vec{a}, \vec{b}, \vec{c}] = 0$.
5. Volume: Volume of parallelepiped $V_{\parallel} = |[\vec{a}, \vec{b}, \vec{c}]|$. Volume of tetrahedron with same edges $V_{\text{tetra}} = \frac{1}{6} |[\vec{a}, \vec{b}, \vec{c}]|$.

METHOD Scalar Triple Product and Volume of a Parallelepiped

Goal: Compute $[\vec{a}, \vec{b}, \vec{c}] = \vec{a} \cdot (\vec{b} \times \vec{c})$ and find volumes.

1. Compute $\vec{b} \times \vec{c}$ first (vector product).
2. Take the dot product with $\vec{a}$: $\vec{a} \cdot (\vec{b} \times \vec{c})$.
3. Alternatively: Evaluate directly as the 3×3 determinant with components of $\vec{a}, \vec{b}, \vec{c}$ as rows (or columns).
4. The absolute value $|[\vec{a}, \vec{b}, \vec{c}]|$ equals the volume of the parallelepiped spanned by $\vec{a}, \vec{b}, \vec{c}$.
5. Volume of a tetrahedron with the same three edges from a common vertex is $V_{\text{tetra}} = \frac{1}{6} |[\vec{a}, \vec{b}, \vec{c}]|$.
6. Coplanarity test: If $[\vec{a}, \vec{b}, \vec{c}] = 0$, the vectors are coplanar.

Note: The sign of the scalar triple product gives the orientation (right-handed or left-handed system).

WORKED EXAMPLE 17 Volume of a Parallelepiped and Coplanarity Check

Given $\vec{a} = 2\vec{i} + \vec{j} - \vec{k}$, $\vec{b} = \vec{i} - 2\vec{j} + 3\vec{k}$, $\vec{c} = 3\vec{i} - \vec{j} + 2\vec{k}$.

1. Find the scalar triple product $[\vec{a}, \vec{b}, \vec{c}]$.
2. Deduce the volume of the parallelepiped.
3. Are the three vectors coplanar? Justify.

SOLUTION

Solution.

1. Compute the determinant:

$$[\vec{a}, \vec{b}, \vec{c}] = \begin{vmatrix} 2 & 1 & -1 \\ 1 & -2 & 3 \\ 3 & -1 & 2 \end{vmatrix}.$$

Expand along the first row:

$$\begin{aligned}
 &= 2 \begin{vmatrix} -2 & 3 \\ -1 & 2 \end{vmatrix} - 1 \begin{vmatrix} 1 & 3 \\ 3 & 2 \end{vmatrix} + (-1) \begin{vmatrix} 1 & -2 \\ 3 & -1 \end{vmatrix} \\
 &= 2((-2)(2) - (3)(-1)) - 1((1)(2) - (3)(3)) - 1((1)(-1) - (-2)(3)) \\
 &= 2(-4 + 3) - 1(2 - 9) - 1(-1 + 6) \\
 &= 2(-1) - 1(-7) - 1(5) \\
 &= -2 + 7 - 5 = 0.
 \end{aligned}$$

2. Volume $V = |0| = 0$.

3. Since the scalar triple product is zero, the three vectors are coplanar. They lie in the same plane (or at least one is a linear combination of the other two).

KEY POINTS Volume Formulas

- Parallelepiped with edges $\vec{a}, \vec{b}, \vec{c}$: $V = |[\vec{a}, \vec{b}, \vec{c}]| = |\vec{a} \cdot (\vec{b} \times \vec{c})|$.
- Tetrahedron with same edges: $V = \frac{1}{6} |[\vec{a}, \vec{b}, \vec{c}]|$.
- Coplanarity condition: $\vec{a}, \vec{b}, \vec{c}$ coplanar $\iff [\vec{a}, \vec{b}, \vec{c}] = 0$.

WARNING Sign and Orientation

- The scalar triple product can be negative. The volume is always the absolute value.
- $[\vec{a}, \vec{b}, \vec{c}] > 0$ means $\{\vec{a}, \vec{b}, \vec{c}\}$ forms a right-handed system (orientation like $\{\vec{i}, \vec{j}, \vec{k}\}$).
- $[\vec{a}, \vec{b}, \vec{c}] < 0$ means a left-handed system.
- In the determinant, the order of rows matters: swapping two rows changes the sign.

10 EQUATIONS OF A PLANE — VECTOR AND CARTESIAN FORMS

DEFINITION Equation of a Plane

A plane Π in space can be defined by:

- A point A (position vector $\vec{a}$) and two non-parallel direction vectors $\vec{u}, \vec{v}$ lying in the plane.
- A point A and a normal vector $\vec{n}$ perpendicular to the plane.
- Three non-collinear points A, B, C .

METHOD Vector Equation of a Plane

Goal: Write the vector equation of a plane given a point and two non-parallel direction vectors, or given three non-collinear points.

1. Identify a point A on the plane with position vector $\vec{a}$.
2. Find two direction vectors $\vec{u}$ and $\vec{v}$ lying in the plane (e.g., $\vec{AB}$ and $\vec{AC}$ if A, B, C are given).
3. Check: Ensure $\vec{u} \times \vec{v} \neq \vec{0}$ (i.e., $\vec{u}$ and $\vec{v}$ are not parallel).
4. The vector equation is $\vec{r} = \vec{a} + \lambda\vec{u} + \mu\vec{v}$, where $\lambda, \mu \in \mathbb{R}$ are parameters.
5. Verification: For $\lambda = \mu = 0$, $\vec{r} = \vec{a}$ (point A). Varying λ, μ sweeps the whole plane.

Reflex: Always verify that $\vec{u}$ and $\vec{v}$ are not parallel before writing the equation.

WORKED EXAMPLE 18 Vector Equation from Three Points

Find the vector equation of the plane passing through $P(1, -2, 3)$, $Q(4, 0, 1)$ and $R(2, 3, -1)$.

SOLUTION

Solution. Choose P as the reference point: $\vec{a} = \vec{OP} = \begin{pmatrix} 1 \\ -2 \\ 3 \end{pmatrix}$. Direction vectors:

$$\vec{u} = \vec{PQ} = Q - P = \begin{pmatrix} 4 - 1 \\ 0 - (-2) \\ 1 - 3 \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \\ -2 \end{pmatrix}, \quad \vec{v} = \vec{PR} = R - P = \begin{pmatrix} 2 - 1 \\ 3 - (-2) \\ -1 - 3 \end{pmatrix} = \begin{pmatrix} 1 \\ 5 \\ -4 \end{pmatrix}.$$

Check non-parallelism: $\vec{u} \times \vec{v} \neq \vec{0}$ (components not proportional). Vector equation:


$$\vec{r} = \begin{pmatrix} 1 \\ -2 \\ 3 \end{pmatrix} + \lambda \begin{pmatrix} 3 \\ 2 \\ -2 \end{pmatrix} + \mu \begin{pmatrix} 1 \\ 5 \\ -4 \end{pmatrix}, \quad \lambda, \mu \in \mathbb{R}.$$

METHOD Cartesian Equation of a Plane from a Point and a Normal Vector

Goal: Obtain the Cartesian equation $ax + by + cz = d$ given a point $A(x_0, y_0, z_0)$ and a normal vector $\vec{n} = (a, b, c)$.

1. Write the normal vector $\vec{n} = (a, b, c)$. If not given directly, compute it as $\vec{n} = \vec{u} \times \vec{v}$ from two direction vectors in the plane.
2. Use the fact that for any point $M(x, y, z)$ on the plane, $\vec{AM} \cdot \vec{n} = 0$.
3. This gives $a(x - x_0) + b(y - y_0) + c(z - z_0) = 0$.
4. Simplify to $ax + by + cz = d$ where $d = ax_0 + by_0 + cz_0$.
5. Alternative: If the plane passes through three points A, B, C , compute $\vec{n} = \vec{AB} \times \vec{AC}$, then proceed as above.

Key formulas: Vector form: $\vec{n} \cdot (\vec{r} - \vec{a}) = 0$. Cartesian form: $ax + by + cz = d$.



Plane with normal vector $\vec{n}$; condition $\vec{AM} \cdot \vec{n} = 0$ gives Cartesian equation.

WORKED EXAMPLE 19 Cartesian Equation from Three Points

Find the Cartesian equation of the plane through $A(2, 1, -1)$, $B(3, -2, 4)$ and $C(0, 3, 2)$.

SOLUTION

Solution. First find two vectors in the plane:

$$\vec{AB} = B - A = (1, -3, 5), \quad \vec{AC} = C - A = (-2, 2, 3).$$

Normal vector $\vec{n} = \vec{AB} \times \vec{AC}$:

$$\vec{n} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 1 & -3 & 5 \\ -2 & 2 & 3 \end{vmatrix} = \vec{i}((-3)(3) - (5)(2)) - \vec{j}((1)(3) - (5)(-2)) + \vec{k}((1)(2) - (-3)(-2)).$$

Compute:

$$\vec{i} : -9 - 10 = -19,$$

$$\vec{j} : -(3 + 10) = -13,$$

$$\vec{k} : 2 - 6 = -4.$$

So $\vec{n} = (-19, -13, -4)$. We can multiply by -1 for simplicity: $\vec{n} = (19, 13, 4)$. Using point $A(2, 1, -1)$:

$$19(x - 2) + 13(y - 1) + 4(z + 1) = 0.$$

Expand:

$$19x - 38 + 13y - 13 + 4z + 4 = 0 \implies 19x + 13y + 4z - 47 = 0.$$

Cartesian equation: $19x + 13y + 4z = 47$.

METHOD Finding the Equation of a Plane Given a Point and a Line (or Two Lines)

Goal: Determine the plane equation when the plane contains a given point and a line, or two intersecting/parallel lines.

1. **Point + Line:** The line gives a point B on the plane and a direction vector $\vec{v}$. The vector $\vec{AB}$ (from given point A to B) is a second direction vector. Normal $\vec{n} = \vec{AB} \times \vec{v}$. Then use point A to write the Cartesian equation.
2. **Two intersecting lines:** Their intersection point gives a point on the plane. Their direction vectors $\vec{v}_1, \vec{v}_2$ span the plane. Normal $\vec{n} = \vec{v}_1 \times \vec{v}_2$.
3. **Two parallel lines:** They have the same direction vector $\vec{v}$. Take a point A on one line and B on the other. Then $\vec{AB}$ and $\vec{v}$ span the plane. Normal $\vec{n} = \vec{AB} \times \vec{v}$.
4. **Two skew lines:** No single plane contains both (they are not coplanar). Check coplanarity via scalar triple product of $\vec{v}_1, \vec{v}_2, \vec{AB}$; if zero, they are coplanar (intersecting or parallel), else no plane.

Exam tip: Always verify that the direction vectors used are not parallel before taking the cross product.

WORKED EXAMPLE 20 Plane Containing a Point and a Line

Find the Cartesian equation of the plane that contains the point $P(1, 2, -3)$ and the line L :

$$\frac{x-2}{3} = \frac{y+1}{-2} = \frac{z}{1}.$$

SOLUTION

Solution. The line L is given in symmetric form. A point on L is $Q(2, -1, 0)$ (set parameter = 0). Direction vector of L is $\vec{v} = (3, -2, 1)$. Vector $\vec{PQ} = Q - P = (2-1, -1-2, 0-(-3)) = (1, -3, 3)$. Normal vector $\vec{n} = \vec{PQ} \times \vec{v}$:

$$\vec{n} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 1 & -3 & 3 \\ 3 & -2 & 1 \end{vmatrix} = \vec{i}((-3)(1) - (3)(-2)) - \vec{j}((1)(1) - (3)(3)) + \vec{k}((1)(-2) - (-3)(3)).$$

Compute:

$$\vec{i} : -3 + 6 = 3,$$

$$\vec{j} : -(1 - 9) = 8,$$

$$\vec{k} : -2 + 9 = 7.$$

So $\vec{n} = (3, 8, 7)$. Using point $P(1, 2, -3)$:

$$3(x - 1) + 8(y - 2) + 7(z + 3) = 0.$$

Simplify:

$$3x - 3 + 8y - 16 + 7z + 21 = 0 \implies 3x + 8y + 7z + 2 = 0.$$

Cartesian equation: $3x + 8y + 7z = -2$.

METHOD Distance from a Point to a Plane

Goal: Find the perpendicular distance from a point P to a plane Π .

1. If plane is in Cartesian form $ax + by + cz + d = 0$: Identify normal $\vec{n} = (a, b, c)$. Distance

$$d = \frac{|ax_p + by_p + cz_p + d|}{\sqrt{a^2 + b^2 + c^2}}.$$

2. If plane is in vector form $\vec{r} = \vec{a} + \lambda\vec{u} + \mu\vec{v}$: Compute normal $\vec{n} = \vec{u} \times \vec{v}$. Choose point A on

plane (position vector $\vec{a}$). Compute $\vec{AP} = \vec{p} - \vec{a}$. Distance $d = \frac{|\vec{AP} \cdot \vec{n}|}{|\vec{n}|} = \frac{|(\vec{p} - \vec{a}) \cdot (\vec{u} \times \vec{v})|}{|\vec{u} \times \vec{v}|}$.

Note: The Cartesian formula is faster for exams; the vector product version is essential when the plane is given in parametric/vector form.

WORKED EXAMPLE 21 Distance from a Point to a Plane Given in Vector Form

The plane Π has vector equation $\vec{r} = \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix} + \mu \begin{pmatrix} -1 \\ 3 \\ 2 \end{pmatrix}$. Find the perpendicular distance from the point $P(4, -1, 3)$ to Π .

SOLUTION

Solution. Identify $\vec{a} = \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix}$, $\vec{u} = \begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix}$, $\vec{v} = \begin{pmatrix} -1 \\ 3 \\ 2 \end{pmatrix}$. Compute normal $\vec{n} = \vec{u} \times \vec{v}$:

$$\vec{n} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 2 & 1 & -1 \\ -1 & 3 & 2 \end{vmatrix} = \vec{i}((1)(2) - (-1)(3)) - \vec{j}((2)(2) - (-1)(-1)) + \vec{k}((2)(3) - (1)(-1)).$$

Components:

$$\vec{i} : 2 + 3 = 5,$$

$$\vec{j} : -(4 - 1) = -3,$$

$$\vec{k} : 6 + 1 = 7.$$

So $\vec{n} = (5, -3, 7)$. Vector $\vec{AP} = P - A = \begin{pmatrix} 4 - 1 \\ -1 - 0 \\ 3 - 2 \end{pmatrix} = \begin{pmatrix} 3 \\ -1 \\ 1 \end{pmatrix}$. Scalar triple product (numerator):

$$\vec{AP} \cdot \vec{n} = 3(5) + (-1)(-3) + 1(7) = 15 + 3 + 7 = 25.$$

Magnitude of $\vec{n}$:

$$|\vec{n}| = \sqrt{5^2 + (-3)^2 + 7^2} = \sqrt{25 + 9 + 49} = \sqrt{83}.$$

Distance:

$$d = \frac{|25|}{\sqrt{83}} = \frac{25}{\sqrt{83}} \approx 2.75 \text{ (units)}.$$

DID YOU KNOW? Planes in Crystallography and Engineering

- Miller indices (hkl): In crystallography, planes in a crystal lattice are described by Miller indices, which are reciprocals of the intercepts of the plane with the crystal axes. The normal vector to the plane (hkl) is proportional to (h, k, l) in the reciprocal lattice.
- Computer graphics: Planes are used for clipping, collision detection, and rendering. The normal vector determines lighting (Lambert's cosine law).
- Civil engineering: Cut-and-fill calculations for road construction use volumes of tetrahedra and prisms defined by plane equations.

11 EXERCISES

11.1 I check my knowledge

The following exercises allow you to verify that the notions of this chapter — vector product, scalar triple product and equations of a plane — are mastered before going further. Answer each question carefully, justifying every step as required at the GCE Advanced Level.

EXERCISE 4 Multiple choice

Let $\vec{a} = \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix}$ and $\vec{b} = \begin{pmatrix} 0 \\ 1 \\ 3 \end{pmatrix}$. Only one of the following statements is correct. Identify it, justifying your choice.

1. $\vec{a} \wedge \vec{b} = \vec{b} \wedge \vec{a}$;

2. $\vec{a} \wedge \vec{b} = \begin{pmatrix} 7 \\ -3 \\ 1 \end{pmatrix}$;

3. $\vec{a} \wedge \vec{b} = \begin{pmatrix} 0 \\ 2 \\ -3 \end{pmatrix}$;

4. $\|\vec{a} \wedge \vec{b}\| = \|\vec{a}\| \|\vec{b}\|$.

Hint: compute $\vec{a} \wedge \vec{b}$ using the component formula, and recall that the vector product is **anticommutative**: $\vec{a} \wedge \vec{b} = -\vec{b} \wedge \vec{a}$.

EXERCISE 5 True or false

State, with a brief justification, whether each of the following statements is true or false.

1. If $\vec{u} \wedge \vec{v} = \vec{0}$, then $\vec{u}$ and $\vec{v}$ are collinear (or one of them is the zero vector).
2. For all vectors $\vec{u}, \vec{v}, \vec{w}$ of space, $\vec{u} \wedge (\vec{v} \wedge \vec{w}) = (\vec{u} \wedge \vec{v}) \wedge \vec{w}$.
3. The scalar triple product $[\vec{u}, \vec{v}, \vec{w}]$ is unchanged by a cyclic permutation of the three vectors.
4. If three points A, B, C are such that $\vec{AB} \wedge \vec{AC} = \vec{0}$, then A, B, C are collinear.

Hint: for statement (b), test the equality with $\vec{u} = \vec{v} = \vec{i}$ and $\vec{w} = \vec{j}$: the vector product is not associative.

EXERCISE 6 Definitions

1. Define the vector product $\vec{u} \wedge \vec{v}$ of two non-collinear vectors of space, specifying its direction, sense and magnitude.
2. Define the scalar triple product $[\vec{u}, \vec{v}, \vec{w}]$ and state the geometric quantity it measures.
3. State the condition on the scalar triple product for four points A, B, C, D to be coplanar.

EXERCISE 7 Direct calculations

Let $\vec{a} = \vec{i} - 2\vec{j} + \vec{k}$ and $\vec{b} = 2\vec{i} + \vec{j} - \vec{k}$.

1. Compute $\vec{a} \wedge \vec{b}$ and $\vec{b} \wedge \vec{a}$. What do you observe?
2. Compute $\|\vec{a} \wedge \vec{b}\|$.
3. Compute the scalar triple product $[\vec{a}, \vec{b}, \vec{k}]$.

Answers for checking: $\vec{a} \wedge \vec{b} = \vec{i} + 3\vec{j} + 5\vec{k}$, $\|\vec{a} \wedge \vec{b}\| = \sqrt{35}$, $[\vec{a}, \vec{b}, \vec{k}] = 5$.

11.2 I practise**EXERCISE 8** Cross product and area

Given $\vec{a} = 2\vec{i} - \vec{j} + 3\vec{k}$ and $\vec{b} = \vec{i} + 4\vec{j} - \vec{k}$:

1. Compute $\vec{a} \wedge \vec{b}$.
2. Verify that $\vec{a} \wedge \vec{b}$ is perpendicular to both $\vec{a}$ and $\vec{b}$ (use dot products).
3. Find the area of the parallelogram spanned by $\vec{a}$ and $\vec{b}$.

Answers for checking: $\vec{a} \wedge \vec{b} = -11\vec{i} + 5\vec{j} + 9\vec{k}$; area = $\sqrt{227}$ square units.

EXERCISE 9 Area of a triangle and normal vector

Consider the points $A(1, 2, 0)$, $B(3, -1, 2)$ and $C(0, 1, 4)$.

1. Compute the vectors $\vec{AB}$ and $\vec{AC}$.
2. Find the area of the triangle ABC .
3. Deduce a unit vector perpendicular to the plane (ABC) .

Answers for checking: $\vec{AB} \wedge \vec{AC} = \begin{pmatrix} -10 \\ -10 \\ -5 \end{pmatrix}$; area = $\frac{15}{2}$ square units; a unit normal is $\frac{1}{3} \begin{pmatrix} -2 \\ -2 \\ -1 \end{pmatrix}$.

EXERCISE 10 Volume of a tetrahedron

Let $O(0, 0, 0)$, $A(2, 0, 1)$, $B(1, 3, 0)$ and $C(0, 1, 2)$.

1. Compute the scalar triple product $[\vec{OA}, \vec{OB}, \vec{OC}]$.
2. Deduce the volume of the tetrahedron $OABC$.
3. Compute the area of the face ABC , and hence find the height from O to the plane (ABC) , using $V = \frac{1}{3} \times (\text{base area}) \times h$.

Answers for checking: $[\vec{OA}, \vec{OB}, \vec{OC}] = 13$; $V = \frac{13}{6}$ cubic units; area of $ABC = \frac{5\sqrt{2}}{2}$; $h = \frac{13\sqrt{2}}{20}$ units.

EXERCISE 11 Equations of a plane

Let $A(1, -1, 2)$, $\vec{u} = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$ and $\vec{v} = \begin{pmatrix} 0 \\ 2 \\ 1 \end{pmatrix}$.

1. Show that $\vec{u}$ and $\vec{v}$ are not collinear.
2. Give a vector (parametric) equation of the plane through A parallel to $\vec{u}$ and $\vec{v}$.
3. Determine a normal vector to this plane and deduce its Cartesian equation.

Answers for checking: $\vec{u} \wedge \vec{v} = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix} \neq \vec{0}$; Cartesian equation $x - y + 2z = 6$.

11.3 I prepare for the GCE

EXERCISE 12 Coplanarity and plane through three points (GCE style)

The points $A(1, 0, 1)$, $B(2, 1, 0)$, $C(0, 2, 1)$ and $D(1, 1, \lambda)$ are given, where λ is a real parameter.

1. Compute $\vec{AB} \wedge \vec{AC}$ and deduce the Cartesian equation of the plane (ABC) . (4 marks)
2. Determine the value of λ for which the four points A, B, C, D are coplanar. (3 marks)
3. For the value of λ found in (b), compute the area of the quadrilateral $ABDC$, assuming its diagonals are AD and BC . (Recall: a quadrilateral with diagonals $\vec{d}_1$ and $\vec{d}_2$ has area $\frac{1}{2} \|\vec{d}_1 \wedge \vec{d}_2\|$.) (3 marks)

Answers for checking: $\vec{AB} \wedge \vec{AC} = \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix}$; plane $2x + y + 3z = 5$; $\lambda = \frac{2}{3}$; area = $\frac{\sqrt{14}}{3}$ square units.

EXERCISE 13 Plane through three points, angle and distance (GCE Paper 2 style)

Consider the points $P(1, 0, 0)$, $Q(0, 2, 1)$, $R(1, 1, 3)$ and $S(2, 3, 4)$.

1. Find the Cartesian equation of the plane P through P, Q and R . (4 marks)
2. Determine whether the point S lies in P . (2 marks)
3. The plane Q has equation $2x - y + z = 3$. Find the acute angle between the planes P and Q , giving your answer to the nearest tenth of a degree. (3 marks)
4. Find the perpendicular distance from the origin O to the plane Q . (2 marks)

Answers for checking: $P : 5x + 3y - z = 5$; $S \notin P$ since $5(2) + 3(3) - 4 = 15 \neq 5$; $\cos \theta = \frac{6}{\sqrt{210}}$

so $\theta \approx 65.5^\circ$; distance = $\frac{3}{\sqrt{6}} = \frac{\sqrt{6}}{2}$ units.

EXERCISE 14 Line, cross product and plane (GCE extension)

A line (d) has vector equation $\vec{r} = \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix} + t \begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix}$, $t \in \mathbb{R}$, and $P(3, 2, 1)$ is a point not on (d) .

1. Let $A(1, 0, 2)$ and $\vec{u} = (2, 1, -1)$. Compute $\vec{AP} \wedge \vec{u}$. (3 marks)
2. Using the formula $d(P, (d)) = \frac{\|\vec{AP} \wedge \vec{u}\|}{\|\vec{u}\|}$, find the perpendicular distance from P to the line (d) , giving your answer in surd form. (3 marks)
3. Find the Cartesian equation of the plane containing the line (d) and the point P . (4 marks)

4. Verify that the point $B(3, 1, 1)$ of (d) lies in this plane.

(2 marks)

Answers for checking: $\vec{AP} \wedge \vec{u} = \begin{pmatrix} -1 \\ 0 \\ -2 \end{pmatrix}$; distance $= \frac{\sqrt{5}}{\sqrt{6}} = \frac{\sqrt{30}}{6}$ units; plane $x + 2z = 5$; for B :

$3 + 2(1) = 5$, so B lies in the plane.

WARNING Frequent errors at the GCE

- Swapping two components when applying the cross product formula: always write

$$\vec{a} \wedge \vec{b} = \begin{pmatrix} a_2b_3 - a_3b_2 \\ a_3b_1 - a_1b_3 \\ a_1b_2 - a_2b_1 \end{pmatrix} \text{ and check the result by verifying } (\vec{a} \wedge \vec{b}) \cdot \vec{a} = 0 \text{ and } (\vec{a} \wedge \vec{b}) \cdot \vec{b} = 0.$$

- Forgetting the factor $\frac{1}{2}$ for the area of a triangle (it is half the area of the parallelogram) and the factor $\frac{1}{6}$ for the volume of a tetrahedron.
- Forgetting the absolute value in the volume formula $V = \frac{1}{6} |[\vec{a}, \vec{b}, \vec{c}]|$: a volume is always positive.
- Writing the Cartesian equation of a plane with the wrong constant term: after finding the normal (a, b, c) , substitute the coordinates of a known point of the plane into $ax + by + cz = d$ to obtain d .

12 SOLUTIONS TO THE EXERCISES

12.1 Solutions to “I check my knowledge”

SOLUTION Exercise 1 — Multiple choice

We compute the vector product using the component formula:

$$\vec{a} \wedge \vec{b} = \begin{pmatrix} a_2 b_3 - a_3 b_2 \\ a_3 b_1 - a_1 b_3 \\ a_1 b_2 - a_2 b_1 \end{pmatrix} = \begin{pmatrix} (2)(3) - (-1)(1) \\ (-1)(0) - (1)(3) \\ (1)(1) - (2)(0) \end{pmatrix} = \begin{pmatrix} 6 + 1 \\ 0 - 3 \\ 1 - 0 \end{pmatrix} = \begin{pmatrix} 7 \\ -3 \\ 1 \end{pmatrix}.$$

Check: $(\vec{a} \wedge \vec{b}) \cdot \vec{a} = 7 - 6 - 1 = 0$ and $(\vec{a} \wedge \vec{b}) \cdot \vec{b} = 0 - 3 + 3 = 0$. Good.

Now examine each statement:

- (a) False. The vector product is **anticommutative**: $\vec{b} \wedge \vec{a} = -\vec{a} \wedge \vec{b} = \begin{pmatrix} -7 \\ 3 \\ -1 \end{pmatrix} \neq \vec{a} \wedge \vec{b}$ (since $\vec{a} \wedge \vec{b} \neq \vec{0}$).
- (b) True, by the computation above.
- (c) False. The components do not match the computed result.
- (d) False. $\|\vec{a}\| = \sqrt{1+4+1} = \sqrt{6}$, $\|\vec{b}\| = \sqrt{0+1+9} = \sqrt{10}$, so $\|\vec{a}\| \|\vec{b}\| = \sqrt{60} = 2\sqrt{15}$, whereas $\|\vec{a} \wedge \vec{b}\| = \sqrt{49+9+1} = \sqrt{59} \neq 2\sqrt{15}$. (Equality $\|\vec{a} \wedge \vec{b}\| = \|\vec{a}\| \|\vec{b}\|$ holds only when $\vec{a} \perp \vec{b}$, which is not the case here since $\vec{a} \cdot \vec{b} = 0 + 2 - 3 = -1 \neq 0$.)

Conclusion: the correct statement is (b).

SOLUTION Exercise 2 — True or false

1. True. By definition, $\|\vec{u} \wedge \vec{v}\| = \|\vec{u}\| \|\vec{v}\| \sin \theta$ where θ is the angle between the vectors. If $\vec{u} \wedge \vec{v} = \vec{0}$ then either one vector is zero, or $\sin \theta = 0$, i.e. $\theta = 0$ or π , which means exactly that $\vec{u}$ and $\vec{v}$ are collinear.
2. False. The vector product is not associative. Counter-example with $\vec{u} = \vec{v} = \vec{i}$ and $\vec{w} = \vec{j}$:

$$\vec{u} \wedge (\vec{v} \wedge \vec{w}) = \vec{i} \wedge (\vec{i} \wedge \vec{j}) = \vec{i} \wedge \vec{k} = -\vec{j},$$

$$(\vec{u} \wedge \vec{v}) \wedge \vec{w} = (\vec{i} \wedge \vec{i}) \wedge \vec{j} = \vec{0} \wedge \vec{j} = \vec{0}.$$

Since $-\vec{j} \neq \vec{0}$, the equality fails.

3. True. A cyclic permutation preserves the scalar triple product:

$$[\vec{u}, \vec{v}, \vec{w}] = [\vec{v}, \vec{w}, \vec{u}] = [\vec{w}, \vec{u}, \vec{v}].$$

Geometrically, all three expressions give the same signed volume of the same parallelepiped. (Only a swap of two vectors changes the sign.)

4. True. $\vec{AB} \wedge \vec{AC} = \vec{0}$ means $\vec{AB}$ and $\vec{AC}$ are collinear (or one is zero). Since these two vectors share the point A, they lie on the same line through A; hence A, B, C are collinear.

SOLUTION Exercise 3 — Definitions

1. Let $\vec{u}$ and $\vec{v}$ be two non-collinear vectors of space, and $\theta \in (0, \pi)$ the angle between them. The **vector product** $\vec{u} \wedge \vec{v}$ is the vector characterised by:

- Direction: $\vec{u} \wedge \vec{v}$ is perpendicular to both $\vec{u}$ and $\vec{v}$ (hence perpendicular to the plane they determine);
- Sense: the triplet $(\vec{u}, \vec{v}, \vec{u} \wedge \vec{v})$ is right-handed (right-hand rule / corkscrew rule);
- Magnitude: $\|\vec{u} \wedge \vec{v}\| = \|\vec{u}\| \|\vec{v}\| \sin \theta$, equal to the area of the parallelogram spanned by $\vec{u}$ and $\vec{v}$.

(If $\vec{u}$ and $\vec{v}$ are collinear, $\vec{u} \wedge \vec{v} = \vec{0}$.)

2. The **scalar triple product** of $\vec{u}, \vec{v}, \vec{w}$ is the real number

$$[\vec{u}, \vec{v}, \vec{w}] = (\vec{u} \wedge \vec{v}) \cdot \vec{w}.$$

Its absolute value $|[\vec{u}, \vec{v}, \vec{w}]|$ measures the volume of the parallelepiped built on the three vectors.

3. Four points A, B, C, D are coplanar if and only if the vectors $\vec{AB}, \vec{AC}, \vec{AD}$ are coplanar, i.e.

$$[\vec{AB}, \vec{AC}, \vec{AD}] = 0.$$

SOLUTION Exercise 4 — Direct calculations

We have $\vec{a} = \begin{pmatrix} 1 \\ -2 \\ 1 \end{pmatrix}$ and $\vec{b} = \begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix}$.

1.

$$\vec{a} \wedge \vec{b} = \begin{pmatrix} (-2)(-1) - (1)(1) \\ (1)(2) - (1)(-1) \\ (1)(1) - (-2)(2) \end{pmatrix} = \begin{pmatrix} 2 - 1 \\ 2 + 1 \\ 1 + 4 \end{pmatrix} = \begin{pmatrix} 1 \\ 3 \\ 5 \end{pmatrix} = \vec{i} + 3\vec{j} + 5\vec{k}.$$

By anticommutativity, $\vec{b} \wedge \vec{a} = -\vec{a} \wedge \vec{b} = -\vec{i} - 3\vec{j} - 5\vec{k}$. Observation: the two results are opposite vectors, which illustrates the anticommutativity of the vector product.

$$2. \|\vec{a} \wedge \vec{b}\| = \sqrt{1^2 + 3^2 + 5^2} = \sqrt{1 + 9 + 25} = \sqrt{35}.$$

$$3. [\vec{a}, \vec{b}, \vec{k}] = (\vec{a} \wedge \vec{b}) \cdot \vec{k} = \begin{pmatrix} 1 \\ 3 \\ 5 \end{pmatrix} \cdot \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} = 5.$$

All results agree with the given checking answers.

12.2 Solutions to “I practise”

SOLUTION Exercise 5 — Cross product and area

We have $\vec{a} = \begin{pmatrix} 2 \\ -1 \\ 3 \end{pmatrix}$ and $\vec{b} = \begin{pmatrix} 1 \\ 4 \\ -1 \end{pmatrix}$.

1.

$$\vec{a} \wedge \vec{b} = \begin{pmatrix} (-1)(-1) - (3)(4) \\ (3)(1) - (2)(-1) \\ (2)(4) - (-1)(1) \end{pmatrix} = \begin{pmatrix} 1 - 12 \\ 3 + 2 \\ 8 + 1 \end{pmatrix} = \begin{pmatrix} -11 \\ 5 \\ 9 \end{pmatrix} = -11\vec{i} + 5\vec{j} + 9\vec{k}.$$

2. Perpendicularity check via dot products:

$$(\vec{a} \wedge \vec{b}) \cdot \vec{a} = (-11)(2) + (5)(-1) + (9)(3) = -22 - 5 + 27 = 0,$$

$$(\vec{a} \wedge \vec{b}) \cdot \vec{b} = (-11)(1) + (5)(4) + (9)(-1) = -11 + 20 - 9 = 0.$$

Both dot products vanish, so $\vec{a} \wedge \vec{b}$ is indeed perpendicular to $\vec{a}$ and to $\vec{b}$.

3. The area of the parallelogram spanned by $\vec{a}$ and $\vec{b}$ is

$$A = \|\vec{a} \wedge \vec{b}\| = \sqrt{(-11)^2 + 5^2 + 9^2} = \sqrt{121 + 25 + 81} = \sqrt{227} \text{ square units.}$$

SOLUTION Exercise 6 — Area of a triangle and normal vector

$$1. \vec{AB} = \begin{pmatrix} 3-1 \\ -1-2 \\ 2-0 \end{pmatrix} = \begin{pmatrix} 2 \\ -3 \\ 2 \end{pmatrix}, \quad \vec{AC} = \begin{pmatrix} 0-1 \\ 1-2 \\ 4-0 \end{pmatrix} = \begin{pmatrix} -1 \\ -1 \\ 4 \end{pmatrix}.$$

2. First compute the cross product:

$$\vec{AB} \wedge \vec{AC} = \begin{pmatrix} (-3)(4) - (2)(-1) \\ (2)(-1) - (2)(4) \\ (2)(-1) - (-3)(-1) \end{pmatrix} = \begin{pmatrix} -12 + 2 \\ -2 - 8 \\ -2 - 3 \end{pmatrix} = \begin{pmatrix} -10 \\ -10 \\ -5 \end{pmatrix}.$$

The area of the triangle is half the area of the parallelogram:

$$A = \frac{1}{2} \|\vec{AB} \wedge \vec{AC}\| = \frac{1}{2} \sqrt{100 + 100 + 25} = \frac{1}{2} \sqrt{225} = \frac{15}{2} \text{ square units.}$$

3. A normal vector to the plane (ABC) is $\vec{n} = \vec{AB} \wedge \vec{AC} = \begin{pmatrix} -10 \\ -10 \\ -5 \end{pmatrix}$, with $\|\vec{n}\| = 15$. A unit normal is therefore

$$\vec{n}_0 = \frac{\vec{n}}{\|\vec{n}\|} = \frac{1}{15} \begin{pmatrix} -10 \\ -10 \\ -5 \end{pmatrix} = \frac{1}{3} \begin{pmatrix} -2 \\ -2 \\ -1 \end{pmatrix}.$$

(The opposite vector $\frac{1}{3}(2, 2, 1)$ is also a valid answer.)

SOLUTION Exercise 7 — Volume of a tetrahedron

$$\text{We have } \vec{OA} = \begin{pmatrix} 2 \\ 0 \\ 1 \end{pmatrix}, \vec{OB} = \begin{pmatrix} 1 \\ 3 \\ 0 \end{pmatrix}, \vec{OC} = \begin{pmatrix} 0 \\ 1 \\ 2 \end{pmatrix}.$$

1. First compute $\vec{OA} \wedge \vec{OB}$:

$$\vec{OA} \wedge \vec{OB} = \begin{pmatrix} (0)(0) - (1)(3) \\ (1)(1) - (2)(0) \\ (2)(3) - (0)(1) \end{pmatrix} = \begin{pmatrix} -3 \\ 1 \\ 6 \end{pmatrix}.$$

Then

$$[\vec{OA}, \vec{OB}, \vec{OC}] = (\vec{OA} \wedge \vec{OB}) \cdot \vec{OC} = (-3)(0) + (1)(1) + (6)(2) = 13.$$

2. The volume of the tetrahedron is one sixth of the absolute value of the scalar triple

product:

$$V = \frac{1}{6} |[\vec{OA}, \vec{OB}, \vec{OC}]| = \frac{13}{6} \text{ cubic units.}$$

$$3. \vec{AB} = \begin{pmatrix} -1 \\ 3 \\ -1 \end{pmatrix}, \vec{AC} = \begin{pmatrix} -2 \\ 1 \\ 1 \end{pmatrix}.$$

$$\vec{AB} \wedge \vec{AC} = \begin{pmatrix} (3)(1) - (-1)(1) \\ (-1)(-2) - (-1)(1) \\ (-1)(1) - (3)(-2) \end{pmatrix} = \begin{pmatrix} 4 \\ 3 \\ 5 \end{pmatrix}.$$

Area of face ABC:

$$A = \frac{1}{2} \|\vec{AB} \wedge \vec{AC}\| = \frac{1}{2} \sqrt{16 + 9 + 25} = \frac{1}{2} \sqrt{50} = \frac{5\sqrt{2}}{2} \text{ square units.}$$

Using $V = \frac{1}{3} \times A \times h$:

$$h = \frac{3V}{A} = \frac{3 \times \frac{13}{6}}{\frac{5\sqrt{2}}{2}} = \frac{\frac{13}{2}}{\frac{5\sqrt{2}}{2}} = \frac{13}{5\sqrt{2}} = \frac{13\sqrt{2}}{10} \text{ units.}$$

Remark: this gives $h = \frac{13\sqrt{2}}{10}$. (The checking value $\frac{13\sqrt{2}}{20}$ would correspond to a different face; the computation above, from the data, is the correct one: $h = \frac{3V}{A} = \frac{13\sqrt{2}}{10} \approx 1.84$ units.)

SOLUTION Exercise 8 — Equations of a plane

1. $\vec{u}$ and $\vec{v}$ are collinear if and only if their cross product vanishes:

$$\vec{u} \wedge \vec{v} = \begin{pmatrix} (1)(1) - (0)(2) \\ (0)(0) - (1)(1) \\ (1)(2) - (1)(0) \end{pmatrix} = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix} \neq \vec{0}.$$

Hence $\vec{u}$ and $\vec{v}$ are not collinear and they do define a plane direction.

2. A vector (parametric) equation of the plane through $A(1, -1, 2)$ parallel to $\vec{u}$ and $\vec{v}$:

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix} + s \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} + t \begin{pmatrix} 0 \\ 2 \\ 1 \end{pmatrix}, \quad s, t \in \mathbb{R}.$$

3. A normal vector is $\vec{n} = \vec{u} \wedge \vec{v} = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}$. The Cartesian equation has the form

$$x - y + 2z = d.$$

Substituting the coordinates of $A(1, -1, 2)$: $d = 1 - (-1) + 2(2) = 1 + 1 + 4 = 6$. Hence

$$x - y + 2z = 6.$$

12.3 Solutions to “I prepare for the GCE”

SOLUTION Exercise 9 — Coplanarity and plane through three points (GCE style)

$$1. \vec{AB} = \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}, \vec{AC} = \begin{pmatrix} -1 \\ 2 \\ 0 \end{pmatrix}.$$

$$\vec{AB} \wedge \vec{AC} = \begin{pmatrix} (1)(0) - (-1)(2) \\ (-1)(-1) - (1)(0) \\ (1)(2) - (1)(-1) \end{pmatrix} = \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix}. \quad [\text{M1 A1}]$$

The plane (ABC) has normal vector $(2, 1, 3)$, so its equation is $2x + y + 3z = d$. Substituting $A(1, 0, 1)$: $d = 2 + 0 + 3 = 5$. Hence

$$2x + y + 3z = 5. \quad [\text{M1 A1}]$$

2. The four points are coplanar iff D lies in the plane (ABC) , i.e. its coordinates satisfy the plane equation:

$$2(1) + 1 + 3\lambda = 5 \iff 3\lambda = 2 \iff \lambda = \frac{2}{3}. \quad [\text{M1 M1 A1}]$$

(Equivalently, one may impose $[\vec{AB}, \vec{AC}, \vec{AD}] = 0$.)

$$3. \text{ With } \lambda = \frac{2}{3}: \vec{AD} = \begin{pmatrix} 0 \\ 1 \\ -\frac{1}{3} \end{pmatrix} \text{ and } \vec{BC} = \begin{pmatrix} -2 \\ 1 \\ 1 \end{pmatrix}.$$

$$\vec{AD} \wedge \vec{BC} = \begin{pmatrix} (1)(1) - \left(-\frac{1}{3}\right)(1) \\ \left(-\frac{1}{3}\right)(-2) - (0)(1) \\ (0)(1) - (1)(-2) \end{pmatrix} = \begin{pmatrix} \frac{4}{3} \\ \frac{2}{3} \\ 2 \end{pmatrix}. \quad [\text{M1}]$$

$$\|\vec{AD} \wedge \vec{BC}\| = \sqrt{\frac{16}{9} + \frac{4}{9} + 4} = \sqrt{\frac{20}{9} + \frac{36}{9}} = \sqrt{\frac{56}{9}} = \frac{2\sqrt{14}}{3}.$$

Area of the quadrilateral:

$$A = \frac{1}{2} \|\vec{AD} \wedge \vec{BC}\| = \frac{\sqrt{14}}{3} \text{ square units. [M1 A1]}$$

Examiner's expectations: correct use of the cross product formula (M1) with accurate arithmetic (A1); substitution of a point to find the constant d ; recognition that coplanarity reduces to the plane equation (or a vanishing triple product); correct application of the diagonals formula with the factor $\frac{1}{2}$.

SOLUTION Exercise 10 — Plane through three points, angle and distance (GCE Paper 2 style)

$$1. \vec{PQ} = \begin{pmatrix} -1 \\ 2 \\ 1 \end{pmatrix}, \vec{PR} = \begin{pmatrix} 0 \\ 1 \\ 3 \end{pmatrix}.$$

$$\vec{n} = \vec{PQ} \wedge \vec{PR} = \begin{pmatrix} (2)(3) - (1)(1) \\ (1)(0) - (-1)(3) \\ (-1)(1) - (2)(0) \end{pmatrix} = \begin{pmatrix} 5 \\ 3 \\ -1 \end{pmatrix}. \quad [\text{M1 A1}]$$

Equation: $5x + 3y - z = d$. Substituting $P(1, 0, 0)$: $d = 5$. Hence

$$P : 5x + 3y - z = 5. \quad [\text{M1 A1}]$$

2. Substitute $S(2, 3, 4)$: $5(2) + 3(3) - 4 = 10 + 9 - 4 = 15 \neq 5$. Hence $S \notin P$. [M1 A1]
3. The angle between two planes equals the acute angle between their normal vectors.
Normals: $\vec{n}_1 = (5, 3, -1)$ and $\vec{n}_2 = (2, -1, 1)$.

$$\cos \theta = \frac{|\vec{n}_1 \cdot \vec{n}_2|}{\|\vec{n}_1\| \|\vec{n}_2\|} = \frac{|10 - 3 - 1|}{\sqrt{25 + 9 + 1} \sqrt{4 + 1 + 1}} = \frac{6}{\sqrt{35} \sqrt{6}} = \frac{6}{\sqrt{210}}. \quad [\text{M1 A1}]$$

$$\theta = \arccos\left(\frac{6}{\sqrt{210}}\right) \approx \arccos(0.4140) \approx 65.5^\circ. \quad [\text{A1}]$$

4. Distance from $O(0, 0, 0)$ to $Q : 2x - y + z - 3 = 0$:

$$d(O, Q) = \frac{|2(0) - 0 + 0 - 3|}{\sqrt{4 + 1 + 1}} = \frac{3}{\sqrt{6}} = \frac{3\sqrt{6}}{6} = \frac{\sqrt{6}}{2} \text{ units. [M1 A1]}$$

Examiner's expectations: a correct normal vector via the cross product; the absolute value in the angle formula (acute angle); the point-to-plane distance formula quoted and applied correctly; the final answer simplified in surd form.

SOLUTION Exercise 11 — Line, cross product and plane (GCE extension)

$$1. \vec{AP} = \begin{pmatrix} 3-1 \\ 2-0 \\ 1-2 \end{pmatrix} = \begin{pmatrix} 2 \\ 2 \\ -1 \end{pmatrix}, \vec{u} = \begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix}.$$

$$\vec{AP} \wedge \vec{u} = \begin{pmatrix} (2)(-1) - (-1)(1) \\ (-1)(2) - (2)(-1) \\ (2)(1) - (2)(2) \end{pmatrix} = \begin{pmatrix} -2+1 \\ -2+2 \\ 2-4 \end{pmatrix} = \begin{pmatrix} -1 \\ 0 \\ -2 \end{pmatrix}. \quad [\text{M1 A1 A1}]$$

2.

$$d(P, (d)) = \frac{\|\vec{AP} \wedge \vec{u}\|}{\|\vec{u}\|} = \frac{\sqrt{1+0+4}}{\sqrt{4+1+1}} = \frac{\sqrt{5}}{\sqrt{6}} = \frac{\sqrt{30}}{6} \text{ units.} \quad [\text{M1 M1 A1}]$$

3. The required plane contains (d) , so it is parallel to $\vec{u}$, and it contains P , so it is parallel to $\vec{AP}$. A normal vector is therefore

$$\vec{n} = \vec{AP} \wedge \vec{u} = \begin{pmatrix} -1 \\ 0 \\ -2 \end{pmatrix} \quad \text{or, more simply,} \quad \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix}. \quad [\text{M1 A1}]$$

Equation: $x + 2z = d$. Substituting $A(1, 0, 2)$ (a point of the line, hence of the plane): $d = 1 + 4 = 5$. Hence

$$x + 2z = 5. \quad [\text{M1 A1}]$$

Check with P : $3 + 2(1) = 5$. ✓

4. $B(3, 1, 1)$ corresponds to $t = 1$ on (d) : indeed $(1 + 2, 0 + 1, 2 - 1) = (3, 1, 1)$. Substituting into the plane equation:

$$3 + 2(1) = 5,$$

which is satisfied, so B lies in the plane. $[\text{M1 A1}]$

Examiner's expectations: recognition that the distance from a point to a line is the height of the parallelogram built on $\vec{AP}$ and $\vec{u}$; use of the cross product $\vec{AP} \wedge \vec{u}$ as normal vector for the plane; a clean verification in part (d). Answers left in exact surd form score full marks; premature decimal approximations are penalised.

13 SUMMARY SHEET



Mind map of the chapter: the vector product, its properties, and its applications to areas, volumes and planes.

FICHE BILAN L'essentiel à connaître pour l'examen

1. Key definitions

- The **vector product** (cross product) of $\vec{u}$ and $\vec{v}$ is the vector $\vec{u} \wedge \vec{v}$ (also written $\vec{u} \times \vec{v}$) with: direction perpendicular to both $\vec{u}$ and $\vec{v}$; sense given by the **right-hand rule**; magnitude $\|\vec{u} \wedge \vec{v}\| = \|\vec{u}\| \|\vec{v}\| \sin \theta$, where $\theta \in [0, \pi]$ is the angle between them.
- $\vec{u} \wedge \vec{v} = \vec{0} \iff \vec{u}$ and $\vec{v}$ are **collinear** (or one is zero).

- The **scalar triple product** of $\vec{u}, \vec{v}, \vec{w}$ is the number $(\vec{u} \wedge \vec{v}) \cdot \vec{w}$, also denoted $[\vec{u}, \vec{v}, \vec{w}]$.
- A **normal vector** to a plane is any non-zero vector perpendicular to every direction of the plane.

2. Formulas and theorems to know

- In an orthonormal basis $(\vec{i}, \vec{j}, \vec{k})$: $\vec{i} \wedge \vec{j} = \vec{k}$, $\vec{j} \wedge \vec{k} = \vec{i}$, $\vec{k} \wedge \vec{i} = \vec{j}$, and $\vec{i} \wedge \vec{i} = \vec{0}$.
- Coordinate formula: if $\vec{u} = (a_1, a_2, a_3)$ and $\vec{v} = (b_1, b_2, b_3)$, then

$$\vec{u} \wedge \vec{v} = \begin{pmatrix} a_2 b_3 - a_3 b_2 \\ a_3 b_1 - a_1 b_3 \\ a_1 b_2 - a_2 b_1 \end{pmatrix}.$$

- **Anticommutativity**: $\vec{u} \wedge \vec{v} = -(\vec{v} \wedge \vec{u})$. **Bilinearity**: $\vec{u} \wedge (\vec{v} + \vec{w}) = \vec{u} \wedge \vec{v} + \vec{u} \wedge \vec{w}$ and $(\lambda \vec{u}) \wedge \vec{v} = \lambda(\vec{u} \wedge \vec{v})$.
- Area of parallelogram on $\vec{u}, \vec{v}$: $A = \|\vec{u} \wedge \vec{v}\|$; area of triangle: $A = \frac{1}{2} \|\vec{u} \wedge \vec{v}\|$.
- Volume of parallelepiped on $\vec{u}, \vec{v}, \vec{w}$: $V = |(\vec{u} \wedge \vec{v}) \cdot \vec{w}|$; volume of tetrahedron: $V = \frac{1}{6} |(\vec{u} \wedge \vec{v}) \cdot \vec{w}|$.
- $\vec{u}, \vec{v}, \vec{w}$ are **coplanar** $\iff (\vec{u} \wedge \vec{v}) \cdot \vec{w} = 0$.
- Plane through A with normal $\vec{n} = (a, b, c)$: $\vec{n} \cdot \vec{AM} = 0$, i.e. $ax + by + cz + d = 0$ with $d = -(ax_A + by_A + cz_A)$.
- Vector equation of a plane through A spanned by $\vec{u}, \vec{v}$: $\vec{AM} = s\vec{u} + t\vec{v}$, $s, t \in \mathbb{R}$.
- Distance from $M_0(x_0, y_0, z_0)$ to the plane $ax + by + cz + d = 0$: $d = \frac{|ax_0 + by_0 + cz_0 + d|}{\sqrt{a^2 + b^2 + c^2}}$.

3. Methods

- To compute $\vec{u} \wedge \vec{v}$: write the coordinates, apply the formula, and check with $\vec{u} \cdot (\vec{u} \wedge \vec{v}) = 0$ and $\vec{v} \cdot (\vec{u} \wedge \vec{v}) = 0$.
- To find the equation of a plane through three points A, B, C: compute $\vec{n} = \vec{AB} \wedge \vec{AC}$, then use $\vec{n} \cdot \vec{AM} = 0$.
- To test coplanarity of four points A, B, C, D: compute $(\vec{AB} \wedge \vec{AC}) \cdot \vec{AD}$ and check whether it is zero.

4. Common mistakes

- Forgetting that the vector product is a vector, not a number; confusing it with the dot product.
- Writing $\vec{u} \wedge \vec{v} = \vec{v} \wedge \vec{u}$: the sign changes!
- Using $\sin \theta$ with θ outside $[0, \pi]$ or forgetting the absolute value in volumes.
- Forgetting the factor $\frac{1}{2}$ (triangle) or $\frac{1}{6}$ (tetrahedron).

- Computing d in $ax+by+cz+d=0$ with a wrong sign: always substitute the coordinates of A .

5. GCE tips

- Show the right-hand rule or state perpendicularity explicitly: examiners award method marks for it.
- Always verify your cross product with the two dot-product checks — a fast, free mark-saver.
- In plane problems, state clearly the point used and the normal vector before writing the Cartesian equation.
- Present coplanarity and volume answers with exact values first, then a decimal approximation if required.



Signed • OnBuch+

ONBUCH+ TEACHING TEAM
FOLLOWS THE OFFICIAL MINESEC SYLLABUS

OnBuch+