

PURE MATHEMATICS WITH STATISTICS • UPPER SIXTH

UPPER SIXTH — PURE MATHEMATICS WITH STATISTICS

CHAPTER 8

DESCRIPTIVE STATISTICS

SUGGESTED TIME: 23 PERIODS

IN THIS CHAPTER

This chapter equips Upper Sixth students with the complete machinery of descriptive statistics: data collection and presentation, measures of central tendency and dispersion, index numbers and skewness. It prepares learners to handle any GCE Advanced Level statistics question with rigour and confidence.

- By the end of this chapter I can define and correctly use the basic vocabulary of statistics (population, sample, variable, parameter, statistic).
- By the end of this chapter I can classify data and choose an appropriate method of collection.
- By the end of this chapter I can construct frequency tables, including grouped frequency distributions with correct class boundaries.
- By the end of this chapter I can represent data graphically using bar charts, pie charts, line graphs, histograms, frequency polygons and ogives.
- By the end of this chapter I can compute and interpret the mode, median, mean, range, quartiles, deciles and percentiles of grouped and ungrouped data.
- By the end of this chapter I can compute and interpret the variance and standard deviation of grouped and ungrouped data, and combine means of several groups.



DOWNLOAD ON- BUCH

Free • Google Play
AI tutor, past papers,
results



GO ONBUCH+

All signed courses, unlimited AI
tutor, PDF sheets — OnBuch+
tab in the app

THIS SIGNED COURSE CONTAINS

COURSE

complete, illustrated,
step by step

METHODS

and worked examples

EXERCISES

exam style + solutions

SUMMARY SHEET

the essentials to
remember



Signed • OnBuch+

ONBUCH+ TEACHING TEAM
FOLLOWS THE OFFICIAL MINESEC SYLLABUS

OnBuch+

CONTENTS

- | | |
|--|--|
| 1. Basic Terminologies and Types of Data | Standard Deviation |
| 2. Tabular Presentation and Grouped Frequency Distributions | 8. Quartiles, Deciles, Percentiles and Interquartile Range |
| 3. Graphical Representation I: Bar, Pie and Line Charts | 9. Index Numbers |
| 4. Graphical Representation II: Histogram, Frequency Polygon and Ogive | 10. Skewness and Shape of Distributions |
| 5. Measures of Central Tendency I: Mode, Median, Range | 11. Integration activity |
| 6. Measures of Central Tendency II: Mean and Combined Mean | 12. Methods and worked examples |
| 7. Measures of Dispersion: Variance and | 13. Exercises |
| | 14. Solutions to the exercises |
| | 15. Summary sheet |

CHAPTER OBJECTIVES

- By the end of this chapter I can define and correctly use the basic vocabulary of statistics (population, sample, variable, parameter, statistic).
- By the end of this chapter I can classify data and choose an appropriate method of collection.
- By the end of this chapter I can construct frequency tables, including grouped frequency distributions with correct class boundaries.
- By the end of this chapter I can represent data graphically using bar charts, pie charts, line graphs, histograms, frequency polygons and ogives.
- By the end of this chapter I can compute and interpret the mode, median, mean, range, quartiles, deciles and percentiles of grouped and ungrouped data.
- By the end of this chapter I can compute and interpret the variance and standard deviation of grouped and ungrouped data, and combine means of several groups.
- By the end of this chapter I can construct and interpret simple, weighted, Laspeyres and Paasche index numbers.
- By the end of this chapter I can identify the shape of a distribution and measure skewness using Pearson's and the quartile coefficients.

PREREQUISITES

- Arithmetic operations with fractions, decimals and percentages (Lower Sixth).
- Reading and plotting points in the Cartesian plane (Lower Sixth).
- Basic algebraic manipulation, including the sigma (Σ) notation for sums (Lower Sixth, Sequences and Series).
- Simple proportional reasoning and ratios (Form 5 / Lower Sixth).
- Square roots and use of a scientific calculator.

1 BASIC TERMINOLOGIES AND TYPES OF DATA

Every morning, the Douala City Council statistician must decide where to send the waste-collection trucks: which neighbourhood produces the most refuse, and how much does it vary from day to day? At the same time, the Ministry of Public Health compares the average weight of babies born in Yaoundé and Bamenda, while the National Institute of Statistics publishes the consumer price index that determines whether your pocket money still buys the same quantity of puff-puff. All these decisions rest on one toolkit: **descriptive statistics**. In this chapter, you will learn to collect, organise, summarise and interpret data exactly as Cameroonian institutions do — and exactly as the GCE Advanced Level examiners expect.

But before we can summarise data, we must answer more basic questions: What exactly is data? Where does it come from? How do we collect it without being misled? A wrong answer at this first stage poisons everything that follows. This section therefore builds the vocabulary and the critical judgement on which the whole chapter stands.

1.1 Statistics as a Science

Intuition. Suppose you record the heights of all 40 students in your Upper Sixth class and compute the average. You have described your class. Now suppose you measure only 40 students chosen at random from a whole school of 1200 students, and you use their average to draw a conclusion about all the students. You have gone beyond description: you have made an inference. This distinction is the first great divide in statistics.

DEFINITION Statistics, Descriptive and Inferential

Statistics is the science of collecting, organising, presenting, analysing and interpreting numerical data in order to make decisions under uncertainty.

- **Descriptive statistics** deals with the collection, organisation, presentation and summary of data (tables, charts, averages, measures of spread) without drawing conclusions beyond the data at hand.
- **Inferential statistics** uses information from a sample to draw conclusions, estimates or predictions about a larger population, together with a measure of the reliability of those conclusions.

This chapter is devoted to descriptive statistics; inference (estimation, hypothesis testing) comes later in your course.

1.2 Population, Census and Sample

Intuition. A cook tasting a pot of ndolé does not eat the whole pot to know whether there is enough salt: she stirs well and tastes one spoonful. The pot is the population; the spoonful is the sample. The tasting works only if the spoonful fairly represents the pot — that is why she stirs first.


DEFINITION Population, Census, Sample

- A **population** is the complete set of all individuals or items under study. It need not be people: the population may be all bags of cement produced by a factory in Douala in June, or all candidates sitting the GCE Advanced Level in a given year.
- A **census** is a study that collects information from every member of the population.
- A **sample** is a subset of the population, chosen to be studied in place of the whole population.
- The **sampling frame** is the actual list (or device) from which the sample is drawn, e.g. the school register, the electoral roll, a list of households.
- A **sampling unit** is a single member of the sampling frame that can be selected.

EXAMPLE Population or Sample?

A market inspector in Bamenda wants to know the average mass of 50 kg bags of rice sold at the main market. There are 800 bags. He weighs 30 bags chosen at random.

- Population: all 800 bags of rice at the market.
- Sampling frame: the list (or physical arrangement) of the 800 bags.
- Sample: the 30 bags actually weighed.
- Sampling unit: one bag of rice.



The sample is drawn from the sampling frame, which should cover the whole population.

WARNING A frame that does not match the population

If the sampling frame omits part of the population, the sample cannot represent it. Sur-

veying only students who own smartphones about school fees excludes poorer families and biases the results. Always check: does my frame cover the whole population?

Why sample at all? A census gives exact information but is expensive, slow, and sometimes destructive (testing the lifetime of a bulb destroys it; checking every bag of cement for strength breaks every bag). A well-chosen sample gives reliable information quickly and cheaply. The simplest way to choose fairly is **simple random sampling**: every member of the population has the same chance of being selected, for example by numbering the sampling frame and drawing numbers from a hat or from a table of random numbers.

1.3 Statistical Variables and Their Types


When we collect data, we record some characteristic of each individual: height, religion, number of siblings, favourite subject. Such a characteristic is called a statistical variable.

DEFINITION Statistical Variable

A **statistical variable** (or simply variable) is a characteristic or attribute that can take different values from one member of the population to another. The set of all recorded values is the **raw data**.

DEFINITION Qualitative and Quantitative Variables

- A variable is **qualitative** (categorical) if its values are qualities or categories, not numbers on which arithmetic makes sense.
 - **Nominal**: categories with no natural order (e.g. region of origin, blood group, religion).
 - **Ordinal**: categories with a natural order but no measurable spacing (e.g. grade A, B, C; level of satisfaction: poor / fair / good).
- A variable is **quantitative** (numerical) if its values are numbers on which arithmetic is meaningful.
 - **Discrete**: takes only isolated values, usually obtained by counting (e.g. number of children per family: 0, 1, 2, ...).
 - **Continuous**: can take any value in an interval, usually obtained by measuring (e.g. height, mass, time, temperature).



Classification of statistical variables.

WORKED EXAMPLE 1 Classifying Variables

Classify each variable as qualitative (nominal or ordinal) or quantitative (discrete or continuous):

1. the number of goals scored by each team in the MTN Elite One championship;
2. the time taken by students to run 100 m;
3. the marital status of teachers in a school;
4. the rank (1st, 2nd, 3rd, ...) of candidates in an examination.

SOLUTION

1. Goals are counted and take whole-number values: quantitative discrete.
2. Time is measured and can take any value in an interval (e.g. 12.37 s): quantitative continuous.
3. Marital status (single, married, widowed, divorced) is a set of unordered categories: qualitative nominal.
4. Ranks have a natural order but the “distance” between 1st and 2nd is not numerically defined: qualitative ordinal.

WARNING Numbers are not always quantitative

A jersey number, a telephone number or a candidate’s index number is made of digits, yet adding two jersey numbers is meaningless: these are nominal labels, not quantitative

variables. The test is always: does arithmetic on the values make sense?

1.4 Parameter versus Statistic; Raw versus Processed Data

DEFINITION Parameter and Statistic

A **parameter** is a numerical summary (mean, proportion, standard deviation, ...) computed from the whole population. A **statistic** is the corresponding summary computed from a sample. We usually use the sample statistic to estimate the unknown population parameter.

WARNING Common mistake: confusing parameter and statistic

The mean mass of all 800 bags of rice is a parameter; the mean mass of the 30 weighed bags is a statistic. In the GCE exam, read the question twice and ask: was this number computed from the whole population or from a sample? Writing “the population mean is the sample mean” is a classic error: the sample mean only estimates the population mean.

Raw data are the values exactly as collected, before any organisation (e.g. a list of 40 heights in centimetres, in the order the students were measured). **Processed data** are raw data that have been organised or summarised: sorted, tallied, grouped into frequency tables, presented in charts, or reduced to averages. The rest of this chapter is precisely the art of turning raw data into processed data without losing the essential information.

1.5 Methods of Data Collection

DEFINITION Primary and Secondary Data

Primary data are collected first-hand by the investigator for the current study. **Secondary data** already exist, having been collected earlier by someone else for another purpose (published reports, hospital records, NIS publications, past examination results).

The main methods of collecting primary data are:

- **Direct observation:** the investigator watches and records events (e.g. counting vehicles passing the Wouri bridge between 7 a.m. and 8 a.m.). Advantage: accurate, no dependence on what people say. Limitation: slow, costly, and unsuitable for opinions or past events.
- **Interview:** a trained enumerator asks questions face-to-face or by telephone. Advantage: high response rate, questions can be explained, suitable for respondents who cannot read. Limitation: expensive, time-consuming, and the interviewer's tone may influence answers.

- **Questionnaire:** a written list of questions distributed to respondents who fill it in themselves. Advantage: cheap, fast, reaches many people, anonymous (encourages honesty). Limitation: low response rate, no chance to clarify misunderstood questions.
- **Experiment:** the investigator controls conditions and records the outcome (e.g. testing two fertilisers on identical plots of maize in Buea). Advantage: can establish cause and effect. Limitation: may be costly, impractical or unethical.
- **Use of existing records** (secondary data): consulting registers, files, publications. Advantage: immediate and cheap. Limitation: definitions may differ, data may be outdated or collected with a different aim.

METHOD Choosing an Appropriate Data-Collection Method

1. Define the variable precisely and the population concerned.
2. Check whether secondary data already exist that answer the question reliably; if yes, use them.
3. Match the method to the variable: opinions and attitudes → questionnaire or interview; observable behaviour → observation; cause-and-effect questions → experiment.
4. Weigh cost, time and accuracy: a census is ideal but often impossible; choose a sample method you can afford that still represents the population.
5. Pilot the instrument: test your questionnaire on a few respondents first and correct ambiguous or leading questions before the full survey.

1.6 Sources of Error and Bias

Even a carefully planned survey can go wrong. The main dangers are:

- **Non-response bias:** people who refuse to answer (or cannot be reached) often differ systematically from those who answer. If only 40 % of questionnaires are returned, the missing 60 % may hold a different opinion.
- **Leading questions:** wording that pushes the respondent towards an answer, e.g. “Do you agree that the excellent new school fees policy is fair?”
- **Ambiguous or double questions:** “Do you like maths and physics?” — a student may like only one.
- **Sensitive questions:** respondents may lie about income, age or health unless anonymity is guaranteed.
- **Faulty sampling frame:** as seen above, a frame that omits part of the population.
- **Measurement and recording errors:** a badly calibrated scale, a mistyped figure.

EXAMPLE A biased versus a well-designed questionnaire

Biased version (circulated in a school): “Don’t you think the boring canteen food should be improved? Tick: Yes / No.” Problems: the question is leading (“boring” pushes towards “Yes”), and it assumes the respondent eats at the canteen.

Improved version: “How often do you eat at the school canteen? Tick one: Daily / Sometimes / Never. If you eat there, how would you rate the food? Tick one: Very good / Good / Fair / Poor.” Each question is neutral, single, and offers ordered response categories.

KEY POINTS Key Points of this Section

- Descriptive statistics summarises data; inferential statistics generalises from a sample to a population.
- Population = all items under study; census = study of all of them; sample = the subset actually studied, drawn from a sampling frame.
- Variables are qualitative (nominal or ordinal) or quantitative (discrete or continuous); digits used as labels are still qualitative.
- A parameter describes the population; a statistic describes a sample and estimates the parameter.
- Primary data are collected first-hand (observation, interview, questionnaire, experiment); secondary data come from existing records.
- Beware of non-response and leading questions: a biased instrument makes all later calculations worthless.

EXERCISE 1 Terminology Check

A GCE Board official wishes to study the performance of all Advanced Level candidates in the Centre Region. She selects 200 candidates at random from the official results database and computes their average mark in Pure Mathematics, obtaining 52.4.

1. Identify the population, the sampling frame, the sample and the sampling unit.
2. Is 52.4 a parameter or a statistic? Justify.
3. Is the average mark of all candidates in the region known? What is it called?
4. State one advantage and one limitation of using the results database (secondary data) instead of interviewing candidates.

SOLUTION Exercise 1

1. Population: all Advanced Level candidates in the Centre Region. Sampling frame: the official results database. Sample: the 200 selected candidates. Sampling unit: one candidate.
2. 52.4 is a statistic, because it was computed from a sample of 200 candidates, not from the whole population.
3. No; it is unknown. It is the population mean, a parameter, which the statistic 52.4 estimates.
4. Advantage: the data already exist, so the study is fast and cheap. Limitation: the official was not in control of how the data were produced and cannot obtain information not recorded in the database (e.g. candidates' study habits).

EXERCISE 2 Variables and Methods

For each study below, state the variable, its type, and the most appropriate method of data collection:

1. a farmer in Ngaoundéré wants to compare the yields of two varieties of maize;
2. a school principal wants to know how students travel to school (on foot, by bike, by taxi, by bus);
3. a nurse records the birth weights of babies delivered in a district hospital during one month.

SOLUTION Exercise 2

1. Variable: yield of maize (e.g. in kg per plot); quantitative continuous. Method: experiment — grow both varieties on identical plots under the same conditions and measure the yields.
2. Variable: means of transport to school; qualitative nominal. Method: questionnaire (or direct observation at the school gate); the categories have no natural order.
3. Variable: birth weight; quantitative continuous. Method: direct measurement/observation recorded in the hospital register (the register itself is a source of secondary data for later studies).

DID YOU KNOW? Statistics in Cameroon

Cameroon's National Institute of Statistics (NIS) coordinates official statistics, conducts the national population and housing census, and publishes the consumer price index that tracks the cost of living. A general census is one of the largest operations in the country: tens of thousands of enumerators visit households across all ten regions, because a census — unlike a survey — aims to reach every member of the population. The results guide the drawing of electoral districts, the siting of schools and hospitals, and the allocation of state resources to the regions.

2 TABULAR PRESENTATION AND GROUPED FREQUENCY DISTRIBUTIONS

In the previous section we learnt to recognise the different **types of data** (qualitative, quantitative, discrete, continuous) and the **methods of collection**. But raw data, as collected, are usually a long, disordered list of numbers: they tell us almost nothing at first sight. Imagine the marks of 40 Upper Sixth students written on a single line. To answer simple questions such as “How many students scored between 50 and 60?” or “What is the most frequent mark?”, we must first **organise** the data. The first and most fundamental tool of organisation is the **frequency table**. This section teaches you how to tally raw data, how to build simple and grouped frequency tables, and how to prepare the cumulative columns that will be used later for the ogive.

2.1 Tallying and the Simple Frequency Table

DEFINITION Frequency, relative frequency, percentage frequency

Let a statistical variable take the values $x_1, x_2, \dots, x_k$ in a population of size N .

- The **frequency** (or absolute frequency) f_i of the value x_i is the number of times the value x_i occurs in the data.
- The **relative frequency** of x_i is the ratio $r_i = \frac{f_i}{N}$.
- The **percentage frequency** of x_i is $r_i \times 100\% = \frac{f_i}{N} \times 100\%$.

PROPERTY Sum of frequencies

The frequencies always satisfy

$$\sum_{i=1}^k f_i = N \quad \text{and} \quad \sum_{i=1}^k r_i = \sum_{i=1}^k \frac{f_i}{N} = \frac{1}{N} \sum_{i=1}^k f_i = 1.$$

Thus the sum of the relative frequencies equals 1 (and the sum of the percentage frequencies equals 100 %). This is an excellent check of any table you construct in the examination.

The practical technique for counting is **tallying**: for each data value read, we draw one stroke in the row of that value. Strokes are grouped in bundles of five, the fifth stroke crossing the first four (|||| crossed), so that counting at the end is fast and reliable.

EXAMPLE Tallying the marks of 40 students

The marks (out of 20) obtained by 40 Upper Sixth students of GBHS Bamenda in a Mathematics test are:

12, 8, 15, 9, 12, 7, 14, 12, 10, 8, 16, 12, 9, 11, 13, 8, 12, 10, 15, 7,
11, 12, 9, 14, 8, 12, 10, 13, 16, 9, 12, 11, 8, 14, 12, 10, 13, 9, 12, 11

Reading the data one by one and tallying gives:

Mark x_i	Tally	f_i	Relative frequency f_i/N	% freq.
7		2	0.050	5.0
8	crossed,	6	0.150	15.0
9	crossed	5	0.125	12.5
10		4	0.100	10.0
11		4	0.100	10.0
12	crossed, crossed	10	0.250	25.0
13		3	0.075	7.5
14		3	0.075	7.5
15		2	0.050	5.0
16		1	0.025	2.5
Total		40	1.000	100.0

Check: $2 + 6 + 5 + 4 + 4 + 10 + 3 + 3 + 2 + 1 = 40 = N$ and the relative frequencies sum to 1. Immediately we see that the mark 12 is the most frequent (it will later be called the mode).

2.2 Why and When to Group Data

The table above works well because the marks take only 10 distinct values. But suppose the

variable is the height of 200 students, measured in centimetres: almost every value would be different, and a simple frequency table would have nearly 200 rows, each with frequency 1. The table would be useless. The remedy is to **group** the values into intervals called **classes**.

KEY POINTS When to group data

Group data into classes when:

- the variable is continuous (heights, masses, times, temperatures), or
- the variable is discrete but takes many distinct values (marks out of 100, ages of a large population, incomes in FCFA).

Grouping loses some detail (individual values disappear) but reveals the shape of the distribution.

How many classes? Too few classes hide the pattern; too many classes recreate the disorder. A common guide is **Sturges' rule**:

$$k \approx 1 + 3.322 \log_{10} N,$$

where N is the number of observations and k the suggested number of classes. For $N = 40$: $k \approx 1 + 3.322 \log_{10} 40 \approx 1 + 3.322(1.602) \approx 6.3$, so about 6 classes. Sturges' rule is only a guide: in practice we choose a convenient number (often between 5 and 15) giving a **convenient class width** (a whole number, a multiple of 5 or 10, etc.).

DID YOU KNOW? Sturges' rule and Cameroon's census

Sturges' rule was published in 1926 by Herbert Sturges. It works well for moderately sized samples ($N < 200$). For very large data sets (like Cameroon's population census conducted by the National Institute of Statistics), other rules such as the Rice rule ($k = 2\sqrt[3]{N}$) or the Freedman–Diaconis rule are often preferred. In GCE examinations, Sturges' rule is the expected reference unless the question specifies otherwise.

2.3 Class Limits, Boundaries, Width and Class Mark

DEFINITION Vocabulary of a class

A class is an interval of values, written for instance 50–59 or [50; 60[.


- The **class limits** are the stated end values: lower class limit 50, upper class limit 59.
- The **class boundaries** are the true separation points between consecutive classes when the variable is continuous or rounded: lower boundary 49.5, upper boundary 59.5.
- The **class width** is the difference between the upper and lower boundaries: $59.5 -$

$$49.5 = 10.$$

- The **class mark** (or mid-point) is the centre of the class:

$$\text{class mark} = \frac{\text{lower limit} + \text{upper limit}}{2} = \frac{50 + 59}{2} = 54.5.$$

Why boundaries? Marks recorded “to the nearest unit” mean that a recorded 59 stands for any true value in $[58.5; 59.5[$. The boundary 59.5 is therefore the genuine frontier with the next class 60–69, whose lower boundary is also 59.5. Boundaries **remove the gaps** between classes and guarantee that every possible value falls in exactly one class.



No gap: boundary 59.5 belongs to both class edges

Class limits (stated) versus class boundaries (true frontiers)
for a continuous variable, with class marks at the centres.

WARNING Common mistake: overlapping classes or gaps

Two classic errors destroy a grouped table:

- Overlapping classes, e.g. 50–60, 60–70: where does a value of 60 go? Use 50–59, 60–69 for discrete data, or the convention $[50; 60[$, $[60; 70[$ (lower limit included, upper excluded) for continuous data.
- Gaps, e.g. 50–59, 61–70: the value 60 has no class. Consecutive classes must touch at a boundary.

Also, never use classes of different widths in the same table unless the syllabus question explicitly does so: unequal widths distort the histogram.

2.4 Constructing a Grouped Frequency Distribution Step by Step

METHOD Constructing a grouped frequency distribution

1. Find the range: $R = x_{\max} - x_{\min}$.
2. Choose the number of classes k (Sturges' rule as a guide, then round sensibly).

3. Compute the class width: $w \approx \frac{R}{k}$, rounded up to a convenient value.
4. Fix the lower limit of the first class (at or just below $x_{\min}$), then write all classes.
5. Write the class boundaries and the class marks.
6. Tally each observation into its class and record the frequencies f_i .
7. Check that $\sum f_i = N$, then add relative and percentage frequency columns if required.

WORKED EXAMPLE 2 Grouping the marks of 40 students

The marks (out of 100) of 40 candidates in a mock GCE paper range from 23 to 88. Construct a grouped frequency distribution with boundaries, class marks, relative frequencies and a “less than” cumulative column.

SOLUTION

Solution. Step 1: range $R = 88 - 23 = 65$. Step 2: Sturges gives $k \approx 6.3$; we take $k = 7$. Step 3: $w \approx 65/7 \approx 9.3$; we round up to the convenient width $w = 10$. Step 4: first class 20–29 (starts just below 23). Steps 5–7: tallying the 40 marks gives:

Class	Boundaries	Mark m_i	Tally	f_i	Rel. freq. $f_i/40$	Cum. <
20–29	19.5–29.5	24.5		3	0.075	3
30–39	29.5–39.5	34.5		4	0.100	7
40–49	39.5–49.5	44.5	crossed,	6	0.150	13
50–59	49.5–59.5	54.5	crossed,	8	0.200	21
60–69	59.5–69.5	64.5	crossed,	7	0.175	28
70–79	69.5–79.5	74.5	crossed,	6	0.150	34
80–89	79.5–89.5	84.5	crossed,	6	0.150	40
Total				40	1.000	

Checks: $\sum f_i = 3 + 4 + 6 + 8 + 7 + 6 + 6 = 40$; relative frequencies sum to 1. Each class mark is the mean of the limits, e.g. $\frac{50+59}{2} = 54.5$; each width is $59.5 - 49.5 = 10$. The last cumulative entry equals $N = 40$, as it must.

2.5 Discrete versus Continuous Grouping

KEY POINTS Two styles of grouping

- Discrete data (counts: number of children per family, marks): classes are written with whole-number limits, 0–4, 5–9, 10–14, and boundaries fall at half-integers: $-0.5-4.5$, $4.5-9.5$, ...
- Continuous data (measurements: height, mass, time): classes are written with the convention $[a; b[$, e.g. $[160; 170[$ cm, $[170; 180[$ cm, where the lower bound is included and the upper bound excluded; limits and boundaries then coincide.

In both cases the class mark is the centre of the class and the width is the difference of the boundaries.

2.6 Cumulative Frequency Columns: Preparing the Ogive**DEFINITION** Cumulative frequencies

- The **less than cumulative frequency** of a class is the total number of observations falling below the upper boundary of that class. It is obtained by adding frequencies from the top of the table downwards.
- The **more than cumulative frequency** of a class is the total number of observations falling at or above the lower boundary of that class. It is obtained by adding frequencies from the bottom upwards.

For the table of the worked example, the “more than” column reads: 40, 37, 33, 27, 19, 12, 6 (starting from 40 and subtracting f_i successively). Note the duality: for every class, (less than) + (more than) = N . These columns, plotted against the upper (respectively lower) boundaries, give the **ogive** studied in Section 4, from which the median and quartiles will be read graphically.

EXERCISE 3 Practice

The masses (in kg, to the nearest kg) of 30 sacks of cocoa at a buying centre in Kumba range from 41 kg to 80 kg.

1. Using Sturges’ rule, suggest a number of classes and a class width.
2. Construct a grouped frequency table with classes 40–44, 45–49, ..., giving boundaries, class marks and frequencies (invent plausible frequencies summing to 30).
3. Add the “less than” and “more than” cumulative columns and verify that their sum in each row equals 30.

SOLUTION Exercise 1

1. $k \approx 1 + 3.322 \log_{10} 30 \approx 1 + 3.322(1.477) \approx 5.9$, so $k = 6$ classes; range = $80 - 41 = 39$, width $\approx 39/6 = 6.5$; the given width 5 with 9 classes (40–44 to 80–84) is also acceptable and more convenient. 2.–3. Example of a valid table (frequencies may vary, only the checks matter): $f = 2, 3, 5, 6, 5, 4, 3, 1, 1$ for the classes 40–44 to 80–84; $\sum f = 30$. Boundaries: 39.5–44.5, 44.5–49.5, ...; marks: 42, 47, 52, ... Less-than column: 2, 5, 10, 16, 21, 25, 28, 29, 30; more-than column: 30, 28, 25, 20, 14, 9, 5, 2, 1; each pair sums to 30. ✓

KEY POINTS Key points of the section

- Tallying in bundles of five gives reliable frequencies; always check $\sum f_i = N$ and $\sum r_i = 1$.
- Group when the variable is continuous or takes many values; Sturges' rule $k \approx 1 + 3.322 \log_{10} N$ guides the number of classes.
- Limits are stated; boundaries are the true frontiers (half a unit beyond the limits for data rounded to the nearest unit); width = difference of boundaries; class mark = centre.
- Classes must neither overlap nor leave gaps.
- "Less than" and "more than" cumulative columns prepare the ogive and satisfy: their sum in each row equals N .

3 GRAPHICAL REPRESENTATION I: BAR, PIE AND LINE CHARTS

In the previous section we learnt how to organise raw data into frequency tables and grouped frequency distributions. A table is precise, but it is not always speaking: a parent reading a school report, a trader in Mokolo market, or a minister presenting a budget rarely has time to scan columns of figures. A well-chosen **diagram** conveys the same information at a glance. In this section we study the three diagrams used for qualitative or discrete data: the **bar chart**, the **pie chart** and the **line graph**. Diagrams for continuous grouped data (histogram, frequency polygon, ogive) will follow in the next section.

3.1 Bar Charts

Intuition.

Suppose a teacher asks the 40 students of an Upper Sixth class their favourite subject. The natural picture is one vertical bar per subject, the height of the bar showing how many students chose it. Longer bar, more students: the comparison is immediate.


DEFINITION Bar chart

A **bar chart** is a diagram in which each category of a qualitative (or discrete) variable is represented by a rectangular bar of equal width, separated by gaps, whose height (or length) is proportional to the frequency of that category.

The two golden rules are: **equal widths** and **gaps between bars**. The gaps are not decorative: they signal that the categories are separate and that there is no continuous scale between them.

EXAMPLE Simple bar chart: favourite subjects

The favourite subjects of 40 Upper Sixth students are: Mathematics 12, Physics 8, Biology 9, Economics 6, Literature 5. The simple bar chart below displays these frequencies directly.



Simple bar chart: favourite subjects of 40 Upper Sixth students.

Multiple (side-by-side) bar charts.


When we want to compare two or more groups across the same categories, we place the bars side by side, one colour per group, with a legend. For instance, comparing boys and girls in the same survey of favourite subjects: for each subject we draw two adjacent bars.

EXAMPLE Multiple bar chart: favourite subjects by gender

The table below shows the favourite subjects of 20 boys and 20 girls.

Subject	Boys	Girls	Total
Mathematics	8	4	12
Physics	5	3	8
Biology	3	6	9
Economics	2	4	6
Literature	2	3	5

The multiple bar chart allows immediate visual comparison: boys prefer Mathematics and Physics, while girls prefer Biology and Economics.



Multiple bar chart: favourite subjects by gender.

Component (stacked) bar charts.

When we want to show both the total of each category and its composition, we stack the components on top of one another inside a single bar per category. For example, one bar per class (Upper Sixth Arts, Upper Sixth Science), each bar split into “pass” and “fail” segments. The total height gives the class size; the segments give the breakdown.

EXAMPLE Reading a stacked bar chart

A stacked bar for Upper Sixth Science has total height 50, with a “pass” segment of height 38. We read off immediately: 50 candidates, 38 passes, hence $50 - 38 = 12$ failures. Extracting frequencies from a diagram is a routine GCE task: always read heights against the axis, and use totals to deduce missing segments.

WARNING Percentage component bar chart

A variant is the **percentage component bar chart** where each bar has the same height (representing 100%) and the segments show the percentage composition. This is useful when totals differ widely but we want to compare proportions.

3.2 Pie Charts**Intuition.**

A pie chart represents the whole (total frequency) by a full circle and each category by a slice (sector) whose angle is proportional to its frequency. It answers the question: “what share of the whole does each category represent?”

PROPERTY Angle of a sector

If a category has frequency f out of a total frequency N , the angle of its sector is

$$\theta = \frac{f}{N} \times 360^\circ.$$

This follows because the full circle (360°) represents the whole frequency N , and angles are shared proportionally. The sum of all sector angles is exactly 360° — a built-in check.

METHOD Computing pie-chart angles

1. Compute the total frequency N .
2. For each category, compute $\theta = \frac{f}{N} \times 360^\circ$.
3. Round each angle (usually to the nearest degree), then check that the rounded angles sum to 360° ; if not, adjust the largest sector by 1° .
4. Draw a circle, mark a starting radius, and construct the sectors one after another with a protractor.
5. Label each sector (or use a legend) with the category and, if useful, its percentage.

WORKED EXAMPLE 3 Household expenditure of a Cameroonian family

Statement. A family in Yaoundé spends its monthly budget of 200 000 FCFA as follows: Food 80 000, Rent 50 000, Transport 20 000, School fees 30 000, Savings 20 000.


Construct the angles of a pie chart representing this budget.

SOLUTION

Solution. Total $N = 200\,000$ FCFA. Applying $\theta = \frac{f}{N} \times 360^\circ$:

Category	Amount (FCFA)	Angle
Food	80 000	$\frac{80000}{200000} \times 360^\circ = 144^\circ$
Rent	50 000	90°
School fees	30 000	54°
Transport	20 000	36°
Savings	20 000	36°
Total	200 000	360° ✓

The angles sum to 360° , so the chart can be drawn as below.



Pie chart of a monthly family budget of 200 000 FCFA.

WARNING Angles and percentages

Do not confuse the angle with the percentage. The percentage is $\frac{f}{N} \times 100$; the angle is $\frac{f}{N} \times 360^\circ$. Also, a pie chart is meaningless if the categories do not form a complete whole (e.g. favourite subjects where students could pick two): the slices would not add up to one pie.

3.3 Line Graphs and Time Series


DEFINITION Time series and line graph

A **time series** is a set of observations of a variable recorded at successive points in time (days, months, years). Its **line graph** is obtained by plotting the values against time and joining consecutive points by straight line segments. Time is always on the horizontal axis.

The line graph is the natural tool for spotting trends (is the quantity rising or falling overall?), seasonal patterns (regular peaks, e.g. rainfall in the rainy season in Buea) and turning points.

EXAMPLE Cocoa prices over 12 months

The average farm-gate price of cocoa (in FCFA per kg) recorded monthly in the South-West Region is plotted below. The rising trend from January to May, the dip around July, and the recovery towards December are visible at a glance — information that a bare table would hide.



Line graph of average cocoa prices over 12 months.

EXAMPLE Reading a line graph: interpolation

From the cocoa price graph, estimate the price in mid-April. April (month 4) is 920, May (month 5) is 980. Mid-April corresponds to $x = 4.5$. Linear interpolation gives $\frac{920+980}{2} = 950$ FCFA/kg. Always state that this is an estimate assuming linear change between recorded points.

3.4 Choosing the Right Diagram

METHOD Choosing an appropriate diagram

1. Identify the type of data: qualitative categories, discrete counts, or a variable evolving in time.
2. To compare frequencies between categories: use a bar chart (simple, multiple or stacked).
3. To show shares of a total: use a pie chart.
4. To show evolution over time: use a line graph.
5. For continuous grouped data: use a histogram or frequency polygon (next section) — never a gapped bar chart.

WARNING Bar chart or histogram?

A very common GCE mistake is to draw bars touching each other for categorical data, or conversely to leave gaps when the data are continuous. Rule: gaps for separate categories (bar chart); touching bars for continuous data grouped into classes (histogram). If your classes are 10–19, 20–29, ..., the bars must touch (using class boundaries 9.5–19.5, ...), and the diagram is a histogram, not a bar chart.

3.5 Reading, Interpreting and Detecting Misleading Graphs

Reading a chart means extracting exact frequencies: read bar heights against the vertical axis, sector angles with the formula $f = \frac{\theta}{360} \times N$, and points of a line graph against both axes. Interpreting means going further: stating the mode category, comparing shares, describing a trend.

WARNING Misleading graphs — GCE exam awareness

Examiners love to show a “bad” graph and ask why it misleads. Watch for:

- Truncated vertical axis: if the frequency axis starts at, say, 90 instead of 0, a bar of height 100 looks twice as tall as a bar of height 95, although the real difference is only 5%. Always check where the axis starts.
- Unequal bar widths: a bar drawn twice as wide and slightly taller suggests (by its area) a difference far greater than the data justify. All bars must have equal width.
- Unequal time intervals on a line graph, which distort the apparent speed of change.
- Missing labels, scales or titles: a graph without a scale cannot be interpreted at all.

The honest remedy: start frequency axes at 0, keep widths and intervals uniform, and label everything.

KEY POINTS Key points

- Bar chart: equal widths, gaps between bars, height = frequency; simple, multiple or stacked versions.
- Pie chart: sector angle $\theta = \frac{f}{N} \times 360^\circ$; the angles must sum to 360° .
- Line graph: the diagram of a time series; time on the horizontal axis; reveals trends and seasonal patterns.
- Choose the diagram according to the type of data and the message to convey.
- Beware of truncated axes and unequal widths: they make small differences look large.

EXERCISE 4 Pie chart and bar chart

The 60 students of a school canteen were asked their preferred meal: ndolé 18, eru 15, rice and beans 12, achu 9, koki 6.

1. Compute the pie-chart angle for each meal and check your answer.
2. Draw a simple bar chart of the same data.
3. Which diagram better shows each meal's share of all preferences? Justify.

SOLUTION Exercise 1

1. $N = 60$, so $\theta = 6f$ degrees: ndolé 108° , eru 90° , rice and beans 72° , achu 54° , koki 36° .
Sum: $108 + 90 + 72 + 54 + 36 = 360^\circ$. ✓
2. Five bars of equal width, separated by gaps, of heights 18, 15, 12, 9, 6 on a frequency axis starting at 0.
3. The pie chart, because it displays shares of a total: each slice is directly the fraction $\frac{f}{60}$ of the whole, e.g. ndolé represents $\frac{18}{60} = 30\%$ of preferences.

4 GRAPHICAL REPRESENTATION II: HISTOGRAM, FREQUENCY POLYGON AND OGIVE

In the previous section we represented data with bar charts, pie charts and line charts. Those tools are excellent for discrete or categorical data: each bar stands apart, each sector of a pie shows a share of a whole. But what happens when the data are **continuous** — heights, masses, examination marks measured on a continuous scale — and have been organised into a grouped frequency distribution? Separated bars would give the false impression that there are gaps between the classes. In this section we meet the three graphical tools designed for grouped continuous data: the **histogram**, the **frequency polygon** and the **ogive** (cumulative frequency curve). The ogive is especially powerful: it allows us to read off the median and the quartiles graphically, anticipating the calculations of Sections 6 and 8.

4.1 The Histogram for Grouped Continuous Data

From observation to idea.

A Biology teacher in Buea measures the heights of 50 Upper Sixth students and groups them into classes 150–155, 155–160, ..., 175–180 cm. If we draw a bar chart with gaps, we suggest that no student can have a height between 155.0 and 155.1 cm — which is absurd, since height is continuous. The natural fix is to let the rectangles touch each other. The resulting diagram is called a **histogram**.

DEFINITION Histogram

A **histogram** is a graphical representation of a grouped frequency distribution of a continuous variable, consisting of adjacent rectangles (no gaps) whose bases are the class intervals on the horizontal axis and whose areas are proportional to the class frequencies.

WARNING Histogram or bar chart?

Do not confuse the two. In a bar chart the bars are separated, the horizontal axis is categorical or discrete, and it is the height of each bar that shows the frequency. In a histogram the bars touch, the horizontal axis is a continuous numerical scale, and it is the area of each rectangle that represents the frequency. When all classes have the same width, height and area are proportional, so the two ideas coincide — but only then.

Equal class widths.

When every class has the same width, we simply take

$$\text{height of rectangle} = \text{frequency},$$


because $\text{area} = \text{width} \times \text{height}$ and the width is constant. The horizontal axis must be a true continuous scale, marked with the class boundaries (150, 155, 160, ...), not the class mid-points.

EXAMPLE Heights of 50 students

The table below gives the grouped heights (in cm) of 50 students.

Height (cm)	150–155	155–160	160–165	165–170	170–175	175–180
Frequency	4	10	16	12	6	2

All classes have width 5, so we plot rectangles of height equal to the frequency over each interval $[150, 155]$, $[155, 160]$, ...



Histogram of the heights of 50 students (equal class widths of 5 cm).

4.2 Unequal Class Widths: Frequency Density

Why height alone can mislead.

Suppose one class is twice as wide as the others. If we draw its rectangle with height equal to its frequency, its area will look twice as big as it should, and the eye judges areas, not heights. The honest solution is to make the **area** of each rectangle equal (or proportional) to the frequency. Since $\text{area} = \text{class width} \times \text{height}$, the height must be adjusted.

DEFINITION Frequency density

For a class in a grouped distribution, the **frequency density** is

$$\text{frequency density} = \frac{\text{class frequency}}{\text{class width}}.$$

In a histogram, the height of each rectangle is the frequency density, so that

$$\text{area of rectangle} = \text{class width} \times \text{frequency density} = \text{frequency}.$$

METHOD Drawing a histogram with unequal class widths

1. Check that the classes are continuous; if necessary, convert class limits to class boundaries (e.g. 10–19, 20–29 become 9.5–19.5, 19.5–29.5).
2. Compute the width of each class.
3. Compute the frequency density = $\frac{\text{frequency}}{\text{class width}}$ for each class.
4. Draw the horizontal axis as a continuous scale covering all boundaries; label the vertical axis “frequency density”.
5. Over each class interval, draw a rectangle whose height is the frequency density. The rectangles must touch.

WORKED EXAMPLE 4 Histogram with unequal classes

The ages of members of a community club in Bamenda are summarised below. Compute the frequency densities and describe the histogram.

Age (years)	10–19	20–29	30–39	40–59
Frequency	20	30	25	15

SOLUTION

Solution. Using boundaries, the classes are 9.5–19.5, 19.5–29.5, 29.5–39.5 and 39.5–59.5.

Class boundaries	Width	Frequency	Frequency density
9.5–19.5	10	20	$20/10 = 2.0$
19.5–29.5	10	30	$30/10 = 3.0$
29.5–39.5	10	25	$25/10 = 2.5$
39.5–59.5	20	15	$15/20 = 0.75$

The histogram has rectangles of heights 2.0, 3.0, 2.5 and 0.75 over the four intervals. Notice the last rectangle: although 15 members is a substantial frequency, the class is twice as wide, so the rectangle is drawn low and wide; its area, $20 \times 0.75 = 15$, correctly represents the frequency.

4.3 The Frequency Polygon

DEFINITION Frequency polygon

A **frequency polygon** is obtained from a histogram by plotting the points


(class mid-point, frequency)

and joining consecutive points by straight line segments. The polygon is usually closed by joining the end points to the mid-points of the empty classes immediately below the first class and above the last class (where the frequency is zero).

The frequency polygon is ideal for **comparing two distributions on the same axes**: two histograms drawn together would overlap confusingly, but two polygons can be drawn in different colours and compared at a glance.

EXAMPLE Comparing two schools

For the heights of the 50 students above, the mid-points are 152.5, 157.5, 162.5, 167.5, 172.5, 177.5 with frequencies 4, 10, 16, 12, 6, 2. Suppose a second school of 50 students gives frequencies 2, 6, 14, 16, 8, 4 over the same classes. Plotting both polygons on the same axes shows immediately that the second school's heights are shifted slightly to the right: its students are on average taller.



Frequency polygons of two schools superimposed on the histogram of School A.

KEY POINTS Frequency polygon — key points

- Points are plotted at class mid-points, joined by straight lines.

- The polygon is closed at both ends on the horizontal axis.
- The area under the frequency polygon equals the area under the corresponding histogram.
- It is the best tool for comparing two (or more) distributions with the same classes.

4.4 Cumulative Frequency and the Ogive

From observation to idea.

A school principal asks: “How many candidates are shorter than 165 cm? Shorter than 170 cm?” To answer such “less than” questions quickly, we accumulate the frequencies as we move up the classes. This gives the **cumulative frequency distribution**.

DEFINITION Cumulative frequency

The **cumulative frequency** of a class is the total frequency of all values less than the upper class boundary of that class. It is obtained by adding successively the class frequencies.

EXAMPLE Cumulative frequency table for the heights

For the 50 students (boundaries 150, 155, ..., 180):

Height (cm)	Frequency	Cumulative frequency
150–155	4	4
155–160	10	14
160–165	16	30
165–170	12	42
170–175	6	48
175–180	2	50

Each cumulative frequency is plotted against the upper class boundary: (155, 4), (160, 14), (165, 30), (170, 42), (175, 48), (180, 50), together with the starting point (150, 0).

DEFINITION Ogive


The **ogive** (or cumulative frequency curve) is the curve obtained by plotting cumulative frequency against the upper class boundaries and joining the points (by straight segments for a polygonal ogive, or by a smooth S-shaped curve).

METHOD Constructing an ogive from a cumulative frequency table

1. Write down the upper class boundary of every class.
2. Build the cumulative frequency column by successive addition; check that the last value equals N , the total frequency.
3. Plot the points (upper boundary, cumulative frequency), and add the point (lower boundary of first class, 0).
4. Join the points with straight segments (or a smooth curve). The ogive always rises from 0 to N and never decreases.
5. Label both axes clearly: the vertical axis is “cumulative frequency”.

WARNING The most common mistake

Never plot cumulative frequency against the class mid-point. A cumulative frequency answers the question “how many observations are less than this value?”, and that value must be the upper class boundary. Plotting against mid-points shifts the whole ogive half a class to the left and gives wrong medians and quartiles. Examiners penalise this heavily.



Ogive of the heights, with readings at $N/4$, $N/2$ and $3N/4$ giving Q_1 , the median Q_2 and Q_3 .

4.5 Reading the Median and Quartiles from the Ogive

The ogive turns “less than” questions around: instead of asking “how many students are below x ?”, we ask “below which height do we find half the students?”. That height is the **median**.

PROPERTY Graphical determination of median and quartiles

For a distribution of N observations, draw the ogive and then:

- locate $\frac{N}{2}$ on the cumulative frequency axis, draw a horizontal line to the ogive, then a vertical line down: the value read is the median Q_2 ;
- locate $\frac{N}{4}$: the value read is the lower quartile Q_1 ;
- locate $\frac{3N}{4}$: the value read is the upper quartile Q_3 .

The interquartile range is then $IQR = Q_3 - Q_1$. (Proof not examinable; the method follows directly from the definitions of Section 8.)

WORKED EXAMPLE 5 Reading the median and quartiles

From the ogive of the 50 heights, estimate the median, the quartiles and the interquartile range.

SOLUTION

Solution. Here $N = 50$, so

$$\frac{N}{4} = 12.5, \quad \frac{N}{2} = 25, \quad \frac{3N}{4} = 37.5.$$

Drawing horizontal lines at 12.5, 25 and 37.5 up to the curve and projecting down to the height axis (see the figure), we read approximately

$$Q_1 \approx 159.3 \text{ cm}, \quad Q_2 \approx 163.4 \text{ cm}, \quad Q_3 \approx 168.1 \text{ cm}.$$

These readings are consistent with linear interpolation on the table: for instance Q_1 lies in the class 155–160 (cumulative frequencies 4 to 14), giving $155 + \frac{12.5 - 4}{10} \times 5 = 159.25 \approx 159.3 \text{ cm}$, and Q_3 lies in the class 165–170 (cumulative frequencies 30 to 42), giving $165 + \frac{37.5 - 30}{12} \times 5 = 168.125 \approx 168.1 \text{ cm}$. Hence

$$IQR = Q_3 - Q_1 \approx 168.1 - 159.3 = 8.8 \text{ cm}.$$

Interpretation: about half of the students have heights between roughly 159.3 cm and 168.1 cm, and the middle 50 % of heights spread over about 8.8 cm. These graphical estimates will be confirmed by calculation in Sections 6 and 8.

WARNING Reading errors to avoid

- Use $N/2$, not $(N + 1)/2$, when reading from the ogive: the ogive is a continuous-curve estimate.
- Always read the horizontal line from the cumulative axis to the curve, then drop vertically to the variable axis — never the reverse.
- State your answers as estimates (use $\approx$): graphical readings are never exact.

DID YOU KNOW? Why “ogive”?

The word ogive comes from architecture: it names the pointed, S-shaped arch used in Gothic cathedrals. Statisticians borrowed the word because cumulative frequency curves have exactly that graceful S-shape — flat at the extremes where few data accumulate, and steep in the dense middle classes. The steeper the ogive over a class, the more concentrated the data are there.

KEY POINTS Chapter summary — what you must master

- A histogram has touching rectangles whose areas represent frequencies; with unequal class widths, plot frequency density = frequency $\div$ class width.
- A frequency polygon joins the points (mid-point, frequency) and is ideal for comparing distributions.
- An ogive plots cumulative frequency against upper class boundaries; it rises from 0 to N .
- From the ogive, read Q_1 at $N/4$, the median at $N/2$ and Q_3 at $3N/4$; then $IQR = Q_3 - Q_1$.

EXERCISE 5 Practice

The masses (in kg) of 60 bags of beans at a market in Mbouda are grouped as follows:

Mass (kg)	40–45	45–50	50–55	55–60	60–70
Frequency	6	15	20	12	7

1. Compute the frequency density of each class and describe the histogram.
2. Construct the cumulative frequency table.
3. State the coordinates of the points to plot for the ogive.

4. Given that the ogive passes through (50, 21) and (55, 41), estimate the median by linear interpolation between the appropriate points.

SOLUTION Practice

- Widths: 5, 5, 5, 5, 10. Frequency densities: $6/5 = 1.2$; $15/5 = 3.0$; $20/5 = 4.0$; $12/5 = 2.4$; $7/10 = 0.7$. The histogram has touching rectangles of these heights; the last rectangle is twice as wide but low, its area $10 \times 0.7 = 7$ correctly showing the frequency.
- Cumulative frequencies: 6, 21, 41, 53, 60.
- Plot (40, 0), (45, 6), (50, 21), (55, 41), (60, 53), (70, 60) — cumulative frequency against upper class boundaries.
- $N/2 = 30$, which lies between 21 (at 50 kg) and 41 (at 55 kg). By linear interpolation,

$$\text{median} \approx 50 + \frac{30 - 21}{41 - 21} \times 5 = 50 + \frac{9}{20} \times 5 = 52.25 \text{ kg.}$$

So the median mass of a bag of beans is approximately 52.25 kg.

5 MEASURES OF CENTRAL TENDENCY I: MODE, MEDIAN, RANGE

In the previous section we learned how to display a distribution: histograms, frequency polygons and ogives give us a picture of the data. But a picture alone is not enough for the GCE examiner, nor for the economist comparing salaries in Douala and Yaoundé. We need numbers that summarise a whole distribution. The first family of such numbers are the **measures of central tendency** (or averages): they tell us where the centre of the data lies. In this section we study three of them: the **mode**, the **median**, and the simplest measure of spread, the **range**.

5.1 The Mode of Ungrouped Data

Imagine a shoe seller at Mokolo market who records the sizes of the last 20 pairs of shoes sold. To restock wisely, she does not need the “average” size in some complicated sense: she needs the size that sells most often. That value is the mode.

DEFINITION Mode

The **mode** of a set of data is the value that occurs with the greatest frequency.

- A distribution with exactly one mode is called **unimodal**.
- A distribution with two values sharing the greatest frequency is called **bimodal**.

- If every value occurs the same number of times (or no value repeats), the data set has no mode.

EXAMPLE Finding the mode

The marks of 15 candidates in a quiz out of 10 are:

4, 7, 5, 7, 6, 7, 5, 8, 7, 4, 6, 7, 5, 9, 7.

Counting frequencies: 4 appears 2 times, 5 appears 3 times, 6 appears 2 times, 7 appears 6 times, 8 once, 9 once. The greatest frequency is 6, so

Mode = 7.

Had both 5 and 7 appeared 6 times each, the data would be bimodal with modes 5 and 7.

WARNING Mode is a value, not a frequency

A very common error is to give the frequency as the answer. In the example above the mode is 7 (the mark), not 6 (its frequency). The mode is always a data value.

5.2 The Modal Class of Grouped Data

When data are grouped into classes, individual values are lost, so we can no longer read off a mode exactly. We proceed in two stages.

DEFINITION Modal class

For grouped data, the **modal class** is the class with the highest frequency (strictly, the highest frequency density when classes have unequal widths; for equal-width classes, the highest frequency suffices).

If a single estimate of the mode is required from grouped data, we may locate it inside the modal class by linear interpolation. The idea: the mode is pulled towards the “stronger” neighbour. If the class before the modal class has frequency f_{-1} and the class after has frequency f_{+1} , the mode divides the modal class in the ratio $(f_m - f_{-1}) : (f_m - f_{+1})$ measured from the lower boundary.

PROPERTY Mode of grouped data by interpolation — GCE extra

For a modal class with lower class boundary L , class width c , modal frequency f_m , and neighbouring frequencies f_{-1} (class before) and f_{+1} (class after):

$$\text{Mode} \approx L + \frac{f_m - f_{-1}}{(f_m - f_{-1}) + (f_m - f_{+1})} \times c.$$

This formula is an estimate; it is not always required at GCE, but it earns marks when “estimate the mode” is asked for grouped data.

EXAMPLE Estimating the mode from grouped data

The heights of 60 students give: modal class 160–165 cm with $f_m = 20$, previous class frequency $f_{-1} = 14$, next class frequency $f_{+1} = 12$, width $c = 5$, lower boundary $L = 159.5$.

$$\text{Mode} \approx 159.5 + \frac{20 - 14}{(20 - 14) + (20 - 12)} \times 5 = 159.5 + \frac{6}{14} \times 5 \approx 161.6 \text{ cm.}$$

5.3 The Median of Ungrouped Data

Suppose nine workers in a small company in Bamenda earn monthly salaries (in thousands of FCFA):

50, 55, 60, 65, 70, 75, 80, 85, 500.

The last value is the manager's salary. The “typical” salary is badly distorted by that extreme value if we average naively. The median solves this: it is simply the **middle value** once the data are arranged in order.

DEFINITION Median

The **median** of a set of n observations arranged in ascending order is the middle value:

- if n is odd, the median is the value in position $\frac{n+1}{2}$;
- if n is even, the median is the mean of the two values in positions $\frac{n}{2}$ and $\frac{n}{2} + 1$.

EXAMPLE Odd and even n

(a) Odd. For the nine salaries above, already ordered, position $\frac{9+1}{2} = 5$ gives median = 70 thousand FCFA. Notice the manager's 500 has no influence.

(b) Even. For the marks 3, 5, 6, 8, 9, 12 ($n = 6$), the middle positions are 3rd and 4th:

$$\text{Median} = \frac{6 + 8}{2} = 7.$$

Note that 7 is not itself a data value — this is perfectly acceptable for a median.

WARNING Order first!

The single most frequent mistake is to pick the middle value of the data as given, without arranging them in ascending order. Always sort first, then locate the middle.

5.4 The Median of Grouped Data by Interpolation

With grouped data we no longer know individual values, but we know how many observations fall below each class boundary — exactly the information contained in the cumulative frequency table and the ogive of the previous section. The median is the value below which 50% of the observations lie, i.e. the value of the $\frac{N}{2}$ -th observation, where N is the total frequency.

METHOD Locating the median and the median class

1. Compute the total frequency N and form the cumulative frequency column.
2. Compute $\frac{N}{2}$ (do not add 1, even for odd N , when data are grouped).
3. Read down the cumulative frequencies until you find the first cumulative frequency $\geq \frac{N}{2}$: the corresponding class is the **median class**.
4. Apply the interpolation formula below.

PROPERTY Median of grouped data by linear interpolation

Let L be the lower class boundary of the median class, F the cumulative frequency up to the end of the class before the median class, f the frequency of the median class, c its width, and N the total frequency. Then

$$\text{Median} = L + \left(\frac{\frac{N}{2} - F}{f} \right) \times c.$$

Justification (examinable as understanding): inside the median class we assume the f observations are spread uniformly. We need to climb from F observations (at the lower boundary L) up to $\frac{N}{2}$ observations, i.e. a fraction $\frac{N/2 - F}{f}$ of the way through the class. Multiplying this fraction by the class width c gives the distance to add to L .

WORKED EXAMPLE 6 Median of grouped data

The table shows the masses (kg) of 50 bags of cement sampled at a depot.

Mass (kg)	Frequency f	Cumulative frequency
45–47	4	4
48–50	10	14
51–53	17	31
54–56	12	43
57–59	7	50

Estimate the median mass.

SOLUTION

Step 1. $N = 50$, so $\frac{N}{2} = 25$.

Step 2. The first cumulative frequency ≥ 25 is 31, so the median class is 51–53.

Step 3. Class boundaries: $L = 50.5$, width $c = 3$; cumulative frequency before the median class $F = 14$; median class frequency $f = 17$.


Step 4. Apply the formula:

$$\begin{aligned}
 \text{Median} &= 50.5 + \left(\frac{25 - 14}{17} \right) \times 3 \\
 &= 50.5 + \frac{11}{17} \times 3 \\
 &= 50.5 + 1.941 \dots \\
 &\approx 52.44 \text{ kg.}
 \end{aligned}$$

Half of the bags weigh less than about 52.44 kg, and half weigh more.

WARNING Using $N/2$ of the number of classes


A classic trap: some candidates compute $\frac{N}{2}$ using the number of classes (here $\frac{5}{2} = 2.5$) or halve the class frequencies. The $\frac{N}{2}$ in the formula is always half the total frequency (here $\frac{50}{2} = 25$). Also, F is the cumulative frequency before the median class, not that of the median class itself, and L is a class boundary (50.5), not a class limit (51).



Interpolation inside the median class: the median sits a fraction $\frac{N/2 - F}{f}$ of the class width above the lower boundary L .

5.5 Graphical Estimation of the Median from the Ogive

Because the ogive plots cumulative frequency against the **upper class boundaries**, the median is read directly at cumulative frequency $\frac{N}{2}$: draw a horizontal line from $\frac{N}{2}$ on the vertical axis to the curve, then drop vertically to the horizontal axis. This is exactly the graphical counterpart of the interpolation formula, and GCE examiners accept either approach — but the graph gives an estimate whose accuracy depends on your drawing.



Reading the median from the ogive: locate $\frac{N}{2} = 25$ on the cumulative axis, meet the curve, and read down to obtain the median ≈ 52.4 kg. Points are plotted at upper class boundaries.

5.6 The Range

DEFINITION Range

The **range** of a set of data is the difference between the greatest and the least values:

$$\text{Range} = \text{maximum value} - \text{minimum value.}$$

The range is a measure of dispersion (spread), not of centre, but it is traditionally studied alongside the mode and median because of its simplicity. For the nine Bamenda salaries, the range is $500 - 50 = 450$ thousand FCFA.

Uses. The range is quick to compute and easy to understand; it is used in quality control (e.g. checking that the masses of bags of cement stay within acceptable limits) and in meteorology (daily temperature range).

Limitations. The range depends only on the two extreme values and ignores everything in between. Worse, it is extremely sensitive to **outliers**: without the manager, the salary range collapses from 450 to 35 thousand FCFA. Two distributions can have the same range and completely different shapes. This is why more robust measures of spread (interquartile range, variance, standard deviation) are developed later in this chapter.

5.7 Comparing the Mode and the Median: Which Average?

Average	Strengths	Best used when
Mode	Easy; unaffected by extreme values; can be used for non-numerical data (most popular shoe size, most common blood group)	The “most typical” or most frequent category is wanted; data are categorical
Median	Unaffected by outliers; always splits data into two equal halves	Data contain extreme values (salaries, house prices) or are skewed

The mode may not exist, or may be far from the centre (a bimodal distribution has no single “typical” value). The median always exists and resists outliers, but it ignores the actual sizes of the values — only their order matters. When every value should count equally, we need the mean, studied in the next section.

KEY POINTS Key points

- Mode = most frequent value (a data value, not a frequency); data may be unimodal, bimodal, or have no mode.
- Modal class = class of highest frequency; mode of grouped data by interpolation: $L + \frac{f_m - f_{-1}}{(f_m - f_{-1}) + (f_m - f_{+1})} C$.
- Median of ungrouped data: middle value after ordering; position $\frac{n+1}{2}$ (odd n), mean of positions $\frac{n}{2}$ and $\frac{n}{2} + 1$ (even n).

- Median of grouped data: $\text{Median} = L + \left(\frac{N/2 - F}{f} \right) c$, where N is the total frequency.
- From the ogive, read the median at cumulative frequency $\frac{N}{2}$ (plot against upper class boundaries).
- Range = max – min; simple but very sensitive to outliers.

EXERCISE 6 Practice

The table gives the times (minutes) taken by 40 students to solve a problem.

Time (min)	5–9	10–14	15–19	20–24	25–29
Frequency	5	9	14	8	4

1. State the modal class and estimate the mode by interpolation.
2. Construct the cumulative frequency table and estimate the median by interpolation.
3. Estimate the median graphically from an ogive and compare with (b).
4. Explain why the range cannot be computed exactly from this table, and give the largest possible value it could take.

SOLUTION Practice

1. Modal class: 15–19 ($f_m = 14$, $f_{-1} = 9$, $f_{+1} = 8$, $L = 14.5$, $c = 5$).

$$\text{Mode} \approx 14.5 + \frac{14 - 9}{(14 - 9) + (14 - 8)} \times 5 = 14.5 + \frac{5}{11} \times 5 \approx 16.8 \text{ min.}$$

2. Cumulative frequencies: 5, 14, 28, 36, 40. $N = 40$, $\frac{N}{2} = 20$. First cumulative frequency ≥ 20 is 28: median class 15–19, with $L = 14.5$, $F = 14$, $f = 14$, $c = 5$.

$$\text{Median} = 14.5 + \frac{20 - 14}{14} \times 5 = 14.5 + \frac{30}{14} \approx 16.6 \text{ min.}$$

3. Plotting (9.5, 5), (14.5, 14), (19.5, 28), (24.5, 36), (29.5, 40) and reading across from 20 on the cumulative axis gives a median of about 16.6 min, agreeing with (b) up to drawing accuracy. 4. Individual values are unknown after grouping, so the exact maximum and minimum are lost. The largest possible range would be (greatest possible value) – (least possible value) = $29 - 5 = 24$ minutes (using class limits; $29.5 - 4.5 = 25$ using boundaries).

6 MEASURES OF CENTRAL TENDENCY II: MEAN AND COMBINED MEAN

In the previous section we met the **mode**, the **median** and the **range**: the mode tells us the most frequent value, the median tells us the middle value, and the range tells us how spread out the data are. Yet none of these measures uses every observation. If a trader in Bamenda records his daily sales for a month, the mode ignores all days except the most frequent one, and the median ignores the actual sizes of the sales. We now introduce the most famous average of all, the **arithmetic mean**, which takes every single value into account — and we study how to combine means of several groups correctly.

6.1 The Arithmetic Mean of Ungrouped Data

Intuition: sharing equally.

Suppose five students score 8, 10, 12, 14 and 16 marks out of 20 in a class test. If the marks were pooled and shared equally among the five students, each would receive

$$\frac{8 + 10 + 12 + 14 + 16}{5} = \frac{60}{5} = 12.$$

This “equal share” is exactly the arithmetic mean. It is the value each observation would have if the total were redistributed fairly.

DEFINITION Arithmetic mean of ungrouped data

The **arithmetic mean** of n observations $x_1, x_2, \dots, x_n$ is

$$\bar{x} = \frac{x_1 + x_2 + \dots + x_n}{n} = \frac{\sum x}{n}.$$

If the value x_i occurs with frequency f_i (discrete frequency table), then


$$\bar{x} = \frac{\sum f_i x_i}{\sum f_i}.$$

The mean as a balance point.

A beautiful way to picture the mean is to imagine the number line as a light bar with equal unit weights placed at each data point. The bar balances exactly at the mean, because the sum of the signed deviations from the mean is zero:

$$\sum (x_i - \bar{x}) = \sum x_i - n\bar{x} = \sum x_i - \sum x_i = 0.$$

This property is fundamental and is the reason the mean appears everywhere in statistics.



The mean as a balance point: deviations to the left and to the right cancel out, since $\sum(x_i - \bar{x}) = 0$.

EXAMPLE Mean of a discrete frequency table

The number of goals scored by a school team in 10 matches is summarised below. Find the mean number of goals per match.

Goals x	0	1	2	3	4
Frequency f	2	3	3	1	1
fx	0	3	6	3	4

Here $\sum f = 10$ and $\sum fx = 16$, so

$$\bar{x} = \frac{\sum fx}{\sum f} = \frac{16}{10} = 1.6 \text{ goals per match.}$$

Note that the mean need not be one of the observed values, nor even a whole number.

METHOD Finding the mean from raw data or a discrete frequency table

1. List the values x_i and their frequencies f_i (if raw data, each value has frequency 1).
2. Compute the product $f_i x_i$ for each row.
3. Find $\sum f_i$ (total number of observations) and $\sum f_i x_i$ (sum of all values).
4. Apply $\bar{x} = \frac{\sum f_i x_i}{\sum f_i}$.

6.2 Mean of Grouped Data Using Class Mid-Points

When data are grouped into classes, the individual values are lost. We therefore estimate the mean by pretending that every observation in a class sits at the **class mid-point** (also called **class mark**). For a class with lower limit L and upper limit U , the mid-point is $x = \frac{L+U}{2}$.

WARNING Class limits vs. class boundaries

If the classes are given as 45–47, 48–50, etc., there is a gap of 1 unit between them. For continuous data (like mass), the true class boundaries are 44.5–47.5, 47.5–50.5, etc. The mid-point calculated from the stated limits $(45 + 47)/2 = 46$ is exactly the same as from the boundaries $(44.5 + 47.5)/2 = 46$. In GCE examinations, always use the stated limits to compute the mid-point unless class boundaries are explicitly given.

PROPERTY Mean of grouped data

For a grouped frequency distribution with class mid-points x_i and frequencies f_i ,

$$\bar{x} = \frac{\sum f_i x_i}{\sum f_i}.$$

This is an estimate of the true mean; the formula itself must be quoted in the examination.

METHOD Estimating the mean from a grouped frequency table

1. For each class, compute the mid-point $x = \frac{\text{lower limit} + \text{upper limit}}{2}$.
2. Multiply each mid-point by its class frequency to get fx .
3. Sum the frequencies ($\sum f$) and the fx column ($\sum fx$).
4. Calculate $\bar{x} = \frac{\sum fx}{\sum f}$.
5. State the units and remember this is an estimate.

WORKED EXAMPLE 7 Mean from a grouped table

The masses (in kg) of 40 bags of cement checked at a depot are:

Mass (kg)	45–47	48–50	51–53	54–56	57–59
Frequency f	4	10	14	8	4

Estimate the mean mass.

SOLUTION

The class mid-points are 46, 49, 52, 55, 58. We build the working table:

Class	Mid-point x	f	fx
45–47	46	4	184
48–50	49	10	490
51–53	52	14	728
54–56	55	8	440
57–59	58	4	232
Totals		$\Sigma f = 40$	$\Sigma fx = 2074$

$$\bar{x} = \frac{\Sigma fx}{\Sigma f} = \frac{2074}{40} = 51.85 \text{ kg.}$$

6.3 The Assumed-Mean (Coding) Method

When the mid-points are large or awkward, the products fx become heavy. The **coding method** (or **assumed-mean method**) lightens the arithmetic by measuring every mid-point from a chosen **assumed mean** A (usually the mid-point of a central class, often the modal class).

METHOD Assumed-mean (coding) technique

1. Choose an assumed mean A , usually the mid-point of the class with the largest frequency.
2. For each class compute the deviation $d = x - A$ (or the step deviation $d = \frac{x - A}{c}$ when classes have common width c).
3. Compute fd for each class and sum to get Σfd .
4. Apply the formula:

$$\bar{x} = A + \frac{\Sigma fd}{\Sigma f} \quad \text{or, with step } c : \quad \bar{x} = A + c \frac{\Sigma fd}{\Sigma f}.$$

Why it works (examinable derivation).

If $d_i = x_i - A$, then $x_i = A + d_i$, so

$$\bar{x} = \frac{\Sigma f_i x_i}{\Sigma f_i} = \frac{\Sigma f_i (A + d_i)}{\Sigma f_i} = \frac{A \Sigma f_i}{\Sigma f_i} + \frac{\Sigma f_i d_i}{\Sigma f_i} = A + \frac{\Sigma f_i d_i}{\Sigma f_i}.$$

With $d_i = \frac{x_i - A}{c}$ we have $x_i = A + cd_i$ and the same algebra gives $\bar{x} = A + c \frac{\Sigma f_i d_i}{\Sigma f_i}$. This derivation is short and can be asked in the examination.

WORKED EXAMPLE 8 Coding method with step deviation

Using the cement data above with $A = 52$ and $c = 3$, estimate the mean.

SOLUTION

Class	x	f	fx	$d = \frac{x-52}{3}$	fd
45–47	46	4	184	-2	-8
48–50	49	10	490	-1	-10
51–53	52	14	728	0	0
54–56	55	8	440	1	8
57–59	58	4	232	2	8
Totals		40	2074		-2

$$\bar{x} = A + c \frac{\sum fd}{\sum f} = 52 + 3 \times \frac{-2}{40} = 52 - 0.15 = 51.85 \text{ kg,}$$

exactly the same answer as before, with much smaller numbers.

WARNING Sign errors in the coding table

Deviations below A are negative; do not drop the signs when summing $\sum fd$. Also, if you use the step deviation $d = \frac{x-A}{c}$, you must multiply back by c at the end — forgetting the factor c is one of the most common errors at the GCE.

WORKED EXAMPLE 9 Finding an unknown frequency

The mean of the following distribution is 25. Find the missing frequency p .

x	10	20	30	40
f	5	p	10	5

SOLUTION

$$\sum f = 20 + p, \sum fx = 50 + 20p + 300 + 200 = 550 + 20p.$$

$$\bar{x} = \frac{550 + 20p}{20 + p} = 25 \implies 550 + 20p = 500 + 25p \implies 5p = 50 \implies p = 10.$$

6.4 Effect of Adding or Multiplying by a Constant

PROPERTY Linear transformations of the mean

Let $\bar{x}$ be the mean of $x_1, \dots, x_n$, and let a and b be constants.

1. If $y_i = x_i + b$ (a constant added to every value), then $\bar{y} = \bar{x} + b$.
2. If $y_i = ax_i$ (every value multiplied by a constant), then $\bar{y} = a\bar{x}$.
3. In general, if $y_i = ax_i + b$, then $\bar{y} = a\bar{x} + b$.

Proof. For (1): $\bar{y} = \frac{\sum(x_i + b)}{n} = \frac{\sum x_i}{n} + \frac{nb}{n} = \bar{x} + b$. For (2): $\bar{y} = \frac{\sum ax_i}{n} = a \frac{\sum x_i}{n} = a\bar{x}$. Part (3) follows by combining the two. ■

EXAMPLE Using the transformation property

The mean mark of a class in a test marked out of 20 is 11.5. The teacher decides to add 2 bonus marks to every script, then the school converts all marks to a percentage (multiply by 5). The new mean is

$$(11.5 + 2) \times 5 = 67.5 \%$$

WORKED EXAMPLE 10 Correcting a mean after detecting an error

The mean of 20 observations is 35. Later it is found that one value, 50, was wrongly recorded as 5. Find the corrected mean.

SOLUTION

The wrong total is $\sum x_{\text{wrong}} = n\bar{x} = 20 \times 35 = 700$. The correct total is $\sum x_{\text{correct}} = 700 - 5 + 50 = 745$. The corrected mean is $\bar{x}_{\text{correct}} = \frac{745}{20} = 37.25$.

6.5 Combined Mean of Two or More Groups

Suppose Form Six Science A has 30 students with mean mark 62, and Form Six Science B has 20 students with mean mark 55. What is the mean of the two classes together? It is not $\frac{62+55}{2} = 58.5$, because the classes have different sizes. We must recover the total marks of each group: since $\bar{x} = \frac{\sum x}{n}$, the total of a group is $n\bar{x}$.

PROPERTY Combined mean of k groups

If group i has n_i observations and mean $\bar{x}_i$ ($i = 1, \dots, k$), the mean of all the observations combined is

$$\bar{x}_c = \frac{n_1\bar{x}_1 + n_2\bar{x}_2 + \dots + n_k\bar{x}_k}{n_1 + n_2 + \dots + n_k}.$$

For two groups: $\bar{x}_c = \frac{n_1\bar{x}_1 + n_2\bar{x}_2}{n_1 + n_2}$. The formula and its derivation from totals are examinable.

Derivation. The total of group i is $n_i\bar{x}_i$. Hence the grand total is $\sum n_i\bar{x}_i$ and the grand number of observations is $\sum n_i$, so $\bar{x}_c = \frac{\sum n_i\bar{x}_i}{\sum n_i}$. ■

METHOD Finding the combined mean

1. Identify the number of observations n_i and the mean $\bar{x}_i$ for each group.
2. Compute the total for each group: $T_i = n_i\bar{x}_i$.
3. Find the grand total $T = \sum T_i$ and the grand number $N = \sum n_i$.
4. The combined mean is $\bar{x}_c = T/N$.

WORKED EXAMPLE 11 Combined mean of two classes

Science A: $n_1 = 30$, $\bar{x}_1 = 62$. Science B: $n_2 = 20$, $\bar{x}_2 = 55$. Find the combined mean.

SOLUTION

$$\bar{x}_c = \frac{n_1\bar{x}_1 + n_2\bar{x}_2}{n_1 + n_2} = \frac{30(62) + 20(55)}{30 + 20} = \frac{1860 + 1100}{50} = \frac{2960}{50} = 59.2.$$

The combined mean 59.2 lies between 55 and 62, and is closer to 62 because Science A is the larger group.

WORKED EXAMPLE 12 Finding an unknown group mean

The mean height of 25 boys in a class is 168 cm. The mean height of all 40 students (boys and girls) is 165 cm. Find the mean height of the 15 girls.

SOLUTION

Let $\bar{x}_g$ be the girls' mean height. Using the combined-mean formula:

$$165 = \frac{25(168) + 15\bar{x}_g}{40} \Rightarrow 6600 = 4200 + 15\bar{x}_g \Rightarrow \bar{x}_g = \frac{2400}{15} = 160 \text{ cm.}$$

WORKED EXAMPLE 13 Combined mean of three groups

A company has three branches. Branch A: 50 staff, mean salary 300000 FCFA. Branch B: 30 staff, mean salary 250000 FCFA. Branch C: 20 staff, mean salary 400000 FCFA. Find the overall mean salary.

SOLUTION

$$\bar{x}_c = \frac{50(300000) + 30(250000) + 20(400000)}{50 + 30 + 20} = \frac{15000000 + 7500000 + 8000000}{100} = \frac{30500000}{100} = 305000$$

WARNING Averaging the averages

Never compute $\frac{\bar{x}_1 + \bar{x}_2}{2}$ when the group sizes differ. The combined mean is a weighted average: each group mean must be multiplied by its group size. The unweighted average is only correct when $n_1 = n_2$.

6.6 Weighted Mean

The combined mean is a special case of a more general idea: the **weighted mean**, used whenever some values “count more” than others.

DEFINITION Weighted mean

If values $x_1, x_2, \dots, x_k$ carry weights $w_1, w_2, \dots, w_k$, the weighted mean is

$$\bar{x}_w = \frac{\sum w_i x_i}{\sum w_i}.$$

The link with the combined mean is immediate: the combined mean is the weighted mean of the group means, with the group sizes n_i as weights. Similarly, the mean of a frequency table

$\bar{x} = \frac{\sum fx}{\sum f}$ is a weighted mean with the frequencies as weights.

EXAMPLE Weighted mean: a student's average

In a school, the term mark counts for 30% and the examination mark for 70%. A student scores 14/20 in the term work and 11/20 in the examination. Her weighted average is

$$\bar{x}_w = \frac{0.3(14) + 0.7(11)}{0.3 + 0.7} = \frac{4.2 + 7.7}{1} = 11.9/20.$$

A plain average would give 12.5/20, overvaluing the term mark.

EXAMPLE Weighted mean: price index preparation

A household buys three commodities. Their prices (FCFA/kg) and quantities bought (kg) are:

Commodity	Price p	Quantity q	pq
Rice	600	50	30000
Beans	800	20	16000
Maize	400	30	12000
Total		100	58000

The weighted mean price per kg (using quantities as weights) is $\frac{58000}{100} = 580$ FCFA/kg. This is the basis for Laspeyres and Paasche index numbers studied later.

6.7 Mean, Median or Mode? Choosing the Right Average

The mean uses every value, which is its strength — and its weakness. A single extreme value can drag the mean far from the bulk of the data.

EXAMPLE Salaries with an extreme value

A small company in Douala pays its nine workers monthly salaries (in thousands of FCFA):

80, 85, 90, 90, 95, 100, 100, 105, 800

where the last figure is the director's salary. The mean is

$$\bar{x} = \frac{1545}{9} \approx 171.7 \text{ thousand FCFA,}$$

yet eight of the nine workers earn less than 110 000 FCFA. The median, 95 000 FCFA, describes a "typical" worker far better. If the company advertises "average salary 171 700 FCFA", it is technically true but misleading.

KEY POINTS Which average to use?

- **Mean:** use when the data are roughly symmetric with no extreme values, and when further calculations (variance, combined means, regression) are needed. It is the only average that can be treated algebraically.
- **Median:** use when the data contain extreme values or are skewed (salaries, house prices, incomes, reaction times). It is robust and represents the “typical” value.
- **Mode:** use for categorical data (favourite colour, most popular shoe size) or when the most popular value matters (stocking inventory). For grouped data, the modal class is the class with highest frequency.

DID YOU KNOW? The mean in national debates

When newspapers report the “average income” of a country, they usually quote the mean, which is pulled upwards by a small number of very high earners. This is why the median income is often much lower, and why economists prefer it when describing the living standard of a typical household. As a statistician, always ask which average is being quoted!

EXERCISE 7 Practice

1. The mean of the numbers 4, 7, 9, x , 15 is 9. Find x .
2. A grouped distribution has $\sum f = 50$ and, with assumed mean $A = 27$ and class width $c = 5$, $\sum fd = 12$. Find the mean.
3. Group X of 40 candidates has mean mark 58; group Y of 60 candidates has mean mark 64. Find the combined mean.
4. The mean of 20 observations is 35. Later it is found that one value, 50, was wrongly recorded as 5. Find the corrected mean.
5. The mean of 10 numbers is 12. If one number is removed, the mean of the remaining 9 numbers is 11. Find the removed number.
6. A student's marks in five subjects are 65, 72, 58, 80, 60. The weights (coefficients) for these subjects are 3, 4, 2, 5, 1 respectively. Calculate the weighted mean.
7. The mean of a grouped frequency distribution with 5 classes is 41. The class mid-points are 30, 35, 40, 45, 50 and the corresponding frequencies are 4, 8, p , 10, 6. Find p .

SOLUTION Exercise 1

1. $\frac{4+7+9+x+15}{5} = 9 \Rightarrow 35+x=45 \Rightarrow x=10.$
2. $\bar{x} = A + c \frac{\sum fd}{\sum f} = 27 + 5 \times \frac{12}{50} = 27 + 1.2 = 28.2.$
3. $\bar{x}_c = \frac{40(58) + 60(64)}{100} = \frac{2320 + 3840}{100} = 61.6.$
4. The wrong total is $20 \times 35 = 700$. Corrected total: $700 - 5 + 50 = 745$. Corrected mean: $\frac{745}{20} = 37.25.$
5. Total of 10 numbers = $10 \times 12 = 120$. Total of remaining 9 numbers = $9 \times 11 = 99$. Removed number = $120 - 99 = 21.$
6. $\bar{x}_w = \frac{3(65)+4(72)+2(58)+5(80)+1(60)}{3+4+2+5+1} = \frac{195+288+116+400+60}{15} = \frac{1059}{15} = 70.6.$
7. $\sum f = 28+p, \sum fx = 120+280+40p+450+300 = 1150+40p. \frac{1150+40p}{28+p} = 41 \Rightarrow 1150+40p = 1148+41p \Rightarrow p=2.$

KEY POINTS Key points of this section

- Ungrouped data: $\bar{x} = \frac{\sum x}{n}$; frequency table: $\bar{x} = \frac{\sum fx}{\sum f}.$
- Grouped data: use class mid-points $x = \frac{L+U}{2}$; the mean is an estimate.
- Coding method: $\bar{x} = A + \frac{\sum fd}{\sum f}$ or $\bar{x} = A + c \frac{\sum fd}{\sum f}$ with $d = \frac{x-A}{c}$. Derivation is examinable.
- If $y = ax + b$, then $\bar{y} = a\bar{x} + b.$
- Combined mean: $\bar{x}_c = \frac{n_1\bar{x}_1 + n_2\bar{x}_2}{n_1 + n_2}$ — always weight by group sizes.
- Weighted mean: $\bar{x}_w = \frac{\sum w_i x_i}{\sum w_i}.$
- The mean is sensitive to extreme values; prefer the median for skewed data such as salaries.

7 MEASURES OF DISPERSION: VARIANCE AND STANDARD DEVIATION

In the previous section we studied the **mean**, and we even saw how to combine the means of several groups into a single combined mean. The mean tells us where the centre of a data set lies. But two data sets can share exactly the same mean and yet behave very differently. Imagine two bakeries in Douala, each producing an average of 500 loaves per day. Bakery A produces 498, 500, 502, 499, 501 over five days; Bakery B produces 200, 750, 480, 620, 450. Both have mean 500, but Bakery A is far more consistent. To distinguish such situations we need **measures of dispersion** — numbers that describe how spread out the data are around the centre.

7.1 Why the Range Is Not Enough

The simplest measure of spread is the **range**: $\text{range} = x_{\max} - x_{\min}$. It is quick, but it uses only two values out of the whole data set and ignores everything in between.

EXAMPLE The weakness of the range

Consider the two sets

$$A : 2, 50, 50, 50, 98 \quad B : 2, 10, 40, 70, 98.$$

Both have range $98 - 2 = 96$ and mean 50. Yet in A most values sit right at 50, while in B the values are scattered across the whole interval. The range cannot see this difference.

WARNING The range is fragile

Because the range depends only on the two extreme values, a single outlier (one unusually large or small value) can inflate it enormously, even when all the other data are tightly clustered. Never use the range alone to compare the spread of two data sets.

A natural first improvement is the **mean deviation**: take the distance of each value from the mean, $|x - \bar{x}|$, and average these distances:

$$\text{Mean deviation} = \frac{1}{n} \sum |x - \bar{x}|.$$

This uses every data value. However, the absolute value is awkward to handle algebraically (it is not differentiable at 0, and it does not combine nicely with other formulas). Statisticians therefore prefer to square the deviations instead of taking absolute values. Squaring also removes the sign, and it penalises large deviations more heavily — which is usually desirable. This leads us to the central notions of this section.

7.2 Variance and Standard Deviation of Ungrouped Data

DEFINITION Variance and standard deviation

Let $x_1, x_2, \dots, x_n$ be n observations with mean $\bar{x}$. The **variance**, denoted σ^2 (or $\text{Var}(x)$), is the mean of the squared deviations from the mean:

$$\sigma^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2.$$

The **standard deviation** σ is the positive square root of the variance:

$$\sigma = \sqrt{\frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2}.$$

The standard deviation is expressed in the same units as the data (the variance is in squared units), which makes σ the preferred measure for interpretation.

WARNING Why divide by n and not $n - 1$?

At GCE Advanced Level we always treat the data as the whole population under study, so we divide by n . (In more advanced work, when the data are only a sample used to estimate a population variance, one divides by $n - 1$ instead; this is not required here.)

PROPERTY Computational form of the variance

The variance can be computed without subtracting the mean from each value:

$$\sigma^2 = \frac{\sum x^2}{n} - \bar{x}^2 \quad (\text{"mean of squares minus square of the mean"}).$$

Proof (examinable). Expanding the square and using $\sum x = n\bar{x}$:

$$\begin{aligned} \sum (x - \bar{x})^2 &= \sum x^2 - 2\bar{x} \sum x + \sum \bar{x}^2 \\ &= \sum x^2 - 2\bar{x}(n\bar{x}) + n\bar{x}^2 \\ &= \sum x^2 - n\bar{x}^2. \end{aligned}$$

Dividing by n gives $\sigma^2 = \frac{\sum x^2}{n} - \bar{x}^2$. ■

WORKED EXAMPLE 14 Variance of ungrouped data

The marks of 6 candidates in a quiz are 4, 6, 7, 8, 9, 8. Find the variance and the standard deviation.

SOLUTION

Solution. First the mean:

$$\bar{x} = \frac{4 + 6 + 7 + 8 + 9 + 8}{6} = \frac{42}{6} = 7.$$

Using the computational form, $\sum x^2 = 16 + 36 + 49 + 64 + 81 + 64 = 310$, so

$$\sigma^2 = \frac{310}{6} - 7^2 = 51.666 \dots - 49 = 2.666 \dots \approx 2.67,$$

$$\sigma = \sqrt{2.666 \dots} \approx 1.63 \text{ marks.}$$

Check by the definition: the deviations are -3, -1, 0, 1, 2, 1; their squares are 9, 1, 0, 1, 4, 1, summing to 16; and $\frac{16}{6} = 2.666 \dots$. Same answer. ■

7.3 Variance and Standard Deviation of Grouped Data

When data are presented in a frequency table, each value x (or class mid-point) occurs f times, so each squared deviation must be weighted by its frequency.

PROPERTY Variance of grouped data

For a frequency distribution with values (or mid-points) x_i and frequencies f_i ,

$$\sigma^2 = \frac{\sum f(x - \bar{x})^2}{\sum f} = \frac{\sum fx^2}{\sum f} - \bar{x}^2, \quad \bar{x} = \frac{\sum fx}{\sum f}.$$

The standard deviation is $\sigma = \sqrt{\sigma^2}$.

METHOD Computing σ^2 from a grouped table

1. Find the mid-point x of each class.
2. Compute the mean $\bar{x} = \frac{\sum fx}{\sum f}$ (add a column fx).
3. Add a column fx^2 (compute x^2 then multiply by f , or compute $(fx) \times x$).
4. Apply $\sigma^2 = \frac{\sum fx^2}{\sum f} - \bar{x}^2$, then take the square root.

WORKED EXAMPLE 15 Variance of grouped data

The table shows the masses (kg) of 50 bags of cement sampled at a depot in Yaoundé. Find the variance and standard deviation.

Class	Mid-point x	f	fx	fx^2
45–47	46	6	276	12 696
48–50	49	18	882	43 218
51–53	52	16	832	43 264
54–56	55	8	440	24 200
57–59	58	2	116	6 728
Totals		$\Sigma f = 50$	$\Sigma fx = 2546$	$\Sigma fx^2 = 130\,106$


SOLUTION

Solution.

$$\bar{x} = \frac{2546}{50} = 50.92 \text{ kg}, \quad \sigma^2 = \frac{130\,106}{50} - (50.92)^2 = 2602.12 - 2592.8464 = 9.2736.$$

$$\sigma = \sqrt{9.2736} \approx 3.05 \text{ kg}.$$

On average, a bag's mass deviates from the mean by roughly 3 kg.



Two factories with the same mean output (50 units) but very different spreads: Factory A has a small standard deviation (consistent), Factory B a large one (erratic).

7.4 The Coding Method for Variance

When the values x are large or the mean is not a whole number, direct computation is heavy. We simplify the arithmetic by a change of variable (**coding**):

$$d = \frac{x - A}{c} \iff x = A + cd,$$

where A is a chosen assumed mean (usually a central mid-point) and c is the class width (or any convenient common divisor).

PROPERTY Coding formula for the variance

With $d = \frac{x - A}{c}$,

$$\sigma^2 = c^2 \left[\frac{\sum fd^2}{\sum f} - \left(\frac{\sum fd}{\sum f} \right)^2 \right].$$

Justification. Since $x = A + cd$, we have $\bar{x} = A + c\bar{d}$ and $x - \bar{x} = c(d - \bar{d})$. Hence

$$\sum f(x - \bar{x})^2 = c^2 \sum f(d - \bar{d})^2,$$

and applying the computational form to the d 's gives the result. ■

METHOD Coding technique for variance

1. Choose an assumed mean A near the centre and the step c (class width).
2. Form the column $d = (x - A)/c$; the d 's should be small integers.
3. Compute columns fd and fd^2 and their totals.
4. Compute $\sigma^2 = c^2 \left[\frac{\sum fd^2}{\sum f} - \left(\frac{\sum fd}{\sum f} \right)^2 \right]$.
5. Take the square root for σ .

WORKED EXAMPLE 16 Variance by coding

Using the cement-bag data above with $A = 52$ and $c = 3$:

x	f	$d = \frac{x-52}{3}$	fd	fd^2
46	6	-2	-12	24
49	18	-1	-18	18
52	16	0	0	0
55	8	1	8	8
58	2	2	4	8
Totals	50		-18	58

Verify the value of σ^2 found earlier.

SOLUTION

Solution.

$$\sigma^2 = 3^2 \left[\frac{58}{50} - \left(\frac{-18}{50} \right)^2 \right] = 9 [1.16 - 0.1296] = 9(1.0304) = 9.2736.$$

Hence $\sigma \approx 3.05$ kg, exactly as before — but with far simpler arithmetic. Note also that $\bar{x} = A + c\bar{d} = 52 + 3 \times \frac{-18}{50} = 52 - 1.08 = 50.92$ kg, confirming the mean.

WARNING Do not forget the factor c^2 !

The most common examination error with the coding method is to compute the variance of the coded values d and to forget to multiply by c^2 at the end. Remember: the d 's are in "step units"; each step is worth c real units, and since variance involves squares, the correction factor is c^2 (not c). For the standard deviation, multiply by c only: $\sigma_x = c \sigma_d$.

7.5 Effect of Linear Transformations

The coding formula is a consequence of a general and very useful result.

PROPERTY Variance under a linear transformation

Let $y = ax + b$ where a and b are constants. Then

$$\text{Var}(y) = a^2 \text{Var}(x), \quad \sigma_y = |a| \sigma_x.$$

In particular:

- Adding a constant does not change the variance: $\text{Var}(x + b) = \text{Var}(x)$. Shifting all the data moves the mean by the same amount, so the deviations $x - \bar{x}$ are unchanged.
- Multiplying by a multiplies the variance by a^2 : $\text{Var}(ax) = a^2 \text{Var}(x)$.

Proof. If $y = ax + b$ then $\bar{y} = a\bar{x} + b$, so $y - \bar{y} = a(x - \bar{x})$ and

$$\text{Var}(y) = \frac{1}{n} \sum a^2(x - \bar{x})^2 = a^2 \text{Var}(x). \quad \blacksquare$$

EXAMPLE Using the transformation property

The marks of a class have mean 52 and variance 16. The teacher decides to scale every mark by $y = 1.5x - 10$. Then

$$\bar{y} = 1.5(52) - 10 = 68, \quad \text{Var}(y) = (1.5)^2(16) = 36, \quad \sigma_y = 6.$$

Note that the shift of -10 has no effect on the variance.

7.6 Interpretation: Comparing Consistency

The standard deviation measures the typical distance of the data from the mean. A small σ means the values cluster tightly around $\bar{x}$ (consistent, reliable, homogeneous); a large σ means wide scatter (erratic, heterogeneous).

EXAMPLE Two factories

Two factories each produce an average of 500 bottles per hour. Factory A has $\sigma = 5$; Factory B has $\sigma = 40$. Factory A's hourly output almost always lies within a few bottles of 500, whereas Factory B's output swings wildly. A manager would judge Factory A far more consistent, even though both have the same mean. The mean alone would have hidden this completely.

DID YOU KNOW? Why statisticians love the standard deviation

For many roughly bell-shaped distributions, about 68% of the data lie within one standard deviation of the mean and about 95% within two. This "empirical rule" makes σ a powerful yardstick: a value more than 2σ from the mean is already unusual.

7.7 Coefficient of Variation

Suppose we want to compare the dispersion of the heights of students (measured in cm, mean around 165) with the dispersion of their masses (in kg, mean around 60). The standard deviations are in different units and refer to different-sized means, so comparing them directly is meaningless. We therefore use a relative, unit-free measure.

DEFINITION Coefficient of variation

The **coefficient of variation** is the standard deviation expressed as a percentage of the mean:

$$CV = \frac{\sigma}{\bar{x}} \times 100\%.$$

The smaller the CV, the more consistent the data set relative to its mean.

WORKED EXAMPLE 17 Comparing data with different units

The heights of a group of students have $\bar{x} = 165$ cm, $\sigma = 8.25$ cm; their masses have $\bar{x} = 60$ kg, $\sigma = 7.2$ kg. Which characteristic is more variable?

SOLUTION

Solution.

$$CV_{\text{height}} = \frac{8.25}{165} \times 100\% = 5\%, \quad CV_{\text{mass}} = \frac{7.2}{60} \times 100\% = 12\%.$$

Although the masses have a smaller standard deviation in absolute terms, they are relatively much more variable (12% against 5%). The CV, being unit-free, allows this fair comparison.

WARNING Trap: CV requires a meaningful zero

The coefficient of variation is only sensible for quantities measured on a scale with a genuine zero (masses, prices, counts). It is meaningless for temperatures in °C, since $\bar{x}$ could be near 0 or even negative, making the CV explode or change sign.

KEY POINTS Key points of this section

- The range uses only two values and is distorted by outliers; dispersion must involve all the data.
- Variance: $\sigma^2 = \frac{\sum(x - \bar{x})^2}{n} = \frac{\sum x^2}{n} - \bar{x}^2$ (ungrouped); $\sigma^2 = \frac{\sum fx^2}{\sum f} - \bar{x}^2$ (grouped).
- Standard deviation $\sigma = \sqrt{\sigma^2}$ is in the same units as the data.
- Coding with $d = \frac{x - A}{c}$: $\sigma^2 = c^2 \left[\frac{\sum fd^2}{\sum f} - \left(\frac{\sum fd}{\sum f} \right)^2 \right]$ — never forget the c^2 .
- $\text{Var}(ax + b) = a^2 \text{Var}(x)$: adding a constant changes nothing; multiplying by a multiplies

the variance by a^2 .

- Small $\sigma \Rightarrow$ consistent data; large $\sigma \Rightarrow$ scattered data.
- Coefficient of variation $CV = \frac{\sigma}{\bar{x}} \times 100\%$ compares dispersion across different units or different-sized means.

EXERCISE 8 Practice

1. Find the variance and standard deviation of 3, 5, 7, 9, 11.
2. A frequency distribution has $\sum f = 40$, $\sum fx = 200$, $\sum fx^2 = 1240$. Find the mean, variance and standard deviation.
3. Using $A = 25$ and $c = 5$, the coded values of a distribution give $\sum f = 60$, $\sum fd = 12$, $\sum fd^2 = 96$. Find $\bar{x}$ and σ .
4. The marks in Mathematics have mean 55 and standard deviation 10; the marks in Physics have mean 40 and standard deviation 8. Using the coefficient of variation, determine in which subject the class is more consistent.

SOLUTION Exercise 1

1. $\bar{x} = 7$; $\sum x^2 = 9 + 25 + 49 + 81 + 121 = 285$; $\sigma^2 = \frac{285}{5} - 49 = 57 - 49 = 8$; $\sigma = \sqrt{8} \approx 2.83$.
2. $\bar{x} = \frac{200}{40} = 5$; $\sigma^2 = \frac{1240}{40} - 25 = 31 - 25 = 6$; $\sigma = \sqrt{6} \approx 2.45$.
3. $\bar{d} = \frac{12}{60} = 0.2$, so $\bar{x} = A + c\bar{d} = 25 + 5(0.2) = 26$. $\sigma^2 = 25 \left[\frac{96}{60} - 0.04 \right] = 25(1.56) = 39$, hence $\sigma = \sqrt{39} \approx 6.24$.
4. $CV_{\text{Maths}} = \frac{10}{55} \times 100\% \approx 18.2\%$; $CV_{\text{Physics}} = \frac{8}{40} \times 100\% = 20\%$. The class is slightly more consistent in Mathematics.

8 QUARTILES, DECILES, PERCENTILES AND INTERQUARTILE RANGE

In the previous section we measured dispersion with the **variance** and the **standard deviation**. Those measures are powerful, but they have two weaknesses: they depend on every value in the data set, so a single extreme value can inflate them enormously, and they require heavy computation. There is another family of tools, based purely on position in ordered data, which

is simpler, more robust, and extremely popular at the GCE: the **quartiles**, **deciles** and **percentiles**. Just as the median splits ordered data into two equal halves, quartiles split it into four equal parts, deciles into ten, and percentiles into a hundred. From these positions we build new measures of spread — the **interquartile range** and the **percentile range** — and a beautiful summary picture, the **box-and-whisker plot**.

8.1 Quartiles of Ungrouped Data

Intuition.

Imagine the 11 best students of an Upper Sixth class in Bamenda ranked by their mock examination mark. The median is the mark of the 6th student: five students are below, five above. Now split each half again: the mark of the 3rd student leaves a quarter of the group below it, and the mark of the 9th student leaves three quarters below it. These two marks are the lower and upper quartiles.

DEFINITION Quartiles, deciles and percentiles

Let a data set of N values be arranged in ascending order.

- The **quartiles** Q_1, Q_2, Q_3 are the values that divide the data into four equal parts. Q_1 (lower quartile) has about 25% of the data below it, Q_2 is the median (50%), and Q_3 (upper quartile) has about 75% below it.
- The **deciles** $D_1, D_2, \dots, D_9$ divide the data into ten equal parts: D_k has about $\frac{k}{10}$ of the data below it.
- The **percentiles** $P_1, P_2, \dots, P_{99}$ divide the data into one hundred equal parts: P_k has about $k\%$ of the data below it.

Note that $Q_2 = D_5 = P_{50} = \text{median}$, and $Q_1 = P_{25}, Q_3 = P_{75}$.

METHOD Quartiles of ungrouped data (Tukey's hinges method)

1. Arrange the N values in ascending order.
2. Locate the median Q_2 :
 - If N is odd, Q_2 is the middle value (position $\frac{N+1}{2}$).
 - If N is even, Q_2 is the mean of the two middle values (positions $\frac{N}{2}$ and $\frac{N}{2} + 1$).
3. Split the ordered data into a lower half and an upper half:
 - If N is odd, exclude the median itself; the lower half consists of the values strictly below Q_2 , the upper half of those strictly above.
 - If N is even, the lower half is the first $N/2$ values, the upper half the last $N/2$ values.
4. Q_1 is the median of the lower half; Q_3 is the median of the upper half (apply the same

rule as in step 2 to each half).

EXAMPLE Quartiles of 11 marks (odd N)

The marks of 11 students are, in order:

4, 7, 8, 10, 12, 15, 18, 20, 22, 25, 30.

$N = 11$ (odd). Median position = $\frac{11+1}{2} = 6$; $Q_2 = 15$ (the 6th value). Lower half (5 values): 4, 7, 8, 10, 12; its median is the 3rd value $\Rightarrow Q_1 = 8$. Upper half (5 values): 18, 20, 22, 25, 30; its median is the 3rd value $\Rightarrow Q_3 = 22$.

EXAMPLE Quartiles of 10 marks (even N)

Marks: 5, 8, 12, 14, 16, 18, 20, 22, 25, 30. $N = 10$ (even). Median $Q_2 = \frac{16+18}{2} = 17$ (average of 5th and 6th). Lower half (first 5): 5, 8, 12, 14, 16; median $Q_1 = 12$ (3rd value). Upper half (last 5): 18, 20, 22, 25, 30; median $Q_3 = 22$ (3rd value).

8.2 Quartiles of Grouped Data by Interpolation

For grouped data we cannot read individual values, so we proceed exactly as for the median: locate the position of the quartile, find the class containing it using cumulative frequencies, then estimate the value by **linear interpolation**.

PROPERTY Interpolation formula for quartiles of grouped data

For grouped data with total frequency N :

$$Q_i = L + \left(\frac{\frac{iN}{4} - F}{f} \right) \times c, \quad i = 1, 2, 3,$$

where

- L = lower class boundary of the quartile class (the first class whose cumulative frequency reaches or exceeds $\frac{iN}{4}$),
- F = cumulative frequency before that class,
- f = frequency of the quartile class,
- c = class width.

The same pattern gives deciles and percentiles:

$$D_k = L + \left(\frac{\frac{kN}{10} - F}{f} \right) c, \quad P_k = L + \left(\frac{\frac{kN}{100} - F}{f} \right) c.$$

WARNING Position is not value!

The most frequent error at the GCE is to confuse the position $\frac{N}{4}$ with the value of Q_1 . If $N = 80$, then $\frac{N}{4} = 20$ is not the lower quartile: it only tells you that Q_1 is the value of the “20th observation”, which you must then find inside its class by interpolation. Always finish the interpolation before writing your answer.

WORKED EXAMPLE 18 Quartiles, a decile and a percentile of grouped data

The marks of 80 candidates in a statistics test are summarised below.

Marks	Frequency f	Cumulative frequency
0–10	8	8
10–20	16	24
20–30	30	54
30–40	18	72
40–50	8	80

Find Q_1 , Q_2 , Q_3 , D_7 and P_{85} .

SOLUTION

Here $N = 80$ and every class has width $c = 10$. Class boundaries are 0, 10, 20, 30, 40, 50. Lower quartile Q_1 . Position $\frac{N}{4} = 20$. The first cumulative frequency reaching 20 is 24, so the quartile class is 10–20, with $L = 10$, $F = 8$, $f = 16$:

$$Q_1 = 10 + \frac{20 - 8}{16} \times 10 = 10 + 7.5 = 17.5.$$

Median Q_2 . Position $\frac{N}{2} = 40$; class 20–30, $L = 20$, $F = 24$, $f = 30$:

$$Q_2 = 20 + \frac{40 - 24}{30} \times 10 = 20 + 5.333 \dots \approx 25.3.$$

Upper quartile Q_3 . Position $\frac{3N}{4} = 60$; class 30–40, $L = 30$, $F = 54$, $f = 18$:

$$Q_3 = 30 + \frac{60 - 54}{18} \times 10 = 30 + 3.333 \dots \approx 33.3.$$

Seventh decile D_7 . Position $\frac{7N}{10} = 56$; class 30–40:


$$D_7 = 30 + \frac{56 - 54}{18} \times 10 = 30 + 1.111 \dots \approx 31.1.$$

85th percentile P_{85} . Position $\frac{85N}{100} = 68$; class 30–40:

$$P_{85} = 30 + \frac{68 - 54}{18} \times 10 = 30 + 7.777 \dots \approx 37.8.$$

8.3 Graphical Determination from the Ogive

Because the ogive (cumulative frequency curve) is precisely a graph of “position against value”, quartiles and percentiles can be read directly from it: locate $\frac{N}{4}$, $\frac{N}{2}$, $\frac{3N}{4}$ on the cumulative frequency axis, project horizontally to the curve, then vertically down to the value axis.



Reading Q_1 , Q_2 , Q_3 and the interquartile range from the ogive ($N = 80$).

8.4 Interquartile Range and Semi-Interquartile Range

The quartiles immediately give a measure of dispersion focused on the middle half of the data, completely unaffected by extreme values.

DEFINITION Interquartile and semi-interquartile range

The **interquartile range** (IQR) is the difference between the upper and lower quartiles:

$$\text{IQR} = Q_3 - Q_1.$$

The **semi-interquartile range** (also called the quartile deviation) is half of it:

$$\text{SIQR} = \frac{Q_3 - Q_1}{2}.$$

The IQR contains the middle 50% of the observations. A small IQR means the central half of the data is tightly packed.


EXAMPLE Computing the IQR

For our 80 candidates: $\text{IQR} = 33.3 - 17.5 = 15.8$ marks, and $\text{SIQR} = \frac{15.8}{2} = 7.9$ marks. For the 11 ungrouped marks: $\text{IQR} = 22 - 8 = 14$.

In the same spirit, the **percentile range** $P_{90} - P_{10}$ measures the spread of the middle 80% of the data, discarding the top and bottom tenths. In our example, P_{10} has position 8, which falls exactly on the boundary 10 (interpolation gives 10), and P_{90} has position 72, falling on the boundary 40; hence $P_{90} - P_{10} = 40 - 10 = 30$ marks. Examiners like it because it resists the influence of one or two abnormal scores even better than the IQR.

8.5 Box-and-Whisker Plots (beyond the strict syllabus)

A **box-and-whisker plot** summarises a distribution with five numbers: minimum, Q_1 , median Q_2 , Q_3 , maximum. The box spans the IQR; the whiskers extend to the extremes. It is not explicitly required by the Cameroon GCE syllabus, but it is an excellent revision tool and frequently appears in international examinations.



Box-and-whisker plot of the 11 marks, aligned with a number line. The box length is the IQR.

8.6 Interpretation: Position within a Group

Percentiles are the natural language of ranking. If a student's mark equals P_{90} of the class, she is in the **top decile**: at least 90% of the candidates scored at or below her mark, and only about 10% scored above. Similarly, scoring below Q_1 places a candidate in the bottom quarter — a signal that remediation is needed — while scoring above Q_3 places him in the top quarter. No-

tice the subtlety: a percentile describes a position relative to the group, not an absolute level of mastery. Being at P_{95} in a weak school may correspond to a lower actual mark than being at P_{60} in a strong one.

KEY POINTS Key points

- Q_1, Q_2, Q_3 split ordered data into quarters; Q_2 is the median; $Q_1 = P_{25}, Q_3 = P_{75}, D_5 = P_{50}$.
- For grouped data: position first $(\frac{iN}{4}, \frac{kN}{10}, \frac{kN}{100})$, then the class, then interpolation $Q_i = L + \frac{iN/4 - F}{f} \times c$.
- $IQR = Q_3 - Q_1$; $SIQR = \frac{Q_3 - Q_1}{2}$; percentile range = $P_{90} - P_{10}$.
- All these measures are robust: extreme values do not affect them.
- Quartiles and percentiles can be read graphically from the ogive.

EXERCISE 9 Practice

Using the table of the 80 candidates above, find: (a) D_3 ; (b) P_{40} ; (c) the interquartile range if the class widths were doubled to 20 with frequencies 24, 48, 8 in classes 0–20, 20–40, 40–50. Comment on the effect of grouping.

SOLUTION Exercise 1

(a) Position $\frac{3N}{10} = 24$: this falls exactly on the boundary of the second class (cumulative frequency 24), so $D_3 = 20$. (Interpolation: class 10–20, $L = 10, F = 8, f = 16, c = 10$ gives $10 + \frac{24-8}{16} \times 10 = 20$.) (b) Position $\frac{40N}{100} = 32$; class 20–30: $P_{40} = 20 + \frac{32-24}{30} \times 10 = 20 + 2.667 \approx 22.7$. (c) New grouping: $N = 80$, classes 0–20 ($f = 24, c = 20$), 20–40 ($f = 48, c = 20$), 40–50 ($f = 8, c = 10$). Q_1 : position 20, class 0–20, $L = 0, F = 0, f = 24$: $Q_1 = 0 + \frac{20}{24} \times 20 \approx 16.7$. Q_3 : position 60, class 20–40, $L = 20, F = 24, f = 48$: $Q_3 = 20 + \frac{60-24}{48} \times 20 = 35$. Hence $IQR \approx 35 - 16.7 = 18.3$ instead of 15.8: wider classes give a coarser (and here larger) estimate of the spread.

9 INDEX NUMBERS

In the previous sections we learnt how to summarise a single set of data at a single moment: quartiles, deciles and percentiles told us where values stand within one distribution. But in real life, the most urgent questions are questions of change. Is garri more expensive in Douala this year than last year? By how much has the cost of living risen in Yaoundé since 2020? Has co-

coconut production in the Centre Region increased or decreased? A single statistic cannot answer such questions, because prices and quantities of many different goods change at the same time and in different proportions. We therefore need a new tool: the **index number**.

9.1 The Idea of an Index Number

EXPERIMENT A market observation

A student notes the price of a basin of garri at Mokolo market: FCFA 8 000 in 2022 and FCFA 10 000 in 2024. The price has risen by FCFA 2 000, that is by $\frac{2000}{8000} \times 100 = 25\%$. We say the price relative is 1.25, or, multiplied by 100, the price index is 125. The number 125 tells us at a glance: “the price is 25% higher than in the reference year”. Index numbers simply generalise this idea to several goods at once.

DEFINITION Index number and base period

An **index number** is a statistical measure that expresses the value of a variable (price, quantity, production, value) at a given period as a percentage of its value at a reference period called the **base period** (or base year). The index at the base period is conventionally 100.

- An index above 100 indicates an increase relative to the base period.
- An index below 100 indicates a decrease relative to the base period.
- An index of 100 indicates no change.

KEY POINTS Reading an index

An index of 118 means an increase of 18% since the base period; an index of 92 means a decrease of 8%. Never say “the price is 118” — the index has no unit; it is a pure percentage relative to the base.

9.2 Simple Index Numbers

DEFINITION Simple price, quantity and value indices

Let p_0, q_0 be the price and quantity of an item in the base year, and p_n, q_n the corresponding values in the current year.

- **Simple price index** (price relative): $I_p = \frac{p_n}{p_0} \times 100$.

- **Simple quantity index:** $I_q = \frac{q_n}{q_0} \times 100$.
- **Simple value index:** $I_v = \frac{p_n q_n}{p_0 q_0} \times 100 = I_p \times I_q / 100$.

EXAMPLE Simple price index

A litre of palm oil cost FCFA 1200 in 2021 (base year) and FCFA 1500 in 2024. Then

$$I_p = \frac{1500}{1200} \times 100 = 125.$$

The price of palm oil has risen by 25% since 2021.

When several commodities are involved, two elementary constructions exist.

DEFINITION Simple aggregate index and average of relatives

For k commodities:

- **Simple aggregate price index:** $I = \frac{\sum_{i=1}^k p_{n,i}}{\sum_{i=1}^k p_{0,i}} \times 100$ (ratio of total prices).
- **Simple average of price relatives:** $I = \frac{1}{k} \sum_{i=1}^k \frac{p_{n,i}}{p_{0,i}} \times 100$ (arithmetic mean of the individual indices).

WARNING Limitations of simple indices

Simple indices treat every item as equally important, which is unrealistic: a household spends far more on garri than on salt. They are also sensitive to the units of measurement (e.g., price per kg vs price per tonne). This motivates weighted indices.

9.3 Weighted Index Numbers

DEFINITION Weighted index number

A **weighted index number** assigns to each commodity a **weight** reflecting its relative importance (usually the quantity consumed, the value spent, or the expenditure share). A

weighted price index has the general form

$$I = \frac{\sum p_n w}{\sum p_0 w} \times 100,$$

where the w are the weights. The whole question is: which quantities should serve as weights — those of the base year or those of the current year?

METHOD Choosing weights in a weighted index

1. Identify the purpose of the index (cost of living, production, sales value).
2. Choose weights that reflect the importance of each item: quantities consumed (q), expenditure shares ($p_0 q_0$), or output volumes.
3. Decide the period of the weights: base-period quantities q_0 give the Laspeyres index; current-period quantities q_n give the Paasche index; geometric mean of both gives Fisher's ideal index.
4. Keep the same weights in the numerator and in the denominator: only the prices must change between top and bottom.
5. State clearly the base year and the weights used, so the index can be interpreted and reproduced.

9.4 Laspeyres and Paasche Price Indices

PROPERTY Laspeyres price index

The **Laspeyres price index** uses base-period quantities q_0 as weights:

$$I_L = \frac{\sum p_n q_0}{\sum p_0 q_0} \times 100.$$

It answers the question: "How much would the old basket of goods cost at new prices, compared with its cost at old prices?" It measures the change in cost of purchasing a fixed base-period basket.

PROPERTY Paasche price index

The **Paasche price index** uses current-period quantities q_n as weights:

$$I_P = \frac{\sum p_n q_n}{\sum p_0 q_n} \times 100.$$

It answers the question: “How much does the present basket cost at new prices, compared with what the same present basket would have cost at old prices?” It measures the change in cost of purchasing the current basket.

PROPERTY Fisher’s ideal price index

The **Fisher’s ideal price index** is the geometric mean of the Laspeyres and Paasche indices:

$$I_F = \sqrt{I_L \times I_P}.$$

It satisfies the time reversal and factor reversal tests, making it theoretically superior, though less used in practice due to computational complexity.

WARNING Do not mix the two periods!

A very common examination error is to write something like $\frac{\sum p_n q_0}{\sum p_0 q_n} \times 100$, mixing base-year quantities in the numerator with current-year quantities in the denominator. This is meaningless: within one formula, the quantities must all come from the same period. Check: Laspeyres $\rightarrow q_0$ everywhere; Paasche $\rightarrow q_n$ everywhere.

WORKED EXAMPLE 19 Laspeyres, Paasche and Fisher indices for a market basket

The table below gives the prices (FCFA per unit) and quantities consumed of four goods in 2022 (base year, index 0) and 2024 (current year, index n).

Good	p_0	q_0	p_n	q_n
Garri (basin)	8 000	12	10 000	10
Plantain (bunch)	2 000	20	2 500	24
Palm oil (litre)	1 200	30	1 500	28
Fish (kg)	2 500	8	3 500	6

Compute the Laspeyres, Paasche and Fisher price indices for 2024 (base 2022 = 100) and interpret the results.

SOLUTION

Solution. We build the four products needed.

$$\begin{aligned}\sum p_0 q_0 &= 8000(12) + 2000(20) + 1200(30) + 2500(8) \\ &= 96\,000 + 40\,000 + 36\,000 + 20\,000 = 192\,000, \\ \sum p_n q_0 &= 10\,000(12) + 2500(20) + 1500(30) + 3500(8) \\ &= 120\,000 + 50\,000 + 45\,000 + 28\,000 = 243\,000, \\ \sum p_0 q_n &= 8000(10) + 2000(24) + 1200(28) + 2500(6) \\ &= 80\,000 + 48\,000 + 33\,600 + 15\,000 = 176\,600, \\ \sum p_n q_n &= 10\,000(10) + 2500(24) + 1500(28) + 3500(6) \\ &= 100\,000 + 60\,000 + 42\,000 + 21\,000 = 223\,000.\end{aligned}$$

Laspeyres index:

$$I_L = \frac{\sum p_n q_0}{\sum p_0 q_0} \times 100 = \frac{243\,000}{192\,000} \times 100 = 126.5625 \approx 126.6.$$


Paasche index:

$$I_P = \frac{\sum p_n q_n}{\sum p_0 q_n} \times 100 = \frac{223\,000}{176\,600} \times 100 = 126.2741 \approx 126.3.$$

Fisher's ideal index:

$$I_F = \sqrt{I_L \times I_P} = \sqrt{126.5625 \times 126.2741} = \sqrt{15980.5} \approx 126.4.$$

Interpretation. All three indices exceed 100: prices have risen by about 26% between 2022 and 2024. Laspeyres says the old basket (q_0) would cost 26.6% more today; Paasche says the current basket (q_n) costs 26.3% more than it would have at 2022 prices; Fisher gives a compromise at 26.4%. The two answers are close here, but they need not be in general.



A consumer price index rising from 100 (base year 2019) to 131 in 2024: an overall increase of 31% in five years.

9.5 Quantity and Value Indices

The same weighting principles apply to quantities and values.

PROPERTY Laspeyres and Paasche quantity indices

- **Laspeyres quantity index:** $I_{L,q} = \frac{\sum q_n p_0}{\sum q_0 p_0} \times 100$ (base-period prices as weights).
- **Paasche quantity index:** $I_{P,q} = \frac{\sum q_n p_n}{\sum q_0 p_n} \times 100$ (current-period prices as weights).

PROPERTY Value index

The **value index** compares total expenditure (or revenue) between two periods:

$$I_v = \frac{\sum p_n q_n}{\sum p_0 q_0} \times 100.$$

Note that $I_v = \frac{I_p \times I_q}{100}$ only when the same weighting system is used for both price and quantity indices (e.g., both Laspeyres or both Paasche). In general, $I_v \neq I_L \times I_{L,q} / 100$.

9.6 Chain-Base vs Fixed-Base Indices

DEFINITION Fixed-base and chain-base indices

- A **fixed-base index** compares every period to the same base period (e.g., 2019 = 100 for all years).
- A **chain-base index** compares each period to the immediately preceding period (link relatives), then chains them together: $I_{0 \rightarrow n} = \frac{I_{0 \rightarrow 1}}{100} \times \frac{I_{1 \rightarrow 2}}{100} \times \dots \times \frac{I_{n-1 \rightarrow n}}{100} \times 100$.

KEY POINTS When to use which

Fixed-base indices are standard for long-term comparisons and official publications (CPI). Chain-base indices are useful when the basket of goods changes frequently, as they allow updating weights each period without breaking the series.

9.7 Comparing Laspeyres and Paasche

KEY POINTS Laspeyres versus Paasche

- Laspeyres (q_0 weights): easy to compute repeatedly, since the basket is fixed once and for all; but it tends to overstate inflation because it ignores the fact that consumers substitute away from goods whose prices rise fastest (substitution bias).
- Paasche (q_n weights): reflects current consumption habits, but requires new quantity data every period and tends to understate inflation because it uses the basket that has already adjusted to price changes.
- Fisher (geometric mean): theoretically ideal, satisfies time and factor reversal tests, but requires both sets of quantities.
- Both Laspeyres and Paasche are valid; they answer different questions. In practice, most national consumer price indices are of Laspeyres type, with the basket updated periodically (e.g., every 5 years).

DID YOU KNOW? Index numbers in Cameroon

In Cameroon, the National Institute of Statistics (INS) publishes a **consumer price index** (CPI) that tracks the cost of a fixed basket of goods and services — food, housing, transport, clothing — in cities such as Yaoundé and Douala. Government and employers use it for **cost-of-living adjustments**: when the CPI rises sharply, salaries and pensions may be revised so that purchasing power is preserved. The CPI is essentially a Laspeyres-type in-

dex: a fixed basket, repriced every month. The INS also produces **producer price indices** (PPI) for agriculture and industry, and **quantity indices** for major export crops like cocoa, coffee, and cotton.

EXERCISE 10 Computing and interpreting indices

The table below shows the prices (FCFA) and quantities of three items bought by a family in 2021 (base) and 2023.

Item	p_0	q_0	p_n	q_n
Rice (kg)	500	40	650	36
Beans (kg)	800	15	1000	18
Oil (litre)	1100	10	1400	12

1. Compute the simple price index of each item for 2023 (base 2021 = 100).
2. Compute the Laspeyres price index.
3. Compute the Paasche price index.
4. Compute Fisher's ideal price index.
5. Interpret the two indices in one sentence each.

SOLUTION Exercise 1

1. Simple price indices:

- Rice: $\frac{650}{500} \times 100 = 130.0$
- Beans: $\frac{1000}{800} \times 100 = 125.0$
- Oil: $\frac{1400}{1100} \times 100 = 127.3$

2. Laspeyres:

$$\begin{aligned}\sum p_0 q_0 &= 500(40) + 800(15) + 1100(10) = 20\,000 + 12\,000 + 11\,000 = 43\,000, \\ \sum p_n q_0 &= 650(40) + 1000(15) + 1400(10) = 26\,000 + 15\,000 + 14\,000 = 55\,000, \\ I_L &= \frac{55\,000}{43\,000} \times 100 = 127.9069 \approx 127.9.\end{aligned}$$

3. Paasche:

$$\begin{aligned}\sum p_0 q_n &= 500(36) + 800(18) + 1100(12) = 18\,000 + 14\,400 + 13\,200 = 45\,600, \\ \sum p_n q_n &= 650(36) + 1000(18) + 1400(12) = 23\,400 + 18\,000 + 16\,800 = 58\,200, \\ I_P &= \frac{58\,200}{45\,600} \times 100 = 127.6316 \approx 127.6.\end{aligned}$$

4. Fisher's ideal index:

$$I_F = \sqrt{127.9069 \times 127.6316} = \sqrt{16325.5} \approx 127.8.$$

5. Laspeyres: the 2021 basket would cost about 27.9% more in 2023. Paasche: the 2023 basket costs about 27.6% more than it would have cost in 2021. Fisher: a compromise estimate of 27.8% increase. All three confirm a strong rise in the cost of living, close to 28%.

WARNING Final warnings for the examination

- Always state the base year and give the index to at least one decimal place.
- An index of 126.6 means a 26.6% increase, not a 126.6% increase.
- Do not add units (FCFA) to an index number.
- Keep quantities consistent within one formula: q_0 for Laspeyres, q_n for Paasche, in both numerator and denominator.
- For Fisher's index, use the exact values of I_L and I_P (not rounded) before taking the square root.
- Distinguish clearly between price, quantity and value indices in your notation and interpretation.

Index numbers thus condense the evolution of a whole economy into a single, readable figure. In the next section, we return to the shape of distributions and learn how to measure their asymmetry through the notion of skewness.

10 SKEWNESS AND SHAPE OF DISTRIBUTIONS

In the previous section we used index numbers to summarise how prices and quantities evolve over time. We now return to the data themselves and ask a final, very visual question: what does the shape of a distribution tell us? Two classes may have the same mean mark of 12/20, yet in one class the marks are balanced around 12 while in the other most students scored below 12 with a few brilliant candidates pulling the average up. The mean alone cannot distinguish these

situations. The missing ingredient is **skewness** — a measure of the asymmetry of a distribution.

10.1 Symmetric and Skewed Distributions

EXPERIMENT Looking at shapes

Consider the marks of two groups of Upper Sixth students in a mock examination.

Group A: 8, 9, 10, 10, 11, 11, 11, 12, 12, 12, 13, 13, 14, 14, 15.

Group B: 4, 5, 6, 7, 8, 8, 9, 9, 10, 10, 11, 12, 14, 16, 19.

For Group A the frequencies rise gently to a peak in the middle and fall off in the same way on both sides: the histogram is a mirror image about its centre. For Group B the bulk of the marks sits at the lower end, but a long tail stretches towards the high marks (14, 16, 19). The two distributions have completely different shapes, and this difference has practical consequences: in Group B, the "typical" student is weaker than the mean suggests. Indeed, both groups have mean 11.5, but the median of Group A is 12 while that of Group B is only 9: the few high marks in Group B drag the mean upwards without representing most students.

DEFINITION Symmetric distribution

A distribution is **symmetric** if its graph (histogram, frequency polygon or frequency curve) can be divided into two halves that are mirror images of each other about a vertical line through the centre. The two tails are of equal length and shape.

DEFINITION Skewness

Skewness is the degree of asymmetry of a distribution. A distribution that is not symmetric is said to be **skewed**.

- A distribution is **positively skewed** (or skewed to the right) if it has a long tail stretching towards the higher values, with most observations concentrated at the lower end.
- A distribution is **negatively skewed** (or skewed to the left) if it has a long tail stretching towards the lower values, with most observations concentrated at the upper end.


WARNING The tail decides — not the peak!

The most common mistake at the GCE is to name the skewness after the side where the bulk of the data lies. This is wrong. The direction of skewness is the direction of the long tail:

- tail to the right (high values) $\Rightarrow$ positive skewness;

- tail to the left (low values) $\Rightarrow$ negative skewness.

A useful memory aid: the skewness "pulls" in the direction of the tail, exactly as the few extreme values pull the mean towards them.



Three typical shapes: a symmetric bell-shaped curve, a positively skewed curve (long right tail) and a negatively skewed curve (long left tail).

10.2 Relative Positions of Mean, Median and Mode

The three measures of central tendency react differently to extreme values. The **mode** sits at the peak of the distribution and ignores the tails completely. The **median** depends only on the order of the data, so it is only slightly affected by a few extreme values. The **mean**, however, uses every value, so it is dragged towards the long tail. This simple observation explains the following fundamental property.

PROPERTY Mean, median and mode according to shape

- For a symmetric (unimodal) distribution:

$$\text{mean} = \text{median} = \text{mode}.$$

- For a positively skewed distribution (tail to the right):

$$\text{mode} < \text{median} < \text{mean}.$$




- For a negatively skewed distribution (tail to the left):

$$\text{mean} < \text{median} < \text{mode}.$$

Moreover, for moderately skewed distributions, the median usually lies about one third of the way from the mode to the mean, which gives the empirical relation

$$\bar{x} - M_o \approx 3(\bar{x} - M_e),$$

and motivates the second Pearson coefficient below. (This property is quoted, not proved, at the GCE.)



Ordering of mean, median and mode for each shape.
The mean is always pulled towards the long tail.

KEY POINTS Why the mean moves most

The mean is the centre of gravity of the distribution: a single very large value (a millionaire in a village, a mark of 19 in a weak class) pulls it strongly. The median only counts positions, and the mode only looks at the peak. Hence: mean is most affected, mode least affected, median in between.

10.3 Pearson's Coefficients of Skewness

Visual inspection is useful, but the examiner expects a numerical measure. Karl Pearson proposed coefficients based on the distance between the mean and the mode (or median), standardised by the standard deviation σ so that the result is a pure number with no units. Dividing by σ is essential: a gap of 2 marks between mean and mode is large when $\sigma = 1$ but negligible when $\sigma = 10$.

PROPERTY Pearson's coefficients of skewness

For a distribution with mean $\bar{x}$, mode M_o , median M_e and standard deviation $\sigma \neq 0$:

$$S_k = \frac{\bar{x} - M_o}{\sigma} \quad (\text{first coefficient})$$

$$S_k = \frac{3(\bar{x} - M_e)}{\sigma} \quad (\text{second coefficient})$$

The second form is preferred when the mode is ill-defined or difficult to estimate (common for grouped data); the factor 3 comes from the empirical relation $\bar{x} - M_o \approx 3(\bar{x} - M_e)$ for moderately skewed distributions. (Formulae quoted; not examinable as a proof.)

Interpretation.

- $S_k > 0$: positively skewed (mean above mode/median, right tail).
- $S_k = 0$: symmetric.
- $S_k < 0$: negatively skewed (left tail).

As a rule of thumb, $|S_k| < 0.5$ indicates a fairly symmetric distribution, $0.5 \leq |S_k| \leq 1$ a moderately skewed one, and $|S_k| > 1$ a highly skewed one.

WORKED EXAMPLE 20 Pearson's coefficient for examination marks

In a mock GCE examination, the marks of a class give: mean $\bar{x} = 11.4$, median $M_e = 12.0$, standard deviation $\sigma = 3.0$. Compute Pearson's second coefficient of skewness and comment.

SOLUTION

Solution.

$$S_k = \frac{3(\bar{x} - M_e)}{\sigma} = \frac{3(11.4 - 12.0)}{3.0} = \frac{3 \times (-0.6)}{3.0} = -0.6.$$

Since $S_k < 0$, the distribution is negatively skewed: most students obtained high marks, with a tail of weaker marks stretching to the left. Since $|S_k| = 0.6$ lies between 0.5 and 1, the skewness is moderate. Note the consistency check: mean < median, exactly as expected for a left-skewed distribution.

WORKED EXAMPLE 21 Pearson's first coefficient from raw data

The daily sales (in thousands of FCFA) of a small shop in Bamenda over one week were:

20, 22, 22, 23, 24, 25, 40.

Compute Pearson's first coefficient of skewness and interpret it.

SOLUTION

Solution. The mode is clearly $M_o = 22$. The mean is

$$\bar{x} = \frac{20 + 22 + 22 + 23 + 24 + 25 + 40}{7} = \frac{176}{7} \approx 25.14.$$

For the standard deviation we compute the squared deviations from $\bar{x} \approx 25.14$:

$$\begin{aligned} \sum (x_i - \bar{x})^2 &\approx (20 - 25.14)^2 + 2(22 - 25.14)^2 + (23 - 25.14)^2 \\ &\quad + (24 - 25.14)^2 + (25 - 25.14)^2 + (40 - 25.14)^2 \\ &\approx 26.42 + 19.72 + 4.58 + 1.30 + 0.02 + 220.90 = 272.94, \end{aligned}$$

so $\sigma = \sqrt{272.94/7} \approx \sqrt{38.99} \approx 6.24$. Hence

$$S_k = \frac{\bar{x} - M_o}{\sigma} \approx \frac{25.14 - 22}{6.24} \approx 0.50.$$

Since $S_k > 0$, the sales are positively skewed: the single exceptional day of 40 000 FCFA pulls the mean above the mode. The shopkeeper should quote the mode or median, not the mean, to describe a "typical" day.

10.4 Quartile Coefficient of Skewness

Pearson's coefficients depend on the mean and standard deviation, which are themselves sensitive to extreme values. A measure based only on quartiles is more robust. Recall that for a symmetric distribution the median Q_2 lies exactly halfway between Q_1 and Q_3 , i.e. $Q_3 - Q_2 = Q_2 - Q_1$. Any imbalance between these two gaps signals skewness: if the upper gap is wider, the right tail is longer and the skew is positive.

PROPERTY Quartile coefficient of skewness (Bowley's coefficient)

For a distribution with quartiles Q_1 , Q_2 (median) and Q_3 :

$$S_{kQ} = \frac{(Q_3 - Q_2) - (Q_2 - Q_1)}{Q_3 - Q_1} = \frac{Q_3 + Q_1 - 2Q_2}{Q_3 - Q_1}.$$

It always satisfies $-1 \leq S_{kQ} \leq 1$. The sign is interpreted exactly as for Pearson's coefficient: positive for right skew, negative for left skew, zero for symmetry. (Formula quoted; not examinable as a proof.)

WORKED EXAMPLE 22 Quartile coefficient for waiting times

The waiting times (in minutes) of patients at a district hospital give $Q_1 = 8$, $Q_2 = 15$, $Q_3 = 34$. Compute the quartile coefficient of skewness and interpret it.

SOLUTION

Solution.

$$S_{kQ} = \frac{Q_3 + Q_1 - 2Q_2}{Q_3 - Q_1} = \frac{34 + 8 - 2(15)}{34 - 8} = \frac{42 - 30}{26} = \frac{12}{26} \approx 0.46.$$

Since $S_{kQ} > 0$, the distribution of waiting times is positively skewed: half the patients wait at most 15 minutes, but a minority waits much longer (the gap $Q_3 - Q_2 = 19$ exceeds $Q_2 - Q_1 = 7$). The skewness is moderate since $|S_{kQ}| < 0.5$.

WARNING**Common mistakes with skewness coefficients**

- Wrong sign convention. $S_k > 0$ means the tail is on the right (positive skew). Do not describe such a distribution as "skewed to the left".
- Forgetting the factor 3 in Pearson's second coefficient: it is $3(\bar{x} - M_e)/\sigma$, not $(\bar{x} - M_e)/\sigma$.
- Mixing the two quartile gaps. In Bowley's formula the numerator is $(Q_3 - Q_2) - (Q_2 - Q_1)$; reversing the order changes the sign and reverses the conclusion.
- Comparing coefficients from different formulae. Pearson's and Bowley's coefficients measure skewness on different scales; do not expect them to be equal for the same data.
- Claiming skewness from the mean alone. You must compare at least two measures (mean vs median, or the two quartile gaps) before concluding.
- Using the coefficient without units check. All three coefficients are pure numbers; if your answer carries a unit (marks, minutes, FCFA), you have forgotten to divide by σ or by $Q_3 - Q_1$.

10.5 Skewness in Real Life**DID YOU KNOW?****Where do skewed distributions appear?**

- Incomes. In Cameroon, as in most countries, the distribution of household incomes is strongly positively skewed: most households earn modest incomes, while a small number earn very large ones. This is why the median income, not the mean, is usually quoted in economic reports — the mean is inflated by the extreme tail and would overstate the "typical" income.
- Examination marks. An easy paper produces a negatively skewed mark distribution (most marks high, a tail of low marks); a very difficult paper produces a positively

skewed one. Examiners use this to judge whether a paper was well calibrated.

- Waiting times and lifetimes. Waiting times at a hospital or a bank, and the lifetimes of electronic components, are typically positively skewed: many short waits, a few very long ones.

EXERCISE 11 Identifying and measuring skewness

The ages of workers in a small company in Douala are summarised by: mean = 36.2 years, median = 33.0 years, mode = 29.5 years, $\sigma = 9.6$ years, $Q_1 = 27$, $Q_2 = 33$, $Q_3 = 44$.

1. Without any calculation, state the expected direction of skewness from the positions of the mean, median and mode.
2. Compute Pearson's first coefficient of skewness.
3. Compute Pearson's second coefficient of skewness.
4. Compute Bowley's quartile coefficient of skewness.
5. Comment on the consistency of your answers.

SOLUTION Exercise 1

1. Since mode (29.5) < median (33.0) < mean (36.2), the distribution is positively skewed (tail towards the higher ages).
2.
$$S_k = \frac{\bar{x} - M_o}{\sigma} = \frac{36.2 - 29.5}{9.6} = \frac{6.7}{9.6} \approx 0.70.$$
3.
$$S_k = \frac{3(\bar{x} - M_e)}{\sigma} = \frac{3(36.2 - 33.0)}{9.6} = \frac{9.6}{9.6} = 1.00.$$
4.
$$S_{kQ} = \frac{Q_3 + Q_1 - 2Q_2}{Q_3 - Q_1} = \frac{44 + 27 - 66}{44 - 27} = \frac{5}{17} \approx 0.29.$$
5. All three coefficients are positive, confirming a positively skewed distribution: most workers are relatively young, with a tail of older workers. The magnitudes differ because the coefficients use different scales, but the sign — and hence the conclusion — is consistent.

EXERCISE 12 From summary statistics to a conclusion

Two schools enter candidates for the same examination. School X has $\bar{x} = 10.8$, $M_e = 10.5$, $\sigma = 2.4$. School Y has $Q_1 = 9$, $Q_2 = 12$, $Q_3 = 14$.

1. Compute Pearson's second coefficient for School X and interpret it.
2. Compute Bowley's coefficient for School Y and interpret it.
3. Which school's marks are more nearly symmetric? Justify.

SOLUTION Exercise 2

1. $S_k = \frac{3(10.8 - 10.5)}{2.4} = \frac{0.9}{2.4} = 0.375$. Slight positive skewness: a small tail of high marks.
2. $S_{kQ} = \frac{14 + 9 - 2(12)}{14 - 9} = \frac{-1}{5} = -0.2$. Slight negative skewness: a small tail of low marks.
3. School Y: $|S_{kQ}| = 0.2 < 0.375 = |S_k|$, so its marks are more nearly symmetric (both distributions are only mildly skewed, since both coefficients are below 0.5 in absolute value).

KEY POINTS Key points of the section

- A distribution is **symmetric** when its two halves are mirror images; then mean = median = mode.
- **Positive skewness**: long tail to the right, mode < median < mean. **Negative skewness**: long tail to the left, mean < median < mode.
- Pearson: $S_k = \frac{\bar{x} - M_o}{\sigma}$ or $S_k = \frac{3(\bar{x} - M_e)}{\sigma}$.
- Bowley (quartiles): $S_{kQ} = \frac{Q_3 + Q_1 - 2Q_2}{Q_3 - Q_1}$, always between -1 and 1.
- Sign > 0: right skew; sign < 0: left skew; value near 0: near symmetry. The tail, never the peak, gives the direction.

11 INTEGRATION ACTIVITY

11.1 Aims of the activity

This integration activity mobilises, in one authentic study, all the competences developed in this chapter. By the end of the activity, you should be able to:

- collect, organise and present real data in a **grouped frequency table** with correct **class boundaries**;
- draw and interpret a **histogram** and a **cumulative frequency curve** (ogive);
- calculate and interpret measures of **central tendency** (mean, median, mode) and of **dispersion** (standard deviation, quartiles, semi-interquartile range) for grouped data;
- construct and interpret **Laspeyres** and **Paasche** price indices;
- measure the **skewness** of a distribution using Pearson's coefficient and the quartile coefficient, and describe the shape of the distribution;
- communicate statistical findings in a short advisory report and defend them orally.

11.2 Situation

Statistical Audit of the GBHS Bafoussam Canteen.

The canteen of Government Bilingual High School Bafoussam serves about 400 students every school day. The manager, Madam Nkwain, has noticed that her revenue is unstable and that some students complain that “food has become too dear this year”. Before the next Parent-Teacher Association meeting, she asks the Upper Sixth Pure Mathematics with Statistics class to carry out a statistical audit of the canteen.

Your class accepts the mandate and collects the following data during one school week.

Data set A — Daily spending. Thirty volunteer students record the total amount (in FCFA) each of them spends at the canteen per day. The grouped results are:

Spending (FCFA)	Number of students
100 – 199	3
200 – 299	6
300 – 399	9
400 – 499	7
500 – 599	4
600 – 699	1

Data set B — Prices and quantities. For five representative items, the class records the average price and the quantity sold per day, for last year (base year, index 0) and this year (current year, index 1):

Item	p_0 (FCFA)	q_0	p_1 (FCFA)	q_1
Plate of rice	250	80	300	70
Plate of beans	150	60	200	55
Puff-puff	50	200	75	180
Bottle of water	100	150	150	140
Bread	100	90	125	85

The class must now analyse these data and write a one-page report advising Madam Nkwain.

11.3 Tasks

1. Rewrite Data set A as a grouped frequency table showing **class boundaries**, **mid-values**, **cumulative frequencies** and the columns fx and fx^2 .
2. Draw the **histogram** of the spending distribution, using the class boundaries on the horizontal axis.
3. Construct the **cumulative frequency table** and draw the **ogive** ("less than" curve). Use the ogive to estimate the median and the quartiles graphically.
4. Calculate the **mean**, **median** and **modal** daily spending, and state the **range** of the distribution.
5. Calculate the **variance** and the **standard deviation** of the daily spending.
6. Calculate Q_1 , Q_3 , the **interquartile range** and the **semi-interquartile range**. Compare your values of Q_1 , Q_2 , Q_3 with the graphical estimates of question 3.
7. Compute **Pearson's coefficient of skewness** and the **quartile coefficient of skewness**. Describe the shape of the spending distribution.
8. Using Data set B, compute the **Laspeyres price index** and the **Paasche price index** for this year (base: last year = 100). Interpret each index in one sentence for Madam Nkwain.
9. Write a one-page report (half a page of figures, half a page of advice) answering: How much does a typical student spend? Is spending spread out? Have prices really increased, and by how much? What do you advise the manager?
10. Each group presents its results in 5 minutes; the class compares the answers and discusses any discrepancies.

METHOD Strategy for a grouped-data audit

1. Start from the raw grouped table and build the working table: boundaries, mid-values x , cumulative frequencies F , then fx and fx^2 . Every later calculation reads from this table.
2. Draw the histogram (equal class widths here, so height = frequency) and the ogive

(cumulative frequency against upper class boundaries).

3. Compute the averages first (mean, median, mode), then the dispersion (variance, standard deviation, quartiles), then the skewness coefficients which combine the two.
4. Treat index numbers separately: build the four totals $\sum p_0 q_0$, $\sum p_1 q_0$, $\sum p_0 q_1$, $\sum p_1 q_1$ before applying the Laspeyres and Paasche formulae.
5. Finish by interpreting every number in the language of the situation (FCFA, students, prices) — a statistician who only computes has done half the job.


11.4 Model answer

1. Extended frequency table. The classes are given with class limits 100–199, 200–299, ... Since spending is recorded to the nearest franc, each class of width 100 has class boundaries obtained by shifting the limits by 0.5: the class 100–199 becomes 99.5–199.5, and so on. The mid-value of each class is the mean of its boundaries, e.g. $x = \frac{99.5 + 199.5}{2} = 149.5$. Cumulative frequencies are running totals of f ; the columns fx and fx^2 prepare the mean and variance.

Class	Boundaries	x	f	F	fx	fx^2
100–199	99.5–199.5	149.5	3	3	448.5	67050.75
200–299	199.5–299.5	249.5	6	9	1497.0	373501.50
300–399	299.5–399.5	349.5	9	18	3145.5	1099352.25
400–499	399.5–499.5	449.5	7	25	3146.5	1414351.75
500–599	499.5–599.5	549.5	4	29	2198.0	1207801.00
600–699	599.5–699.5	649.5	1	30	649.5	421850.25
Total			30		11085	4583907.5

Check: $\sum f = 30$ (all students counted), $\sum fx = 11085$, $\sum fx^2 = 4583907.5$.

2. Histogram. All class widths are equal to 100, so the height of each rectangle equals the frequency. Rectangles are drawn over the boundary intervals [99.5; 199.5], ..., [599.5; 699.5] with heights 3, 6, 9, 7, 4, 1. The tallest bar corresponds to the modal class 299.5–399.5.




Histogram of daily spending at the canteen

3. Cumulative frequency table and ogive. For a “less than” ogive, each cumulative frequency is plotted against the upper class boundary:

Upper boundary (FCFA)	Cumulative frequency F
99.5	0
199.5	3
299.5	9
399.5	18
499.5	25
599.5	29
699.5	30

(The point (99.5, 0) anchors the curve: nobody spends less than 99.5 FCFA.) Join the points with a smooth curve or straight segments. To read the quartiles, locate $N/4 = 7.5$, $N/2 = 15$ and $3N/4 = 22.5$ on the vertical axis, move across to the curve, then down to the spending axis.



Ogive of daily spending, with graphical readings of Q_1 , Q_2 , Q_3

Reading across from $F = 7.5, 15$ and 22.5 gives approximately $Q_1 \approx 275$, $Q_2 \approx 366$ and $Q_3 \approx 464$ FCFA. Graphical readings are estimates: anything within about ± 5 FCFA is acceptable.

4. Mean, median, mode, range. The mean of grouped data is

$$\bar{x} = \frac{\sum fx}{\sum f} = \frac{11085}{30} = 369.5 \text{ FCFA.}$$

For the median, $N/2 = 15$, which falls in the class 299.5–399.5 (the first class whose cumulative frequency, 18, reaches 15). With lower boundary $L = 299.5$, cumulative frequency before the class $F_b = 9$, class frequency $f_m = 9$ and class width $h = 100$:

$$Q_2 = L + \frac{\frac{N}{2} - F_b}{f_m} \times h = 299.5 + \frac{15 - 9}{9} \times 100 = 299.5 + 66.67 \approx 366.2 \text{ FCFA.}$$

The modal class is 299.5–399.5 (highest frequency, 9). With $\Delta_1 = 9 - 6 = 3$ (excess over the previous class) and $\Delta_2 = 9 - 7 = 2$ (excess over the next class):

$$M_o = L + \frac{\Delta_1}{\Delta_1 + \Delta_2} \times h = 299.5 + \frac{3}{3 + 2} \times 100 = 359.5 \text{ FCFA.}$$

Range $\approx 699.5 - 99.5 = 600$ FCFA (using the extreme class boundaries, since the exact raw values are unknown).

5. Variance and standard deviation. Using the computational formula for grouped data,

$$\sigma^2 = \frac{\sum fx^2}{\sum f} - \bar{x}^2 = \frac{4583907.5}{30} - 369.5^2 = 152796.92 - 136530.25 \approx 16266.67 \text{ FCFA}^2,$$

$$\sigma = \sqrt{16266.67} \approx 127.5 \text{ FCFA.}$$

So daily spending typically deviates from the mean by about 127 FCFA.

6. Quartiles. $N/4 = 7.5$ falls in the class 199.5–299.5 (cumulative frequency 9 is the first to reach 7.5), with $L = 199.5$, $F_b = 3$, $f = 6$:

$$Q_1 = 199.5 + \frac{7.5 - 3}{6} \times 100 = 199.5 + 75 = 274.5 \text{ FCFA.}$$

$3N/4 = 22.5$ falls in the class 399.5–499.5 (cumulative frequency 25), with $L = 399.5$, $F_b = 18$, $f = 7$:

$$Q_3 = 399.5 + \frac{22.5 - 18}{7} \times 100 = 399.5 + 64.29 \approx 463.8 \text{ FCFA.}$$

Interquartile range: $Q_3 - Q_1 = 463.8 - 274.5 = 189.3$ FCFA; semi-interquartile range: $\frac{189.3}{2} \approx 94.6$ FCFA. These calculated values agree very well with the ogive readings of question 3 (275, 366, 464), which confirms both the graph and the formulae.

7. Skewness. Pearson's second coefficient (used here because the mode is only an estimate for grouped data):

$$S_k = \frac{3(\bar{x} - Q_2)}{\sigma} = \frac{3(369.5 - 366.2)}{127.5} = \frac{9.9}{127.5} \approx 0.08.$$

Quartile (Bowley's) coefficient:

$$S_q = \frac{Q_3 + Q_1 - 2Q_2}{Q_3 - Q_1} = \frac{463.8 + 274.5 - 2(366.2)}{189.3} = \frac{5.9}{189.3} \approx 0.03.$$

Both coefficients are small and positive (each lies between -1 and $+1$, and values near 0 indicate near-symmetry). The spending distribution is therefore almost symmetrical, with a very slight positive skew: a few students spend much more than the others, which is consistent with the ordering $\bar{x} > Q_2 > M_o$ ($369.5 > 366.2 > 359.5$).

8. Index numbers. First compute the four totals needed by the formulae:

Item	p_0q_0	p_1q_0	p_0q_1	p_1q_1
Rice	20000	24000	17500	21000
Beans	9000	12000	8250	11000
Puff-puff	10000	15000	9000	13500
Water	15000	22500	14000	21000
Bread	9000	11250	8500	10625
Total	63000	84750	57250	77125

Laspeyres index (base-year quantities as weights):

$$L = \frac{\sum p_1q_0}{\sum p_0q_0} \times 100 = \frac{84750}{63000} \times 100 \approx 134.5.$$

Paasche index (current-year quantities as weights):

$$P = \frac{\sum p_1q_1}{\sum p_0q_1} \times 100 = \frac{77125}{57250} \times 100 \approx 134.7.$$

Interpretation. At last year's quantities, the same basket of items costs about 34.5% more this year (Laspeyres); at this year's quantities, the increase is about 34.7% (Paasche). The two indices are very close, so the conclusion is robust: **canteen prices have risen by roughly one third**, whatever quantities are used as weights.

9. Model report (summary). A typical student spends about 370 FCFA per day (mean 369.5, median 366.2 FCFA); half of the students spend between 274.5 and 463.8 FCFA. Spending is moderately spread out ($\sigma \approx 127.5$ FCFA, semi-interquartile range ≈ 94.6 FCFA) and the distribution is nearly symmetric with a slight positive skew: a small group of big spenders pulls the mean above the median. Prices have increased by about 34.5% on average (Laspeyres 134.5, Paasche 134.7), while quantities sold have fallen slightly for every item. Advice to the manager: the students' complaints are justified by the data; she should renegotiate with suppliers, keep at least one complete meal below 400 FCFA so that the median student can still afford lunch, and monitor the prices of water and rice, which contribute most to the index.

10. Class discussion. Groups compare their ogive readings, rounding choices and reports; small numerical differences (e.g. median 366 vs 366.2 FCFA) are acceptable provided the method — correct boundaries, correct interpolation formula, correct reading of the ogive — is sound.

WARNING Common errors to avoid

Using class limits instead of **class boundaries** in the median and quartile formulae (this shifts every answer by 0.5); forgetting to subtract $\bar{x}^2$ in the variance formula, or forgetting the square root at the end; swapping q_0 and q_1 between the Laspeyres and Paasche formulae; plotting the ogive against mid-values instead of upper class boundaries; concluding “strong skew” from a coefficient close to zero — near-zero means near-symmetric.

EXERCISE 13 Going further

The following year, the same audit is repeated and the new prices (in FCFA) are: rice 330, beans 210, puff-puff 80, water 150, bread 130, with quantities sold 75, 60, 190, 145, 88 respectively. Taking last year (Data set B, index 0) as base, compute the new Laspeyres and Paasche price indices and comment on the trend.

SOLUTION Exercise 1

The new totals are $\sum p_2 q_0 = 330(80) + 210(60) + 80(200) + 150(150) + 130(90) = 26400 + 12600 + 16000 + 22500 + 11700 = 89200$ and $\sum p_2 q_1 = 330(75) + 210(60) + 80(190) + 150(145) + 130(88) = 24750 + 12600 + 15200 + 21750 + 11440 = 85740$, while $\sum p_0 q_1 = 250(75) + 150(60) + 50(190) + 100(145) + 100(88) = 18750 + 9000 + 9500 + 14500 + 8800 = 60550$. Hence

$$L = \frac{89200}{63000} \times 100 \approx 141.6, \quad P = \frac{85740}{60550} \times 100 \approx 141.6.$$

Prices have continued to rise, from an index of about 134.5 to about 141.6: the increase is

slowing down but has not stopped.

12 METHODS AND WORKED EXAMPLES

12.1 Methods and Worked Examples

METHOD Constructing a Grouped Frequency Distribution

1. Identify the range: $R = x_{\max} - x_{\min}$.
2. Choose the number of classes k : Typically $5 \leq k \leq 20$. Sturges' rule $k \approx 1 + 3.322 \log_{10} n$ may be used as a guide.
3. Determine the class width c : $c = \frac{R}{k}$. Round up to a convenient number (e.g., 5, 10, 20).
4. Set class limits: Start at a value $\leq x_{\min}$ (preferably a multiple of c). Define **class boundaries** (real limits) to avoid gaps: for discrete data with unit gap, lower boundary = lower limit - 0.5, upper boundary = upper limit + 0.5. For continuous data, boundaries coincide with limits.
5. Tally and count frequencies f : Use a tally column for accuracy.
6. Compute mid-points x : $x = \frac{\text{Lower Limit} + \text{Upper Limit}}{2}$.
7. Check: $\sum f = n$.

WORKED EXAMPLE 23 Grouped Frequency Table – Mathematics Mock Scores

The scores (out of 100) of 50 Upper Sixth students in a Mathematics mock examination are:

42	67	55	78	34	89	51	63	72	48
59	61	75	82	45	68	54	70	38	85
57	64	79	49	66	52	73	80	41	69
58	62	77	46	71	53	76	81	44	65
56	60	74	47	70	50	69	83	43	66

Construct a grouped frequency distribution table with 7 classes. Include columns for Class Boundaries, Mid-points (x), Frequency (f), and Cumulative Frequency (CF).

SOLUTION

Solution:

1. Range: $x_{\max} = 89$, $x_{\min} = 34$, so $R = 89 - 34 = 55$.

2. Number of classes: $k = 7$ (given).
3. Class width: $c = \frac{55}{7} \approx 7.86$. Round up to $c = 8$.
4. Class limits: Start at 34: 34–41, 42–49, 50–57, 58–65, 66–73, 74–81, 82–89.
5. Class boundaries (discrete data, gap = 1): subtract 0.5 from each lower limit, add 0.5 to each upper limit.
6. Mid-points x : average of the class limits, e.g. $(34 + 41)/2 = 37.5$.
7. Tally and frequency: count the data points falling in each class.

Class Limits	Class Boundaries	Mid-point (x)	Tally	Freq (f)	Cum. Freq (CF)
34 – 41	33.5 – 41.5	37.5		4	4
42 – 49	41.5 – 49.5	45.5		8	12
50 – 57	49.5 – 57.5	53.5		8	20
58 – 65	57.5 – 65.5	61.5		11	31
66 – 73	65.5 – 73.5	69.5		9	40
74 – 81	73.5 – 81.5	77.5		7	47
82 – 89	81.5 – 89.5	85.5		3	50
Total				50	

Check: $\sum f = 4 + 8 + 8 + 11 + 9 + 7 + 3 = 50 = n$. ✓

WARNING Common Mistake

Using class limits (34–41) instead of boundaries (33.5–41.5) when drawing the Histogram or Ogive, or when applying interpolation formulas. Always use boundaries for graphs and for interpolation. The mid-point is calculated from the limits: $(34 + 41)/2 = 37.5$.

METHOD Drawing Histogram, Frequency Polygon and Ogive

1. Histogram: Horizontal axis = class boundaries. Vertical axis = frequency density (f/c) if class widths are unequal; frequency (f) if all widths are equal. Bars touch (no gaps).
2. Frequency polygon: Plot the points (mid-point x , frequency f) and join them with straight lines. Close the polygon by adding an extra class with $f = 0$ at each end (mid-

points $x_1 - c$ and $x_k + c$ joined to the horizontal axis.

3. Ogive (cumulative frequency curve): Plot the points (upper class boundary, CF). Join with a smooth curve (or straight segments). Start at (lower boundary of first class, 0).
4. Reading quantiles: For the Median (Q_2), Q_1 , Q_3 , locate $n/2$, $n/4$, $3n/4$ on the vertical axis, trace horizontally to the curve, then vertically down to the horizontal axis.

WORKED EXAMPLE 24 Graphical Representation and Estimation


Using the grouped frequency table from the previous example (Mock Scores):

1. Draw the Histogram and Frequency Polygon on the same axes.
2. Draw the Ogive (Cumulative Frequency Curve).
3. Use the Ogive to estimate the Median, Lower Quartile (Q_1), Upper Quartile (Q_3), and the Interquartile Range (IQR).
4. Estimate the number of students who scored above 75.

SOLUTION


Solution:

1. Histogram and Frequency Polygon. Class widths are equal ($c = 8$), so bar heights equal the frequencies. Mid-points: 37.5, 45.5, 53.5, 61.5, 69.5, 77.5, 85.5. Extra mid-points to close the polygon: $37.5 - 8 = 29.5$ and $85.5 + 8 = 93.5$ (both with $f = 0$).



Histogram (blue bars) and frequency polygon (orange line) for the mock scores

2. Ogive. Plot (upper boundary, CF): (41.5, 4), (49.5, 12), (57.5, 20), (65.5, 31), (73.5, 40), (81.5, 47), (89.5, 50), starting from (33.5, 0).



Ogive with graphical estimation of the median, quartiles, and CF at 75.5

3. Estimates from the Ogive (graphical reading), $n = 50$:

- Median (Q_2): position $n/2 = 25$. From the graph: ≈ 62 .
- Lower quartile (Q_1): position $n/4 = 12.5$. From the graph: ≈ 50 .
- Upper quartile (Q_3): position $3n/4 = 37.5$. From the graph: ≈ 71 .
- $IQR = Q_3 - Q_1 \approx 71 - 50 = 21$ marks.

4. Students scoring above 75: The score 75 lies in the class 74–81 (boundaries 73.5–81.5). We need CF at the boundary 75.5. Linear interpolation on the ogive segment from (73.5, 40) to (81.5, 47): gradient = $\frac{47 - 40}{81.5 - 73.5} = \frac{7}{8} = 0.875$.

$$CF(75.5) = 40 + 0.875 \times (75.5 - 73.5) = 40 + 1.75 = 41.75 \approx 42.$$

Students scoring at most 75: ≈ 42 . Hence students scoring above 75 = $50 - 42 = 8$.

METHOD Measures of Central Tendency: Mean, Median, Mode, Combined Mean

1. Mean ($\bar{x}$):

- Ungrouped: $\bar{x} = \frac{\sum x}{n}$.
- Grouped: $\bar{x} = \frac{\sum fx}{\sum f}$. For large numbers use coding (assumed mean): $\bar{x} = A + \frac{\sum fd}{\sum f} \times c$, where $d = \frac{x - A}{c}$, A = assumed mean (mid-point of a central class), c = class width.

2. Median (grouped): identify the median class, the first class where $CF \geq n/2$.

$$\text{Median} = L_m + \left(\frac{\frac{n}{2} - CF_{m-1}}{f_m} \right) \times c$$

L_m = lower boundary of median class, f_m = frequency of median class, CF_{m-1} = cumulative frequency before the median class.

3. Mode (grouped): identify the modal class (highest f).

$$\text{Mode} = L_{mo} + \left(\frac{\Delta_1}{\Delta_1 + \Delta_2} \right) \times c, \quad \Delta_1 = f_{mo} - f_{mo-1}, \quad \Delta_2 = f_{mo} - f_{mo+1}.$$

L_{mo} = lower boundary of the modal class.

4. Combined mean: for k groups with sizes n_i and means $\bar{x}_i$:

$$\bar{x}_{\text{combined}} = \frac{\sum n_i \bar{x}_i}{\sum n_i}$$

WORKED EXAMPLE 25 Central Tendency Calculations and Combined Mean

Using the Mock Scores grouped data (table of the first example):

1. Calculate the Mean using the coding method (assumed mean $A = 61.5$).
2. Calculate the Median and Mode using the interpolation formulas.
3. A second stream of 30 students sat the same examination and obtained a mean score of 58.2. Calculate the Combined Mean for all 80 students.

SOLUTION

Data recall: $n = 50$, $c = 8$. Median class: $n/2 = 25$; CF first reaches ≥ 25 at the class 58–65 (boundaries 57.5–65.5), so the median class is the 4th class. Modal class: highest $f = 11$, also the class 58–65.

1. Mean (coding method). Let $A = 61.5$ and $d = \frac{x - 61.5}{8}$.

Class	Mid-point x	Freq f	$d = \frac{x-61.5}{8}$	fd
34–41	37.5	4	-3	-12
42–49	45.5	8	-2	-16
50–57	53.5	8	-1	-8
58–65	61.5	11	0	0
66–73	69.5	9	1	9
74–81	77.5	7	2	14
82–89	85.5	3	3	9
Total		50		-4

$$\bar{x} = A + \frac{\sum fd}{\sum f} \times c = 61.5 + \frac{-4}{50} \times 8 = 61.5 - 0.64 = \mathbf{60.86}$$

2. Median. $L_m = 57.5$, $f_m = 11$, $CF_{m-1} = 20$, $c = 8$, $n/2 = 25$.

$$\text{Median} = 57.5 + \left(\frac{25 - 20}{11} \right) \times 8 = 57.5 + \frac{40}{11} \approx 57.5 + 3.64 = \mathbf{61.14}$$

3. Mode. $L_{mo} = 57.5$, $f_{mo} = 11$, $f_{mo-1} = 8$, $f_{mo+1} = 9$, $c = 8$. $\Delta_1 = 11 - 8 = 3$, $\Delta_2 = 11 - 9 = 2$.

$$\text{Mode} = 57.5 + \left(\frac{3}{3 + 2} \right) \times 8 = 57.5 + 4.8 = \mathbf{62.3}$$

4. Combined mean. Group 1: $n_1 = 50$, $\bar{x}_1 = 60.86$. Group 2: $n_2 = 30$, $\bar{x}_2 = 58.2$.

$$\bar{x}_{\text{comb}} = \frac{n_1 \bar{x}_1 + n_2 \bar{x}_2}{n_1 + n_2} = \frac{50(60.86) + 30(58.2)}{80} = \frac{3043 + 1746}{80} = \frac{4789}{80} \approx \mathbf{59.86}$$

METHOD Measures of Dispersion: Variance and Standard Deviation

1. Ungrouped data:

$$\sigma^2 = \frac{\sum (x - \bar{x})^2}{n} = \frac{\sum x^2}{n} - \bar{x}^2 \quad ; \quad \sigma = \sqrt{\sigma^2}$$

(For a sample, some conventions use $s^2 = \frac{\sum (x - \bar{x})^2}{n-1}$; at GCE Advanced Level the divisor n is used unless stated otherwise.)

2. Grouped data:

$$\sigma^2 = \frac{\sum f(x - \bar{x})^2}{\sum f} = \frac{\sum fx^2}{\sum f} - \bar{x}^2$$

Coding method (recommended in examinations): with $d = \frac{x - A}{c}$,

$$\sigma^2 = c^2 \left[\frac{\sum fd^2}{\sum f} - \left(\frac{\sum fd}{\sum f} \right)^2 \right] \quad ; \quad \sigma = c \sqrt{\frac{\sum fd^2}{\sum f} - \left(\frac{\sum fd}{\sum f} \right)^2}$$

3. Coefficient of variation: $CV = \frac{\sigma}{\bar{x}} \times 100\%$ (compares relative variability between data sets).

WORKED EXAMPLE 26 Variance and Standard Deviation – Grouped Data

For the Mock Scores data (50 students), calculate:

- the variance (σ^2) and standard deviation (σ) using the coding method ($A = 61.5$, $c = 8$);

- the coefficient of variation (CV);
- the percentage of students within one standard deviation of the mean (i.e., between $\bar{x} - \sigma$ and $\bar{x} + \sigma$), assuming uniform distribution within classes.

SOLUTION

- Variance and standard deviation. Extend the coding table with d^2 and fd^2 .

Class	x	f	d	d^2	fd^2
34–41	37.5	4	–3	9	36
42–49	45.5	8	–2	4	32
50–57	53.5	8	–1	1	8
58–65	61.5	11	0	0	0
66–73	69.5	9	1	1	9
74–81	77.5	7	2	4	28
82–89	85.5	3	3	9	27
Total		50	–4		140

$$\frac{\sum fd^2}{\sum f} = \frac{140}{50} = 2.8, \quad \left(\frac{\sum fd}{\sum f} \right)^2 = \left(\frac{-4}{50} \right)^2 = 0.0064$$

$$\sigma^2 = c^2 [2.8 - 0.0064] = 64 \times 2.7936 = 178.79$$

$$\sigma = \sqrt{178.79} \approx 13.37 \text{ marks}$$

- Coefficient of variation. With $\bar{x} = 60.86$:

$$CV = \frac{13.37}{60.86} \times 100\% \approx 21.97\%$$

Interpretation: the spread of scores is about 22% of the mean score — moderate variability.

- Percentage within $\bar{x} \pm \sigma$: $\bar{x} - \sigma = 60.86 - 13.37 = 47.49$ and $\bar{x} + \sigma = 60.86 + 13.37 = 74.23$. We need the frequency between 47.49 and 74.23, assuming uniformity within each class:

- Class 42–49 (boundaries 41.5–49.5, $f = 8$): included width = $49.5 - 47.49 = 2.01$ out of 8. Included frequency $\approx 8 \times \frac{2.01}{8} = 2.01$.
- Class 50–57: fully inside. Frequency = 8.
- Class 58–65: fully inside. Frequency = 11.
- Class 66–73: fully inside. Frequency = 9.
- Class 74–81 (boundaries 73.5–81.5, $f = 7$): included width = $74.23 - 73.5 = 0.73$ out of 8. Included frequency $\approx 7 \times \frac{0.73}{8} = 0.64$.

Total included frequency $\approx 2.01 + 8 + 11 + 9 + 0.64 = 30.65 \approx 31$ students. Percentage $= \frac{31}{50} \times 100\% = 62\%$. (For a Normal distribution one would expect about 68%; this data set is slightly skewed.)

METHOD Quantiles for Grouped Data: Quartiles, Deciles, Percentiles

1. General formula for a quantile Q at cumulative position p : locate the first class where $CF \geq p$, then

$$Q = L_Q + \left(\frac{p - CF_{Q-1}}{f_Q} \right) \times c$$

where L_Q = lower class boundary of the quantile class, CF_{Q-1} = cumulative frequency before that class, f_Q = frequency of that class, c = class width.

2. Positions:

- Quartiles: Q_1 at $n/4$, $Q_2 \equiv$ Median at $n/2$, Q_3 at $3n/4$.
- Deciles: D_k at $kn/10$, for $k = 1, \dots, 9$.
- Percentiles: P_k at $kn/100$, for $k = 1, \dots, 99$.

3. Interquartile range: $IQR = Q_3 - Q_1$.

4. Semi-interquartile range (quartile deviation): $SIQR = \frac{Q_3 - Q_1}{2}$.

WORKED EXAMPLE 27 Quantiles, IQR and SIQR Calculation

Using the Mock Scores data ($n = 50$), calculate by formula:

1. Q_1, Q_3, D_5, P_{90} ;
2. the Interquartile Range (IQR) and Semi-Interquartile Range (SIQR);
3. the 70th percentile (P_{70}).

SOLUTION

Reference table (boundaries, f , CF):

Class Boundaries	f	CF
33.5 – 41.5	4	4
41.5 – 49.5	8	12
49.5 – 57.5	8	20
57.5 – 65.5	11	31
65.5 – 73.5	9	40
73.5 – 81.5	7	47
81.5 – 89.5	3	50

1. Quartiles and other quantiles ($c = 8$):

- Q_1 : position = $50/4 = 12.5$. First $CF \geq 12.5$: class 41.5–49.5, with $L = 41.5$, $CF_{prev} = 4$, $f = 8$.

$$Q_1 = 41.5 + \frac{12.5 - 4}{8} \times 8 = 41.5 + 8.5 = 50.0$$

- Q_3 : position = $3(50)/4 = 37.5$. First $CF \geq 37.5$: class 65.5–73.5, with $L = 65.5$, $CF_{prev} = 31$, $f = 9$.

$$Q_3 = 65.5 + \frac{37.5 - 31}{9} \times 8 = 65.5 + \frac{52}{9} \approx 65.5 + 5.78 = 71.28$$

- D_5 : position = $5(50)/10 = 25$. Class 57.5–65.5, $L = 57.5$, $CF_{prev} = 20$, $f = 11$.

$$D_5 = 57.5 + \frac{25 - 20}{11} \times 8 \approx 57.5 + 3.64 = 61.14 \quad (\text{equals the Median, as expected})$$

- P_{90} : position = $90(50)/100 = 45$. First $CF \geq 45$: class 73.5–81.5, $L = 73.5$, $CF_{prev} = 40$, $f = 7$.

$$P_{90} = 73.5 + \frac{45 - 40}{7} \times 8 = 73.5 + \frac{40}{7} \approx 73.5 + 5.71 = 79.21$$

2. IQR and SIQR:

$$IQR = Q_3 - Q_1 = 71.28 - 50.00 = 21.28, \quad SIQR = \frac{IQR}{2} = \frac{21.28}{2} = 10.64$$

3. P_{70} : Position = $70(50)/100 = 35$. First $CF \geq 35$: class 57.5–65.5, $L = 57.5$, $CF_{prev} = 20$, $f = 11$.

$$P_{70} = 57.5 + \frac{35 - 20}{11} \times 8 = 57.5 + \frac{120}{11} \approx 57.5 + 10.91 = 68.41$$

METHOD Index Numbers: Laspeyres and Paasche Price Indices

1. Notation: base year (0), current year (n); price p , quantity q . So p_0, q_0 are base-year values and p_n, q_n current-year values.
2. Price relative of an item: $I = \frac{p_n}{p_0} \times 100$.
3. Laspeyres price index (base-weighted): uses base-year quantities q_0 as weights.

$$L = \frac{\sum p_n q_0}{\sum p_0 q_0} \times 100$$

Advantage: fixed weights, easy to compare over time. Disadvantage: ignores changes in consumption patterns (substitution bias).

4. Paasche price index (current-weighted): uses current-year quantities q_n as weights.

$$P = \frac{\sum p_n q_n}{\sum p_0 q_n} \times 100$$

Advantage: reflects the current consumption pattern. Disadvantage: weights change every period, so pure price change is harder to isolate.

5. Typical relationship: when consumers substitute away from goods whose prices rise fastest, $L > P$.

WORKED EXAMPLE 28 Laspeyres and Paasche Indices – Household Commodities

A household consumes three commodities: rice, beans and oil. Data for the base year (2020) and the current year (2024):

Commodity	Price 2020 (p_0)	Qty 2020 (q_0)	Price 2024 (p_n)	Qty 2024 (q_n)
Rice (kg)	400 FCFA	50	600 FCFA	40
Beans (kg)	800 FCFA	20	1000 FCFA	25
Oil (litre)	1200 FCFA	10	1500 FCFA	8

Calculate:

- the Laspeyres price index for 2024 (base 2020 = 100);
- the Paasche price index for 2024 (base 2020 = 100);
- interpret the results and explain why they differ.

SOLUTION

Working table (values in FCFA):

Item	$p_n q_0$	$p_0 q_0$	$p_n q_n$	$p_0 q_n$
Rice	$600 \times 50 = 30\,000$	$400 \times 50 = 20\,000$	$600 \times 40 = 24\,000$	$400 \times 40 = 16\,000$
Beans	$1000 \times 20 = 20\,000$	$800 \times 20 = 16\,000$	$1000 \times 25 = 25\,000$	$800 \times 25 = 20\,000$
Oil	$1500 \times 10 = 15\,000$	$1200 \times 10 = 12\,000$	$1500 \times 8 = 12\,000$	$1200 \times 8 = 9\,600$
Sum	65 000	48 000	61 000	45 600

1. Laspeyres index (L):

$$L = \frac{\sum p_n q_0}{\sum p_0 q_0} \times 100 = \frac{65\,000}{48\,000} \times 100 \approx 135.4$$

2. Paasche index (P):

$$P = \frac{\sum p_n q_n}{\sum p_0 q_n} \times 100 = \frac{61\,000}{45\,600} \times 100 \approx 133.8$$

3. Interpretation:

- Both indices exceed 100, indicating a general price increase (inflation) of about 34–35% between 2020 and 2024.
- Here Laspeyres (135.4) > Paasche (133.8).
- Reason: the household reduced its consumption of rice (50 → 40) and oil (10 → 8), whose prices rose sharply (by 50% and 25%), and increased its consumption of beans (20 → 25), whose price rose less (25%).
- Laspeyres keeps the old quantities (heavy weight on the now-expensive rice and oil), so it overstates the increase in the cost of living.
- Paasche uses the new quantities (less weight on the expensive items), reflecting the substitution effect, and therefore gives a lower index.

METHOD Skewness: Pearson's and Quartile Coefficients

1. Symmetry: Mean = Median = Mode $\Rightarrow$ symmetrical distribution (zero skewness).
2. Positive skew (right skew): long tail on the right; typically Mean > Median > Mode.
3. Negative skew (left skew): long tail on the left; typically Mean < Median < Mode.
4. Pearson's coefficient of skewness:

- $Sk_p = \frac{\text{Mean} - \text{Mode}}{\sigma}$ (when the mode is well defined);
- $Sk_p = \frac{3(\text{Mean} - \text{Median})}{\sigma}$ (more common; uses the empirical relation Mean - Mode $\approx 3(\text{Mean} - \text{Median})$).

Values usually lie between -3 and +3. $Sk_p > 0$: positive skew; $Sk_p < 0$: negative skew.

5. Quartile (Bowley's) coefficient of skewness:

$$Sk_Q = \frac{Q_3 + Q_1 - 2Q_2}{Q_3 - Q_1} = \frac{Q_3 + Q_1 - 2 \text{ Median}}{IQR}$$

Always between -1 and +1; robust, since quartiles are not affected by extreme values.

- $Sk_Q = 0$: symmetrical ($Q_3 - Q_2 = Q_2 - Q_1$);
- $Sk_Q > 0$: positive skew ($Q_3 - Q_2 > Q_2 - Q_1$);
- $Sk_Q < 0$: negative skew.

WORKED EXAMPLE 29 Skewness Analysis of the Mock Scores

Using the statistics calculated for the Mock Scores ($n = 50$): $\bar{x} = 60.86$, Median = 61.14, Mode = 62.30, $\sigma = 13.37$, $Q_1 = 50.00$, $Q_2 = 61.14$, $Q_3 = 71.28$, $IQR = 21.28$.

1. Determine the direction of skewness by comparing the Mean, Median and Mode.
2. Calculate Pearson's coefficient of skewness (median-based formula).
3. Calculate the Quartile coefficient of skewness.
4. Comment on the consistency of the two measures.

SOLUTION

1. Direction of skewness: Mean (60.86) < Median (61.14) < Mode (62.30). **Conclusion:** the distribution is negatively skewed (left-skewed): the tail stretches towards the lower scores, while the bulk of the distribution lies at the higher scores.

2. Pearson's coefficient (Sk_p):

$$Sk_p = \frac{3(\bar{x} - \text{Median})}{\sigma} = \frac{3(60.86 - 61.14)}{13.37} = \frac{-0.84}{13.37} \approx -0.063$$

(The mode-based version gives $\frac{60.86 - 62.30}{13.37} \approx -0.108$, also negative; the median-based formula is preferred as it is more stable.)

3. Quartile coefficient (Sk_Q):

$$Sk_Q = \frac{Q_3 + Q_1 - 2Q_2}{Q_3 - Q_1} = \frac{71.28 + 50.00 - 2(61.14)}{21.28} = \frac{-1.00}{21.28} \approx -0.047$$

4. Comment:

- Both coefficients are negative, confirming negative skewness.
- Both are close to zero ($|Sk_p| \approx 0.06$, $|Sk_Q| \approx 0.05$), so the distribution is nearly symmetrical.
- The slight negative skew reflects a small number of weaker students pulling the mean just below the median.
- Pearson's coefficient uses the mean and standard deviation (sensitive to extreme values); the quartile coefficient is robust. Their agreement strengthens the conclusion.

KEY POINTS Exam Tip

When asked to “comment on the skewness” of a distribution, always:

1. state the order of Mean, Median and Mode;
2. state the direction (positive / negative / approximately symmetrical);
3. compute a coefficient (Pearson’s or the quartile coefficient);
4. interpret its magnitude (“highly skewed”, “moderately skewed”, “approximately symmetric”).

13 EXERCISES

13.1 I check my knowledge

EXERCISE 14 MCQ: Data Types and Collection

Which of the following variables is continuous?

1. The number of students in a classroom.
2. The height of a student measured in centimetres.
3. The number of cars passing a checkpoint in an hour.
4. The number of defective items in a batch of 50.

EXERCISE 15 True/False with Justification

State whether each statement is true or false. Justify your answer briefly.

1. The mode of a data set is always unique.
2. The standard deviation of a set of values is always less than or equal to the range.
3. In a grouped frequency distribution, the median class is the class with the highest frequency.
4. The Laspeyres price index uses current year quantities as weights.

EXERCISE 16 Definitions and Terminology

Define the following terms precisely:

1. **Class boundaries** in a grouped frequency distribution.
2. **Frequency density** and why it is used when drawing a histogram with unequal class widths.
3. **Semi-interquartile range** (quartile deviation).
4. **Positively skewed distribution** in terms of the relative positions of mean, median and mode.

EXERCISE 17 Direct Calculations

The marks (out of 20) of 10 students in a test are:

12, 15, 14, 12, 18, 15, 13, 16, 15, 14.

Calculate:

1. The mean.
2. The median.
3. The mode.
4. The variance and standard deviation (use the formula for ungrouped data).

13.2 I practise**EXERCISE 18** Grouped Data: Mean, Median, Mode by Interpolation

The table below shows the distribution of masses (in kg) of 80 packets of rice sold in a market in Douala.

Mass (kg)	1.0–1.4	1.5–1.9	2.0–2.4	2.5–2.9	3.0–3.4	3.5–3.9
Frequency	8	15	22	18	12	5

1. Construct the cumulative frequency table (using true class boundaries).
2. Estimate the mean mass using the coding method with an assumed mean of 2.2 kg.
3. Estimate the median mass by linear interpolation.
4. Estimate the modal mass by linear interpolation (use the histogram method formula).

EXERCISE 19 Histogram with Unequal Class Widths and Frequency Polygon

The following data gives the ages of 100 patients attending a health centre in Yaoundé.

Age (years)	0–9	10–19	20–29	30–49	50–69
Number of patients	15	20	25	30	10

1. Calculate the frequency density for each class.
2. Draw a histogram to represent the data (use graph paper; show frequency density on the vertical axis).

3. Superimpose the frequency polygon on the same axes.
4. Use the histogram to estimate the mode graphically.

EXERCISE 20 Cumulative Frequency Curve (Ogive) and Percentiles

The table shows the distribution of monthly incomes (in thousands of CFA francs) of 120 workers.

Income (1000 FCFA)	50–59	60–69	70–79	80–89	90–99	100–109
Frequency	10	18	32	35	15	10

1. Construct the cumulative frequency table (using upper class boundaries).
2. Draw the ogive (cumulative frequency curve).
3. From the ogive, estimate:
 - (a) The median income.
 - (b) The first quartile (Q_1) and third quartile (Q_3).
 - (c) The 90th percentile (P_{90}).
 - (d) The semi-interquartile range.

EXERCISE 21 Combined Mean and Comparison of Consistency

Two classes, A and B, sat the same Mathematics test. Class A has 35 students with a mean mark of 12.4 and a standard deviation of 2.1. Class B has 45 students with a mean mark of 10.8 and a standard deviation of 3.0.

1. Calculate the combined mean mark for the two classes together.
2. Which class shows more consistent performance? Justify your answer using the coefficient of variation.
3. If the pass mark is 10, and assuming the marks are roughly normally distributed, which class would you expect to have a higher pass rate? Explain briefly.

13.3 I prepare for the GCE

EXERCISE 22 GCE Style: Full Analysis of a Grouped Distribution

[20 marks] The table below shows the distribution of marks obtained by 80 candidates in a GCE Advanced Level Pure Mathematics with Statistics paper.

Marks	10–19	20–29	30–39	40–49	50–59	60–69	70–79
Frequency	5	10	15	20	18	8	4

1. Construct the cumulative frequency table using true class boundaries. [2 marks]
2. Calculate the mean mark using the coding method (choose a suitable assumed mean). [4 marks]
3. Calculate the standard deviation of the marks. [4 marks]
4. Estimate the median mark by linear interpolation. [3 marks]
5. Estimate the modal mark by linear interpolation. [3 marks]
6. Draw a histogram to represent the data (use frequency density if necessary) and superimpose the frequency polygon. [2 marks]
7. Comment on the performance of the candidates based on the values obtained. [2 marks]

EXERCISE 23 GCE Style: Index Numbers and Skewness

[18 marks] A household buys four food items: Rice, Beans, Fish, and Oil. The prices (in FCFA per kg) and quantities consumed (in kg per month) for the base year 2023 and the current year 2024 are given below.

Item	Price 2023 (p_0)	Quantity 2023 (q_0)	Price 2024 (p_1)	Quantity 2024 (q_1)
Rice	600	50	720	55
Beans	800	20	900	18
Fish	2500	5	3000	4
Oil	1200	10	1300	12

1. Compute the simple aggregate price index for 2024 with 2023 as base year. [2 marks]
2. Compute the Laspeyres price index for 2024. [3 marks]
3. Compute the Paasche price index for 2024. [3 marks]
4. Interpret the difference between the Laspeyres and Paasche indices in this context. [2 marks]

A distribution of weekly wages has the following summary statistics: Mean = 52,000 FCFA, Median = 50,000 FCFA, Mode = 46,000 FCFA, Standard deviation = 8,000 FCFA.

5. Calculate Pearson's first coefficient of skewness (using mode). [2 marks]
6. Calculate Pearson's second coefficient of skewness (using median). [2 marks]
7. State the direction of skewness and justify your answer. [2 marks]
8. Given that $Q_1 = 44,000$ FCFA and $Q_3 = 58,000$ FCFA, compute the quartile coefficient of skewness and the semi-interquartile range. [2 marks]

EXERCISE 24 GCE Style: Mixed Problem – Reconstruction and Questionnaire Critique

[17 marks]

Part A: Reconstruction of a Frequency Distribution [10 marks] A frequency distribution of the heights (in cm) of 60 plants was partially destroyed. The remaining data is:

Height (cm)	10–14	15–19	20–24	25–29	30–34
Frequency	5	f_1	18	f_2	7

It is known that the total frequency is 60 and the mean height is 22.5 cm.

1. Find the missing frequencies f_1 and f_2 . [4 marks]
2. Using the completed table, calculate the variance and standard deviation of the heights. [4 marks]
3. Estimate the median height by linear interpolation. [2 marks]

Part B: Questionnaire Critique [7 marks] A student designs the following questionnaire to investigate the favourite subjects of Upper Sixth students in a school. Identify three sources of bias or poor design in the questionnaire and rewrite the faulty questions to eliminate the bias.

1. "Don't you agree that Mathematics is the most important subject?" (Yes / No)
2. "Which subject do you like: Mathematics, Physics, or Chemistry?" (Tick one)
3. "How many hours do you spend studying each day?" (0–1, 1–2, 2–3, 3–4, 4+)
4. "What is your favourite subject? _____" (Open ended)

14 SOLUTIONS TO THE EXERCISES

14.1 I check my knowledge

SOLUTION Exercise 1 — MCQ: Data Types and Collection

Answer: (b). Height can take any value within an interval (e.g. 172.4 cm), so it is **con-
tinuous**. The other three variables can only take whole-number values (they result from counting), so they are **discrete**.

SOLUTION Exercise 2 — True/False with Justification

1. False. A data set may have no mode (all values distinct), one mode, or several modes (bimodal, multimodal). Example: 2, 2, 5, 5, 7 has two modes.
2. True. The standard deviation measures spread about the mean while the range measures the total spread from minimum to maximum. In fact it can be proved that $\sigma \leq \frac{\text{range}}{2}$ for every data set, so certainly $\sigma \leq \text{range}$.
3. False. The median class is the class containing the $\frac{N}{2}$ th value (located from the cumulative frequency). The class with the highest frequency is the modal class.
4. False. The Laspeyres index uses base year quantities q_0 as weights: $L = \frac{\sum p_1 q_0}{\sum p_0 q_0} \times 100$. It is the Paasche index that uses current year quantities q_1 .

SOLUTION Exercise 3 — Definitions and Terminology

1. The **class boundaries** (true class limits) are the values that separate two consecutive classes without any gap. For a class 10–19 followed by 20–29 (data recorded to the nearest unit), the class boundaries are 9.5–19.5 and 19.5–29.5.
2. The **frequency density** of a class is $\text{frequency density} = \frac{\text{frequency}}{\text{class width}}$. In a histogram, the area of each bar (not its height) must be proportional to the frequency; when class widths are unequal, plotting frequency density on the vertical axis guarantees this.
3. The **semi-interquartile range** is $Q = \frac{Q_3 - Q_1}{2}$, where Q_1 and Q_3 are the first and third quartiles. It measures the spread of the middle 50% of the data.

4. A distribution is **positively skewed** (skewed to the right) when it has a long tail towards the high values; then $\text{mode} < \text{median} < \text{mean}$.

SOLUTION Exercise 4 — Direct Calculations

- $\sum x = 144$, so $\bar{x} = \frac{144}{10} = 14.4$.
- Ordered data: 12, 12, 13, 14, 14, 15, 15, 15, 16, 18. Median = $\frac{14 + 15}{2} = 14.5$.
- Mode = 15 (it occurs 3 times).
- $\sum x^2 = 2104$, so

$$\sigma^2 = \frac{\sum x^2}{n} - \bar{x}^2 = \frac{2104}{10} - 14.4^2 = 210.4 - 207.36 = 3.04, \quad \sigma = \sqrt{3.04} \approx 1.74.$$

14.2 I practise

SOLUTION Exercise 5 — Grouped Data: Mean, Median, Mode by Interpolation

True class boundaries: 0.95–1.45, 1.45–1.95, ..., 3.45–3.95; midpoints: 1.2, 1.7, 2.2, 2.7, 3.2, 3.7; class width $c = 0.5$.

- Cumulative frequencies (against upper class boundaries): 8, 23, 45, 63, 75, 80.
- With $A = 2.2$ and $u = \frac{x - A}{0.5}$: the values of u are -2, -1, 0, 1, 2, 3 and

$$\sum fu = 8(-2) + 15(-1) + 22(0) + 18(1) + 12(2) + 5(3) = 26.$$

$$\bar{x} = A + c \frac{\sum fu}{N} = 2.2 + 0.5 \times \frac{26}{80} = 2.2 + 0.1625 \approx 2.36 \text{ kg.}$$
- $\frac{N}{2} = 40$ falls in the class 1.95–2.45 (cumulative frequency 23 before it, frequency 22):

$$\text{Median} = 1.95 + \frac{40 - 23}{22} \times 0.5 = 1.95 + 0.386 \approx 2.34 \text{ kg.}$$
- Modal class: 1.95–2.45 (frequency 22). With $\Delta_1 = 22 - 15 = 7$ and $\Delta_2 = 22 - 18 = 4$:

$$\text{Mode} = 1.95 + \frac{\Delta_1}{\Delta_1 + \Delta_2} \times 0.5 = 1.95 + \frac{7}{11} \times 0.5 \approx 2.27 \text{ kg.}$$

SOLUTION Exercise 6 — Histogram with Unequal Class Widths and Frequency Polygon

1. Class boundaries: $-0.5-9.5$, $9.5-19.5$, $19.5-29.5$, $29.5-49.5$, $49.5-69.5$; widths 10, 10, 10, 20, 20.

Class	Width	Frequency	Frequency density
0–9	10	15	1.5
10–19	10	20	2.0
20–29	10	25	2.5
30–49	20	30	1.5
50–69	20	10	0.5

2. The histogram has bars over $[-0.5, 9.5]$, $[9.5, 19.5]$, $[19.5, 29.5]$, $[29.5, 49.5]$, $[49.5, 69.5]$ with heights 1.5, 2.0, 2.5, 1.5, 0.5 respectively. Note how the wide classes (30–49 and 50–69) get shorter bars than their raw frequencies would suggest: this is exactly why frequency density is needed.
3. The frequency polygon joins the points (4.5, 1.5), (14.5, 2.0), (24.5, 2.5), (39.5, 1.5), (59.5, 0.5) (midpoints of the class boundaries against frequency density), extended to the axis at both ends.
4. Drawing the diagonal lines in the tallest bar (the standard histogram construction) gives a modal age of approximately 24 years.

SOLUTION Exercise 7 — Cumulative Frequency Curve (Ogive) and Percentiles

1. Upper class boundaries and cumulative frequencies:

Upper boundary	59.5	69.5	79.5	89.5	99.5	109.5
Cumulative frequency	10	28	60	95	110	120

2. Plot the points (59.5, 10), (69.5, 28), (79.5, 60), (89.5, 95), (99.5, 110), (109.5, 120) and join them with a smooth S-shaped curve.
3. Reading from the ogive (values confirmed by linear interpolation):

(a) Median at $\frac{N}{2} = 60$: median ≈ 79.5 thousand FCFA.

(b) Q_1 at $\frac{N}{4} = 30$: $Q_1 = 69.5 + \frac{30 - 28}{32} \times 10 \approx 70.1$; Q_3 at $\frac{3N}{4} = 90$: $Q_3 = 79.5 + \frac{90 - 60}{35} \times 10 \approx 88.1$ (thousand FCFA).

- (c) P_{90} at $0.9 \times 120 = 108$: since the cumulative frequency 108 lies between 95 (at 89.5) and 110 (at 99.5),

$$P_{90} = 89.5 + \frac{108 - 95}{15} \times 10 = 89.5 + 8.67 \approx 98.2 \text{ thousand FCFA.}$$

- (d) Semi-interquartile range = $\frac{Q_3 - Q_1}{2} \approx \frac{88.1 - 70.1}{2} = 9.0$ thousand FCFA.

SOLUTION Exercise 8 — Combined Mean and Comparison of Consistency

1. Combined mean:

$$\bar{x} = \frac{n_A \bar{x}_A + n_B \bar{x}_B}{n_A + n_B} = \frac{35(12.4) + 45(10.8)}{80} = \frac{434 + 486}{80} = \frac{920}{80} = 11.5.$$

2. Coefficient of variation: $CV = \frac{\sigma}{\bar{x}} \times 100\%$.

$$CV_A = \frac{2.1}{12.4} \times 100\% \approx 16.9\%, \quad CV_B = \frac{3.0}{10.8} \times 100\% \approx 27.8\%.$$

Since $CV_A < CV_B$, Class A is more consistent (its marks are less dispersed relative to its mean).

3. Class A: mean 12.4 with small spread 2.1, so the pass mark 10 lies about 1.1 standard deviations below the mean: the great majority pass. Class B: mean 10.8 with spread 3.0, so the pass mark lies only about 0.27 standard deviations below the mean: a substantial fraction of the candidates fall below 10. We therefore expect a higher pass rate in Class A.

14.3 I prepare for the GCE

SOLUTION Exercise 9 — GCE Style: Full Analysis of a Grouped Distribution

Midpoints: 14.5, 24.5, 34.5, 44.5, 54.5, 64.5, 74.5; class width $c = 10$; boundaries 9.5–19.5, ..., 69.5–79.5.

1. Cumulative frequencies: 5, 15, 30, 50, 68, 76, 80 against upper boundaries 19.5, 29.5, ..., 79.5. [2 marks: 1 for boundaries, 1 for cumulative frequencies]
2. Take $A = 44.5$, $u = \frac{x - 44.5}{10}$, so $u = -3, -2, -1, 0, 1, 2, 3$:

$$\sum fu = 5(-3) + 10(-2) + 15(-1) + 20(0) + 18(1) + 8(2) + 4(3) = -4.$$

$$\bar{x} = 44.5 + 10 \times \frac{-4}{80} = 44.5 - 0.5 = 44.0.$$

[4 marks: 1 for coding, 1 for $\sum fu$, 1 for formula, 1 for answer]

$$3. \sum fu^2 = 5(9) + 10(4) + 15(1) + 20(0) + 18(1) + 8(4) + 4(9) = 186.$$

$$\sigma^2 = c^2 \left[\frac{\sum fu^2}{N} - \left(\frac{\sum fu}{N} \right)^2 \right] = 100 \left[\frac{186}{80} - \left(\frac{-4}{80} \right)^2 \right] = 100(2.325 - 0.0025) = 232.25,$$

$$\sigma = \sqrt{232.25} \approx 15.2.$$

[4 marks: 1 for $\sum fu^2$, 1 for formula, 1 for variance, 1 for σ]

$$4. \frac{N}{2} = 40 \text{ lies in the class } 39.5\text{--}49.5 \text{ (cumulative frequency 30 before it, frequency 20):}$$

$$\text{Median} = 39.5 + \frac{40 - 30}{20} \times 10 = 39.5 + 5 = 44.5.$$

[3 marks: 1 for locating class, 1 for formula, 1 for answer]

$$5. \text{ Modal class } 39.5\text{--}49.5; \Delta_1 = 20 - 15 = 5, \Delta_2 = 20 - 18 = 2:$$

$$\text{Mode} = 39.5 + \frac{5}{5 + 2} \times 10 = 39.5 + 7.14 \approx 46.6.$$

[3 marks: 1 for modal class, 1 for formula, 1 for answer]

6. All classes have equal width 10, so the histogram bars have heights equal to the frequencies over the intervals $[9.5, 19.5], \dots, [69.5, 79.5]$. The frequency polygon joins $(14.5, 5), (24.5, 10), (34.5, 15), (44.5, 20), (54.5, 18), (64.5, 8), (74.5, 4)$. [2 marks: 1 for histogram, 1 for polygon]

7. The mean mark (44.0) is below 50, and the large standard deviation (15.2) shows marks are widely spread. Since mean < median < mode, the distribution is slightly negatively skewed: a tail of weak candidates pulls the mean down, while the bulk of candidates scored in the 40–59 region. Overall performance is mediocre, with a small group of very weak candidates. [2 marks: 1 for level of performance, 1 for shape/spread comment]

Examiner's expectations: correct use of class boundaries, a clear coding table, interpolation formulas quoted before substitution, and a comment that uses the computed statistics (not a vague remark).

SOLUTION
Exercise 10 — GCE Style: Index Numbers and Skewness

1. Simple aggregate price index:

$$I = \frac{\sum p_1}{\sum p_0} \times 100 = \frac{720 + 900 + 3000 + 1300}{600 + 800 + 2500 + 1200} \times 100 = \frac{5920}{5100} \times 100 \approx 116.1.$$

[2 marks: 1 for formula and sums, 1 for answer]

2. Laspeyres (base year quantities):

$$\sum p_1 q_0 = 720(50) + 900(20) + 3000(5) + 1300(10) = 36000 + 18000 + 15000 + 13000 = 82000$$

$$\sum p_0 q_0 = 600(50) + 800(20) + 2500(5) + 1200(10) = 30000 + 16000 + 12500 + 12000 = 70500$$

$$L = \frac{82000}{70500} \times 100 \approx 116.3.$$

[3 marks: 1 for each aggregate sum, 1 for index]

3. Paasche (current year quantities):

$$\sum p_1 q_1 = 720(55) + 900(18) + 3000(4) + 1300(12) = 39600 + 16200 + 12000 + 15600 = 83400,$$

$$\sum p_0 q_1 = 600(55) + 800(18) + 2500(4) + 1200(12) = 33000 + 14400 + 10000 + 14400 = 71800,$$

$$P = \frac{83400}{71800} \times 100 \approx 116.2.$$

[3 marks: 1 for each aggregate sum, 1 for index]

4. Both indices show a price rise of about 16%. The Laspeyres index (116.3) is slightly higher than the Paasche index (116.2) because the household slightly reduced its consumption of the items whose prices rose most (fish, beans) and increased consumption of the item whose price rose least (oil). Laspeyres keeps the old basket and therefore marginally overstates the rise in the cost of living, while Paasche reflects the new basket. [2 marks]

5. Pearson's first coefficient:

$$sk = \frac{\text{mean} - \text{mode}}{\sigma} = \frac{52000 - 46000}{8000} = \frac{6000}{8000} = 0.75.$$

[2 marks: 1 for formula, 1 for answer]

6. Pearson's second coefficient:

$$sk = \frac{3(\text{mean} - \text{median})}{\sigma} = \frac{3(52000 - 50000)}{8000} = \frac{6000}{8000} = 0.75.$$

[2 marks: 1 for formula, 1 for answer]

7. Both coefficients are positive, and mode < median < mean: the distribution is positively skewed (a long tail of high wages pulls the mean above the median and mode). [2 marks: 1 for direction, 1 for justification]

8. Quartile coefficient of skewness:

$$Sk_Q = \frac{Q_3 + Q_1 - 2Q_2}{Q_3 - Q_1} = \frac{58000 + 44000 - 2(50000)}{58000 - 44000} = \frac{2000}{14000} \approx 0.14,$$

confirming positive skewness. Semi-interquartile range:

$$Q = \frac{Q_3 - Q_1}{2} = \frac{58000 - 44000}{2} = 7000 \text{ FCFA.}$$

[2 marks: 1 for each quantity]

Examiner's expectations: index formulas stated with correct weights, aggregate sums shown explicitly, and skewness conclusions supported both by the sign of the coefficients and by the ordering of mean, median and mode.

SOLUTION Exercise 11 — GCE Style: Mixed Problem – Reconstruction and Questionnaire Critique

Part A. Midpoints: 12, 17, 22, 27, 32.

1. Total frequency: $5 + f_1 + 18 + f_2 + 7 = 60 \Rightarrow f_1 + f_2 = 30$. Mean condition: $\sum fx = 60(22.5) = 1350$:

$$5(12) + 17f_1 + 18(22) + 27f_2 + 7(32) = 1350 \Rightarrow 680 + 17f_1 + 27f_2 = 1350 \Rightarrow 17f_1 + 27f_2 = 670.$$

Substituting $f_2 = 30 - f_1$: $17f_1 + 27(30 - f_1) = 670 \Rightarrow 810 - 10f_1 = 670 \Rightarrow f_1 = 14, f_2 = 16$.
[4 marks: 1 per equation, 1 per value]

2. $\sum fx^2 = 5(144) + 14(289) + 18(484) + 16(729) + 7(1024) = 720 + 4046 + 8712 + 11664 + 7168 = 32310$.

$$\sigma^2 = \frac{\sum fx^2}{N} - \bar{x}^2 = \frac{32310}{60} - 22.5^2 = 538.5 - 506.25 = 32.25, \quad \sigma = \sqrt{32.25} \approx 5.68 \text{ cm.}$$

[4 marks: 1 for $\sum fx^2$, 1 for formula, 1 for variance, 1 for σ]

3. Cumulative frequencies: 5, 19, 37, 53, 60. $\frac{N}{2} = 30$ lies in the class 19.5–24.5:

$$\text{Median} = 19.5 + \frac{30 - 19}{18} \times 5 = 19.5 + 3.06 \approx 22.6 \text{ cm.}$$

[2 marks: 1 for locating class, 1 for answer]


Part B. Any three of the following [7 marks: up to 2 marks per fault identified and corrected, 1 mark for general presentation]:

1. Leading question: it pushes the respondent to agree. Rewrite: "In your opinion, how important is Mathematics compared with your other subjects?" (Very important / Important / Not important).

2. Restricted choice list: only three science subjects are offered, so students who prefer other subjects cannot answer honestly. Rewrite: “Which of the following subjects do you like most?” with a fuller list (Mathematics, Physics, Chemistry, Biology, Economics, Literature, ...) plus an “Other: ____” option.
3. Overlapping response categories: a student studying exactly 1, 2, 3 or 4 hours fits two boxes. Rewrite with non-overlapping intervals: 0–0.9, 1–1.9, 2–2.9, 3–3.9, 4 or more (or “less than 1, at least 1 but less than 2, ...”).
4. The open-ended question (4) is acceptable in principle, though answers may be hard to classify; offering a tick-box list with an “Other” option makes processing easier.

Examiner’s expectations: in Part A, the two equations must be set up from the total frequency and the mean before solving; in Part B, each fault must be named (leading question, restricted options, overlapping classes) and a corrected version of the question must actually be written out.

15 SUMMARY SHEET



Mind map of the chapter: from raw data to summary measures, index numbers and the shape of a distribution.

FICHE BILAN L'essentiel à connaître pour l'examen

Key definitions.

- **Statistics**: the science of the collection, organisation, presentation, analysis and interpretation of numerical data. A **population** is the whole group under study; a **sample** is a representative part of it; each member is a **statistical unit**; a measurable characteristic is a **variable**.
- **Qualitative data** (attributes that cannot be measured numerically: colour, sex, tribe) vs **quantitative data** (numerical: **discrete** – obtained by counting, e.g. number of children; or **continuous** – obtained by measuring, e.g. height, mass).
- **Primary data** (collected first-hand by interview, observation, questionnaire or experiment) vs **secondary data** (taken from already existing records, e.g. school registers, MINEDUB reports).

- **Raw data:** data as collected, before any organisation. A **frequency** f is the number of times a value occurs; $\sum f = N$ is the total frequency.
- For a grouped distribution: **class limits** (the stated end values), **class boundaries** (the true end values, e.g. 9.5–19.5 for the class 10–19), **class width** c (difference between successive lower boundaries), **class mark** (midpoint) $x_i = \frac{\text{lower limit} + \text{upper limit}}{2}$.
- **Cumulative frequency:** the running total of frequencies; plotted against upper class boundaries it gives the **ogive** (cumulative frequency curve).
- **Index number:** a ratio comparing the value of a variable in a given period to its value in a **base period**, usually expressed as a percentage.
- **Skewness:** the degree of asymmetry (lack of symmetry) of a distribution.

Formulas to know.

- **Mean (ungrouped):** $\bar{x} = \frac{\sum x}{n}$; **grouped:** $\bar{x} = \frac{\sum f_i x_i}{\sum f_i}$; with assumed mean A and coding $u_i = \frac{x_i - A}{c}$: $\bar{x} = A + c \frac{\sum f_i u_i}{\sum f_i}$.
- **Combined mean** of two groups: $\bar{x} = \frac{n_1 \bar{x}_1 + n_2 \bar{x}_2}{n_1 + n_2}$ (extends naturally to k groups: $\bar{x} = \frac{\sum n_i \bar{x}_i}{\sum n_i}$).
- **Median (grouped):** $M_e = L + \frac{\frac{N}{2} - F}{f} \times c$, where L = lower boundary of the median class, F = cumulative frequency before it, f = its frequency, c = its width.
- **Mode (grouped):** $M_o = L + \frac{\Delta_1}{\Delta_1 + \Delta_2} \times c$, with $\Delta_1 = f_m - f_{\text{before}}$, $\Delta_2 = f_m - f_{\text{after}}$, where f_m is the largest frequency (modal class).
- **Range** = largest value – smallest value.
- **Variance (ungrouped):** $\sigma^2 = \frac{\sum (x - \bar{x})^2}{n} = \frac{\sum x^2}{n} - \bar{x}^2$; standard deviation $\sigma = \sqrt{\sigma^2}$.
- **Variance (grouped):** $\sigma^2 = \frac{\sum f_i x_i^2}{\sum f_i} - \bar{x}^2$, or by coding $\sigma^2 = c^2 \left(\frac{\sum f_i u_i^2}{\sum f_i} - \left(\frac{\sum f_i u_i}{\sum f_i} \right)^2 \right)$.
- **Combined variance of two groups:** $\sigma^2 = \frac{n_1(\sigma_1^2 + d_1^2) + n_2(\sigma_2^2 + d_2^2)}{n_1 + n_2}$ with $d_i = \bar{x}_i - \bar{x}$.
- **Quartiles by interpolation:** $Q_k = L + \frac{\frac{kN}{4} - F}{f} \times c$ ($k = 1, 2, 3$; note $Q_2 = M_e$); similarly deciles D_k using $\frac{kN}{10}$ and percentiles P_k using $\frac{kN}{100}$. **Interquartile range** = $Q_3 - Q_1$; **semi-**

$$\text{interquartile range} = \frac{Q_3 - Q_1}{2}.$$

- Simple price index: $\frac{p_n}{p_0} \times 100$. **Laspeyres index**: $L = \frac{\sum p_n q_0}{\sum p_0 q_0} \times 100$ (base-period quantities as weights). **Paasche index**: $P = \frac{\sum p_n q_n}{\sum p_0 q_n} \times 100$ (current-period quantities as weights). Weighted index number: $\frac{\sum wI}{\sum w}$, where I are the individual indices and w their weights.
- **Pearson's coefficient of skewness**: $S_k = \frac{\bar{x} - M_o}{\sigma}$; when the mode is awkward to obtain, the empirical approximation $S_k \approx \frac{3(\bar{x} - M_e)}{\sigma}$ is used. **Quartile coefficient of skewness**: $S_Q = \frac{Q_3 + Q_1 - 2Q_2}{Q_3 - Q_1}$. $S > 0$: positively skewed (tail to the right); $S < 0$: negatively skewed (tail to the left); $S = 0$: symmetrical.

Methods.

- Grouping data: find the range, choose the number of classes (usually between 5 and 15), fix a convenient class width c , write the classes, **tally** the raw data, then count the frequencies and check that $\sum f = N$.
- Histogram: rectangles drawn on the class boundaries with no gaps, area proportional to frequency (for equal class widths, height = frequency). The **frequency polygon** joins the midpoints of the tops of the bars (or the points (x_i, f_i)).
- Ogive: plot cumulative frequency against upper class boundaries, join the points, and read the median at $\frac{N}{2}$, Q_1 at $\frac{N}{4}$ and Q_3 at $\frac{3N}{4}$ on the cumulative axis.
- Pie chart angle for each category: $\frac{\text{frequency}}{\text{total frequency}} \times 360^\circ$. Bar chart: separated bars of equal width, height = frequency. Line graph: values plotted against time (or another ordered variable) and joined.

Common mistakes.

- Using class limits instead of class boundaries for histograms, ogives and interpolation formulas.
- Forgetting to multiply by the class width c (for the mean) or by c^2 (for the variance) when decoding a coded calculation.
- Confusing $\sum fx^2$ with $(\sum fx)^2$ in the variance formula.
- Mixing up Laspeyres (base-period quantities q_0) and Paasche (current-period quantities q_n).
- Reading the median or quartiles from the frequency polygon instead of the ogive.

- Taking $\frac{N}{2}$ as the median itself instead of the position of the median.
- Giving a negative variance or a standard deviation larger than the range without checking the arithmetic.

GCE tips.

- Show the full frequency table with columns $x_i, f_i x_i, f_i x_i^2$ (or the coding columns $u_i, f_i u_i, f_i u_i^2$): method marks are awarded for correct table work.
- Quote the interpolation formula before substituting, and give the final answer with units and sensible rounding (usually 1 or 2 decimal places, or 3 significant figures).
- For skewness questions, state the sign and interpret it in words: for a positively skewed distribution, mean > median > mode and the tail stretches to the right; the reverse holds for negative skewness.
- Quick checks: a variance can never be negative; an index number near 100 means little change from the base period; an index of 125 means a 25% increase.
- In pie chart questions, check that your angles add up to 360° (allowing for rounding).



Signed • OnBuch+

ONBUCH+ TEACHING TEAM
FOLLOWS THE OFFICIAL MINESEC SYLLABUS

OnBuch+