

BIOLOGY • UPPER SIXTH

UPPER SIXTH — BIOLOGY

CHAPTER 7

BIOTECHNOLOGY

SUGGESTED TIME: 6 PERIODS

IN THIS CHAPTER

Ce chapitre couvre l'ensemble du programme de biotechnologie pour la Terminale : des procédés traditionnels de fermentation aux technologies de l'ADN recombinant, en passant par la thérapie génique et les applications médicales, agricoles et environnementales. Il prépare l'élève à analyser, évaluer et concevoir des applications biotechnologiques pertinentes pour le contexte camerounais et mondial.

- Expliquer les principes biochimiques des fermentations traditionnelles et identifier les micro-organismes impliqués.
- Décrire les mécanismes d'homéostasie et illustrer comment la biotechnologie intervient dans le soutien et la locomotion (prothèses, biomatériaux).
- Analyser les stratégies de correction génique (thérapie génique, édition CRISPR) et en évaluer les avantages et limites.
- Maîtriser les étapes clés de la technologie de l'ADN recombinant : enzymes de restriction, vecteurs, clonage, sélection et amplification par PCR.
- Évaluer les applications majeures de la biotechnologie (médicaments, OGM, bioremédiation, biocarburants) sous l'angle scientifique, éthique et réglementaire.
- Concevoir un protocole simple de clonage ou de détection d'un gène d'intérêt en utilisant les outils moléculaires standards.



DOWNLOAD ON- BUCH

Free • Google Play
AI tutor, past papers,
results



GO ONBUCH+

All signed courses, unlimited AI
tutor, PDF sheets — OnBuch+
tab in the app

THIS SIGNED COURSE CONTAINS

COURSE

complete, illustrated,
step by step

METHODS

and worked examples

EXERCISES

exam style + solutions

SUMMARY SHEET

the essentials to
remember



Signed • OnBuch+

ONBUCH+ TEACHING TEAM
FOLLOWS THE OFFICIAL MINESEC SYLLABUS

OnBuch+

CONTENTS

- | | |
|--|---|
| 1. 1. Biotechnologie traditionnelle : fermentation et transformation alimentaire | 5. 5. Applications majeures de la biotechnologie : santé, agriculture, environnement, industrie |
| 2. 2. Biotechnologie au service de l'homéostasie, du soutien et de la locomotion | 6. Integration activity |
| 3. 3. Correction et remédiation génique : thérapie génique et édition du génome | 7. Methods and worked examples |
| 4. 4. Technologie de l'ADN recombinant et altération génique | 8. Exercises |
| | 9. Solutions to the exercises |
| | 10. Summary sheet |

CHAPTER OBJECTIVES

- Expliquer les principes biochimiques des fermentations traditionnelles et identifier les micro-organismes impliqués.
- Décrire les mécanismes d'homéostasie et illustrer comment la biotechnologie intervient dans le soutien et la locomotion (prothèses, biomatériaux).
- Analyser les stratégies de correction génique (thérapie génique, édition CRISPR) et en évaluer les avantages et limites.
- Maîtriser les étapes clés de la technologie de l'ADN recombinant : enzymes de restriction, vecteurs, clonage, sélection et amplification par PCR.
- Évaluer les applications majeures de la biotechnologie (médicaments, OGM, bioremédiation, biocarburants) sous l'angle scientifique, éthique et réglementaire.
- Concevoir un protocole simple de clonage ou de détection d'un gène d'intérêt en utilisant les outils moléculaires standards.
- Argumenter sur les enjeux de biosécurité et de propriété intellectuelle liés aux biotechnologies modernes.

PREREQUISITES

- Structure et fonction des cellules procaryotes et eucaryotes (Lower Sixth).
- Bases de la génétique mendélienne et moléculaire (réplication, transcription, traduction).
- Cinétique enzymatique et rôle des enzymes dans le métabolisme.
- Notions de microbiologie : culture, identification, croissance bactérienne.
- Principes de base de la chimie organique (glucides, acides aminés, acides nucléiques).

1 BIOTECHNOLOGIE TRADITIONNELLE : FERMENTATION ET TRANSFORMATION ALIMENTAIRE

Dans les marchés de Yaoundé, l'odeur du vin de palme et du foutou fermenté rappelle que la biotechnologie fait partie du quotidien camerounais depuis des siècles. Aujourd'hui, les mêmes principes permettent de produire de l'insuline humaine dans des bactéries et de créer des plantes résistantes à la sécheresse. Comment passer de la fermentation artisanale à l'édition du génome ? Ce chapitre explore cette évolution, des méthodes traditionnelles aux outils moléculaires de pointe.

Le problème central de cette première section est le suivant : comment des micro-organismes invisibles transforment-ils nos aliments, et comment l'Homme a-t-il appris, d'abord empiriquement puis scientifiquement, à contrôler ces transformations ?

1.1 Définitions : biotechnologie traditionnelle et moderne

Observez un vendeur de vin de palme à Buea : il récolte la sève du palmier, la laisse reposer quelques heures, et le liquide sucré devient pétillant et alcoolisé. Personne n'a ajouté de levure : les micro-organismes présents naturellement dans l'air et sur l'écorce ont fait le travail. C'est cela, la biotechnologie traditionnelle : une utilisation **empirique** du vivant, transmise de génération en génération, sans connaissance des mécanismes moléculaires sous-jacents.

DEFINITION Biotechnologie

La **biotechnologie** est l'utilisation d'organismes vivants (micro-organismes, cellules végétales ou animales) ou de leurs composants (enzymes, gènes) pour produire des biens et des services utiles à l'Homme : aliments, boissons, médicaments, procédés industriels.

DEFINITION Biotechnologie traditionnelle vs moderne

- La **biotechnologie traditionnelle** repose sur des pratiques empiriques ancestrales : fermentation spontanée des aliments et boissons, sélection paysanne des semences, fabrication artisanale de fromages. Elle ne modifie pas directement le patrimoine génétique des organismes.
- La **biotechnologie moderne** utilise les outils de la biologie moléculaire : ADN recombinant, génie génétique, culture cellulaire, édition du génome. Elle permet de modifier de façon ciblée le génome des organismes (insuline humaine produite par *Escherichia coli*, plantes transgéniques).

KEY POINTS Lien entre les deux

La biotechnologie moderne n'est pas une rupture mais un approfondissement : elle repose sur les mêmes phénomènes biologiques (métabolisme microbien, reproduction, expression des gènes) que la tradition a exploités aveuglément pendant des millénaires.


1.2 La fermentation : une voie métabolique anaérobie

Pourquoi la sève de palme devient-elle alcoolisée ? Parce que les levures, en l'absence d'oxygène, doivent tout de même produire de l'ATP pour survivre. Elles utilisent alors une voie de secours : la fermentation.

DEFINITION Fermentation


La **fermentation** est le catabolisme anaérobie incomplet d'un substrat organique (généralement un glucide) par des micro-organismes. Elle se déroule entièrement dans le cytoplasme, sans chaîne respiratoire, et ne produit que peu d'ATP ; le substrat n'est que partiellement oxydé, d'où la formation de produits finaux encore riches en énergie (éthanol, acide lactique).

Étape commune : la glycolyse. Quel que soit le type de fermentation, le glucose est d'abord dégradé par la glycolyse :




Le problème est le suivant : la glycolyse consomme du NAD^+ et produit du NADH, H^+ . Or la cellule ne possède qu'un stock limité de NAD^+ . En anaérobiose, la chaîne respiratoire ne peut pas régénérer le NAD^+ . La fermentation résout ce problème en **réoxydant le NADH, H^+ au niveau du pyruvate**.

Fermentation alcoolique (levures, ex. *Saccharomyces cerevisiae*) :




Bilan global : $\text{C}_6\text{H}_{12}\text{O}_6 \rightarrow 2 \text{C}_2\text{H}_5\text{OH} + 2 \text{CO}_2 + 2 \text{ATP}$.

Fermentation lactique homolactique (bactéries lactiques, ex. *Lactobacillus*, *Streptococcus*) :



Bilan global : $\text{C}_6\text{H}_{12}\text{O}_6 \rightarrow 2 \text{C}_3\text{H}_6\text{O}_3 + 2 \text{ATP}$.



Voies métaboliques de la fermentation alcoolique et lactique à partir du glucose. Dans les deux cas, la régénération du NAD⁺ permet à la glycolyse de se poursuivre.

PROPERTY Rendement énergétique de la fermentation

Une molécule de glucose ne fournit que 2 ATP en fermentation (glycolyse seule), contre environ 30 à 32 ATP en respiration aérobie complète. La fermentation est donc une voie énergétiquement peu rentable : les micro-organismes fermentaires doivent consommer de grandes quantités de substrat, ce qui explique l'accumulation rapide des produits finaux (alcool, acide) dans le milieu.

WARNING Erreur fréquente

Ne dites jamais que la fermentation « produit de l'énergie sans oxygène donc sans ATP » : elle produit bien 2 ATP par glucose, par phosphorylation au niveau du substrat (glycolyse), et non par phosphorylation oxydative. Autre piège : le CO₂ n'est produit que par la fermentation alcoolique, pas par la fermentation lactique homolactique.

1.3 Les grands procédés fermentaires camerounais

Le Cameroun possède un patrimoine fermentaire remarquable, qui illustre parfaitement la diversité des micro-organismes et des substrats.

Produit	Substrat et principe	Micro-organismes dominants
Vin de palme	Sève de palmier riche en sucres ; fermentation alcoolique spontanée rapide	Levures (<i>Saccharomyces</i> spp.), puis bactéries acétiques si exposition à l'air
Foléré (bissap)	Calices d'hibiscus ; boisson fermentée légèrement acide	Bactéries lactiques, levures
Foutou / bâton de manioc	Tubercules de manioc fermentés en eau (réttage) puis pilés	Bactéries lactiques (<i>Lactobacillus</i> , <i>Leuconostoc</i>) ; détoxification de l'acide cyanhydrique
Ngondo	Graines de courge fermentées, condiment	Bactéries lactiques, moisissures de surface
Fromage de soja	Lait de soja caillé puis fermenté	Moisissures (<i>Rhizopus</i> , <i>Aspergillus</i>), bactéries lactiques

Trois rôles essentiels de la fermentation alimentaire sont à retenir :

- **Conservation** : l'acide lactique et l'éthanol abaissent le pH et inhibent les germes de détérioration.
- **Amélioration nutritionnelle et gustative** : arômes nouveaux, vitamines synthétisées par les micro-organismes, prédigestion des protéines et glucides.
- **Détoxification** : la fermentation du manioc élimine les hétérosides cyanogénétiques (qui libèrent de l'acide cyanhydrique, toxique), rendant le tubercule consommable.




Diagramme de flux du procédé traditionnel de transformation du manioc en foutou.

DID YOU KNOW? Le manioc, un aliment rendu sûr par la biotechnologie traditionnelle

Le manioc contient naturellement des composés cyanogénétiques. Sans fermentation ni trempage, sa consommation régulière peut provoquer des intoxications chroniques. Les procédés traditionnels camerounais (réttage, fermentation) constituent donc une véritable biotechnologie de sécurité alimentaire, mise au point empiriquement bien avant que la chimie n'explique le phénomène.

1.4 Facteurs influençant la fermentation

La fermentation est une réaction enzymatique conduite par des êtres vivants : elle dépend donc étroitement des conditions du milieu.

- **Température** : chaque micro-organisme possède un optimum (environ 25–30 °C pour les levures du vin de palme). Une chaleur excessive dénature les enzymes ; le froid ralentit le métabolisme.
- **pH** : les bactéries lactiques acidifient le milieu, ce qui finit par inhiber leur propre croissance (autolimitation) et protège le produit des contaminants.
- **Concentration en substrat** : un excès de sucre crée une pression osmotique élevée qui peut inhiber les levures ; un déficit limite la production.
- **Anaérobiose** : la fermentation alcoolique exige l'absence d'oxygène ; à l'air libre, l'éthanol est oxydé en acide acétique par les bactéries acétiques — c'est pourquoi le vin de palme « tourne au vinaigre » en un ou deux jours.
- **Contamination** : des micro-organismes indésirables (moisissures toxigènes, bactéries pathogènes) peuvent s'installer si l'hygiène est mauvaise.

WARNING Risque de contamination

En l'absence de contrôle hygiénique, des pathogènes dangereux peuvent se développer dans les aliments fermentés ou mal conservés : *Clostridium botulinum* (toxine botulique, mortelle, en milieu anaérobie mal acidifié), *Salmonella*, moisissures productrices d'aflatoxines. L'acidification rapide par les ferments lactiques est la principale barrière naturelle : un produit qui ne s'acidifie pas correctement ne doit jamais être consommé.

1.5 Du contrôle artisanal au contrôle quantitatif

Traditionnellement, le producteur juge sa fermentation par les **organes des sens** : le goût (acidité, douceur résiduelle), l'odeur (arômes alcooliques ou acides), l'aspect (mousse, dégagement gazeux) et la densité perçue du liquide. Ces méthodes sont rapides mais subjectives et non reproductibles.

Les premières approches quantitatives ont introduit des instruments simples :

- le **densimètre** (ou pèse-mout) : la densité du moût diminue au fur et à mesure que le sucre est converti en éthanol ;
- la **titration acide** : dosage de l'acidité totale par une solution de soude de concentration connue, en présence d'un indicateur coloré, pour suivre la fermentation lactique.

METHOD Dosage de l'éthanol par distillation suivie de densimétrie

1. Prélever un volume connu V de la boisson fermentée (ex. 100 mL de vin de palme).
2. Distiller : l'éthanol, plus volatil que l'eau, s'évapore en premier ; recueillir le distillat.
3. Mesurer la densité ρ du distillat avec un densimètre (ou pycnomètre).
4. Convertir la densité en degré alcoolique à l'aide d'une table de correspondance éthanol-eau (tables officielles).
5. Calculer le titre alcoométrique volumique : $T = \frac{V_{\text{éthanol}}}{V_{\text{échantillon}}} \times 100$ (en % vol).

WORKED EXAMPLE 1 Calcul d'un titre alcoolique

Un élève prélève 200 mL de vin de palme. Après distillation et densimétrie, il détermine que le distillat contient 8 mL d'éthanol pur rapportés à 100 mL de boisson. Quel est le titre alcoolique du vin de palme ?

SOLUTION

Le titre alcoométrique volumique est le volume d'éthanol pur contenu dans 100 volumes de boisson :

$$T = \frac{V_{\text{éthanol}}}{V_{\text{boisson}}} \times 100 = \frac{8}{100} \times 100 = 8 \% \text{ vol.}$$

Le vin de palme titre donc 8 % vol, valeur cohérente pour un vin de palme bien fermenté (la teneur augmente avec la durée de fermentation, puis l'acétification peut la faire baisser).

1.6 De l'artisanat à l'industrie : standardisation et bioréacteurs

Les procédés industriels modernes reprennent exactement les mêmes réactions, mais en les **contrôlant** :

Critère	Procédé traditionnel	Procédé industriel
Micro-organismes	Flore spontanée, variable	Souches sélectionnées, pures, conservées en banques de souches
Conditions	Ambiantes, non contrôlées	Température, pH, oxygénation régulés en bioréacteur
Qualité du produit	Variable d'un lot à l'autre	Standardisée, contrôlée (densité, titration, chromatographie)
Rendement et échelle	Faible, familial ou artisanal	Élevé, cuves de plusieurs milliers de litres
Sécurité	Empirique	Contrôles microbiologiques systématiques

KEY POINTS À retenir de cette section

- La biotechnologie traditionnelle exploite empiriquement le métabolisme microbien ; la biotechnologie moderne le pilote au niveau moléculaire.
- La fermentation est un catabolisme anaérobie incomplet : 2 ATP par glucose, avec régénération du NAD^+ par réduction du pyruvate (éthanol + CO_2 chez les levures, lactate chez les bactéries lactiques).
- Les produits camerounais fermentés (vin de palme, foutou, foléré, ngondo, fromage de soja) illustrent la conservation, l'amélioration gustative et la détoxification des aliments.
- La maîtrise de la température, du pH, de l'anaérobiose et de l'hygiène conditionne la qualité et la sécurité du produit ; l'industrie ajoute souches sélectionnées et bioréacteurs.

EXERCISE 1 Fermentation et rendement énergétique

1. Une levure dégrade 5 moles de glucose par fermentation alcoolique. Combien de moles d'ATP, d'éthanol et de CO_2 sont produites ?
2. Comparer avec la dégradation complète de 5 moles de glucose par respiration aérobie (rendement : 30 ATP par glucose).
3. Expliquer pourquoi un moût très sucré peut ralentir la fermentation.

SOLUTION Exercice 1

1. D'après le bilan $C_6H_{12}O_6 \rightarrow 2 C_2H_5OH + 2 CO_2 + 2 ATP$: pour 5 mol de glucose, on obtient $5 \times 2 = 10$ mol d'ATP, 10 mol d'éthanol et 10 mol de CO_2 .
2. Respiration : $5 \times 30 = 150$ mol d'ATP, soit 15 fois plus que la fermentation. La fermentation est donc très peu rentable énergétiquement.
3. Une concentration élevée en sucre augmente la pression osmotique du milieu : l'eau sort des cellules de levure (plasmolyse), ce qui inhibe leur métabolisme et ralentit la fermentation.

2 BIOTECHNOLOGIE AU SERVICE DE L'HOMÉOSTASIE, DU SOUTIEN ET DE LA LOCOMOTION

Dans la section précédente, nous avons vu comment l'être humain exploite des cellules vivantes depuis des millénaires pour fermenter et transformer les aliments. La biotechnologie traditionnelle utilise des organismes entiers dans leur état naturel. La biotechnologie moderne va bien plus loin : elle ne se contente pas de transformer la nourriture, elle peut réparer, remplacer ou assister le corps humain lui-même. Lorsqu'un patient diabétique à Yaoundé s'injecte de l'insuline produite par des bactéries génétiquement modifiées, lorsqu'un amputé à Douala remarcher grâce à une prothèse myoélectrique, ou lorsqu'un ligament déchiré est reconstruit sur un échafaudage de polymère biodégradable, la biotechnologie agit directement sur trois fonctions fondamentales étudiées plus tôt dans ce cours : l'**homéostasie**, le **soutien** et la **locomotion**. Cette section montre comment.

2.1 Rappel sur l'homéostasie : les boucles de rétroaction négative

Intuition première. Imaginez le système de climatisation d'une salle de classe à Yaoundé. Lorsque la température dépasse la valeur de consigne, un capteur la détecte, un contrôleur déclenche le refroidissement, et la température redescend. Lorsqu'elle devient trop basse, le refroidissement s'arrête. Le système s'oppose à tout écart. Votre corps fonctionne exactement de la même manière pour la glycémie, la température corporelle et le pH sanguin.

DEFINITION Homéostasie


L'**homéostasie** est le maintien d'un milieu intérieur stable (le milieu intérieur : plasma sanguin et liquide interstitiel) dans des limites étroites, malgré les variations du milieu extérieur ou de l'activité métabolique. Elle est réalisée par des mécanismes d'autorégulation, principalement des **boucles de rétroaction négative**, impliquant des récepteurs (capteurs), un centre de coordination et des effecteurs.

PROPERTY **Rétroaction négative**

Dans une **boucle de rétroaction négative**, la réponse réduit ou annule le stimulus initial : la sortie du système inhibe sa propre production. Exemple : une hausse de la glycémie déclenche la sécrétion d'insuline ; l'insuline abaisse la glycémie ; la baisse de la glycémie stoppe alors la sécrétion d'insuline. (Le mécanisme détaillé est exigible à l'examen et doit être décrit avec précision au niveau Advanced Level.)

Trois boucles à maîtriser parfaitement.

- **Glycémie (taux de glucose sanguin).** La glycémie à jeun normale est d'environ $0,8-1,0 \text{ g L}^{-1}$ (soit environ $4,4-5,5 \text{ mmol L}^{-1}$). Après un repas riche en glucides (pensez à un plat de nouilles avec du mince), la glycémie augmente : les cellules β des îlots de Langerhans sécrètent de l'**insuline**, qui stimule la capture du glucose par les muscles et le tissu adipeux et son stockage sous forme de glycogène dans le foie (glycogénèse). Au cours du jeûne, les cellules α sécrètent du **glucagon**, qui déclenche la glycogénolyse et la néoglucogénèse, remontant la glycémie. Les deux hormones sont antagonistes.
- **Température corporelle.** L'hypothalamus agit comme un thermostat. La surchauffe déclenche la vasodilatation des artérioles cutanées et la transpiration (l'évaporation évacue la chaleur) ; le refroidissement déclenche la vasoconstriction, les frissons (contractions musculaires libérant de la chaleur) et une augmentation de la thyroxine et de l'adrénaline qui élève le métabolisme de base.
- **pH sanguin.** Le pH sanguin est maintenu près de 7,4 par des systèmes tampons (le couple $\text{H}_2\text{CO}_3/\text{HCO}_3^-$ est le principal), par les poumons (l'expiration de CO_2 élimine de l'acide) et par les reins (excrétion de H^+ et réabsorption de HCO_3^-).



Rétroaction négative du contrôle de la glycémie. L'insuline abaisse la glycémie ; le glucagon la remonte. La réponse s'oppose toujours à l'écart initial.

Jeûne : cellule α
→ glycogène
remonte (bo)

WARNING **Confusion entre les deux hormones**

Une erreur classique au GCE est d'écrire que le glucagon abaisse la glycémie, ou que l'insuline est sécrétée pendant le jeûne. Retenir : l'insuline met le glucose inside les cellules et en réserve (glycémie baisse) ; le glucagon est libéré quand le glucose est parti (gone en anglais, glycémie monte). De plus, la rétroaction négative ne signifie pas que la variable est constante : elle oscille autour d'un point de consigne.

2.2 Insuline humaine recombinante : du gène à la protéine thérapeutique

Lorsque la boucle de l'insuline fait défaut, le résultat est le **diabète sucré**. Dans le **diabète de type 1**, les cellules β sont détruites (réaction auto-immune) et aucune insuline n'est produite ; la survie impose des injections quotidiennes d'insuline. Pendant des décennies, l'insuline était extraite de pancréas de porcs ou de bovins provenant d'abattoirs. L'insuline animale fonctionne, mais elle diffère légèrement de l'insuline humaine (l'insuline bovine diffère de trois acides aminés), provoquant des réactions allergiques et une production d'anticorps chez certains patients, et l'approvisionnement était limité.

La technologie de l'ADN recombinant (détaillée en Section 4) a résolu ce problème. La logique est simple : donner à une bactérie le gène humain de l'insuline, et la bactérie devient une usine microscopique.

METHOD Production d'insuline humaine recombinante

1. Isoler le gène. Le gène humain de l'insuline est obtenu (ou synthétisé à partir de l'ARNm des cellules β grâce à la transcriptase inverse, donnant un ADNc sans introns).
2. Insérer dans un vecteur. Le gène est coupé et ligaturé dans un plasmide à l'aide d'enzymes de restriction et d'ADN ligase, placé sous le contrôle d'un promoteur bactérien fort.
3. Transformer l'hôte. Le plasmide recombinant est introduit dans *Escherichia coli* (ou la levure *Saccharomyces cerevisiae*).
4. Sélectionner et cultiver. Les cellules transformées sont sélectionnées (marqueur de résistance aux antibiotiques) et cultivées en grands fermenteurs industriels sous température, pH et aération contrôlés.
5. Extraire et purifier. Les chaînes d'insuline sont récoltées, les deux chaînes (A et B) sont repliées correctement et reliées par des ponts disulfure, puis purifiées et formulées pour injection.

DID YOU KNOW? Insuline et Cameroun

Le diabète est un problème de santé publique croissant au Cameroun, favorisé par

l'urbanisation et le changement des régimes alimentaires. L'insuline humaine recombinante, produite industriellement depuis le début des années 1980, a été la première protéine thérapeutique jamais obtenue par génie génétique. Comme elle est identique à l'hormone humaine, elle provoque beaucoup moins de réactions immunitaires que l'insuline animale, et sa production peut être montée à l'échelle pour répondre à la demande mondiale — un avantage décisif là où l'accès au traitement est déjà difficile.

2.3 Biomatériaux pour le soutien tissulaire

L'homéostasie n'est pas la seule fonction que la biotechnologie peut restaurer. Os, cartilage, ligaments et peau assurent le **soutien** ; lorsqu'ils sont détruits par un traumatisme, une infection ou un cancer, les ingénieurs les remplacent aujourd'hui par des **biomatériaux**.

DEFINITION Biomatériau

Un **biomatériau** est un matériau (naturel ou synthétique) conçu pour interagir avec un système biologique à des fins thérapeutiques : remplacer, soutenir ou régénérer un tissu ou un organe, tout en étant toléré par l'organisme (biocompatibilité).

Trois familles à connaître.

- **Hydrogels.** Réseaux de polymères hydrophiles gonflés d'eau (jusqu'à 90 % d'eau). Leur souplesse et leur teneur en eau rappellent le tissu vivant, ils servent donc pour les lentilles de contact, les pansements qui maintiennent les brûlures humides, et comme échafaudages libérant des cellules ou des facteurs de croissance vers du cartilage lésé.
- **Échafaudages de collagène.** Le collagène est la protéine structurelle la plus abondante de l'os, de la peau et des tendons. Du collagène purifié (souvent d'origine bovine, traité pour éliminer les parties antigéniques) est façonné en éponges ou membranes qui servent de squelette temporaire : les propres cellules du patient le colonisent, sécrètent leur matrice, et l'échafaudage est progressivement remplacé par du tissu naturel. C'est le principe de l'**ingénierie tissulaire**.
- **Polymères biodégradables : PLA et PGA.** L'acide polylactique (PLA) et l'acide polyglycolique (PGA) sont des polyestères synthétiques qui s'hydrolysent dans l'organisme en acide lactique et acide glycolique — métabolites naturels éliminés par la respiration et les urines. Ils servent pour les fils de suture résorbables, les broches et vis de fixation osseuse, et les échafaudages poreux pour la régénération osseuse : l'implant soutient le tissu pendant la cicatrisation, puis disparaît, évitant une seconde chirurgie pour l'enlever.

WARNING Rejet et infection à l'interface


Aucun biomatériau n'est parfaitement inerte. Les deux grands dangers à l'interface biomatériau-tissu sont : (i) le rejet immunitaire ou l'inflammation chronique si le matériau

est reconnu comme étranger (l'organisme l'isole par du tissu fibreux, altérant la fonction) ; (ii) l'infection, car les bactéries peuvent former un biofilm à la surface de l'implant que les antibiotiques pénètrent mal. D'où la nécessité absolue de tests de biocompatibilité, de conditions chirurgicales stériles, et parfois d'immunosuppression. N'écrivez jamais en examen qu'un matériau artificiel est "accepté automatiquement" par le corps.

2.4 Prothèses et exosquelettes : restaurer la locomotion

Le soutien donne au corps son cadre ; la **locomotion** résulte de muscles tirant sur ce cadre sous contrôle nerveux. Lorsqu'un membre est perdu — après des accidents de la route, hélas fréquents sur les routes camerounaises, ou après amputation pour gangrène diabétique — la biotechnologie propose des **prothèses myoélectriques**.

Principe. Lorsqu'un muscle se contracte, même un muscle résiduel de moignon, il génère un minuscule signal électrique (quelques millivolts) : le **signal électromyographique (EMG)**. Des électrodes de surface placées sur la peau du moignon détectent ce signal ; un microprocesseur (le contrôleur) l'interprète et commande des moteurs électriques (actionneurs) qui bougent l'articulation artificielle. Le patient commande la prothèse simplement en contractant les muscles qui actionnaient autrefois le membre manquant. Les systèmes plus avancés ajoutent une **interface neurale-machine** : électrodes connectées aux nerfs périphériques, voire implantées dans le cortex moteur, qui lisent l'intention de mouvement directement, et un retour sensoriel (capteurs de pression stimulant la peau) qui restaure une forme de toucher. Les **exosquelettes** appliquent la même logique à l'extérieur : un cadre motorisé attaché aux jambes d'un patient paraplégique détecte les mouvements du tronc et commande les articulations de hanche et de genou pour reproduire la marche.



Vue éclatée d'une prothèse de genou myoélectrique. Les signaux EMG des muscles du moignon sont décodés par le contrôleur, qui commande l'actionneur du genou ; les capteurs de force au pied fournissent un retour pour une marche stable.

2.5 Évaluation fonctionnelle : le dispositif marche-t-il vraiment ?

Une prothèse ou un implant n'a de valeur que s'il améliore la vie du patient mesurablement. Biotechnologues et kinésithérapeutes utilisent donc des tests fonctionnels standardisés.

METHOD Protocole d'évaluation d'une prothèse de membre inférieur

1. Test de marche de six minutes (6MWT). Le patient marche le plus loin possible en 6 minutes sur un couloir plat ; la distance totale (en mètres) mesure l'endurance. Le test est répété avant l'appareillage et après rééducation pour quantifier le progrès.
2. Analyse de la marche. Caméras et marqueurs réfléchissants enregistrent les angles articulaires ; une plateforme de force intégrée au sol mesure les forces de réaction du sol, révélant les asymétries entre le membre sain et la prothèse.
3. Capteurs de force. Semelles instrumentées mesurent la distribution des pressions plantaires, détectant les zones de surcharge pouvant causer des ulcères cutanés.
4. Score de satisfaction. Un questionnaire standardisé (confort, douleur, facilité d'utilisation, image corporelle) est noté, car un dispositif techniquement parfait que le patient refuse de porter est un échec.
5. Comparer et décider. Les résultats sont comparés aux valeurs de référence et à la ligne de base du patient ; les réglages du contrôleur sont alors ajustés.

WORKED EXAMPLE 2 Interprétation d'un test de marche de six minutes

Un patient appareillé d'une prothèse de genou myoélectrique parcourt 210 m au 6MWT lors du premier appareillage. Après 8 semaines de rééducation à la marche, il parcourt 330 m. (a) Calculer le pourcentage d'amélioration. (b) Sa vitesse moyenne lors du second test, en m s^{-1} . (c) Pourquoi un questionnaire de satisfaction reste-t-il nécessaire ?

SOLUTION

(a) Amélioration = $\frac{330 - 210}{210} \times 100 = \frac{120}{210} \times 100 \approx 57\%$.

(b) Vitesse = $\frac{\text{distance}}{\text{temps}} = \frac{330 \text{ m}}{360 \text{ s}} \approx 0,92 \text{ m s}^{-1}$.

(c) La distance seule ne dit rien sur la douleur, le confort de l'emboîture ou la confiance. Un patient peut marcher plus loin tout en souffrant d'irritations cutanées ou en trouvant l'appareil trop lourd ; seule l'évaluation combinée objective (distance, symétrie de la marche) et subjective (score de satisfaction) valide la prothèse.

2.6 Considérations éthiques et réglementaires

Ces technologies touchent directement le corps humain, elles soulèvent donc des questions

éthiques spécifiques que l'examineur du GCE attend que vous discutiez de manière équilibrée.

- **Consentement éclairé.** Le patient doit comprendre les bénéfices, les risques (infection, rejet, défaillance du dispositif) et les alternatives avant de recevoir un implant ou de rejoindre un essai clinique. Le consentement doit être libre, éclairé et révoquant.
- **Accessibilité et équité.** Les prothèses myoélectriques et l'insuline recombinante sont coûteuses. Est-il acceptable qu'un dispositif changeant la vie soit accessible aux patients aisés mais pas à un paysan de la région du Nord-Ouest ? La biotechnologie doit s'accompagner de politiques (subventions, production locale, insulines génériques) qui réduisent les inégalités.
- **Brevets sur les dispositifs médicaux.** Les brevets récompensent l'innovation et financent la recherche, mais ils peuvent maintenir les prix élevés pendant des années. Le débat entre protection des inventeurs et protection des patients est central en biotechnologie de santé.
- **Sécurité et régulation.** Tout biomatériau ou dispositif doit passer des tests de biocompatibilité, des essais cliniques et l'approbation des autorités de régulation avant mise sur le marché ; une surveillance à long terme se poursuit après commercialisation.
- **Augmentation humaine.** Si un exosquelette peut un jour permettre à une personne saine de courir plus vite, où se situe la frontière entre thérapie (restaurer une fonction perdue) et augmentation (améliorer une fonction normale) ? La société, pas les seuls scientifiques, doit répondre.

KEY POINTS Points clés de cette section

- L'homéostasie (glycémie, température, pH sanguin) est maintenue par des **boucles de rétroaction négative** : la réponse annule le stimulus.
- L'insuline humaine recombinante, produite par des bactéries ou levures génétiquement modifiées, a restauré une boucle homéostatique défaillante et transformé le traitement du diabète de type 1.
- Les **biomatériaux** (hydrogels, échafaudages de collagène, polymères biodégradables PLA/PGA) remplacent ou régénèrent les tissus de soutien ; les risques majeurs sont le rejet immunitaire et l'infection.
- Les prothèses myoélectriques et les exosquelettes convertissent les **signaux EMG** en mouvement via électrodes, contrôleur et actionneurs ; les interfaces neurales vont plus loin en lisant directement les signaux nerveux.
- La fonction est évaluée objectivement (test de marche 6 min, analyse de la marche, capteurs de force) et subjectivement (scores de satisfaction).
- Consentement, accès équitable, brevets et régulation de sécurité encadrent l'usage éthique de ces technologies.

EXERCISE 2 Vérifiez votre compréhension

1. Dessinez et annotez la boucle de rétroaction négative qui restaure la glycémie après un repas riche en glucides.
2. Expliquez pourquoi l'insuline humaine recombinante provoque moins de réactions immunitaires que l'insuline bovine.
3. Donnez un avantage et un risque de l'utilisation d'une vis en PLA biodégradable au lieu d'une vis métallique pour fixer une fracture osseuse.
4. Tracez le chemin de l'information dans une prothèse myoélectrique, de l'intention du patient au mouvement du genou.
5. Un patient parcourt 250 m au 6MWT avant rééducation et 325 m après. Calculez le pourcentage d'amélioration et commentez pourquoi l'analyse de la marche doit compléter ce résultat.

SOLUTION Exercice 2

1. Repas → glycémie monte → cellules β îlots Langerhans stimulées → insuline sécrétée → capture glucose muscle/adipose et glycogénèse foie → glycémie baisse → sécrétion insuline stoppée (rétroaction).
2. L'insuline recombinante a exactement la séquence d'acides aminés de l'insuline humaine, alors que l'insuline bovine diffère de trois acides aminés ; ces séquences étrangères peuvent être reconnues comme antigènes, déclenchant production d'anticorps et allergie.
3. Avantage : elle s'hydrolyse en acide lactique, métabolite naturel, donc pas de seconde opération pour l'enlever. Risque : inflammation ou infection à l'interface tissu-matériau, ou rupture mécanique avant consolidation complète.
4. Intention → contraction muscles moignon → signal EMG détecté électrodes surface → microprocesseur decode signal → commande envoyée actionneur genou (moteur) → mouvement articulation ; capteurs force fournissent retour pour ajustement.
5. Amélioration = $\frac{325 - 250}{250} \times 100 = 30 \%$. Le 6MWT mesure l'endurance seulement ; l'analyse de la marche révèle la qualité de la marche (symétrie, angles articulaires, boiterie), que la distance seule ne montre pas.

3 CORRECTION ET REMÉDIATION GÉNIQUE : THÉRAPIE GÉNIQUE ET ÉDITION DU GÉNOME

In the previous section, we explored how biotechnology provides external support for homeostasis, mobility and structural repair — from biosensors that monitor blood glucose to engineered prostheses and tissue scaffolds. We now turn to a more radical approach: instead of managing the symptoms of a genetic defect, we correct the defect at its source, the DNA itself. This is the promise of **gene therapy** and **genome editing**. Imagine a patient with sickle cell disease, a condition highly prevalent in Cameroon, whose red blood cells sickle because of a single nucleotide mutation in the HBB gene. What if we could precisely edit that mutation back to the normal sequence, or introduce a functional copy of the gene? This section unpacks the molecular tools, the clinical realities, and the ethical boundaries of rewriting the human genome.

3.1 Monogenic Diseases and Therapeutic Strategies

Many debilitating inherited disorders are **monogenic diseases** — caused by mutations in a single gene. Classic examples include sickle cell disease (mutation in HBB, β -globin), cystic fibrosis (mutation in CFTR, chloride channel), Duchenne muscular dystrophy (DMD, dystrophin), and Leber congenital amaurosis (RPE65, retinal pigment epithelium). The therapeutic logic is straightforward: if a disease results from a loss-of-function mutation, restoring functional protein should alleviate the phenotype. Two broad strategies exist:

DEFINITION Gene Therapy

Gene therapy is the introduction of nucleic acids (DNA or RNA) into the cells of a patient to treat or prevent a disease. It aims to modify the genetic information of target cells to achieve a therapeutic effect.


- **Gene addition (augmentation):** A functional copy of the defective gene (**transgene**) is delivered into the nucleus, often remaining episomal (non-integrating) or integrating randomly. The endogenous mutant allele is untouched. This suits recessive loss-of-function disorders.
- **Gene editing (correction):** The endogenous mutated sequence is directly modified — corrected, disrupted, or rewritten — at its native chromosomal locus. This preserves physiological regulation and avoids insertional mutagenesis. The breakthrough tool is **CRISPR-Cas9**.

3.2 Gene Addition Therapy: Viral and Non-Viral Vectors

Delivering a transgene (often 2–5 kb of cDNA plus promoter and polyA signal) into the nucleus of specific cells in a living patient is a formidable delivery problem. The vehicle is called a **vector**.

DEFINITION Viral Vector

A viral vector is a virus that has been genetically modified to remove its pathogenic genes and replication capacity, while retaining its ability to enter target cells and deliver a therapeutic transgene cassette.



Life cycle of an AAV vector in a target cell. AAV remains predominantly episomal, making it ideal for non-dividing cells (neurons, muscle, retina) but diluted in dividing tissues.


Vector	Genome / Capacity	Integration	Key Features / Use Cases
Adenovirus (up to 8–10 kb insert)	dsDNA, ~36 kb Episomal (non-integrating)	High immunogenicity (strong innate/adaptive response). Transient expression. Used in vaccines (e.g. COVID-19 AstraZeneca, J&J) and some cancer therapies.	
AAV (Adeno-Associated Virus) (max ~4.5 kb insert)	ssDNA, ~4.7 kb Mostly episomal (concaten-ers); low frequency integration at AAVS1	Low immunogenicity, many serotypes for tissue tropism (AAV2–retina, AAV8/9–liver, AAV9–CNS). Gold standard for in vivo gene therapy (Luxturna, Zolgensma).	
Lentivirus (HIV-1 based) (up to 8 kb insert)	ssRNA, ~9 kb Integrating (random in active genes)	Transduces dividing & non-dividing cells. Stable expression in progeny. Used ex vivo for HSPCs (Strimvelis, Zynteglo, Casgevy).	

METHOD Choosing a Vector for Gene Therapy

Step 1: Transgene size. > 4.5 kb → exclude AAV; consider Lentivirus, Adenovirus, or non-viral. Step 2: Target tissue & division status. Non-dividing (neurons, muscle, retina) → AAV ideal. Dividing (HSPCs, T cells) → Lentivirus (integrating) or ex vivo editing. Step 3: Duration of expression needed. Lifelong → Integrating (Lentivirus) or stable episome in non-dividing cells (AAV). Transient → Adenovirus, mRNA-LNP. Step 4: Immunogenicity constraints. Pre-existing neutralizing antibodies (NABs) against AAV serotypes are common (30–60% of humans). Screen patients; consider alternate serotypes or immunosuppression. Step 5: Safety profile. Insertional mutagenesis risk (Lentivirus) vs. immune toxicity (Adenovirus) vs. off-target editing (CRISPR). Regulatory precedent matters.

3.3 Genome Editing: The CRISPR–Cas9 Revolution

While gene addition adds a functional copy, **genome editing** rewrites the existing sequence. The discovery of the bacterial adaptive immune system CRISPR–Cas (Clustered Regularly Interspaced Short Palindromic Repeats, CRISPR-associated proteins) and its repurposing by Doudna, Charpentier, and Zhang (2012–2013) transformed biology.



CRISPR–Cas9 mechanism. The gRNA spacer (20 nt) base-pairs with the target DNA strand adjacent to a PAM (NGG for *S. pyogenes* Cas9). Cas9 undergoes conformational change; HNH domain cuts the target (complementary) strand, RuvC domain cuts the non-target strand, generating a blunt double-strand break (DSB) 3 bp upstream of the PAM.

PROPERTY DNA Repair Pathways After CRISPR–Cas9 Cleavage

A double-strand break (DSB) is repaired by two main cellular pathways:

1. NHEJ (Non-Homologous End Joining): Error-prone, ligates ends directly. Frequent in all cell types, especially G1. Creates small insertions/deletions (indels) → gene

knockout (frameshift). High frequency, imprecise.

2. HDR (Homology-Directed Repair): Uses a homologous template (sister chromatid or exogenous donor DNA). Precise correction/insertion possible. Restricted to S/G2 phases; low frequency in vivo; requires donor template.

Examinable: You must be able to explain why HDR is inefficient in non-dividing cells (neurons, muscle) and why NHEJ dominates in vivo.

WARNING Off-Target Effects of CRISPR

CRISPR-Cas9 can cleave DNA at sites with sequence similarity to the target (mismatches tolerated, especially in the PAM-distal "seed" region). These **off-target mutations** are random, potentially disrupting tumor suppressor genes or activating oncogenes. Mitigation: High-fidelity Cas9 variants (eSpCas9, HiFi Cas9), truncated gRNAs (17–18 nt), RNP delivery (short exposure), and rigorous off-target screening (GUIDE-seq, CIRCLE-seq, DISCOVER-seq) in preclinical studies.

3.4 Clinical Case Studies

EXAMPLE Leber Congenital Amaurosis Type 2 (LCA2) — RPE65 Gene Addition

Disease: Autosomal recessive blindness from infancy. RPE65 encodes an isomerase essential for the visual cycle. Mutations → no functional visual pigment regeneration. Therapy: **Voretigene neparvovec (Luxturna[®])**, AAV2 vector carrying human RPE65 cDNA. Subretinal injection (detaches retina locally, delivers vector to RPE cells). Outcome: Improved light sensitivity, navigational ability in low light. First FDA-approved in vivo gene therapy (2017). Demonstrates: AAV + non-dividing target + subretinal immune privilege = success.

EXAMPLE Sickle Cell Disease (SCD) / β -Thalassemia — **CTX001 (Casgevy[®])** — Genome Editing

Disease: β -globin mutation (HbS: Glu6Val). Polymerization of deoxy-HbS → sickling, vaso-occlusion, anemia. Strategy: Not correcting HBB directly (hard in HSPCs). Instead: Reactivate fetal hemoglobin (HbF). HbF ($\alpha_2\gamma_2$) does not sickle. The BCL11A erythroid enhancer represses γ -globin. Disrupt this enhancer → HbF ↑. Procedure (Ex vivo):

1. Mobilize CD34+ hematopoietic stem/progenitor cells (HSPCs) from patient (G-CSF + plerixafor).

2. Electroporate Cas9–gRNA RNP targeting BCL11A erythroid enhancer.
3. Myeloablative conditioning (busulfan) to make niche space.
4. Autologous infusion of edited HSPCs.

Results (CLIMB-111/121 trials): > 90% patients free of vaso-occlusive crises (VOCs) or transfusion independence for β -thalassemia. HbF ~20–40%. Approved UK/US/EU 2023–2024. First CRISPR therapy approved. Significance: Proof that ex vivo editing of HSPCs with NHEJ-mediated disruption is curative for hemoglobinopathies.

WORKED EXAMPLE 3 Designing a CRISPR Strategy for a Hypothetical Mutation

Statement: A patient has a recessive disease caused by a nonsense mutation (CAG → TAG, Gln → Stop) in exon 4 of gene X. The genomic target sequence (non-target strand) is: 5'-GGACTG**TAG**GGTACC-3' (PAM underlined). Design a gRNA spacer (20 nt) to correct this via HDR using a single-stranded oligodeoxynucleotide (ssODN) donor. Explain the PAM requirement and the donor design.

Solution: 1. Identify PAM and Protospacer. *S. pyogenes* Cas9 requires 5'-NGG-3' on the non-target strand. The sequence shows TGG at positions 10–12 (if we number from 1). Wait, the mutation is at position 7–9 (TAG). We need a PAM downstream (3') of the target site on the non-target strand. Let's scan 3' of the mutation: ...TAGGGT... No GG. ...TAGGTA... No. ...TAGGTACC... No. Scan 5' (upstream): GGACTG... No. GACTG... No. ACTG... No. CTG... No. TGA... No. Problem: There may be no suitable NGG PAM near the mutation for SpCas9. Alternative: Use a different Cas ortholog (e.g. *S. aureus* Cas9: NNGRRT; Cpf1/Cas12a: TTV on 5' end). Assume we find a PAM: Suppose sequence is 5'-...CTGTAGGGTAGG...-3' (PAM = AGG at end). The protospacer is the 20 nt immediately 5' of PAM: 5'-CTGTAGGGTAGGCTGAA???-3' (need 20 nt total). 2. gRNA Spacer Sequence: The gRNA spacer is identical to the non-target strand (except U for T). If protospacer = 5'-ATGCTGTAGGGTAGGCTGAA-3' (example 20 nt), gRNA = 5'-AUGUUGUAGGGUAGGCUGAA-3'? Wait: DNA: A T G C T G T A G G G T A G G C T G A A. RNA: A U G C U G U A G G G U A G G C U G A A. 3. Donor ssODN Design:

- Length: ~100–200 nt total, symmetric homology arms (~50–90 nt each side).
- Correction: Change TAG (Stop) back to CAG (Gln).
- Silent PAM/Seed mutation: Introduce a synonymous mutation in the PAM or seed region (e.g. AGG → AGA, still Arg) to prevent re-cutting of the corrected allele by Cas9–gRNA.
- Strand: Use the strand complementary to the non-target strand (i.e. same as target strand) for better HDR efficiency with Cas9.

Key Point: PAM availability dictates targetability. Base editors (CBEs/ABEs) or Prime Editing are often preferred for point mutations as they don't require DSBs or donor tem-

plates.

3.5 Evaluation of Efficacy and Safety

Moving from bench to bedside requires rigorous biomarkers.

- **Vector Copy Number (VCN) / Transduction Efficiency:** qPCR or ddPCR on genomic DNA from target tissue (or blood for HSPCs) measuring transgene vs. single-copy reference gene. Target VCN: 1–3 copies/cell for lentiviral; vector genomes (vg) per cell for AAV.
- **Transgene Expression:** mRNA (RT-qPCR, RNA-seq) and protein (ELISA, Western blot, flow cytometry, functional assay e.g. Factor IX activity for hemophilia B).
- **Off-Target Analysis (Editing):** Targeted deep sequencing of predicted off-target sites (in silico prediction + GUIDE-seq). Whole-genome sequencing (WGS) for unbiased detection. Threshold: typically < 0.1% indel frequency at off-targets.
- **Insertional Site Analysis (Integrating vectors):** LAM-PCR or nrLAM-seq to map integration sites. Check for clonal dominance (enrichment near oncogenes like LMO2, CCND2).
- **Immunogenicity:** Neutralizing antibodies (NAbs) against capsid (AAV) or transgene product (if non-self). T-cell responses to capsid peptides (monitor liver enzymes for AAV-hepatotropism).

3.6 Ethical and Legal Framework

DID YOU KNOW?

Bioethics in Cameroon and Internationally

International: The Declaration of Helsinki (WMA, 1964, revised 2013) is the cornerstone for human research: informed consent, risk/benefit assessment, independent ethics committee (IEC/IRB) review. The UNESCO Universal Declaration on the Human Genome and Human Rights (1997) states the genome is "heritage of humanity" and opposes germline modifications "contrary to human dignity". Europe/West: Strict distinction: Somatic gene therapy (treats patient, not inherited) = allowed with oversight. Germline editing (embryos, gametes, heritable) = banned (EU Convention on Human Rights and Biomedicine, Oviedo 1997). Cameroon: Law No. 2002/003 of 19 April 2002 relating to the protection of persons undergoing biomedical research. Requires authorization from the National Ethics Committee for Human Health Research (CNERSH). Gene therapy trials would fall under this. No specific "bioethics law" on germline editing yet, but the Constitution protects human dignity and the Helsinki principles are ratified. The Ministry of Public Health (MINSANTÉ) oversees clinical trials. He Jiankui Affair (2018): Chinese scientist created CRISPR-edited babies (CCR5 Δ 32 for HIV resistance). Universal condemnation: premature, inadequate consent, off-target risks, germline modification. Highlights need for global governance (WHO Expert Advisory Committee on Developing Global Standards for Governance and Oversight of Human Genome Editing, 2019–2021).

KEY POINTS Key Points for Revision

- Gene Therapy = Nucleic acid delivery for therapeutic effect. Two paradigms: Addition (viral/non-viral vectors) vs Editing (CRISPR-Cas9, Base/Prime editors).
- Vectors: AAV (episomal, non-dividing cells, low immunogenicity, ~4.5 kb limit). Lentivirus (integrating, dividing/non-dividing, ~8 kb, ex vivo HSPCs). Non-viral/LNP (scalable, transient, improving).
- CRISPR-Cas9: gRNA (spacer + scaffold) guides Cas9 to target DNA adjacent to PAM (NGG). DSB → NHEJ (indels, knockout) or HDR (precise edit, needs donor, low efficiency).
- Clinical successes: Luxturna (AAV-RPE65, retinal), Zolgensma (AAV9-SMN1, SMA), Casgevy (CRISPR-BCL11A enhancer, SCD/Thal).
- Safety: Off-target edits (oncogenic risk), insertional mutagenesis (lentivirus), immune responses (AAV capsid, transgene). Monitoring: VCN, expression, off-target seq, integration site analysis.
- Ethics: Somatic = therapeutic, regulated. Germline = heritable, internationally prohibited. Cameroon: Law 2002/003, CNERSH oversight, Helsinki Declaration.

EXERCISE 3 Gene Therapy Design Challenge

A 6-year-old Cameroonian boy is diagnosed with X-linked Severe Combined Immunodeficiency (SCID-X1) due to a null mutation in IL2RG (common γ chain). He has no matched sibling donor for HSCT. You are the molecular biologist designing a gene therapy protocol.

1. Choose the optimal vector (AAV, Lentivirus, Adenovirus, LNP) and justify your choice based on: target cell type (HSPCs), need for lifelong correction, transgene size (IL2RG cDNA ~1.2 kb), and division status of target cells.
2. Outline the ex vivo manufacturing steps from leukapheresis to cryopreserved drug product.
3. Identify the major historical safety concern for integrating vectors in SCID-X1 trials (early 2000s) and how modern vectors mitigate it.
4. Propose a monitoring plan for the first 2 years post-infusion (efficacy + safety biomarkers).

SOLUTION Exercise: Gene Therapy Design Challenge

1. Vector: Lentivirus (SIN, self-inactivating). Justification: Target = CD34+ HSPCs (dividing, long-lived). Need stable integration for lifelong multi-lineage reconstitution. Transgene small (fits easily). AAV is episomal → lost on HSPC division. Adenovirus immunogenic, transient. LNP DNA delivery to HSPCs inefficient.
2. Steps: (1) Mobilization (G-CSF/plerixafor) → Leukapheresis. (2) CD34+ selection (CliniMACS). (3) Activation/Prestimulation (cytokines: SCF, TPO, Flt3L, IL-6). (4) Transduction (LV vector, MOI 5–20, retronectin/fibronectin). (5) Wash, formulation (cryoprotectant DMSO/albumin). (6) QC: Sterility, VCN (ddPCR), viability, potency (colony-forming units), replication-competent lentivirus (RCL) test. (7) Cryopreservation. (8) Patient conditioning (busulfan). (9) Thaw → Infusion.
3. Historical concern: Insertional oncogenesis. First SCID-X1 trials (gamma-retrovirus, LTR-driven) → T-cell leukemia in ~25% patients due to LTR enhancer activating LMO2 oncogene. Mitigation: SIN-LTRs (deleted U3 enhancer), internal promoters (e.g. EFS, PGK) weaker/physiological, chromatin insulators (chs4), integration-deficient lentiviral vectors (IDLV) for non-dividing targets (not here).
4. Monitoring: Months 1–3: Engraftment (chimerism), lymphocyte subsets (CD3, CD4, CD8, CD19, NK), Ig levels, VCN in PBMCs (qPCR/ddPCR), integration site analysis (clonal tracking), RCL testing. Months 6–24: Immune reconstitution (vaccine responses, thymic output TRECs), clinical infections, VCN stability, clonal diversity (no dominant clone > 30%), oncogene proximity screening. Long-term: Annual follow-up 15 years (FDA/EMA).

4 TECHNOLOGIE DE L'ADN RECOMBINANT ET ALTÉRATION GÉNIQUE

In Section 3 we saw how a defective gene can be corrected or replaced inside a patient — the logic of gene therapy and genome editing. But before any gene can be corrected, delivered or edited, molecular biologists must first be able to cut DNA at precise positions, paste fragments together, copy a gene millions of times, and check that the final construct is exactly what was intended. These four abilities — cut, paste, amplify, verify — are the heart of **recombinant DNA technology**. This section presents the enzymatic toolbox (restriction enzymes, ligases, polymerases), the vehicles that carry foreign DNA into cells (cloning vectors), the classical cloning workflow, the polymerase chain reaction (PCR), modern editing tools such as CRISPR–Cas9, and the quality-control methods that validate every construct.

4.1 Restriction enzymes: the molecular scissors

Observation. Bacteria are constantly attacked by viruses (bacteriophages). Many bacteria

defend themselves by producing enzymes that chop the invading viral DNA into pieces while leaving their own DNA untouched (their own recognition sites are protected by methylation). These enzymes, purified and used in the laboratory, became the first great tool of genetic engineering.

DEFINITION Recombinant DNA


Recombinant DNA is a DNA molecule formed by joining together fragments of DNA from different sources (for example a human insulin gene inserted into a bacterial plasmid). Its construction requires enzymes able to cut DNA at defined sequences and to reseal the sugar–phosphate backbone.

DEFINITION Restriction endonuclease

A **restriction enzyme** (restriction endonuclease) is an enzyme that recognises a specific short DNA sequence, the **recognition site**, and cuts both strands of the double helix. Type II enzymes, the most useful in the laboratory, cut within or immediately next to their recognition site.

Nomenclature. The name encodes the organism of origin: the first letter of the genus, the first two letters of the species, the strain, and a Roman numeral for the order of discovery. Thus **EcoRI** comes from *Escherichia coli*, strain R, enzyme I; **HindIII** from *Haemophilus influenzae* strain d; **BamHI** from *Bacillus amyloliquefaciens* strain H.

Palindromic recognition sites. Most recognition sites are **palindromes**: the sequence reads identically in the 5' → 3' direction on both strands. For example **EcoRI** recognises:



Reading the top strand 5' → 3' gives GAATTC; reading the bottom strand 5' → 3' also gives GAATTC. This symmetry matches the dimeric structure of the enzyme.

Sticky versus blunt ends. Depending on where the two strands are cut, two kinds of ends are produced:

- **Cohesive (sticky) ends:** the cuts are staggered, leaving short single-stranded overhangs (**EcoRI** leaves a 5'–AATT overhang). Two fragments cut with the same enzyme have complementary overhangs and can pair by hydrogen bonding — this makes ligation efficient and directional cloning possible when two different enzymes are used.
- **Blunt ends:** both strands are cut at the same position (e.g. **SmaI**, CCC↓GGG). Any blunt end can join any other blunt end, but ligation is far less efficient and orientation cannot be controlled.

PROPERTY Type II restriction enzymes

Type II restriction endonucleases cut DNA within or very close to their recognition sequence, generating predictable, reproducible ends (cohesive or blunt). Consequently, digestion of a given DNA molecule by a given enzyme always yields the same set of fragments — the basis of restriction mapping and cloning. (Statement examinable; no formal proof required.)

WARNING Star activity and methylation

Under non-optimal conditions (high glycerol, wrong pH or ionic strength), some enzymes show “star activity” and cut at sites resembling the true site, giving extra bands on gels. Also, DNA extracted from *dam*⁺ *E. coli* strains is methylated and may resist enzymes such as *Cl*I or *Xba*I. Always check the enzyme data sheet before designing a digest.

4.2 Cloning vectors: vehicles for foreign DNA

A restriction fragment alone cannot replicate inside a cell. It must be joined to a **cloning vector**: a small, autonomously replicating DNA molecule that carries the insert into a host cell and amplifies it.


DEFINITION Cloning vector

A **cloning vector** is a DNA molecule (plasmid, phage-derived or artificial chromosome) capable of independent replication in a host cell, used to carry and amplify a foreign DNA insert. Every useful vector possesses three essential elements:

1. an **origin of replication** (*ori*), recognised by the host replication machinery;
2. a **selectable marker** (e.g. an antibiotic-resistance gene such as *amp*^R) allowing survival of only those cells that received the vector;
3. a **multiple cloning site** (MCS or polylinker): a short region containing many unique restriction sites where the insert is placed.

Vector	Insert capacity	Typical use
Plasmid (pUC19, pBR322)	up to ~ 10 kb	Routine cloning, sub-cloning, expression
Cosmid (plasmid + phage λ cos sites)	35–45 kb	Genomic libraries of moderate-size genomes
BAC (bacterial artificial chromosome)	100–300 kb	Large genomic fragments, genome projects
YAC (yeast artificial chromosome)	250 kb – 1 Mb	Very large fragments; contains yeast ori, centromere and telomeres

The plasmid **pUC19** is the classic teaching example: it carries the ori, the amp^R gene (ampicillin resistance), and the MCS inserted inside the $\text{lacZ}\alpha$ gene — an arrangement exploited by blue/white screening (Section 4.3).



Simplified map of the cloning plasmid pUC19: origin of replication (ori), ampicillin-resistance marker (amp^R), and the multiple cloning site (MCS) embedded in the $\text{lacZ}\alpha$ reporter gene.

4.3 The classical cloning workflow

METHOD Classical gene cloning: the five steps

1. Digestion. Cut both the vector and the source DNA with the same restriction enzyme(s) so that compatible ends are generated. Dephosphorylate the vector ends (alkaline phosphatase) to prevent self-ligation.

2. Ligation. Mix insert and vector with T4 DNA ligase, which reseals the sugar–phosphate backbone (see the ligation protocol box below).
3. Transformation. Introduce the recombinant plasmid into competent *E. coli* cells, either by **heat shock** (cells made competent with CaCl_2 , chilled on ice, then pulsed at 42°C for ~ 45 s) or by **electroporation** (a brief electric pulse opens transient pores in the membrane; higher efficiency).
4. Selection. Spread the cells on agar containing ampicillin: only bacteria carrying a plasmid (with amp^R) form colonies.
5. Screening. Distinguish recombinant from non-recombinant plasmids:
 - **Blue/white screening**: the MCS lies inside $\text{lacZ}\alpha$. An intact $\text{lacZ}\alpha$ produces β -galactosidase, which cleaves X-gal into a blue product. An insert disrupts $\text{lacZ}\alpha$, so recombinant colonies stay white while empty-vector colonies turn blue.
 - **Colony PCR**: a tiny amount of each colony is used directly as PCR template with primers flanking the MCS; the size of the product reveals the presence and size of the insert.

METHOD Ligation protocol

1. Prepare the reaction with an **insert : vector molar ratio of 3:1** (excess insert favours productive joining over vector self-circularisation). Compute masses with

$$m_{\text{insert}} = 3 \times m_{\text{vector}} \times \frac{\text{size of insert (bp)}}{\text{size of vector (bp)}}$$

2. Add T4 DNA ligase and its ATP-containing buffer.
3. Incubate at 16°C overnight (a compromise: ends anneal best at low temperature, but the ligase works faster at higher temperature).
4. Transform 1–2 μL of the ligation mixture into competent cells.

WORKED EXAMPLE 4 Calculating insert mass for a ligation

We wish to clone a 750 bp insert into plasmid pUC19 (2686 bp). We use 50 ng of vector. What mass of insert gives a 3:1 molar ratio?

SOLUTION

Solution. Equal numbers of molecules require masses proportional to length. For a 3:1

insert:vector molar ratio:

$$m_{\text{insert}} = 3 \times 50 \text{ ng} \times \frac{750}{2686} \approx 150 \times 0.279 \approx 42 \text{ ng}.$$

We therefore mix about 42 ng of insert with 50 ng of cut, dephosphorylated vector before adding T4 DNA ligase.

WARNING Antibiotic-resistance markers and biosafety

Resistance markers such as amp^R are convenient, but their release into the environment could spread resistance genes to pathogenic bacteria — a real public-health concern, including in Cameroon where antibiotic resistance is already a serious problem. Good practice: never discard recombinant cultures untreated (autoclave or bleach them), and prefer alternative selection systems where possible — **auxotrophic complementation** (the vector restores a metabolic gene the host lacks), **fluorescent markers** (GFP) or toxin-antitoxin systems.

4.4 DNA amplification by PCR

Cloning amplifies DNA inside cells. The **polymerase chain reaction** (PCR), conceived by Kary Mullis in the 1980s, amplifies a chosen DNA region in vitro, in a tube, in a few hours.


DEFINITION PCR

PCR is a cyclic in vitro reaction that doubles a target DNA sequence at each cycle, using: (i) a template DNA; (ii) two short **primers** (amorces) flanking the target, one per strand; (iii) the four dNTPs; (iv) a **thermostable DNA polymerase** (Taq polymerase, from the hot-spring bacterium *Thermus aquaticus*) that survives repeated heating; (v) a buffer containing Mg^{2+} .

Each cycle has three steps:

1. **Denaturation** (94–98 °C, 20–30 s): the double helix separates into single strands.
2. **Annealing** (typically 50–65 °C, 20–40 s, depending on the primers): primers hybridise to their complementary sequences.
3. **Extension (elongation)** (72 °C, ~ 1 min per kb): Taq polymerase synthesises the new strands from the primers in the 5' → 3' direction.

After n cycles the target is amplified theoretically 2^n -fold: 30 cycles give ~ 10^9 copies.



Temperature chronogram of a typical PCR programme: initial denaturation, then repeated cycles of denaturation (94 °C), primer annealing (55 °C) and extension by Taq polymerase (72 °C).

Applications. PCR is used for **genotyping** (e.g. detecting the sickle-cell allele Hb^S from a drop of blood), diagnosis of infections (malaria parasites, HIV, SARS-CoV-2), forensic identification, and cloning. In **quantitative PCR (qPCR)**, a fluorescent dye or probe reports product accumulation after each cycle: the earlier the fluorescence crosses a threshold (the C_q value), the more target was initially present — allowing precise quantification of viral load or gene expression.

WORKED EXAMPLE 5 How many copies after 25 cycles?

A PCR starts from a single DNA molecule. Assuming perfect efficiency, how many copies of the target exist after 25 cycles? If the efficiency is only 90% (each cycle multiplies by 1.9), how many copies result?

SOLUTION

Solution. At 100% efficiency, $N = 2^{25} = 33\,554\,432 \approx 3.4 \times 10^7$ copies.

At 90% efficiency, $N = 1.9^{25}$. Taking logarithms: $\log_{10} N = 25 \times \log_{10} 1.9 = 25 \times 0.2788 = 6.97$, so $N \approx 10^{6.97} \approx 9.3 \times 10^6$ copies.

Comment. A 10% drop in efficiency reduces the final yield almost four-fold over 25 cycles — this is why primer design and Mg^{2+} optimisation matter, and why qPCR efficiency must be measured before quantification.

4.5 Modern genome-editing tools

PCR and cloning manipulate DNA in a tube; **genome editing** modifies a chosen sequence inside the living genome.

CRISPR–Cas9. Derived from a bacterial immune system, the **CRISPR–Cas9** complex uses a **guide RNA (gRNA)** of about 20 nucleotides, complementary to the target, to lead the Cas9 nuclease to the matching genomic site, which must sit next to a short **PAM** motif (NGG for the *Streptococcus pyogenes* enzyme). Cas9 cuts both strands; the cell then repairs the break either by error-prone non-homologous end joining (gene knockout) or, if a donor template is provided, by homology-directed repair (precise correction — the strategy behind the sickle-cell therapies discussed in Section 3).

Base editors and prime editing. **Base editors** fuse a catalytically impaired Cas9 to a deaminase and convert one base into another (C→T or A→G) without cutting both strands. **Prime editing** uses a Cas9 nickase fused to a reverse transcriptase and an extended guide RNA (pegRNA) that both specifies the target and encodes the desired edit, allowing virtually any small substitution, insertion or deletion with minimal off-target damage.

METHOD Designing a good guide RNA

1. Choose a 20-nt target sequence adjacent to a PAM (NGG) within the region to be edited.
2. Check specificity with bioinformatics tools such as **CRISPOR** or **Benchling**: reject any gRNA whose sequence shows more than ~ 80% homology with another genomic site, especially sites differing by fewer than 3 mismatches.
3. Prefer guides with a GC content of 40–60% and avoid long poly-T stretches (which can terminate transcription).
4. Order the top-scoring candidates and validate them experimentally (e.g. by a T7E1 assay or targeted sequencing) before committing to the full experiment.

4.6 Quality control of constructs

No construct is trusted until it has been verified by independent methods:

- **Agarose gel electrophoresis:** DNA fragments migrate towards the positive electrode at a speed inversely related to their size; staining (ethidium bromide or safer dyes) reveals bands whose sizes are compared with a molecular-weight ladder. A quick check of digestions, ligations and PCR products.
- **Restriction analysis:** the purified plasmid is re-digested with diagnostic enzymes; the observed band pattern must match the pattern predicted from the map.
- **Sanger sequencing:** the gold standard. A primer anneals to the construct and a polymerase extends it in the presence of dNTPs plus small amounts of chain-terminating dideoxynucleotides, each labelled with a different fluorescent colour. The resulting fragments, separated by capillary electrophoresis, give the exact sequence, base by base, confirming that the insert is present, correctly oriented and free of mutations.

KEY POINTS Key points of Section 4

- Type II restriction enzymes cut at palindromic sites, giving predictable sticky or blunt ends; T4 DNA ligase reseals compatible ends (insert:vector = 3 : 1, 16 °C overnight).
- A cloning vector needs an ori, a selectable marker and an MCS; capacity increases from plasmids to cosmids, BACs and YACs.
- Cloning workflow: digestion → ligation → transformation (heat shock or electroporation) → antibiotic selection → screening (blue/white, colony PCR).
- PCR amplifies a target exponentially (2^n) using primers and a thermostable polymerase; qPCR quantifies the starting amount.
- CRISPR–Cas9, base editors and prime editing modify genomes at chosen sites; gRNAs must be designed with tools such as CRISPOR or Benchling to avoid off-targets (> 80% homology rejected).
- Every construct is validated by gel electrophoresis, restriction analysis and Sanger sequencing.

EXERCISE 4 Restriction mapping

A linear DNA fragment of 6000 bp is digested with EcoRI alone, giving fragments of 4000 bp and 2000 bp. Digestion with BamHI alone gives 4500 bp and 1500 bp. The double digest EcoRI + BamHI gives three fragments: 2000, 2500 and 1500 bp. Draw the restriction map showing the relative positions of the EcoRI (E) and BamHI (B) sites.

SOLUTION Restriction mapping

Let the left end be position 0 and the right end 6000 bp. EcoRI alone gives 4000 and 2000 bp ⇒ EcoRI site (E) is at 2000 bp or 4000 bp from the left. BamHI alone gives 4500 and 1500 bp ⇒ BamHI site (B) is at 1500 bp or 4500 bp from the left. Double digest gives three fragments: 2000, 2500, 1500 bp (sum = 6000). These are the intervals between 0, E, B, 6000.

Test the four combinations:

- E=2000, B=1500: intervals 1500, 500, 4000 ≠ given.
- E=2000, B=4500: intervals 2000, 2500, 1500 ✓ matches double digest.
- E=4000, B=1500: intervals 1500, 2500, 2000 ✓ same set, but order differs.
- E=4000, B=4500: intervals 4000, 500, 1500 ≠ given.

Both (E=2000, B=4500) and (E=4000, B=1500) produce the same three fragment sizes. However, the single digests distinguish them:

- If E=2000, B=4500: EcoRI fragments = 2000 and 4000 ✓; BamHI fragments = 4500 and 1500 ✓.
- If E=4000, B=1500: EcoRI fragments = 4000 and 2000 ✓; BamHI fragments = 1500 and 4500 ✓.

Both are consistent with all data; the map is symmetric. We can draw either orientation. Conventionally, we place the smaller coordinate first:

$$O \xrightarrow{2000} E \xrightarrow{2500} B \xrightarrow{1500} 6000$$

Check: EcoRI at 2000 → 2000 + 4000. BamHI at 4500 → 4500 + 1500. Double digest: O–E = 2000, E–B = 2500, B–6000 = 1500. All consistent.

5 APPLICATIONS MAJEURES DE LA BIOTECHNOLOGIE : SANTÉ, AGRICULTURE, ENVIRONNEMENT, INDUSTRIE

In the previous sections we mastered the tools of modern biotechnology: restriction enzymes, vectors, PCR, recombinant DNA, and gene editing with CRISPR–Cas9. A tool, however, is only as valuable as what it builds. In this final section we answer the question every examiner loves: what has biotechnology actually done for humanity? We shall tour the four great fields of application — **health**, **agriculture**, **environment** and **industry/energy** — and finish with the regulatory framework that governs all of this in Cameroon.

5.1 Pharmaceutical applications: medicines made by living cells


The intuition. A diabetic patient needs insulin every day. Before 1982, insulin was extracted from the pancreases of pigs and cattle slaughtered in abattoirs: it was expensive, scarce, and sometimes triggered allergic reactions because animal insulin differs slightly from human insulin in its amino–acid sequence. The biotechnological idea is disarmingly simple: why not give the human insulin gene to a microbe, and let the microbe — which doubles every 20 minutes — manufacture the hormone for us?

EXAMPLE Recombinant human insulin

The gene coding for human insulin is inserted into a plasmid and introduced into *Escherichia coli* (or the yeast *Saccharomyces cerevisiae*). The bacteria are grown in huge fermenters; they produce the insulin chains, which are extracted, folded, purified and formulated. Since 1982 (first commercial product: Humulin), virtually all insulin used worldwide is recombinant human insulin: cheaper, unlimited in supply, and immunologi–

cally identical to the patient's own hormone.

The same logic gives us **recombinant human growth hormone** (somatotropin), used to treat growth hormone deficiency in children. Previously it had to be extracted from the pituitary glands of human cadavers — a practice abandoned after it transmitted a fatal prion disease (Creutzfeldt–Jakob disease) to some treated patients. Recombinant production eliminated this risk entirely. Other recombinant pharmaceuticals include **factor VIII** for haemophiliacs (formerly prepared from pooled blood, with a risk of HIV and hepatitis transmission), **erythropoietin** (EPO) for anaemia, and **hepatitis B vaccine**, produced as a recombinant surface antigen in yeast.



Production flow of recombinant human insulin: from the human gene to the purified pharmaceutical product.

Monoclonal antibodies. A **monoclonal antibody** (mAb) is a single, identical antibody species directed against one precise epitope, produced industrially. Classical production uses hybridoma cells, obtained by fusing an antibody-producing B lymphocyte with an immortal myeloma (cancer) cell: the hybrid combines specific antibody production with unlimited division. Modern industrial production uses genetically engineered **CHO cells** (Chinese Hamster Ovary cells), which perform correct human-like protein folding and glycosylation. mAbs treat cancers (e.g. trastuzumab against certain breast cancers), autoimmune diseases, and serve in diagnostics. A promising low-cost alternative is the production of antibodies in transgenic plants (“plantibodies”), for example in tobacco.

PROPERTY mRNA vaccines

An **mRNA vaccine** delivers, inside **lipid nanoparticles**, a messenger RNA molecule that codes for a pathogen's antigen (e.g. the spike protein of SARS-CoV-2). The patient's own cells translate this mRNA and display the antigen, triggering immunity. The mRNA works in the cytoplasm, is rapidly degraded, and never enters the nucleus nor integrates into the host genome. Advantages: extremely rapid design (days once the sequence is known) and no risk of infection since no live pathogen is used.

WARNING A common misconception

mRNA vaccines do not modify your DNA. mRNA cannot integrate into the genome: it lacks the enzymes (integrase, reverse transcriptase) required, and it never crosses the nuclear envelope. Confusing mRNA vaccines with gene therapy is a classic exam error.

5.2 Agricultural applications: GM crops and gene editing


DEFINITION GMO

A **Genetically Modified Organism** (GMO) is an organism whose genetic material has been modified by genetic engineering techniques (introduction, deletion or targeted modification of one or more genes), in a way that does not occur naturally by mating or natural recombination.

Three flagship examples dominate world agriculture:

- **Bt cotton** and Bt maize carry a gene from the bacterium *Bacillus thuringiensis* coding for a protein toxic to insect larvae (bollworms, stem borers) but harmless to humans. Result: drastic reduction of insecticide spraying — a major benefit for smallholder farmers, including in West and Central Africa.
- **Herbicide-tolerant maize and soybean** carry a gene (e.g. a modified EPSPS enzyme) allowing them to survive glyphosate, so weeds can be controlled without killing the crop.
- **Golden Rice** is engineered to produce β -carotene (provitamin A) in the grain, to fight vitamin A deficiency, which causes blindness in hundreds of thousands of children yearly in developing countries.

Beyond transgenesis, **gene editing** (CRISPR–Cas9) now allows precise modification of a plant's own genes, without introducing foreign DNA. For example, editing rice genes that control stomatal closure in response to the stress hormone ABA (abscisic acid) improves **drought tolerance** — a crucial trait as rainfall becomes erratic in the Sudano-Saharan zone of northern Cameroon.



Simplified world map of GM crop adoption. The Americas dominate cultivation; Africa (including Cameroon) proceeds cautiously under biosafety law.

METHOD Risk assessment of a GMO

Before any release, a GMO is evaluated in three steps:

1. Comparative analysis (principle of substantial equivalence): compare the GM plant with its conventional counterpart — composition, nutrients, toxins, allergens. Any significant difference triggers deeper study.
2. Environmental impact study: gene flow to wild relatives, effects on non-target organisms (e.g. pollinators), persistence in soil.
3. Management plan: monitoring after release, refuge zones to delay resistance, traceability and labelling.

WARNING Resistance evolution

Widespread planting of Bt crops exerts strong selection pressure: rare resistant caterpillars survive, reproduce, and can produce fully resistant populations within years. The same applies to weeds exposed repeatedly to one herbicide. This is evolution by natural selection in action — the remedy is the refuge strategy (non-Bt plots where susceptible insects survive, diluting resistance genes). Never write that “the insecticide creates the mutation”: mutations arise randomly; the insecticide only selects them.

5.3 Environmental applications: cleaning up with life

DEFINITION Bioremediation

Bioremediation is the use of living organisms — micro-organisms (bacteria, fungi) or plants — to degrade, transform or immobilise pollutants in contaminated soils, water or sediments, restoring the environment.

How it works. Many bacteria possess enzymes that oxidise hydrocarbons and use them as carbon sources. After an oil spill, one can either seed specialised bacteria (bioaugmentation) or add nutrients (nitrogen, phosphorus) to stimulate indigenous oil-degrading bacteria such as *Pseudomonas* species (biostimulation). Other micro-organisms transform toxic heavy metals: for example, certain bacteria reduce the highly toxic and mobile Cr(VI) to the less toxic, less mobile Cr(III).

Phytoremediation. Plants can also clean up: hyperaccumulator plants (e.g. some Brassica species) absorb heavy metals such as cadmium, zinc or nickel from contaminated soils into their harvestable shoots; the plants are then collected and treated. This is slow but cheap and aesthetically acceptable — relevant for mining areas in Cameroon.

Bioindicators. Living organisms also reveal pollution: the absence of sensitive aquatic larvae (e.g. mayflies) in a stream, or the accumulation of pollutants in lichens, signals environmental degradation before chemical instruments detect it.

5.4 Energy applications: biofuels of the new generations

First-generation biofuels (ethanol from sugarcane or maize starch) compete with food crops. Biotechnology now targets:

- **Second-generation bioethanol:** lignocellulosic biomass (straw, bagasse, wood waste) is broken down by engineered enzymes (cellulases) into fermentable sugars — no competition with food.
- **Biodiesel from microalgae:** microalgae can accumulate oils (up to about 50% of dry mass in some species) and grow on non-arable land with far higher yields per hectare than oil palms.
- **Biohydrogen:** certain algae and bacteria produce H_2 by photosynthesis or fermentation, a clean fuel whose combustion releases only water.

5.5 Diagnostic applications: seeing the invisible

- **Real-time PCR (qPCR):** amplifies and quantifies pathogen DNA/RNA as the reaction proceeds (fluorescence measured each cycle). It detects viruses (HIV viral load, SARS-CoV-2) with extreme sensitivity.
- **ELISA:** enzyme-linked immunosorbent assay; an antigen-antibody reaction coupled to an enzyme producing a coloured signal proportional to the target (used for HIV screening).
- **Lateral flow rapid tests:** the principle of pregnancy tests and of the malaria RDTs used daily in Cameroonian health centres — a drop of blood migrates along a strip; a coloured line appears if the target antigen (e.g. Plasmodium proteins) is present. Result in about 15 minutes,

no laboratory needed.

- **Next-generation sequencing (NGS)**: millions of DNA fragments sequenced simultaneously, allowing identification of new pathogens, tracking of variants, and personalised medicine.

WORKED EXAMPLE 6 Interpreting a qPCR result

In a real-time PCR test for a virus, the C_t (cycle threshold) is the cycle number at which fluorescence exceeds the background. Sample A gives $C_t = 20$, sample B gives $C_t = 30$. Each cycle doubles the DNA. Which patient has the higher initial viral load, and by approximately what factor?

SOLUTION

Solution. A lower C_t means the signal was reached earlier, so the starting material was more abundant: patient A has the higher viral load. The difference is $\Delta C_t = 30 - 20 = 10$ cycles. Since each cycle doubles the DNA:

$$\text{factor} = 2^{\Delta C_t} = 2^{10} = 1024.$$

Patient A therefore started with approximately 1024 times more viral genetic material than patient B. Common mistake: thinking a high C_t means high viral load — it is the opposite.

5.6 The Cameroonian and international regulatory framework

Powerful technologies require governance. Cameroon is a Party to the **Cartagena Protocol on Biosafety** (an international agreement under the Convention on Biological Diversity) which regulates the transboundary movement of living modified organisms, based on the precautionary principle and on Advance Informed Agreement: an exporting country must inform and obtain consent from the importing country.

At national level, **Law No. 2003/006 of 21 April 2003** lays down the safety rules governing modern biotechnology in Cameroon. It established the **National Biosafety Committee**, a multidisciplinary body (scientists, administrators, civil society) that examines applications for contained use, field trials or import of GMOs, conducts risk assessments, and advises the government before any authorisation. Thus, while Cameroon develops biotechnological capacity (e.g. molecular diagnosis of malaria and COVID-19, research on improved crop varieties), every step is framed by biosafety rules protecting human health and our exceptional biodiversity.

KEY POINTS Key points of Section 5

- Recombinant DNA technology produces human insulin, growth hormone, factor VIII, monoclonal antibodies (hybridomas, CHO cells, plants) and mRNA vaccines (lipid nanoparticles, no genomic integration).

- GM crops (Bt cotton, herbicide-tolerant maize, Golden Rice) and CRISPR-edited crops (drought-tolerant rice) improve yields and nutrition, but require rigorous risk assessment and management of resistance evolution.
- Bioremediation and phytoremediation use microbes and plants to clean polluted environments; bioindicators monitor pollution.
- Advanced biofuels (cellulosic ethanol, algal biodiesel, biohydrogen) avoid competition with food.
- Diagnostics (qPCR, ELISA, lateral flow tests, NGS) revolutionise disease detection — vital for malaria and emerging diseases in Cameroon.
- In Cameroon, biotechnology is regulated by Law No. 2003/006 and the National Biosafety Committee, under the Cartagena Protocol.

EXERCISE 5 Practice questions

1. Explain why recombinant human insulin is preferred to insulin extracted from animal pancreases (three reasons).
2. A farmer plants Bt cotton on his entire farm for ten consecutive years. Predict, with justification, the likely evolution of the bollworm population, and propose a strategy to prevent it.
3. Distinguish biostimulation from bioaugmentation in the treatment of an oil-polluted mangrove.
4. In a qPCR test, sample X has $C_t = 18$ and sample Y has $C_t = 24$. Calculate the ratio of their initial template quantities.

SOLUTION Exercise 5

1. Recombinant insulin is (i) identical to human insulin, so no allergic reactions; (ii) available in unlimited quantities from fermenters, independent of animal supply; (iii) cheaper and free of animal pathogens.
2. Continuous exposure to the Bt toxin selects rare resistant caterpillars; they survive and reproduce, so the frequency of resistance alleles rises until the population becomes largely resistant (directional selection). Prevention: maintain refuge plots of non-Bt cotton nearby, where susceptible insects survive and mate with resistant ones, diluting resistance alleles.
3. Biostimulation = adding nutrients (N, P) to stimulate indigenous pollutant-degrading microbes already present. Bioaugmentation = introducing specialised degrading micro-

organisms from outside into the polluted site.

4. $\Delta C_t = 24 - 18 = 6$; ratio = $2^6 = 64$. Sample X contained about 64 times more initial template than sample Y.

6 INTEGRATION ACTIVITY

7 INTEGRATION ACTIVITY: CASSAVA VALORISATION BY RECOMBINANT YEAST

7.1 Aims of the activity

This integration activity invites you to act as a junior biotechnologist in a Cameroonian agro-food research team. By the end of the activity, you should be able to:

- **Analyse** the scientific and technical limitations of a traditional biotechnological process (artisanal cassava fermentation) using your knowledge of microbial metabolism;
- **Design** a recombinant DNA construct (promoter – gene of interest – terminator – selectable marker) suitable for expressing a thermostable α -amylase in the yeast *Saccharomyces cerevisiae*;
- **Plan** a screening and assay strategy (starch-iodine test, reducing-sugar assay) to select and characterise transformed clones;
- **Evaluate** a pilot-scale bioprocess quantitatively (yield, productivity) and critically (biosafety, cost, social acceptability);
- **Communicate** scientific results through a poster and an oral defence, as expected at the GCE Advanced Level.

The activity is carried out in **groups of four** and mobilises the core lessons of this chapter: traditional biotechnology, recombinant DNA technology, gene alteration, screening methods, and the applications and limits of biotechnology.

7.2 Situation

Cassava in Cameroon: an opportunity and a bottleneck.

Cassava (*Manihot esculenta*) is one of the most important food crops in Cameroon. In the Littoral, Centre and South regions, tubers are processed by artisanal fermentation into products such as bâton de manioc (bobolo/miondo), gari and kumkum. Fermentation is carried out by naturally occurring micro-organisms (mainly lactic acid bacteria and yeasts) present on the tubers and in the environment. This is a typical example of **traditional biotechnology**: it softens the tissue, develops flavour and, crucially, helps to degrade the **cyanogenic glucosides** (linamarin and lotaustralin) which release toxic hydrogen cyanide (HCN) when the tissue is damaged.

However, a field survey conducted by a (fictional) cooperative in Édéa, “Coopérative Manioc Plus”, revealed the following problems:

Problem observed	Data from the survey
Residual cyanide	Some batches of bobolo still contain HCN levels considered unsafe when fermentation is too short (less than 3 days).
Variable quality	Fermentation time varies from 3 to 7 days depending on ambient temperature and the chance micro-organisms present; taste and texture are inconsistent.
Low starch valorisation	The cooperative wants to produce glucose syrup from cassava starch, but the starch must first be hydrolysed. Industrial hydrolysis uses purified α -amylase bought abroad, which is expensive.
Yield	Only about 60 kg of marketable product is obtained per 100 kg of fresh tubers.

The project.

The cooperative contacts your team, based at a university biotechnology laboratory in Douala. Your mission: design a **mini-project** to improve cassava valorisation using modern biotechnology. The central idea is to construct a strain of the baker's yeast *Saccharomyces cerevisiae* that secretes a **thermostable α -amylase** from the bacterium *Bacillus licheniformis*. Such a yeast could, in a single step, liquefy cassava starch at high temperature (90–95 °C for the enzyme step) and then ferment the released sugars, reducing the need to import purified enzyme.

Data provided to your team.

- The *B. licheniformis* α -amylase gene (*amyL*) is about 1.5 kb long; its sequence is available in public databases.
- You have a **shuttle plasmid** (replicating in both *E. coli* and yeast) carrying: an origin of replication for *E. coli*, a yeast 2μ origin, a strong constitutive yeast promoter (P_{PGK}), a multiple cloning site (MCS), a yeast terminator (T_{CYC1}), an ampicillin-resistance gene (*bla*, for selection in *E. coli*) and the *URA3* gene (selection marker in yeast: allows growth on medium lacking uracil).
- Restriction sites available in the MCS: *EcoRI*, *BamHI*, *HindIII*. The *amyL* gene can be amplified by PCR with *BamHI* and *HindIII* sites added at its two ends.
- Starch-iodine test: iodine gives a blue-black colour with starch; a clear halo around a colony on starch agar indicates starch hydrolysis.
- Reducing sugars can be assayed colorimetrically (DNS method): the absorbance at 540 nm is proportional to the glucose concentration.
- Pilot trial target: convert 10 kg of cassava starch per batch; native (non-transformed) yeast hydrolyses less than 5 % of the starch, while a good engineered strain should hydrolyse at least 80 %.

7.3 Tasks

Answer the following questions progressively. Questions 1–3 concern traditional biotechnology; questions 4–7 concern recombinant DNA technology; questions 8–10 concern evaluation and critical discussion.

1. Explain why artisanal cassava fermentation is described as biotechnology, and identify the type of metabolism (respiration or fermentation) mainly used by the micro-organisms involved.
2. Using the survey data, state **three** limitations of the artisanal process and, for each one, explain its biological or technological cause.
3. Explain the biochemical role of fermentation in the detoxification of cassava (cyanogenic glucosides → HCN). Why is an uncontrolled, spontaneous fermentation risky from a public-health point of view?
4. The team chooses *S. cerevisiae* as the host rather than *E. coli*. Give **two** reasons that justify this choice for a food application.
5. Draw a labelled diagram of the final expression cassette as it will appear in the shuttle plasmid, showing: promoter, *amyL* gene, terminator, yeast selectable marker, and the two restriction sites used for cloning. Indicate the orientation of each element.
6. Describe, in numbered steps, how you would obtain the recombinant plasmid, from the *B. licheniformis* chromosome to a transformed yeast colony. Name the key enzymes used at each step.
7. Explain the role of the *URA3* marker during the selection of transformed yeasts. What culture medium would you use, and why can untransformed yeasts not grow on it?
8. Describe the **screening test** on starch agar that distinguishes yeast clones secreting active α -amylase from negative clones. What result do you expect for a positive clone, and why?
9. In the pilot trial, a transformed strain hydrolyses 82 % of the 10 kg of starch per batch, and 1 kg of starch yields theoretically 1.1 kg of glucose. Calculate the mass of glucose expected per batch, then the mass actually recovered if the measured recovery is 90 % of the theoretical value. Compare with the native strain (5 % hydrolysis) and conclude.
10. Discuss **three** points that must be examined before the engineered yeast is used at village scale: one biosafety point, one economic point, and one social/ethical point. Suggest a precaution for each.

7.4 Model answer

Question 1. Artisanal cassava fermentation is biotechnology because it uses **living micro-organisms** (lactic acid bacteria and yeasts naturally present on the tubers) to transform a raw material into a useful product with improved properties (texture, flavour, safety, preservation). The micro-organisms mainly use **anaerobic metabolism**: yeasts perform alcoholic fermentation (glucose → ethanol + CO₂) and lactic acid bacteria perform lactic fermentation (glucose → lactic acid), which acidifies the product and inhibits spoilage organisms.

Question 2. Three limitations:

- Residual cyanide: detoxification depends on the enzyme linamarase, produced by the plant tissue and by some fermenting microbes; if fermentation is too short, linamarin is not fully degraded and HCN remains in the food.
- Variable quality: the process is spontaneous — the microbial community is not controlled,

so its composition, and therefore the duration (3–7 days) and the sensory result, vary with temperature and chance contamination.


- Low valorisation of starch: the artisanal microbes do not produce enough amylase to hydrolyse starch into fermentable or marketable sugars, so glucose-syrup production is impossible without imported enzymes, which raises costs.

Question 3. When cassava cells are crushed, the glucoside linamarin comes into contact with the enzyme linamarase (β -glucosidase), which hydrolyses it into glucose and acetone cyanohydrin; the cyanohydrin then decomposes spontaneously, releasing HCN. Prolonged fermentation, combined with pressing and drying/cooking, allows the volatile HCN to escape. A spontaneous, uncontrolled fermentation is risky because neither the duration nor the microbial enzymatic activity is guaranteed: a short or failed fermentation leaves dangerous residual cyanide, which can cause acute poisoning or, chronically, neurological disorders (e.g. konzo).

Question 4. Two reasons for choosing *S. cerevisiae*:

1. It has a long history of safe use in food (bread, beer, palm wine) and is **Generally Recognised As Safe (GRAS)**, unlike *E. coli*, which is associated with disease and may produce endotoxins (lipopolysaccharides) undesirable in food.
2. As a eukaryote, it performs proper protein folding, post-translational modifications and **secretion** via the ER-Golgi pathway; moreover, it naturally ferments sugars to ethanol, so the same organism can both help hydrolyse starch and ferment the released sugars in a consolidated bioprocess.

Question 5. Expression cassette in the shuttle plasmid (orientation 5' → 3'):



Map of the recombinant shuttle plasmid: the *amyL* gene is inserted between the *Bam*HI and *Hind*III sites of the MCS, under the control of the yeast promoter P_{PGK} and terminator T_{CYC1} ; *URA3* selects transformed yeasts.

Question 6. Steps to obtain the recombinant plasmid:

1. Extract chromosomal DNA from *B. licheniformis*.

2. Amplify the amyL gene by **PCR** using Taq DNA polymerase and primers carrying a BamHI site (forward) and a HindIII site (reverse).
3. Digest the purified PCR product and the shuttle plasmid separately with the **restriction enzymes** BamHI and HindIII; the two different sticky ends ensure directional cloning (correct orientation).
4. Join the insert and the vector with **T4 DNA ligase**.
5. Transform competent E. coli (e.g. by heat shock or electroporation); select on LB-ampicillin medium (only bacteria carrying the plasmid grow).
6. Screen colonies (colony PCR or restriction analysis of extracted plasmids) to confirm the 1.5 kb insert; verify by DNA sequencing.
7. Amplify the verified plasmid in E. coli, purify it (miniprep), and transform S. cerevisiae (e.g. by lithium acetate/PEG method).
8. Plate the yeasts on synthetic defined medium **lacking uracil** (SD-Ura): only transformed clones grow.

Question 7. URA3 encodes orotidine-5'-phosphate decarboxylase, an enzyme required for uracil biosynthesis. The host yeast strain is $ura3^-$ (auxotrophic for uracil): it cannot synthesise its own uracil. On a medium lacking uracil, only yeasts that have taken up the plasmid (and therefore a functional URA3) can synthesise uracil and grow; untransformed yeasts die. This is **selection** (survival based on a vital function), distinct from **screening** (detection of a specific phenotype like enzyme activity).

Question 8. Screening on starch agar: replica-plate or streak the selected yeast colonies on agar containing soluble starch, incubate at 30 °C for 24–48 h, then flood the plate with iodine solution (iodine-potassium iodide). Starch forms a blue-black complex with iodine everywhere except around colonies that secrete active α -amylase: there, starch has been hydrolysed to maltose/glucose, so a **clear halo** appears. Positive clone = colony surrounded by a clear halo; the larger the halo, the higher the amylase activity. The best clones are then confirmed by the quantitative DNS assay of reducing sugars (absorbance at 540 nm).

Question 9. Calculations for one 10 kg batch of starch:

$$\begin{aligned}
 m_{\text{starch hydrolysed}} &= 10 \times 0.82 = 8.2 \text{ kg} \\
 m_{\text{glucose, theoretical}} &= 8.2 \times 1.1 = 9.02 \text{ kg} \\
 m_{\text{glucose, recovered}} &= 9.02 \times 0.90 = 8.118 \text{ kg} \approx 8.1 \text{ kg}
 \end{aligned}$$

For the native strain (5 % hydrolysis, same recovery):

$$m_{\text{glucose, native}} = 10 \times 0.05 \times 1.1 \times 0.90 = 0.495 \text{ kg} \approx 0.50 \text{ kg}$$

The engineered strain therefore recovers about **16 times more glucose** per batch (8.1 kg versus 0.50 kg) and exceeds the 80 % hydrolysis target: the pilot trial is successful.

Question 10. Discussion before village-scale use:

- Biosafety: the engineered yeast must be used in closed vessels to avoid uncontrolled release into the environment; the final food strain carries no antibiotic-resistance gene (bla is

only for the *E. coli* cloning step) and URA3 is a harmless metabolic gene. **Precaution:** contain the process physically and inactivate the yeast by heat treatment ($> 70^{\circ}\text{C}$) before the product is sold.

- Economic: producing glucose syrup locally with the engineered yeast removes the recurring cost of imported purified α -amylase, but the upstream laboratory steps (PCR, sequencing, strain verification) are capital-intensive. **Precaution:** once the strain is built and validated, maintain it as a stable starter culture that the cooperative can multiply cheaply on-site (like a bread-yeast or palm-wine starter), avoiding repeated lab interventions.
- Social/ethical: consumers may fear “genetically modified” food and reject the product. **Precaution:** engage the community early through the cooperative, explain that the yeast is a **processing aid** removed/inactivated before consumption (similar in spirit to traditional fermentation starters), and seek regulatory approval from the relevant Cameroonian authorities (e.g. MINSANTE, MINADER) for transparent labelling.

WARNING Common mistakes to avoid

Do not confuse selection (a marker that keeps only transformed cells alive, e.g. URA3 on uracil-free medium) with screening (a test that reveals which survivors actually express the desired activity, e.g. the starch-iodine halo). In the cloning scheme, the insert and the vector must be cut with the same restriction enzymes, and the promoter must be placed upstream (5') of the gene, in the correct orientation. In yield calculations, always apply the percentages sequentially (hydrolysis $\times$ theoretical yield $\times$ recovery) and state your units at every step.

KEY POINTS What this activity consolidates

Traditional fermentation is real biotechnology but suffers from variability and safety limits; recombinant DNA technology (PCR, restriction enzymes, ligase, shuttle vector, selectable marker) allows a precise, heritable improvement of a food micro-organism; screening assays verify enzyme activity; and a responsible biotechnologist always evaluates yield and biosafety, cost and social acceptability before scaling up.

DID YOU KNOW? Link to medical biotechnology (Homeostasis)

The same recombinant DNA principles used here to produce a thermostable enzyme are applied to produce human proteins for therapy: **human insulin** (diabetes homeostasis), **growth hormone**, **clotting factors** and **vaccines** (e.g. hepatitis B surface antigen in yeast). These are major applications of biotechnology in the service of human homeostasis, directly addressing Lesson 84 of the syllabus.

8 METHODS AND WORKED EXAMPLES

9 BIOTECHNOLOGY




9.1 Skill 1: Analysing a traditional fermentation process

METHOD How to describe and explain a traditional fermentation

When a GCE question asks you to describe a traditional biotechnological process (palm wine, fufu water, burukutu, yoghurt, bread), proceed in numbered steps:

1. Identify the substrate (the raw material: palm sap, milk, cassava, dough) and the micro-organism responsible (yeast *Saccharomyces cerevisiae*, lactic acid bacteria *Lactobacillus* spp., *Acetobacter* spp.).
2. State the type of respiration: fermentation is **anaerobic** (occurs without oxygen).
3. Write the balanced summary equation. For alcoholic fermentation: $C_6H_{12}O_6 \longrightarrow 2C_2H_5OH + 2CO_2 + \text{small amount of energy (2 ATP)}$. For lactic fermentation: $C_6H_{12}O_6 \longrightarrow 2CH_3CHOHCOOH + 2ATP$.
4. Explain the conditions: warmth (optimum about 25–35 °C), absence of air, presence of fermentable sugar, time, pH near neutral initially.
5. Link the products to the observed changes: bubbles = CO_2 ; sour taste = lactic acid; alcohol smell = ethanol; preservation = acid/alcohol inhibits spoilage microbes.
6. Conclude with the definition: traditional biotechnology is the use of living micro-organisms or their enzymes to transform raw materials into useful products, without altering their genetic material.

Reflex: always mention that fermentation yields only 2 ATP per glucose (versus about 36–38 in aerobic respiration), because the electron transport chain is not used and glucose is only partially oxidised.

Alcoholic fermentation:**Lactic fermentation:****Acetic oxidation (aerobic):**

Traditional palm wine fermentation in a calabash: spontaneous mixed-culture fermentation.


WORKED EXAMPLE 7**Fermentation of palm wine in the South–West Region**

A student in Buea collects fresh palm sap in a clean calabash. After 6 hours the sap is sweet and slightly fizzy; after 24 hours it tastes strongly alcoholic and sour. (a) Name the micro-organism mainly responsible for the first stage. (b) Write the equation of the reaction occurring. (c) Explain why the sap becomes fizzy. (d) Explain why the drink becomes sour after 24 hours. (e) State two reasons why palm wine spoils faster than bottled beer.

SOLUTION

(a) The first stage is carried out by yeasts (mainly *Saccharomyces* species) naturally present on the palm tree bark, in the air and in the calabash. No starter culture is added: this is a spontaneous fermentation, characteristic of traditional biotechnology.


(b) The yeasts perform alcoholic fermentation of the sugars (mainly sucrose, glucose and fructose) in the sap:



This is anaerobic respiration: glucose is only partially oxidised, so most of the chemical energy remains locked in the ethanol.

(c) The fizziness is caused by the carbon dioxide released by the fermentation. The gas dissolves in the sap and escapes as bubbles, exactly as in bread dough rising or in the brewing of burukutu (millet beer) in the North.

(d) After about 24 hours, lactic acid bacteria (*Lactobacillus*) and acetic acid bacteria (*Acetobacter*) multiply in the wine. Lactic acid bacteria convert remaining sugars to lactic acid (lactic fermentation), and acetic acid bacteria oxidise ethanol to ethanoic (acetic) acid in the presence of air:



The accumulation of these acids lowers the pH and gives the sour taste. This is why palm wine left overnight is used like vinegar in some households.

(e) Palm wine spoils faster than bottled beer because:

- it is not pasteurised or filtered, so the mixed microbial population (yeasts, lactic and acetic bacteria) remains alive and active;
- it is kept in open or non-sterile containers at ambient tropical temperature (25–30 °C), which favours rapid microbial growth and exposure to oxygen (needed by acetic acid bacteria).


Examiner's tip: in part (b) do not forget to state that the process is anaerobic and releases only 2 ATP; this is the marking point most often missed.

9.2 Skill 2: Interpreting a fermentation experiment (data analysis)

METHOD How to analyse gas-production or growth data in biotechnology

1. Read the axes and units first; identify the independent variable (time, temperature, sugar concentration, pH) and the dependent variable (volume of CO₂, optical density, alcohol content, pH change).
2. Describe the curve in phases: lag phase (slow start, cells adapting), exponential/active phase (rapid gas production or growth), stationary phase (substrate exhausted or product inhibiting), death/decline phase.
3. Explain each phase biologically, not just descriptively: e.g. stationary phase because sugar is finished or ethanol has reached a toxic concentration for the yeast.
4. Compare treatments by quoting figures (rates = gradient of the curve in the exponential phase).
5. Draw a conclusion that answers the aim, and state one control variable that had to be kept constant (temperature, volume, yeast mass, sugar concentration, pH, anaerobiosis).

Reflex: rate = $\frac{\text{change in dependent variable}}{\text{time}}$; always give units (e.g. cm³ min⁻¹).



Typical fermentation curves at different temperatures. At 37°C: rapid exponential phase, high yield. At 10°C: slow throughout. At 70°C: no activity (enzymes denatured).

WORKED EXAMPLE 8 Effect of temperature on yeast fermentation

Equal masses of yeast and glucose solution were placed in four water baths and the volume of CO₂ collected after 20 minutes was measured:

Temperature (°C)	10	25	37	70
Volume of CO ₂ (cm ³)	4	22	35	0

(a) Calculate the rate of CO₂ production at 37 °C. (b) Explain the results at 10 °C and at 70 °C. (c) State two variables that must be controlled. (d) Suggest the optimum temperature and justify.

SOLUTION

(a) Rate = $\frac{\text{volume of CO}_2}{\text{time}} = \frac{35 \text{ cm}^3}{20 \text{ min}} = 1.75 \text{ cm}^3 \text{ min}^{-1}$.

(b) At 10 °C the rate is low (0.2 cm³ min⁻¹) because the enzymes of the yeast (zymase complex) have low kinetic energy: few enzyme–substrate collisions occur per unit time, so fermentation is slow but the yeast is not dead. At 70 °C no gas is produced because the high temperature has denatured the enzymes: the tertiary structure and the active site are destroyed irreversibly, and the yeast cells are killed. No fermentation can occur.

(c) Variables to control: the mass/concentration of yeast, the concentration and volume of glucose solution, the duration of the experiment, pH, and absence of oxygen (anaerobic conditions).

(d) The optimum is around 37 °C (possibly slightly below or above), because this temperature gave the greatest volume of gas in the same time, i.e. the highest rate. To confirm, the experiment should be repeated at 30, 35, 40 and 45 °C to narrow down the peak.

Examiner's tip: "the yeast is killed by cold" is a classic error: cold only slows enzyme activity; only heat denatures enzymes irreversibly.

9.3 Skill 3: Biotechnology at the service of homeostasis, support and locomotion

DEFINITION Biotechnology for homeostasis, support and locomotion

This refers to the application of biotechnological products and techniques to maintain internal stability (homeostasis), provide structural support, and enable movement (locomotion) in humans and animals. Key examples include recombinant hormones, biomaterials, tissue engineering, and assistive devices.

PROPERTY Major applications

- Glucose homeostasis: Recombinant human insulin (produced in *E. coli* or yeast) and glucagon for diabetes mellitus treatment. Continuous glucose monitors (CGMs) and insulin pumps (biotechnology + engineering).
- Calcium/phosphate homeostasis: Recombinant calcitonin and parathyroid hormone analogues for osteoporosis and hypercalcaemia.
- Growth and metabolism: Recombinant human growth hormone (somatropin) for growth hormone deficiency, Turner syndrome, chronic renal failure.
- Red blood cell homeostasis: Recombinant erythropoietin (EPO) for anaemia in chronic kidney disease and chemotherapy.
- Support (skeletal repair): Biomaterials – hydroxyapatite coatings, collagen scaffolds, biodegradable polymers (PLA, PGA) for bone grafts and fracture fixation. Tissue engineering of cartilage and bone using mesenchymal stem cells on 3D scaffolds.
- Locomotion (joint/muscle repair): Viscosupplementation with hyaluronic acid (produced by bacterial fermentation) for osteoarthritis. Monoclonal antibodies (e.g. anti-TNF α like adalimumab) for rheumatoid arthritis. Myostatin inhibitors (experimental) for muscular dystrophies.
- Prosthetics and interfaces: Osseointegrated implants (titanium with bioactive coatings), neural interfaces for bionic limbs.

EXAMPLE Insulin and glucose homeostasis

In type 1 diabetes, pancreatic β -cells are destroyed $\rightarrow$ no insulin $\rightarrow$ chronic hyperglycaemia. Before 1982, insulin was extracted from pig/cow pancreas (slightly different amino acid sequence $\rightarrow$ allergic reactions, limited supply). Recombinant DNA technology allowed production of identical human insulin in bacteria/yeast: unlimited supply, no immune reaction, consistent quality. This is a direct application of biotechnology to re-

store glucose homeostasis.

EXPERIMENT Testing the effect of insulin on blood glucose (simulation)

Aim: Demonstrate the role of insulin in glucose uptake by cells. **Materials:** Yeast suspension (model for cells), glucose solution, recombinant insulin, Benedict's reagent, water bath. **Method:** Set up three tubes: (1) yeast + glucose, (2) yeast + glucose + insulin, (3) yeast + water (control). Incubate at 37 °C for 30 min. Test supernatant for remaining glucose with Benedict's reagent. **Expected:** Tube 2 shows greatest glucose uptake (least reducing sugar) because insulin stimulates glucose transporters. Tube 1 shows moderate uptake. Tube 3 shows no uptake. **Conclusion:** Insulin enhances cellular glucose uptake, lowering blood glucose – a homeostatic mechanism restored by biotechnological insulin.

9.4 Skill 4: Correction and gene remediation – gene therapy and genome editing

METHOD How to structure an answer on gene correction (gene therapy / CRISPR)

1. Define: gene therapy is the introduction of a normal functional allele into a patient's cells to correct a genetic defect; genome editing (e.g. CRISPR–Cas9) directly modifies the faulty DNA sequence.
2. Distinguish the two types: somatic gene therapy (body cells; affects only the patient; not inherited) and germ-line therapy (gametes/early embryo; heritable; ethically controversial and generally prohibited internationally).
3. Give the steps for somatic therapy: isolate the normal gene → insert into a vector (often a disabled virus: adenovirus, adeno-associated virus AAV, lentivirus/retrovirus, or non-viral liposomes/nanoparticles) → deliver to target cells (ex vivo or in vivo) → gene expressed → functional protein produced.
4. For CRISPR–Cas9: a guide RNA (gRNA) directs the Cas9 nuclease to the target sequence; Cas9 cuts both strands; the cell repairs the break via NHEJ (error-prone) or HDR (homology-directed repair with a correct template), allowing the faulty base to be corrected.
5. Evaluate: advantages (treats the cause, not symptoms; potential cure) and limitations (delivery efficiency, immune reaction to vector, risk of off-target cuts/insertional mutagenesis, high cost, ethical issues for germ-line).

Reflex: always state whether the correction is heritable or not – this is the key marking point.

Somatic gene therapy: Normal gene therapy. Inserted into Ex vivo: patient's haematopoietic cells. Transfused. Corrected stem cells infused.

CRISPR-Cas9 genome editing: gRNA + Cas9 protein. Complex enters nucleus. Cas9 cuts both DNA strands. HDR repair with correct template.

Comparison of somatic gene therapy (ex vivo lentiviral) and CRISPR-Cas9 genome editing for sickle cell disease.

WORKED EXAMPLE 9 Gene therapy for sickle cell disease

Sickle cell disease is common in Cameroon. Scientists propose two strategies: (A) introducing a normal β -globin gene into the patient's bone-marrow stem cells using a modified virus; (B) editing the faulty gene directly in embryos using CRISPR-Cas9. (a) Classify each strategy as somatic or germ-line. (b) Explain why bone-marrow stem cells are chosen in strategy A. (c) State the role of the guide RNA in strategy B. (d) Give one ethical objection to strategy B.

SOLUTION

(a) Strategy A targets the body (somatic) cells of an already-born patient: it is somatic gene therapy. Its effects are limited to the treated individual and are not transmitted to offspring. Strategy B modifies an embryo, whose cells include the future germ line: it is germ-line editing, and the change would be heritable.

(b) Bone-marrow haematopoietic stem cells are chosen because they are the cells that continuously divide and differentiate to produce all red blood cells throughout life. Correcting the gene in these stem cells means that all red blood cells produced afterwards will contain normal haemoglobin, giving a long-lasting (potentially permanent) treatment rather than a temporary one.

(c) The guide RNA (gRNA) has a sequence complementary to the target DNA (the mutated region of the β -globin gene, specifically the A $\rightarrow$ T transversion at codon 6). It binds the target by base pairing and thereby directs the Cas9 enzyme to exactly the right position, where Cas9 cuts both DNA strands. The cell's repair machinery then fixes the break using a correct template (provided or endogenous), restoring the normal sequence (GAG $\rightarrow$ Glu instead of GTG $\rightarrow$ Val).

(d) Ethical objections to embryo editing include: the modification is heritable and affects future generations without their consent; off-target cuts could introduce new harmful mutations into the human gene pool; and it opens the way to "designer babies" (selection for non-medical traits), which raises issues of justice, equity and human dignity.

Examiner's tip: note the link with malaria: the Hb^S allele is maintained in Central Africa because heterozygotes (Hb^AHb^S) are more resistant to severe malaria – an example of bal-

anced polymorphism that examiners love to combine with this topic.


9.5 Skill 5: Describing the steps of recombinant DNA technology

METHOD How to present the production of a transgenic organism (e.g. insulin)

Learn the sequence as a numbered chain and be able to name the enzyme or tool at each step:

1. Isolate the gene of interest (e.g. human insulin gene) from donor DNA, or obtain its mRNA and make cDNA using reverse transcriptase (avoids introns).
2. Cut the donor DNA (or cDNA) and the vector (usually a bacterial plasmid) with the same restriction endonuclease, producing complementary sticky ends (or blunt ends).
3. Join the gene into the plasmid using DNA ligase → recombinant DNA (rDNA).
4. Introduce the recombinant plasmid into a host cell (e.g. *E. coli*) by transformation (heat shock or electroporation).
5. Select the transformed cells using a marker gene (e.g. antibiotic resistance, fluorescence) carried by the plasmid.
6. Screen/identify clones containing the correct insert (e.g. blue-white screening, PCR, restriction digest).
7. Culture the host cells in a fermenter (large-scale cloning = gene amplification); the cells express the gene (transcription + translation).
8. Extract and purify the product (e.g. insulin) for medical use (downstream processing).

Reflex: restriction enzyme = “molecular scissors”; ligase = “molecular glue”; plasmid = “vector/vehicle”; host bacterium = “factory”; marker gene = “selection tool”.



Step-by-step production of recombinant human insulin in *E. coli*.

WORKED EXAMPLE 10 Producing human insulin in bacteria

(a) Explain why the same restriction enzyme must be used to cut both the human DNA and the plasmid. (b) Name the enzyme that seals the insulin gene into the plasmid. (c) The plasmid carries a gene for resistance to the antibiotic ampicillin. Explain how this is used to identify bacteria that took up the plasmid. (d) Give two advantages of bacterial insulin over insulin extracted from pig pancreas.

SOLUTION

(a) A restriction endonuclease cuts DNA at a specific recognition sequence (e.g. EcoRI: 5'-GAATTC-3') and leaves characteristic single-stranded projections called sticky ends. Using the same enzyme on both the human DNA (or cDNA) and the plasmid produces complementary sticky ends whose base sequences can pair by hydrogen bonding (A with T, G with C). This allows the insulin gene to fit precisely into the opened plasmid. If different enzymes were used, the ends would not match and ligation would fail or be inefficient.

(b) DNA ligase seals the gene into the plasmid by forming phosphodiester bonds in the sugar-phosphate backbone, producing stable recombinant DNA.

(c) The bacteria are spread on agar containing ampicillin. Only bacteria that have taken up the plasmid carry the ampicillin-resistance marker gene (*bla*, encoding β -lactamase) and can survive and form colonies; bacteria without the plasmid are killed. The surviving

colonies are therefore the transformed bacteria, which can then be screened further for the insulin gene itself (e.g. by colony PCR or restriction analysis).

(d) Advantages of bacterial (recombinant) human insulin:

- it is identical to human insulin (same amino acid sequence), so it causes fewer allergic reactions and no immune rejection, unlike pig insulin which differs by one amino acid (Thr → Ala at B30);
- it can be produced in large, continuous quantities in fermenters, independent of animal supply, and is free from the risk of transmitting animal pathogens (e.g. prions, viruses).


Examiner's tip: always name the tool (restriction enzyme, ligase, plasmid, marker gene) and its role; a step without its enzyme earns only half the marks.

9.6 Skill 6: Reading restriction maps and gel electrophoresis

METHOD How to interpret a restriction digest and a gel

1. Restriction digest: count the number of cut sites for the enzyme; n cuts in a linear DNA molecule give $n + 1$ fragments; a circular plasmid gives n fragments (if $n > 0$).
2. Fragment sizes add up to the total DNA length; use this to check your answer.
3. Gel electrophoresis: DNA is negatively charged (phosphate groups), so it migrates to the positive electrode (anode); smaller fragments move faster and further through the agarose gel matrix.
4. Match bands to sizes: the band nearest the bottom (furthest from the wells) is the smallest fragment. Use a DNA ladder (marker) of known sizes to estimate fragment lengths.
5. Compare lanes (e.g. suspect vs crime-scene DNA, or normal vs mutant allele) to draw the conclusion.

Reflex: a mutation that destroys a restriction site gives fewer, larger fragments; a mutation that creates a site gives more, smaller fragments. This is the basis of RFLP analysis for sickle cell diagnosis.



RFLP analysis of β -globin gene with MstII. Normal allele (HbA) cut into 1.15 kb + 0.20 kb; sickle allele (HbS) uncut at 1.35 kb. Father and mother are carriers; child has sickle cell anaemia.

WORKED EXAMPLE 11 Diagnosing sickle cell anaemia by RFLP

The normal β -globin allele (Hb^A) contains a restriction site for the enzyme MstII that is destroyed by the sickle cell mutation (Hb^S). Digestion of the normal allele gives two fragments of 1.15 kb and 0.20 kb; the mutant allele gives a single fragment of 1.35 kb. DNA from a father (lane 1), mother (lane 2) and their child (lane 3) is digested and run on a gel. Lane 1 shows bands at 1.35 and 1.15 kb; lane 2 shows bands at 1.35, 1.15 and 0.20 kb; lane 3 shows a single band at 1.35 kb. (a) State the genotype of each person. (b) Explain why the 0.20 kb band may be faint or missing on a real gel. (c) What is the probability that a second child of this couple is also affected?

SOLUTION

(a) Interpretation of the bands:

- Lane 1 (father): bands at 1.35 kb (mutant allele, uncut) and 1.15 kb (cut normal allele; the 0.20 kb fragment runs off or is too faint). He carries both alleles: genotype Hb^AHb^S – a carrier (heterozygous), phenotypically normal.
- Lane 2 (mother): bands at 1.35, 1.15 and 0.20 kb: again both alleles present, genotype Hb^AHb^S – carrier.
- Lane 3 (child): only the 1.35 kb band: both alleles are mutant, genotype Hb^SHb^S – the child has sickle cell anaemia.

(b) The 0.20 kb fragment is very small: it binds little DNA stain (ethidium bromide or SYBR Safe), so it fluoresces weakly, and it migrates so fast that it may run off the end of the gel. Hence it is often faint or absent, and diagnosis relies on the 1.35/1.15 kb bands.

(c) Cross: Hb^AHb^S × Hb^AHb^S.

Gametes	Hb ^A	Hb ^S
Hb ^A	Hb ^A Hb ^A	Hb ^A Hb ^S
Hb ^S	Hb ^A Hb ^S	Hb ^S Hb ^S

$P(\text{affected child, Hb}^S\text{Hb}^S) = \frac{1}{4} = 0.25$, i.e. 25% for each pregnancy, independent of the first child's genotype.

Examiner's tip: this technique (RFLP + electrophoresis) is exactly how prenatal diagnosis of sickle cell disease is performed – a very relevant application in Cameroon, where carrier frequency is high (up to 25% in some zones). Do not confuse “carrier” (healthy heterozygote) with “affected”.

9.7 Skill 7: Evaluating applications of biotechnology (structured essay)

METHOD How to write a balanced “applications and implications” essay

1. Define biotechnology in one sentence: the use of living organisms, cells or their components (enzymes, DNA) to produce goods and services useful to humans.
2. Organise by sector – health, agriculture, environment, industry – with at least one precise example each:
 - Health: recombinant insulin, growth hormone, vaccines (hepatitis B, HPV), monoclonal antibodies (cancer, autoimmune), gene therapy, prenatal diagnosis of sickle cell disease, PCR diagnosis of HIV/malaria/COVID-19.
 - Agriculture: GM crops (Bt cotton resistant to insects, Golden Rice enriched in β -carotene/provitamin A), disease-resistant plantain/banana varieties (e.g. against BXW, Fusarium wilt), biofertilisers (Rhizobium, Azotobacter), marker-assisted selection (MAS) for cassava mosaic resistance.
 - Environment: bioremediation (bacteria degrading oil spills in the Niger Delta, heavy metal biosorption), sewage treatment (activated sludge, anaerobic digesters for biogas), biofuels (bioethanol from sugarcane/molasses, biodiesel from Jatropha), biosensors for pollution monitoring.
 - Industry: enzymes in detergents (proteases, lipases, amylases), brewing and food processing (rennin/chymosin for cheese, pectinases for fruit juice), bioplastics (PHA, PLA from microbial fermentation).
3. Discuss risks/concerns: gene flow to wild relatives (e.g. Bt gene to wild cotton), antibiotic-resistance markers (horizontal transfer risk), allergenicity of novel proteins, reduction of biodiversity (monocultures), cost and dependence of smallholder farmers on proprietary seeds, ethical issues (GMOs, germ-line editing, patenting)

life).

4. Conclude with a balanced judgement and the need for regulation/biosafety frameworks (Cartagena Protocol, national biosafety laws).

Reflex: one well-explained example per sector scores more than a long list of names. Use Cameroonian context where natural.

WORKED EXAMPLE 12 GM crops for Cameroonian agriculture?

“Genetically modified crops could improve food security in Cameroon, but their introduction must be carefully controlled.” Discuss this statement, giving at least two potential benefits and two potential risks.

SOLUTION

Introduction. Biotechnology offers tools, such as recombinant DNA technology, to introduce useful genes into crops. Whether Cameroon should adopt GM crops requires weighing benefits against risks within a national biosafety framework.

Potential benefits.

- Pest resistance: Bt crops carry a gene from the bacterium *Bacillus thuringiensis* coding for a Cry protein toxic to certain insect larvae (Lepidoptera, Coleoptera). Bt cotton or Bt maize would reduce losses to stem borers (*Busseola fusca*, *Chilo partellus*) and cut pesticide use, lowering costs for smallholder farmers and reducing chemical pollution of watercourses (e.g. Wouri, Sanaga).
- Improved nutrition and stress tolerance: biofortified crops (e.g. vitamin-A-enriched “Golden Rice”, provitamin-A cassava developed by IITA) can combat vitamin A deficiency, a real public-health problem in the North and Far North; drought-tolerant maize (WEMA project) could secure yields in the Sudano-Sahelian zone where rainfall is unreliable.
- Disease resistance: engineering plantains and cassava resistant to viral (e.g. banana bunchy top virus, cassava mosaic virus) and bacterial diseases (*Xanthomonas* wilt of banana) would protect staple foods of the forest and humid savanna zones.

Potential risks.

- Environmental: transgenes may spread by pollen to wild relatives (gene flow), possibly creating resistant weeds (e.g. wild cotton in the North); Bt toxin may affect non-target insects such as pollinators or beneficial predators, and pests may evolve resistance to the toxin (requiring refuge strategies).
- Health and socio-economic: possible allergenicity of new proteins must be tested rigorously (Codex Alimentarius guidelines); seed patents could make farmers dependent on companies for seed each season, threatening local seed-saving traditions

and agrobiodiversity (landraces of maize, sorghum, millet).

Conclusion. GM crops could indeed contribute to food security in Cameroon, but each variety must be assessed case by case under a biosafety framework (confined field trials, environmental risk assessment, food safety assessment, labelling, post-release monitoring), combined with protection of local varieties and support for agroecology. Biotechnology is a powerful tool, but not a substitute for good agronomy, infrastructure, and equitable policies.


Examiner's tip: a "discuss" question demands both sides plus a personal, justified conclusion; a one-sided answer cannot score full marks. Mention the Cartagena Protocol on Biosafety (Cameroon is a party) and the national biosafety law.

9.8 Skill 8: Calculations in biotechnology – PCR and microbial growth kinetics

METHOD How to calculate DNA amplification and microbial population growth

1. PCR: each cycle doubles the number of DNA molecules (assuming 100% efficiency). After n cycles from N_0 starting molecules: $N = N_0 \times 2^n$.
2. Bacterial growth (exponential phase): if the generation (doubling) time is g minutes, the number of generations in time t is $n = \frac{t}{g}$, and the final population is $N = N_0 \times 2^n$.
3. Solve for the unknown using powers of 2 or logarithms: $n = \frac{\log(N/N_0)}{\log 2}$.
4. Check units (minutes vs hours) and state assumptions (exponential phase, unlimited nutrients, no inhibition, synchronous division).

Reflex: PCR needs a thermostable DNA polymerase (Taq polymerase from *Thermus aquaticus*) because each cycle includes a $\sim 94^\circ\text{C}$ denaturation step; primers define the region amplified.



Bacterial growth curve in a batch fermenter. Exponential phase: constant generation time g . Stationary phase: nutrients exhausted, waste accumulated.

WORKED EXAMPLE 13 PCR amplification and fermenter culture

(a) Starting from a single DNA molecule, how many molecules are present after 30 PCR cycles? Give your answer in standard form to 3 significant figures. (b) A recombinant *E. coli* culture starts with 1.0×10^3 cells and divides every 20 minutes. Calculate the population after 4 hours, assuming exponential growth. (c) In practice the population stops growing exponentially. Give two reasons.

SOLUTION

(a) Each PCR cycle doubles the DNA, so after n cycles:

$$N = N_0 \times 2^n = 1 \times 2^{30} = 1\,073\,741\,824 \approx 1.07 \times 10^9 \text{ molecules (3 s.f.)}$$

This enormous amplification from a single molecule is why PCR is used in diagnosis (e.g. detecting HIV, malaria parasite *Plasmodium falciparum*, or SARS-CoV-2) and forensic analysis from tiny samples.

(b) Number of generations in 4 hours:

$$n = \frac{t}{g} = \frac{4 \times 60 \text{ min}}{20 \text{ min}} = 12 \text{ generations}$$

Final population:

$$N = N_0 \times 2^n = 1.0 \times 10^3 \times 2^{12} = 10^3 \times 4096 = 4.096 \times 10^6 \approx 4.10 \times 10^6 \text{ cells (3 s.f.)}$$

(c) Exponential growth stops because:

- nutrients (glucose, amino acids, phosphate) become exhausted in the fermenter;
- toxic waste products accumulate (e.g. organic acids lowering the pH, ethanol, ammonia), and oxygen may become limiting for aerobic cultures, so cells enter the stationary phase.

In industrial fermenters, sensors continuously monitor and adjust pH, temperature, dissolved oxygen and nutrient supply (fed-batch or continuous culture) to keep the culture productive for as long as possible.

Examiner's tip: the most frequent errors are forgetting to convert hours to minutes and writing $N = N_0 \times 2t$ instead of $N_0 \times 2^{t/g}$. Always compute the number of generations first.

9.9 Skill 9: Evaluation and remediation – integrated review

KEY POINTS

Key points for GCE Advanced Level Biotechnology

- Traditional biotechnology: uses wild-type micro-organisms, no genetic modification. Examples: palm wine, fufu, burukutu, yoghurt, bread, cheese. Fermentation = anaerobic, 2 ATP/glucose.
- Modern biotechnology: involves genetic manipulation (recombinant DNA, gene editing). Tools: restriction enzymes, ligase, vectors (plasmids, viruses), PCR, electrophoresis, CRISPR-Cas9.
- Recombinant DNA steps: isolate gene → cut vector/gene with same RE → ligate → transform → select (marker) → screen → scale up in fermenter → purify product.
- Gene therapy: somatic (non-heritable, e.g. lentiviral β -globin for sickle cell) vs germ-line (heritable, embryo editing, prohibited). CRISPR: gRNA guides Cas9 to target.
- RFLP diagnosis: mutation alters restriction site → fragment size change → gel band pattern reveals genotype. Sickle cell: MstII site lost in HbS.
- Applications: health (insulin, vaccines, mAbs, gene therapy), agriculture (Bt crops, biofortification, disease resistance), environment (bioremediation, biofuels), industry (enzymes, bioplastics).
- Risks/ethics: gene flow, resistance evolution, allergenicity, farmer dependence, germ-line heritability, equity. Regulation: biosafety laws, Cartagena Protocol.
- Calculations: PCR $N = N_0 2^n$; bacterial growth $N = N_0 2^{t/g}$. Always convert time to same units as generation time.
- Cameroon context: sickle cell high prevalence (balanced polymorphism with malaria), palm wine/burukutu traditional fermentation, need for disease-resistant plantain/cassava, vitamin A deficiency in North, oil spill bioremediation in coastal regions.

EXERCISE 6 GCE-style practice questions

1. Traditional fermentation: Describe the changes that occur when cassava tubers are soaked in water to produce fufu. Name the micro-organisms involved, the type of respiration, the main products, and how they contribute to detoxification and texture change. (8 marks)
2. Data analysis: A student investigated the effect of pH on yeast fermentation. The volume of CO_2 produced in 30 min at different pH values was: pH 3: 5 cm^3 ; pH 5: 38 cm^3 ; pH 7: 22 cm^3 ; pH 9: 8 cm^3 . (a) Plot a graph. (b) State the optimum pH. (c) Explain the low rate at pH 3 and pH 9. (d) Name two controlled variables. (7 marks)
3. Recombinant DNA: Outline the steps to produce a transgenic maize plant expressing the Bt toxin gene. Include the roles of restriction enzyme, DNA ligase, *Agrobacterium tumefaciens*, and a selectable marker. (10 marks)
4. RFLP: The β -globin gene is 1.35 kb. MstII cuts the normal allele (HbA) into 1.15 kb and 0.20 kb fragments. The sickle allele (HbS) lacks the site. Draw the expected gel bands for: (i) HbA HbA, (ii) HbA HbS, (iii) HbS HbS. Explain why the 0.20 kb band is often not seen. (6 marks)
5. Gene therapy: Distinguish between somatic and germ-line gene therapy. Explain why somatic therapy for sickle cell disease targets haematopoietic stem cells. State two risks of using lentiviral vectors. (7 marks)
6. Calculations: (a) How many PCR cycles are needed to obtain at least 10^9 molecules from 10 starting molecules? (b) A culture of *Bacillus thuringiensis* doubles every 30 min. If the fermenter is inoculated with 5×10^4 spores/mL, what is the concentration after 5 hours? (5 marks)
7. Essay: "Modern biotechnology offers solutions to Cameroon's agricultural challenges, but its adoption carries significant risks." Discuss this statement with reference to at least two specific crop examples. (15 marks)

SOLUTION Exercise 1 – Fufu fermentation

Micro-organisms: Lactic acid bacteria (*Lactobacillus*, *Leuconostoc*) and sometimes yeasts. Respiration: Anaerobic (lactic fermentation). Equation: $\text{C}_6\text{H}_{12}\text{O}_6 \longrightarrow 2 \text{CH}_3\text{CHOHCOOH} + 2 \text{ATP}$. Detoxification: Cassava contains cyanogenic glycosides (linamarin, lotaustralin). Soaking allows endogenous enzyme linamarase and microbial enzymes to hydrolyse them, releasing volatile HCN which diffuses out. Lactic acid lowers pH, inhibiting putrefactive bacteria. Texture: Pectinases and cellulases from microbes soften cell walls; starch granules swell and ferment, giving the characteristic doughy texture.

SOLUTION Exercise 2 – pH effect

(a) Graph: pH on x-axis (3.5, 7.9), CO₂ volume on y-axis. Peak at pH 5. (b) Optimum pH = 5. (c) pH 3: excess H⁺ denatures enzymes (zymase), disrupts tertiary structure. pH 9: excess OH⁻ similarly denatures enzymes; also yeast membrane transport disrupted. (d) Temperature, yeast mass, glucose concentration, volume, time, anaerobiosis.

SOLUTION Exercise 3 – Bt maize

1. Isolate cry gene from *Bacillus thuringiensis*. 2. Cut gene and plant expression vector (Ti plasmid derivative) with same restriction enzyme → sticky ends. 3. Ligate with DNA ligase → recombinant Ti plasmid. 4. Transform into *Agrobacterium tumefaciens*. 5. Co-cultivate *Agrobacterium* with maize embryogenic callus → T-DNA transfer. 6. Select transformed cells on herbicide/antibiotic medium (marker gene, e.g. bar for phosphinothricin resistance). 7. Regenerate whole plants via tissue culture. 8. Screen for cry expression (ELISA/PCR). 9. Field trials, biosafety assessment.

SOLUTION Exercise 4 – RFLP gel

(i) HbA HbA: two bands at 1.15 kb and 0.20 kb (0.20 kb may be faint/absent). (ii) HbA HbS: three bands at 1.35 kb, 1.15 kb, 0.20 kb (or two visible: 1.35, 1.15). (iii) HbS HbS: single band at 1.35 kb. Reason: 0.20 kb fragment binds little stain, runs fast, may run off gel.

SOLUTION Exercise 5 – Gene therapy

Somatic: targets body cells, not inherited, treats patient only. Germ-line: targets gametes/embryo, heritable, affects offspring, ethically prohibited. Stem cells: self-renew and differentiate into all blood lineages → lifelong supply of corrected RBCs. Lentiviral risks: insertional mutagenesis (oncogene activation), immune response to vector, generation of replication-competent virus.

SOLUTION Exercise 6 – Calculations

(a) $10 \times 2^n \geq 10^9 \Rightarrow 2^n \geq 10^8 \Rightarrow n \geq \log_2(10^8) = 8 \times \log_2(10) \approx 8 \times 3.322 = 26.57 \Rightarrow n = 27$ cycles. (b) $t = 5 \text{ h} = 300 \text{ min}$, $g = 30 \text{ min}$, $n = 300/30 = 10$ generations. $N = 5 \times 10^4 \times 2^{10} = 5 \times 10^4 \times 1024 = 5.12 \times 10^7$ spores/mL.

SOLUTION Exercise 7 – Essay outline

Intro: Define modern biotechnology, relevance to Cameroon (agriculture = 70% employment, food security challenges). Benefits: (1) Bt cotton/maize – reduce pesticide use, increase yield, farmer income; (2) Disease-resistant plantain/banana (BXW, Fusarium) – staple food security; (3) Biofortified cassava (provitamin A) – nutrition; (4) Drought-tolerant maize – climate adaptation in North. Risks: (1) Gene flow to wild relatives (wild cotton, wild banana); (2) Pest resistance evolution (refuge compliance difficult for small-holders); (3) Seed dependency/patents vs farmer-saved seeds; (4) Biodiversity loss (displacement of landraces); (5) Socio-economic: cost, access, trade implications (EU GMO regulations). Regulation: Cameroon Biosafety Law (2003), Cartagena Protocol, case-by-case risk assessment, labelling, public participation. Conclusion: Potential is real but requires strong regulatory capacity, farmer involvement, coexistence measures, and integration with agroecology. Not a silver bullet.

DID YOU KNOW? Biotechnology in Cameroon – did you know?

- The Institute of Agricultural Research for Development (IRAD) has developed improved plantain varieties (e.g. FHIA hybrids) using conventional breeding and tissue culture, and is researching GM banana for BXW resistance.
- Centre Pasteur du Cameroun and Centre de Recherche sur les Maladies Infectieuses (CREMI) use PCR for diagnosis of HIV, tuberculosis, malaria, and COVID-19 – modern biotechnology in public health.
- Sickle cell screening: Neonatal screening using HPLC or electrophoresis is being piloted in Yaoundé and Douala; RFLP/PCR prenatal diagnosis is available in specialised centres.
- Bioremediation: After oil spills in the Rio del Rey and Kribi areas, *Pseudomonas* and *Bacillus* strains are tested for hydrocarbon degradation.
- Traditional biotechnology economy: Palm wine, bilibili (millet beer), fufu, eru fermentation support thousands of rural livelihoods – a cultural heritage and economic asset.

WARNING Common GCE traps in Biotechnology

- Confusing traditional (no genetic alteration) vs modern (recombinant DNA, gene editing) biotechnology.
- Forgetting that fermentation yields 2 ATP (not 36–38) and is anaerobic.
- Writing “yeast is killed by cold” – cold inactivates/slows enzymes; heat denatures

them.

- Omitting the enzyme/tool at each recombinant DNA step (restriction enzyme, ligase, reverse transcriptase, Taq polymerase).
- Misinterpreting gel bands: smallest fragment runs furthest (not highest).
- Confusing carrier (heterozygous, healthy) with affected (homozygous recessive, diseased).
- Not distinguishing somatic (non-heritable) vs germ-line (heritable) gene therapy.
- Calculation errors: not converting time units, using $2t$ instead of $2^{t/g}$.
- One-sided essays: must give balanced arguments (pros + cons) + justified conclusion.

10 EXERCISES

10.1 I check my knowledge

EXERCISE 7 MCQ: Foundations of biotechnology

For each question, choose the single correct answer.

1. The production of palm wine from raffia sap in the South–West Region of Cameroon is an example of:
 - (a) recombinant DNA technology
 - (b) traditional biotechnology
 - (c) gene therapy
 - (d) genome editing
2. During alcoholic fermentation by *Saccharomyces cerevisiae*, the role of NAD^+ is to:
 - (a) accept electrons during glycolysis and be regenerated in the reduction of acetaldehyde
 - (b) serve as the final electron acceptor of the respiratory chain
 - (c) fix carbon dioxide into organic matter
 - (d) hydrolyse sucrose into glucose and fructose
3. A restriction endonuclease such as EcoRI:
 - (a) joins two DNA fragments by forming phosphodiester bonds
 - (b) cuts DNA at a specific recognition sequence
 - (c) copies DNA into messenger RNA
 - (d) amplifies a DNA fragment in vitro
4. In gene therapy, a vector is:
 - (a) a defective gene to be replaced
 - (b) a carrier used to deliver a functional gene into target cells
 - (c) an enzyme that cuts the chromosome
 - (d) a protein that inhibits transcription

EXERCISE 8 True or false, with justification

State whether each statement is true or false, and justify your answer in one or two sentences.

1. Fermentation is an aerobic process that completely oxidises glucose to carbon dioxide and water.
2. The bacterium *Agrobacterium tumefaciens* can be used as a natural vector to transfer genes into plant cells.
3. CRISPR–Cas9 genome editing always modifies every cell of the organism with absolute specificity and no off-target effects.
4. Golden Rice has been engineered to produce β -carotene, a precursor of vitamin A, in its grains.
5. Recombinant human insulin is extracted directly from the pancreas of pigs.


EXERCISE 9 Key definitions

Define each of the following terms precisely, in two or three sentences each, and give one concrete example for each:

1. Traditional biotechnology
2. Recombinant DNA
3. Transgenic organism (GMO)
4. Somatic gene therapy
5. Bioremediation

EXERCISE 10 Direct calculation: theoretical ethanol yield

Saccharomyces cerevisiae ferments glucose according to the overall equation:



A brewer ferments a wort containing 100 g of glucose. (Molar masses: glucose = 180 g mol^{-1} ; ethanol = 46 g mol^{-1} .)

1. Calculate the number of moles of glucose fermented.
2. Deduce the theoretical mass of ethanol produced.
3. If the wort volume is 1 L, express the theoretical ethanol yield in g L^{-1} .

4. The brewer actually recovers 40 g of ethanol. Calculate the percentage yield and suggest one biological reason why the real yield is lower than the theoretical yield.

10.2 I practise

EXERCISE 11 Restriction mapping and gel analysis

A circular plasmid of 5.4 kb carries a 1.2 kb insert. The plasmid has one EcoRI site and one HindIII site, located 2.0 kb apart on the map, both outside the insert.

1. Predict the number and sizes of the fragments obtained after:
 - (a) single digestion with EcoRI;
 - (b) double digestion with EcoRI and HindIII.
2. Draw the expected banding pattern on a 1% agarose gel, indicating the position of each band relative to a molecular weight marker.
3. Explain how you would verify, from this gel, that the insert is present.

EXERCISE 12 Steps of a recombinant DNA experiment

A Cameroonian research team wants to produce human insulin in *Escherichia coli*. Rearrange the following steps in the correct chronological order and justify the role of each step:

1. Transform competent bacteria and select recombinant colonies on a medium containing an antibiotic.
2. Isolate the human insulin gene (or synthesise its cDNA from pancreatic mRNA).
3. Extract the plasmids and verify the insert by restriction digestion.
4. Cut the plasmid vector and the insert with the same restriction enzyme.
5. Culture the transformed bacteria in a fermenter and purify the insulin produced.
6. Ligate the insert into the plasmid using DNA ligase.

EXERCISE 13 Fermentation monitoring in a local brewery

During the production of a traditional sorghum beer, a student measures the sugar con-

centration and the ethanol concentration of the must over five days:

Day	0	1	2	3	4
Sugar (g L^{-1})	120	95	60	25	8
Ethanol (g L^{-1})	0	12	30	48	55

1. Plot the two curves on the same graph (time on the x-axis).
2. Describe and explain the shape of each curve in relation to yeast metabolism.
3. Explain why ethanol production slows down after day 3, giving two possible reasons.
4. State two precautions a local producer should take to avoid contamination of the brew by unwanted microorganisms.

EXERCISE 14 Biotechnology for support and locomotion

Tissue engineering is increasingly used to repair damaged cartilage and bone.

1. Explain the principle of producing a cartilage implant from a patient's own stem cells cultured on a biodegradable scaffold.
2. State two advantages of using the patient's own cells rather than a donor's cells, in relation to the immune system.
3. Name one biotechnological product (other than cell therapy) used to support bone repair, and briefly describe how it is produced.

10.3 I prepare for the GCE

EXERCISE 15 CRISPR–Cas9 therapy for sickle cell disease

(20 marks)

Sickle cell disease is frequent in Cameroon. A new treatment uses CRISPR–Cas9 to switch off the BCL11A gene in the patient's haematopoietic stem cells, which reactivates the production of fetal haemoglobin (HbF).

1. Describe the molecular basis of sickle cell disease, indicating the gene concerned and the type of mutation involved. (4 marks)
2. Explain step by step how the CRISPR–Cas9 system can be guided to the BCL11A gene and what happens after the cut. (6 marks)
3. Explain why reactivating HbF compensates for the defective adult haemoglobin. (3 marks)

4. Identify two risks associated with this approach, including off-target effects, and propose one molecular follow-up strategy to monitor treated patients. (4 marks)
5. Discuss one ethical or social issue raised by the availability of such an expensive therapy in Cameroon. (3 marks)

EXERCISE 16 Golden Rice versus vitamin A supplementation**(20 marks)**

Vitamin A deficiency causes blindness and increases child mortality in many tropical countries. Golden Rice is a transgenic rice engineered to accumulate β -carotene in the grain.

1. Outline the biotechnological steps used to obtain Golden Rice, naming the type of vector commonly used for monocot or dicot transformation. (5 marks)
2. Compare Golden Rice and periodic vitamin A capsule supplementation under the following headings: nutritional effectiveness, agronomic constraints, economic cost for families and the State, and ethical/environmental concerns. Present your comparison in a table. (10 marks)
3. As an adviser to the Cameroonian Ministry of Public Health, write a short argued paragraph (8–10 lines) recommending a strategy that combines both approaches for rural areas. (5 marks)

EXERCISE 17 Risk assessment for a GM biosurfactant bacterium**(20 marks)**

A company proposes to clean up an oil-polluted site in the Littoral Region of Cameroon by releasing a genetically modified bacterium that produces a biosurfactant degrading petroleum hydrocarbons.

1. Explain the principle of bioremediation and the role of the biosurfactant in making hydrocarbons accessible to the bacteria. (4 marks)
2. Draw up a risk assessment sheet identifying: (i) two risks for the local ecosystems (including horizontal gene transfer), (ii) two risks for human populations living near the site, and (iii) two containment or monitoring measures for each category of risk. Present your answer as a structured table. (10 marks)
3. Propose an experimental protocol, carried out before any field release, to test the survival and spread of the GM bacterium in soil microcosms. (4 marks)
4. State one argument in favour of, and one against, authorising this release, and conclude with a justified personal position. (2 marks)

11 SOLUTIONS TO THE EXERCISES

SOLUTION Exercise 1 — MCQ: Foundations of biotechnology

1. Correct answer: (b) traditional biotechnology. Palm wine production exploits the natural fermentation of raffia sap sugars by wild yeasts, an ancestral practice that does not involve any manipulation of DNA. It therefore belongs to **traditional biotechnology**, unlike recombinant DNA technology, gene therapy or genome editing, which all require direct modification of genetic material.
2. Correct answer: (a). During glycolysis, NAD^+ accepts electrons (and H^+) from glyceraldehyde-3-phosphate and is reduced to NADH. In alcoholic fermentation, NADH is re-oxidised to NAD^+ when acetaldehyde is reduced to ethanol, allowing glycolysis to continue in the absence of oxygen. Oxygen, not NAD^+ , is the final electron acceptor of the respiratory chain.
3. Correct answer: (b). A restriction endonuclease such as EcoRI recognises a specific palindromic sequence (5'-GAATTC-3') and cuts the DNA backbone at that site. Joining fragments is the role of DNA ligase; copying DNA into mRNA is transcription; in vitro amplification is PCR.
4. Correct answer: (b). In gene therapy, a **vector** (modified virus, liposome, plasmid) is a carrier used to deliver a functional copy of a gene into the patient's target cells. It is neither the defective gene, nor a cutting enzyme, nor a transcription inhibitor.

SOLUTION Exercise 2 — True or false, with justification

1. False. Fermentation is an anaerobic process (it occurs without oxygen) and is an incomplete oxidation of glucose: the end products (ethanol, lactate) still contain most of the chemical energy. Complete oxidation to CO_2 and H_2O is aerobic respiration.
2. True. *Agrobacterium tumefaciens* naturally transfers a fragment of its Ti plasmid (T-DNA) into the genome of plant cells, causing crown gall disease. Biotechnologists disarm the Ti plasmid and replace the tumour genes with a gene of interest, making it a natural vector for plant genetic engineering (mainly dicots).
3. False. CRISPR-Cas9 is highly specific but not absolute: **off-target** cuts can occur at sequences similar to the target, and editing efficiency varies between cells (mosaicism), so not every cell is necessarily modified. This is why edited cells must be screened and sequenced.
4. True. Golden Rice was engineered by introducing genes (e.g. psy from daffodil/maize

and crtI from a bacterium) so that the endospore accumulates β -carotene (provitamin A), which the human body converts into vitamin A.

5. False. Recombinant human insulin is produced by bacteria (*E. coli*) or yeast carrying the human insulin gene inserted into a plasmid; it is not extracted from pig pancreas. Animal (porcine/bovine) insulin was the older, pre-recombinant source.

SOLUTION Exercise 3 — Key definitions

1. Traditional biotechnology: the use of living organisms or their enzymes, based on empirical ancestral know-how, to produce or transform food and other goods, without any direct manipulation of DNA. Example: fermentation of raffia sap into palm wine, or of corn/cassava dough into “nkui”-type products or sorghum beer in Cameroon.
2. Recombinant DNA: a DNA molecule artificially constructed in vitro by joining DNA fragments from different sources (e.g. a human gene inserted into a bacterial plasmid) using restriction enzymes and DNA ligase. Example: a plasmid carrying the human insulin gene used to transform *E. coli*.
3. Transgenic organism (GMO): an organism whose genome has been stably modified by the introduction of one or more genes from another species (a transgene), which is expressed and transmitted to offspring. Example: Bt maize carrying a *Bacillus thuringiensis* toxin gene, or Golden Rice.
4. Somatic gene therapy: the introduction of a functional gene into the somatic (body, non-reproductive) cells of a patient to correct or compensate for a genetic defect; the modification is not transmitted to descendants. Example: delivery of a functional CFTR gene to lung cells of a cystic fibrosis patient using a viral vector.
5. Bioremediation: the use of living organisms (bacteria, fungi, plants) to degrade, transform or immobilise pollutants in a contaminated environment (soil, water). Example: hydrocarbon-degrading bacteria used to clean up oil-polluted soils in the Littoral Region.

SOLUTION Exercise 4 — Direct calculation: theoretical ethanol yield

1. Number of moles of glucose:

$$n(\text{glucose}) = \frac{m}{M} = \frac{100 \text{ g}}{180 \text{ g mol}^{-1}} \approx 0.556 \text{ mol.}$$

2. From the equation $\text{C}_6\text{H}_{12}\text{O}_6 \longrightarrow 2 \text{C}_2\text{H}_5\text{OH} + 2 \text{CO}_2$, one mole of glucose gives two

moles of ethanol:

$$n(\text{ethanol}) = 2 \times 0.556 = 1.111 \text{ mol},$$

$$m(\text{ethanol}) = n \times M = 1.111 \times 46 \approx 51.1 \text{ g}.$$

3. For a wort volume of 1 L, the theoretical yield is:

$$\frac{51.1 \text{ g}}{1 \text{ L}} = 51.1 \text{ g L}^{-1}.$$

4. Percentage yield:

$$\text{yield} = \frac{40}{51.1} \times 100 \approx 78.3 \text{ \%}.$$


Biological reason for the lower real yield (any one): part of the glucose is used by the yeast for its own growth and respiration (biomass, ATP production) rather than for fermentation; some ethanol evaporates or remains dissolved in the lees; fermentation may stop before all sugar is consumed because ethanol becomes toxic to the yeast.

SOLUTION Exercise 5 — Restriction mapping and gel analysis

1. Predicted fragments:

- (a) Single digestion with EcoRI: a circular plasmid cut at one site is simply linearised, giving one linear fragment of 5.4 kb.
- (b) Double digestion EcoRI + HindIII: two cuts on a circular molecule give two fragments: one of 2.0 kb (the distance between the two sites) and one of $5.4 - 2.0 = 3.4$ kb.

2. Expected banding pattern: on a 1% agarose gel, small fragments migrate faster (further towards the anode, bottom of the gel).



Simulated agarose gel: marker (left), EcoRI single digest (middle), EcoRI+HindIII double digest (right).

3. Verifying the insert is present: the undigested/linearised recombinant plasmid is 5.4 kb, whereas the empty vector would be only $5.4 - 1.2 = 4.2$ kb; the size shift of 1.2 kb relative to the empty-vector control shows the insert is present. To confirm directly, one can digest with an enzyme that cuts on each side of the insert to release a 1.2 kb band, or use the 1.2 kb insert as a probe in a Southern blot.

SOLUTION Exercise 6 — Steps of a recombinant DNA experiment


Correct chronological order: (b) → (d) → (f) → (a) → (c) → (e).

1. (b) Isolate the human insulin gene (or synthesise its cDNA from pancreatic mRNA using reverse transcriptase). Role: obtain the gene of interest; cDNA is preferred because bacteria cannot remove introns.
2. (d) Cut the plasmid vector and the insert with the same restriction enzyme. Role: generate compatible (complementary sticky) ends so that the insert can be joined to the vector.
3. (f) Ligate the insert into the plasmid using DNA ligase. Role: DNA ligase forms phosphodiester bonds, sealing the insert into the vector to create a **recombinant plasmid**.
4. (a) Transform competent bacteria and select recombinant colonies on an antibiotic medium. Role: introduce the plasmid into *E. coli*; the plasmid's antibiotic-resistance marker allows only transformed bacteria to grow.
5. (c) Extract the plasmids and verify the insert by restriction digestion. Role: quality control — check that the selected colonies carry the correct insert (expected fragment sizes on a gel).
6. (e) Culture the transformed bacteria in a fermenter and purify the insulin. Role: large-

scale production, then extraction and purification of the recombinant human insulin for medical use.

SOLUTION Exercise 7 — Fermentation monitoring in a local brewery

1. Graph:



Sugar consumption and ethanol production during alcoholic fermentation.

- Shape of the curves: the sugar concentration decreases continuously because the yeasts consume sugars as their respiratory/fermentative substrate; the ethanol concentration increases because ethanol is the end product of alcoholic fermentation. The two curves are roughly mirror images: ethanol is produced as sugar is consumed, according to $\text{C}_6\text{H}_{12}\text{O}_6 \longrightarrow 2 \text{C}_2\text{H}_5\text{OH} + 2 \text{CO}_2$.
- Slowing of ethanol production after day 3 (two reasons):
 - the sugar substrate is becoming scarce (25 g L⁻¹ then 8 g L⁻¹), so fermentation slows down by substrate limitation;
 - ethanol accumulates and becomes toxic to the yeasts (it inhibits their enzymes and eventually kills the cells), reducing the fermenting population.
- Precautions against contamination (any two): wash and, where possible, boil/sterilise the vessels and utensils; cover the brew to keep out dust, insects (e.g. vinegar flies that bring acetic acid bacteria) and airborne microbes; use clean water; inoculate with a healthy starter from a previous good batch rather than relying on chance contamination.

SOLUTION Exercise 8 — Biotechnology for support and locomotion

1. Principle of the cartilage implant: stem cells (e.g. mesenchymal stem cells from the patient's bone marrow) are harvested, multiplied in vitro, then seeded onto a **biodegradable scaffold** (e.g. collagen or polylactic acid) that reproduces the shape of the damaged cartilage. In a bioreactor, with appropriate growth factors, the cells differentiate into chondrocytes and secrete extracellular matrix. The scaffold progressively degrades while the new tissue forms, and the construct is then implanted at the lesion site.
2. Advantages of autologous (own) cells:
 - No immune rejection: the cells carry the patient's own HLA (self) antigens, so the immune system does not recognise them as foreign;
 - No need for immunosuppressive treatment and no risk of transmitting donor-borne infections (e.g. viral contamination of the graft).
3. Another biotechnological product for bone repair: recombinant Bone Morphogenetic Protein (rhBMP-2), a growth factor that stimulates osteoblast differentiation and bone formation. It is produced by recombinant DNA technology: the human BMP gene is inserted into host cells (e.g. mammalian cell cultures), which synthesise the protein, later purified and applied on a collagen sponge at the fracture site.

SOLUTION Exercise 9 — CRISPR-Cas9 therapy for sickle cell disease (20 marks)

1. Molecular basis (4 marks): sickle cell disease is caused by a point mutation (base substitution: A → T) in the gene coding for the β -chain of haemoglobin (HBB gene, chromosome 11). The codon GAG (glutamic acid) becomes GTG (valine) at position 6 of the β -globin chain. The abnormal haemoglobin HbS polymerises when deoxygenated, deforming red blood cells into sickle shapes that block capillaries and are destroyed prematurely (chronic anaemia, painful crises). Examiner expects: gene named, type of mutation, amino-acid substitution, one phenotypic consequence.
2. CRISPR-Cas9 mechanism (6 marks):
 - (a) a guide RNA (sgRNA) is synthesised, complementary to a specific sequence of the BCL11A gene;
 - (b) the sgRNA associates with the Cas9 nuclease and guides it, by complementary base pairing, to the target site (next to a PAM sequence);
 - (c) Cas9 cuts both DNA strands at that precise site;
 - (d) the cell repairs the break by non-homologous end joining (NHEJ), which introduces small insertions/deletions that inactivate BCL11A;

- (e) the edited haematopoietic stem cells are reinfused into the patient, where they produce red blood cells expressing fetal haemoglobin.
3. Why HbF compensates (3 marks): fetal haemoglobin (HbF, $\alpha_2\gamma_2$) does not contain β -globin, so it is unaffected by the sickle mutation; it transports oxygen normally and dilutes/inhibits the polymerisation of HbS, preventing sickling. Since BCL11A normally represses the γ -globin gene after birth, switching it off reactivates HbF production.
 4. Risks and follow-up (4 marks): risks — off-target cuts elsewhere in the genome (possible mutations, cancer risk); incomplete editing (mosaicism) or genotoxicity of the conditioning chemotherapy; long-term effects of BCL11A inactivation unknown. Follow-up: regular molecular monitoring by PCR and DNA sequencing of the target locus and predicted off-target sites, plus measurement of HbF levels and blood counts.
 5. Ethical/social issue (3 marks): the therapy is extremely expensive and requires highly specialised centres, so very few Cameroonian patients could access it, raising a problem of equity of access to care; resources might alternatively fund prevention (premarital screening, genetic counselling). Any reasoned, balanced argument is credited.

SOLUTION**Exercise 10 — Golden Rice versus vitamin A supplementation (20 marks)**

1. Obtaining Golden Rice (5 marks): (i) identify and isolate the genes of the β -carotene pathway: psy (phytoene synthase, from daffodil or maize) and crtI (bacterial carotene desaturase); (ii) place them under an endosperm-specific promoter so β -carotene accumulates in the grain; (iii) insert them into a suitable vector — the disarmed Ti plasmid of *Agrobacterium tumefaciens* (standard for dicots, and adapted strains for monocots such as rice) or use biolistics (gene gun) for monocots; (iv) transform rice cells, regenerate whole plants by tissue culture on selective medium; (v) verify insertion and expression (PCR, carotenoid assays) and select stable lines.
2. Comparison (10 marks):

Criterion	Golden Rice	Vitamin A capsules
Nutritional effectiveness	Continuous daily supply of provitamin A with the staple food; dose depends on consumption and variety	Precise, high dose but only 1–2 times per year; risk of missed children
Agronomic constraints	Requires adapted varieties, seed distribution, farmer adoption; yields must equal local varieties	None: no change to farming practices
Economic cost	Low recurrent cost for families once seeds available; high initial research/seed-system cost for the State	Recurrent cost of purchase, storage and distribution campaigns, often donor-dependent
Ethical/environmental concerns	GMO acceptance, possible gene flow to wild relatives, dependence on seed suppliers	Few ethical concerns; risk of overdose rare; no environmental impact

3. Argued recommendation (5 marks) — model paragraph: I recommend a combined strategy. In the short term, capsule supplementation campaigns must continue, because they reach children immediately and require no change in farming. In parallel, the State should support field trials of biofortified varieties (Golden Rice or, more realistically for Cameroon, biofortified orange-fleshed sweet potato and vitamin-A maize/cassava) adapted to local conditions, with free seed distribution to rural farmers. Biofortification then provides a sustainable, low-cost daily supply integrated into the diet, while capsules cover emergencies and populations not yet reached. Public information campaigns are essential to ensure acceptance of biofortified foods. This progressive combination guarantees both immediate coverage and long-term food autonomy.

SOLUTION Exercise 11 — Risk assessment for a GM biosurfactant bacterium (20 marks)

1. Principle (4 marks): **bioremediation** uses microorganisms to degrade pollutants into harmless products (CO_2 , H_2O , biomass). Petroleum hydrocarbons are hydrophobic


and poorly soluble in water, which limits their uptake by bacteria. The biosurfactant is an amphiphilic molecule that emulsifies the hydrocarbons, increasing their dispersion and bioavailability, so the bacteria can absorb and oxidise them efficiently.

2. Risk assessment sheet (10 marks):

Category	Risks	Containment / monitoring measures
Ecosystems	(i) Horizontal gene transfer of the biosurfactant/antibiotic-resistance gene to native soil bacteria; (ii) disruption of indigenous microbial communities and soil food chains	(i) Use a strain without antibiotic-resistance markers, with a “suicide gene” limiting survival; (ii) confine treatment to the polluted plot with physical barriers and monitor soil microbiota by regular sampling
Human populations	(i) Allergenicity/toxicity of the bacterium or its metabolites for workers and residents; (ii) spread of the GMO to water sources used for drinking or fishing	(i) Toxicity tests before release and protective equipment for workers; (ii) buffer zones around wells/streams and regular microbiological testing of water

3. Microcosm protocol (4 marks): fill sealed soil columns (microcosms) with non-sterile local soil, some polluted with oil and some not; inoculate half with the GM bacterium (marked, e.g. with a fluorescent gfp marker) and keep uninoculated controls; incubate under field-like temperature and humidity; sample at regular intervals (days 0, 7, 14, 30, 60) and quantify the GM strain by selective plating and PCR of its marker gene; measure hydrocarbon degradation and check for gene transfer to native bacteria; all manipulations in a contained laboratory (biosafety level 2) with autoclaving of all waste.
4. Arguments and position (2 marks): For: the method is cheaper and more ecological than excavation/incineration and can restore farmland and fisheries. Against: once released, a GMO cannot be recalled, and long-term ecological effects are uncertain. Personal position (example): authorisation only after successful microcosm and greenhouse trials, with a marker-free, survival-limited strain and a strict monitoring plan — a precautionary but not prohibitive stance.

12 SUMMARY SHEET



Mind map of Chapter BIO07: Biotechnology

FICHE BILAN L'essentiel à connaître pour l'examen

Key Definitions

- **Biotechnology**: Use of living organisms or their components to make or modify products for specific use.
- **Traditional biotechnology**: Ancient processes like fermentation (e.g., kwacha production, palm wine, cassava fermentation) relying on natural microbes.
- **Recombinant DNA technology**: Artificial combination of DNA from different sources to create transgenic organisms.
- **Gene therapy**: Introduction of functional genes into patient cells to treat genetic disorders (e.g., sickle cell disease trials).
- **Genome editing**: Precise alteration of DNA sequences using tools like CRISPR-Cas9.
- **Homeostasis in biotech**: Use of biotechnological devices (biosensors, artificial organs) to maintain internal stability.

Laws, Formulas & Principles to Know

- **Chargaff's rules** and **Watson-Crick base pairing** underpin DNA manipulation.
- **Restriction enzyme kinetics**: $v = \frac{V_{\max}[S]}{K_m + [S]}$ (Michaelis-Menten) for predicting digestion

efficiency.

- **PCR amplification:** $N = N_0 \times 2^n$ where n = number of cycles.
- **Plasmid copy number** and **selection marker** principles for cloning vectors.
- **Hardy–Weinberg equilibrium** ($p^2 + 2pq + q^2 = 1$) for population genetics in GMO risk assessment.

Methods (Step-by-Step)

- Fermentation monitoring: (1) Inoculate sterile substrate; (2) Control pH, temperature, anaerobiosis; (3) Measure CO₂ evolution or ethanol yield; (4) Harvest product.
- Recombinant plasmid construction: (1) Choose vector & insert; (2) Digest both with compatible restriction enzymes; (3) Ligate with DNA ligase; (4) Transform competent *E. coli*; (5) Screen colonies (blue–white, PCR, sequencing).
- CRISPR–Cas9 editing: (1) Design sgRNA targeting gene; (2) Clone sgRNA into expression plasmid; (3) Deliver Cas9–sgRNA complex (electroporation, viral vector); (4) Verify edit by sequencing; (5) Assess off–target effects.
- Biosensor calibration: (1) Immobilise enzyme/antibody; (2) Record signal vs known analyte concentrations; (3) Plot calibration curve; (4) Use for real–time homeostasis monitoring (e.g., glucose sensor).

Common Mistakes (Traps)

- Confusing traditional (spontaneous, mixed cultures) with modern (defined strains, controlled bioreactors) fermentation.
- Forgetting that restriction enzymes cut at specific palindromic sites; using incompatible ends prevents ligation.
- Assuming PCR doubles DNA every cycle indefinitely — plateau phase occurs due to reagent depletion.
- Overlooking ethical & regulatory frameworks (Cameroon’s biosafety law, Cartagena Protocol) when discussing GMO release.
- Misapplying Hardy–Weinberg: it assumes random mating, no selection, large population — rarely true for transgenic crops in field.

GCE Advanced Level Tips

- Definitions: Write concise, exam–style definitions; mention at least one Cameroonian example (e.g., foléré fermentation, sickle cell gene therapy trials in Yaoundé).
- Diagrams: Draw labelled plasmid map (origin, selectable marker, MCS, insert) and CRISPR mechanism (Cas9, sgRNA, PAM, target DNA). Use TikZ–style sketches in answers.
- Calculations: Show all steps for PCR yield, restriction fragment sizes, allele frequencies. Units must be included.

- Essay questions: Structure: Introduction → Traditional vs Modern → Recombinant DNA steps → Applications (health, agriculture, environment) → Ethical/safety concerns → Conclusion.
- Practicals: Know how to interpret gel electrophoresis bands, fermentation graphs (lag, log, stationary phases), and biosensor calibration curves.

**Signed • OnBuch+**ONBUCH+ TEACHING TEAM
FOLLOWS THE OFFICIAL MINESEC SYLLABUS**OnBuch+**