

BIOLOGY • UPPER SIXTH

UPPER SIXTH — BIOLOGY

CHAPTER 8

AGRICULTURE

SUGGESTED TIME: 5 PERIODS

IN THIS CHAPTER

This chapter explores the scientific foundations of agriculture: the types and properties of soils, the nutritional requirements of plants and animals, and the techniques used to cultivate crops and raise livestock. It equips learners with the knowledge needed to improve agricultural productivity and to tackle GCE Advanced Level questions on agriculture with confidence.

- By the end of this chapter I can describe the main types of soils and relate their properties to agricultural use.
- By the end of this chapter I can explain soil formation, soil profile and the factors affecting soil fertility.
- By the end of this chapter I can state the mineral elements required by plants and describe deficiency symptoms and methods of correcting them.
- By the end of this chapter I can distinguish between food and non-food crops and describe the cultivation techniques of selected Cameroonian crops.
- By the end of this chapter I can describe the nutritional requirements of farm animals and the classes of feedstuffs.
- By the end of this chapter I can explain the principles of ration formulation and feeding regimes for livestock.



DOWNLOAD ON- BUCH

Free • Google Play
AI tutor, past papers,
results



GO ONBUCH+

All signed courses, unlimited AI
tutor, PDF sheets — OnBuch+
tab in the app

THIS SIGNED COURSE CONTAINS

COURSE

complete, illustrated,
step by step

METHODS

and worked examples

EXERCISES

exam style + solutions

SUMMARY SHEET

the essentials to
remember



Signed • OnBuch+

ONBUCH+ TEACHING TEAM
FOLLOWS THE OFFICIAL MINESEC SYLLABUS

OnBuch+

CONTENTS

- | | |
|--|--------------------------------|
| 1. Soils: Types, Formation and Fertility | 5. Integration activity |
| 2. Plant Nutritional Requirements | 6. Methods and worked examples |
| 3. Cultivation Techniques of Food and Non-Food Crops | 7. Exercises |
| 4. Animal Nutrition and Animal Production | 8. Solutions to the exercises |
| | 9. Summary sheet |

CHAPTER OBJECTIVES

- By the end of this chapter I can describe the main types of soils and relate their properties to agricultural use.
- By the end of this chapter I can explain soil formation, soil profile and the factors affecting soil fertility.
- By the end of this chapter I can state the mineral elements required by plants and describe deficiency symptoms and methods of correcting them.
- By the end of this chapter I can distinguish between food and non-food crops and describe the cultivation techniques of selected Cameroonian crops.
- By the end of this chapter I can describe the nutritional requirements of farm animals and the classes of feedstuffs.
- By the end of this chapter I can explain the principles of ration formulation and feeding regimes for livestock.
- By the end of this chapter I can describe the systems and techniques of animal production, including housing, breeding, health and record keeping.
- By the end of this chapter I can apply agricultural knowledge to solve practical farm problems and answer GCE-style structured and essay questions.

PREREQUISITES

- Plant nutrition and photosynthesis (Lower Sixth): autotrophic nutrition, mineral absorption by roots

- Basic ecology: ecosystems, nutrient cycling (nitrogen and carbon cycles)
- Chemical composition of living matter: carbohydrates, proteins, lipids, vitamins and minerals
- Digestion in mammals and the structure of the alimentary canal (including ruminants)
- Reproduction in plants and animals (Lower Sixth)

1 SOILS: TYPES, FORMATION AND FERTILITY

In the village of Bafut, two farmers plant maize on neighbouring plots during the same week; at harvest, one fills ten bags while the other barely fills three, despite using the same seed variety. Meanwhile, a poultry keeper in Bamenda loses half her broilers before week six, while her neighbour's birds reach market weight on schedule. What hidden factors in the soil, the feed and the farming methods explain such dramatic differences? This chapter takes you into the science of agriculture — understanding soils, feeding crops and animals correctly, and mastering production techniques — so that farming in Cameroon becomes a predictable, profitable science rather than a gamble. We begin at the very foundation of it all: the soil beneath our feet.

1.1 What is Soil and Why Does it Matter?

Walk into any farm around Buea, Ngaoundéré or Bafoussam and scoop up a handful of earth. What you hold is not simply “dirt”. It is a living, complex mixture of rock fragments, decaying organisms, water, air and countless microbes. Every crop a farmer plants — maize, cassava, cocoa, plantain — depends on this thin layer of material for anchorage, water and mineral nutrients.

DEFINITION Soil

Soil is the uppermost layer of the Earth's crust, formed by the weathering of rocks and the decomposition of organic matter, consisting of mineral particles, organic matter (humus), water, air and living organisms, and capable of supporting plant growth.

Soil is the **basis of agriculture** because it performs four essential functions for the farmer:

- **Anchorage:** roots grip soil particles, holding the plant upright against wind and rain.
- **Water supply:** soil stores rain and irrigation water and releases it to roots.
- **Mineral nutrition:** soil supplies essential elements such as nitrogen, phosphorus and potassium.
- **Habitat:** soil hosts organisms (earthworms, bacteria, fungi) that recycle nutrients and maintain fertility.

DID YOU KNOW? Soil is a non-renewable resource on a human timescale

It can take several hundred years for just a few centimetres of topsoil to form naturally, yet a single season of careless farming on a steep slope in the North West Region can wash it away. This is why soil conservation is a matter of national food security.

1.2 Soil Formation (Pedogenesis)

Soil does not appear from nowhere. It forms slowly from **parent rock** through a process called

pedogenesis, driven by weathering and the action of living organisms.

Weathering of parent rock

Weathering is the breakdown of solid rock into smaller and smaller particles. Three types act together:

- **Physical (mechanical) weathering:** rock is broken into fragments without any change in chemical composition. In Cameroon this includes the alternate heating and cooling of rocks under the hot sun of the Far North (thermal expansion and contraction), and the wedging action of water and growing roots in rock cracks.
- **Chemical weathering:** the minerals in rock are chemically changed. Rainwater, slightly acidified by dissolved carbon dioxide, dissolves limestone (solution); oxygen oxidises iron minerals, giving tropical soils their red colour; water reacts with feldspar minerals to form clay (hydrolysis). Chemical weathering is fastest in hot, wet climates — which is why tropical soils in the South of Cameroon are deeply weathered.
- **Biological weathering:** living organisms break down rock. Plant roots exert pressure in cracks; burrowing animals mix and fragment material; lichens and bacteria secrete acids that dissolve minerals.


Factors controlling soil formation

The nature of the soil produced depends on five factors, easily remembered as **CLORPT**:

- **Climate:** rainfall and temperature control the speed of weathering and leaching.
- **Organisms:** vegetation, microbes and animals add organic matter and mix the soil.
- **Relief (topography):** steep slopes lose soil by erosion; valleys accumulate deep soils.
- **Parent material:** the original rock determines mineral content (e.g. basalt gives fertile volcanic soils).
- **Time:** the longer the process, the deeper and more developed the soil.

1.3 The Soil Profile

If you dig a pit about one metre deep on undisturbed land, the wall of the pit reveals distinct horizontal layers called **horizons**. The vertical sequence of horizons is called the **soil profile**.



A typical soil profile showing the horizons O, A, B, C and the bedrock.

Horizon	Characteristics
O (organic)	Surface layer of dead leaves and decomposing remains; source of humus.
A (topsoil)	Dark, rich in humus mixed with mineral particles; most biological activity; most roots found here; loses minerals by leaching (eluviation).
B (subsoil)	Lighter in colour; accumulates clay, iron and aluminium oxides washed down from A (illuviation); fewer organisms.
C	Partly weathered parent rock fragments; little organic matter.
Bedrock (R)	Solid, unweathered rock from which the soil formed.

1.4 Components of Soil

A fertile soil is a four-part mixture. By volume, an ideal mineral soil contains roughly 45% mineral particles, 5% organic matter, 25% water and 25% air.


- **Mineral particles:** weathered rock fragments (sand, silt, clay) supplying mineral nutrients.
- **Organic matter (humus):** dark, decomposed plant and animal remains. **Humus** improves structure, holds water and releases nutrients slowly.
- **Soil water:** a dilute solution of mineral salts; the medium through which roots absorb nutrients.
- **Soil air:** occupies the pore spaces; supplies oxygen for root respiration and for aerobic microbes.
- **Soil organisms:** earthworms, termites, bacteria and fungi decompose organic matter, fix nitrogen and aerate the soil.

1.5 Soil Texture

DEFINITION Soil texture

Soil texture is the relative proportion by mass of the different-sized mineral particles — sand (0.05–2 mm), silt (0.002–0.05 mm) and clay (less than 0.002 mm) — present in a soil.

Texture controls almost everything a farmer cares about: how fast water drains, how much water is stored, how easily roots penetrate, and how well nutrients are held. The **soil texture triangle** is used to classify any soil once the percentages of sand, silt and clay are known.



The soil texture triangle. Reading the percentages of sand, silt and clay locates the soil class; the shaded central zone is loam.

METHOD Determining soil texture by the sedimentation (jar) method

1. Collect a sample of dry topsoil, remove stones and crush lumps.
2. Half-fill a tall transparent jar (or measuring cylinder) with the soil, then add water and a spoonful of salt or dispersing agent.
3. Shake vigorously for about two minutes to separate all particles.
4. Stand the jar undisturbed. Sand settles within about one minute, silt within a few hours, and clay may take a day or more, forming distinct layers.

5. Measure the thickness of each layer and calculate its percentage of the total settled depth.
6. Plot the three percentages on the texture triangle to name the soil class.

Texture can also be estimated **by feel**: a moist sandy soil feels gritty and will not form a ball; a clay soil feels sticky and can be rolled into a long, flexible ribbon; a loam forms a ball that breaks easily.

1.6 Main Soil Types and their Distribution in Cameroon

Soil type	Properties	Occurrence in Cameroon
Sandy	Large particles; drains fast, warms quickly, low nutrient and water retention	Parts of the Far North (sahelian zone)
Clay	Tiny particles; holds water and nutrients well but poorly aerated, hard when dry, waterlogged when wet	Low-lying plains, e.g. Logone flood-plain
Loamy	Balanced sand-silt-clay mixture with humus; fertile and easy to work	Western highlands, many farmed areas
Lateritic	Red, rich in iron and aluminium oxides; heavily leached, low fertility unless manured	Humid South, Centre and East
Alluvial	Deposited by rivers; deep, fertile, regularly renewed by floods	Benue and Logone valleys, coastal plains
Volcanic	From weathered basalt; dark, deep, very fertile	Mount Cameroon and the Bamenda highlands
Peaty	Very rich in organic matter; acidic and often waterlogged	Waterlogged depressions and swamps

PROPERTY Loamy soil: the ideal agricultural soil

A **loam** (roughly 40% sand, 40% silt, 20% clay, plus humus) is the best general-purpose agricultural soil because it combines good drainage (from sand), good water and nutrient retention (from clay and humus) and good aeration and easy root penetration (from its crumb structure). Proof is not examinable; you must simply be able to justify the superiority of loam in terms of drainage, aeration and nutrient retention.

1.7 Soil Structure

DEFINITION Soil structure

Soil structure is the way in which soil particles are grouped and bound together into larger units called aggregates (or peds), and the arrangement of the pore spaces between them.

WARNING Texture ≠ structure


A very common GCE mistake is to confuse the two. Texture is about the sizes of individual particles (sand, silt, clay) and cannot be changed by cultivation. Structure is about how those particles are arranged into aggregates; it can be improved by adding humus or destroyed by cultivating wet soil. A clay soil (texture) can have an excellent crumb structure.

The main structural types are:

- **Crumb:** small, rounded, porous aggregates; ideal for seedbeds — excellent aeration, drainage and root growth.
- **Blocky:** larger, roughly cube-shaped aggregates; moderately good, common in subsoils.
- **Platy:** thin horizontal plates; impedes drainage and root penetration.
- **Prismatic:** vertical columns; restricts root spread and water movement.

1.8 Physical and Chemical Properties of Soil

Porosity is the percentage of soil volume occupied by pore spaces; it determines how much air and water the soil can hold. **Water-holding capacity** is the amount of water a soil retains against gravity — highest in clay and humus-rich soils, lowest in sands. **Capillarity** is the upward movement of water through fine pores, which can bring stored water up to roots but also causes water loss by evaporation from bare soil.



Comparison of water-holding capacity and drainage rate in sandy, loamy and clay soils (relative values). Loam offers the best balance.

Soil pH measures acidity or alkalinity. Most Cameroonian crops grow best between pH 5.5 and 7.0. Heavy tropical rainfall leaches away basic cations (Ca^{2+} , Mg^{2+} , K^+), leaving soils acidic. Acidity is corrected by **liming** — adding agricultural lime (calcium carbonate), which neutralises excess H^+ ions and also supplies calcium.

Cation exchange capacity (CEC) is the soil's ability to hold positively charged nutrient ions (K^+ , Ca^{2+} , Mg^{2+} , NH_4^+) on the surfaces of clay and humus particles, preventing them from being leached while keeping them available to roots. Clay and humus have high CEC; sand has very low CEC — one more reason sandy soils are infertile.

WORKED EXAMPLE 1 Choosing a crop for a given soil

A farmer near Maroua has a light, gritty soil that dries out two days after rain and gives low yields of maize despite heavy fertiliser use. (a) Identify the soil texture. (b) Explain the poor response to fertiliser. (c) Suggest one suitable crop and one soil improvement measure.

SOLUTION

- (a) Rapid drainage and gritty feel indicate a sandy soil.
- (b) Sand has very low cation exchange capacity and low water-holding capacity. Soluble fertiliser ions (e.g. K^+ , NO_3^-) are rapidly leached below the rooting zone by rain, so the plants cannot use them and the fertiliser is wasted.
- (c) A drought-tolerant crop such as groundnuts, millet or cowpea suits sandy soil. The soil can be improved by adding organic manure or compost, which raises water-holding capacity and CEC, and by applying fertiliser in small, frequent doses.

1.9 Soil Fertility and its Maintenance

DEFINITION Soil fertility

Soil fertility is the ability of a soil to supply plants with adequate water, air and mineral nutrients in the right proportions for healthy growth and good yields.

Fertility is not permanent: continuous cropping, leaching and erosion exhaust it. Good farmers maintain fertility by:

- **Crop rotation:** growing different crops in sequence on the same plot (e.g. maize → ground-nuts → cassava). Legumes restore nitrogen through their root-nodule bacteria, and pests specific to one crop are starved out.
- **Fallowing:** resting the land under natural vegetation for some years so that humus and nutrients accumulate again — the principle of traditional shifting cultivation.
- **Organic manures:** farmyard manure, compost and green manure add humus, improve structure and release nutrients slowly.
- **Inorganic (chemical) fertilisers:** supply concentrated nutrients quickly (e.g. NPK 20–10–10 for maize), but add no humus and can acidify soil if misused.
- **Cover cropping and mulching:** covering the soil with living plants (e.g. Pueraria) or dead material (grass, plantain leaves) protects against erosion, conserves moisture, suppresses weeds and adds organic matter.

Soil erosion is the removal of topsoil by running water or wind — the fastest way to destroy fertility, and a serious problem on the steep cultivated slopes of the North West and South West Regions. **Conservation methods** include:

- **Contour ploughing:** ploughing along lines of equal height so each furrow checks run-off.
- **Terracing:** cutting slopes into level steps to slow water flow (seen on the hills around Dschang).
- **Windbreaks:** rows of trees or tall crops that reduce wind speed and protect soil in the dry North.
- **Strip cropping, cover crops and mulching,** which keep the soil permanently covered.

KEY POINTS Key points

- Soil forms by weathering of parent rock plus organic activity, controlled by climate, organisms, relief, parent material and time.
- The profile shows horizons O, A, B, C above bedrock; the A horizon is the fertile topsoil.
- Texture = particle size proportions; structure = arrangement of aggregates. Do not confuse them.

- Loam is the ideal soil: balanced drainage, aeration and nutrient retention.
- Fertility is maintained by rotation, fallowing, manures, mulching and erosion control.

EXAMPLE Exam tip

GCE questions frequently ask you to relate soil properties to crop choice (e.g. “Explain why rice is grown in the clay soils of the Logone plain while groundnuts thrive in the sandy soils of the Far North”). Always structure your justification around the same three pillars: drainage, aeration and nutrient content (including CEC and pH). A property named without its consequence for the crop earns little credit.

EXERCISE 1 Test your understanding

1. Define soil texture and soil structure, giving one example of each.
2. A soil sample settles in a jar into layers of 6 cm sand, 3 cm silt and 1 cm clay. Calculate the percentage of each fraction and name the likely soil class.
3. Explain three ways in which adding farmyard manure improves a sandy soil.
4. A farmer on a steep slope near Bamenda loses topsoil every rainy season. Describe two conservation methods he could adopt, explaining how each works.

SOLUTION Exercise 1


1. Texture: the proportions of sand, silt and clay particles in a soil (e.g. a loam with 40% sand, 40% silt, 20% clay). Structure: the arrangement of particles into aggregates (e.g. crumb structure).
2. Total depth = $6 + 3 + 1 = 10$ cm. Sand = 60%, silt = 30%, clay = 10%. This is a sandy loam — free-draining but reasonably workable.
3. Manure (i) increases water-holding capacity, (ii) raises cation exchange capacity so nutrients are retained, and (iii) improves crumb structure and aeration as it decomposes to humus.
4. Contour ploughing: furrows along contour lines trap run-off and let water soak in instead of flowing downhill. Terracing: level steps reduce slope length and water speed, so soil is deposited rather than carried away. (Windbreaks or mulching are also acceptable with correct explanations.)

2 PLANT NUTRITIONAL REQUIREMENTS

In the previous section we saw that a fertile soil is not merely a pile of particles: it is a living, chemically active medium whose texture, structure, organic matter content and pH determine how well it can support a crop. But why does a maize plant in a red lateritic soil of the West Region turn pale and stunted, while the same variety thrives a few kilometres away on a well-manured plot? The answer lies in **plant nutrition**: the soil is the pantry, and the plant must draw from it a precise menu of chemical elements. This section studies that menu — which elements a plant needs, in what quantities, what happens when one is missing, and how the farmer can correct shortages with fertilisers.

2.1 Autotrophic Nutrition: The Basis of Crop Production

Green plants are **autotrophs**: they build their own organic matter from simple inorganic raw materials. The fundamental process is **photosynthesis**, summarised by:



Carbon dioxide provides the carbon skeleton, water provides hydrogen and electrons, and light provides the energy. The glucose produced is then converted into starch (the yam tuber, the cassava root, the maize grain we harvest), cellulose (stems, cotton fibre), proteins and oils.

WARNING What photosynthesis does NOT provide

Photosynthesis supplies carbon, hydrogen and oxygen — about 95% of a plant's dry mass. The remaining 5%, the mineral fraction, must come from the soil solution through the roots. A plant can photosynthesise perfectly and still die if a single essential mineral is missing. This is why soil fertility (Section 1) and mineral nutrition are inseparable.

Even with all minerals present, the **yield** of a crop is limited by whichever factor is in shortest supply: light intensity, CO_2 concentration, water availability, temperature, or mineral nutrients. This is the **law of the minimum** (associated with the chemist Liebig): growth is controlled not by the total of resources available, but by the scarcest resource — just as the water level in a leaking barrel can rise no higher than the shortest stave. In most Cameroonian farms, after water, the limiting factors are mineral nutrients — especially nitrogen, phosphorus and potassium.

2.2 Essential Mineral Elements

An element is **essential** if it satisfies three **criteria of essentiality**:

1. The plant cannot complete its life cycle (germination → growth → flowering → seed production) without it.
2. Its role is specific and cannot be replaced by any other element.
3. It is directly involved in plant metabolism (a constituent of a molecule or an enzyme activator), not merely improving growth indirectly.

Essential elements are divided into two groups according to the quantities required.

DEFINITION **Macronutrient and micronutrient**

A **macronutrient** is an essential element required by the plant in relatively large amounts (typically more than 1 g per kg of dry matter). The mineral macronutrients absorbed from the soil are: nitrogen (N), phosphorus (P), potassium (K), calcium (Ca), magnesium (Mg) and sulphur (S). (Carbon, hydrogen and oxygen, obtained from CO₂ and H₂O, are also macronutrients.)

A **micronutrient** (trace element) is an essential element required in very small amounts (milligrams per kg of dry matter): iron (Fe), manganese (Mn), boron (B), zinc (Zn), copper (Cu), molybdenum (Mo) and chlorine (Cl). Despite the tiny quantities, a deficiency of any one of them is just as damaging as a deficiency of a macronutrient.


2.3 Functions and Deficiency Symptoms of the Major Elements

Each element has precise roles, and its absence produces characteristic, diagnosable symptoms. A useful diagnostic clue is **element mobility** within the plant: mobile elements (N, P, K, Mg) are moved from old leaves to young growing points when scarce, so symptoms appear on older leaves first; immobile elements (Ca, Fe) cannot be relocated, so symptoms appear on young leaves first.

Element	Main functions	Deficiency symptoms
Nitrogen (N)	Constituent of proteins, nucleic acids, chlorophyll	General yellowing (chlorosis) of older leaves; stunted growth; thin, weak stems
Phosphorus (P)	ATP, nucleic acids, phospholipids; energy transfer; root development	Poor root growth; delayed maturity; dark green or purplish leaves (anthocyanin accumulation)
Potassium (K)	Enzyme activation; stomatal opening; osmoregulation; disease resistance	Leaf scorch : yellowing then browning of leaf margins of older leaves; weak stems, lodging in cereals
Magnesium (Mg)	Central atom of the chlorophyll molecule	Interveneal chlorosis of older leaves (veins stay green, tissue between them yellows)
Calcium (Ca)	Cell wall cement (calcium pectate); membrane stability	Death of growing points and root tips; blossom-end rot in tomato; symptoms on young tissues
Sulphur (S)	Constituent of some amino acids (cysteine, methionine)	General chlorosis of young leaves (resembles N deficiency but on young leaves)
Iron (Fe)	Chlorophyll synthesis (cofactor); electron transport	Interveneal chlorosis of young leaves; common in alkaline/calcareous soils

PROPERTY NPK: the elements most often limiting yield

Nitrogen, phosphorus and potassium are the three elements removed from the soil in the largest quantities by harvested crops, and they are therefore the ones most often deficient in cultivated soils. They form the basis of compound fertilisers labelled NPK, whose three numbers give the percentage by mass of N, P_2O_5 and K_2O respectively. For example, a bag marked 20–10–10 contains 20% nitrogen, 10% phosphorus pentoxide and 10% potassium oxide by mass. (Knowledge of this property is examinable; no proof is required.)



Stylised leaf showing the three classic chlorosis patterns: uniform yellowing (N), interveinal yellowing with green veins (Mg), and marginal scorch (K).

WARNING Not all yellow leaves mean nitrogen deficiency

The most common diagnostic error — in the field and in the GCE exam — is to attribute every yellowing to lack of nitrogen. Magnesium deficiency causes interveinal chlorosis of older leaves, and iron deficiency causes interveinal chlorosis of young leaves. Always note which leaves are affected (old or young) and where the yellowing occurs (whole leaf, between veins, or margins) before concluding.

2.4 Absorption of Minerals by Roots

Minerals are absorbed from the soil solution as ions (NO_3^- , NH_4^+ , $H_2PO_4^-$, K^+ , Ca^{2+} , Mg^{2+} ...), mainly by the **root hairs** just behind the root tip. Root hairs enormously increase the absorptive surface: a single maize plant may carry millions of them.

Absorption occurs by two mechanisms:

- **Passive uptake** (diffusion, mass flow with the transpiration stream) when the ion concen-

tration in the soil solution exceeds that in the root cells.

- **Active transport**, the dominant mechanism: carrier proteins in the root-cell membranes pump ions against their concentration gradient, using energy from ATP supplied by respiration. This is why waterlogged, airless soils (poor in O_2 for respiration) reduce mineral uptake even when minerals are abundant.

Two further points are essential:


Mycorrhizal associations. The roots of most crop plants form a mutualistic symbiosis with soil fungi (**mycorrhizae**). The fungal hyphae extend far beyond the root hairs, greatly increasing the absorbing surface and improving uptake of poorly mobile ions such as phosphate; in return the fungus receives carbohydrates from the plant.

Effect of soil pH. Nutrient availability depends strongly on pH (link with Section 1). In strongly acid soils ($pH < 5$, common in the humid forest zone of the South and Centre), phosphorus becomes fixed as insoluble iron and aluminium phosphates, and calcium and magnesium are leached away. In alkaline soils ($pH > 7.5$), iron, manganese, zinc and boron become unavailable, producing chlorosis even when the elements are present in the soil. Maximum overall availability occurs around $pH\ 6.0-7.0$, which is why liming acid soils is a standard corrective practice.

2.5 The Nitrogen Cycle and Biological Nitrogen Fixation

Nitrogen is the element needed in the greatest quantity, yet although air is about 78% N_2 , plants cannot use gaseous nitrogen directly. They depend on the **nitrogen cycle**, driven largely by micro-organisms:

1. **Nitrogen fixation:** conversion of atmospheric N_2 into ammonium compounds, either industrially (Haber process), by lightning, or biologically by bacteria such as free-living *Azotobacter* and symbiotic *Rhizobium*.
2. **Ammonification:** decomposition of organic nitrogen (proteins, urea in manure and dead matter) into ammonium ions NH_4^+ by saprotrophic bacteria and fungi.
3. **Nitrification:** oxidation of NH_4^+ to nitrites NO_2^- (by *Nitrosomonas*) then to nitrates NO_3^- (by *Nitrobacter*). Nitrate is the main form absorbed by crop roots.
4. **Plant uptake and assimilation:** NO_3^- is absorbed and built into amino acids and proteins, which pass along food chains.
5. **Denitrification:** under waterlogged, anaerobic conditions, denitrifying bacteria convert NO_3^- back to N_2 , a loss of fertility.



The nitrogen cycle: fixation, ammonification, nitrification, plant uptake and denitrification.

DID YOU KNOW?

Rhizobium and the Cameroonian mixed farm

Legumes such as groundnut, cowpea (niébé), soya and beans host Rhizobium bacteria in swellings on their roots called **root nodules**. The bacterium fixes atmospheric N_2 and passes ammonium compounds to the plant; the plant supplies the bacterium with carbohydrates. When the legume residues are ploughed in, the soil is enriched in nitrogen — which is why Cameroonian farmers traditionally intercrop maize with cowpea or rotate groundnut with cereals. This natural fertilisation reduces the need for expensive nitrogen fertiliser and is a classic GCE example of applied symbiosis.

2.6 Fertilisers: Organic and Inorganic

DEFINITION Fertiliser

A **fertiliser** is any material, organic or inorganic, added to the soil (or to a nutrient solution) to supply one or more essential plant nutrients and so maintain or increase crop yield.

Type	Advantages	Disadvantages
Organic: farmyard manure, compost, green manure (ploughed-in legumes), crop residues	Release nutrients slowly; improve soil structure, water retention and humus content; feed soil organisms; cheap and locally available	Bulky to transport; nutrient content low and variable; slow action; may carry weed seeds or pathogens if not well rotted
Inorganic (mineral): NPK compounds, urea (46% N), ammonium sulphate (21% N + S), single superphosphate	Concentrated and fast-acting; precise, known composition; easy to store and apply; doses can match crop needs	Expensive; no improvement of soil structure; easily leached (nitrates); risk of over-use: eutrophication, soil acidification (ammonium sulphate), salt damage

Correct application methods.

- **Broadcasting:** spreading evenly over the whole field, usually before planting.
- **Placement (localised) application:** putting the fertiliser in the planting hole or in a band a few centimetres beside and below the seed — more efficient, and avoids scorching the seed.
- **Top-dressing:** applying nitrogen (e.g. urea) to a growing crop, e.g. maize at knee-height; should be done on moist soil and lightly covered to limit volatilisation losses.
- **Timing and dose:** match the crop's demand curve; split nitrogen applications rather than applying one heavy dose.

WORKED EXAMPLE 2 Reading an NPK label and calculating a dose

A farmer buys a 50 kg bag of compound fertiliser labelled 15–15–15. (a) What mass of nitrogen does the bag contain? (b) The recommended nitrogen dose for his maize crop is 60 kg of N per hectare. How many bags must he apply per hectare?

SOLUTION

(a) The first number of the label gives the percentage of nitrogen by mass:


$$m(\text{N}) = 50 \text{ kg} \times \frac{15}{100} = 7.5 \text{ kg of N per bag.}$$

(The bag also contains 7.5 kg of P_2O_5 and 7.5 kg of K_2O .)

(b) Number of bags required:

$$n = \frac{60 \text{ kg N ha}^{-1}}{7.5 \text{ kg N per bag}} = 8 \text{ bags per hectare.}$$

The farmer must therefore apply 8 bags (400 kg) of 15–15–15 per hectare to supply 60 kg of nitrogen.



Yield versus fertiliser rate: each additional unit of fertiliser gives a smaller yield increase (law of diminishing returns) until a ceiling is reached where another factor becomes limiting.

The curve above illustrates the **law of diminishing returns**: beyond the economic optimum, extra fertiliser costs more than the extra yield is worth, and the excess leaches into rivers.

WARNING Dangers of over-fertilisation

Applying too much mineral fertiliser does not give unlimited yield. Excess nitrate leaches into streams and lakes, causing **eutrophication**: algal blooms, deoxygenation and death of aquatic life. Continuous use of ammonium sulphate acidifies the soil; heavy urea or salt-based fertiliser doses can “burn” seedlings. Fertiliser is a medicine: the right dose cures, the wrong dose poisons.

2.7 Experimental Evidence: Water Culture (Hydroponics)

How do we know that, say, magnesium is essential? By growing plants in a solution containing everything except magnesium and observing the result. This is the **culture solution** (water culture) technique, and **hydroponics** is its commercial extension.

METHOD Setting up a culture-solution experiment to test essentiality

1. Prepare a complete culture solution containing known salts supplying all essential elements (e.g. potassium nitrate, calcium phosphate, magnesium sulphate, an iron salt and a trace-element mixture) in distilled water.
2. Prepare a series of deficient solutions, each identical to the complete solution but lacking one element (e.g. replace magnesium sulphate by potassium sulphate to re-

move Mg while keeping S).

3. Place seedlings of the same species, age and size (e.g. young maize or tomato) in each solution, supporting them so that only the roots are immersed; wrap the containers in foil to exclude light and prevent algal growth.
4. Aerate the solutions continuously (aquarium pump) so that roots can respire; renew the solutions regularly to keep composition and pH stable.
5. Grow all plants under identical light, temperature and humidity for several weeks.
6. Observe and record: plants in the complete solution grow normally; each deficient solution produces its characteristic symptom (e.g. interveinal chlorosis of old leaves without Mg). Confirm by re-adding the missing element: recovery proves essentiality.

WORKED EXAMPLE 3 Diagnosing a field deficiency

A farmer near Bafoussam notices that the older leaves of his maize turn yellow between the veins while the veins remain green; the young leaves look normal. (a) Name the likely deficient element and justify. (b) State the element's function. (c) Suggest two corrections.

SOLUTION

(a) The symptom is interveinal chlorosis on older leaves. Because the element concerned is mobile in the plant, it is withdrawn from old leaves to supply young ones, so old leaves suffer first. Interveinal chlorosis of old leaves is characteristic of magnesium deficiency (nitrogen would yellow the whole leaf uniformly; iron would attack young leaves).

(b) Magnesium is the central atom of the chlorophyll molecule; without it chlorophyll cannot be formed, hence the yellowing and reduced photosynthesis.

(c) Corrections: apply a magnesium-containing fertiliser (e.g. magnesium sulphate / kieserite, or dolomitic lime if the soil is also acid); and add organic matter (compost or farmyard manure), which releases Mg slowly and improves its retention in the soil. Checking and correcting soil acidity by liming also improves Mg availability.

EXERCISE 2 Deficiency symptoms table

Copy and complete a table with three columns — Element / Function / Deficiency symptom — for N, P, K, Mg, Ca and Fe. Then answer: a tomato grower observes death of the shoot tips and rotting of the blossom end of young fruits. Which element is deficient, and why do the symptoms appear on the youngest tissues?

SOLUTION Exercise 1

Table: see the summary table in this section (N: protein/chlorophyll, uniform chlorosis of old leaves; P: ATP/nucleic acids, poor roots and purpling; K: enzyme activation/stomata, marginal scorch of old leaves; Mg: chlorophyll centre, interveinal chlorosis of old leaves; Ca: cell walls, death of growing points; Fe: chlorophyll synthesis, interveinal chlorosis of young leaves).

The tomato grower's plants lack calcium. Calcium is immobile in the plant (it is fixed in cell walls as calcium pectate), so it cannot be moved from old tissues to new ones; the youngest tissues — shoot tips and developing fruits — are therefore the first to show deficiency, giving death of growing points and blossom-end rot.

EXERCISE 3 Fertiliser calculation

A bag of urea is labelled 46% N and has a mass of 50 kg. (a) Calculate the mass of nitrogen in one bag. (b) A yam grower wishes to supply 92 kg of nitrogen per hectare. How many bags of urea does he need per hectare? (c) Give one reason why he should split this dose into two or three applications rather than applying it all at planting.

SOLUTION Exercise 2

(a) $m(\text{N}) = 50 \times \frac{46}{100} = 23 \text{ kg of N per bag.}$

(b) $n = \frac{92}{23} = 4 \text{ bags per hectare.}$

(c) Nitrogen from urea is rapidly converted to nitrate, which is easily leached by heavy rain before the crop can absorb it; splitting the dose matches supply to the crop's demand, reduces leaching losses (and the risk of eutrophication) and avoids scorching young plants with a single concentrated dose.

KEY POINTS Key points for the GCE

- Plants are autotrophs: CO_2 and water give C, H, O; all other elements come from the soil solution.
- Macronutrients: N, P, K, Ca, Mg, S. Micronutrients: Fe, Mn, B, Zn, Cu, Mo, Cl. Essentiality is proved by culture-solution experiments.
- Learn deficiency symptoms in table form (element – function – symptom): this is a frequent GCE short-answer item. Remember mobility: mobile elements → old leaves first; immobile → young leaves first.

- N, P and K most often limit yield $\Rightarrow$ NPK fertilisers; the label numbers give % N, % P_2O_5 , % K_2O ; yield follows the law of diminishing returns.
- Rhizobium in legume nodules fixes atmospheric nitrogen — the biological basis of intercropping and rotation in Cameroonian mixed farming.
- Over-fertilisation causes eutrophication and soil acidification: correct diagnosis and correct dose are the farmer's (and the candidate's) watchwords.

3 CULTIVATION TECHNIQUES OF FOOD AND NON-FOOD CROPS

In the previous section we established that a crop can only yield well if the soil supplies it with the right mineral elements in the right amounts. But even the most fertile soil in the Centre Region will give a poor harvest if the farmer plants the wrong variety, at the wrong time, at the wrong spacing, and never weeds. Feeding the plant is necessary; managing the plant is equally essential. This section takes you through the complete **cultivation** sequence — from the choice of crop and the preparation of the land to harvesting, storage and marketing — using Cameroonian crops as our working examples.

3.1 Classification of Crops

Farmers and agricultural officers group crops according to their main use.

DEFINITION Food crop, cash crop

A **food crop** is a crop grown primarily for human (or animal) consumption: maize, cassava, plantain, yam, cocoyam, groundnut, cowpea, rice, millet, sorghum. A **cash crop** (or non-food/industrial crop) is grown primarily for sale — for export or for industry — rather than for direct consumption: cocoa, coffee, cotton, oil palm, rubber, tobacco, banana (export), tea, sugar cane. Note that the distinction is about purpose, not biology: a farmer may sell part of his maize and eat part of his cocoa-bean income.

Criterion	Food crops	Cash crops
Main purpose	Feeding the family / local markets	Sale, export, agro-industry
Examples in Cameroon	Maize, cassava, plantain, yam, groundnut, cowpea, rice, millet, sorghum	Cocoa, coffee, cotton, oil palm, rubber, tobacco, banana (export), tea, sugar cane
Typical scale	Smallholdings (0.5–3 ha)	Smallholdings or large plantations (CDC, SOCAPALM, HEVE-CAM)
Life cycle	Mostly annual or short-cycle	Mostly perennial tree crops

3.2 Farming Systems in Cameroon

A **farming system** is the overall way a farmer organises land, labour and crops.


- **Shifting cultivation:** a piece of forest or bush is cleared, burned, cropped for 2–3 years, then abandoned when fertility falls; the farmer moves to a new plot. Simple but destructive when population pressure shortens the fallow cycle below the threshold needed for soil recovery.
- **Bush fallowing:** an improvement on shifting cultivation — plots are cultivated in rotation and left to fallow (rest under natural vegetation) for several years so that soil fertility is restored before re-cropping. Common in the humid forest zone where land is still relatively abundant.
- **Subsistence farming:** production aimed mainly at feeding the farmer's household, with small surpluses sold locally. Typical of food-crop farms in the West and North-West Regions.
- **Plantation agriculture:** large estates growing a single cash crop (monoculture) with hired labour and processing facilities, e.g. CDC banana and rubber estates in the South-West, SOCAPALM oil palm estates in the Littoral and South-West, HEVECAM rubber in the Centre and South.
- **Mixed farming:** crops and livestock are kept on the same farm; crop residues feed the animals and manure fertilises the fields — common in the Adamawa and North-West highlands.
- **Market gardening (maraîchage):** intensive production of vegetables (tomato, cabbage, carrot, onion, green beans, huckleberry) on small, well-watered plots near towns, e.g. the valleys around Bafoussam, Santa, Dschang and the Wouri lowlands near Douala.
- **Agroforestry systems:** deliberate integration of trees with crops and/or livestock — e.g. cocoa under shade trees, coffee under Albizia or Leucaena, or Acacia alley cropping in the Sudan-Sahel zone.

3.3 Land Preparation

Land preparation creates a suitable **seedbed**: loose, aerated, weed-free soil in which roots can penetrate and water can infiltrate.

1. **Clearing:** removal of trees, shrubs and previous crop residues (cutlass, axe, chainsaw, or bulldozer on plantations). In forest zones clearing is done at the end of the rainy season (November–January); debris may be burned (adds ash/potash but destroys organic matter and soil organisms) or left as mulch (slash-and-mulch).
2. **Ploughing:** the primary tillage — turning over the topsoil (15–25 cm depth) with a hoe, animal-drawn plough or tractor to bury weeds and loosen the soil. On steep slopes, contour ploughing or minimum tillage reduces erosion.
3. **Harrowing:** secondary tillage — breaking the clods left by ploughing into a fine tilth and levelling the surface. Done with a harrow, rake or hoe.
4. **Ridging and bed making:** soil is heaped into ridges (yam, groundnut, potato, cassava in wet areas) or raised beds (vegetables, nursery) to improve drainage and give tubers room to swell. In the Sudan-Sahel, tied ridges (with cross-ties) conserve runoff water.

Timing with the seasons. In the humid forest zone (bimodal rainfall), the main planting follows the onset of the heavy rains (March–April) with a second season around August–September. In the savanna zones (unimodal rainfall), land is prepared at the end of the dry season (February–April) and planting waits for the first reliable rains (May–June). Planting too early exposes seeds to drought; too late shortens the growing season and increases pest pressure.



Production cycle of maize, from land preparation to storage. Part of the harvest is kept as seed for the next season.


3.4 Propagation Methods

Crops are multiplied either by **seeds** (sexual propagation) or by plant parts (asexual/vegetative propagation).

DEFINITION Vegetative propagation

Vegetative propagation is the production of new plants from vegetative parts — stems (cuttings, suckers), underground organs (tubers, rhizomes, corms, setts) — or by artificial joining of plant parts (grafting, budding, layering) — without seeds. The offspring are genetically identical to the parent (a clone), so desirable characters are preserved; the drawback is that diseases of the parent are transmitted and there is no genetic variation for adaptation.

Method	Principle	Cameroonian examples
Seed (sexual)	Sowing of fertilised ovules	Maize, cowpea, groundnut, rice, millet, sorghum, cocoa, oil palm (nursery)
Stem cutting	A stem piece (20–30 cm, 5–7 nodes) roots and sprouts	Cassava, sugarcane, sweet potato vines, Gmelina, Leucaena
Tuber / sett	Underground organ with buds (“eyes”)	Yam (setts 200–500 g), potato (seed tubers), cocoyam (corms/cormels)
Sucker	Shoot from the base of the mother plant	Plantain, banana, pineapple, sisal
Rhizome division	Underground stem with buds	Ginger, turmeric
Grafting	A scion is joined onto a rooted stock (whip, cleft, side)	Mango, citrus, avocado, improved rubber (GT1, PB series)
Budding	A single bud is inserted under the stock bark (T-bud, patch bud)	Rubber, citrus, cocoa (improved clones from IRAD/SODECAO)
Layering	Stem rooted while still attached to mother	Citrus, mango, some ornamentals



Three vegetative propagation methods: stem cutting, grafting (scion joined to stock) and budding (a single bud inserted in a T-shaped cut).

Seed selection and treatment. Only bold, clean, undamaged seeds of a good variety should be planted. Diseased or weevil-bored grains give weak seedlings. Seeds may be treated before sowing: dressing with fungicide/insecticide against soil pests (e.g. maize seed dressed with thiram + imidacloprid), or soaking (e.g. rice for 24 h, maize for 6–12 h) to speed germination. For legumes, inoculation with *Rhizobium* strains specific to the crop enhances nitrogen fixation.

METHOD Carrying out a seed germination test

1. Take a representative sample of the seed lot and count out 100 seeds (or 4 replicates of 25).

2. Place them on moist cotton wool, blotting paper or sand in a covered dish; keep warm (about 25–30 °C) and moist, in the dark or diffuse light.
3. After the normal germination period for the species (e.g. 5–7 days for maize, 7–10 days for cocoa), count the seeds that have produced a normal radicle and plumule (discard abnormal seedlings).
4. Calculate the percentage germination:

$$\text{Germination (\%)} = \frac{\text{number of normal seedlings}}{\text{number of seeds sown}} \times 100$$

5. Decide: a lot giving less than about 80 % germination (for most cereals and legumes) should be rejected or sown at a higher rate. For cocoa, minimum is 90 %.

WORKED EXAMPLE 4 Germination test and seed requirement

A farmer in Ngaoundéré tests 50 maize seeds; 43 germinate normally. He wants about 40 000 plants per hectare. (a) Calculate the percentage germination. (b) How many viable seeds per hectare must he sow, assuming one seed per stand?

SOLUTION

(a) Germination = $\frac{43}{50} \times 100 = 86\%$. The lot is acceptable (above 80 % for maize).

(b) Only 86 % of sown seeds will give a plant, so:

$$\text{Seeds to sow} = \frac{40\,000}{0.86} \approx 46\,512 \text{ seeds per hectare.}$$

He should sow about 46 500 seeds per hectare (roughly 12 kg at about 3 900 seeds/kg) to obtain the target stand.

3.5 Planting Practices

Spacing and depth. Each crop has an optimum spacing determined by its growth habit, root system and target yield. Sowing depth must suit seed size and soil type: small seeds shallow (1–2 cm), large seeds or cuttings deeper (5–10 cm for maize; cassava stakes buried two-thirds of their length, buds upward).

Crop	Spacing (cm)	Depth (cm)	Plant population (s
Maize (grain)	75 × 25 (1 seed/stand)	5–7	53 333
Maize (green)	50 × 25 (2 seeds/stand)	5–7	160 000
Cassava	100 × 100	stake 2/3 buried	10 000
Yam	100 × 100	sett 5–10	10 000
Plantain / Banana	300 × 300	sucker in 30 cm hole	1 111
Cocoa	300 × 300	seedling in 40 × 40 × 40 cm hole	1 111
Oil palm	900 × 900 × 900 (triangular)	seedling in 60 cm cube hole	143
Groundnut	50 × 15 (1 seed/stand)	5–6	133 333
Rice (lowland)	20 × 20 (3–4 seedlings/hill)	2–3 (transplanted)	250 000 h
Cotton	90 × 30 (2–3 seeds/stand)	3–5	74 000

PROPERTY Spacing and yield

Correct spacing maximises yield per hectare because it minimises **competition** between plants for light, water and mineral nutrients while using the land fully. Too dense a stand gives many small, etiolated plants and low yield per plant; too wide a stand wastes land and lets weeds invade. (Not examinable as a proof, but you must be able to explain it in terms of intraspecific competition and light interception.)

Nursery techniques. Some crops are first raised in a **nursery** — a small, sheltered, well-watered seedbed — then transplanted. This allows close care (watering, shading, protection from pests) during the delicate seedling stage and gives a head start on weeds.

- **Cocoa:** seeds sown in shaded nursery beds or polythene bags (4 × 6 cm); transplanted at 4–6 months (4–6 leaves) into field at 3 × 3 m.
- **Oil palm:** pre-nursery (3–4 months in small bags) → main nursery (12–15 months in large bags) → field at 9 × 9 × 9 m triangular.
- **Vegetables** (tomato, cabbage, onion, pepper): raised in nursery beds (1 m wide, raised 15 cm, shaded) and transplanted at 3–5 weeks (4–6 true leaves).
- **Coffee:** seeds in nursery beds, transplanted at 6–9 months.

WARNING Planting cuttings the right way up

A classic and costly mistake: cassava stakes and yam setts planted upside down or too deep sprout poorly or rot. Cassava stakes must be planted with the buds (nodes) pointing upward; yam setts with the sprouting end up, about 5–10 cm deep. Reversed or deeply buried cuttings give weak, late sprouts and a sharply reduced yield.

DID YOU KNOW?**Improved planting material in Cameroon**

IRAD (Institute of Agricultural Research for Development) and SODECAO (Cocoa Development Corporation) distribute improved varieties: maize (CMS 8704, CMS 9015, TZEE-Y), cassava (TMS 30572, TMS 4(2)1425 resistant to mosaic), cocoa (SNKA 12, SNKA 13, SNKA 14 — high yield, black pod tolerant), oil palm (Tenera hybrids from Dami and La Mé). Always source seed from certified multipliers to ensure genetic purity and phytosanitary quality.

3.6 Crop Maintenance

After planting, the farmer must protect the crop's access to light, water and nutrients:

- **Weeding:** removes competitors; usually 2–3 weeding per season for annual crops (e.g. maize at 2–3 and 6–7 weeks after planting). Methods: hand hoe, cutlass, animal-drawn cultivator, or selective herbicides (pre-emergence: atrazine + metolachlor for maize; post-emergence: nicosulfuron for grasses).
- **Mulching:** covering the soil with dry grass, leaves, crop residues or plastic film to conserve moisture, suppress weeds, moderate soil temperature and add organic matter (standard practice in plantain, cocoa young plantations and market-garden plots).
- **Watering / irrigation:** essential for nurseries, market gardening and dry-season cropping. Methods: watering can, hose, sprinkler, drip irrigation (increasingly used for high-value vegetables), flood irrigation (irrigated rice at Yagoua, Maga, SEMRY schemes).
- **Staking:** supporting climbing or heavy-bearing plants (yam vines — 2 m stakes, tomato — stakes or trellis, passion fruit) to keep fruits off the soil and improve air circulation.
- **Pruning:** removing dead, diseased or excess branches (cocoa — chupons and low branches; coffee — multiple stem pruning, capping at 1.5–2 m; fruit trees — formative and maintenance pruning) to improve aeration, light penetration and yield.
- **Thinning:** removing excess seedlings so the remaining ones have the correct spacing (maize: thin to 1 plant/stand at 2 weeks; vegetables: thin to final spacing at 2–3 true leaves).
- **Earthing-up:** heaping soil around the base of plants (maize at 6–7 weeks to support stems and cover brace roots; groundnut at flowering to encourage peg penetration; potato to prevent tuber greening; cassava in waterlogged areas).
- **Fertiliser application:** basal (NPK at planting) and top-dressing (urea at knee-high for maize, at flowering for groundnut). Rates based on soil test; typical: maize 200 kg/ha NPK 20–10–10 + 100 kg/ha urea; cocoa 300 kg/ha NPK 12–12–17+2MgO+Te per year split in two applications.

3.7 Pests, Diseases and Weeds of Major Crops

A **pest** is any organism that damages the crop; control methods fall into three groups:

- **Cultural control:** practices that prevent or reduce attack — crop rotation, early planting, resistant/tolerant varieties, destruction of crop residues, clean planting material, proper

spacing, field hygiene, trap crops (e.g. Desmodium for stem borer push-pull).

- **Biological control:** use of natural enemies — e.g. parasitoid wasp *Anagyrus lopezi* against cassava mealybug (classical success in 1980s), *Bacillus thuringiensis* (Bt) against caterpillars, *Trichoderma* spp. against soil-borne fungi, neem (*Azadirachta indica*) extracts as botanical insecticide.
- **Chemical control:** pesticides (insecticides, fungicides, herbicides, nematicides). Effective but costly, polluting and dangerous if misused; always a last resort, applied at the correct dose with protective equipment (PPE), respecting pre-harvest intervals (PHI).

DEFINITION Integrated pest management (IPM)

Integrated pest management is the combined, rational use of cultural, biological and (minimal, well-targeted) chemical methods to keep pest populations below the economic damage threshold, while protecting the farmer, the consumer and the environment. Key steps: monitoring → threshold decision → selective intervention → evaluation.

Crop	Important enemies	Main IPM components
Maize	Stem borers (<i>Busseola</i> , <i>Chilo</i>), fall armyworm (<i>Spodoptera frugiperda</i>); maize streak virus (MSV); striga (<i>Striga hermonthica</i>) parasitic weed	Early planting, resistant varieties (TZEE-Y, CMS 9015), rotation with legumes, push-pull (<i>Desmodium</i> /Napier), targeted Bt or emamectin benzoate for FAW
Cassava	Cassava mealybug (<i>Phenacoccus manihoti</i>), green mite (<i>Mononychellus tanajoa</i>); cassava mosaic disease (CMD), cassava brown streak disease (CBSD)	Clean cuttings (certified), resistant varieties (TMS 30572, TME 419), biological control (mealybug), roguing infected plants
Cocoa	Capsid bugs (<i>Sahlbergella</i> , <i>Distantiella</i>); black pod disease (<i>Phytophthora megakarya</i> , <i>P. palmivora</i>); swollen shoot virus (CSSV)	Regular harvesting of diseased pods (sanitation), pruning for aeration, shade management, resistant clones (SNKA series), copper-based fungicide sprays (8–10 rounds/yr), vector control for CSSV
Groundnut	Aphids (<i>Aphis craccivora</i>) vector of rosette virus; leaf spots (<i>Cercospora</i> early/late); termites	Resistant varieties (ICGV-SM 90704, RMP 12), early planting (escape aphids), seed treatment, fungicide (chlorothalonil) for leaf spot, crop rotation
Plantain / Banana	Banana weevil (<i>Cosmopolites sordidus</i>), nematodes (<i>Radopholus</i> , <i>Pratylenchus</i>); black Sigatoka (<i>Mycosphaerella fijiensis</i>)	Clean planting material (boiling water treatment 50°C 20 min or tissue culture), mulching, nematicide (if economic), resistant hybrids (FHIA, PITA), deleafing for Sigatoka
Cotton	Bollworms (<i>Helicoverpa</i> , <i>Diparopsis</i>), aphids, jassids; bacterial blight	Early planting, resistant varieties (IRMA 1243, FK 37), threshold-based sprays (pyrethroids + organophosphates rotation), destruction of ratoon cotton

3.8 Harvesting, Processing and Storage

Maturity indices tell the farmer when to harvest. Harvesting too early gives immature, watery produce that rots; too late invites field pests, shattering losses and quality decline.

Crop	Maturity indices
Maize (grain)	Husks dry and brown, grains hard (dent stage), moisture content 20–25%
Cassava	9–18 months after planting (variety dependent), tubers full size, leaves yellowing
Yam	7–10 months, vines yellowing and drying, tuber skin set
Plantain / Banana	Fingers plump, angularities disappear, colour change (green to light green/yellow), 12–15 weeks after flowering
Cocoa	Pods turn yellow/orange (depending on variety), 5–6 months after pollination
Groundnut	Leaves yellow, pods well-filled with veined interior, 100–130 days after planting
Rice	Grains hard, 80–85% straw coloured, moisture 20–22%
Cotton	Bolls fully open, lint fluffy, 150–180 days after planting

Harvesting techniques: hand-picking (cocoa pods with cutlass or hooked knife, taking care not to damage the flower cushion), pulling (groundnut, cotton), digging (yam, cassava, potato — careful to avoid tuber damage), cutting (maize cobs, rice panicles), mechanical combine (large-scale rice, maize in some schemes).

Processing and drying:

- **Maize:** shelled (by hand, stick, or sheller) and dried on clean surfaces (mats, tarpaulins, raised platforms) to 12–13 % moisture; winnowed to remove chaff.
- **Cassava:** highly perishable (48–72 h); processed into gari (fermented, roasted granules), flour (dried chips milled), water-fufu (fermented paste), or chips (dried). Detoxification (cyanogenic glucosides) by fermentation, boiling, roasting.
- **Cocoa:** beans + pulp removed from pods, fermented 5–6 days in heaps (covered with banana leaves) or wooden boxes (turned every 2 days) — fermentation develops chocolate flavour precursors, kills embryo, reduces bitterness; then sun-dried on raised platforms to 7–8 % moisture.
- **Coffee:** wet processing (pulping → fermentation 12–36 h → washing → drying) for arabica; dry processing (whole cherry dried) for robusta.
- **Oil palm:** fresh fruit bunches (FFB) processed within 24 h (sterilisation → stripping → digestion → pressing → clarification) to crude palm oil (CPO) and palm kernel oil (PKO).

Storage: dried grains are kept in granaries (traditional raised cribs), jute or polythene bags on pallets (not directly on floor), or airtight metal silos / hermetic bags (PICS bags), in a dry, ventilated store. The great enemies of stored produce are **storage pests**:

- **Primary insects:** weevils (*Sitophilus zeamais* in maize, *S. oryzae* in rice) and lesser grain borer (*Rhyzopertha dominica*) bore into whole grains. **Secondary insects:** flour beetles (*Tribolium*), saw-toothed grain beetle (*Oryzaephilus*) feed on damaged grain/dust.
- **Pulse beetles:** bruchids (*Callosobruchus maculatus* in cowpea, *Acanthoscelides obtectus* in beans) lay eggs on pods/seeds; larvae develop inside.

- **Control:** thorough drying (< 13% mc), clean stores (hygiene), insecticide dusting (actellic dust, pirimiphos-methyl) or fumigation (phosphine — certified operators only), airtight storage (hermetic bags, metal silos), use of ash, neem leaves or Ocimum in traditional stores.
- **Moulds** (*Aspergillus flavus*, *A. parasiticus*) produce dangerous aflatoxins in groundnut, maize, cottonseed when stored wet (> 14% mc). Control: dry properly before storage, ventilate, avoid re-wetting, sort mouldy grains, use aflasafe (biocontrol *A. flavus* atoxigenic strains) in field.

WARNING Aflatoxin — silent killer

Aflatoxins are potent carcinogens (liver cancer) and immunosuppressants. They are invisible, odourless and heat-stable (not destroyed by cooking). Groundnut and maize in Cameroon frequently exceed the EU limit (4 ppb total aflatoxins). Prevention starts in the field: avoid drought stress at pod-filling, harvest promptly, dry fast on clean surfaces, store dry. Never blend mouldy grain with clean grain — it contaminates the whole lot.

3.9 Case Studies: Maize and Cocoa

Maize (food crop, *Zea mays*).


1. **Site selection:** well-drained loam, pH 5.5–7.0, rainfall 600–1200 mm.
2. **Land preparation:** clear, plough, harrow at end of dry season (Feb–Mar in bimodal, Apr–May in unimodal). Conservation agriculture: minimum tillage + cover crop (*Mucuna*) in some zones.
3. **Planting:** at first effective rains. Selected seed (germination > 80 %, dressed) sown 75 × 25 cm, 5–7 cm deep, 1 seed/stand (grain) or 2 seeds/stand (green maize). Basal fertiliser: NPK 20–10–10 at 200 kg/ha in band beside row.
4. **Maintenance:** 1st weeding at 2–3 weeks; 2nd weeding + earthing-up + top-dress urea (100 kg/ha) at 6–7 weeks (knee-high). Monitor for fall armyworm (scout 20 plants/ha, spray if > 20% plants with fresh damage).
5. **Harvest:** 90–120 days (grain), 60–70 days (green). Cobs harvested when husks dry, shelled, dried to 12–13% mc.
6. **Storage:** in PICS bags or treated bags on pallets; monitor for weevils.
7. **Marketing:** sold to traders, cooperatives, or used for poultry feed (major demand from poultry industry in Yaoundé, Douala, Bafoussam).

Cocoa (cash crop, *Theobroma cacao*). Perennial crop of the humid forest zone (rainfall 1500–2500 mm, well-distributed).

1. **Nursery:** seeds from approved seed gardens (SODECAO/IRAD) sown in polythene bags (4 × 6 cm) under 50–60% shade; watered daily; transplanted at 4–6 months (4–6 hardened leaves).
2. **Field establishment:** holes 40 × 40 × 40 cm at 3 × 3 m (1111 trees/ha), lined with topsoil + 5

kg compost + 100 g NPK. Plant at onset of rains (Mar–Apr or Aug–Sep). Temporary shade: plantain (1:1) or Leucaena; permanent shade: Albizia, Terminalia, Milicia (iroko) at 12–15 m spacing.

- Young plantation (0–3 yr):** weeding (ring weeding 1 m radius), mulching, shade management, fertiliser (NPK 12–12–17+2MgO at 100 g/tree/yr split), pest monitoring.
- Mature plantation (3+ yr):** pruning (remove chupons, dead wood, low branches — once/yr after main harvest), sanitation harvesting (remove black pods weekly), fertiliser (300 kg/ha/yr NPK 12–12–17+2MgO split Mar/Apr and Sep/Oct), capsid control (8–10 sprays/yr with approved insecticides, rotation of modes of action), black pod control (copper hydroxide/fosetyl-Al sprays at pod set).
- Harvest:** main crop Oct–Mar, light crop May–Aug. Pods cut with knife/hook (no damage to cushion), opened with wooden mallet, beans + pulp heaped/boxed for fermentation.
- Fermentation & drying:** 5–6 days fermentation (turn day 2 and 4), sun-dried on raised platforms to 7–8% mc (beans rattle, no mould).
- Marketing:** sold to licensed buying agents (LBAs) or cooperatives; quality checked (bean count/100g < 100, moisture < 8%, mould < 4%, slaty < 8%). Price set by National Cocoa and Coffee Board (NCCB) based on world market.



Seasonal calendar of cropping activities in Cameroon. Dark bars: main season; pale bars: secondary season. The humid forest zone has two rainy seasons (two maize plantings possible); the savanna zone has one.

KEY POINTS Key points

- Food crops feed people; cash crops are grown for sale — the distinction is one of purpose, not biology.
- The cultivation sequence is always: land preparation → propagation/planting → maintenance → protection → harvesting → processing/storage.
- Vegetative propagation gives uniform clones but transmits diseases; seeds give vari-

ation and are easier to store and transport.

- Correct spacing, depth and orientation of planting material determine germination, sprouting and final yield.
- IPM combines cultural, biological and chemical control; proper drying is the key to safe storage and aflatoxin prevention.
- Improved varieties (IRAD, SODECAO, IITA, CIRAD) and certified planting material are essential for productivity and disease management.

WARNING Exam tip — structuring a crop essay

For any GCE essay asking you to “describe the cultivation of a named crop”, organise your answer in the chronological order of the production cycle: land preparation → planting (propagation material, spacing, depth) → maintenance (weeding, fertiliser, pest and disease control) → harvesting (maturity index, technique) → processing and storage. Add the climatic/soil requirements at the start and marketing at the end for full marks. Always name a real crop (maize, cocoa, cassava, plantain) and give real figures (spacing, duration, fertiliser rates, plant population).

EXERCISE 4 Spacing and plant population

A farmer plants cassava at a spacing of $1\text{ m} \times 1\text{ m}$ on a square field of side 100 m . (a) How many stands does he plant? (b) If each stand yields on average 4 kg of tubers, what is the expected yield in tonnes per hectare? (c) Explain why doubling the number of stands by planting at $0.5 \times 0.5\text{ m}$ would probably reduce the yield per stand.

SOLUTION Exercise 1

(a) Area of field = $100 \times 100 = 10\,000\text{ m}^2 = 1\text{ ha}$. Each stand occupies 1 m^2 , so he plants $10\,000$ stands.

(b) Yield = $10\,000 \times 4 = 40\,000\text{ kg} = 40\text{ tonnes per hectare}$.

(c) At $0.5 \times 0.5\text{ m}$, each stand has only 0.25 m^2 . Intraspecific competition for light, water and mineral nutrients becomes severe: the plants shade one another, roots overlap, and each plant produces fewer and smaller tubers. Yield per stand falls, and total yield per hectare usually falls too — this illustrates why correct spacing, not maximum density, maximises yield.

EXERCISE 5 Seed rate calculation

A farmer wants to plant 1 hectare of maize at 75 × 25 cm spacing (1 seed per stand). The seed lot has a germination percentage of 85% and a 1000-grain weight of 320 g. Calculate: (a) the number of stands per hectare; (b) the number of seeds to sow per hectare; (c) the weight of seed required in kg.

SOLUTION Exercise 2

$$(a) \text{ Stands per hectare} = \frac{10\,000 \text{ m}^2}{0.75 \times 0.25 \text{ m}^2} = \frac{10\,000}{0.1875} = 53\,333 \text{ stands/ha.}$$

$$(b) \text{ Seeds to sow} = \frac{53\,333}{0.85} = 62\,745 \text{ seeds/ha.}$$

$$(c) \text{ Weight} = \frac{62\,745 \times 320 \text{ g}}{1000} = 20.08 \text{ kg} \approx 20.1 \text{ kg/ha.}$$

EXERCISE 6 Integrated pest management

A cocoa farmer in the Centre Region observes black pod disease on 15% of pods in July. He has been spraying a copper fungicide every 3 weeks since May. (a) Why might the disease still be spreading despite spraying? (b) Propose an IPM strategy to reduce black pod incidence for the next season, listing at least four cultural/biological measures and one chemical adjustment.

SOLUTION Exercise 3

(a) Possible reasons: (i) poor spray coverage (dense canopy, no pruning); (ii) wrong timing (spraying after infection established); (iii) resistance development (same mode of action used repeatedly); (iv) high inoculum from mummified pods on trees/ground; (v) excessive shade/humidity.

(b) IPM strategy for next season:

1. Sanitation: complete removal and burial/burning of all mummified pods (source of inoculum) before flowering.
2. Pruning: structural pruning after main harvest to open canopy, improve air flow and spray penetration.
3. Shade management: thin permanent shade trees to reduce humidity below 80% in canopy.
4. Resistant clones: gradually replant gaps with SNKA 12/13/14 (black pod tolerant).

5. Chemical adjustment: rotate copper (M1) with fosetyl-Al (33) or metalaxyl-M (4) + mancozeb (M3) to manage resistance; apply at pod set (first cherelles) and repeat at 3-week intervals during high humidity; use motorised mist-blower for better coverage.

4 ANIMAL NUTRITION AND ANIMAL PRODUCTION

In the previous section we studied how crops are cultivated to feed people and to supply raw materials. But crops also feed animals, and animals in turn supply meat, milk, eggs, hides, draught power and manure. From the cattle herds of the Adamawa Plateau to the poultry farms around Bafoussam and the pigsties of the West Region, animal production is a pillar of the Cameroonian rural economy. Just as a crop only yields well when the soil supplies the right nutrients, a farm animal only produces well when its diet supplies the right nutrients in the right proportions. This section therefore answers two linked questions: what do farm animals need to eat? and how do we organise their production efficiently and profitably?

4.1 Classes of Nutrients Required by Farm Animals

Observe a goat tethered by the roadside in Bamenda. It eats grass all day, drinks water, and yet grows, produces milk and kids once a year. Hidden inside that grass are six classes of **nutrients**, each with a distinct role.

DEFINITION Nutrients required by farm animals

A **nutrient** is any chemical substance in feed that is used by the animal's body for energy, growth, repair or regulation. Farm animals require six classes:

- Carbohydrates: the main **energy** source (starches, sugars, cellulose). Energy fuels movement, body heat, milk synthesis and egg formation. Excess is stored as fat.
- Proteins: supply **amino acids** for growth of muscle, wool, feathers, milk and eggs, and for enzymes and hormones. Essential for young, pregnant and lactating animals. Ten amino acids are essential for non-ruminants; ruminants can synthesise them via rumen microbes if nitrogen is available.
- Lipids (fats and oils): concentrated energy (about 2.25 times the energy of carbohydrates per gram), carriers of fat-soluble vitamins (A, D, E, K), and sources of essential fatty acids (linoleic, linolenic).
- Vitamins: needed in tiny amounts for regulation; fat-soluble (A, D, E, K) and water-soluble (B-complex, C). Examples: vitamin A (vision, reproduction, epithelial integrity), vitamin D (calcium/phosphorus metabolism, bone formation), vitamin E (antioxidant), B-vitamins (coenzymes in metabolism).

- **Minerals:** macro-minerals (calcium, phosphorus, sodium, chlorine, magnesium, potassium, sulphur) and micro-minerals (iron, copper, zinc, manganese, iodine, cobalt, selenium). Calcium and phosphorus for bones, eggshells and milk; sodium and chlorine (common salt) for osmotic balance; iron and copper for haemoglobin; iodine for thyroid hormones.
- **Water:** the most abundant and most neglected nutrient. A lactating cow may drink 40–60 litres per day; without water, feed intake and production collapse within hours. Water regulates temperature, transports nutrients, and participates in hydrolysis reactions.

WARNING Water is a nutrient

Many farmers in the dry season of the Far North provide feed but forget water. A cow can survive weeks without feed but only a few days without water. Always provide clean water *ad libitum* (freely available). Water quality matters: high salinity or microbial contamination reduces intake and causes disease.

4.2 Feedstuffs: Roughages, Concentrates, Supplements and Additives

Feeds are grouped according to their fibre content and nutrient density.

DEFINITION Roughage and concentrate

A **roughage** (or forage) is a bulky feed with high crude fibre content (more than about 18% of dry matter) and low digestible energy per kilogram: grasses (e.g. *Panicum maximum*, *Brachiaria*), legume fodder (e.g. *Stylosanthes*, *Calliandra*, *Leucaena*), hay (dried forage) and silage (forage fermented and stored in the absence of air). A **concentrate** is a feed with low fibre and high nutrient density, fed in small quantities: energy concentrates (maize, sorghum, cassava flour, wheat bran, rice bran) and protein concentrates (cottonseed cake, groundnut cake, soya bean meal, fish meal, bone meal, blood meal, brewer's dried grains).

DEFINITION Ration and balanced diet

A **ration** is the total quantity of feed offered to an animal in 24 hours. A **balanced ration** (balanced diet) is a ration that supplies all the nutrients required by the animal, in the correct amounts and proportions, for its physiological state (maintenance, growth, pregnancy, lactation or egg production), at minimum cost. It must be palatable, non-toxic, and provide adequate dry matter intake.

Supplements and additives. A **supplement** corrects a specific deficiency: mineral licks (salt, calcium, phosphorus) for grazing cattle, or bone meal / oyster shell added to poultry feed for eggshell formation. **Additives** are non-nutritive substances added in small amounts: vitamin premixes, antibiotics or coccidiostats (disease prevention), enzymes (phytase, xylanase) and growth promoters. Their misuse (e.g. antibiotic residues in meat, antimicrobial resistance) is a public health concern; withdrawal periods must be respected.

DID YOU KNOW? Local feed resources in Cameroon


Cameroon has abundant agro-industrial by-products for animal feed: brewer's dried grains (from breweries in Douala and Yaoundé), cottonseed cake (from SODECOTON in the North), palm kernel cake (from palm oil mills in the South-West and Littoral), wheat bran (from flour mills), and cassava peels (from processing units). Research at IRAD (Institute of Agricultural Research for Development) evaluates their nutritive value and inclusion levels.

4.3 Digestion in Ruminants versus Non-Ruminants

Why can a cow thrive on grass alone while a pig or a chicken cannot? The answer lies in the stomach.

PROPERTY Microbial fermentation in the rumen

Ruminants (cattle, goats, sheep) possess a four-chambered stomach. The largest chamber, the **rumen**, hosts billions of micro-organisms (bacteria, protozoa, fungi) which ferment cellulose and other fibres, producing **volatile fatty acids** (VFAs: acetate, propionate, butyrate) that the animal absorbs through the rumen wall and uses for energy. The microbes also synthesise microbial protein (from non-protein nitrogen like urea) and B-vitamins, which the animal digests later in the abomasum and small intestine. Non-ruminants (pigs, poultry, humans) lack a functional rumen and cannot digest cellulose efficiently; they therefore require concentrated, low-fibre, high-quality feeds.



The four chambers of the ruminant stomach and the flow of digesta: oesophagus → rumen → reticulum → omasum → abomasum (the “true stomach” with gastric juice) → small intestine.

Feed passes from the oesophagus into the **rumen** and **reticulum** (often called the reticulorumen), where microbes ferment fibre; coarse material is regurgitated and re-chewed (rumination). It then flows to the **omasum** (water absorption, many folds) and the **abomasum**, the true stomach where acid (HCl) and enzymes (pepsin) act as in non-ruminants.

Practical implications. Cattle and goats can be raised on pasture and crop residues (maize stover, groundnut haulms) with only small concentrate supplements. Pigs and poultry must be fed concentrates rich in energy and good-quality protein, because they cannot extract enough from fibre.

WARNING Do not overfeed concentrates to ruminants

A common and costly mistake: suddenly giving cattle or goats large amounts of maize or bran. Rapid fermentation produces excess gas trapped in the rumen (**bloat**) and excess lactic acid (**ruminal acidosis**), both potentially fatal. Always introduce concentrates gradually (over 2–3 weeks) and ensure roughage forms at least about 40% of the ration dry matter to keep the rumen functioning.

4.4 Nutritional Requirements by Physiological State

An animal's needs change with what its body is doing. Every ration is built in two layers: a **maintenance ration** (to keep the animal alive, warm and moving without gaining or losing weight) plus a **production ration** added on top.

Physiological state	Extra requirements above maintenance
Maintenance (adult, idle)	Energy and protein just to maintain body weight and temperature
Growth (young animals)	High protein (essential amino acids) and minerals (Ca, P) for muscle and bone
Pregnancy (last third)	Extra energy, protein and minerals for the foetus; rapid foetal growth in last trimester
Lactation	Very high demand: milk exports protein, fat, lactose, calcium and water daily; peak yield at 4–8 weeks post-partum
Egg production (layers)	High calcium (shells $\approx$ 2 g Ca/egg), protein and energy; about 110–120 g feed/bird/day for modern hybrids

EXAMPLE Nutrient partitioning in a dairy cow

A 500 kg Holstein cow producing 25 kg milk/day (3.5% fat, 3.2% protein) requires approximately: 160 MJ metabolisable energy, 1.8 kg crude protein, 90 g calcium, 65 g phosphorus. Maintenance alone needs 60 MJ ME and 600 g CP. The production ration must supply the difference. Underfeeding in early lactation causes excessive body condition loss and ketosis.

4.5 Ration Formulation: the Pearson Square

To formulate a balanced ration we compare the **dry matter** (DM, feed minus water), **crude protein** (CP) and energy content of available feeds, then mix them in calculated proportions. The simplest tool is the **Pearson square**, used to blend two feeds to a target protein percentage. It assumes linear mixing and is valid for one nutrient (usually CP) at a time; energy and minerals must be checked separately.

METHOD Using the Pearson square

1. Write the target CP percentage (on dry matter basis) in the centre of a square.
2. Write the CP of the two feeds at the left corners (high-protein feed top, low-protein feed bottom).
3. Subtract diagonally (larger minus smaller) and write the differences at the right corners.
4. The right-hand values give the parts of each feed in the mixture: the value opposite a feed is the proportion of that feed.
5. Convert parts to percentages of the total.
6. Verify by calculating the weighted average CP of the mixture.

WORKED EXAMPLE 5 Formulating a 16% CP broiler ration

A farmer near Dschang wants to mix maize (9% CP) and soya bean meal (44% CP) to obtain a 16% CP ration. In what proportions should he mix them?


SOLUTION

Pearson square: parts of maize = $44 - 16 = 28$; parts of soya bean meal = $16 - 9 = 7$. Total parts = $28 + 7 = 35$.

$$\text{Maize} = \frac{28}{35} \times 100 = 80\%$$

$$\text{Soya bean meal} = \frac{7}{35} \times 100 = 20\%$$

Check: $(80 \times 9 + 20 \times 44) / 100 = (720 + 880) / 100 = 16\%$ CP. The farmer mixes 80 kg of maize with 20 kg of soya bean meal for every 100 kg of feed (then adds minerals and vitamins).



Subtract diagonally: $44 - 16 = 28$; $16 - 9 = 7$

Pearson square for a 16% crude protein ration: 7 parts soya bean meal to 28 parts maize, i.e. 20% soya and 80% maize.

WORKED EXAMPLE 6 Formulating a 17% CP dairy supplement with three ingredients

A farmer wants a 17% CP supplement using cottonseed cake (41% CP), maize (9% CP) and wheat bran (15% CP). Use Pearson square twice: first combine maize and wheat bran to make a 12% CP energy mix, then combine that with cottonseed cake.

SOLUTION

Step 1: Maize (9%) + Wheat bran (15%) → target 12%. Parts: maize = 15–12 = 3; wheat bran = 12–9 = 3. Mix = 50% maize, 50% wheat bran. CP = 12%. Step 2: Energy mix (12%) + Cottonseed cake (41%) → target 17%. Parts: energy mix = 41–17 = 24; cottonseed cake = 17–12 = 5. Total = 29. Final mix: Energy mix = 24/29 ≈ 82.8%; Cottonseed cake = 5/29 ≈ 17.2%. Within energy mix: Maize = 41.4%, Wheat bran = 41.4%, Cottonseed cake = 17.2%. Check: $(41.4 \times 9 + 41.4 \times 15 + 17.2 \times 41)/100 = 17\%$.

4.6 Systems of Animal Production

System	Features	Advantages / limitations in Cameroon
Extensive	Free range, pastoralism, transhumance; animals find their own feed; low inputs; indigenous breeds	Low cost; uses marginal land; but low productivity, high disease exposure, farmer–grazer conflicts (North West, Adamawa), seasonal feed shortage
Semi-intensive	Animals graze by day, housed at night; some supplementary feeding and health care; crossbred or improved breeds	Good compromise for village poultry and goats; moderate cost and output; manure collected for crops
Intensive	Animals permanently housed; all feed, water and health care provided; high stocking density; exotic/hybrid breeds	High output per animal (broilers, layers, pigs); efficient feed conversion; but high capital, feed cost (70–80% of total cost) and disease risk; requires reliable electricity and water

DID YOU KNOW? Transhumance in the Adamawa

The Adamawa Plateau hosts the largest cattle population in Cameroon. Pastoralists (Mbororo) practice transhumance: moving herds between dry-season lowlands (river valleys) and wet-season highlands. This traditional system is under pressure from crop expansion, climate variability, and land tenure insecurity. Improved corridors and water points are policy priorities.


4.7 Housing and Equipment

Good housing protects animals from predators, rain, heat and thieves, and makes feeding, cleaning and disease control easy.

- **Poultry house:** orientation east–west to avoid direct sun; open sides with wire mesh for **ventilation**; deep litter (wood shavings, sawdust, rice husks) kept dry; density about 8–10 broilers or 4–5 layers per m²; perches (for layers), nest boxes (1 per 4–5 hens), feeders and drinkers (1 per 50 birds for tube feeders, 1 per 80 for bell drinkers).
- **Pigsty:** concrete floor sloping to a drain for easy washing; separate pens for boar, pregnant sows, farrowing sows (with a creep area protecting piglets from crushing) and fatteners; shade and wallow or sprinklers, because pigs cannot sweat and are heat-stressed above 25°C.
- **Cattle shed:** roofed, well-drained floor (earth or concrete with grooves), feeding trough and water trough, milking area; space about 3.5–4 m² per adult cow; calf pens separate.

PROPERTY Hygiene and ventilation

Damp, dirty, poorly ventilated housing concentrates ammonia and pathogens and is the single greatest cause of respiratory disease and parasite build-up in intensive systems. Dry litter, regular removal of dung, disinfection and free air movement are non-negotiable. (Not examinable as a proof, but frequently examined as short-answer reasoning.)



Plan of a poultry house for 500 broilers: east-west orientation, wire-mesh ventilation on both long sides, evenly spaced feeders and drinkers, door with footbath. Deep litter floor (wood shavings).

4.8 Breeding and Reproduction

Improvement begins with **selection** of breeding stock: choose parents with good growth rate, milk or egg yield, disease resistance and body conformation, from reliable records. **Mating systems** include natural mating (one bull or boar for a group of females) and **artificial insemination** (AI), in which semen from a superior male is collected, stored (fresh or frozen) and deposited in the female by a technician; AI spreads superior genes cheaply and avoids keeping dangerous males. **Crossbreeding** (e.g. Holstein $\times$ Gudali for dairy, Large White $\times$ local pig for growth) exploits heterosis (hybrid vigour).

Gestation lengths to remember: cow about 283 days, goat and ewe about 150 days, sow about 114 days ("3 months, 3 weeks, 3 days"), hen incubation 21 days. Care of the young includes colostrum (first milk, rich in antibodies) within the first 6 hours, warmth (brooder for chicks, heat lamp for piglets), and creep feed for early weaning.

EXAMPLE Artificial insemination in Cameroon

AI is promoted by MINEPIA (Ministry of Livestock, Fisheries and Animal Industries) for dairy improvement. The Centre for Artificial Insemination (CIAC) in Ngaoundéré produces and

distributes semen from improved dairy breeds (Holstein, Montbéliarde) and beef breeds. Farmers in the West and North West use AI to upgrade local cattle without keeping a bull.

4.9 Animal Health: Diseases, Parasites and Prevention

Animal	Common diseases / parasites	Prevention
Poultry	Newcastle disease, fowl pox, coccidiosis, Gumboro (IBD); lice, worms (Ascaridia)	Vaccination (Newcastle: Hitchner B1 at day 7, La Sota at day 21; Gumboro at day 14), coccidiostats in feed, clean dry litter, deworming
Cattle	Trypanosomiasis (tsetse fly), tick-borne diseases (anaplasmosis, babesiosis, heartwater), foot-and-mouth disease (FMD); ticks, worms, liver fluke	Trypanocidal drugs (prophylactic/isometamidium), dipping/spraying against ticks (acaricides), strategic deworming (albendazole, ivermectin), FMD vaccination in risk zones
Pigs	African swine fever (ASF), worms, mange (Sarcoptes)	Strict biosecurity (fencing, no swill feeding without boiling 30 min), deworming, acaricide spray for mange; no vaccine for ASF
Goats/sheep	Peste des petits ruminants (PPR), worms (Haemonchus), mange, foot rot	PPR vaccination (at 3 months, then annually), deworming (FAMACHA-guided for Haemonchus), hoof trimming, clean pens

Prevention rests on three pillars: **vaccination** (following a calendar, e.g. Newcastle vaccine in chicks), regular **deworming** and tick control, and **biosecurity** (quarantine of new animals for 21 days, footbaths at entrances, restricting visitors, isolating sick animals, proper carcass disposal).

WARNING Exam tip

GCE questions on animal production very often ask for a labelled housing plan (poultry house or pigsty) or a vaccination schedule presented as a table. Practise drawing a simple, clearly labelled plan and tabulating: age of animal – vaccine – route of administration (e.g. eye drop, drinking water, intramuscular).

4.10 Records, Economics and Improvement

Without records, a farmer cannot know whether the enterprise is making a profit. Essential **farm records** include: inventory records (stock of animals, feed, equipment), production records (milk per cow per day, eggs per day per hen house), breeding records (service dates, birth dates,

parentage), health records (treatments, vaccinations, mortality), and financial records (variable costs: feed, drugs, vaccines; fixed costs: housing, equipment; revenue: sales of animals, products, manure).

Key **production indices**:

$$\text{Average daily gain (ADG)} = \frac{\text{final weight} - \text{initial weight}}{\text{number of days}} \quad (\text{kg/day or g/day})$$

$$\text{Feed conversion ratio (FCR)} = \frac{\text{feed consumed (kg)}}{\text{weight gained (kg)}} \quad (\text{dimensionless; lower is better})$$

$$\text{Egg production (\%)} = \frac{\text{eggs collected per day}}{\text{number of hens present}} \times 100$$

$$\text{Mortality rate (\%)} = \frac{\text{number of deaths}}{\text{number at start}} \times 100$$

WORKED EXAMPLE 7 Calculating ADG and FCR for broilers


A batch of broilers weighs 0.2 kg at 2 weeks and 2.0 kg at 8 weeks (42 days between weighings). Over this period each bird consumed 3.36 kg of feed. Calculate the ADG and the FCR.

SOLUTION

$$\text{ADG} = \frac{2.0 - 0.2}{42} = \frac{1.8}{42} \approx 0.043 \text{ kg/day} = 43 \text{ g/day}$$

$$\text{FCR} = \frac{3.36}{1.8} \approx 1.87$$

An FCR of 1.87 means 1.87 kg of feed produced 1 kg of live weight gain — a good performance for broilers (lower FCR = better efficiency). Target for modern broilers: FCR < 1.7 at 35 days.



Growth curves of broiler chickens: live weight versus age. A balanced ration (high crude protein and energy) gives faster growth and earlier market weight; poor feed delays slaughter age and raises costs.

Marketing. Profit depends as much on selling as on producing. Farmers should time sales to periods of high demand (end-of-year festivities, Ramadan for sheep and goats, Easter for poultry), add value where possible (eggs graded and packed, chickens dressed and chilled), and keep transport costs low by grouping sales through cooperatives or farmer associations. Knowledge of live weight vs. carcass weight (dressing percentage: broilers $\approx 70\%$, pigs $\approx 75\%$, cattle $\approx 50\%$) helps negotiate prices.

KEY POINTS Key points

- Six nutrient classes: carbohydrates, proteins, lipids, vitamins, minerals, water.
- Roughages (fibre-rich) versus concentrates (nutrient-dense); ruminants digest fibre thanks to rumen microbes, non-ruminants cannot.
- Rations are built for maintenance plus production; the Pearson square balances two feeds to a target protein level (check energy and minerals separately).
- Production systems range from extensive (pastoralism) to intensive (housed, high input); each has trade-offs in Cameroon.
- Success rests on housing hygiene, selection and breeding (AI, crossbreeding), vaccination and deworming, and record-keeping with indices such as ADG, FCR and egg production percentage.

EXERCISE 7 Practice

1. A farmer mixes cottonseed cake (41% CP) with maize (9% CP) to obtain a 17% CP supplement for dairy cows. Using the Pearson square, find the proportions of each feed.
2. A layer flock of 200 hens produces 150 eggs per day. Calculate the percentage egg production and comment.
3. Draw a labelled plan of a poultry house for 500 broilers, indicating orientation, ventilation openings, feeders and drinkers.
4. Explain why a sudden change from pasture to a maize-rich ration can kill a goat.
5. A piglet weighs 1.5 kg at weaning (28 days) and 30 kg at 112 days. It consumed 85 kg of feed over this period. Calculate ADG and FCR for the growing-finishing phase.

SOLUTION Exercise 1

1. Pearson square: parts of cottonseed cake = $17 - 9 = 8$; parts of maize = $41 - 17 = 24$; total = 32. Cottonseed cake = $8/32 \times 100 = 25\%$; maize = $24/32 \times 100 = 75\%$. Check: $(25 \times 41 + 75 \times 9)/100 = (1025 + 675)/100 = 17\%$.
2. Egg production = $150/200 \times 100 = 75\%$. This is a good laying rate for a well-managed flock at peak production (target > 80% for modern hybrids at peak).
3. The plan should show: east-west orientation, wire-mesh side walls for ventilation, roof overhang, deep litter floor, feeders and drinkers distributed evenly, a door and a footbath at the entrance; stocking density about 10 birds/m² gives 50 m² of floor space (e.g. 10 × 5 m).
4. Sudden excess of maize causes rapid microbial fermentation in the rumen: gas accumulates (bloat) and lactic acid lowers rumen pH (acidosis), damaging the rumen wall and killing rumen microbes; both conditions can be fatal. Concentrates must be introduced gradually with adequate roughage.
5. $ADG = (30 - 1.5)/(112 - 28) = 28.5/84 \approx 0.339 \text{ kg/day} = 339 \text{ g/day}$. $FCR = 85/28.5 \approx 2.98$. This FCR is acceptable for growing-finishing pigs (target < 3.0).

5 INTEGRATION ACTIVITY

6 AGRICULTURE

6.1 Integration Activity: The Nkeng Family Mixed Farm Project

KEY POINTS Aims of the Activity

This integration activity brings together all concepts from Chapter BIO08 (Lessons 88–92). By the end, you should be able to:

- assess the properties of a given soil and propose scientifically justified measures to improve its **fertility**;
- design a rational **land-use plan** including **crop rotation** and soil conservation on sloping land;
- construct a **cropping calendar** adapted to the bimodal rainfall regime of the Cameroonian highlands;
- apply the **Pearson square** method to formulate a balanced poultry ration from local feedstuffs;
- plan **housing**, a **vaccination** and **deworming** schedule, and a **record book** for small livestock;
- integrate plant and animal production into a sustainable **mixed farming** system and defend your choices orally.

WARNING Skills Assessed

This activity mobilises the GCE Advanced Level competences of analysing data, applying knowledge to a novel situation, carrying out calculations and communicating scientific arguments. These are exactly the skills tested in Paper 2 (structured questions) and Paper 3 (practical/problem-solving).

6.2 Situation

EXPERIMENT The Nkeng Family Farm Project

The Nkeng family lives near Dschang in the West Region of Cameroon, at an altitude of about 1400 m. The climate is of the highland type, with a long rainy season from mid-March to mid-November (peak rainfall July–September) and a dry season from mid-November to mid-March. Mean temperatures range from 18 °C to 24 °C.

The family owns 2 hectares of gently sloping land. A soil survey and simple field tests gave the following results:

Property	Observation
Texture	Lateritic sandy loam; reddish colour
pH (water)	5.2 (acidic)
Organic matter	Low; topsoil only about 10 cm deep
Drainage	Good on the slope; a small marshy patch at the bottom
Erosion	Rills visible after heavy storms on the upper slope

The family wishes to produce, for home consumption and for sale at the Dschang market:

- maize and cassava (staple food crops);
- vegetables (cabbage, carrots, tomatoes) on the more fertile lowland patch;
- 100 broilers and 50 layers per production cycle;
- 2 dairy goats for milk.

Available local feedstuffs and their approximate crude protein (CP) contents:

Feedstuff	Crude protein (%)
Maize grain (energy feed)	9
Cottonseed cake (protein concentrate)	41

A broiler finisher ration should contain about 20 % CP. The family has access to poultry manure, kitchen waste, and can grow *Calliandra* and *leucaena* as fodder shrubs along the boundaries.

DID YOU KNOW? Cameroon Context

The West Region (Dschang, Bafoussam) is a major agricultural zone with volcanic and lateritic soils. The bimodal rainfall pattern (though here described as a long single season at 1400 m) allows two maize crops per year. *Calliandra calothyrsus* and *Leucaena leucocephala* are widely promoted by IRAD and NGOs for alley cropping and fodder banks because they fix nitrogen, tolerate acid soils, and provide high-protein browse for ruminants.

6.3 Tasks

Work in groups of four to six. Answer the following tasks in order; each one builds on the previous.

1. **Soil assessment.** From the soil data, identify **two strengths** and **three limitations** of this soil for crop production. For each limitation, propose one corrective measure and justify it biologically or chemically.
2. **Land-use plan.** Draw a sketch plan of the 2 ha farm, dividing it into at least four plots. Position the crops, the poultry house, the goat shed and the fodder shrubs. Justify the position of each element (think of slope, drainage, wind, hygiene and labour).
3. **Crop rotation.** Propose a **three-year rotation** for the main upland plot, including a legume. Explain two advantages of your rotation.
4. **Cropping calendar.** Using the rainfall data, construct a seasonal calendar (table) showing land preparation, planting, weeding/fertiliser application and harvesting for maize, cassava and one vegetable.
5. **Feed formulation.** Using the **Pearson square**, calculate the proportions of maize grain and cottonseed cake needed to prepare 100 kg of a broiler finisher ration at 20 % CP. Show all working.
6. **Animal husbandry plan.** For the poultry and the goats, propose:
 - (a) one housing requirement and the reason for it;
 - (b) a simple health calendar (vaccination of chicks against Newcastle disease; deworming of goats at the start of the rains);
 - (c) three records the family should keep, and why.
7. **Integration and sustainability.** Show with a simple flow diagram how wastes and products circulate between the crop and animal components (e.g. manure → crops; crop residues → animals). State one reason why this integration improves sustainability.
8. **Presentation.** Prepare a 10-minute group presentation defending your plan. The class will vote for the most **sustainable and profitable** design.

6.4 Model Answers

Task 1 — Soil Assessment.


Strengths:

1. The sandy loam texture is **well aerated** and easy to work, suitable for root crops like cassava which require loose soil for tuber expansion.
2. Drainage is good on the slope, preventing waterlogging of maize roots (maize is sensitive to anaerobic conditions).

Limitations and Corrections:

- **Acidity (pH 5.2):** At this pH, aluminium and manganese become soluble and toxic to roots, while phosphorus availability is sharply reduced due to fixation by iron/aluminium oxides. **Correction:** Apply agricultural lime (CaCO_3) at a rate determined by a lime requirement test (typically 2 t/ha to 4 t/ha for such soils) to raise the pH towards 6.0–6.5. Liming also supplies Ca^{2+} and Mg^{2+} , improves microbial activity, and enhances P availability.
- **Low organic matter:** Poor soil structure, low water-holding capacity, low cation exchange capacity (CEC), and limited nutrient supply. **Correction:** Incorporate compost, farmyard manure from the poultry and goats, and practise green manuring with a legume (e.g., *Mucuna* or *Crotalaria*) during the rotation. This builds humus, improves aggregation, and increases CEC.
- **Erosion on the slope:** Loss of the thin fertile topsoil (only 10 cm deep) by surface run-off during heavy storms. **Correction:** Construct contour ridges (tied ridges) and plant grass strips (*Pennisetum purpureum*) or fodder shrubs (*Calliandra*) along contour lines to slow run-off, increase infiltration, and trap sediment.

Task 2 – Land-Use Plan.



Model land-use plan for the 2 ha mixed farm near Dschang (not to scale).

Justifications:

- **Buildings (poultry house, goat shed, compost pit)** are placed on well-drained ground (mid-slope, away from the marshy patch) near the homestead for easy surveillance, security, and to minimise labour for carrying feed and manure. The east–west orientation reduces direct solar radiation on the long walls, preventing overheating. They are positioned downwind of the vegetable plot (prevailing wind from the west in this region) to avoid contamination of crops by dust and odours.
- **Vegetables (Plot C)** occupy the moist, relatively fertile lowland patch where water is accessible for irrigation in the dry season. Raised beds improve drainage for cabbage and carrots.
- **Maize/Legume rotation (Plot A, upper slope)** benefits from contour ridges to control erosion on the steepest part.

- **Cassava + fodder shrubs (Plot B, mid slope):** Cassava tolerates poorer soils and drought; Calliandra hedgerows on the contour serve simultaneously as **erosion barriers** and as **protein-rich fodder** for the goats (cut-and-carry system).

Task 3 — Three-Year Rotation (Plot A).

Year 1: Maize (heavy feeder, high N demand) → **Year 2:** Groundnuts (*Arachis hypogaea*) or cow-pea (*Vigna unguiculata*) (legume, N-fixing) → **Year 3:** Cassava (tolerates lower fertility, deep roots) or a green manure fallow (*Mucuna pruriens*).

Advantages:

1. The legume fixes atmospheric nitrogen through *Rhizobium* bacteria in its root nodules, adding 50–100 kg N ha⁻¹ to the system and reducing the need for mineral nitrogen fertiliser for the subsequent crop.
2. Rotating crops with different botanical families, rooting depths, and pest hosts breaks **pest and disease cycles** (e.g., maize streak virus, cassava mosaic virus, nematodes) and exploits nutrients from different soil layers (shallow maize roots vs. deep cassava roots).

Task 4 — Cropping Calendar.

Crop	Land Preparation	Planting	Maintenance / Harvest
Maize (1st season)	Feb – early Mar: clearing, burning (if needed), contour ridging, basal fertiliser (NPK 20–10–10)	Mid-Mar (onset of rains): 2 seeds/hole, 80×40 cm	Weed at 3 and 6 weeks; top-dress urea at 6 weeks (50 kg/ha); harvest Jul–Aug (green maize) or Aug–Sep (dry grain)
Maize (2nd season)	Late Aug: minimal tillage on residues	Early Sep	Weed + top-dress; harvest Dec–Jan
Cassava	Feb–Mar: mounds or ridges on contour	Mar–Apr: stem cuttings 25 cm, 1×1 m	Weed at 1, 3, 6 months; harvest 10–12 months later (Jan–Mar Year 2)
Cabbage (Plot C)	Mar: raised beds, incorporate poultry manure (10 t/ha)	Nursery: Mar; Transplant: Apr (45×45 cm)	Regular watering; side-dress urea at 3 weeks; harvest Jun–Jul. Dry-season crop: nursery Oct, transplant Nov, harvest Feb–Mar (with irrigation)

Task 5 — Pearson Square for Broiler Finisher Ration.

METHOD Applying the Pearson Square

1. Draw a square. Write the CP content of each feedstuff at the left corners (Maize = 9, Cottonseed cake = 41) and the target CP (20) in the centre.
2. Subtract diagonally (always larger minus smaller):
 - $41 - 20 = 21$ parts of maize (energy feed)
 - $20 - 9 = 11$ parts of cottonseed cake (protein concentrate)
3. Total parts = $21 + 11 = 32$.
4. Calculate the weight of each ingredient for 100 kg of ration:

$$\begin{aligned}\text{Maize} &= \frac{21}{32} \times 100 \text{ kg} = 65.625 \text{ kg} \approx 65.6 \text{ kg} \\ \text{Cottonseed cake} &= \frac{11}{32} \times 100 \text{ kg} = 34.375 \text{ kg} \approx 34.4 \text{ kg}\end{aligned}$$

Verification:

$$(65.6 \text{ kg} \times 0.09) + (34.4 \text{ kg} \times 0.41) = 5.904 \text{ kg} + 14.104 \text{ kg} = 20.008 \text{ kg CP per 100 kg ration} = 20.0 \% \text{ CP. } \checkmark$$

WARNING Practical Note


This ration provides energy and protein only. A complete commercial broiler finisher ration must also include: a vitamin–mineral premix (0.3 % to 0.5 %), salt (0.25 %), lysine/methionine if needed, and a coccidiostat. The cottonseed cake must be from a source with low gossypol (or treated) to avoid toxicity.

Task 6 — Animal Husbandry Plan.

- **Housing:**
 - Poultry: The house must be **well ventilated but draught-free**, with a deep litter of dry wood shavings or rice husks (10 cm deep) and an east–west orientation. **Reason:** Damp litter favours coccidiosis (sporulation of oocysts) and respiratory diseases (ammonia from uric acid decomposition); good ventilation removes moisture and ammonia while preventing cold draughts that stress chicks.
 - Goats: The shed must be **raised 60 cm above ground** with a slatted floor (bamboo or timber, gaps 1.5 cm) and a roof overhang of 50 cm. **Reason:** Droppings fall through, keeping the floor dry and clean; this drastically reduces re-infection by gastrointestinal nematodes (worm larvae develop in moist faeces on the ground) and prevents foot rot.
- **Health Calendar:**

- Poultry (Newcastle disease): Vaccinate day-old chicks (intra-ocular or spray), booster at 4 weeks (drinking water or eye drop), then every 3–4 months for layers. Use a thermostable vaccine (e.g., I-2) if cold chain is unreliable.
 - Goats (Strategic deworming): Treat with a broad-spectrum anthelmintic (e.g., albendazole or levamisole) at the **start of the rains** (March) when infective larvae on pasture surge, and again at **mid-rains** (July). Rotate drug classes annually to delay resistance. Check faecal egg counts if possible.
 - General: Isolate and treat any sick animal immediately; disinfect houses between batches; control rodents and wild birds.
- **Records (three essential ones):**
 1. **Production records:** Daily egg count per layer batch; weekly broiler weights; daily milk yield per goat. Why: Early detection of disease, nutritional deficiency, or management problems; basis for culling decisions.
 2. **Health records:** Dates of vaccination, deworming, treatments (drug, dose, withdrawal period), mortalities with suspected cause. Why: Ensures timely boosters, avoids drug residues in meat/milk, tracks disease patterns.
 3. **Financial records:** Costs of feed, chicks, drugs, equipment; revenue from sales of eggs, broilers, milk, manure, crops. Why: Calculates gross margin, identifies profitable enterprises, supports loan applications.

Task 7 — Integration Diagram.



Nutrient and energy flows in the Nkeng family mixed farming system.

Sustainability Statement: Recycling manure and compost as fertiliser reduces the need to purchase mineral fertiliser, maintains soil organic matter and cation exchange capacity, and closes the nutrient loop. This makes the farm productive without degrading the soil resource, embodying the principle of **sustainable intensification**.

EXERCISE 8 Extension Task

The family later considers adding a fish pond on the marshy patch. Suggest one advantage and one risk of this addition, and state how pond mud could be used in the farm's nutrient cycle.

SOLUTION Extension Task

Advantage: The pond converts an unproductive marshy patch into a source of high-quality animal protein (fish, e.g., *Oreochromis niloticus*) and income. Pond water can also be used for supplementary irrigation of vegetables during dry spells, increasing crop yields.

Risk: Stagnant water can favour breeding of *Anopheles* mosquitoes (malaria vectors) and *Simulium* (onchocerciasis); predators (birds, snakes, monitor lizards) may cause losses; poor embankment construction can lead to flooding and escape of fish. Proper design (deep water, steep banks, predator nets) and management (larvivorous fish, regular monitoring) are essential.

Pond mud: During pond draining (every 1–2 years), the accumulated mud is rich in nitrogen, phosphorus, and organic matter from fish waste and uneaten feed. It can be spread on vegetable beds (Plot C) or composted with crop residues as an excellent organic fertiliser, further closing the nutrient cycle and reducing external input dependence.

7 METHODS AND WORKED EXAMPLES

7.1 Method 1: Identifying a Soil Type from its Physical Properties

METHOD How to Identify and Classify a Soil

To determine the nature of a soil sample in the field or in the laboratory, proceed in order:

1. Observe the colour: dark soils are usually rich in **humus** (decomposed organic matter); red or yellow colours indicate iron oxides (**lateritic soils**, very common in the forest zone of Cameroon); grey-blue colours suggest poor drainage and waterlogging (**gley soils**).
2. Feel the texture (the “sausage test”): moisten a small sample and try to roll it into a cylinder. If it cannot be rolled, the soil is **sandy**; if it rolls but cracks, it is **loamy**; if it rolls into a long flexible sausage that can be bent into a ring without breaking, it is **clayey**.
3. Estimate particle sizes by sedimentation: shake the soil thoroughly with water in a measuring cylinder and leave it to settle. Sand settles first (bottom layer), then silt, then clay (top mineral layer); humus floats or remains suspended. Measure the thickness of each layer and express it as a percentage of the total mineral depth.
4. Test drainage and water retention: pour a known volume of water through a known mass of soil and measure the volume collected. Sandy soils drain fast and retain little water; clay soils retain much water but may become waterlogged and poorly aerated.
5. Conclude by matching your results to the properties of the main soil types: sandy, clay, loamy, humus-rich, lateritic, alluvial. A **loam** (a roughly balanced mixture of sand, silt and clay, with humus) is generally the best agricultural soil.

Reflex: always link the soil type to its consequences for agriculture: aeration, drainage, water retention, ease of cultivation (tilth), and suitability for particular crops.

WORKED EXAMPLE 8 Identifying Soil Samples from a School Farm

A student in Buea collected three soil samples (A, B, C). The results of simple tests are summarised below:

Test	Sample A	Sample B	Sample C
Sausage test	Cannot be rolled	Rolls into a long flexible ring	Rolls but cracks easily
Water drained through 100 g of soil (out of 50 mL poured)	45 mL	5 mL	25 mL
Colour	Pale yellow	Reddish	Dark brown

Identify each soil type and state which is most suitable for growing maize, justifying your answer.

SOLUTION

Step 1 — Analyse Sample A. It cannot be rolled into a sausage, so its particles do not stick together: they are coarse. 45 mL out of 50 mL of water drained through, i.e. very rapid drainage and poor water retention. The pale colour indicates little humus. Sample A is a sandy soil.

Step 2 — Analyse Sample B. It forms a long flexible ring, showing very fine, sticky particles. Only 5 mL of water passed through: drainage is very poor and water retention is very high, so the soil risks waterlogging and poor root aeration. The reddish colour indicates iron oxides. Sample B is a clay soil (lateritic clay).

Step 3 — Analyse Sample C. It rolls but cracks: an intermediate texture, i.e. a balanced mixture of sand, silt and clay. Drainage is moderate (25 mL out of 50 mL), so it retains enough water without waterlogging. The dark brown colour shows it is rich in humus. Sample C is a loamy soil (humus-rich loam).

Step 4 — Conclusion. Sample C (loam) is the most suitable for maize because it combines good drainage and aeration (roots respire easily) with adequate water and nutrient retention; it is easy to till, and its humus releases mineral salts as it decomposes. Sample A would need frequent irrigation and manuring; Sample B would be heavy to work and poorly aerated.

7.2 Method 2: Diagnosing Plant Mineral Deficiencies

METHOD How to Diagnose a Mineral Deficiency in a Crop Plant

1. Locate the symptoms: do they appear first on old (lower) leaves or on young (upper) leaves? Mobile elements (N, P, K, Mg) are withdrawn from old leaves and moved to

young ones when scarce, so their deficiency shows on old leaves first. Immobile elements (Ca, Fe) cannot be relocated, so their deficiency shows on young leaves first.

2. Describe the symptom precisely: uniform yellowing (**chlorosis**), yellowing between green veins (**interveinal chlorosis**), purple coloration, scorched leaf edges, stunted growth, poor roots, etc.
3. Match the symptom to the element:
 - Nitrogen (N): uniform yellowing of old leaves, stunted growth (N is needed for proteins and chlorophyll).
 - Phosphorus (P): dark green or purplish leaves, poor root development, poor seed/fruit formation.
 - Potassium (K): yellowing then scorching (browning) of the margins of old leaves; weak stems.
 - Magnesium (Mg): interveinal chlorosis of old leaves (Mg is the central atom of chlorophyll).
 - Calcium (Ca): death of growing points, distorted young leaves, blossom-end rot of fruits.
 - Iron (Fe): interveinal chlorosis of young leaves.
4. Propose a remedy: apply the appropriate fertiliser (e.g. NPK 20–10–10, urea for N, superphosphate for P, muriate of potash for K), compost or farmyard manure, and correct the soil pH (liming) if necessary.

Reflex: in the exam, always justify the diagnosis with TWO arguments: (i) where the symptom appears and (ii) its precise form.

WORKED EXAMPLE 9 Deficiency in a Tomato Field

A farmer near Bafoussam observes that the older, lower leaves of his tomato plants turn yellow while the veins remain green; the young leaves at the top look normal. The soil is slightly acidic.

- a) Identify the deficient element, justifying your answer.
- b) Suggest two corrective measures.

SOLUTION

a) Diagnosis. Two observations must be combined.

(i) Location of symptoms. The symptoms appear on the old leaves first, while young leaves remain healthy. The element concerned is therefore mobile in the plant: when it becomes scarce, the plant withdraws it from old leaves and transports it to the growing

points. This eliminates calcium and iron, which are immobile and whose deficiency appears on young leaves.

(ii) Form of the symptom. The yellowing is interveinal: the tissue between the veins turns yellow while the veins stay green. This is the classic sign of magnesium deficiency, because magnesium is the central atom of the chlorophyll molecule; without it, chlorophyll cannot be synthesised, while the veins (mostly vascular tissue with little chlorophyll) remain green.

Conclusion: the crop suffers from magnesium (Mg) deficiency.

b) Corrective measures.

- Apply a magnesium-containing fertiliser, e.g. magnesium sulphate (Epsom salt) or dolomitic limestone, which supplies both Mg and Ca.
- Correct the soil acidity by liming: in acidic soils, Mg^{2+} ions are easily leached and their uptake is hindered by excess H^+ ions; adding lime (CaCO_3) raises the pH towards neutrality and improves Mg availability.
- (Also acceptable: adding well-decomposed compost or farmyard manure, which releases Mg and other minerals slowly and improves soil structure.)

7.3 Method 3: Calculating Fertiliser Requirements (NPK)

METHOD How to Calculate the Quantity of Fertiliser to Apply

1. Read the fertiliser grade: a bag labelled NPK 15–15–15 contains 15 % nitrogen (N), 15 % phosphorus pentoxide (P_2O_5) and 15 % potassium oxide (K_2O) by mass.
2. Write the key formula:

$$\text{Mass of fertiliser} = \frac{\text{mass of nutrient required}}{\text{percentage of nutrient in fertiliser}} \times 100$$

3. Scale to the field area: nutrient requirement per hectare $\times$ number of hectares.
4. Check units throughout (kg, ha) and round sensibly to whole bags (usually 50 kg bags).

Reflex: the percentage refers to the nutrient, not the whole fertiliser. Never multiply when you should divide: the fertiliser mass must be larger than the nutrient mass.

WORKED EXAMPLE 10 Fertilising a Maize Field

An agricultural extension officer recommends that a maize crop receives 90 kg of nitro-

gen (N) per hectare. The farmer has two fertilisers available: urea (46 % N) and NPK 20–10–10 (20 % N). His field covers 2.5 ha.

- Calculate the mass of urea needed for the whole field.
- Calculate the mass of NPK 20–10–10 that would be needed to supply the same amount of nitrogen.
- How many 50 kg bags of urea must the farmer buy?

SOLUTION

a) Mass of urea.

Nitrogen required per hectare: 90 kg. For 2.5 ha:

$$90 \times 2.5 = 225 \text{ kg of N required.}$$

Urea contains 46 % N, i.e. 46 kg of N per 100 kg of urea. Therefore:

$$\text{Mass of urea} = \frac{225}{46} \times 100 = 489.1 \text{ kg} \approx 489 \text{ kg of urea.}$$

b) Mass of NPK 20–10–10.

This fertiliser contains only 20 % N:

$$\text{Mass of NPK} = \frac{225}{20} \times 100 = 1125 \text{ kg of NPK 20–10–10.}$$

Comment: much more NPK is needed because it is less concentrated in nitrogen; however, it also supplies phosphorus and potassium, which urea does not. The choice depends on the soil analysis.

c) Number of 50 kg bags of urea.

$$\frac{489.1}{50} = 9.78 \text{ bags.}$$

Since bags cannot be bought in fractions, the farmer must buy 10 bags of 50 kg of urea. Check: 10 bags = 500 kg of urea, supplying $500 \times 0.46 = 230$ kg of N, which slightly exceeds the 225 kg required — acceptable.

7.4 Method 4: Calculating Plant Density and Expected Yield

METHOD How to Use Spacing to Compute Plant Population and Yield

- Convert all measurements to the same unit (metres and hectares: 1 ha = 10 000 m²).
- Area occupied by one plant = row spacing × spacing within the row.

3. Plant population:

$$\text{Population} = \frac{\text{area of field}}{\text{area per plant}}$$

4. Expected yield = population × mean yield per plant (or per stand, if several seeds are planted per hole).

5. Discuss: closer spacing increases population but increases competition for light, water and nutrients; wider spacing wastes land. Correct spacing is a key **cultivation technique**.

WORKED EXAMPLE 11 Groundnut Field in the North Region

A farmer plants groundnuts at a spacing of 50 cm between rows and 25 cm within the row, on a field of 1.5 ha. Each stand produces on average 40 g of dry pods.

- Calculate the plant population of the field.
- Estimate the total expected harvest in kg and in 100 kg bags.
- State two advantages of correct spacing.

SOLUTION

a) Plant population.

Convert spacings to metres: 50 cm = 0.5 m; 25 cm = 0.25 m.

Area per plant:

$$0.5 \times 0.25 = 0.125 \text{ m}^2.$$

Area of field:

$$1.5 \text{ ha} = 1.5 \times 10\,000 = 15\,000 \text{ m}^2.$$

Population:

$$\frac{15\,000}{0.125} = 120\,000 \text{ plants.}$$

b) Expected harvest.

Total yield:

$$120\,000 \times 40 \text{ g} = 4\,800\,000 \text{ g} = 4800 \text{ kg.}$$

Number of 100 kg bags:

$$\frac{4800}{100} = 48 \text{ bags.}$$

c) Advantages of correct spacing.

- Each plant receives adequate light, water and mineral salts, reducing competition and giving uniform, high yields.
- It facilitates weeding, pest control and harvesting (passage between rows), and ensures good aeration, which limits fungal diseases.

7.5 Method 5: Choosing and Justifying Cultivation Techniques

METHOD How to Answer a Question on Crop Cultivation Techniques

For any named crop (food crop: maize, cassava, plantain; cash/non-food crop: cocoa, oil palm, cotton, rubber), structure your answer along the production cycle:

1. Choice of site and land preparation: suitable soil and climate; clearing, ploughing/tilling, ridging or mounding.
2. Planting material: improved/certified seeds, cuttings (cassava), suckers (plantain), budded/grafted seedlings (cocoa, rubber); treatment of seeds against pests.
3. Planting: correct season (onset of the rains), depth, spacing.
4. Maintenance (cultural practices): weeding, mulching, thinning, staking, pruning, earthing-up, irrigation/drainage, fertiliser/manure application, pest and disease control (integrated pest management).
5. Harvesting and post-harvest: at correct maturity; drying, storage in cribs/barns, processing.
6. For perennial cash crops: add the nursery stage, shading of young plants, and the immature period before production (e.g. cocoa 3–5 years, oil palm about 3–4 years, rubber about 6–7 years).

Reflex: always link each technique to its purpose (e.g. “mulching conserves soil moisture and suppresses weeds”).

WORKED EXAMPLE 12 Comparing Cassava and Cocoa Cultivation

- a) Describe the propagation and planting of cassava.
- b) Give three ways in which the cultivation of cocoa differs from that of cassava, explaining each difference.

SOLUTION

- a) Propagation and planting of cassava.

Cassava is propagated vegetatively by stem cuttings. Mature stems (from plants 8–12 months old) are cut into pieces about 20–25 cm long, each bearing several nodes. The land is cleared and tilled, often into ridges or mounds. Cuttings are planted at the onset of the rains, inserted vertically or slanting with about two-thirds of their length buried, at a spacing of about 1 m × 1 m (roughly 10 000 stands/ha). The field is weeded regularly (2–3 times) and the tubers are ready for harvest 9–12 months after planting.

- b) Differences with cocoa cultivation.

1. Mode of propagation: cocoa is grown from seeds first raised in a nursery (or from budded/grafted seedlings), whereas cassava is planted directly as stem cuttings. Reason: cocoa seeds lose viability quickly and young seedlings are delicate, so they need nursery protection before transplanting.
2. Life cycle and waiting period: cocoa is a perennial tree crop that starts bearing pods only after 3–5 years and produces for decades; cassava is an annual/semi-perennial food crop harvested after less than a year. Cocoa therefore requires long-term investment and permanent management of the plantation.
3. Shade requirement: young cocoa plants require shade (provided by plantain/banana or shade trees) because direct strong sunlight damages them; cassava is a sun-loving crop planted in open fields.

(Also acceptable: cocoa requires a humid equatorial climate with well-distributed rain-fall, while cassava tolerates poorer soils and drier conditions.)

7.6 Method 6: Formulating a Balanced Animal Ration (Pearson's Square)

METHOD How to Balance a Ration Using Pearson's Square

To mix two feedstuffs (an energy feed and a protein supplement) to obtain a desired protein percentage:

1. Write the target protein % in the centre of a square.
2. Write the protein % of the energy feed (top left) and of the protein supplement (bottom left).
3. Subtract diagonally (larger minus smaller) to obtain the parts of each feed on the right side.
4. Convert parts to percentages of the total, then to masses for the quantity of feed required.
5. Check: the protein contributed by each component must sum to the target.

Reflex: an animal ration must supply energy (carbohydrates/fats), protein, minerals, vitamins and water, adjusted to the animal's age, mass and production (growth, lactation, egg-laying). Pearson's square balances one nutrient (usually crude protein).

WORKED EXAMPLE 13 Ration for Broiler Chickens


A poultry farmer in Bamenda wants to prepare 100 kg of a broiler feed containing 20 %

crude protein (CP) by mixing maize (9 % CP) and soybean meal (45 % CP).

- Use Pearson's square to find the proportions of maize and soybean meal.
- Calculate the mass of each ingredient in the 100 kg mixture.
- Verify your answer.

SOLUTION

a) Pearson's square.



Total parts: $25 + 11 = 36$ parts.

Proportions:

$$\text{Maize} = \frac{25}{36} \times 100 = 69.4\% \quad \text{Soybean meal} = \frac{11}{36} \times 100 = 30.6\%$$

b) Masses for 100 kg of feed.

$$\text{Maize} = \frac{25}{36} \times 100 = 69.4 \text{ kg}; \quad \text{Soybean meal} = \frac{11}{36} \times 100 = 30.6 \text{ kg}.$$

c) Verification.

Protein supplied by maize: $69.4 \times 0.09 = 6.25 \text{ kg}$.

Protein supplied by soybean: $30.6 \times 0.45 = 13.77 \text{ kg}$.

Total protein: $6.25 + 13.77 = 20.02 \approx 20 \text{ kg per } 100 \text{ kg of feed}$, i.e. 20 % CP. The mixture is correct.

Practical note: in a real ration, a mineral–vitamin premix, salt and a source of calcium (e.g. oyster shell) would also be added, and the maize/soybean proportions slightly reduced to accommodate them.

7.7 Method 7: Evaluating Animal Production Performance

METHOD

How to Calculate Growth Rate and Feed Conversion Ratio

1. Average daily gain (ADG):

$$\text{ADG} = \frac{\text{final mass} - \text{initial mass}}{\text{number of days}}$$

2. Feed conversion ratio (FCR):

$$\text{FCR} = \frac{\text{total feed consumed (kg)}}{\text{total weight gain (kg)}}$$

A low FCR means efficient conversion (broilers typically around 1.6–2; pigs around 3; cattle considerably higher).

3. Cost of production per kg of gain = FCR × cost of 1 kg of feed.

4. Interpret and advise: compare with standard values; suggest improvements (balanced ration, disease control, improved breeds, good housing and hygiene).

Reflex: in production questions, always end with an interpretation — a number alone earns only part of the marks.

WORKED EXAMPLE 14 Performance of a Piggery

A farmer fattens a batch of pigs. At the start, the average mass per pig is 25 kg; after 90 days the average mass is 85 kg. Each pig consumed on average 180 kg of feed costing 250 FCFA per kg.

- Calculate the average daily gain (ADG).
- Calculate the feed conversion ratio (FCR) and comment.
- Calculate the feed cost per kg of weight gain.
- Suggest two ways of improving the FCR.

SOLUTION

a) Average daily gain.

$$\text{ADG} = \frac{85 - 25}{90} = \frac{60}{90} = 0.67 \text{ kg/day } (\approx 670 \text{ g/day}).$$

b) Feed conversion ratio.

$$\text{FCR} = \frac{\text{feed consumed}}{\text{weight gain}} = \frac{180}{60} = 3.0.$$

Comment: the pig eats 3 kg of feed to produce 1 kg of weight gain. This is a typical, acceptable value for pigs, though well-managed improved breeds on balanced rations can do slightly better (around 2.5–2.8).

c) Feed cost per kg of gain.

$$\text{Cost} = \text{FCR} \times \text{price per kg of feed} = 3.0 \times 250 = 750 \text{ FCFA per kg of weight gain.}$$

The farmer can compare this with the selling price of pork per kg live weight to judge profitability.

d) Improving the FCR.

- Provide a balanced ration with adequate protein, energy, minerals and vitamins (e.g. correctly formulated using Pearson's square), and clean water ad libitum.
- Improve health and husbandry: regular deworming and vaccination, clean dry housing, reduced stress; and/or use improved breeds with better genetic potential for growth.

7.8 Method 8: Interpreting an Animal Production System

METHOD How to Analyse an Animal Production System (Extensive vs Intensive)

1. Identify the system from the description: **extensive** (large area, animals graze freely, low inputs, low output per animal — e.g. traditional cattle herding on the Adamawa Plateau) or **intensive** (animals housed/confined, high inputs of feed, veterinary care and labour, high output per animal — e.g. modern poultry or piggery units near Yaoundé and Douala). **Semi-intensive** systems are intermediate.
2. List inputs and outputs: land, labour, capital, feed, veterinary care → meat, milk, eggs, hides, manure.
3. Compare in a table: stocking rate, productivity per animal, disease risk, cost, environmental impact.
4. Conclude with a justified judgement: which system suits which context (land availability, capital, market).

WORKED EXAMPLE 15 Choosing a Production System for Dairy Cattle

A group of farmers on the Bamenda Highlands wants to increase milk production. They currently keep local zebu cattle on open range (extensive system), obtaining about 2 litres of milk per cow per day. An adviser proposes an intensive zero-grazing unit with improved (exotic) dairy cows.

- a) Give three advantages the intensive system could bring.
- b) Give three constraints or risks the farmers must consider.
- c) Propose one intermediate (semi-intensive) measure.

SOLUTION

- a) Advantages of the intensive system.

- Higher productivity per animal: improved dairy breeds, well fed, can produce several times more milk per day than local zebus on range.
- Better health control and management: housed animals are easily inspected, treated, vaccinated and milked hygienically; breeding (e.g. artificial insemination) is controlled.
- Efficient land use: zero grazing produces more milk per hectare, which is important where land is limited, and manure is easily collected for crop fields or biogas.

b) Constraints and risks.

- High capital and running costs: construction of housing, purchase of improved cows, concentrates and veterinary products; a market failure or disease outbreak can cause heavy losses.
- Feed supply must be guaranteed: zero-grazed animals depend entirely on cut fodder and purchased feed; fodder must be grown or conserved (silage, hay) for the dry season.
- Exotic breeds are less hardy: they are generally more susceptible to heat, parasites (e.g. ticks) and diseases than local zebus, requiring skilled management and reliable veterinary services.

c) Semi-intensive measure.

The farmers could keep their cattle in fenced paddocks or night kraals, supplementing grazing with cultivated fodder (e.g. elephant grass, Calliandra) and mineral licks, while introducing improved bulls or artificial insemination to upgrade the herd gradually. This raises productivity at moderate cost and risk, allowing a progressive transition towards intensification.

8 EXERCISES

9 I CHECK MY KNOWLEDGE

EXERCISE 9 Multiple choice questions

For each question, choose the single correct answer.

1. The soil horizon richest in humus and most important for crop growth is:
 - (a) the A horizon
 - (b) the B horizon
 - (c) the C horizon
 - (d) the bedrock
2. The element whose deficiency causes interveinal chlorosis in older leaves and which is a component of chlorophyll is:
 - (a) nitrogen
 - (b) phosphorus
 - (c) magnesium
 - (d) potassium
3. In the Pearson square method, a ration of 16% crude protein is to be made from maize (9% CP) and groundnut cake (45% CP). The proportion of groundnut cake needed is approximately:
 - (a) 19%
 - (b) 36%
 - (c) 64%
 - (d) 81%
4. Newcastle disease in poultry is caused by:
 - (a) a bacterium
 - (b) a virus
 - (c) a fungus
 - (d) a protozoan

EXERCISE 10 True or false — justify

State whether each statement is true or false, and justify your answer in one or two sentences.

1. A clay soil is always more fertile than a sandy soil and therefore always better for farming.
2. Legumes such as groundnut and cowpea can grow well without nitrogen fertiliser because they fix atmospheric nitrogen through *Rhizobium* bacteria in their root nodules.
3. Ruminants such as cattle can digest cellulose because they themselves secrete the enzyme cellulase.
4. In extensive animal production, output per animal is generally lower than in intensive production.

EXERCISE 11 Definitions

Define each of the following terms precisely, as expected at the GCE Advanced Level:

1. soil profile;
2. humus;
3. macronutrient (in plant nutrition);
4. ration (in animal nutrition);
5. biosecurity (in animal production).

EXERCISE 12 Direct calculations

1. A soil sample contains 40 % sand, 40 % silt and 20 % clay. Calculate the sand-to-clay ratio and state, with a reason, whether this soil is likely to drain quickly or slowly.
2. A farmer applies urea (46% N) at a rate of 100 kg per hectare. Calculate the mass of nitrogen applied per hectare.
3. A broiler chick consumes 150 g of feed per day containing 21% crude protein. Calculate the mass of protein consumed per day.

10 I PRACTISE

EXERCISE 13 Using the texture triangle

A soil sample from a farm near Buea contains 70% sand, 20% silt and 10% clay.

1. Use the texture triangle to identify the textural class of this soil.
2. State three physical properties of this soil type (drainage, water retention, ease of tillage).
3. Suggest two crops suited to this soil, giving a reason for each choice.
4. Propose two measures the farmer could take to improve the fertility of this soil.

EXERCISE 14 Mineral elements in plants

Copy and complete the following table by giving, for each mineral element, one function in the plant and one symptom of its deficiency.

Element	One function	One deficiency symptom
Nitrogen (N)		
Phosphorus (P)		
Potassium (K)		
Magnesium (Mg)		
Iron (Fe)		

METHOD Balancing a ration using the Pearson square

The Pearson square is a simple tool to calculate the proportions of two feeds needed to obtain a mixture with a desired nutrient concentration.

1. Write the desired nutrient percentage in the centre of the square.
2. Write the nutrient percentages of the two available feeds at the top left and bottom left corners.
3. Subtract diagonally (larger minus smaller) to get the parts of each feed required. The parts are always positive.
4. The sum of the parts gives the total parts of the mixture.

5. Calculate the percentage of each feed: $(\text{parts of feed} / \text{total parts}) \times 100\%$.

EXERCISE 15 Balancing a ration with the Pearson square

A poultry farmer in Bafoussam wants to prepare a broiler starter ration containing 20% crude protein using maize (9% crude protein) and fish meal (60% crude protein).

1. Draw a Pearson square and calculate the proportions of maize and fish meal required.
2. Calculate the mass of each ingredient needed to prepare 100 kg of the ration.
3. Comment on the result: why is fish meal used only in a limited proportion, apart from the protein requirement? (Think of cost and palatability.)

EXERCISE 16 Cultivation of a food crop

A farmer in the West Region wishes to establish a one-hectare maize farm.

1. List, in chronological order, the main cultivation operations from land preparation to storage of the harvest.
2. For each operation, state its purpose.
3. Indicate two common pests or diseases of maize in Cameroon and one control measure for each.

11 I PREPARE FOR THE GCE

EXERCISE 17 Structured question: soils and fertility (20 marks)

The table below gives the results of the analysis of two soil samples, A and B, taken from two farms.

Sample	Sand (%)	Silt (%)	Clay (%)	Humus content
A	85	10	5	Low
B	20	25	55	Moderate

1. Identify the textural class of each soil. (4 marks)
2. Compare the drainage, aeration and water-holding capacity of the two soils. (6)

marks)

3. Explain why soil A warms up quickly in the dry season but loses nutrients easily by leaching. (4 marks)
4. For each soil, suggest one suitable crop and one method of improving its fertility, justifying your answers. (6 marks)

EXERCISE 18 Essay: cultivation of cocoa (20 marks)

Cocoa is one of the main export crops of Cameroon.

Describe the cultivation of cocoa from land preparation to the marketing of dried beans, indicating the importance of each operation. Your answer should include: choice of site and soil, land preparation, planting and shade management, maintenance operations (weeding, pruning, pest and disease control, fertiliser application), harvesting, fermentation, drying and marketing.

(Marks will be awarded for the correct sequence of operations, the technical accuracy of each step, and the stated importance of each operation.)

EXERCISE 19 Problem situation and essay: animal production (20 marks)

Part A. A poultry farmer near Yaoundé records high mortality among week-old chicks showing respiratory signs (gasping, nasal discharge) and twisted necks.

1. Identify the most likely disease and its causal agent. (2 marks)
2. Describe how the disease spreads within a flock. (3 marks)
3. Outline a vaccination and biosecurity programme that the farmer should adopt to protect the farm. (5 marks)

Part B. Discuss the advantages and disadvantages of intensive and extensive systems of animal production, with reference to cattle rearing in the Adamawa Region of Cameroon. Conclude by suggesting how the two systems could be improved or combined to increase meat and milk supply in Cameroon. (10 marks)

DID YOU KNOW? Cameroon context: Agriculture

Cameroon is often called "Africa in miniature" due to its diverse agro-ecological zones. The volcanic soils around Mount Cameroon and in the West Region (Bamenda Highlands) are

highly fertile and support intensive cultivation of crops like tea, coffee, and market gardening. The Adamawa plateau is the heart of extensive cattle ranching, while the Sudano-Saharan zone in the Far North relies on drought-tolerant crops like millet, sorghum, and cotton. Cocoa, coffee, cotton, and bananas are major export earners. The government's "Second Generation Agriculture" policy aims to modernise the sector through mechanisation, improved seeds, and value chain development.

WARNING Common mistakes in GCE Agriculture

- Confusing soil texture (mineral particle size) with soil structure (arrangement of particles into aggregates).
- Stating that legumes "do not need nitrogen" — they fix their own but still require nitrogen for growth; the fixation supplies it.
- Forgetting that ruminants rely on microbial cellulase, not their own enzyme.
- In Pearson square, subtracting the wrong way (always subtract diagonally: desired minus feed, taking absolute values).
- Listing cultivation operations out of chronological order (e.g., fertiliser before planting, or weeding after harvest).
- Vague disease control measures: "use chemicals" instead of naming specific fungicides/insecticides and application timing.

12 SOLUTIONS TO THE EXERCISES

SOLUTION Exercise 1 — Multiple choice questions

1. Answer: a) the A horizon.

The A horizon (topsoil) is the zone where humus accumulates and mixes with mineral particles. It is dark in colour, biologically very active (roots, earthworms, micro-organisms) and supplies most of the nutrients and water absorbed by crops. The B horizon is mainly a zone of accumulation of leached minerals, the C horizon is weathered parent rock, and the bedrock is unweathered rock.

2. Answer: c) magnesium.

Magnesium is the central atom of the chlorophyll molecule. Because Mg is mobile in the plant, it is withdrawn from older leaves to supply young ones, so deficiency appears first as **interveinal chlorosis** (yellowing between the veins, which remain green) on the older leaves. Nitrogen deficiency also causes yellowing of older leaves, but it is general, not interveinal, and N is not a component of the chlorophyll ring itself.

3. Answer: a) 19%.

Pearson square: desired CP = 16%; feeds 9% (maize) and 45% (groundnut cake).

$$\text{Parts of maize} = 45 - 16 = 29$$

$$\text{Parts of groundnut cake} = 16 - 9 = 7$$

$$\text{Total parts} = 29 + 7 = 36$$

$$\text{Groundnut cake} = \frac{7}{36} \times 100 \approx 19.4\% \approx 19\%$$

4. Answer: b) a virus.

Newcastle disease of poultry is caused by an avian paramyxovirus (a virus). This is why antibiotics are useless against it and control relies on vaccination and biosecurity.

SOLUTION Exercise 2 — True or false — justify

1. False. Clay soils retain water and nutrients well, but they are heavy, poorly drained, poorly aerated and difficult to till, and they become waterlogged in the rains and hard in the dry season. A loam (balanced mixture of sand, silt and clay) is generally the most fertile and easiest soil to farm, so a clay soil is not always better than a sandy one.
2. True. Legumes live in symbiosis with *Rhizobium* bacteria housed in root nodules; the bacteria reduce atmospheric nitrogen (N_2) to ammonium compounds that the plant

can use. The legume therefore obtains its nitrogen supply without mineral nitrogen fertiliser (though it still needs the other mineral elements, especially P and K).

3. False. Ruminants do not secrete cellulase themselves. Cellulose digestion is carried out by symbiotic micro-organisms (bacteria, protozoa, fungi) living in the rumen, which produce cellulase and ferment cellulose into volatile fatty acids that the animal absorbs.
4. True. In extensive production (e.g. free-range grazing on the Adamawa plateau) animals receive little supplementary feed and limited health care, so growth rate, milk yield and offtake per animal are lower than in intensive systems, where feed, housing and veterinary care are optimised. (However, output per hectare of labour/capital invested may be lower in extensive systems, which are cheaper to run.)

SOLUTION Exercise 3 — Definitions

1. Soil profile: a vertical section through the soil, from the surface down to the parent rock, showing the sequence of horizontal layers called **horizons** (O, A, B, C) which differ in colour, texture, structure and composition.
2. Humus: the dark, colloidal, partially decomposed organic matter of the soil, formed by the action of soil organisms on dead plant and animal remains; it improves soil structure, water retention and nutrient supply.
3. Macronutrient: a mineral element required by plants in relatively large quantities for normal growth, e.g. nitrogen, phosphorus, potassium, calcium, magnesium and sulphur (as opposed to micronutrients/trace elements such as Fe, Mn, Zn, Cu, B, Mo, needed in very small amounts).
4. Ration: the total quantity of feed given to an animal in a specified period (usually 24 hours), formulated to supply all the nutrients (energy, protein, minerals, vitamins, water) needed for maintenance and production (growth, milk, eggs).
5. Biosecurity: the set of preventive measures (quarantine of new animals, disinfection, control of visitors and vehicles, vaccination, hygiene of housing and equipment, safe disposal of carcasses) designed to prevent the introduction and spread of disease-causing agents on a farm.

SOLUTION Exercise 4 — Direct calculations

1. Sand-to-clay ratio:

$$\frac{\text{sand}}{\text{clay}} = \frac{40}{20} = 2 \quad (\text{i.e. } 2 : 1).$$

With 40% sand and only 20% clay, the soil is a loam tending to a sandy loam. It will drain moderately to fairly quickly: the high sand fraction creates large pores allowing water to percolate, while the 20% clay still retains some water and nutrients.

2. Mass of nitrogen applied:

$$m(\text{N}) = 100 \text{ kg} \times \frac{46}{100} = 46 \text{ kg of N per hectare.}$$

3. Mass of protein consumed per day:

$$m(\text{protein}) = 150 \text{ g} \times \frac{21}{100} = 31.5 \text{ g of crude protein per day.}$$

SOLUTION Exercise 5 — Using the texture triangle

- On the texture triangle, the point corresponding to 70% sand, 20% silt and 10% clay falls in the class sandy loam (close to the boundary with loamy sand). The soil is therefore a sandy loam.
- Physical properties of a sandy loam:
 - Drainage: good to rapid — the large pores between sand particles let excess water percolate freely, so waterlogging is rare.
 - Water retention: low to moderate — little water is held against gravity, so the soil dries out quickly and crops may suffer in dry spells.
 - Ease of tillage: very easy — the soil is light, crumbly and workable even with simple hand tools or animal traction; it also warms up quickly.
- Suitable crops (any two):
 - Groundnut: needs a light, well-drained soil so that the pegs can penetrate and the pods develop without rotting.
 - Cassava: tolerates low-fertility, well-drained sandy soils and gives reasonable tuber yields where heavier crops would fail. (Accept also: sweet potato, water-melon, pineapple, early maize with fertiliser.)
- Measures to improve fertility (any two):
 - Add organic matter (farmyard manure, compost, mulch, green manure) to increase water and nutrient retention and feed soil organisms.
 - Apply mineral fertilisers (NPK) in split doses to compensate for leaching losses.

- Practise crop rotation with a legume (cowpea, groundnut) to enrich the soil in nitrogen.


SOLUTION Exercise 6 — Mineral elements in plants

Element	One function	One deficiency symptom
Nitrogen (N)	Component of proteins, nucleic acids and chlorophyll; promotes vegetative (leafy) growth	General yellowing (chlorosis) of older leaves; stunted growth
Phosphorus (P)	Component of ATP, nucleic acids and membranes; promotes root development, flowering and seed formation	Poor root growth; dark green or purplish coloration of older leaves; delayed maturity
Potassium (K)	Activates enzymes; regulates stomatal opening and water balance; improves disease resistance and fruit quality	Yellowing and scorching (necrosis) of the margins of older leaves; weak stems
Magnesium (Mg)	Central atom of the chlorophyll molecule; enzyme activator	Interveneal chlorosis of older leaves (veins stay green)
Iron (Fe)	Needed for chlorophyll synthesis (component of electron carriers such as cytochromes)	Interveneal chlorosis of young leaves (Fe is immobile in the plant)

Examiner's note: the key distinction is the position of the symptoms: deficiencies of mobile elements (N, P, K, Mg) appear first on old leaves, whereas deficiencies of immobile elements (Fe, Ca) appear first on young leaves.

SOLUTION Exercise 7 — Balancing a ration with the Pearson square

1. Pearson square (desired CP = 20%; maize = 9%; fish meal = 60%):



Subtractions are done diagonally, taking positive differences:

$$\begin{aligned}\text{Parts of maize} &= 60 - 20 = 40 \\ \text{Parts of fish meal} &= 20 - 9 = 11 \\ \text{Total parts} &= 40 + 11 = 51 \\ \text{Maize} &= \frac{40}{51} \times 100 \approx 78.4\% \\ \text{Fish meal} &= \frac{11}{51} \times 100 \approx 21.6\%\end{aligned}$$

2. For 100 kg of ration:

$$\begin{aligned}m(\text{maize}) &= \frac{40}{51} \times 100 \approx 78.4 \text{ kg} \\ m(\text{fish meal}) &= \frac{11}{51} \times 100 \approx 21.6 \text{ kg}\end{aligned}$$

Check: protein supplied = $78.4 \times 0.09 + 21.6 \times 0.60 \approx 7.06 + 12.96 = 20.0$ kg, i.e. 20% of 100 kg. ✓

3. Fish meal is used in a limited proportion because:

- it is expensive — increasing its share would raise the cost of the ration and reduce profit;
- in excess it can give an undesirable fishy taste to the meat and reduce palatability of the feed;
- it also supplies minerals (Ca, P), so beyond the requirement it would unbalance the mineral content of the ration.

SOLUTION Exercise 8 — Cultivation of a food crop (maize)

1. and 2. Operations in chronological order, with their purposes:

Operation	Purpose
1. Land clearing	Remove bushes and previous crop residues to expose the soil
2. Ploughing /tilling	Loosen the soil, bury

2. Two common pests/diseases of maize in Cameroon (any two):

- Fall armyworm / stem borer (pests): control by early planting, field hygiene and application of a recommended insecticide into the funnel.
- Maize streak virus (disease): control by using resistant/tolerant varieties and controlling the leafhopper vector.
- Storage weevils: control by proper drying and use of a grain protectant or airtight storage.

SOLUTION Exercise 9 — Structured question: soils and fertility (20 marks)

Indicative mark scheme and model answers.

1. Textural classes (4 marks — 2 per soil). Using the texture triangle:

- Soil A (85% sand, 10% silt, 5% clay): sand (sandy soil). (2)
- Soil B (20% sand, 25% silt, 55% clay): clay (clay soil). (2)

2. Comparison (6 marks — 1 per valid comparative point).

- Drainage: A drains rapidly (large pores between sand grains); B drains slowly and is easily waterlogged.
- Aeration: A is well aerated; B is poorly aerated, especially when wet.
- Water-holding capacity: A holds little water and dries quickly; B holds a large amount of water (small pores retain it by capillarity).

3. Explanation (4 marks).

- A warms quickly because it holds little water: little heat is wasted evaporating water, so solar energy rapidly raises the temperature of the dry sand particles. (2)
- It loses nutrients by leaching because its large pores let rainwater percolate rapidly, and with very little clay and low humus there are few negatively charged colloids to retain nutrient cations (K^+ , Ca^{2+} , Mg^{2+} , NH_4^+), which are therefore washed down beyond the root zone. (2)

4. Crop and improvement for each soil (6 marks — 1.5 per item, justification required).

- Soil A: suitable crop: groundnut, cassava or watermelon, which tolerate light, well-drained, low-fertility soils (groundnut pods also need loose soil to develop). Improvement: regular addition of organic matter (compost/manure) to raise water and nutrient retention, plus split applications of NPK fertiliser to limit leaching losses.
- Soil B: suitable crop: swamp rice (paddy), which tolerates — indeed requires — waterlogged conditions, or yam/maize on ridges. Improvement: drainage or

ridging to improve aeration, addition of organic matter and liming if acid, to improve structure and workability.

Examiner expectations: correct use of the texture triangle; comparative (not just descriptive) statements in (b); a causal link between low water content / low colloid content and the two properties in (c); justified choices in (d) — a crop or method without justification earns only half the marks.

SOLUTION Exercise 10 — Essay: cultivation of cocoa (20 marks)

Indicative mark scheme: sequence and coverage 8 marks; technical accuracy 8 marks; stated importance of each operation 4 marks.

Model plan and content.

1. Choice of site and soil: cocoa requires a hot, humid climate (about 1500–2000 mm of well-distributed rainfall) and a deep, well-drained, humus-rich soil (loamy or volcanic), slightly acidic. Importance: waterlogging causes root rot and black pod, while shallow soils dry out and stress the trees.
2. Land preparation: clear the bush but retain or plant some tall trees for shade; line and peg at about 3 × 3 m. Importance: shade protects young plants from sun scorch and reduces evaporation; correct spacing ensures good canopy development and easy maintenance.
3. Planting and shade management: plant seedlings (from nursery) or seeds at stake at the start of the rains, with temporary shade (plantain, banana, Gliricidia); reduce shade progressively as the canopy closes. Importance: good establishment determines the future yield of the plantation.
4. Maintenance:
 - Weeding (2–3 times per year): removes competition for water and nutrients.
 - Pruning: removes dead, diseased and chupon (vertical) shoots to shape the tree, aerate the canopy and ease harvesting.
 - Pest and disease control: capsids (mirids) controlled with recommended insecticides; black pod disease (*Phytophthora*) controlled by regular harvesting of ripe pods, removal of diseased pods, pruning for aeration and copper-based fungicides.
 - Fertiliser application: NPK fertiliser maintains yield on old or exhausted soils.
5. Harvesting: ripe pods (yellow/orange) are cut with a sharp knife or hooked pole every 2–3 weeks, without damaging the cushion. Importance: regular, careful harvesting limits black pod losses and protects future flowers on the same cushion.
6. Breaking and fermentation: pods are broken and the beans (with mucilage) heaped in

boxes or banana-leaf piles for about 5–6 days, with turning. Importance: fermentation kills the embryo, removes the mucilage and develops the chocolate flavour and brown colour — it determines the commercial quality (grade).

7. Drying: beans are dried in the sun on trays/mats (or in dryers) to about 7–8% moisture, with regular turning. Importance: proper drying prevents mould and off-flavours during storage and transport.
8. Marketing: dried beans are sorted, bagged and sold to cooperatives or licensed buying agents, then exported (e.g. through the port of Douala). Importance: grading and proper storage preserve quality and secure the best price.

Examiner expectations: operations in the correct chronological order; at least one precise technical detail per step (spacing, fermentation duration, moisture target, named pest/disease); the importance of each operation explicitly stated.

SOLUTION Exercise 11 — Problem situation and essay: animal production (20 marks)

Part A (10 marks).


1. Most likely disease (2 marks): Newcastle disease, caused by an avian paramyxovirus (a virus). The combination of respiratory signs (gasping, nasal discharge), nervous signs (twisted neck / torticollis) and high mortality in young chicks is characteristic. (2)
2. Spread within the flock (3 marks — any three points):
 - by direct contact between sick and healthy birds;
 - through contaminated air/droplets (respiratory secretions);
 - through contaminated feed, water, litter and equipment (faeces contain the virus);
 - by people (attendants, visitors), vehicles and introduction of infected birds from other farms or live-bird markets.
3. Vaccination and biosecurity programme (5 marks):
 - Vaccination: vaccinate all chicks against Newcastle disease (e.g. first dose in the first days of life by eye drop or drinking water, with boosters according to the veterinary schedule); vaccinate against other local diseases (Gumboro, fowl pox) as advised; keep vaccines cold (cold chain) and follow the calendar strictly.
 - Biosecurity: quarantine new birds before introduction; restrict and disinfect visitors, vehicles and equipment (footbaths at the entrance); provide clean water and feed; clean and disinfect the house between batches (all-in/all-out system); isolate sick birds and burn/bury dead birds; avoid mixing ages and species.

Part B (10 marks). Indicative content:

- Extensive system (e.g. Fulani cattle herds grazing the Adamawa plateau): advantages — low capital and feed costs (natural pasture), hardy local breeds (e.g. Gudali) adapted to the environment, large areas usable, animals convert grass inedible to humans into meat and milk. Disadvantages — low productivity per animal (slow growth, low milk yield, late maturity), losses to diseases (trypanosomiasis, tick-borne diseases) and parasites, overgrazing and soil degradation, seasonal weight loss in the dry season, farmer-grazer conflicts, difficulty of veterinary follow-up.
- Intensive system (zero-grazing, feedlots, dairy units): advantages — high output per animal (improved breeds, balanced rations), better disease control and record keeping, production all year round, efficient use of small land areas. Disadvantages — high investment and running costs (feed, housing, veterinary care), need for skilled labour, rapid spread of disease in dense housing if biosecurity fails, dependence on purchased inputs.
- Improvement / combination (conclusion): adopt semi-intensive systems: keep herds on improved or sown pastures with supplementary feeding (crop residues, agro-industrial by-products, mineral licks) especially in the dry season; provide vaccination and dipping against ticks; improve breeds by crossing local hardy cattle with higher-yielding breeds; develop water points and fodder banks on the Adamawa; organise ranching with paddocks to prevent overgrazing; strengthen veterinary services and farmer training. This combines the low cost of extensive grazing with the productivity gains of intensive management, increasing meat and milk supply sustainably.

Examiner expectations: Part A — correct identification of the disease and agent, concrete modes of transmission, and a programme mentioning both a vaccination schedule and named biosecurity measures. Part B — balanced discussion (at least two advantages and two disadvantages per system), explicit reference to the Adamawa context, and a reasoned, realistic conclusion.

13 SUMMARY SHEET



Mind map of Chapter BIO08 – Agriculture: five interconnected branches from soil to animal production.

FICHE BILAN L'essentiel à connaître pour l'examen

1. Key definitions

- **Soil:** the upper layer of the Earth's crust, formed by weathering of parent rock plus decomposition of organic matter, which supports plant growth.
- **Soil profile:** vertical section showing horizons O (organic), A (topsoil), B (subsoil), C (weathered parent material), R (bedrock).
- **Soil texture:** relative proportions of sand, silt and clay particles; determines

drainage, aeration and water retention.

- **Loam**: balanced mixture of sand, silt and clay with humus – the most fertile soil for most crops.
- **Soil fertility**: capacity of a soil to supply water, air, anchorage and mineral nutrients for good crop growth.
- **Macronutrients**: elements needed in large amounts (N, P, K, Ca, Mg, S); **micronutrients** (trace elements) needed in tiny amounts (Fe, Mn, B, Zn, Cu, Mo, Cl).
- **Fertiliser**: material (organic – manure, compost; or inorganic – NPK, urea) added to soil to restore nutrients.
- **Food crop**: grown mainly for human consumption (maize, cassava, plantain, yam); **cash (non-food) crop**: grown for sale/industry (cocoa, coffee, cotton, oil palm).
- **Balanced ration**: feed supplying all an animal's nutrients in correct proportions for maintenance and production.
- **Extensive system**: animals range freely with low inputs; **intensive system**: animals confined, high inputs, high yield per head.

2. Facts, comparisons and “laws” to know

- Soil formation = **weathering** (physical: temperature, rain, wind; chemical: hydrolysis, oxidation; biological: roots, microbes, earthworms) + humus accumulation.
- Texture comparison: sandy soils drain fast, warm quickly, poor nutrient retention; clay soils retain water and nutrients but are heavy, poorly aerated, waterlog easily; loam is intermediate and best.
- Soil pH: most crops prefer 6.0–7.0; acidity corrected by **liming** (adding CaCO_3).
- Roles of key elements: N – leaf/vegetative growth (deficiency: yellowing, chlorosis); P – roots, energy transfer ATP, flowering (deficiency: purpling, poor roots); K – enzyme activation, water regulation, disease resistance (deficiency: scorched leaf edges); Mg – chlorophyll component (deficiency: interveinal chlorosis); Fe – chlorophyll synthesis (deficiency: chlorosis of young leaves).
- Legumes (groundnut, cowpea, soya) fix atmospheric nitrogen via Rhizobium in root nodules – basis of crop rotation.
- Fertility maintenance: crop rotation, fallowing, mulching, manuring, mixed cropping, contour ploughing against erosion.
- Cultivation steps: land clearing → ploughing/tilling → harrowing → ridging → planting (correct spacing/depth) → weeding, fertilising, pest control → harvesting → storage.
- Animal feed classes: carbohydrates (energy), proteins (growth, repair, milk/egg production), lipids (concentrated energy), vitamins and minerals (health, bones: Ca and P), water (essential, often forgotten).

- Ruminants (cattle, goats, sheep) digest cellulose via rumen microbes; non-ruminants (pigs, poultry) need higher-quality protein and cannot use roughage efficiently.
- Animal production management: selection and breeding, housing/hygiene, vaccination and deworming, balanced feeding, record keeping.

3. Methods to master

- Soil texture test (feel method): moisten a sample, rub between fingers – gritty = sandy; smooth and sticky, moulds into a ball = clay; slightly gritty, holds shape loosely = loam.
- Sedimentation test: shake soil with water in a jar, let settle – sand sinks first (bottom layer), then silt, clay stays suspended longest; humus floats.
- Water retention/drainage experiment: equal volumes of sand, clay, loam in funnels with filter paper; pour equal water; compare volume drained and time.
- Planning a crop enterprise: choose site and crop suited to soil/climate → prepare land → select viable seed → plant at correct season → manage → harvest and store.

4. Common mistakes to avoid

- Confusing **soil texture** (particle sizes) with **soil structure** (arrangement into crumbs).
- Saying fertilisers “feed” plants – plants make their own food by photosynthesis; fertilisers supply mineral elements.
- Believing more fertiliser always means more yield – excess causes toxicity, leaching and water pollution (eutrophication).
- Confusing macronutrients with “most important” – trace elements are equally essential, only needed in smaller quantities.
- Mixing up extensive and intensive systems, or assuming intensive is always better (cost, disease risk, waste disposal).
- Forgetting water as a nutrient for animals, or ignoring the special digestion of ruminants.

5. GCE tips

- Draw and label the **soil profile** (O, A, B, C, R) neatly – a frequent diagram question.
- Learn deficiency symptoms of N, P, K, Mg, Fe in a table – examiners love “state and explain” questions.
- Use Cameroonian examples: cocoa and coffee in the South-West and Centre, cotton in the North, maize in the West highlands, cattle on the Adamawa plateau.
- In essays on soil fertility, always mention both organic methods (manure, rotation, fallow) and inorganic methods (NPK, liming).

- For animal production questions, structure answers under: feeding, housing, health, breeding, records.
- Define every technical term you use (e.g. ration, weeding, mulching) – marks are awarded for precise vocabulary.

**Signed • OnBuch+**ONBUCH+ TEACHING TEAM
FOLLOWS THE OFFICIAL MINESEC SYLLABUS**OnBuch+**