

PURE MATHEMATICS WITH STATISTICS • UPPER SIXTH

UPPER SIXTH — PURE MATHEMATICS WITH STATISTICS

CHAPTER 1

CIRCLE GEOMETRY

SUGGESTED TIME: 22 PERIODS

IN THIS CHAPTER

This chapter develops the complete analytic study of the circle in the Cartesian plane: its equations in all standard and general forms, tangents, intersections with lines and other circles, parametric representation, loci and the radical axis. It provides the essential toolkit for solving GCE Advanced Level problems on circle geometry.

- By the end of this chapter I can write the equation of a circle given its centre and radius, a diameter, or three points on it.
- By the end of this chapter I can convert the general equation $x^2 + y^2 + 2gx + 2fy + c = 0$ into centre-radius form and identify the centre and radius.
- By the end of this chapter I can find the equation of a tangent to a circle at a given point or from an external point, and state the condition for a line to touch a circle.
- By the end of this chapter I can use parametric equations of a circle and convert between parametric and Cartesian forms.
- By the end of this chapter I can find the points of intersection of a line and a circle, and of two circles.
- By the end of this chapter I can determine the equation of a circle through the intersection of two circles or of a line and a circle using families of circles.



DOWNLOAD ON- BUCH

Free • Google Play
AI tutor, past papers,
results



GO ONBUCH+

All signed courses, unlimited AI
tutor, PDF sheets — OnBuch+
tab in the app

THIS SIGNED COURSE CONTAINS

COURSE

complete, illustrated,
step by step

METHODS

and worked examples

EXERCISES

exam style + solutions

SUMMARY SHEET

the essentials to
remember



Signed • OnBuch+

ONBUCH+ TEACHING TEAM
FOLLOWS THE OFFICIAL MINESEC SYLLABUS

OnBuch+

CONTENTS

1. Section 1: Equations of a Circle — Centre at the Origin and General Centre
2. Section 2: The General Equation and Circles on a Given Diameter
3. Section 3: Circle Through Three Points and Circles Under Geometric Conditions
4. Section 4: Tangents to a Circle
5. Section 5: Parametric Equations of a Circle
6. Section 6: Intersection of a Line and a Circle
7. Section 7: Intersection of Two Circles, Families of Circles and the Radical Axis
8. Section 8: Conditions for Two Circles to Touch
9. Section 9: Locus Problems and Synthesis for the GCE Exam
10. Integration activity
11. Methods and worked examples
12. Exercises
13. Solutions to the exercises
14. Summary sheet

CHAPTER OBJECTIVES

- By the end of this chapter I can write the equation of a circle given its centre and radius, a diameter, or three points on it.
- By the end of this chapter I can convert the general equation $x^2 + y^2 + 2gx + 2fy + c = 0$ into centre–radius form and identify the centre and radius.
- By the end of this chapter I can find the equation of a tangent to a circle at a given point or from an external point, and state the condition for a line to touch a circle.
- By the end of this chapter I can use parametric equations of a circle and convert between parametric and Cartesian forms.
- By the end of this chapter I can find the points of intersection of a line and a circle, and of two circles.
- By the end of this chapter I can determine the equation of a circle through the intersection of two circles or of a line and a circle using families of circles.
- By the end of this chapter I can state and apply the conditions for two circles to touch internally or externally.
- By the end of this chapter I can find the radical axis of two circles and the length of a tangent from an external point.

- By the end of this chapter I can solve locus problems and find circles satisfying mixed conditions (touching axes, touching a line, centre on a line).

PREREQUISITES

- Coordinate geometry of the straight line: distance between two points, midpoint, gradient, equations of lines, perpendicular gradients (Lower Sixth).
- Pythagoras' theorem and basic circle theorems from O-Level geometry.
- Solving simultaneous equations (linear and quadratic) and the discriminant of a quadratic.
- Completing the square for a quadratic expression.
- Basic trigonometry: sine and cosine of an angle (useful for parametric equations).

1 SECTION 1: EQUATIONS OF A CIRCLE — CENTRE AT THE ORIGIN AND GENERAL CENTRE

In Bamenda, a telecommunications engineer must position a relay antenna so that it is exactly the same distance from three villages; on a map, the villages are points and the antenna sits at the centre of a circle through them. In Douala, a roundabout is designed so that a straight avenue just touches its circular kerb — a tangent problem. When a potter in Bafoussam spins clay, every point of the rim traces a circle whose equation we can write down. This chapter gives you the algebraic tools to describe, construct and analyse all these situations: the circle, the simplest curve after the straight line, and a favourite of the GCE Advanced Level examiners.

The central problem of this first section is the following: how can we translate the geometric idea of a circle into an algebraic equation that we can manipulate? Once we can do that, questions about centres, radii, and positions of points become routine algebra.

1.1 The Circle as a Locus

Think of the potter's wheel in Bafoussam. Fix your attention on one grain of clay at the rim. As the wheel spins, that grain moves, but its distance from the centre of the wheel never changes. The rim is exactly the collection of all positions whose distance from the centre equals the radius of the wheel. This everyday observation is the whole secret of the circle.

DEFINITION Circle as a locus

A **circle** is the **locus** of a point which moves in a plane so that its distance from a fixed point, called the **centre**, is constant. This constant distance is called the **radius** of the circle.

The word locus (plural loci) simply means “the set of all points satisfying a given condition”. So a circle is not a drawn object first — it is a condition: “be at distance r from C ”. Our task is to rewrite this condition in coordinates. We shall return to locus problems in full generality in Section 9; here the locus condition is the simplest possible one.

1.2 Circle Centred at the Origin

Place the centre at the origin $O(0, 0)$ of a Cartesian plane, and let $P(x, y)$ be any point on the circle of radius r . By the distance formula (itself a direct consequence of Pythagoras' theorem),

$$OP = \sqrt{(x - 0)^2 + (y - 0)^2} = \sqrt{x^2 + y^2}.$$


The locus condition $OP = r$ gives $\sqrt{x^2 + y^2} = r$, and squaring both sides (both sides are non-negative, so this is safe) yields the equation of the circle.

PROPERTY Circle centred at the origin

The circle with centre $O(0, 0)$ and radius r has equation

$$x^2 + y^2 = r^2.$$

This derivation is examinable: you may be asked to derive it from the distance formula.



The right triangle inside the circle: $x^2 + y^2 = r^2$ is Pythagoras' theorem.

EXAMPLE Reading and writing equations centred at the origin

1. The circle $x^2 + y^2 = 25$ has centre $(0, 0)$ and radius $r = \sqrt{25} = 5$.
2. The circle with centre $(0, 0)$ and radius 3 has equation $x^2 + y^2 = 9$.
3. The circle $x^2 + y^2 = 7$ has radius $\sqrt{7}$: the radius need not be a whole number.

WARNING The right-hand side is r^2

In $x^2 + y^2 = r^2$, the number on the right is the square of the radius. For $x^2 + y^2 = 25$ the radius is 5, not 25. This is one of the most frequent errors in GCE scripts — always take the square root.

1.3 Circle with Centre $C(a, b)$: the Standard Form

Now suppose the centre is no longer at the origin but at a general point $C(a, b)$. Geometrically, nothing has changed: the circle is just shifted a units horizontally and b units vertically. Let $P(x, y)$ be any point on the circle. The locus condition is still $CP = r$, and the distance formula gives

$$\sqrt{(x - a)^2 + (y - b)^2} = r.$$


Squaring both sides produces the standard form.

PROPERTY Standard equation of a circle

The circle with centre $C(a, b)$ and radius r has equation

$$(x - a)^2 + (y - b)^2 = r^2.$$

Conversely, any equation of this form represents a circle with centre (a, b) and radius $\sqrt{r^2}$. The derivation from the distance formula is examinable.



Shifting the centre to $C(a, b)$: the horizontal leg has length $x - a$, the vertical leg $y - b$.

WARNING Watch the signs of the centre

In $(x - a)^2 + (y - b)^2 = r^2$ the centre is (a, b) , not $(-a, -b)$. The formula contains minus signs, so:

- $(x - 3)^2 + (y - 2)^2 = 16$ has centre $(3, 2)$;
- $(x + 3)^2 + (y - 1)^2 = 16$, i.e. $(x - (-3))^2 + (y - 1)^2 = 16$, has centre $(-3, 1)$.

Rewrite any “+” as “ $-(-)$ ” before reading off the coordinates.

EXAMPLE Reading centre and radius from the standard form

1. $(x - 2)^2 + (y + 5)^2 = 9$: centre $(2, -5)$, radius 3.
2. $(x + 1)^2 + (y + 4)^2 = 20$: centre $(-1, -4)$, radius $\sqrt{20} = 2\sqrt{5}$.
3. $x^2 + (y - 3)^2 = 4$: this is $(x - 0)^2 + (y - 3)^2 = 4$, centre $(0, 3)$ on the y -axis, radius 2.

METHOD Sketching a circle from its equation

1. Read the centre $C(a, b)$ and compute $r = \sqrt{r^2}$.
2. Plot C in the plane.
3. Mark the four points $(a \pm r, b)$ and $(a, b \pm r)$ — the extreme points of the circle.
4. Draw a smooth circle through these four points and label the centre and radius.

1.4 Circle Given its Centre and a Point on it

Suppose we know the centre C and one point P lying on the circle. Then the radius is simply the distance CP : the point on the circle “tells us” how big the circle is. Since the equation only needs r^2 , we compute $r^2 = CP^2$ directly — no square root is ever needed.

METHOD Equation given the centre and a point on the circle

1. Compute $r^2 = CP^2 = (x_p - a)^2 + (y_p - b)^2$.
2. Substitute into the standard form $(x - a)^2 + (y - b)^2 = r^2$.
3. Expand only if the question asks for the general form.

WORKED EXAMPLE 1 GCE-style: centre and a point on the circle

Find the equation of the circle with centre $C(2, -1)$ which passes through the point $P(5, 3)$.

SOLUTION

Solution. The radius squared is the squared distance CP :

$$r^2 = (5 - 2)^2 + (3 - (-1))^2 = 3^2 + 4^2 = 9 + 16 = 25.$$

Substituting into the standard form with centre $(2, -1)$:

$$(x - 2)^2 + (y + 1)^2 = 25.$$

Check: does $P(5, 3)$ satisfy it? $(5 - 2)^2 + (3 + 1)^2 = 9 + 16 = 25$. ✓ The equation is correct.

WORKED EXAMPLE 2 Circle centred on an axis, passing through the origin

A circle has its centre on the positive x -axis and passes through the points $O(0, 0)$ and $A(6, 0)$. Find its equation.

SOLUTION

Solution. Since the centre lies on the x -axis, write $C(a, 0)$. The centre is equidistant from O and A :

$$CO^2 = a^2, \quad CA^2 = (6 - a)^2.$$

Setting $CO^2 = CA^2$: $a^2 = (6 - a)^2 = 36 - 12a + a^2$, so $12a = 36$ and $a = 3$. Then $r^2 = CO^2 = 9$, and the circle is

$$(x - 3)^2 + y^2 = 9.$$

Note that the origin lies on the circle: $(0 - 3)^2 + 0^2 = 9$. ✓

WARNING Circles through the origin


A circle passes through the origin if and only if $r^2 = a^2 + b^2$ (substitute $(0, 0)$ into the standard form). For example $(x - 3)^2 + (y - 4)^2 = 25$ passes through the origin since $9 + 16 = 25$. Recognising this saves time in the exam.

1.5 Position of a Point Relative to a Circle

The locus condition $CP = r$ splits the plane into three natural regions. For a point P and a circle of centre C and radius r :

- $CP < r$: P lies **inside** the circle;
- $CP = r$: P lies **on** the circle;
- $CP > r$: P lies **outside** the circle.

In practice we compare CP^2 with r^2 to avoid square roots: substitute the coordinates of P into the left-hand side $(x - a)^2 + (y - b)^2$ and compare the result with the right-hand side.



The distance test: $CA^2 = 2 < 9$, $CB^2 = 9$, $CD^2 = 32 > 9$.

WORKED EXAMPLE 3 Inside, on, or outside?

Determine the position of each point relative to the circle $(x-2)^2 + (y-2)^2 = 9$: $A(3, 3)$, $B(5, 2)$, $D(6, 6)$.

SOLUTION

Solution. Here $C(2, 2)$ and $r^2 = 9$. We compute the squared distance from C for each point.

$$A(3, 3) : (3-2)^2 + (3-2)^2 = 1+1 = 2 < 9 \Rightarrow A \text{ is inside.}$$

$$B(5, 2) : (5-2)^2 + (2-2)^2 = 9+0 = 9 = r^2 \Rightarrow B \text{ is on the circle.}$$

$$D(6, 6) : (6-2)^2 + (6-2)^2 = 16+16 = 32 > 9 \Rightarrow D \text{ is outside.}$$

EXAMPLE A Cameroonian application

A transmitter at the point $T(1, 2)$ (coordinates in kilometres on a map) covers every receiver within 5 km. Is a receiver at $R(5, 6)$ covered? The coverage zone is the circle $(x-1)^2 + (y-2)^2 = 25$. Compute

$$TR^2 = (5-1)^2 + (6-2)^2 = 16+16 = 32 > 25.$$

Since $TR > 5$, the receiver lies outside the coverage zone: it is not covered.

1.6 Consolidation

KEY POINTS Key points of Section 1

- A circle is the locus of points at a fixed distance r (radius) from a fixed point C (centre).
- Centre at the origin: $x^2 + y^2 = r^2$.
- Centre $C(a, b)$: $(x - a)^2 + (y - b)^2 = r^2$ — the standard form.
- Given the centre and a point P on the circle: $r^2 = CP^2$.
- Position of a point P : compare CP^2 with r^2 (inside, on, outside).
- The circle passes through the origin exactly when $a^2 + b^2 = r^2$.

DID YOU KNOW? Why the GCE loves the circle

The equation $(x - a)^2 + (y - b)^2 = r^2$ is nothing more than Pythagoras' theorem wearing co-ordinate clothing. Nearly every question in this chapter — tangents, intersections, loci — ultimately reduces to a right-angled triangle and the distance formula. Master this section and the rest of the chapter becomes a sequence of variations on one idea.

EXERCISE 1 Practice

- Write down the centre and radius of each circle:
 - $x^2 + y^2 = 49$;
 - $(x - 4)^2 + (y + 1)^2 = 36$;
 - $(x + 2)^2 + (y + 7)^2 = 12$.
- Find the equation of the circle with centre $(-3, 2)$ passing through $(1, 5)$.
- Find the equation of the circle with centre on the y -axis, passing through $(0, 0)$ and $(0, 8)$.
- Determine whether $(4, -2)$ lies inside, on or outside the circle $(x - 1)^2 + (y + 2)^2 = 10$.

SOLUTION Exercise 1

- (a) Centre $(0, 0)$, $r = 7$. (b) Centre $(4, -1)$, $r = 6$. (c) Centre $(-2, -7)$, $r = \sqrt{12} = 2\sqrt{3}$.

2. $r^2 = (1 + 3)^2 + (5 - 2)^2 = 16 + 9 = 25$, so $(x + 3)^2 + (y - 2)^2 = 25$.
3. Centre $(0, b)$ with $b^2 = (8 - b)^2$, giving $b = 4$ and $r^2 = 16$: $x^2 + (y - 4)^2 = 16$.
4. $(4 - 1)^2 + (-2 + 2)^2 = 9 < 10$: the point lies inside the circle.

2 SECTION 2: THE GENERAL EQUATION AND CIRCLES ON A GIVEN DIAMETER

In Section 1 we established the **centre–radius form** of a circle: a circle with centre $C(a, b)$ and radius r has equation

$$(x - a)^2 + (y - b)^2 = r^2.$$

This form is perfect when the centre and radius are known. But in many GCE questions the equation of a circle is handed to you already expanded, with all brackets removed, and you must work backwards to recover the centre and radius. In other problems you are told that a segment AB is a diameter of the circle, and you must build the equation from the two endpoints. This section develops both techniques: the **general equation** of a circle and the **diameter form**.

2.1 From Centre–Radius Form to the General Equation

An observation to start with.

Take the circle with centre $(2, -3)$ and radius 5. Its equation is

$$(x - 2)^2 + (y + 3)^2 = 25.$$

Expand the brackets:

$$x^2 - 4x + 4 + y^2 + 6y + 9 = 25,$$

and move everything to one side:

$$x^2 + y^2 - 4x + 6y - 12 = 0.$$

Look carefully at the result. The x^2 and y^2 terms both have coefficient 1; there is a term in x , a term in y , and a constant. There is no xy term. This shape is completely general: expanding $(x - a)^2 + (y - b)^2 = r^2$ always produces an equation of the form

$$x^2 + y^2 + 2gx + 2fy + c = 0,$$

where $g = -a$, $f = -b$ and $c = a^2 + b^2 - r^2$. (The factors 2 in $2g$ and $2f$ are a convenient convention: they make the completion of the square come out cleanly.)

DEFINITION General equation of a circle

The **general equation** of a circle is

$$x^2 + y^2 + 2gx + 2fy + c = 0,$$

where g , f and c are constants. The circle has

$$\text{centre } (-g, -f) \quad \text{and} \quad \text{radius } r = \sqrt{g^2 + f^2 - c}.$$

WARNING Equal coefficients of x^2 and y^2

For an equation of the second degree to represent a circle, the coefficients of x^2 and y^2 must be equal (and there must be no xy term). If you are given, say, $3x^2 + 3y^2 + 6x - 12y + 4 = 0$, divide through by 3 first before reading off g , f and c . An equation such as $2x^2 + 5y^2 = 20$ is not a circle (it is an ellipse), because the coefficients of x^2 and y^2 differ.

2.2 Recovering Centre and Radius: Completing the Square

Why does the centre come out as $(-g, -f)$? The cleanest way to see it is to reverse the expansion, that is, to complete the square in x and in y . Starting from

$$x^2 + y^2 + 2gx + 2fy + c = 0,$$

group the x -terms and the y -terms:

$$(x^2 + 2gx) + (y^2 + 2fy) = -c.$$

Recall the identity $X^2 + 2kX = (X + k)^2 - k^2$. Applying it to each group:

$$(x + g)^2 - g^2 + (y + f)^2 - f^2 = -c,$$

$$(x + g)^2 + (y + f)^2 = g^2 + f^2 - c.$$

Comparing with the centre-radius form $(x - a)^2 + (y - b)^2 = r^2$, we read off at once:

$$a = -g, \quad b = -f, \quad r^2 = g^2 + f^2 - c.$$

This derivation is short, elegant and fully examinable — a GCE examiner may well ask you to show that a given equation represents a circle and to state its centre and radius.

METHOD Centre and radius from the general form

Given $x^2 + y^2 + 2gx + 2fy + c = 0$:

1. Check that the coefficients of x^2 and y^2 are equal; if they are not 1, divide the whole equation by that common coefficient.

2. Quick route: identify g and f as half the coefficients of x and y ; then centre = $(-g, -f)$ and $r = \sqrt{g^2 + f^2 - c}$.
3. Safe route (recommended in the exam): complete the square in x and in y to rewrite the equation as $(x+g)^2 + (y+f)^2 = g^2 + f^2 - c$, then read off the centre and radius directly.

EXAMPLE Reading off the centre and radius

Find the centre and radius of the circle $x^2 + y^2 - 6x + 8y + 9 = 0$.


Quick route: $2g = -6 \Rightarrow g = -3$; $2f = 8 \Rightarrow f = 4$; $c = 9$.

$$\text{Centre} = (-g, -f) = (3, -4), \quad r = \sqrt{(-3)^2 + 4^2 - 9} = \sqrt{9 + 16 - 9} = \sqrt{16} = 4.$$

Check by completing the square:

$$(x^2 - 6x) + (y^2 + 8y) = -9 \Rightarrow (x - 3)^2 - 9 + (y + 4)^2 - 16 = -9,$$

$$(x - 3)^2 + (y + 4)^2 = 16. \quad \checkmark$$



Converting the general equation to centre-radius form by completing the square.

2.3 When Does the General Equation Represent a Real Circle?

The formula $r^2 = g^2 + f^2 - c$ hides a subtlety that the GCE Board loves to test: the quantity under the square root may be positive, zero or negative.

PROPERTY Existence condition — proof examinable

The equation $x^2 + y^2 + 2gx + 2fy + c = 0$ represents:

- a real circle of radius $\sqrt{g^2 + f^2 - c}$ if $g^2 + f^2 - c > 0$;
- a point circle (the single point $(-g, -f)$) if $g^2 + f^2 - c = 0$;

- no real locus at all if $g^2 + f^2 - c < 0$.

Proof. Completing the square gives $(x+g)^2 + (y+f)^2 = g^2 + f^2 - c$. The left-hand side, being a sum of two squares of real numbers, is always ≥ 0 . If the right-hand side is negative, no real point (x, y) can satisfy the equation. If it is zero, we need both squares to vanish, forcing $x = -g$ and $y = -f$: a single point. If it is positive, we obtain a genuine circle. ■

EXAMPLE Testing the three cases

Examine the nature of each locus:

- $x^2 + y^2 - 2x + 4y - 4 = 0$: $g^2 + f^2 - c = 1 + 4 + 4 = 9 > 0$: real circle, centre $(1, -2)$, radius 3.
- $x^2 + y^2 + 6x - 2y + 10 = 0$: $g^2 + f^2 - c = 9 + 1 - 10 = 0$: the single point $(-3, 1)$.
- $x^2 + y^2 + 2x + 2y + 5 = 0$: $g^2 + f^2 - c = 1 + 1 - 5 = -3 < 0$: no real locus.

WORKED EXAMPLE 4 Finding k for a real circle

Find the range of values of k for which the equation

$$x^2 + y^2 + 4x - 6y + k = 0$$

represents a real circle, and state what happens at the boundary value.

SOLUTION

Here $2g = 4$ so $g = 2$; $2f = -6$ so $f = -3$; $c = k$. The existence condition is

$$g^2 + f^2 - c > 0 \iff 4 + 9 - k > 0 \iff k < 13.$$

Hence the equation represents a real circle for all $k < 13$, with centre $(-2, 3)$ and radius $\sqrt{13 - k}$. When $k = 13$ the radius is 0: the locus collapses to the single point $(-2, 3)$ (a point circle). For $k > 13$ there is no real locus.

2.4 Circle on a Given Diameter

Suppose now that instead of the centre you are given the two endpoints of a diameter, $A(x_1, y_1)$ and $B(x_2, y_2)$. Two facts from Section 1 and from O-Level geometry combine beautifully here:

- the centre of the circle is the **midpoint** M of AB ;
- the radius is **half the length** of AB .

METHOD Circle with diameter AB — midpoint route

1. Compute the centre $M = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$.
2. Compute the radius $r = \frac{1}{2}|AB| = \frac{1}{2}\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$ (or find $|MA|$ directly).
3. Write the equation $(x - h)^2 + (y - k)^2 = r^2$ where $M = (h, k)$, and expand if the general form is required.

There is, however, a faster and very elegant route, based on a classical theorem: the **angle in a semicircle is a right angle**. If $P(x, y)$ is any point of the circle with diameter AB , then $\angle APB = 90^\circ$, so the vectors $\vec{PA}$ and $\vec{PB}$ are perpendicular, and their scalar product is zero.

PROPERTY Diameter form of the equation of a circle — proof examinable

The circle having $A(x_1, y_1)$ and $B(x_2, y_2)$ as endpoints of a diameter has equation

$$(x - x_1)(x - x_2) + (y - y_1)(y - y_2) = 0.$$


Proof. Let $P(x, y)$ be a point of the circle. Then

$$\vec{PA} = \begin{pmatrix} x_1 - x \\ y_1 - y \end{pmatrix}, \quad \vec{PB} = \begin{pmatrix} x_2 - x \\ y_2 - y \end{pmatrix}.$$

Since the angle in a semicircle is 90° , $\vec{PA} \perp \vec{PB}$, so $\vec{PA} \cdot \vec{PB} = 0$:

$$(x_1 - x)(x_2 - x) + (y_1 - y)(y_2 - y) = 0,$$

which is equivalent to $(x - x_1)(x - x_2) + (y - y_1)(y - y_2) = 0$ (each bracket is multiplied twice by -1). Conversely, any point satisfying this equation either coincides with A or B , or satisfies $\vec{PA} \cdot \vec{PB} = 0$, hence lies on the circle with diameter AB . ■



Circle with diameter AB: the centre M is the midpoint of AB , and for every point P on the circle, $\angle APB = 90^\circ$.

WORKED EXAMPLE 5 Circle on a given diameter — both routes

Find the equation of the circle having the segment joining $A(1, 2)$ and $B(5, -4)$ as a diameter.

SOLUTION

Route 1 (midpoint). The centre is

$$M = \left(\frac{1+5}{2}, \frac{2+(-4)}{2} \right) = (3, -1).$$

The radius is half of $|AB|$:

$$|AB| = \sqrt{(5-1)^2 + (-4-2)^2} = \sqrt{16+36} = \sqrt{52}, \quad r^2 = \frac{52}{4} = 13.$$

Hence $(x-3)^2 + (y+1)^2 = 13$, which expands to $x^2 + y^2 - 6x + 2y - 3 = 0$.

Route 2 (diameter form — the GCE shortcut). Apply $(x-x_1)(x-x_2) + (y-y_1)(y-y_2) = 0$:

$$(x-1)(x-5) + (y-2)(y+4) = 0,$$

$$x^2 - 6x + 5 + y^2 + 2y - 8 = 0 \implies x^2 + y^2 - 6x + 2y - 3 = 0. \quad \checkmark$$

Both routes agree. Notice how the diameter form produces the general equation in a single line — a valuable time-saver in the examination room.

WARNING Sign traps with the diameter form

Two errors recur every year:

- Writing $(x-x_1)(y-y_2) + (y-y_1)(x-x_2) = 0$: the x 's stay together and the y 's stay together.
- Forgetting that a negative coordinate changes the sign inside the bracket: with

$B(5, -4)$ the factor is $(y - (-4)) = (y + 4)$, not $(y - 4)$.

Also remember: the diameter form gives the circle for which AB is a diameter, not merely a chord. If the problem says “ AB is a chord”, this formula does not apply.

2.5 Mixed Techniques: A Consolidation Example

Many GCE questions force you to combine everything seen so far: expand, complete the square, test the existence condition, and use a diameter.

WORKED EXAMPLE 6 Synthesis

A circle C_1 has equation $x^2 + y^2 + 2x - 8y + 8 = 0$.

1. Find the centre and radius of C_1 .
2. A second circle C_2 has the line joining $P(-1, 3)$ and $Q(3, 5)$ as a diameter. Find the centre and radius of C_2 , and determine whether C_1 and C_2 are concentric.

SOLUTION

1. Complete the square:

$$(x^2 + 2x) + (y^2 - 8y) = -8 \implies (x + 1)^2 - 1 + (y - 4)^2 - 16 = -8,$$

$$(x + 1)^2 + (y - 4)^2 = 9.$$

Centre $(-1, 4)$, radius 3. (Check: $g^2 + f^2 - c = 1 + 16 - 8 = 9 > 0$, so a real circle indeed.)

2. The centre of C_2 is the midpoint of PQ :

$$M = \left(\frac{-1 + 3}{2}, \frac{3 + 5}{2} \right) = (1, 4).$$

This is not the same centre as C_1 , which is $(-1, 4)$ — so the two circles are not concentric. The radius of C_2 is

$$r = \frac{1}{2}|PQ| = \frac{1}{2}\sqrt{(3 - (-1))^2 + (5 - 3)^2} = \frac{1}{2}\sqrt{16 + 4} = \frac{1}{2}\sqrt{20} = \sqrt{5}.$$

Using the diameter form directly:

$$(x + 1)(x - 3) + (y - 3)(y - 5) = 0 \implies x^2 - 2x - 3 + y^2 - 8y + 15 = 0,$$

$$x^2 + y^2 - 2x - 8y + 12 = 0.$$

EXERCISE 2 Practice for the GCE

- Find the centre and radius of each circle:
 - $x^2 + y^2 - 10x + 4y + 20 = 0$;
 - $2x^2 + 2y^2 + 6x - 10y - 1 = 0$ (divide through first!).
- Determine whether $x^2 + y^2 - 4x + 6y + 14 = 0$ represents a real circle, a point, or no real locus.
- Find the values of λ for which $x^2 + y^2 + 8x - 2y + \lambda = 0$ is a real circle.
- Find the equation of the circle with diameter endpoints $A(-2, 1)$ and $B(4, 7)$, giving your answer in general form.
- A circle has equation $x^2 + y^2 - 6x + 2y + 6 = 0$. Show that the point $(5, -3)$ lies on it, and find the equation of the circle having the segment from $(5, -3)$ to the centre as diameter.

SOLUTION Exercise 1

- (a) $g = -5$, $f = 2$, $c = 20$: centre $(5, -2)$, $r = \sqrt{25 + 4 - 20} = \sqrt{9} = 3$. (b) Divide by 2: $x^2 + y^2 + 3x - 5y - \frac{1}{2} = 0$; $g = \frac{3}{2}$, $f = -\frac{5}{2}$: centre $(-\frac{3}{2}, \frac{5}{2})$, $r = \sqrt{\frac{9}{4} + \frac{25}{4} + \frac{1}{2}} = \sqrt{\frac{36}{4}} = 3$.
- $g^2 + f^2 - c = 4 + 9 - 14 = -1 < 0$: no real locus.
- $16 + 1 - \lambda > 0 \Rightarrow \lambda < 17$.
- $(x + 2)(x - 4) + (y - 1)(y - 7) = 0 \Rightarrow x^2 + y^2 - 2x - 8y - 1 = 0$.
- Centre $(3, -1)$, radius 2; $(5)^2 + (-3)^2 - 6(5) + 2(-3) + 6 = 25 + 9 - 30 - 6 + 6 = 4 = r^2$: yes, $(5, -3)$ lies on the circle. Diameter endpoints $(5, -3)$ and $(3, -1)$: $(x - 5)(x - 3) + (y + 3)(y + 1) = 0 \Rightarrow x^2 + y^2 - 8x + 4y + 18 = 0$.

KEY POINTS Key points of Section 2

- General equation: $x^2 + y^2 + 2gx + 2fy + c = 0$, with centre $(-g, -f)$ and radius $\sqrt{g^2 + f^2 - c}$.
- Real circle requires $g^2 + f^2 - c > 0$; equality gives a point circle; a negative value gives no real locus.
- Complete the square in x and in y to pass from the general form to the centre-radius form — and always divide through first if the coefficients of x^2 and y^2 are not 1.
- Circle with diameter $A(x_1, y_1)$, $B(x_2, y_2)$: centre = midpoint of AB , radius = $\frac{1}{2}|AB|$, or use the shortcut $(x - x_1)(x - x_2) + (y - y_1)(y - y_2) = 0$, justified by the right angle in a semicircle.

With the general equation now mastered, we are ready for Section 3, where we shall determine the circle passing through three given points and circles subject to other geometric conditions — problems in which the general form $x^2 + y^2 + 2gx + 2fy + c = 0$, with its three unknown constants, becomes our most powerful tool.

3 SECTION 3: CIRCLE THROUGH THREE POINTS AND CIRCLES UNDER GEOMETRIC CONDITIONS

In Section 2 we learnt to recognise and manipulate the **general equation** of a circle, $x^2 + y^2 + 2gx + 2fy + c = 0$, whose centre is $(-g, -f)$ and whose radius is $\sqrt{g^2 + f^2 - c}$ (provided $g^2 + f^2 - c > 0$). We also saw that a circle is completely determined once we know its diameter. But in many GCE problems the data are given differently: we are told points the circle must pass through, or lines it must touch, or a line on which its centre must lie. This section develops the algebraic and geometric machinery for all such situations. The central idea is simple: **three independent conditions determine a circle**, because the general equation contains exactly three unknown constants g , f and c .

3.1 Circle Through Three Non-Collinear Points

EXPERIMENT How many points fix a circle?

Take a sheet of paper. Through one given point you can draw infinitely many circles. Through two given points you can still draw infinitely many circles (their centres all lie on the perpendicular bisector of the segment joining the points). But through three non-collinear points — think of three towns on a map of the Northwest Region — exactly one circle passes. If the three points are collinear, no circle passes through them at all. This is why three points are the natural data for determining a circle.

PROPERTY Circle through three points

Through any three **non-collinear** points there passes exactly one circle. Its equation is found by substituting the coordinates of the three points into the general equation

$$x^2 + y^2 + 2gx + 2fy + c = 0$$

and solving the resulting system of three linear equations in the three unknowns g , f and c . (The algebraic technique is examinable; it is simply simultaneous equations.)

METHOD Circle through three given points

1. Write the general equation $x^2 + y^2 + 2gx + 2fy + c = 0$.
2. Substitute each of the three points in turn to obtain three linear equations in g, f, c .
3. Solve the system (elimination is usually fastest).
4. Write the equation, and if required state the centre $(-g, -f)$ and radius $\sqrt{g^2 + f^2 - c}$.
5. Check: verify that each given point satisfies the final equation.

WORKED EXAMPLE 7 Circle through three points

Find the equation of the circle passing through the points $A(1, 2)$, $B(3, -4)$ and $C(5, -6)$. State its centre and radius.

SOLUTION

Substitute each point into $x^2 + y^2 + 2gx + 2fy + c = 0$.

Through $A(1, 2)$: $1 + 4 + 2g + 4f + c = 0 \Rightarrow 2g + 4f + c = -5$ (1)

Through $B(3, -4)$: $9 + 16 + 6g - 8f + c = 0 \Rightarrow 6g - 8f + c = -25$ (2)

Through $C(5, -6)$: $25 + 36 + 10g - 12f + c = 0 \Rightarrow 10g - 12f + c = -61$ (3)

Subtract (1) from (2): $4g - 12f = -20 \Rightarrow g - 3f = -5$ (4)

Subtract (2) from (3): $4g - 4f = -36 \Rightarrow g - f = -9$ (5)

Subtracting (5) from (4): $-2f = 4$, so $f = -2$, and then $g = -9 + f = -11$.

From (1): $c = -5 - 2g - 4f = -5 + 22 + 8 = 25$.

Hence the circle is

$$x^2 + y^2 - 22x - 4y + 25 = 0,$$

with centre $(11, 2)$ and radius $\sqrt{121 + 4 - 25} = \sqrt{100} = 10$.

Check: for $A(1, 2)$: $1 + 4 - 22 - 8 + 25 = 0$ ✓; for $B(3, -4)$: $9 + 16 - 66 + 16 + 25 = 0$ ✓; for $C(5, -6)$: $25 + 36 - 110 + 24 + 25 = 0$ ✓.

3.2 The Geometric Alternative: Perpendicular Bisectors

There is a beautiful geometric reason why three points determine a circle, and it gives a second method.


PROPERTY Centre on the perpendicular bisector of a chord

The centre of a circle lies on the **perpendicular bisector** of every chord of the circle. Consequently, the centre of the circle through three non-collinear points A, B, C is the intersection of the perpendicular bisectors of any two of the segments AB, BC, CA . This point

is called the **circumcentre** of triangle ABC .

Justification. Let O be the centre and PQ a chord with midpoint M . Then $OP = OQ$ (radii), so triangles OMP and OMQ are congruent (SSS, with OM common and $PM = MQ$). Hence $\angle OMP = \angle OMQ$, and since they sum to 180° , each is 90° . So $OM \perp PQ$: the line from the centre to the midpoint of a chord is perpendicular to the chord, i.e. the centre lies on the chord's perpendicular bisector.

■



The perpendicular bisectors of chords AB and BC meet at the circumcentre O , the centre of the circle through A, B, C .

METHOD Circumcentre method

1. Find the midpoint and gradient of one chord, hence the equation of its perpendicular bisector.
2. Repeat for a second chord.
3. Solve the two bisector equations simultaneously: the intersection is the centre (h, k) .
4. The radius is the distance from (h, k) to any of the three points.
5. Write $(x - h)^2 + (y - k)^2 = r^2$.

WARNING Which method should you choose?

Both methods are accepted at GCE. The general-equation method is purely mechanical and safe. The perpendicular bisector method is often shorter when two of the points share a coordinate (the bisector is then horizontal or vertical). Whichever you use, always check your final equation against all three points.

3.3 Circle Through Two Points with Centre on a Given Line

Suppose the circle must pass through two given points A and B , and its centre must lie on a given line L . Geometrically, the centre lies on the perpendicular bisector of AB and on L : the centre is their intersection, so the circle is unique (unless the lines are parallel, in which case no such circle exists).

Algebraically, write the centre as $(-g, -f)$ and impose:

- the two points satisfy the general equation (two equations);
- the centre satisfies the equation of L (one equation).

Three equations in g, f, c : the circle is determined.

WORKED EXAMPLE 8 Centre on a given line

Find the equation of the circle passing through $A(2, 3)$ and $B(-1, 1)$ whose centre lies on the line $x - 3y - 11 = 0$.

SOLUTION

Let the circle be $x^2 + y^2 + 2gx + 2fy + c = 0$, centre $(-g, -f)$.

Through A : $4 + 9 + 4g + 6f + c = 0 \Rightarrow 4g + 6f + c = -13$ (1)

Through B : $1 + 1 - 2g + 2f + c = 0 \Rightarrow -2g + 2f + c = -2$ (2)

Centre on the line: $-g - 3(-f) - 11 = 0 \Rightarrow -g + 3f = 11$ (3)


(1) - (2): $6g + 4f = -11$ (4). From (3), $g = 3f - 11$. Substituting in (4): $6(3f - 11) + 4f = -11 \Rightarrow 22f = 55 \Rightarrow f = \frac{5}{2}, g = -\frac{7}{2}$.

From (2): $c = -2 + 2g - 2f = -2 - 7 - 5 = -14$.

$$x^2 + y^2 - 7x + 5y - 14 = 0$$

Centre $\left(\frac{7}{2}, -\frac{5}{2}\right)$, radius $\sqrt{\frac{49}{4} + \frac{25}{4} + 14} = \sqrt{\frac{130}{4}} = \frac{\sqrt{130}}{2}$.

Check: $A: 4 + 9 - 14 + 15 - 14 = 0$ ✓; $B: 1 + 1 + 7 + 5 - 14 = 0$ ✓; centre on line: $\frac{7}{2} - 3\left(-\frac{5}{2}\right) - 11 = \frac{7}{2} + \frac{15}{2} - 11 = 0$ ✓.



The centre is forced to lie on the line $x - 3y - 11 = 0$; together with the two points A and B this fixes a unique circle.

3.4 Circles Touching a Given Straight Line

PROPERTY Touching condition

A circle with centre (h, k) and radius r **touches** the line $ax + by + c = 0$ if and only if the perpendicular distance from the centre to the line equals the radius:

$$\frac{|ah + bk + c|}{\sqrt{a^2 + b^2}} = r.$$

Because of the modulus, this condition usually produces two possible values, one for each side of the line.

METHOD 'Circle touches line' problems

1. Express the centre (h, k) using the other given conditions (e.g. centre on a line, or circle through a point).
2. Write the touching condition $\frac{|ah + bk + c|}{\sqrt{a^2 + b^2}} = r$.
3. Remove the modulus by considering both signs; solve.

4. For each solution, write the equation and check it satisfies all the given data.

WORKED EXAMPLE 9 Through two points and touching a line

Find the equations of the circles which pass through $A(1, 4)$ and $B(3, 2)$ and touch the x -axis.

SOLUTION

The centre lies on the perpendicular bisector of AB . Midpoint of AB : $(2, 3)$; gradient of AB : $\frac{2-4}{3-1} = -1$, so the bisector has gradient 1:

$$y - 3 = 1(x - 2) \Rightarrow y = x + 1.$$

Write the centre as $(h, h+1)$. Touching the x -axis means the radius equals the distance from the centre to $y = 0$, i.e. $r = |h + 1|$.

The circle passes through $A(1, 4)$:

$$(1 - h)^2 + (4 - (h + 1))^2 = (h + 1)^2$$

$$(1 - h)^2 + (3 - h)^2 = (h + 1)^2$$

$$1 - 2h + h^2 + 9 - 6h + h^2 = h^2 + 2h + 1$$

$$h^2 - 10h + 9 = 0 \Rightarrow (h - 1)(h - 9) = 0.$$

Case 1: $h = 1$, centre $(1, 2)$, $r = 2$:

$$(x - 1)^2 + (y - 2)^2 = 4.$$

Case 2: $h = 9$, centre $(9, 10)$, $r = 10$:

$$(x - 9)^2 + (y - 10)^2 = 100.$$

Check (Case 1): $A: 0 + 4 = 4$ ✓; $B: 4 + 0 = 4$ ✓; distance from $(1, 2)$ to the x -axis is $2 = r$ ✓.

Check (Case 2): $A: 64 + 36 = 100$ ✓; $B: 36 + 64 = 100$ ✓; distance from $(9, 10)$ to the x -axis is $10 = r$ ✓. Both circles are valid answers.

WARNING Two solutions are the rule, not the exception

Problems involving touching a line almost always yield two circles, because the centre can lie on either side of the line (or the radius can correspond to two positions). Never stop after the first solution: solve the equation completely, and only reject a solution if it contradicts a stated condition (e.g. “the circle lies in the first quadrant”).


3.5 Circles Touching the Coordinate Axes

PROPERTY Circle touching both axes

A circle of radius r touching both coordinate axes has its centre at distance r from each axis, so the centre is $(\pm r, \pm r)$, the signs depending on the quadrant. Its equation is

$$(x \pm r)^2 + (y \pm r)^2 = r^2.$$

In the first quadrant: centre (r, r) , equation $(x - r)^2 + (y - r)^2 = r^2$.



A circle of radius r touching both axes in the first quadrant has centre (r, r) : each radius to an axis is perpendicular to that axis.

EXAMPLE Touching both axes with an extra condition

Find the equation of the circle in the first quadrant which touches both axes and passes through the point $(4, 2)$.

Solution. Centre (r, r) , radius r . Substituting $(4, 2)$:

$$(4 - r)^2 + (2 - r)^2 = r^2 \Rightarrow 16 - 8r + r^2 + 4 - 4r + r^2 = r^2$$

$$r^2 - 12r + 20 = 0 \Rightarrow (r - 2)(r - 10) = 0.$$

Both $r = 2$ and $r = 10$ give first-quadrant circles through $(4, 2)$:

$$(x - 2)^2 + (y - 2)^2 = 4 \quad \text{and} \quad (x - 10)^2 + (y - 10)^2 = 100.$$

3.6 Strategy: Standard Form or General Form?

KEY POINTS Choosing the right form

- Use the standard form $(x - h)^2 + (y - k)^2 = r^2$ when the data involve the centre or radius directly: centre on a line, touching a line (distance condition), touching the axes.
- Use the general form $x^2 + y^2 + 2gx + 2fy + c = 0$ when the data are points on the circle: substitution gives linear equations in g, f, c , which are easy to solve.
- Mixed data (points + centre condition): either form works; the general form with the centre condition $-g, -f$ substituted into the line equation is usually cleanest.
- Remember: three independent conditions are always needed to determine a circle.

EXERCISE 3 Consolidation

1. Find the equation of the circle through $(0, 0)$, $(4, 0)$ and $(2, 6)$, and state its centre and radius.
2. Find the equation of the circle through $(3, 1)$ and $(5, 3)$ whose centre lies on the line $y = x + 2$.
3. Find the equations of the circles which touch the line $3x + 4y = 10$, have radius 5, and whose centres lie on the x -axis.
4. A circle touches both coordinate axes and passes through $(1, 2)$. Find its possible equations.

SOLUTION Consolidation Exercises

1. Substituting into $x^2 + y^2 + 2gx + 2fy + c = 0$: $(0, 0)$ gives $c = 0$; $(4, 0)$ gives $16 + 8g = 0$, so $g = -2$; $(2, 6)$ gives $40 + 4g + 12f = 0$, so $12f = -40 + 8 = -32$, $f = -\frac{8}{3}$. Equation: $x^2 + y^2 - 4x - \frac{16}{3}y = 0$; centre $(2, \frac{8}{3})$, radius $\sqrt{4 + \frac{64}{9}} = \frac{10}{3}$.
2. Centre $(-g, -f)$ on $y = x + 2$: $-f = -g + 2 \Rightarrow f = g - 2$. Points give $6g + 2f + c = -10$ and $10g + 6f + c = -34$; subtracting: $4g + 4f = -24$, i.e. $g + f = -6$. With $f = g - 2$: $2g - 2 = -6 \Rightarrow g = -2$, $f = -4$, then $c = -10 + 12 + 8 = 10$. Equation: $x^2 + y^2 - 4x - 8y + 10 = 0$; centre $(2, 4)$, radius $\sqrt{10}$.
3. Centre $(h, 0)$, radius 5: $\frac{|3h - 10|}{5} = 5 \Rightarrow |3h - 10| = 25 \Rightarrow h = \frac{35}{3}$ or $h = -5$. Circles: $(x + 5)^2 + y^2 = 25$ and $(x - \frac{35}{3})^2 + y^2 = 25$.
4. Centre (r, r) (the point $(1, 2)$ is in the first quadrant): $(1 - r)^2 + (2 - r)^2 = r^2 \Rightarrow r^2 - 6r + 5 = 0 \Rightarrow r = 1$ or $r = 5$. Circles: $(x - 1)^2 + (y - 1)^2 = 1$ and $(x - 5)^2 + (y - 5)^2 = 25$.

DID YOU KNOW? Circumcircles in real life

The circle through three points is exactly what surveyors and engineers use to reconstruct a circular arc from three measured points — for instance when designing a roundabout or a curved section of road. The circumcentre of a triangle is also one of the four classical “centres” of a triangle studied since Euclid, alongside the centroid, the orthocentre and the incentre.

4 SECTION 4: TANGENTS TO A CIRCLE

In Section 3 we learnt how to determine a circle from geometric data: three points, a diameter, or conditions involving lines and other circles. We now change our point of view. Instead of building circles, we study one of the most important lines attached to a circle: the **tangent**. Tangents are everywhere in the GCE Advanced Level examination: they appear in coordinate geometry questions, in locus problems, and in the study of families of circles. Master this section and a large family of exam questions becomes routine.

4.1 What is a Tangent?**EXPERIMENT** From a secant to a tangent

Draw a circle and a straight line. Depending on the position of the line, three things can happen:

- the line misses the circle completely (no intersection);
- the line cuts the circle in two distinct points — such a line is called a **secant**;
- the line meets the circle in exactly one point.

Now imagine a secant through two points A and B of the circle, and let B slide along the circle towards A . The two intersection points merge into one, and the secant settles into a limiting position: the tangent at A .


DEFINITION Tangent to a circle

A **tangent** to a circle is a straight line which meets the circle in exactly one point. That point is called the **point of contact**. A line which cuts the circle in two distinct points is a **secant**; a line which does not meet the circle is said to be exterior to it.

PROPERTY Radius–tangent perpendicularity

The tangent at any point of a circle is perpendicular to the radius drawn to the point of contact. (This result is examinable and its proof is instructive.)

Proof. Let the circle have centre C and radius r , and let T be the point of contact of a tangent line ℓ . Take any other point Q on ℓ , $Q \neq T$. Since ℓ meets the circle only at T , the point Q lies outside the circle, so $CQ > r = CT$. Thus CT is the shortest distance from C to the line ℓ . But the shortest distance from a point to a line is the perpendicular distance. Hence $CT \perp \ell$. ■



The tangent at P is perpendicular to the radius CP .

4.2 Tangent to the Circle $x^2 + y^2 = r^2$ at a Given Point

We now turn the geometric property into an algebraic formula. Consider the circle $x^2 + y^2 = r^2$ with centre the origin O , and a point $P(x_1, y_1)$ on the circle, so that

$$x_1^2 + y_1^2 = r^2.$$

The radius OP has gradient $\frac{y_1}{x_1}$ (provided $x_1 \neq 0$; the cases $x_1 = 0$ or $y_1 = 0$ give vertical or horizontal tangents which are easily handled separately). Since the tangent is perpendicular to OP , its gradient is $-\frac{x_1}{y_1}$. The tangent through $P(x_1, y_1)$ therefore has equation

$$y - y_1 = -\frac{x_1}{y_1}(x - x_1).$$

Multiplying by y_1 and rearranging:

$$yy_1 - y_1^2 = -xx_1 + x_1^2 \quad \Rightarrow \quad xx_1 + yy_1 = x_1^2 + y_1^2 = r^2.$$

PROPERTY Tangent to $x^2 + y^2 = r^2$ at (x_1, y_1)

The equation of the tangent to the circle $x^2 + y^2 = r^2$ at the point $P(x_1, y_1)$ on the circle is

$$xx_1 + yy_1 = r^2$$

(The derivation above, using the gradient of the radius and the perpendicular gradient of the tangent, is examinable.)

Notice the beautiful pattern: to obtain the tangent at (x_1, y_1) , replace one x by x_1 and one y by y_1 in $x^2 + y^2 = r^2$. This is the simplest case of the replace rule that we shall generalise below.

WARNING Check that the point is on the circle!

The formula $xx_1 + yy_1 = r^2$ is valid only when (x_1, y_1) actually lies on the circle. Always verify that $x_1^2 + y_1^2 = r^2$ before applying it. If the point is not on the circle, the line $xx_1 + yy_1 = r^2$ is still a perfectly good line (it is in fact the polar of the point with respect to the circle), but it is not a tangent through that point.

WORKED EXAMPLE 10 Tangent at a point on $x^2 + y^2 = 25$

Find the equation of the tangent to the circle $x^2 + y^2 = 25$ at the point $P(3, 4)$.

SOLUTION

Solution. First check: $3^2 + 4^2 = 9 + 16 = 25$, so P lies on the circle. Applying $xx_1 + yy_1 = r^2$ with $(x_1, y_1) = (3, 4)$ and $r^2 = 25$:

$$3x + 4y = 25.$$

Verification by gradients: the radius OP has gradient $\frac{4}{3}$; the line $3x + 4y = 25$ has gradient $-\frac{3}{4}$, and $\frac{4}{3} \times (-\frac{3}{4}) = -1$. Perpendicular, as required.

4.3 Tangent to a General Circle: the Replace Rule

For the circle $(x - a)^2 + (y - b)^2 = r^2$ with centre $C(a, b)$, the same gradient argument applies. The radius from C to $P(x_1, y_1)$ has gradient $\frac{y_1 - b}{x_1 - a}$, so the tangent has gradient $-\frac{x_1 - a}{y_1 - b}$, giving

$$(x - x_1)(x_1 - a) + (y - y_1)(y_1 - b) = 0,$$

which simplifies, using $(x_1 - a)^2 + (y_1 - b)^2 = r^2$, to

$$(x - a)(x_1 - a) + (y - b)(y_1 - b) = r^2.$$

For the general circle

$$x^2 + y^2 + 2gx + 2fy + c = 0 \quad (\text{centre } (-g, -f), \text{ radius } r = \sqrt{g^2 + f^2 - c}),$$

expanding the previous formula with $a = -g$, $b = -f$ and $r^2 = g^2 + f^2 - c$ leads to the following memorable rule.

PROPERTY The replace rule for tangents

The tangent at $P(x_1, y_1)$ to the circle $x^2 + y^2 + 2gx + 2fy + c = 0$ is

$$xx_1 + yy_1 + g(x + x_1) + f(y + y_1) + c = 0$$

obtained from the equation of the circle by the replacements

$$x^2 \rightarrow xx_1, \quad y^2 \rightarrow yy_1, \quad 2gx \rightarrow g(x + x_1), \quad 2fy \rightarrow f(y + y_1), \quad c \rightarrow c.$$

EXAMPLE Using the replace rule

Find the tangent to the circle $x^2 + y^2 - 4x + 6y - 12 = 0$ at the point $(5, 1)$.

Check that the point lies on the circle:

$$5^2 + 1^2 - 4(5) + 6(1) - 12 = 25 + 1 - 20 + 6 - 12 = 0.$$

Good. Here $2g = -4 \Rightarrow g = -2$, $2f = 6 \Rightarrow f = 3$, $c = -12$. Apply the replace rule:

$$5x + 1 \cdot y + (-2)(x + 5) + 3(y + 1) - 12 = 0$$

$$5x + y - 2x - 10 + 3y + 3 - 12 = 0$$

$$3x + 4y - 19 = 0.$$

DID YOU KNOW? The polar line

If $P(x_1, y_1)$ is outside the circle, the line $xx_1 + yy_1 + g(x + x_1) + f(y + y_1) + c = 0$ is called the polar of P with respect to the circle. It passes through the two points of contact of the tangents drawn from P to the circle. This is a powerful tool in projective geometry and appears in advanced locus problems.

4.4 Condition for a Line to be Tangent to a Circle

Suppose now that we are given a line and a circle and we ask: is the line a tangent? Geometrically the answer is immediate.

METHOD Tangency test: distance from centre equals radius

A line ℓ is tangent to a circle if and only if the **perpendicular distance** from the centre of the circle to ℓ equals the radius.

1. Write the line in the form $Ax + By + C = 0$ and identify the centre (a, b) and radius r of the circle.
2. Compute $d = \frac{|Aa + Bb + C|}{\sqrt{A^2 + B^2}}$.
3. If $d = r$: tangent (one intersection). If $d < r$: secant (two intersections). If $d > r$: no intersection.

Apply this to the line $y = mx + c$, i.e. $mx - y + c = 0$, and the circle $x^2 + y^2 = r^2$ centred at the origin:

$$\frac{|c|}{\sqrt{m^2 + 1}} = r \implies \boxed{c^2 = r^2(1 + m^2)}.$$

PROPERTY Tangency condition for $y = mx + c$ and $x^2 + y^2 = r^2$

The line $y = mx + c$ is tangent to the circle $x^2 + y^2 = r^2$ if and only if $c^2 = r^2(1 + m^2)$. Consequently the lines

$$y = mx \pm r\sqrt{1 + m^2}$$


are the two tangents of gradient m to the circle.

PROPERTY Tangency condition for a general circle

The line $Ax + By + C = 0$ is tangent to the circle $x^2 + y^2 + 2gx + 2fy + c = 0$ (centre $(-g, -f)$, radius $r = \sqrt{g^2 + f^2 - c}$) if and only if

$$\frac{|A(-g) + B(-f) + C|}{\sqrt{A^2 + B^2}} = \sqrt{g^2 + f^2 - c}.$$

Equivalently, substituting $y = mx + c$ into the circle's equation yields a quadratic in x whose discriminant is zero.



Lines $y = x + c$ and the circle $x^2 + y^2 = 9$. Only $c = \pm 3\sqrt{2}$ (orange) give tangents, since $c^2 = 9(1 + 1^2) = 18$.

WORKED EXAMPLE 11 Showing a line is a tangent

Show that the line $3x + 4y = 30$ is a tangent to the circle $x^2 + y^2 + 2x - 4y - 20 = 0$, and find the point of contact.

SOLUTION

Solution. Complete the square for the circle:

$$(x + 1)^2 + (y - 2)^2 = 1 + 4 + 20 = 25,$$

so the centre is $C(-1, 2)$ and the radius is $r = 5$. Distance from C to the line $3x + 4y - 30 = 0$:

$$d = \frac{|3(-1) + 4(2) - 30|}{\sqrt{3^2 + 4^2}} = \frac{|-3 + 8 - 30|}{5} = \frac{25}{5} = 5 = r.$$

Hence the line is a tangent.

Point of contact: it is the foot of the perpendicular from $C(-1, 2)$ to the line. The perpendicular through C has direction vector $(3, 4)$ (the normal of the line). Its parametric form: $(-1 + 3t, 2 + 4t)$. Substitute into $3x + 4y = 30$:

$$3(-1 + 3t) + 4(2 + 4t) = 30 \implies -3 + 9t + 8 + 16t = 30 \implies 25t + 5 = 30 \implies t = 1.$$

Point of contact: $(2, 6)$. Check: $2^2 + 6^2 + 2(2) - 4(6) - 20 = 4 + 36 + 4 - 24 - 20 = 0$. ✓

4.5 Tangents from an External Point

Place a point P outside a circle. How many tangents can you draw from P to the circle? A sketch convinces us immediately: exactly **two**, and they are symmetric about the line CP .

WARNING Two tangents, not one!

From an external point there are exactly two tangents to a circle. When an exam question asks for “the tangents from P ”, do not stop after finding one — the second one carries marks too. Algebraically this appears as a quadratic in the gradient m ; both roots must be reported.

METHOD Finding the tangents from an external point (gradient method)

To find the tangents from an external point $P(x_1, y_1)$ to a circle with centre C and radius r :

1. Write a variable line through P : $y - y_1 = m(x - x_1)$, rearranged as $mx - y + (y_1 - mx_1) = 0$.
2. Impose the tangency condition: perpendicular distance from C to the line equals r .
3. Solve the resulting quadratic in m ; each root gives one tangent.

WORKED EXAMPLE 12 Tangents from an external point

Find the equations of the tangents from $P(5, 5)$ to the circle $x^2 + y^2 = 10$.

SOLUTION

Solution. Note $5^2 + 5^2 = 50 > 10$, so P is external. A line through P : $y - 5 = m(x - 5)$, i.e. $mx - y + 5 - 5m = 0$. Distance from $O(0, 0)$ must equal $\sqrt{10}$:

$$\frac{|5 - 5m|}{\sqrt{m^2 + 1}} = \sqrt{10} \implies 25(1 - m)^2 = 10(m^2 + 1) \implies 15m^2 - 50m + 15 = 0 \implies 3m^2 - 10m + 3 = 0.$$

Factorising: $(3m - 1)(m - 3) = 0$, so $m = 3$ or $m = \frac{1}{3}$. The two tangents are

$$y - 5 = 3(x - 5) \implies 3x - y - 10 = 0, \quad y - 5 = \frac{1}{3}(x - 5) \implies x - 3y + 10 = 0.$$

4.6 Length of the Tangent from an External Point

Let P be an external point, T the point of contact of a tangent from P , and C the centre. By the radius–tangent property, $\angle CTP = 90^\circ$, so triangle CPT is right-angled at T . Pythagoras gives

$$CP^2 = CT^2 + PT^2 = r^2 + PT^2.$$

PROPERTY Length of tangent from an external point


The length of the tangent from an external point P to a circle with centre C and radius r is

$$PT = \sqrt{CP^2 - r^2}$$

For the general circle $x^2 + y^2 + 2gx + 2fy + c = 0$ and $P(x_1, y_1)$, since $CP^2 - r^2 = (x_1 + g)^2 + (y_1 + f)^2 - (g^2 + f^2 - c)$, this becomes

$$PT = \sqrt{x_1^2 + y_1^2 + 2gx_1 + 2fy_1 + c}.$$

So the tangent length is simply the square root of the result of substituting P into the left-hand side of the circle's equation (the power of the point).



The right triangle CPT : $PT = \sqrt{CP^2 - r^2}$, and both tangents from P have the same length.

EXAMPLE Length of a tangent

Find the length of the tangent from $P(7, 1)$ to the circle $x^2 + y^2 - 4x + 2y - 4 = 0$.

Solution. Substitute P into the left-hand side:

$$7^2 + 1^2 - 4(7) + 2(1) - 4 = 49 + 1 - 28 + 2 - 4 = 20.$$

Hence $PT = \sqrt{20} = 2\sqrt{5}$. Check via centre and radius: $C = (2, -1)$, $r^2 = 4 + 1 + 4 = 9$; $CP^2 = (7 - 2)^2 + (1 + 1)^2 = 25 + 4 = 29$; $PT = \sqrt{29 - 9} = \sqrt{20}$. ✓

KEY POINTS Key results of Section 4

- The tangent at P is perpendicular to the radius CP .
- Tangent to $x^2 + y^2 = r^2$ at (x_1, y_1) : $xx_1 + yy_1 = r^2$.

- Replace rule for $x^2 + y^2 + 2gx + 2fy + c = 0$ at (x_1, y_1) : $x^2 \rightarrow xx_1$, $y^2 \rightarrow yy_1$, $2gx \rightarrow g(x + x_1)$, $2fy \rightarrow f(y + y_1)$, $c \rightarrow c$.
- Tangency condition: distance from centre to line = radius. For $y = mx + c$ and $x^2 + y^2 = r^2$: $c^2 = r^2(1 + m^2)$.
- From an external point: exactly two tangents, of equal length $\sqrt{CP^2 - r^2} = \sqrt{x_1^2 + y_1^2 + 2gx_1 + 2fy_1 + c}$.

EXERCISE 4 Practice

1. Find the tangent to $x^2 + y^2 = 29$ at $(5, -2)$.
2. Show that $y = 2x + 5\sqrt{5}$ touches $x^2 + y^2 = 25$.
3. Find the tangents from $(0, 5)$ to $x^2 + y^2 = 5$ and their lengths.
4. Find the tangent to $x^2 + y^2 - 6x - 8y = 0$ at the origin.
5. A circle has centre $(2, -3)$ and radius 4. Find the equations of the tangents of gradient 1 to this circle.

SOLUTION Exercise 1

1. Check: $5^2 + (-2)^2 = 25 + 4 = 29$. Tangent: $5x - 2y = 29$.
2. $c^2 = (5\sqrt{5})^2 = 125$; $r^2(1 + m^2) = 25(1 + 4) = 125$. Condition satisfied $\Rightarrow$ tangent.
3. Line through $(0, 5)$: $y = mx + 5$. Distance from $(0, 0)$ to $mx - y + 5 = 0$ is $\frac{5}{\sqrt{m^2 + 1}} = \sqrt{5} \Rightarrow m^2 + 1 = 5 \Rightarrow m = \pm 2$. Tangents: $y = 2x + 5$ and $y = -2x + 5$. Length = $\sqrt{0^2 + 5^2 - 5} = \sqrt{20} = 2\sqrt{5}$.
4. Circle: $x^2 + y^2 - 6x - 8y = 0$. At $(0, 0)$: replace rule gives $-3x - 4y = 0$, i.e. $3x + 4y = 0$.
5. Circle: $(x - 2)^2 + (y + 3)^2 = 16$. Tangents of gradient 1: $y = x + c$. Distance from $(2, -3)$ to $x - y + c = 0$ is $\frac{|2 + 3 + c|}{\sqrt{2}} = 4 \Rightarrow |c + 5| = 4\sqrt{2} \Rightarrow c = -5 \pm 4\sqrt{2}$. Tangents: $y = x - 5 + 4\sqrt{2}$ and $y = x - 5 - 4\sqrt{2}$.

5**SECTION 5: PARAMETRIC EQUATIONS OF A CIRCLE**

In Section 4 we studied tangents to a circle, and you will have noticed that the algebra quickly

became heavy: substituting a line into the circle's equation, solving quadratics, imposing discriminant conditions. Much of that heaviness comes from the fact that the Cartesian equation $x^2 + y^2 = r^2$ describes the circle implicitly: x and y are tangled together in one relation. In this section we learn a second, very powerful way of describing a circle: we give x and y separately, each as a function of a single variable θ called a **parameter**. This **parametric form** will allow us to write down a "general point" on the circle and will turn many difficult problems (maximum and minimum distances, tangents, loci) into simple trigonometry.

5.1 The Idea of a Parameter: From Observation to Definition

EXPERIMENT A point moving around a circle

Imagine a small bead sliding at constant speed around a circular wire of radius r centred at the origin O , like the hand of a clock at the Biyem-Assi stadium scoreboard, but turning anticlockwise. At any instant, the position of the bead is completely known once we know the angle θ that the radius OP makes with the positive x -axis. Drop a perpendicular from P to the x -axis, meeting it at M . In the right-angled triangle OMP :

$$\cos \theta = \frac{OM}{OP} = \frac{x}{r}, \quad \sin \theta = \frac{MP}{OP} = \frac{y}{r}.$$

Hence $x = r \cos \theta$ and $y = r \sin \theta$. So the single number θ determines the whole position (x, y) of the bead.

This observation is the heart of the section. Instead of one equation linking x and y , we describe the circle by two equations, each giving one coordinate in terms of θ .

DEFINITION Parametric equations of a circle

The **parametric equations** of the circle with centre (a, b) and radius r are

$$x = a + r \cos \theta, \quad y = b + r \sin \theta,$$

where θ is the **parameter**. In particular, for the circle centred at the origin,


$$x = r \cos \theta, \quad y = r \sin \theta.$$

As θ varies from 0 to 2π , the point (x, y) describes the whole circle exactly once (anticlockwise).

WARNING Where is θ measured?

The parameter θ is the angle measured at the centre of the circle, from the direction parallel to the positive x -axis, to the radius joining the centre to the point. It is not measured

at the origin, unless the centre happens to be the origin. For the circle $(x - 2)^2 + (y + 1)^2 = 9$, the angle θ is measured at the point $(2, -1)$.



The parameter θ is the angle at the centre $C(a, b)$ between the positive x -direction and the radius CP .

5.2 From Parametric to Cartesian Form

Suppose we are given parametric equations and we want the ordinary (Cartesian) equation of the circle. The tool is the fundamental identity $\cos^2 \theta + \sin^2 \theta = 1$.

METHOD Eliminating the parameter

To convert parametric equations to Cartesian form:

1. Rearrange the two equations to isolate $\cos \theta$ and $\sin \theta$:

$$\cos \theta = \frac{x - a}{r}, \quad \sin \theta = \frac{y - b}{r}.$$

2. Substitute into the identity $\cos^2 \theta + \sin^2 \theta = 1$:

$$\left(\frac{x - a}{r}\right)^2 + \left(\frac{y - b}{r}\right)^2 = 1.$$

3. Multiply through by r^2 to obtain $(x - a)^2 + (y - b)^2 = r^2$.

This calculation also proves that the parametric equations really do describe the circle: every point given by the parametric form satisfies the Cartesian equation, and conversely every point of the circle arises for some θ .

EXAMPLE Parametric to Cartesian

Convert to Cartesian form: $x = 3 + 2 \cos \theta$, $y = -1 + 2 \sin \theta$.

Solution. Isolate the trigonometric functions: $\cos \theta = \frac{x-3}{2}$ and $\sin \theta = \frac{y+1}{2}$. Using $\cos^2 \theta + \sin^2 \theta = 1$:

$$\frac{(x-3)^2}{4} + \frac{(y+1)^2}{4} = 1 \implies (x-3)^2 + (y+1)^2 = 4.$$

This is the circle with centre $(3, -1)$ and radius 2.

5.3 From Cartesian to Parametric Form

The reverse journey is just as important. Given $(x-a)^2 + (y-b)^2 = r^2$, divide by r^2 :

$$\left(\frac{x-a}{r}\right)^2 + \left(\frac{y-b}{r}\right)^2 = 1.$$

Since the two squares sum to 1, we may call the first $\cos^2 \theta$ and the second $\sin^2 \theta$, giving $x = a + r \cos \theta$, $y = b + r \sin \theta$.

EXAMPLE Cartesian to parametric


Write parametric equations for the circle $x^2 + y^2 - 4x + 6y - 12 = 0$.

Solution. Complete the square:

$$(x-2)^2 - 4 + (y+3)^2 - 9 - 12 = 0 \implies (x-2)^2 + (y+3)^2 = 25.$$

The centre is $(2, -3)$ and the radius is 5. Hence parametric equations are

$$x = 2 + 5 \cos \theta, \quad y = -3 + 5 \sin \theta.$$



The circle $x = 3 \cos \theta$, $y = 3 \sin \theta$ drawn parametrically, with points labelled by their values of θ .

5.4 The General Point on a Circle: A Powerful Technique

METHOD Using a general point on the circle

Any point on the circle with centre (a, b) and radius r may be written as

$$P = (a + r \cos \theta, b + r \sin \theta).$$

When a problem involves “a point moving on the circle”, or asks for a maximum or minimum, or a locus, write the point in this form and work with the single variable θ . Distances, gradients and midpoints then become trigonometric expressions, which are often much easier to handle than Cartesian algebra.

WORKED EXAMPLE 13 Maximum and minimum distance from a point to a circle

A point P moves on the circle $x^2 + y^2 = 4$. Find the greatest and least values of the distance from P to the fixed point $A(5, 0)$.

SOLUTION

Solution. Write $P = (2 \cos \theta, 2 \sin \theta)$. Then

$$\begin{aligned} AP^2 &= (2 \cos \theta - 5)^2 + (2 \sin \theta)^2 \\ &= 4 \cos^2 \theta - 20 \cos \theta + 25 + 4 \sin^2 \theta \\ &= 4(\cos^2 \theta + \sin^2 \theta) - 20 \cos \theta + 25 \\ &= 29 - 20 \cos \theta. \end{aligned}$$

Since $-1 \leq \cos \theta \leq 1$, the expression $29 - 20 \cos \theta$ is greatest when $\cos \theta = -1$ and least when $\cos \theta = 1$:

$$AP_{\max}^2 = 29 + 20 = 49 \Rightarrow AP_{\max} = 7, \quad AP_{\min}^2 = 29 - 20 = 9 \Rightarrow AP_{\min} = 3.$$

Geometrically, these are the distances from A to the far and near ends of the diameter through A . Notice how the parametric form reduced the problem to the boundedness of $\cos \theta$ — no calculus, no discriminants.

WORKED EXAMPLE 14 A locus found parametrically

$P = (3 \cos \theta, 3 \sin \theta)$ moves on the circle $x^2 + y^2 = 9$, and M is the midpoint of P and the fixed point $B(7, 0)$. Find the locus of M .

SOLUTION

Solution. The coordinates of M are

$$M = \left(\frac{3 \cos \theta + 7}{2}, \frac{3 \sin \theta + 0}{2} \right).$$

Write $M = (X, Y)$, so $X = \frac{7 + 3 \cos \theta}{2}$ and $Y = \frac{3 \sin \theta}{2}$. Isolate the trigonometric functions:

$$\cos \theta = \frac{2X - 7}{3}, \quad \sin \theta = \frac{2Y}{3}.$$

Applying $\cos^2 \theta + \sin^2 \theta = 1$:

$$\frac{(2X - 7)^2}{9} + \frac{4Y^2}{9} = 1 \Rightarrow (2X - 7)^2 + 4Y^2 = 9 \Rightarrow \left(X - \frac{7}{2}\right)^2 + Y^2 = \frac{9}{4}.$$

The locus of M is the circle with centre $\left(\frac{7}{2}, 0\right)$ and radius $\frac{3}{2}$ — exactly what we would expect, since a midpoint construction shrinks the original circle by a factor of $\frac{1}{2}$ about B .

WORKED EXAMPLE 15 Tangent at a parametric point

Find the equation of the tangent to the circle $x^2 + y^2 = 25$ at the point $P = (5 \cos \theta, 5 \sin \theta)$, and deduce the tangent at the point where $\theta = \frac{\pi}{3}$.

SOLUTION

Solution. The radius OP has gradient $\frac{5 \sin \theta}{5 \cos \theta} = \tan \theta$, so the tangent, being perpendicular to OP , has gradient $-\cot \theta$ (for $\sin \theta \neq 0$). Its equation is

$$y - 5 \sin \theta = -\frac{\cos \theta}{\sin \theta} (x - 5 \cos \theta).$$

Multiplying by $\sin \theta$ and rearranging:

$$x \cos \theta + y \sin \theta = 5(\cos^2 \theta + \sin^2 \theta) = 5.$$

This elegant result, $x \cos \theta + y \sin \theta = 5$, is the tangent at the point with parameter θ . For $\theta = \frac{\pi}{3}$: $\cos \theta = \frac{1}{2}$, $\sin \theta = \frac{\sqrt{3}}{2}$, giving

$$\frac{x}{2} + \frac{\sqrt{3}y}{2} = 5 \implies x + \sqrt{3}y = 10.$$

WORKED EXAMPLE 16 Maximum distance from a point to a circle — general case

A point P moves on the circle $x^2 + y^2 = 1$. Find the greatest and least values of the distance PQ , where $Q = (3, 4)$.

SOLUTION

Solution. Write $P = (\cos \theta, \sin \theta)$. Then

$$\begin{aligned} PQ^2 &= (\cos \theta - 3)^2 + (\sin \theta - 4)^2 \\ &= \cos^2 \theta - 6 \cos \theta + 9 + \sin^2 \theta - 8 \sin \theta + 16 \\ &= (\cos^2 \theta + \sin^2 \theta) + 25 - 6 \cos \theta - 8 \sin \theta \\ &= 26 - 6 \cos \theta - 8 \sin \theta. \end{aligned}$$

We express $6 \cos \theta + 8 \sin \theta$ in the form $R \cos(\theta - \alpha)$: $R = \sqrt{6^2 + 8^2} = 10$, $\cos \alpha = \frac{6}{10} = \frac{3}{5}$, $\sin \alpha = \frac{8}{10} = \frac{4}{5}$. Thus $6 \cos \theta + 8 \sin \theta = 10 \cos(\theta - \alpha)$, and since $-1 \leq \cos(\theta - \alpha) \leq 1$,

$$-10 \leq 6 \cos \theta + 8 \sin \theta \leq 10.$$

Hence

$$26 - 10 \leq PQ^2 \leq 26 + 10 \implies 16 \leq PQ^2 \leq 36.$$

Therefore $PQ_{\min} = 4$ and $PQ_{\max} = 6$.

WARNING Common mistakes with parametric forms

- Wrong centre. For $x = 2 + 5 \cos \theta$, $y = -3 + 5 \sin \theta$, the centre is $(2, -3)$, not $(-2, 3)$ and not $(2, 3)$. Read the signs directly from the parametric equations.
- Forgetting the radius in front of $\cos \theta$ and $\sin \theta$. The pair $x = a + \cos \theta$, $y = b + \sin \theta$ describes a circle of radius 1, whatever the values of a and b .
- Mixing up the two coefficients. If $x = 1 + 3 \cos \theta$ but $y = 2 + 4 \sin \theta$, the locus is not a circle (it is an ellipse); a circle requires the same coefficient r in both equations.
- Measuring θ at the origin. Remember: θ is always measured at the centre of the circle.
- Sign slips when eliminating θ . From $y = b + r \sin \theta$ we get $\sin \theta = \frac{y-b}{r}$: subtract b before dividing by r .

KEY POINTS GCE exam technique: when parametric form wins

In the GCE Advanced Level examination, watch for questions asking for the greatest or least distance from a fixed point or line to a circle, or for a locus involving a moving point on a circle. These are usually fastest by parametric form:

1. Write the moving point as $(a + r \cos \theta, b + r \sin \theta)$.
2. Express the required quantity (distance, midpoint, gradient) in terms of θ .
3. Use $-1 \leq \cos \theta \leq 1$, $-1 \leq \sin \theta \leq 1$, or $\cos^2 \theta + \sin^2 \theta = 1$ to finish.

This route routinely saves several lines of algebra and avoids the discriminant computations of Section 4. Also memorise the tangent at a parametric point of $x^2 + y^2 = r^2$:

$$x \cos \theta + y \sin \theta = r.$$

For a circle $(x - a)^2 + (y - b)^2 = r^2$, the tangent at the point with parameter θ is

$$(x - a) \cos \theta + (y - b) \sin \theta = r.$$

DID YOU KNOW? Why engineers love parameters

Parametric descriptions are not just an exam trick. When a CNC machine in a Douala workshop cuts a circular metal plate, its controller does not use $x^2 + y^2 = r^2$; it drives the cutting head by computing $x = a + r \cos \theta$, $y = b + r \sin \theta$ as θ increases steadily with time. The parameter θ plays the role of a clock: it tells the machine where to be at each instant. The same idea steers satellite orbits, car sat-nav curves and the rounded letters on this page.

EXERCISE 5 Practice

- Write down the centre and radius of each circle given parametrically:
 - $x = 4 \cos \theta$, $y = 4 \sin \theta$;
 - $x = -1 + 3 \cos \theta$, $y = 5 + 3 \sin \theta$.
- Convert to Cartesian form: $x = 2 + 7 \cos \theta$, $y = -3 + 7 \sin \theta$.
- Write parametric equations for the circle $x^2 + y^2 + 6x - 2y + 6 = 0$.
- A point P moves on the circle $x^2 + y^2 = 1$. Find the greatest and least values of the distance PQ , where $Q = (3, 4)$.
- $P = (\cos \theta, \sin \theta)$ moves on the unit circle and M is the midpoint of P and $(4, 0)$. Find the Cartesian equation of the locus of M .
- Find the equation of the tangent to the circle $x^2 + y^2 = 16$ at the point corresponding to $\theta = \frac{\pi}{6}$.
- A circle has parametric equations $x = 1 + 2 \cos \theta$, $y = -2 + 2 \sin \theta$. Find the Cartesian equation of the tangent at the point where $\theta = \frac{\pi}{4}$.

SOLUTION Exercise 1

- (a) Centre $(0, 0)$, radius 4. (b) Centre $(-1, 5)$, radius 3.
- $\cos \theta = \frac{x-2}{7}$, $\sin \theta = \frac{y+3}{7}$; hence $(x-2)^2 + (y+3)^2 = 49$.
- Completing the square: $(x+3)^2 + (y-1)^2 = 4$, so $x = -3 + 2 \cos \theta$, $y = 1 + 2 \sin \theta$.
- $PQ^2 = (\cos \theta - 3)^2 + (\sin \theta - 4)^2 = 26 - 6 \cos \theta - 8 \sin \theta$. Write $6 \cos \theta + 8 \sin \theta = 10 \cos(\theta - \alpha)$ with $\cos \alpha = \frac{3}{5}$, $\sin \alpha = \frac{4}{5}$. Then $PQ_{\max}^2 = 36$, $PQ_{\min}^2 = 16$, so $PQ_{\max} = 6$ and $PQ_{\min} = 4$.
- $M = \left(\frac{\cos \theta + 4}{2}, \frac{\sin \theta}{2} \right)$; eliminating θ gives $(2X-4)^2 + (2Y)^2 = 1$, i.e. $(X-2)^2 + Y^2 = \frac{1}{4}$: a circle of centre $(2, 0)$, radius $\frac{1}{2}$.
- For $x^2 + y^2 = 16$, tangent at θ is $x \cos \theta + y \sin \theta = 4$. With $\theta = \frac{\pi}{6}$: $\frac{\sqrt{3}}{2}x + \frac{1}{2}y = 4 \Rightarrow \sqrt{3}x + y = 8$.
- Circle centre $(1, -2)$, radius 2. Tangent at θ : $(x-1) \cos \theta + (y+2) \sin \theta = 2$. With $\theta = \frac{\pi}{4}$: $\frac{\sqrt{2}}{2}(x-1) + \frac{\sqrt{2}}{2}(y+2) = 2 \Rightarrow x + y + 1 = 2\sqrt{2}$.

With the parametric form in hand, we now return to Cartesian methods to study how a line meets

a circle — the subject of Section 6 — where the number of intersection points will be decided by a discriminant condition.

6 SECTION 6: INTERSECTION OF A LINE AND A CIRCLE

6.1 Introduction: Intersection of a Line and a Circle — an overview

In Section 5, we described a circle using parametric equations $x = a + r \cos \theta$, $y = b + r \sin \theta$. That description tells us where the points of the circle are. We now ask a new question: where does a given straight line meet a given circle? Think of a football field drawn on the ground at your school in Bamenda: the centre circle is painted, and the halfway line is a straight line crossing it. Where exactly does the halfway line enter and leave the circle? Answering that question precisely is the goal of this section. We shall see that the whole story is controlled by a single number: the **discriminant** of a quadratic equation.

6.2 Finding the Points of Intersection by Substitution

EXPERIMENT A first encounter

Consider the circle $x^2 + y^2 = 25$ (centre the origin, radius 5) and the line $y = x + 1$. A point lies on both curves exactly when its coordinates satisfy both equations at the same time. So we substitute $y = x + 1$ into the circle equation:

$$x^2 + (x + 1)^2 = 25 \implies 2x^2 + 2x - 24 = 0 \implies x^2 + x - 12 = 0.$$

Factorising, $(x - 3)(x + 4) = 0$, so $x = 3$ or $x = -4$. Using $y = x + 1$, the intersection points are $(3, 4)$ and $(-4, -3)$. Two points: the line is a **secant**.

This example contains the whole method. Geometrically, a line and a circle can meet in **two points** (the line is a **secant**), in **one point** (the line is a **tangent**), or in **no point at all** (the line misses the circle). Algebraically, substitution always produces a **quadratic equation** in one variable, and the three geometric situations correspond exactly to the three possible signs of its discriminant $\Delta = b^2 - 4ac$.


METHOD Intersection of a line and a circle

Given a circle $S = 0$ and a line $L = 0$:

1. From the line equation, express one variable in terms of the other (e.g. $y = mx + c$).
2. Substitute into the circle equation and simplify to a quadratic $ax^2 + bx + c = 0$.
3. Compute the discriminant $\Delta = b^2 - 4ac$.
4. Classify: $\Delta > 0$: two distinct intersection points (secant); $\Delta = 0$: one repeated point

(tangent); $\Delta < 0$: no real intersection (line misses the circle).

5. If $\Delta \geq 0$, solve the quadratic and find the corresponding values of the other variable to obtain the coordinates of the intersection points.



The circle $x^2 + y^2 = 16$ with a secant (two points), a tangent (one point) and a line that misses the circle (no point).

WORKED EXAMPLE 17 Classifying three lines

For the circle $x^2 + y^2 = 25$, determine the nature of the intersection with each line, and find the intersection points when they exist:

- (a) $y = x + 1$;
- (b) $3x + 4y = 25$;
- (c) $y = x + 8$.

SOLUTION

(a) Substituting $y = x + 1$ into $x^2 + y^2 = 25$:

$$x^2 + (x + 1)^2 = 25 \implies 2x^2 + 2x - 24 = 0 \implies x^2 + x - 12 = 0.$$

$\Delta = 1 + 48 = 49 > 0$: secant. Roots $x = 3$ and $x = -4$, giving the points $(3, 4)$ and $(-4, -3)$.

(b) From $3x + 4y = 25$, $y = \frac{25 - 3x}{4}$. Substituting:

$$x^2 + \frac{(25 - 3x)^2}{16} = 25 \implies 16x^2 + 9x^2 - 150x + 625 = 400 \implies 25x^2 - 150x + 225 = 0,$$

i.e. $x^2 - 6x + 9 = 0$, so $(x - 3)^2 = 0$. $\Delta = 0$: the line is a tangent, touching the circle at the single point $(3, 4)$.

(c) Substituting $y = x + 8$:

$$x^2 + (x + 8)^2 = 25 \implies 2x^2 + 16x + 39 = 0.$$

$\Delta = 256 - 312 = -56 < 0$: no real intersection; the line misses the circle completely.

WARNING Check before you conclude

A negative discriminant does not mean you made an algebra error — it is a genuine geometric answer: the line and the circle simply do not meet. But do check your substitution arithmetic first; a single sign slip can flip the sign of Δ .

6.3 The Distance Method: Comparing d with r

There is a second, purely geometric way to classify the intersection, which is often faster. Compute the perpendicular distance d from the centre $C(a, b)$ of the circle to the line $px + qy + s = 0$:

$$d = \frac{|pa + qb + s|}{\sqrt{p^2 + q^2}}.$$

Then compare d with the radius r :

- $d < r$: the line cuts the circle in **two points** (secant);
- $d = r$: the line **touches** the circle (tangent);
- $d > r$: the line **misses** the circle.

This is exactly the algebraic classification in disguise: one can prove that Δ (after substitution) has the same sign as $r^2 - d^2$, up to a positive factor.

EXAMPLE Distance method in action

Circle $(x - 1)^2 + (y - 2)^2 = 9$, so $C = (1, 2)$, $r = 3$. Line: $4x + 3y = 5$, i.e. $4x + 3y - 5 = 0$.

$$d = \frac{|4(1) + 3(2) - 5|}{\sqrt{16 + 9}} = \frac{|10 - 5|}{5} = 1.$$

Since $d = 1 < 3 = r$, the line is a secant: two intersection points.

6.4 Chords: Length and Midpoint

PROPERTY The perpendicular from the centre bisects the chord


Let AB be a chord of a circle with centre M , and let N be the foot of the perpendicular from M to the line AB . Then N is the midpoint of AB : the perpendicular from the centre of a circle to a chord bisects the chord. (Proof: triangles MNA and MNB are right-angled at N , share the side MN , and $MA = MB = r$; by Pythagoras $NA = NB$.)

PROPERTY Length of a chord

If a line is at perpendicular distance d from the centre of a circle of radius r (with $d < r$), then the length of the chord it cuts off is

$$AB = 2\sqrt{r^2 - d^2}.$$

This follows at once from Pythagoras in the right triangle MNA : $NA^2 = MA^2 - MN^2 = r^2 - d^2$, and $AB = 2 NA$.



The right triangle MNA : radius r , distance d from centre to chord, and half-chord $\sqrt{r^2 - d^2}$.

WORKED EXAMPLE 18 Length of a chord

The line $3x + 4y = 20$ cuts the circle $x^2 + y^2 = 25$. Find the length of the chord cut off, and the coordinates of its midpoint.

SOLUTION

The circle has centre $M = (0, 0)$ and radius $r = 5$. Distance from M to the line $3x + 4y - 20 = 0$:

$$d = \frac{|-20|}{\sqrt{9 + 16}} = \frac{20}{5} = 4.$$

Chord length:

$$AB = 2\sqrt{r^2 - d^2} = 2\sqrt{25 - 16} = 2 \times 3 = 6.$$

The midpoint N is the foot of the perpendicular from M to the line. The perpendicular through $(0, 0)$ has direction $(3, 4)$, so points on it are $(3t, 4t)$. Substituting into $3x + 4y = 20$: $9t + 16t = 20$, so $t = \frac{4}{5}$, giving $N = \left(\frac{12}{5}, \frac{16}{5}\right)$.

WARNING Do not confuse d with the chord length

The number d is the distance from the centre to the line, not the distance between the intersection points. The chord length is $2\sqrt{r^2 - d^2}$, and the half-chord is $\sqrt{r^2 - d^2}$. Forgetting the factor 2 is one of the most frequent errors at the GCE.

6.5 Family of Circles Through the Intersection of a Circle and a Line

Suppose the circle $S = 0$ and the line $L = 0$ meet at two points P and Q . We want all circles passing through P and Q without solving for P and Q explicitly.

PROPERTY The family $S + \lambda L = 0$

Let $S = x^2 + y^2 + 2gx + 2fy + c = 0$ be a circle and $L = px + qy + s = 0$ a line which intersect it. Then for every real number λ , the equation

$$S + \lambda L = 0$$

represents a circle passing through the intersection points of $S = 0$ and $L = 0$. Conversely, every circle through those two points (except S itself, in some formulations) is obtained for a suitable λ . Justification (examinable): $S + \lambda L = 0$ has the form $x^2 + y^2 + (\text{linear terms}) = 0$, so it is a circle. If (x_0, y_0) lies on both $S = 0$ and $L = 0$, then $S(x_0, y_0) + \lambda L(x_0, y_0) = 0 + \lambda \cdot 0 = 0$, so the point lies on every member of the family.

WARNING Normalise S first

Before writing $S + \lambda L = 0$, make sure the circle equation is written with coefficient 1 for both x^2 and y^2 . If you are given $2x^2 + 2y^2 + 4x - 3 = 0$, divide by 2 first. Otherwise $S + \lambda L$ will not have equal coefficients of x^2 and y^2 and will not even be a circle.

METHOD Using the family $S + \lambda L = 0$

1. Write $S = 0$ with unit coefficients of x^2 and y^2 , and $L = 0$ in the form $px + qy + s = 0$.
2. Write the family $S + \lambda L = 0$.
3. Use the extra condition (the circle passes through a given point, has a given radius, a given centre line, etc.) to form an equation in λ .
4. Solve for λ and substitute back to obtain the required circle.

WORKED EXAMPLE 19 Circle through the intersection of a circle and a line

Find the equation of the circle which passes through the points of intersection of the circle $x^2 + y^2 - 4x - 2y - 4 = 0$ and the line $x + 2y - 4 = 0$, and which also passes through the point $(2, 3)$.

SOLUTION

The family of circles through the intersection is

$$x^2 + y^2 - 4x - 2y - 4 + \lambda(x + 2y - 4) = 0.$$

The point (2, 3) lies on the required circle, so substitute $x = 2, y = 3$:

$$4 + 9 - 8 - 6 - 4 + \lambda(2 + 6 - 4) = 0 \implies -5 + 4\lambda = 0 \implies \lambda = \frac{5}{4}.$$

Substituting back and multiplying by 4 to clear fractions:

$$4x^2 + 4y^2 - 16x - 8y - 16 + 5x + 10y - 20 = 0,$$

$$4x^2 + 4y^2 - 11x + 2y - 36 = 0.$$

Check: the coefficients of x^2 and y^2 are equal, so this is indeed a circle, and one may verify that (2, 3) satisfies it: $16 + 36 - 22 + 6 - 36 = 0$. ✓

WORKED EXAMPLE 20 Determining λ from a radius condition

Find the circle through the intersection of $x^2 + y^2 = 4$ and the line $x - y = 0$ which has radius 2.

SOLUTION

Family: $x^2 + y^2 - 4 + \lambda(x - y) = 0$. Completing the square:

$$\left(x + \frac{\lambda}{2}\right)^2 + \left(y - \frac{\lambda}{2}\right)^2 = 4 + \frac{\lambda^2}{2}.$$

The radius squared is $4 + \frac{\lambda^2}{2}$. Setting this equal to $2^2 = 4$ gives $\lambda^2 = 0$, so $\lambda = 0$: the only such circle is the original circle $x^2 + y^2 = 4$ itself. (Any other member of the family has a strictly larger radius — a nice geometric remark: the common chord is a diameter of the smallest circle of the family.)

6.6 Consolidation Exercises**EXERCISE 6** Exercise 1

Find the coordinates of the points where the line $y = 2x - 3$ meets the circle $x^2 + y^2 - 4x + 2y - 5 = 0$.

EXERCISE 7 Exercise 2

Show that the line $x + y = 5$ is a tangent to the circle $x^2 + y^2 - 2x - 4y + 3 = 0$, and find the point of contact.

EXERCISE 8 Exercise 3

The line $4x - 3y = 10$ cuts the circle $x^2 + y^2 - 2x + 4y - 20 = 0$ at A and B. Find the length AB and the midpoint of AB.

EXERCISE 9 Exercise 4

Find the equation of the circle passing through the intersection of $x^2 + y^2 - 6x - 8y + 9 = 0$ and the line $x - y + 1 = 0$, and through the point (1, 2).

SOLUTION Exercise 1

Substitute $y = 2x - 3$: $x^2 + (2x - 3)^2 - 4x + 2(2x - 3) - 5 = 0$ gives $5x^2 - 12x - 2 = 0$... expanding carefully: $x^2 + 4x^2 - 12x + 9 - 4x + 4x - 6 - 5 = 5x^2 - 12x - 2 = 0$. Then $x = \frac{12 \pm \sqrt{144 + 40}}{10} = \frac{12 \pm \sqrt{184}}{10} = \frac{6 \pm \sqrt{46}}{5}$. The points are $\left(\frac{6 + \sqrt{46}}{5}, \frac{-3 + 2\sqrt{46}}{5}\right)$ and $\left(\frac{6 - \sqrt{46}}{5}, \frac{-3 - 2\sqrt{46}}{5}\right)$.

SOLUTION Exercise 2

Circle: centre (1, 2), radius $\sqrt{1 + 4 - 3} = \sqrt{2}$. Distance from (1, 2) to $x + y - 5 = 0$: $d = \frac{|1 + 2 - 5|}{\sqrt{2}} = \frac{2}{\sqrt{2}} = \sqrt{2} = r$. Hence tangent. Point of contact: foot of perpendicular from (1, 2) along direction (1, 1): $(1 + t, 2 + t)$ on the line gives $1 + t + 2 + t = 5$, $t = 1$; contact point (2, 3).

SOLUTION Exercise 3

Centre (1, -2), radius $\sqrt{1 + 4 + 20} = 5$. Distance to line $4x - 3y - 10 = 0$: $d = \frac{|4 + 6 - 10|}{5} = 0$. The line passes through the centre, so the chord is a diameter: $AB = 10$, midpoint = (1, -2).

SOLUTION Exercise 4

Family: $x^2 + y^2 - 6x - 8y + 9 + \lambda(x - y + 1) = 0$. At $(1, 2)$: $1 + 4 - 6 - 16 + 9 + \lambda(1 - 2 + 1) = -8 + 0 = -8 \neq 0$. Since $L(1, 2) = 0$ but $S(1, 2) = -8 \neq 0$, no value of λ works: the point $(1, 2)$ lies on the line L but not on the circle S , so no circle of the family passes through it. The problem as stated has no solution — always check that the given point does not lie on the line $L = 0$ outside the circle!

KEY POINTS Key points of Section 6

- Substituting the line equation into the circle equation gives a quadratic; its discriminant classifies the intersection: $\Delta > 0$ secant, $\Delta = 0$ tangent, $\Delta < 0$ no meeting.
- Equivalently, compare the distance d from the centre to the line with the radius r : $d < r$, $d = r$, $d > r$.
- The perpendicular from the centre to a chord bisects the chord; the chord length is $2\sqrt{r^2 - d^2}$.
- The family of circles through the intersection of $S = 0$ and $L = 0$ is $S + \lambda L = 0$ — with S normalised so that x^2 and y^2 both have coefficient 1.

7 SECTION 7: INTERSECTION OF TWO CIRCLES, FAMILIES OF CIRCLES AND THE RADICAL AXIS

In Section 6 we studied how a line meets a circle: we substituted the equation of the line into the equation of the circle and analysed the resulting quadratic. We now go one step further and ask: what happens when two circles meet? The beautiful surprise is that the hard work is again done by a straight line. By simply subtracting the equations of the two circles, we obtain a line which contains their common points — the **common chord**. This single idea unlocks three powerful tools of the GCE syllabus: the intersection points of two circles, the **family of circles** through those points, and the **radical axis**.

Throughout this section we write a circle in general form

$$S : x^2 + y^2 + 2gx + 2fy + c = 0,$$

and we use the shorthand $S(x, y) = 0$ or simply $S = 0$. Recall that the centre is $(-g, -f)$ and the radius is $\sqrt{g^2 + f^2 - c}$.

7.1 Points of Intersection of Two Circles and the Common Chord

Observation. Let two circles

$$S_1 : x^2 + y^2 + 2g_1x + 2f_1y + c_1 = 0, \quad S_2 : x^2 + y^2 + 2g_2x + 2f_2y + c_2 = 0$$

intersect at points A and B. If (x, y) is a common point, then both equations hold, so their difference also holds:

$$S_1 - S_2 : 2(g_1 - g_2)x + 2(f_1 - f_2)y + (c_1 - c_2) = 0.$$

The x^2 and y^2 terms cancel, leaving the equation of a straight line. Since A and B both satisfy $S_1 - S_2 = 0$, this line passes through both intersection points: it is the line of the **common chord** AB.

METHOD Intersecting two circles

To find the points of intersection of $S_1 = 0$ and $S_2 = 0$:

1. Write both equations in general form and compute $S_1 - S_2 = 0$: this is the line of the common chord.
2. Express y (or x) from the chord equation and substitute into either circle.
3. Solve the resulting quadratic for the remaining coordinate.
4. Write down both points of intersection and check them in the other circle.

WORKED EXAMPLE 21 Intersection of two circles

Find the points of intersection of the circles

$$S_1 : x^2 + y^2 - 4x - 2y - 4 = 0 \quad \text{and} \quad S_2 : x^2 + y^2 + 2x - 6y + 6 = 0.$$

SOLUTION

Step 1: common chord. Subtracting,

$$S_1 - S_2 : -6x + 4y - 10 = 0 \iff 3x - 2y + 5 = 0, \quad \text{so } y = \frac{3x+5}{2}.$$

Step 2: substitute into S_1 .

$$x^2 + \left(\frac{3x+5}{2}\right)^2 - 4x - 2\left(\frac{3x+5}{2}\right) - 4 = 0.$$


Multiplying by 4:

$$4x^2 + (9x^2 + 30x + 25) - 16x - (12x + 20) - 16 = 0 \implies 13x^2 + 2x - 11 = 0.$$

Step 3: solve. $13x^2 + 2x - 11 = 0$ factorises as $(13x - 11)(x + 1) = 0$, giving $x = -1$ or $x = \frac{11}{13}$.

For $x = -1$: $y = \frac{-3+5}{2} = 1$. For $x = \frac{11}{13}$: $y = \frac{1}{2}\left(\frac{33}{13} + 5\right) = \frac{49}{13}$.

Conclusion. The circles meet at $(-1, 1)$ and $\left(\frac{11}{13}, \frac{49}{13}\right)$. Both points satisfy $S_2 = 0$, as required.



Two intersecting circles: the common chord AB is perpendicular to the line of centres C_1C_2 .

7.2 The Family of Circles Through the Intersection of Two Circles

Observation. Suppose we want the circle through the two intersection points of $S_1 = 0$ and $S_2 = 0$ which also passes through a third given point. Solving from scratch (three points, three unknowns g, f, c) is long. There is a far more elegant route: build a whole family of circles through A and B with one parameter, then choose the parameter to fit the third point.

PROPERTY Family of circles through two intersection points — proof examinable

Let $S_1 = 0$ and $S_2 = 0$ be two intersecting circles. Then for every real $\lambda \neq -1$, the equation

$$S_1 + \lambda S_2 = 0$$

represents a circle passing through all the common points of S_1 and S_2 . Conversely, every circle through those common points (other than S_2 itself) has this form for some λ .

Proof. First, $S_1 + \lambda S_2 = 0$ is

$$(1 + \lambda)x^2 + (1 + \lambda)y^2 + 2(g_1 + \lambda g_2)x + 2(f_1 + \lambda f_2)y + (c_1 + \lambda c_2) = 0.$$

Since $\lambda \neq -1$, we may divide by $1 + \lambda$: the coefficients of x^2 and y^2 become 1 and there is no xy term, so this is indeed a circle. Secondly, if P is a common point of the two circles, then $S_1(P) = 0$ and $S_2(P) = 0$, hence $S_1(P) + \lambda S_2(P) = 0$: the point P lies on every member of the family. ■

WARNING The case $\lambda = -1$ and the missing circle

When $\lambda = -1$, the quadratic terms vanish and $S_1 - S_2 = 0$ is the common chord (a line, not a circle). Also, the family $S_1 + \lambda S_2 = 0$ never produces S_2 itself (that would require $1 + \lambda = 0$). If a problem's answer turns out to be S_2 , check it separately.

METHOD Circle through the intersection of two circles and a given point

1. Write the family $S_1 + \lambda S_2 = 0$.
2. Substitute the coordinates of the given point: this gives a linear equation in λ .
3. Solve for λ (reject $\lambda = -1$), substitute back and simplify.

WORKED EXAMPLE 22 Circle through two intersection points and a given point

Find the equation of the circle which passes through the points of intersection of

$$S_1 : x^2 + y^2 - 4x - 2y - 4 = 0, \quad S_2 : x^2 + y^2 + 2x - 6y + 6 = 0$$

and through the point $(1, 2)$.

SOLUTION

The family is $S_1 + \lambda S_2 = 0$:

$$(1 + \lambda)x^2 + (1 + \lambda)y^2 + (-4 + 2\lambda)x + (-2 - 6\lambda)y + (-4 + 6\lambda) = 0.$$


Substituting $(1, 2)$: note $S_1(1, 2) = 1 + 4 - 4 - 4 - 4 = -7$ and $S_2(1, 2) = 1 + 4 + 2 - 12 + 6 = 1$. Hence

$$-7 + \lambda(1) = 0 \implies \lambda = 7.$$

Substituting back:

$$8x^2 + 8y^2 + 10x - 44y + 38 = 0 \iff 4x^2 + 4y^2 + 5x - 22y + 19 = 0.$$

Check: at $(1, 2)$: $4 + 16 + 5 - 44 + 19 = 0$. ✓



Members of the family $S_1 + \lambda S_2 = 0$: every circle passes through the same two points A and B.

7.3 The Radical Axis of Two Circles

We now give the line $S_1 - S_2 = 0$ a name and a meaning that remain valid even when the circles do not intersect.

Recall from Section 4 that the length of the tangent from an external point $P(x_1, y_1)$ to the circle $S = 0$ is $\sqrt{S(x_1, y_1)}$, provided S is written with unit coefficients of x^2 and y^2 .

DEFINITION Radical axis

The **radical axis** of two circles $S_1 = 0$ and $S_2 = 0$ is the locus of points P from which the tangent lengths to the two circles are equal:

$$\sqrt{S_1(P)} = \sqrt{S_2(P)} \iff S_1(P) = S_2(P) \iff S_1 - S_2 = 0.$$

Its equation is therefore

$$2(g_1 - g_2)x + 2(f_1 - f_2)y + (c_1 - c_2) = 0.$$

PROPERTY The radical axis is perpendicular to the line of centres — proof examinable

For two non-concentric circles, the radical axis is perpendicular to the line joining their centres.


Proof. The centres are $C_1(-g_1, -f_1)$ and $C_2(-g_2, -f_2)$, so a direction vector of the line of centres is

$$\vec{C_1C_2} = \begin{pmatrix} -(g_2 - g_1) \\ -(f_2 - f_1) \end{pmatrix}.$$

The radical axis $2(g_1 - g_2)x + 2(f_1 - f_2)y + (c_1 - c_2) = 0$ has normal vector $\begin{pmatrix} g_1 - g_2 \\ f_1 - f_2 \end{pmatrix}$, which is parallel to $\vec{C_1C_2}$. A line whose normal is parallel to C_1C_2 is perpendicular to C_1C_2 . ■

KEY POINTS The radical axis in the three configurations

- Intersecting circles: the radical axis is the line of the **common chord**.
- Touching circles: the radical axis is the **common tangent** at the point of contact.
- Non-intersecting circles: the radical axis is the line of points with equal tangent lengths; it lies between the circles, perpendicular to the line of centres.



Non-intersecting circles: from any point P of the radical axis, the tangent segments PT_1 and PT_2 have equal length.

WORKED EXAMPLE 23 Radical axis and equal tangent lengths

Find the radical axis of the circles

$$S_1 : x^2 + y^2 - 2x - 4y - 4 = 0, \quad S_2 : x^2 + y^2 + 4x - 2y + 1 = 0,$$

and verify that the point $P(1, 4)$ on it has equal tangent lengths to both circles.

SOLUTION

Radical axis:

$$S_1 - S_2 : -6x - 2y - 5 = 0 \iff 6x + 2y + 5 = 0.$$

Check that P lies on it: $6(1) + 2(4) + 5 = 19 \neq 0$ — so in fact $P(1, 4)$ is not on the axis. Take

instead the point where the axis meets $x = 1$: $2y = -11$, $y = -\frac{11}{2}$; call $Q\left(1, -\frac{11}{2}\right)$. Then

$$S_1(Q) = 1 + \frac{121}{4} - 2 + 22 - 4 = \frac{189}{4}, \quad S_2(Q) = 1 + \frac{121}{4} + 4 + 11 + 1 = \frac{189}{4}.$$

Hence the tangent lengths are equal: $\ell_1 = \ell_2 = \sqrt{\frac{189}{4}} = \frac{3\sqrt{21}}{2}$. This illustrates the defining property of the radical axis.

DID YOU KNOW? The radical centre of three circles

Given three circles with non-collinear centres, the three radical axes taken in pairs are concurrent: they meet at a single point called the **radical centre**. Indeed, if P satisfies $S_1 = S_2$ and $S_2 = S_3$, then automatically $S_1 = S_3$. The radical centre is the unique point with equal tangent lengths to all three circles — a favourite one-line justification in GCE questions.

7.4 Locus Problems Involving Two Circles

The radical axis answers a whole class of locus questions.

WORKED EXAMPLE 24 Locus of points with equal tangent lengths

A point P moves so that the lengths of the tangents from P to the circles

$$x^2 + y^2 - 6x - 8y = 0 \quad \text{and} \quad x^2 + y^2 + 2x - 4y - 3 = 0$$

are equal. Find the locus of P .

SOLUTION

Let $P(x, y)$. Equal tangent lengths means $S_1(P) = S_2(P)$:

$$x^2 + y^2 - 6x - 8y = x^2 + y^2 + 2x - 4y - 3 \implies 8x + 4y - 3 = 0.$$

The locus is the straight line $8x + 4y - 3 = 0$ — precisely the radical axis of the two circles.

A second classical type asks for a constant ratio or sum of squared distances.

WORKED EXAMPLE 25 Constant sum of squared distances

Find the locus of a point P such that the sum of the squares of its distances from $A(2, 0)$ and $B(-2, 0)$ is 20.

SOLUTION

Let $P(x, y)$. Then

$$PA^2 + PB^2 = (x - 2)^2 + y^2 + (x + 2)^2 + y^2 = 2x^2 + 2y^2 + 8.$$

Setting this equal to 20: $2x^2 + 2y^2 = 12$, i.e.

$$x^2 + y^2 = 6,$$

a circle centred at the origin (the midpoint of AB) with radius $\sqrt{6}$.

WARNING Common mistakes in this section

- Subtracting the equations when the coefficients of x^2 and y^2 are not both 1: first divide each circle through by the coefficient of x^2 , otherwise $S_1 - S_2$ is not the radical axis.
- Forgetting that $S_1 + \lambda S_2 = 0$ with $\lambda = -1$ gives a line, not a circle: always state $\lambda \neq -1$.
- Claiming the radical axis passes through the intersection points when the circles do not intersect: it is still well defined, but it is no longer a chord.
- Sign errors when computing $S_1 - S_2$: subtract every term, including the constants.

EXERCISE 10

The circles $S_1 : x^2 + y^2 - 2x - 6y + 6 = 0$ and $S_2 : x^2 + y^2 + 4x + 2y - 4 = 0$ are given.

1. Find the equation of their radical axis.
2. Find the coordinates of their points of intersection.
3. Find the equation of the circle through these intersection points and the point $(0, 2)$.

SOLUTION Exercise 1

1. $S_1 - S_2 : -6x - 8y + 10 = 0$, i.e. $3x + 4y - 5 = 0$.

2. From the axis, $y = \frac{5 - 3x}{4}$. Substituting into S_1 and multiplying by 16:

$$16x^2 + (25 - 30x + 9x^2) - 32x - 24(5 - 3x) + 96 = 0$$

$$\Rightarrow 25x^2 + 10x + 1 = 0 \Rightarrow (5x + 1)^2 = 0.$$

Thus $x = -\frac{1}{5}$ (double root). Then $y = \frac{5-3(-1/5)}{4} = \frac{5+3/5}{4} = \frac{28/5}{4} = \frac{7}{5}$. The circles touch externally at $(-\frac{1}{5}, \frac{7}{5})$. (The discriminant is zero, confirming they touch.)

3. The family of circles through the common point (the point of contact) is $S_1 + \lambda S_2 = 0$, $\lambda \neq -1$. Substituting $(0, 2)$: $S_1(0, 2) = 4 - 12 + 6 = -2$, $S_2(0, 2) = 4 + 4 - 4 = 4$, so $-2 + 4\lambda = 0$, $\lambda = \frac{1}{2}$. Substituting back:

$$S_1 + \frac{1}{2}S_2 = 0 \implies 2S_1 + S_2 = 0 \implies 3x^2 + 3y^2 - 10y + 8 = 0.$$

This circle passes through the point of contact $(-\frac{1}{5}, \frac{7}{5})$ and through $(0, 2)$.

EXERCISE 11

Find the radical centre of the three circles

$$x^2 + y^2 - 4x - 2y - 4 = 0, \quad x^2 + y^2 + 2x - 6y + 6 = 0, \quad x^2 + y^2 - 8x + 4y + 4 = 0.$$

SOLUTION Exercise 2

Radical axis of the first pair: $-6x + 4y - 10 = 0$, i.e. $3x - 2y + 5 = 0$. Radical axis of circles 1 and 3: $4x - 6y - 8 = 0$, i.e. $2x - 3y - 4 = 0$. Solving: from the first, $y = \frac{3x+5}{2}$; substituting: $2x - \frac{9x+15}{2} - 4 = 0$, so $4x - 9x - 15 - 8 = 0$, $x = -\frac{23}{5}$, $y = \frac{1}{2}(-\frac{69}{5} + 5) = -\frac{22}{5}$. The radical centre is $(-\frac{23}{5}, -\frac{22}{5})$; one may verify it also lies on the radical axis of circles 2 and 3.

KEY POINTS Key points of Section 7

- Subtracting two circle equations gives a line: the common chord (intersecting case), the common tangent (touching case), or the radical axis in general.
- The family of circles through the intersections of $S_1 = 0$ and $S_2 = 0$ is $S_1 + \lambda S_2 = 0$, $\lambda \neq -1$; find λ by substituting a given point.
- The radical axis $S_1 - S_2 = 0$ is the locus of points with equal tangent lengths and is perpendicular to the line of centres.
- The three radical axes of three circles concur at the radical centre.

8 SECTION 8: CONDITIONS FOR TWO CIRCLES TO TOUCH

In Section 7 we studied what happens when two circles intersect: we found their common points, the family of circles through those points, and the radical axis $S_1 - S_2 = 0$. But between “two circles that cut each other in two points” and “two circles that do not meet at all” there is a remarkable boundary case: the two circles meet in exactly one point. We then say that the circles **touch** each other, or that they are **tangent** to one another. This section gives simple algebraic tests, in terms of the distance between the centres and the radii, to decide in which configuration two given circles lie, and how to find the point of contact and the common tangent when they touch. This is a classic GCE Advanced Level question.

8.1 The Five Relative Positions of Two Circles

Consider two circles C_1 and C_2 , with centres C_1, C_2 and radii r_1, r_2 . Let

$$d = |C_1C_2|$$

be the distance between the centres. Imagine holding C_1 fixed and sliding C_2 towards it from far away. At first the circles are separate; then they just touch; then they overlap and cut in two points; then, if C_2 is small enough, it slips inside C_1 , touches it from the inside, and finally lies entirely inside without touching. The whole geometry is controlled by comparing d with the two numbers


$$r_1 + r_2 \quad \text{and} \quad |r_1 - r_2|.$$

PROPERTY Relative positions of two circles

Let $C_1(C_1, r_1)$ and $C_2(C_2, r_2)$ be two circles and $d = |C_1C_2|$. Then:

1. $d > r_1 + r_2$: the circles are **separate** (external to each other), no common point;
2. $d = r_1 + r_2$: the circles **touch externally**, exactly one common point;
3. $|r_1 - r_2| < d < r_1 + r_2$: the circles **intersect** in two distinct points;
4. $d = |r_1 - r_2|$ (with $d \neq 0$): the circles **touch internally**, exactly one common point;
5. $d < |r_1 - r_2|$: one circle lies **inside the other** without touching, no common point.

(The degenerate case $d = 0$ with $r_1 = r_2$ gives two identical circles; if $d = 0$ and $r_1 \neq r_2$ the circles are concentric and have no common points.)



The five relative positions of two circles, classified by comparing d with $r_1 + r_2$ and $|r_1 - r_2|$.

8.2 Conditions for Two Circles to Touch

PROPERTY Touching conditions

Two circles with centres C_1, C_2 , radii r_1, r_2 , and $d = |C_1C_2|$:

- touch externally if and only if $d = r_1 + r_2$;
- touch internally if and only if $d = |r_1 - r_2|$ (with $d \neq 0$).

In both cases the two circles have exactly one common point, called the **point of contact**. (The proof of the “if” direction, by exhibiting the common point on the line of centres, is instructive and sketched below; the statement itself must be known for the examination.)

Why these conditions are natural. Suppose the circles touch externally at T . Since T lies on both circles, $|C_1T| = r_1$ and $|C_2T| = r_2$, and T lies between C_1 and C_2 on the line of centres, so

$$d = |C_1C_2| = |C_1T| + |TC_2| = r_1 + r_2.$$

If they touch internally (say C_2 inside C_1), then C_2 and T are on the same side of C_1 and C_1, C_2, T are collinear with T furthest from C_1 , so

$$d = |C_1C_2| = |C_1T| - |C_2T| = r_1 - r_2 = |r_1 - r_2|.$$

Conversely, if $d = r_1 + r_2$, the point T on segment $[C_1C_2]$ with $|C_1T| = r_1$ satisfies $|TC_2| = d - r_1 = r_2$, so T lies on both circles; any other common point P would give a triangle C_1PC_2 with $|C_1P| + |PC_2| = r_1 + r_2 = d = |C_1C_2|$, forcing P onto the segment $[C_1C_2]$ by the triangle inequality, hence $P = T$. The internal case is similar.

PROPERTY The point of contact lies on the line of centres

If two circles touch, their point of contact T lies on the line joining their centres. Moreover T divides C_1C_2 in the ratio of the radii:

- external contact: T divides $[C_1C_2]$ internally in the ratio $r_1 : r_2$, so

$$T = \frac{r_2 C_1 + r_1 C_2}{r_1 + r_2} \quad \text{i.e.} \quad T = C_1 + \frac{r_1}{r_1 + r_2} \vec{C_1C_2};$$


- internal contact: T divides the line C_1C_2 externally in the ratio $r_1 : r_2$; the vector formula

$$T = C_1 + \frac{r_1}{r_1 - r_2} \vec{C_1C_2}$$

works in all cases (if $r_1 < r_2$ the coefficient is negative, placing T on the ray opposite to $\vec{C_1C_2}$).

METHOD Finding the point of contact

1. Compute the centres C_1, C_2 and radii r_1, r_2 ; compute $d = |C_1C_2|$.
2. Test $d = r_1 + r_2$ (external) and $d = |r_1 - r_2|$ (internal).
3. Use the vector formula:
 - External: $T = C_1 + \frac{r_1}{r_1 + r_2} \vec{C_1C_2}$.
 - Internal: $T = C_1 + \frac{r_1}{r_1 - r_2} \vec{C_1C_2}$ (this automatically handles whether $r_1 > r_2$ or $r_1 < r_2$).



Two circles touching externally at T : C_1, T, C_2 are collinear, $d = r_1 + r_2$, and the common tangent at T is perpendicular to the line of centres.

8.3 The Common Tangent at the Point of Contact

When two circles touch at T , the tangent to C_1 at T and the tangent to C_2 at T are both perpendicular to the line of centres C_1C_2 at the same point T : they are therefore the same line. Two touching circles always share a **common tangent** at their point of contact.

Recall from Section 7 that the radical axis of two circles $S_1 = 0$ and $S_2 = 0$ is the line $S_1 - S_2 = 0$. When the circles intersect, this is the line through the two intersection points. When the circles touch, the two intersection points merge into one, and the radical axis becomes the common tangent at that point.

METHOD Common tangent of two touching circles

Write the equations of the two circles in general form with the coefficients of x^2 and y^2

equal to 1 in both (if necessary, divide the whole equation by the coefficient of x^2):

$$S_1 = x^2 + y^2 + 2g_1x + 2f_1y + c_1 = 0, \quad S_2 = x^2 + y^2 + 2g_2x + 2f_2y + c_2 = 0.$$

The common tangent at the point of contact is

$$S_1 - S_2 = 0,$$

$$\text{i.e. } 2(g_1 - g_2)x + 2(f_1 - f_2)y + (c_1 - c_2) = 0.$$

WORKED EXAMPLE 26 Two circles touching externally

Show that the circles

$$x^2 + y^2 - 4x - 6y + 9 = 0 \quad \text{and} \quad x^2 + y^2 + 2x + 2y - 7 = 0$$

touch each other. Find the point of contact and the equation of the common tangent at that point.

SOLUTION

Centres and radii. For C_1 : centre $C_1 = (2, 3)$, radius $r_1 = \sqrt{4 + 9 - 9} = 2$. For C_2 : centre $C_2 = (-1, -1)$, radius $r_2 = \sqrt{1 + 1 + 7} = 3$.

Distance between centres.

$$d = |C_1C_2| = \sqrt{(2 - (-1))^2 + (3 - (-1))^2} = \sqrt{9 + 16} = \sqrt{25} = 5.$$

Since $d = 5 = 2 + 3 = r_1 + r_2$, the circles touch externally. ■

Point of contact. T lies on $[C_1C_2]$ with $|C_1T| = r_1 = 2$ and $d = 5$:

$$T = C_1 + \frac{r_1}{d} \vec{C_1C_2} = (2, 3) + \frac{2}{5}(-3, -4) = \left(2 - \frac{6}{5}, 3 - \frac{8}{5}\right) = \left(\frac{4}{5}, \frac{7}{5}\right).$$

$$\text{Check: } |C_2T| = \sqrt{\left(\frac{4}{5} + 1\right)^2 + \left(\frac{7}{5} + 1\right)^2} = \sqrt{\frac{81}{25} + \frac{144}{25}} = \sqrt{9} = 3 = r_2. \quad \checkmark$$

Common tangent. Using $S_1 - S_2 = 0$:

$$(-4x - 6y + 9) - (2x + 2y - 7) = 0 \implies -6x - 8y + 16 = 0 \implies 3x + 4y - 8 = 0.$$

$$\text{Check that } T \text{ lies on it: } 3 \cdot \frac{4}{5} + 4 \cdot \frac{7}{5} - 8 = \frac{12+28-40}{5} = 0. \quad \checkmark$$

8.4 Determining Unknown Constants so that Two Circles Touch

A typical GCE question gives two circles, one of which contains an unknown constant k , and asks for the value(s) of k for which the circles touch. The strategy is always the same: express d, r_1, r_2 in terms of k and solve both equations $d = r_1 + r_2$ and $d = |r_1 - r_2|$.

WORKED EXAMPLE 27 Finding k so that two circles touch

Find the values of k for which the circle

$$x^2 + y^2 - 6x - 8y + k = 0$$

touches the circle $x^2 + y^2 = 4$.

SOLUTION

C_2 : centre $C_2 = (0, 0)$, radius $r_2 = 2$. C_1 : centre $C_1 = (3, 4)$, radius $r_1 = \sqrt{9 + 16 - k} = \sqrt{25 - k}$ (we need $k < 25$ for a real circle).

Distance between centres: $d = \sqrt{9 + 16} = 5$.

External contact: $d = r_1 + r_2$ gives

$$5 = \sqrt{25 - k} + 2 \implies \sqrt{25 - k} = 3 \implies 25 - k = 9 \implies k = 16.$$

Internal contact: $d = |r_1 - r_2|$ gives

$$5 = |\sqrt{25 - k} - 2|.$$

Either $\sqrt{25 - k} - 2 = 5$, i.e. $\sqrt{25 - k} = 7$, giving $25 - k = 49$, i.e. $k = -24$; or $\sqrt{25 - k} - 2 = -5$, i.e. $\sqrt{25 - k} = -3$, impossible.

Hence the circles touch when $k = 16$ (external contact) or $k = -24$ (internal contact, with C_2 inside C_1 since then $r_1 = 7 > r_2 = 2$).

WARNING “Touch” may be internal or external

Unless the question says externally or internally, the word **touch** covers both cases. Always test $d = r_1 + r_2$ and $d = |r_1 - r_2|$. In the example above, a candidate who only tries external contact finds $k = 16$ and loses the second answer $k = -24$ — typically half the marks.

WARNING Internal contact is not “one inside the other”

Do not confuse $d = |r_1 - r_2|$ (the circles touch internally, one common point) with $d < |r_1 - r_2|$ (one circle lies strictly inside the other, no common point). Similarly $d > r_1 + r_2$ means separate, not touching. The equality sign is essential: touching is always an equality between d and one of $r_1 + r_2$, $|r_1 - r_2|$.

8.5 Number of Common Tangents (Useful for Multiple-Choice Questions)

A quick way to visualise each configuration is to count the **common tangents** — lines tangent to both circles at once:

Configuration	Condition	Common tangents
Separate	$d > r_1 + r_2$	4
Touch externally	$d = r_1 + r_2$	3
Intersecting	$ r_1 - r_2 < d < r_1 + r_2$	2
Touch internally	$d = r_1 - r_2 $	1
One inside the other	$d < r_1 - r_2 $	0

Notice the pattern: each time the circles pass from one configuration to the next, one pair of common tangents merges into a single tangent and then disappears. The tangent that survives the “touching” cases is precisely the common tangent at the point of contact, $S_1 - S_2 = 0$.

EXERCISE 12 Practice

- Determine the relative position of the circles $x^2 + y^2 = 9$ and $x^2 + y^2 - 8x + 6y + 9 = 0$. If they touch, find the point of contact and the common tangent.
- Find the values of a for which the circle $x^2 + y^2 - 2ax = 0$ touches the circle $x^2 + y^2 - 4y = 0$.
- Two circles have centres $(1, 2)$ and $(4, 6)$ and touch externally. If the radius of the first is 2, find the radius of the second and the equation of their common tangent.

SOLUTION Exercise 1

$C_1: C_1 = (0, 0), r_1 = 3$. $C_2: C_2 = (4, -3), r_2 = \sqrt{16 + 9 - 9} = 4$. Then $d = \sqrt{16 + 9} = 5$. Since $|r_1 - r_2| = 1 < 5 < 7 = r_1 + r_2$, the circles intersect in two points; they do not touch.

SOLUTION Exercise 2

C_1 : centre $(a, 0)$, radius $|a|$. C_2 : centre $(0, 2)$, radius 2. Then $d = \sqrt{a^2 + 4}$. External contact: $\sqrt{a^2 + 4} = |a| + 2 \Rightarrow a^2 + 4 = a^2 + 4|a| + 4 \Rightarrow |a| = 0 \Rightarrow a = 0$, but then C_1 is the point circle $x^2 + y^2 = 0$ (radius 0) — rejected for a proper circle. Internal contact: $\sqrt{a^2 + 4} = ||a| - 2| \Rightarrow a^2 + 4 = a^2 - 4|a| + 4 \Rightarrow |a| = 0$, again rejected. For $a \neq 0$, we have $(|a| - 2)^2 < a^2 + 4 < (|a| + 2)^2$, i.e. $|r_1 - r_2| < d < r_1 + r_2$, so the circles always intersect in two distinct points. Hence there is no value of a for which the two proper circles touch.

SOLUTION Exercise 3

$d = \sqrt{(4-1)^2 + (6-2)^2} = \sqrt{9+16} = 5$. External contact: $r_2 = d - r_1 = 5 - 2 = 3$. Point of contact: $T = (1, 2) + \frac{2}{5}(3, 4) = (\frac{11}{5}, \frac{18}{5})$. The common tangent is perpendicular to C_1C_2 (gradient $\frac{4}{3}$) at T , so its gradient is $-\frac{3}{4}$: $y - \frac{18}{5} = -\frac{3}{4}(x - \frac{11}{5})$, i.e. $3x + 4y - 21 = 0$.

KEY POINTS Key points of Section 8

- Compare $d = |C_1C_2|$ with $r_1 + r_2$ and $|r_1 - r_2|$ to classify the position of two circles.
- Touch externally $\iff d = r_1 + r_2$; touch internally $\iff d = |r_1 - r_2|$.
- The point of contact lies on the line of centres and divides it in the ratio of the radii (internally for external contact, externally for internal contact).
- The common tangent at the point of contact is $S_1 - S_2 = 0$ (with x^2, y^2 coefficients equal to 1).
- “Touch” without qualification means: test both conditions.
- Common tangents: 4, 3, 2, 1, 0 as the circles move from separate to nested.

9 SECTION 9: LOCUS PROBLEMS AND SYNTHESIS FOR THE GCE EXAM

In Section 8 we studied the conditions for two circles to touch, and throughout this chapter we have repeatedly answered the question: given an equation, what curve does it represent? In this final section we reverse the question: given a geometric condition, what is the equation — and the nature — of the curve traced out? This is the idea of a **locus**, one of the most elegant and most frequently examined topics at the GCE Advanced Level. We close the chapter with synthesis problems in full GCE style, together with advice on exam technique.

9.1 The Concept of a Locus

Imagine a goat tethered by a rope of length 5 m to a peg in a field near Bafoussam. As the goat moves while keeping the rope taut, the tip of its path traces a circle of radius 5 m about the peg. The circle is the locus of the goat: the set of all positions satisfying the condition “distance from the peg equals 5 m”.


DEFINITION Locus

A **locus** is the set of all points satisfying a given geometric condition. Finding a locus means determining the equation (and the geometric description) of the curve formed by all such points.

The word locus is Latin for “place”: the locus is literally “the place” where the moving point is allowed to be. The power of coordinate geometry is that it converts a geometric condition into an algebraic equation.

METHOD The Four-Step Locus Strategy

1. Name the moving point. Let $P(x, y)$ be a general point of the locus.
2. Translate the condition. Write the given geometric condition as an algebraic equation involving x and y (distance formula, gradient condition, etc.).
3. Simplify. Square where necessary, expand, and reduce to a recognisable form.
4. Identify and describe. State clearly what curve the equation represents (line, circle with centre and radius, etc.), and check for excluded points.



The four-step method for every locus problem.

9.2 Elementary Loci: Fixed Distance from a Point or a Line

The two simplest loci already contain an important contrast.

Fixed distance from a fixed point. The locus of P with $|OP| = r$ for a fixed point O is, by definition, a **circle** of centre O and radius r . With $O = (a, b)$ the equation is $(x - a)^2 + (y - b)^2 = r^2$.

Fixed distance from a fixed line. The locus of P at a fixed distance d from a fixed line ℓ is a pair of parallel lines, one on each side of ℓ , each at distance d from ℓ . For instance, the locus of points at distance 2 from the line $y = 1$ is the pair $y = 3$ and $y = -1$.

WARNING A circle, not a single line

Do not confuse the two situations: fixed distance from a point gives one circle; fixed dis-

tance from a line gives two parallel lines (unless the problem restricts the point to one side). Forgetting one of the two lines is a classic loss of a mark.

9.3 Equidistant Points and the Circle of Apollonius

Equal distances: the perpendicular bisector. Let $A(x_1, y_1)$ and $B(x_2, y_2)$ be fixed. The condition $PA = PB$ gives

$$(x - x_1)^2 + (y - y_1)^2 = (x - x_2)^2 + (y - y_2)^2,$$

and after squaring out, the quadratic terms cancel, leaving a linear equation: the locus is the **perpendicular bisector** of AB .

Constant ratio of distances. Now replace equality by a ratio.

PROPERTY Circle of Apollonius

Let A and B be two fixed points and let $k > 0$, $k \neq 1$. The locus of points P such that

$$\frac{PA}{PB} = k$$

is a **circle**, called the circle of Apollonius. (For $k = 1$ the locus degenerates to the perpendicular bisector of AB .) The derivation below is instructive; the result itself is a GCE-useful extra.

Derivation. Take $A = (-c, 0)$ and $B = (c, 0)$ (we may always choose axes this way). Then $PA^2 = k^2 PB^2$ gives

$$(x + c)^2 + y^2 = k^2[(x - c)^2 + y^2].$$

Expanding both sides,

$$x^2 + 2cx + c^2 + y^2 = k^2x^2 - 2k^2cx + k^2c^2 + k^2y^2,$$


and collecting like terms:

$$(1 - k^2)x^2 + (1 - k^2)y^2 + 2c(1 + k^2)x + c^2(1 - k^2) = 0.$$

Since $k \neq 1$ we may divide by $1 - k^2$:

$$x^2 + y^2 + \frac{2c(1 + k^2)}{1 - k^2}x + c^2 = 0,$$

which has the form $x^2 + y^2 + 2gx + 2fy + c' = 0$ with $f = 0$: a circle whose centre lies on the line AB . ■



Circle of Apollonius for $A = (-2, 0)$, $B = (2, 0)$ with ratio $k = 2$: centre $(\frac{10}{3}, 0)$, radius $\frac{8}{3}$; every point P of the circle satisfies $PA = 2PB$.

WORKED EXAMPLE 28 An Apollonius locus

$A = (0, 0)$ and $B = (6, 0)$. Find the locus of P such that $PA = 2PB$.

SOLUTION

Let $P(x, y)$. The condition $PA^2 = 4PB^2$ gives

$$x^2 + y^2 = 4[(x - 6)^2 + y^2].$$

Expanding: $x^2 + y^2 = 4x^2 - 48x + 144 + 4y^2$, so

$$3x^2 + 3y^2 - 48x + 144 = 0 \implies x^2 + y^2 - 16x + 48 = 0.$$

Completing the square: $(x - 8)^2 + y^2 = 16$. Description: the locus is a circle of centre $(8, 0)$ and radius 4.

9.4 Loci Defined by $PA^2 + PB^2$ or $PA^2 - PB^2$ Constant

These two conditions appear regularly at the GCE and reward algebraic fluency.

EXAMPLE Sum of squares constant

Let $A = (-a, 0)$, $B = (a, 0)$ and suppose $PA^2 + PB^2 = 2c$ (constant). Then

$$(x + a)^2 + y^2 + (x - a)^2 + y^2 = 2c \implies 2x^2 + 2y^2 + 2a^2 = 2c,$$

so $x^2 + y^2 = c - a^2$: a circle centred at the midpoint of AB , with radius $\sqrt{c - a^2}$ (real only if

$$c \geq a^2).$$

EXAMPLE Difference of squares constant

With the same points, suppose $PA^2 - PB^2 = k$ (constant). Then

$$[(x+a)^2 + y^2] - [(x-a)^2 + y^2] = 4ax = k,$$


so $x = \frac{k}{4a}$: a straight line perpendicular to AB . Note the contrast: the sum gives a circle, the difference gives a line.

9.5 Locus of the Midpoint of a Chord Through a Fixed Point

A beautiful result links loci to the circle geometry of this chapter. Let C be a circle with centre O , and let F be a fixed point inside it. Draw any chord of C passing through F , and let M be its midpoint. Since the line from the centre to the midpoint of a chord is perpendicular to the chord, $\hat{OMF} = 90^\circ$. By the converse of the angle-in-a-semicircle theorem, M lies on the circle with diameter OF .

PROPERTY Locus of midpoints of chords through a fixed point

The locus of the midpoints of the chords of a circle C (centre O) which pass through a fixed interior point F is the circle with diameter OF (restricted to the interior of C).



Chords of C through the fixed point F (dashed) have midpoints M lying on the circle with diameter OF .

WORKED EXAMPLE 29 Midpoint locus by algebra

Find the locus of the midpoints of chords of the circle $x^2 + y^2 = 25$ that pass through $F(3, 0)$.

SOLUTION

Let $M(h, k)$ be the midpoint of a chord. The chord with midpoint (h, k) of the circle $x^2 + y^2 = 25$ has equation $T = S_1$:

$$hx + ky = h^2 + k^2.$$

Since the chord passes through $F(3, 0)$: $3h = h^2 + k^2$. Replacing (h, k) by (x, y) , the locus is

$$x^2 + y^2 - 3x = 0 \iff \left(x - \frac{3}{2}\right)^2 + y^2 = \frac{9}{4},$$

the circle with diameter OF : centre $\left(\frac{3}{2}, 0\right)$, radius $\frac{3}{2}$. (Only the part inside the given circle is attained.)

9.6 Loci via Parametric Elimination

Sometimes the moving point is given parametrically: its coordinates depend on a parameter t (or θ). The locus is found by eliminating the parameter — exactly the technique of Lesson 10.

WORKED EXAMPLE 30 Eliminating the parameter

A point moves so that its position is $P = (3 + 2 \cos \theta, -1 + 2 \sin \theta)$. Find its locus.

SOLUTION

We have $\cos \theta = \frac{x-3}{2}$ and $\sin \theta = \frac{y+1}{2}$. Using $\cos^2 \theta + \sin^2 \theta = 1$:

$$(x-3)^2 + (y+1)^2 = 4.$$

The locus is the circle of centre $(3, -1)$ and radius 2.

METHOD Parametric locus problems

1. Express $\cos \theta$ and $\sin \theta$ (or the parameter t) in terms of x and y .
2. Eliminate using an identity ($\cos^2 + \sin^2 = 1$, or substitution for t).
3. Simplify and describe the resulting curve completely.

9.7 Synthesis: GCE-Style Problems and Exam Technique

GCE questions rarely test one idea in isolation. A typical Part B question might give a circle through three points, ask for a tangent, then impose a touching condition or a locus. The following exercise mixes several lessons of this chapter.

EXERCISE 13 GCE synthesis

The circle C has equation $x^2 + y^2 - 4x - 6y + 9 = 0$.

1. Find the centre and radius of C .
2. Show that the line $y = 1$ is a tangent to C and find the point of contact.
3. A point P moves so that its distance from the centre of C equals its distance from the line $x = 0$. Show that the locus of P is a parabola and find its equation.
4. Find the equation of the circle which passes through the points of intersection of C and the circle $x^2 + y^2 = 9$ and also through the point $(0, 0)$.

SOLUTION Exercise 1

1. Complete the squares: $x^2 - 4x + y^2 - 6y + 9 = 0$ gives $(x - 2)^2 - 4 + (y - 3)^2 - 9 + 9 = 0$, i.e. $(x - 2)^2 + (y - 3)^2 = 4$. Centre $C(2, 3)$, radius $r = 2$.
2. The distance from $C(2, 3)$ to the line $y = 1$ is $|3 - 1| = 2 = r$. Hence $y = 1$ is a tangent to C . The point of contact is the foot of the perpendicular from C to the line, namely $(2, 1)$.
3. Condition: $\sqrt{(x - 2)^2 + (y - 3)^2} = |x|$. Squaring: $(x - 2)^2 + (y - 3)^2 = x^2$, i.e. $x^2 - 4x + 4 + (y - 3)^2 = x^2$, so $(y - 3)^2 = 4x - 4 = 4(x - 1)$: a parabola with vertex $(1, 3)$ and axis $y = 3$.
4. Family through the intersections: $x^2 + y^2 - 4x - 6y + 9 + \lambda(x^2 + y^2 - 9) = 0$. Imposing $(0, 0)$: $9 - 9\lambda = 0 \Rightarrow \lambda = 1$. Hence $2x^2 + 2y^2 - 4x - 6y = 0$, i.e. $x^2 + y^2 - 2x - 3y = 0$.

WARNING State the full description — and check restrictions

Two final warnings for the examination:

- Never leave the answer as a raw equation. Write: “the locus is a circle of centre $(8, 0)$ and radius 4” (or a line, a pair of lines, a parabola, with its key data).
- Check for restrictions and degenerate cases: the ratio $k = 1$ in Apollonius gives a line, not a circle; $PA^2 + PB^2 = 2c$ gives a real circle only if $c \geq a^2$; a midpoint locus may be limited to the interior of the given circle. Examiners award a specific mark for these observations.

KEY POINTS Key points of Section 9

- A **locus** is the set of all points satisfying a geometric condition; find it by the four-step method: name $P(x, y)$, translate, simplify, identify.
- Fixed distance from a point $\Rightarrow$ circle; from a line $\Rightarrow$ two parallel lines; equidistant from two points $\Rightarrow$ perpendicular bisector.
- $PA/PB = k \neq 1 \Rightarrow$ circle of Apollonius; $PA^2 + PB^2$ constant $\Rightarrow$ circle; $PA^2 - PB^2$ constant $\Rightarrow$ line.
- Midpoints of chords of a circle through a fixed interior point F lie on the circle of diameter OF .
- Parametric loci: eliminate the parameter using $\cos^2 \theta + \sin^2 \theta = 1$ or substitution.
- In the exam: describe the curve completely, check restrictions, and manage your time — a locus part is usually short once the method is launched.

DID YOU KNOW? Apollonius of Perga

The circle of Apollonius is named after the Greek geometer Apollonius of Perga (3rd century BC), whose treatise *Conics* also gave us the names ellipse, parabola and hyperbola. Nearly two thousand years later, Descartes' coordinate geometry — the tool you have used throughout this chapter — turned his purely geometric results into simple algebraic manipulations. The same circle of Apollonius appears today in physics (field patterns between two charges) and in telecommunications (locating a receiver from signal-strength ratios).

Exam technique and time management.

In the GCE Advanced Level paper, circle geometry questions are usually structured: early parts earn quick marks (centre, radius, tangent verification), while the final part is often a locus or a family-of-circles demand. Budget your time accordingly: secure the routine marks first, and do not spend more than a few minutes stuck on algebra — write down the condition $PA^2 = k^2 PB^2$ (or the family equation $S_1 + \lambda S_2 = 0$) immediately, since method marks are awarded for correct set-up even if the simplification is incomplete. Finally, re-read the question: if it asks for “the locus”, a bare equation without a geometric description will cost you the last mark.

This completes our study of Circle Geometry. You now command every tool of the chapter: equations of circles in all their forms, tangents and their lengths, parametric representations, intersections of lines and circles, families of circles and the radical axis, touching conditions, and loci. Practise the mixed exercises of this section under timed conditions, and you will approach the GCE with confidence.

10 INTEGRATION ACTIVITY

10.1 Aims of the activity

By the end of this group activity, you should be able to:

- determine the equation of a circle passing through three given points, using both the general equation $x^2 + y^2 + 2gx + 2fy + c = 0$ and the **perpendicular bisector** method;
- interpret the centre of that circle as the point **equidistant** from three given locations, and its radius as a coverage distance;
- use the **perpendicular distance from a point to a line** to decide whether a line cuts, touches or misses a circle, and compute a **chord length**;
- form the **radical axis** of two circles by eliminating the quadratic terms, and give it a physical meaning;
- identify the **locus** of a point equidistant from two fixed points and compare it with the radical axis;
- communicate a complete mathematical solution, supported by an accurate sketch on grid paper.

KEY POINTS Skills mobilised

Circle through three points • perpendicular bisectors • line–circle intersection • chord length $2\sqrt{r^2 - d^2}$ • radical axis $S_1 - S_2 = 0$ • locus (perpendicular bisector).

10.2 Situation

The North–West Regional delegation of telecommunications wishes to improve mobile coverage between three villages situated on the highland plateau. On a simplified map drawn on squared paper, the planner chooses axes along two perpendicular dirt tracks and uses the kilometre as unit. The three villages are modelled by the points

$$A(1, 2), \quad B(7, 4), \quad C(3, 9) \quad (\text{coordinates in km}).$$

A first relay antenna, called **Antenna 1**, must be placed so that its signal reaches the three villages equally well: its position must therefore be equidistant from A , B and C , and its coverage zone is the circle through the three villages. Recall the key geometric fact that makes this possible:

PROPERTY Circle through three points


Through any three **non-collinear** points there passes exactly one circle. Its centre is the

point equidistant from the three points, found as the intersection of the perpendicular bisectors of any two of the segments joining them.

A straight tarred road is modelled by the line of equation $y = x - 1$. For safety and engineering reasons, the delegation wants to know whether this road enters the coverage zone, and if so, over what length.

Later, a second relay, **Antenna 2**, is installed at the point $P(10, 8)$ with a coverage radius of 4 km. The engineers need to know along which line the signals of the two antennas balance (a point where a phone could switch from one network to the other), and a technician walking so that he stays at equal distance from the two antennas wants to describe his path.

Your group is the consultancy team. You must produce a full report: all equations, all decisions justified by calculation, and an accurate sketch on grid paper.



Map of the situation: villages A, B, C, coverage circle of Antenna 1 (blue), Antenna 2 (orange), the road (green) and the radical axis (dashed).

10.3 Tasks

Part A — Positioning Antenna 1.

1. Write the general equation of a circle $x^2 + y^2 + 2gx + 2fy + c = 0$ and explain why substituting A, B and C gives a system of three linear equations in g, f, c .
2. Solve the system and write the equation of the coverage circle.
3. Deduce the coordinates of the centre O_1 (the antenna position) and the coverage radius, to 2 decimal places.
4. Check: verify your centre by finding the perpendicular bisectors of $[AB]$ and $[AC]$ and inter-

secting them.

Part B — The road and the coverage zone.

5. Compute the perpendicular distance d from O_1 to the road $y = x - 1$.
6. Comparing d with the radius, decide whether the road cuts, touches or misses the coverage circle.
7. If the road cuts the circle, calculate the length of road lying inside the coverage zone.

Part C — Two antennas.

8. Write the equation of the coverage circle of Antenna 2 (centre $(10, 8)$, radius 4).
9. Find the radical axis of the two circles and interpret it.
10. Find the locus of the technician who moves so that his distances to the two antennas are equal. Compare this locus with the radical axis: what do you notice, and why are they not the same line?

Part D — Presentation. Produce the final report with a TikZ-style sketch on grid paper showing the villages, both coverage circles, the road, the radical axis and the technician's path.

10.4 Model answer

Part A. The general equation $x^2 + y^2 + 2gx + 2fy + c = 0$ contains three unknown constants g, f, c . Each point (x_0, y_0) lying on the circle must satisfy the equation, and since $x_0^2 + y_0^2$ is a known number, each substitution produces one linear equation in g, f, c . Three non-collinear points therefore give three independent linear equations, which determine g, f, c uniquely. Substituting:

$$\begin{array}{lll} A(1, 2) : & 1 + 4 + 2g + 4f + c = 0 & \Rightarrow 2g + 4f + c = -5, \\ B(7, 4) : & 49 + 16 + 14g + 8f + c = 0 & \Rightarrow 14g + 8f + c = -65, \\ C(3, 9) : & 9 + 81 + 6g + 18f + c = 0 & \Rightarrow 6g + 18f + c = -90. \end{array}$$

Subtracting the first equation from the other two eliminates c :

$$12g + 4f = -60 \Rightarrow 3g + f = -15, \quad 4g + 14f = -85.$$

From the first of these, $f = -15 - 3g$; substituting into the second:

$$4g + 14(-15 - 3g) = -85 \Rightarrow -38g - 210 = -85 \Rightarrow -38g = 125,$$

giving

$$\begin{aligned} g &= -\frac{125}{38} \approx -3.29, & f &= -15 - 3g = -\frac{195}{38} \approx -5.13, \\ c &= -5 - 2g - 4f = -5 + \frac{250}{38} + \frac{780}{38} = \frac{420}{19} \approx 22.11. \end{aligned}$$

The coverage circle is

$$x^2 + y^2 - \frac{125}{19}x - \frac{195}{19}y + \frac{420}{19} = 0,$$

with centre $(-g, -f)$ and radius $\sqrt{g^2 + f^2 - c}$:

$$O_1\left(\frac{125}{38}, \frac{195}{38}\right) \approx (3.29, 5.13), \quad r = \sqrt{\frac{53650}{1444} - \frac{420}{19}} = \sqrt{\frac{10865}{722}} \approx 3.88 \text{ km}.$$

Check by bisectors. Midpoint of $[AB]$: $(4, 3)$; gradient of AB is $\frac{4-2}{7-1} = \frac{1}{3}$, so the perpendicular bisector has gradient -3 and equation $y - 3 = -3(x - 4)$, i.e. $y = -3x + 15$. Midpoint of $[AC]$: $(2, \frac{11}{2})$; gradient of AC is $\frac{9-2}{3-1} = \frac{7}{2}$, so the bisector has gradient $-\frac{2}{7}$ and equation $y - \frac{11}{2} = -\frac{2}{7}(x - 2)$. Solving the two bisector equations simultaneously:

$$-3x + 15 = \frac{11}{2} - \frac{2}{7}x + \frac{4}{7} \Rightarrow -\frac{19}{7}x = -\frac{125}{14} \Rightarrow x = \frac{125}{38}, \quad y = \frac{195}{38}.$$

Same centre as before. ✓

Part B. Write the road in the standard form $x - y - 1 = 0$. The perpendicular distance from $O_1(\frac{125}{38}, \frac{195}{38})$ to the line is

$$d = \frac{\left| \frac{125}{38} - \frac{195}{38} - 1 \right|}{\sqrt{1^2 + (-1)^2}} = \frac{\left| -\frac{108}{38} \right|}{\sqrt{2}} = \frac{54}{19\sqrt{2}} \approx 2.01 \text{ km.}$$

Since $d \approx 2.01 < r \approx 3.88$, the road **cuts** the coverage circle in two points: a portion of the road lies inside the zone. That portion is a chord, and by Pythagoras' theorem applied to the right triangle formed by the centre, the foot of the perpendicular and an endpoint of the chord, half the chord has length $\sqrt{r^2 - d^2}$. Hence

$$L = 2\sqrt{r^2 - d^2} = 2\sqrt{\frac{10865}{722} - \frac{2916}{722}} = 2\sqrt{\frac{7949}{722}} \approx 6.64 \text{ km.}$$

About 6.64 km of the road lies inside the coverage zone.

Part C. Antenna 2: $(x - 10)^2 + (y - 8)^2 = 16$, which expands to

$$S_2 : x^2 + y^2 - 20x - 16y + 148 = 0.$$

Writing $S_1 : x^2 + y^2 - \frac{125}{19}x - \frac{195}{19}y + \frac{420}{19} = 0$, the radical axis is $S_1 - S_2 = 0$ (the quadratic terms cancel, leaving a straight line):

$$\left(-\frac{125}{19} + 20\right)x + \left(-\frac{195}{19} + 16\right)y + \left(\frac{420}{19} - 148\right) = 0,$$

and since $-\frac{125}{19} + 20 = \frac{255}{19}$, $-\frac{195}{19} + 16 = \frac{109}{19}$ and $\frac{420}{19} - 148 = -\frac{2392}{19}$, multiplying through by 19 gives

$$255x + 109y - 2392 = 0.$$

Interpretation: along this line the powers of a point with respect to both circles are equal — equivalently, tangent segments drawn from a point of this line to the two circles have equal length. Physically, the signals of the two antennas balance along this line; a phone crossing it switches network.

For the technician, equidistance from O_1 and O_2 means his locus is the **perpendicular bisector** of $[O_1O_2]$. The midpoint is $(\frac{505}{76}, \frac{499}{76})$; the gradient of O_1O_2 is $\frac{8 - \frac{195}{38}}{10 - \frac{125}{38}} = \frac{109}{255}$, so the locus has gradient $-\frac{255}{109}$ and equation

$$255x + 109y = 255 \cdot \frac{505}{76} + 109 \cdot \frac{499}{76} = \frac{91583}{38} \approx 2410.1.$$

Comparison: the locus has the same coefficients of x and y as the radical axis, so the two lines are **parallel** — but distinct, since $2392 \neq 2410.1$. The radical axis balances the powers of a point with respect to the two circles (equal tangent lengths), whereas the technician's path balances the distances to the centres; the two notions coincide only when the circles have equal radii.

WARNING Pitfalls to avoid in your report

- Forgetting to halve the coefficients: the centre of $x^2 + y^2 + 2gx + 2fy + c = 0$ is $(-g, -f)$, not $(-2g, -2f)$.
- Using $d > r$ to conclude “the line cuts the circle” — it is the opposite: the line cuts the circle when $d < r$, touches it when $d = r$ and misses it when $d > r$.
- Confusing the radical axis (equal powers) with the perpendicular bisector of the centres (equal distances).
- Rounding too early: keep exact fractions throughout the working and round only the final answers.

EXERCISE 14 Extension — for the fastest groups

A third antenna of radius 3 km is proposed at $(0, 11)$. Show that the three radical axes (taken two circles at a time) are concurrent, and find their common point (the **radical centre**). What could this point mean for the network engineer?

SOLUTION Extension exercise

The third circle is $S_3 : x^2 + (y - 11)^2 = 9$, i.e. $S_3 : x^2 + y^2 - 22y + 112 = 0$. The radical axis of S_1 and S_3 is $S_1 - S_3 = 0$:

$$-\frac{125}{19}x + \left(-\frac{195}{19} + 22\right)y + \left(\frac{420}{19} - 112\right) = 0 \Rightarrow -125x + 223y - 1708 = 0,$$

after multiplying by 19 (using $-\frac{195}{19} + 22 = \frac{223}{19}$ and $\frac{420}{19} - 112 = -\frac{1708}{19}$). The radical axis of S_1 and S_2 was $255x + 109y - 2392 = 0$. Solving these two linear equations simultaneously: from the second, $x = \frac{223y - 1708}{125}$; substituting into the first,

$$255 \cdot \frac{223y - 1708}{125} + 109y = 2392 \Rightarrow 14098y = 146908 \Rightarrow y = \frac{73454}{7049} \approx 10.42,$$

and then $x \approx 4.93$. The third radical axis, $S_2 - S_3 = 0$, is $-20x + 6y + 36 = 0$, i.e. $10x - 3y - 18 = 0$; substituting $x \approx 4.93$, $y \approx 10.42$ gives $10(4.93) - 3(10.42) - 18 \approx 0$, so the third axis passes through the same point: the three radical axes are concurrent at the radical centre $(4.93, 10.42)$ (to 2 d.p.). This concurrence is general: any point on two of the radical axes has equal powers with respect to all three circles, so it lies on the third axis too. For the engineer, the radical centre is the unique point where the three networks balance — a natural location for a shared switching station.

11 METHODS AND WORKED EXAMPLES


12 EQUATIONS OF A CIRCLE — CENTRE AT THE ORIGIN AND GENERAL CENTRE

12.1 Introduction and Geometric Definition

A **circle** is the locus of a point $P(x, y)$ which moves in a plane such that its distance from a fixed point $C(a, b)$ (the **centre**) is always equal to a constant r (the **radius**). Using the distance formula:

$$PC = \sqrt{(x - a)^2 + (y - b)^2} = r \quad \Rightarrow \quad (x - a)^2 + (y - b)^2 = r^2.$$

This is the **standard form** (or centre-radius form) of the equation of a circle.



The standard circle with centre $C(a, b)$ and radius r .

DEFINITION Circle Centred at the Origin

If the centre is the origin $O(0, 0)$, the equation reduces to:

$$x^2 + y^2 = r^2.$$

PROPERTY Centre and Radius from Standard Form

For the circle $(x - a)^2 + (y - b)^2 = r^2$:

- Centre: $C(a, b)$.
- Radius: $r = \sqrt{r^2}$ (must have $r^2 > 0$ for a real circle).

EXAMPLE Identifying Centre and Radius

Find the centre and radius of: (i) $x^2 + y^2 = 16$; (ii) $(x + 2)^2 + (y - 3)^2 = 10$.

Solution. (i) Centre $(0, 0)$, radius $r = 4$. (ii) Rewrite as $(x - (-2))^2 + (y - 3)^2 = (\sqrt{10})^2$. Centre $(-2, 3)$, radius $\sqrt{10}$.

12.2 Equation Given Centre and a Point on the Circle

If the centre $C(a, b)$ and a point $P(x_1, y_1)$ on the circumference are known, the radius is $r = CP = \sqrt{(x_1 - a)^2 + (y_1 - b)^2}$. Substitute into standard form.

WORKED EXAMPLE 31 Centre and Point on Circumference

Find the equation of the circle with centre $C(2, -3)$ passing through $P(5, 1)$.

SOLUTION

Solution. $r^2 = (5 - 2)^2 + (1 - (-3))^2 = 3^2 + 4^2 = 25$. Equation: $(x - 2)^2 + (y + 3)^2 = 25$ or $x^2 + y^2 - 4x + 6y - 12 = 0$.

13 THE GENERAL EQUATION AND CIRCLES ON A GIVEN DIAMETER**13.1 General Equation of a Circle**

Expanding the standard form $(x - a)^2 + (y - b)^2 = r^2$ gives:

$$x^2 + y^2 - 2ax - 2by + (a^2 + b^2 - r^2) = 0.$$

Letting $-2a = 2g$, $-2b = 2f$, and $a^2 + b^2 - r^2 = c$, we obtain the **general equation**:

$$x^2 + y^2 + 2gx + 2fy + c = 0 \quad (1)$$

with centre $(-g, -f)$ and radius $r = \sqrt{g^2 + f^2 - c}$, provided $g^2 + f^2 - c > 0$.

PROPERTY Conditions for a Real Circle

The equation $x^2 + y^2 + 2gx + 2fy + c = 0$ represents:

- A real circle if $g^2 + f^2 - c > 0$ (radius $r = \sqrt{g^2 + f^2 - c}$).

- A point circle (radius zero) if $g^2 + f^2 - c = 0$.
- No real locus (imaginary circle) if $g^2 + f^2 - c < 0$.

METHOD Completing the Square: General $\rightarrow$ Standard

To find the centre and radius from $x^2 + y^2 + 2gx + 2fy + c = 0$:

1. Group x and y terms: $(x^2 + 2gx) + (y^2 + 2fy) = -c$.
2. Complete squares: Add g^2 and f^2 to both sides: $(x^2 + 2gx + g^2) + (y^2 + 2fy + f^2) = g^2 + f^2 - c$.
3. Write as: $(x + g)^2 + (y + f)^2 = g^2 + f^2 - c$.
4. Centre: $(-g, -f)$; Radius: $\sqrt{g^2 + f^2 - c}$.

WORKED EXAMPLE 32 General to Standard Form

Find the centre and radius of $x^2 + y^2 - 6x + 4y - 12 = 0$.

SOLUTION

Solution. $(x^2 - 6x) + (y^2 + 4y) = 12$. $(x - 3)^2 - 9 + (y + 2)^2 - 4 = 12$. $(x - 3)^2 + (y + 2)^2 = 25$. Centre $(3, -2)$, Radius 5.

13.2 Circle on a Given Diameter

If $A(x_1, y_1)$ and $B(x_2, y_2)$ are endpoints of a diameter, the centre is the midpoint and the radius is half the length AB .

PROPERTY Equation of Circle on Diameter AB

Centre: $\left(\frac{x_1+x_2}{2}, \frac{y_1+y_2}{2}\right)$. Radius: $\frac{1}{2}\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$. Equation: $(x - x_1)(x - x_2) + (y - y_1)(y - y_2) = 0$ (diameter form).

WORKED EXAMPLE 33 Circle on Diameter

Find the equation of the circle with $A(2, -1)$ and $B(4, 3)$ as diameter endpoints.

SOLUTION

Solution. Centre $C\left(\frac{2+4}{2}, \frac{-1+3}{2}\right) = (3, 1)$. $AB = \sqrt{(4-2)^2 + (3+1)^2} = \sqrt{20} = 2\sqrt{5} \implies r = \sqrt{5}$.
 Standard form: $(x-3)^2 + (y-1)^2 = 5$. General form: $x^2 + y^2 - 6x - 2y + 5 = 0$. Check using
 diameter form: $(x-2)(x-4) + (y+1)(y-3) = 0 \implies x^2 - 6x + 8 + y^2 - 2y - 3 = 0 \implies x^2 + y^2 - 6x - 2y + 5 = 0$. ✓

WARNING Common Mistake: Sign of Centre

In $x^2 + y^2 + 2gx + 2fy + c = 0$, the centre is $(-g, -f)$, not (g, f) . Example: $x^2 + y^2 - 4x + 6y = 0 \implies g = -2, f = 3 \implies \text{Centre } (2, -3)$.

14 CIRCLE THROUGH THREE POINTS AND CIRCLES UNDER GEOMETRIC CONDITIONS

14.1 Circle Through Three Non-Collinear Points

Three non-collinear points uniquely determine a circle. We use the general form $x^2 + y^2 + 2gx + 2fy + c = 0$ and substitute the coordinates to get three linear equations in g, f, c .

METHOD Circle Through Three Points P_1, P_2, P_3

1. Let the circle be $x^2 + y^2 + 2gx + 2fy + c = 0$.
2. Substitute each point (x_i, y_i) : $2gx_i + 2fy_i + c = -(x_i^2 + y_i^2)$ for $i = 1, 2, 3$.
3. Solve the 3×3 system for g, f, c (elimination or determinants).
4. Write the equation. Centre $= (-g, -f)$, Radius $= \sqrt{g^2 + f^2 - c}$.
5. Alternative (Perpendicular Bisectors): Find equations of perpendicular bisectors of chords P_1P_2 and P_2P_3 . Their intersection is the centre. Radius = distance to any point. Use as a check.

WORKED EXAMPLE 34 Circle Through Three Points

A circle passes through $P(1, 2)$, $Q(3, -4)$, $R(5, 2)$. Find its equation, centre and radius.

SOLUTION

Solution. Let circle be $x^2 + y^2 + 2gx + 2fy + c = 0$. $P(1, 2) : 1 + 4 + 2g + 4f + c = 0 \Rightarrow 2g + 4f + c = -5$
 $(1) Q(3, -4) : 9 + 16 + 6g - 8f + c = 0 \Rightarrow 6g - 8f + c = -25$ $(2) R(5, 2) : 25 + 4 + 10g + 4f + c = 0 \Rightarrow 10g + 4f + c = -29$ (3)
 $(3) - (1) : 8g = -24 \Rightarrow g = -3$. Sub $g = -3$ in $(1) : -6 + 4f + c = -5 \Rightarrow 4f + c = 1$ (4) Sub $g = -3$ in $(2) : -18 - 8f + c = -25 \Rightarrow -8f + c = -7$ (5)
 $(4) - (5) : 12f = 8 \Rightarrow f = \frac{2}{3}$. From $(4) : 4(\frac{2}{3}) + c = 1 \Rightarrow c = -\frac{5}{3}$.
Equation: $x^2 + y^2 - 6x + \frac{4}{3}y - \frac{5}{3} = 0$ or $3x^2 + 3y^2 - 18x + 4y - 5 = 0$. Centre: $(3, -\frac{2}{3})$. Radius: $\sqrt{9 + \frac{4}{9} + \frac{5}{3}} = \sqrt{\frac{100}{9}} = \frac{10}{3}$.

14.2 Circles Touching the Coordinate Axes

A circle touches the x -axis if $|y_{\text{centre}}| = r$. It touches the y -axis if $|x_{\text{centre}}| = r$. If it touches both axes, centre is $(\pm r, \pm r)$ (four quadrants). Equation: $(x \mp r)^2 + (y \mp r)^2 = r^2$.

WORKED EXAMPLE 35 Circle Touching Both Axes

Find the equation of the circle of radius 3 touching both axes in the second quadrant.

SOLUTION

Solution. Second quadrant: centre $(-3, 3)$. Equation: $(x + 3)^2 + (y - 3)^2 = 9 \Rightarrow x^2 + y^2 + 6x - 6y + 9 = 0$.

14.3 Circle Through Two Points with Centre on a Given Line

The centre lies on the perpendicular bisector of the chord joining the two points and on the given line. Intersection gives the centre.

WORKED EXAMPLE 36 Two Points, Centre on Line

Find the circle through $A(1, 2)$, $B(3, 4)$ with centre on $y = 2x - 1$.

SOLUTION

Solution. Midpoint of AB : $M(2, 3)$. Slope $AB = 1 \Rightarrow$ perp. bisector slope $= -1$. Eq. perp. bisector: $y - 3 = -(x - 2) \Rightarrow y = -x + 5$. Intersect with $y = 2x - 1$: $-x + 5 = 2x - 1 \Rightarrow 3x = 6 \Rightarrow x = 2, y = 3$. Centre $C(2, 3)$. Radius $r = CA = \sqrt{(2 - 1)^2 + (3 - 2)^2} = \sqrt{2}$. Equation: $(x - 2)^2 + (y - 3)^2 = 2$ or $x^2 + y^2 - 4x - 6y + 11 = 0$.

14.4 Circle Through Two Points and Touching a Given Line

Let the line be $L : ax + by + c = 0$. Centre (h, k) satisfies:

1. Equidistant from the two points (lies on perp. bisector).
2. Distance to line = r = distance to either point.

$$\frac{|ah+bk+c|}{\sqrt{a^2+b^2}} = \sqrt{(h-x_1)^2 + (k-y_1)^2}. \text{ Solve for } (h, k) \text{ (usually two solutions).}$$

WORKED EXAMPLE 37 Two Points, Touching a Line

Find circles through $A(1, 1)$, $B(2, 2)$ touching the line $x + y = 0$.

SOLUTION

Solution. Midpoint $M(1.5, 1.5)$. Slope $AB = 1 \Rightarrow$ perp. bisector: $y - 1.5 = -1(x - 1.5) \Rightarrow x + y = 3$. Centre (h, k) lies on $h + k = 3 \Rightarrow k = 3 - h$. Distance to line $x + y = 0$: $r = \frac{|h+k|}{\sqrt{2}} = \frac{3}{\sqrt{2}}$. Distance to $A(1, 1)$: $r^2 = (h - 1)^2 + (k - 1)^2 = (h - 1)^2 + (2 - h)^2 = 2h^2 - 6h + 5$. Equate r^2 : $\frac{9}{2} = 2h^2 - 6h + 5 \Rightarrow 4h^2 - 12h + 1 = 0$. $h = \frac{12 \pm \sqrt{144 - 16}}{8} = \frac{3 \pm \sqrt{8}}{2} = \frac{3 \pm 2\sqrt{2}}{2}$. Two circles exist. Centres $\left(\frac{3+2\sqrt{2}}{2}, \frac{3-2\sqrt{2}}{2}\right)$ and $\left(\frac{3-2\sqrt{2}}{2}, \frac{3+2\sqrt{2}}{2}\right)$, radius $\frac{3}{\sqrt{2}}$.

14.5 Circle Touching a Given Line (at a Given Point or Not)

If the point of contact $T(x_1, y_1)$ on line L is given: Centre lies on normal to L through T . Radius = CT . If only the line is given (and other conditions like radius or centre on line), use distance formula $r = \frac{|ah+bk+c|}{\sqrt{a^2+b^2}}$.


15 TANGENTS TO A CIRCLE

15.1 Equation of Tangent at a Point on the Circle

For the circle $S : x^2 + y^2 + 2gx + 2fy + c = 0$ and point $P(x_1, y_1)$ on the circle ($S_{11} = 0$), the tangent is:

$$T = 0 \quad \text{where} \quad T : xx_1 + yy_1 + g(x + x_1) + f(y + y_1) + c = 0.$$

For circle centred at origin $x^2 + y^2 = r^2$: $xx_1 + yy_1 = r^2$.



Tangent at P is perpendicular to radius CP .

PROPERTY Tangent–Radius Perpendicularity

The tangent at any point P on a circle is perpendicular to the radius CP . Gradient of radius $m_{CP} = \frac{y_1 - b}{x_1 - a} \implies$ Gradient of tangent $m_T = -\frac{1}{m_{CP}} = -\frac{x_1 - a}{y_1 - b}$ (if $y_1 \neq b$).

15.2 Condition for a Line to be Tangent

Line $y = mx + c$ (or $y = mx + d$ to avoid confusion with circle constant c) and circle $x^2 + y^2 = r^2$ (centre origin). Substitute: $x^2 + (mx + d)^2 = r^2 \implies (1 + m^2)x^2 + 2mdx + (d^2 - r^2) = 0$. Tangency $\iff$ Discriminant $\Delta = 0$:

$$(2md)^2 - 4(1 + m^2)(d^2 - r^2) = 0 \implies d^2 = r^2(1 + m^2).$$

For circle $(x - a)^2 + (y - b)^2 = r^2$, substitute line equation, set $\Delta = 0$.

METHOD Tangent Condition and Equation

1. Given line $y = mx + d$ and circle centre (a, b) radius r : Substitute y , get quadratic in x . Set $\Delta = 0$ to find d (or m).
2. Given point $P(x_1, y_1)$ on circle: Use $T = 0$ formula or gradient method.
3. Given external point $P(x_1, y_1)$: Find tangents by solving for m in $y - y_1 = m(x - x_1)$ with $\Delta = 0$, or use chord of contact $T = 0$ (where T uses external point).

WORKED EXAMPLE 38 Tangent Condition: Line $y = 2x + k$ tangent to $x^2 + y^2 - 4x + 2y - 4 = 0$

Find k and points of contact.

SOLUTION

Solution. Circle: $(x - 2)^2 + (y + 1)^2 = 9$. Centre $C(2, -1)$, $r = 3$. Substitute $y = 2x + k$: $(x - 2)^2 + (2x + k + 1)^2 = 9$. $5x^2 + 4kx + (k^2 + 2k - 4) = 0$. $\Delta = (4k)^2 - 4(5)(k^2 + 2k - 4) = 16k^2 - 20k^2 - 40k + 80 = -4k^2 - 40k + 80 = 0$. $k^2 + 10k - 20 = 0 \implies k = -5 \pm 3\sqrt{5}$. Points of contact (double root $x = -\frac{4k}{10} = -\frac{2k}{5}$): For $k_1 = -5 + 3\sqrt{5}$: $x_1 = 2 - \frac{6\sqrt{5}}{5}$, $y_1 = -1 + \frac{3\sqrt{5}}{5}$. For $k_2 = -5 - 3\sqrt{5}$: $x_2 = 2 + \frac{6\sqrt{5}}{5}$, $y_2 = -1 - \frac{3\sqrt{5}}{5}$.

15.3 Length of Tangent from an External Point

For point $P(x_1, y_1)$ outside circle $S : x^2 + y^2 + 2gx + 2fy + c = 0$, the length of tangent PT is:

$$PT = \sqrt{x_1^2 + y_1^2 + 2gx_1 + 2fy_1 + c} = \sqrt{S_{11}}.$$

This is the square root of the **Power of the Point** S_{11} . If $S_{11} > 0$: Point outside, two real tangents. If $S_{11} = 0$: Point on circle, tangent length 0. If $S_{11} < 0$: Point inside, no real tangents.

WORKED EXAMPLE 39 Length of Tangent

Find the length of the tangent from $P(3, 4)$ to the circle $x^2 + y^2 - 2x - 4y - 4 = 0$.


SOLUTION

Solution. $S_{11} = 3^2 + 4^2 - 2(3) - 4(4) - 4 = 9 + 16 - 6 - 16 - 4 = -1 < 0$. Point is inside the circle. No real tangent exists.

16 PARAMETRIC EQUATIONS OF A CIRCLE**16.1 Standard Parametrisation**

A circle with centre (a, b) and radius r can be described by the parameter θ (angle with positive x -axis):

$$x = a + r \cos \theta, \quad y = b + r \sin \theta, \quad \theta \in [0, 2\pi).$$



Parametric coordinates: θ is the angle CP makes with the horizontal.

16.2 Conversion: Parametric $\rightarrow$ Cartesian

From $x = a + r \cos \theta$, $y = b + r \sin \theta$: $\cos \theta = \frac{x-a}{r}$, $\sin \theta = \frac{y-b}{r}$. Using $\cos^2 \theta + \sin^2 \theta = 1$:

$$\left(\frac{x-a}{r}\right)^2 + \left(\frac{y-b}{r}\right)^2 = 1 \implies (x-a)^2 + (y-b)^2 = r^2.$$

WORKED EXAMPLE 40 Parametric to Cartesian and Tangent

Circle: $x = 1 + 3 \cos \theta$, $y = -2 + 3 \sin \theta$. (i) Cartesian equation. (ii) Tangent at $\theta = \frac{\pi}{3}$.

SOLUTION

Solution. (i) $\cos \theta = \frac{x-1}{3}$, $\sin \theta = \frac{y+2}{3}$. $\left(\frac{x-1}{3}\right)^2 + \left(\frac{y+2}{3}\right)^2 = 1 \implies (x-1)^2 + (y+2)^2 = 9$. Centre $(1, -2)$, $r = 3$.

(ii) At $\theta = \frac{\pi}{3}$: $\cos \frac{\pi}{3} = \frac{1}{2}$, $\sin \frac{\pi}{3} = \frac{\sqrt{3}}{2}$. Point P : $x_1 = 1 + 3 \cdot \frac{1}{2} = \frac{5}{2}$, $y_1 = -2 + 3 \cdot \frac{\sqrt{3}}{2} = \frac{3\sqrt{3}-4}{2}$. Method 1 (Gradient): $m_{CP} = \frac{y_1+2}{x_1-1} = \frac{3\sqrt{3}/2}{1/2} = \sqrt{3}$. $m_T = -\frac{1}{\sqrt{3}} = -\frac{\sqrt{3}}{3}$. Tangent: $y - \frac{3\sqrt{3}-4}{2} = -\frac{\sqrt{3}}{3} \left(x - \frac{5}{2}\right)$. Multiply by 6: $6y - 9\sqrt{3} + 12 = -2\sqrt{3}(2x - 5) = -4\sqrt{3}x + 10\sqrt{3}$. $4\sqrt{3}x + 6y + 12 - 19\sqrt{3} = 0$ (or simplified $x + \sqrt{3}y = 7 - 2\sqrt{3}$).

Method 2 (Formula $T = 0$): Circle: $x^2 + y^2 - 2x + 4y - 4 = 0$ ($g = -1, f = 2, c = -4$). $xx_1 + yy_1 - 1(x + x_1) + 2(y + y_1) - 4 = 0$. $x \cdot \frac{5}{2} + y \cdot \frac{3\sqrt{3}-4}{2} - x - \frac{5}{2} + 2y + 3\sqrt{3} - 4 - 4 = 0$. Multiply by 2: $5x + (3\sqrt{3}-4)y - 2x - 5 + 4y + 6\sqrt{3} - 16 = 0$. $3x + 3\sqrt{3}y + 6\sqrt{3} - 21 = 0 \implies x + \sqrt{3}y = 7 - 2\sqrt{3}$. ✓

17 INTERSECTION OF A LINE AND A CIRCLE

17.1 Algebraic Method and Discriminant Analysis

Substitute line equation into circle equation $\rightarrow$ quadratic in x (or y). Let quadratic be $Ax^2 + Bx + C =$

0, discriminant $\Delta = B^2 - 4AC$.

PROPERTY Line–Circle Intersection

- $\Delta > 0$: Line cuts circle in two distinct points (chord).
- $\Delta = 0$: Line touches circle (tangent), one repeated root (point of contact).
- $\Delta < 0$: Line does not meet the circle.

METHOD Chord Length and Midpoint

If line $y = mx + d$ cuts circle giving roots x_1, x_2 :

1. Chord length $AB = \sqrt{1+m^2} |x_1 - x_2| = \frac{\sqrt{1+m^2}\sqrt{\Delta}}{|A|}$.
2. Midpoint M : $x_M = \frac{x_1+x_2}{2} = -\frac{B}{2A}$, $y_M = mx_M + d$.
3. Geometric property: The perpendicular from centre to chord bisects the chord. Distance from centre to line $d = \frac{|Ax_c + By_c + C|}{\sqrt{A^2+B^2}}$. Half-chord length $= \sqrt{r^2 - d^2}$.

WORKED EXAMPLE 41 Chord Length and Midpoint

Line $y = x - 2$ cuts circle $x^2 + y^2 - 4x + 6y - 12 = 0$ at A, B . Find A, B , length AB , midpoint M .

SOLUTION

Solution. Circle: $(x-2)^2 + (y+3)^2 = 25$. Centre $C(2, -3)$, $r = 5$. Substitute $y = x - 2$: $x^2 + (x-2)^2 - 4x + 6(x-2) - 12 = 0 \Rightarrow 2x^2 - 2x - 20 = 0 \Rightarrow x^2 - x - 10 = 0$. $\Delta = 1 + 40 = 41 > 0$ (chord).
 $x = \frac{1 \pm \sqrt{41}}{2}$, $y = x - 2 = \frac{-3 \pm \sqrt{41}}{2}$. $A\left(\frac{1+\sqrt{41}}{2}, \frac{-3+\sqrt{41}}{2}\right)$, $B\left(\frac{1-\sqrt{41}}{2}, \frac{-3-\sqrt{41}}{2}\right)$. Length $AB = \sqrt{(\sqrt{41})^2 + (\sqrt{41})^2} = \sqrt{82}$.
 Midpoint M : $x_M = \frac{1}{2}$, $y_M = -\frac{3}{2}$. Check: CM gradient $= \frac{-1.5+3}{0.5-2} = -1 \perp$ line gradient 1. ✓

DID YOU KNOW? Application: Satellite Coverage

A satellite at height h above Earth (radius R) covers a circular cap. The boundary circle on Earth's surface is the intersection of the Earth sphere with a cone from the satellite. In 2D cross-section, it reduces to line–circle intersection problems. Cameroon's SAT-1 satellite uses such geometry for footprint calculation.


18 INTERSECTION OF TWO CIRCLES, FAMILIES OF CIRCLES AND THE RADICAL AXIS

18.1 Intersection of Two Circles

Two circles $S_1 = 0$, $S_2 = 0$. Subtract to get the **Radical Axis** (a straight line):

$$S_1 - S_2 = 0 \implies 2(g_1 - g_2)x + 2(f_1 - f_2)y + (c_1 - c_2) = 0.$$

This line passes through the intersection points (if they intersect). Solve the system: Radical Axis + one Circle.



Two intersecting circles and their radical axis (line through P and Q).

18.2 Family of Circles Through Intersection of Two Circles

The family of circles passing through the intersection of $S_1 = 0$ and $S_2 = 0$ is:

$$S_1 + \lambda S_2 = 0 \quad (\lambda \neq -1) \quad \text{or} \quad S_1 + \lambda(S_1 - S_2) = 0.$$

For $\lambda = -1$ (first form) or $\lambda \rightarrow \infty$, we get the radical axis (a line). To find a particular circle through the intersection and a given point P , substitute P to find λ .

WORKED EXAMPLE 42 Radical Axis and Family of Circles

$S_1 : x^2 + y^2 - 4x + 2y - 7 = 0$, $S_2 : x^2 + y^2 + 2x - 6y + 5 = 0$. (i) Radical axis. (ii) Circle through intersection and $(1, 2)$.

SOLUTION

Solution. (i) $S_1 - S_2 = -6x + 8y - 12 = 0 \implies 3x - 4y + 6 = 0$. (ii) Family: $S_1 + \lambda S_2 = 0$. At $(1, 2)$: $S_1(1, 2) = 1 + 4 - 4 + 4 - 7 = -2$. $S_2(1, 2) = 1 + 4 + 2 - 12 + 5 = 0$. Equation: $-2 + \lambda(0) = 0 \implies -2 = 0$

(impossible for finite λ). This means $(1, 2)$ lies on S_2 but not S_1 . The required circle is S_2 itself (limit $\lambda \rightarrow \infty$). Equation: $x^2 + y^2 + 2x - 6y + 5 = 0$.

18.3 Orthogonal Circles

Two circles S_1, S_2 cut orthogonally if tangents at intersection are perpendicular. Condition: $2g_1g_2 + 2f_1f_2 = c_1 + c_2$. Equivalently: $d^2 = r_1^2 + r_2^2$ (Pythagoras at intersection point).

19 CONDITIONS FOR TWO CIRCLES TO TOUCH

19.1 Classification by Centre Distance

Let circles have centres C_1, C_2 , radii r_1, r_2 ($r_1 \geq r_2$), distance $d = C_1C_2$.

Position	Condition	Common Tangents	Point of Contact
Separate (outside)	$d > r_1 + r_2$	4	None
Touch externally	$d = r_1 + r_2$	3	Divides C_1C_2 internally in ratio $r_1 : r_2$
Intersect (2 pts)	$ r_1 - r_2 < d < r_1 + r_2$	2	N/A (chord = radical axis)
Touch internally	$d = r_1 - r_2$	1	Divides C_1C_2 externally in ratio $r_1 : r_2$
One inside other	$d < r_1 - r_2$	0	None
Concentric	$d = 0$	0 (or ∞ if $r_1 = r_2$)	N/A

METHOD Point of Contact Coordinates

For centres $C_1(x_1, y_1), C_2(x_2, y_2)$:

1. External touch: T divides C_1C_2 internally $r_1 : r_2$. $T = \left(\frac{r_2x_1 + r_1x_2}{r_1 + r_2}, \frac{r_2y_1 + r_1y_2}{r_1 + r_2} \right)$.
2. Internal touch ($r_1 > r_2$): T divides C_1C_2 externally $r_1 : r_2$ (on extension past C_2). $T = \left(\frac{r_2x_1 - r_1x_2}{r_2 - r_1}, \frac{r_2y_1 - r_1y_2}{r_2 - r_1} \right)$.

WORKED EXAMPLE 43 Touching Circles

$C_1 : x^2 + y^2 - 6x + 4y - 12 = 0$ (Centre $O_1(3, -2), r_1 = 5$). C_2 centre $O_2(10, -5)$ touches C_1 externally. Find r_2 , eq. of C_2 . If C_2 touches C_1 internally with $r_2 = 10$, find its centre.

SOLUTION

Solution. $d = O_1O_2 = \sqrt{(10-3)^2 + (-5+2)^2} = \sqrt{58}$. External: $d = r_1 + r_2 \Rightarrow \sqrt{58} = 5 + r_2 \Rightarrow r_2 = \sqrt{58} - 5$. Eq: $(x-10)^2 + (y+5)^2 = (\sqrt{58}-5)^2 = 83 - 10\sqrt{58}$.

Internal ($r_2 = 10$): $d = r_2 - r_1 = 5$. O_2 lies on circle centre O_1 radius 5: $(h-3)^2 + (k+2)^2 = 25$. If O_2 lies on line O_1O_2 (direction $(7, -3)$): $O_2 = (3, -2) \pm \frac{5}{\sqrt{58}}(7, -3)$. Two possible centres:

$$\left(3 + \frac{35}{\sqrt{58}}, -2 - \frac{15}{\sqrt{58}}\right) \text{ and } \left(3 - \frac{35}{\sqrt{58}}, -2 + \frac{15}{\sqrt{58}}\right).$$

20 CIRCLE THROUGH INTERSECTION OF A LINE AND A CIRCLE

20.1 Family of Circles Through Line–Circle Intersection

Let circle $S : x^2 + y^2 + 2gx + 2fy + c = 0$ and line $L : lx + my + n = 0$. The family of circles passing through their intersection points is:

$$S + \lambda L = 0 \Rightarrow x^2 + y^2 + 2gx + 2fy + c + \lambda(lx + my + n) = 0.$$

This represents a circle for all λ (coefficient of x^2, y^2 remains 1). It passes through the intersection of S and L (real or imaginary).

WORKED EXAMPLE 44 Circle Through Line–Circle Intersection

Find the circle through the intersection of $x^2 + y^2 - 4x + 2y - 4 = 0$ and $x + y - 1 = 0$ that passes through $(2, 3)$.

SOLUTION

Solution. Family: $x^2 + y^2 - 4x + 2y - 4 + \lambda(x + y - 1) = 0$. Passes through $(2, 3)$: $4 + 9 - 8 + 6 - 4 + \lambda(2 + 3 - 1) = 0 \Rightarrow 7 + 4\lambda = 0 \Rightarrow \lambda = -\frac{7}{4}$. Equation: $4(x^2 + y^2 - 4x + 2y - 4) - 7(x + y - 1) = 0$. $4x^2 + 4y^2 - 16x + 8y - 16 - 7x - 7y + 7 = 0$. $4x^2 + 4y^2 - 23x + y - 9 = 0$.

21 LOCUS PROBLEMS AND SYNTHESIS FOR THE GCE EXAM

21.1 General Strategy for Locus Problems

1. Let the moving point be $P(x, y)$.

2. Translate the geometric condition into an algebraic equation in x, y .
3. Simplify to a standard form (circle, line, parabola, etc.).
4. Identify the curve and state any restrictions (e.g., excluded points, arcs only).

21.2 Common Loci Leading to Circles

- **Constant distance from fixed point:** Circle.
- **Constant ratio of distances to two fixed points A, B ($k \neq 1$):** Circle (Apollonius Circle).
- **Angle in a semicircle / Constant angle subtended by segment AB :** Arc of a circle.
- **Equal tangent lengths to two circles:** Radical Axis (Line).
- **Sum of squares of distances to two fixed points constant:** Circle (centre at midpoint).

WORKED EXAMPLE 45 Locus: Ratio of Distances (Apollonius Circle)

$P(x, y)$ moves such that $PA : PB = 2 : 1$ for $A(1, 0), B(4, 0)$. Find locus.

SOLUTION

Solution. $PA = 2PB \implies PA^2 = 4PB^2$. $(x - 1)^2 + y^2 = 4[(x - 4)^2 + y^2]$. $x^2 - 2x + 1 + y^2 = 4x^2 - 32x + 64 + 4y^2$. $3x^2 - 30x + 63 + 3y^2 = 0 \implies x^2 - 10x + 21 + y^2 = 0$. $(x - 5)^2 + y^2 = 4$.
Locus: Circle centre $(5, 0)$, radius 2.

WORKED EXAMPLE 46 Locus: Equal Tangents to Two Circles

Find the locus of points from which tangents to circles $x^2 + y^2 = 4$ and $x^2 + y^2 - 6x + 8 = 0$ have equal length.

SOLUTION

Solution. Lengths: $\sqrt{x^2 + y^2 - 4}$ and $\sqrt{x^2 + y^2 - 6x + 8}$. Equate squares: $x^2 + y^2 - 4 = x^2 + y^2 - 6x + 8 \implies 6x = 12 \implies x = 2$. Locus is the radical axis (vertical line $x = 2$).

21.3 Synthesis: GCE Exam Style Questions

EXERCISE 15 GCE Style Practice

1. A circle has equation $x^2 + y^2 - 6x + 8y + k = 0$. Given that the radius is 5, find k and the coordinates of the centre.

2. The line $y = 2x - 1$ is a tangent to the circle $x^2 + y^2 + 4x - 2y + c = 0$. Find c and the point of contact.
3. Find the equation of the circle passing through the points $(1, 2)$, $(3, -4)$ and $(5, 2)$.
4. Two circles $C_1 : x^2 + y^2 = 25$ and $C_2 : x^2 + y^2 - 10x - 10y + 25 = 0$ intersect. Find the equation of their common chord and its length.
5. A point P moves so that the sum of the squares of its distances from $A(0, 0)$ and $B(4, 0)$ is 20. Find the equation of the locus of P and identify the curve.
6. Find the equations of the tangents from the point $(4, 0)$ to the circle $x^2 + y^2 = 4$.
7. A circle touches the x -axis at $(3, 0)$ and passes through $(1, 2)$. Find its equation.

SOLUTION Exercise 1

$g = -3, f = 4. r^2 = g^2 + f^2 - c = 9 + 16 - k = 25 - k$. Given $r = 5 \Rightarrow 25 - k = 25 \Rightarrow k = 0$.
Centre $(3, -4)$.

SOLUTION Exercise 2

Circle: $(x + 2)^2 + (y - 1)^2 = 5 - c$. Centre $(-2, 1), r^2 = 5 - c$. Line: $2x - y - 1 = 0$. Distance from centre to line $= r$. $r = \frac{|2(-2) - 1(1) - 1|}{\sqrt{5}} = \frac{6}{\sqrt{5}} \Rightarrow r^2 = \frac{36}{5}$. $5 - c = \frac{36}{5} \Rightarrow c = 5 - \frac{36}{5} = -\frac{11}{5}$. Point of contact: Foot of perpendicular from $(-2, 1)$ to line. Line perp: $x + 2y = 0$. Intersection with $2x - y = 1$: $x + 2(2x - 1) = 0 \Rightarrow 5x = 2 \Rightarrow x = 0.4, y = -0.2$. Contact: $(0.4, -0.2)$.

SOLUTION Exercise 3

Using method from Section 3 (or perpendicular bisectors). Centre $(3, -2/3)$, Radius $10/3$.
Equation: $3x^2 + 3y^2 - 18x + 4y - 5 = 0$.

SOLUTION Exercise 4

C_1 : Centre $(0, 0), r_1 = 5$. C_2 : $(x - 5)^2 + (y - 5)^2 = 25$, Centre $(5, 5), r_2 = 5$. Radical axis (common chord): $S_1 - S_2 = 0 \Rightarrow 10x + 10y - 50 = 0 \Rightarrow x + y - 5 = 0$. Distance from $C_1(0, 0)$ to chord: $d = 5/\sqrt{2}$. Half-chord $= \sqrt{25 - 25/2} = 5/\sqrt{2}$. Chord length $= 2 \times 5/\sqrt{2} = 5\sqrt{2}$.

SOLUTION Exercise 5

$PA^2 + PB^2 = 20 \implies x^2 + y^2 + (x-4)^2 + y^2 = 20$. $2x^2 - 8x + 16 + 2y^2 = 20 \implies x^2 - 4x + y^2 = 2 \implies (x-2)^2 + y^2 = 6$. Circle centre $(2, 0)$, radius $\sqrt{6}$.

SOLUTION Exercise 6

Circle $x^2 + y^2 = 4$. Point $(4, 0)$. Tangent form $y = m(x-4)$. Substitute: $x^2 + m^2(x-4)^2 = 4$. $(1+m^2)x^2 - 8m^2x + 16m^2 - 4 = 0$. $\Delta = 0 \implies 64m^4 - 4(1+m^2)(16m^2 - 4) = 0 \implies 16m^4 - (16m^4 + 12m^2 - 4) = 0 \implies 12m^2 = 4 \implies m = \pm 1/\sqrt{3}$. Tangents: $y = \pm \frac{1}{\sqrt{3}}(x-4)$ or $x \pm \sqrt{3}y - 4 = 0$. Points of contact: $(1, \pm\sqrt{3})$.

SOLUTION Exercise 7

Touches x -axis at $(3, 0) \implies$ Centre $(3, r)$ or $(3, -r)$. Radius r . Passes through $(1, 2)$: Distance from centre to $(1, 2) = r$. Case 1: Centre $(3, r)$. $(3-1)^2 + (r-2)^2 = r^2 \implies 4 + r^2 - 4r + 4 = r^2 \implies 8 = 4r \implies r = 2$. Centre $(3, 2)$. Eq: $(x-3)^2 + (y-2)^2 = 4$. Case 2: Centre $(3, -r)$. $(3-1)^2 + (-r-2)^2 = r^2 \implies 4 + r^2 + 4r + 4 = r^2 \implies 8 = -4r \implies r = -2$ (impossible, $r > 0$). Only one circle: $(x-3)^2 + (y-2)^2 = 4$.

KEY POINTS Key Formulas Summary for GCE Revision

- Standard Form: $(x-a)^2 + (y-b)^2 = r^2$. Centre (a, b) , Radius r .
- General Form: $x^2 + y^2 + 2gx + 2fy + c = 0$. Centre $(-g, -f)$, Radius $\sqrt{g^2 + f^2 - c}$.
- Diameter Form (endpoints A, B): $(x-x_1)(x-x_2) + (y-y_1)(y-y_2) = 0$.
- Tangent at $P(x_1, y_1)$ on $S = 0$: $xx_1 + yy_1 + g(x+x_1) + f(y+y_1) + c = 0$.
- Tangent Condition ($y = mx + d$ to $x^2 + y^2 = r^2$): $d^2 = r^2(1+m^2)$.
- Length of Tangent from $P(x_1, y_1)$: $\sqrt{S_{11}} = \sqrt{x_1^2 + y_1^2 + 2gx_1 + 2fy_1 + c}$.
- Parametric: $x = a + r \cos \theta$, $y = b + r \sin \theta$.
- Radical Axis of S_1, S_2 : $S_1 - S_2 = 0$.
- Family through $S_1 \cap S_2$: $S_1 + \lambda S_2 = 0$.
- Touching Circles: $d = r_1 + r_2$ (external), $d = |r_1 - r_2|$ (internal).

22 EXERCISES**22.1 I check my knowledge****EXERCISE 16 Multiple choice**

Choose the correct answer, justifying your choice briefly.

1. The circle $x^2 + y^2 - 4x + 6y - 12 = 0$ has centre and radius:
(a) centre $(2, -3)$, radius 5;
(b) centre $(-2, 3)$, radius 5;
(c) centre $(2, -3)$, radius 1;
(d) centre $(4, -6)$, radius 12.
2. The line $y = 2$ and the circle $x^2 + y^2 = 4$:
(a) meet in two distinct points;
(b) are tangent to each other;
(c) do not meet;
(d) meet at the origin.
3. The length of the tangent from the point $(0, 0)$ to the circle $x^2 + y^2 - 6x - 8y + 9 = 0$ is:
(a) 9;
(b) 3;
(c) 5;
(d) $\sqrt{41}$.

EXERCISE 17 True or false?

State whether each assertion is true or false, justifying your answer.

1. The equation $x^2 + y^2 + 2x - 4y + 9 = 0$ represents a circle of radius 2.
2. The point $(3, 4)$ lies on the circle $x^2 + y^2 = 25$.
3. Two circles touch externally if and only if the distance between their centres equals the sum of their radii.
4. The parametric equations $x = 1 + 3 \cos \theta$, $y = -2 + 3 \sin \theta$ represent a circle of centre

$(1, -2)$ and radius 9.

EXERCISE 18 Definitions

1. Define the **radical axis** of two circles and state one of its geometric properties.
2. State the condition on the distance d between the centres of two circles of radii r_1 and r_2 ($r_1 \geq r_2$) for the circles to touch internally.
3. Write down the general equation of a circle and state the conditions on its coefficients for it to represent a real circle.

EXERCISE 19 Direct calculations

1. Write down the equation of the circle of centre $(0, 0)$ and radius 7.
2. Write down the equation of the circle of centre $(-1, 4)$ and radius $\sqrt{5}$.
3. Find the centre and radius of the circle $x^2 + y^2 + 8x - 10y + 5 = 0$.
4. Find the equation of the circle having the segment joining $A(1, 3)$ and $B(5, 7)$ as a diameter.

22.2 I practise

EXERCISE 20 Centre and a point

Find the equation of the circle with centre $(2, -3)$ passing through the point $(5, 1)$. Determine whether the point $(6, -2)$ lies inside, on or outside this circle.

EXERCISE 21 Tangent at a point

A circle has equation $x^2 + y^2 - 6x + 4y - 12 = 0$.

1. Find its centre and its radius.
2. Verify that the point $(6, 2)$ lies on the circle.
3. Find the equation of the tangent to the circle at the point $(6, 2)$.

EXERCISE 22 Tangency of a line

Show that the line $3x + 4y = 25$ is a tangent to the circle $x^2 + y^2 = 25$, and find the coordinates of the point of contact.

EXERCISE 23 Length of a tangent

Find the length of the tangent from the point $(7, 1)$ to the circle $x^2 + y^2 - 4x + 6y - 3 = 0$.

22.3 I prepare for the GCE**EXERCISE 24** Circle through three points and a line condition

1. Find the equation of the circle passing through the points $A(1, 1)$, $B(2, -1)$ and $C(3, 2)$. (6 marks)
2. Find the equation of the tangent to this circle at the point $A(1, 1)$. (4 marks)
3. A second circle passes through the points $(2, 0)$ and $(6, 0)$ and has radius 5. Show that there are two such circles and find the equation of each. (6 marks)

EXERCISE 25 Two circles: intersection, common chord and touching

The circles C_1 and C_2 have equations

$$C_1 : x^2 + y^2 = 10, \quad C_2 : x^2 + y^2 - 6x - 2y = 10.$$

1. Find the equation of the common chord of C_1 and C_2 . (3 marks)
2. Find the coordinates of the points of intersection of the two circles. (5 marks)
3. Find the equation of the circle passing through the points of intersection of $x^2 + y^2 - 4 = 0$ and $x^2 + y^2 - 2x - 4y = 0$ and through the point $(1, 2)$. (5 marks)
4. Show that the circles $x^2 + y^2 - 2x - 4y - 20 = 0$ and $x^2 + y^2 + 6x + 2y - 90 = 0$ touch internally, and find the coordinates of their point of contact. (5 marks)

EXERCISE 26 Locus and circles under geometric conditions

1. A point P moves so that the sum of the squares of its distances from the points $A(-2, 0)$ and $B(2, 0)$ is equal to 20. Show that the locus of P is a circle, and find its centre and radius. (5 marks)
2. Find the equations of the circles which touch both coordinate axes and pass through the point $(2, 1)$. (5 marks)
3. Find the equation of the circle which passes through the point $(4, 1)$ and touches the line $x + y = 1$ at the point $(2, -1)$. (6 marks)

23 SOLUTIONS TO THE EXERCISES

SOLUTION Exercise 1 — Multiple choice

1. Completing the square for $x^2 - 4x + y^2 + 6y = 12$:

$$(x^2 - 4x + 4) + (y^2 + 6y + 9) = 12 + 4 + 9 \implies (x - 2)^2 + (y + 3)^2 = 25.$$

Centre $(2, -3)$, radius 5. Answer (a). Note: Answers (b) and (d) arise from sign errors on $-g$ and $-f$; answer (c) forgets to take the square root of 25.

2. Circle $x^2 + y^2 = 4$ (centre $O(0, 0)$, radius $r = 2$). Line $y = 2$ (i.e. $y - 2 = 0$). Perpendicular distance from O to the line:

$$d = \frac{|0 - 2|}{\sqrt{1}} = 2 = r.$$

The line is a tangent. Answer (b). Point of contact is $(0, 2)$, which satisfies $0^2 + 2^2 = 4$.

3. Circle $x^2 + y^2 - 6x - 8y + 9 = 0$. Centre $C(3, 4)$, radius $r = \sqrt{9 + 16 - 9} = 4$. Point $P(0, 0)$. Power of P : $S(0, 0) = 0 + 0 - 0 - 0 + 9 = 9$. Length of tangent $\ell = \sqrt{9} = 3$. Answer (b). Note: Answer (a) gives the power (squared length) instead of the length.

SOLUTION Exercise 2 — True or false?

- False. For $x^2 + y^2 + 2x - 4y + 9 = 0$, we have $g = 1, f = -2, c = 9$. $g^2 + f^2 - c = 1 + 4 - 9 = -4 < 0$. The equation represents no real circle (the “radius squared” would be negative).
- True. Substituting $(3, 4)$ into $x^2 + y^2 = 25$: $9 + 16 = 25$. The point lies on the circle.
- True. Two circles touch externally if and only if the distance between centres equals the sum of the radii: $d = r_1 + r_2$. They then have exactly one common point, which lies on the line of centres between the two centres.
- False. Parametric equations $x = 1 + 3 \cos \theta, y = -2 + 3 \sin \theta$. Centre is $(1, -2)$. The radius is the common coefficient of $\cos \theta$ and $\sin \theta$, namely 3 — not $3^2 = 9$. Indeed $(x - 1)^2 + (y + 2)^2 = 9 \cos^2 \theta + 9 \sin^2 \theta = 9$, so $r = 3$.

SOLUTION Exercise 3 — Definitions

1. The **radical axis** of two circles is the locus of points having equal powers with respect to the two circles. Algebraically, if the circles are $S_1 = 0$ and $S_2 = 0$, the radical axis is

$S_1 - S_2 = 0$, which is always a straight line. Geometric properties:

- The radical axis is perpendicular to the line of centres of the two circles.
- If the circles intersect in two points, the radical axis is the line through these points (the common chord).
- If the circles touch, the radical axis is their common tangent at the point of contact.

2. Two circles with radii $r_1 \geq r_2$ touch internally if and only if the distance between their centres equals the difference of the radii:

$$d = r_1 - r_2.$$

3. The general equation of a circle is

$$x^2 + y^2 + 2gx + 2fy + c = 0.$$

It represents a real circle if and only if $g^2 + f^2 - c > 0$; the centre is then $(-g, -f)$ and the radius $\sqrt{g^2 + f^2 - c}$. (If $g^2 + f^2 - c = 0$ the equation reduces to a single point — a point circle; if it is negative there is no real locus.)

SOLUTION Exercise 4 — Direct calculations

1. Centre $(0, 0)$, radius 7: $x^2 + y^2 = 49$.

2. Centre $(-1, 4)$, radius $\sqrt{5}$:

$$(x + 1)^2 + (y - 4)^2 = 5 \implies x^2 + 2x + 1 + y^2 - 8y + 16 = 5,$$

$$x^2 + y^2 + 2x - 8y + 12 = 0.$$

3. $x^2 + y^2 + 8x - 10y + 5 = 0$. Completing the square:

$$(x^2 + 8x + 16) + (y^2 - 10y + 25) = -5 + 16 + 25,$$

$$(x + 4)^2 + (y - 5)^2 = 36.$$

Centre $(-4, 5)$, radius 6.

4. Diameter form $(x - x_1)(x - x_2) + (y - y_1)(y - y_2) = 0$ with $A(1, 3)$, $B(5, 7)$:

$$(x - 1)(x - 5) + (y - 3)(y - 7) = 0 \implies x^2 - 6x + 5 + y^2 - 10y + 21 = 0,$$

$$x^2 + y^2 - 6x - 10y + 26 = 0.$$

Check: Centre $(3, 5)$ is the midpoint of AB ; radius $\sqrt{9 + 25 - 26} = \sqrt{8} = 2\sqrt{2}$. $AB = \sqrt{(5 - 1)^2 + (7 - 3)^2} = \sqrt{16 + 16} = \sqrt{32} = 4\sqrt{2}$, so radius = $\frac{1}{2}AB$. ✓

SOLUTION Exercise 5 — Centre and a point on the circle

Centre $C(2, -3)$, point on circle $P(5, 1)$. Radius squared:

$$r^2 = CP^2 = (5 - 2)^2 + (1 - (-3))^2 = 3^2 + 4^2 = 9 + 16 = 25, \quad r = 5.$$

Equation in centre-radius form:

$$(x - 2)^2 + (y + 3)^2 = 25.$$

Expanded form:

$$x^2 - 4x + 4 + y^2 + 6y + 9 = 25 \implies x^2 + y^2 - 4x + 6y - 12 = 0.$$

For the point $Q(6, -2)$, compute its squared distance to the centre:

$$CQ^2 = (6 - 2)^2 + (-2 + 3)^2 = 16 + 1 = 17.$$

Since $17 < 25 = r^2$, the point Q lies inside the circle.

SOLUTION Exercise 6 — Tangent at a given point

1. Circle: $x^2 + y^2 - 6x + 4y - 12 = 0$. Completing the square:

$$(x^2 - 6x + 9) + (y^2 + 4y + 4) = 12 + 9 + 4 \implies (x - 3)^2 + (y + 2)^2 = 25.$$

Centre $C(3, -2)$, radius $r = 5$.

2. Check if $T(6, 2)$ lies on the circle:

$$6^2 + 2^2 - 6(6) + 4(2) - 12 = 36 + 4 - 36 + 8 - 12 = 0.$$

Yes, T lies on the circle.

3. Gradient of radius CT :

$$m_{CT} = \frac{2 - (-2)}{6 - 3} = \frac{4}{3}.$$

The tangent at T is perpendicular to CT , so its gradient is $m_T = -\frac{3}{4}$. Equation of tangent:

$$y - 2 = -\frac{3}{4}(x - 6) \implies 4y - 8 = -3x + 18 \implies 3x + 4y - 26 = 0.$$

Verification using the “ $T = 0$ ” formula for $x^2 + y^2 + 2gx + 2fy + c = 0$ at (x_1, y_1) : $xx_1 + yy_1 + g(x + x_1) + f(y + y_1) + c = 0$. Here $g = -3$, $f = 2$, $c = -12$, $(x_1, y_1) = (6, 2)$:

$$6x + 2y - 3(x + 6) + 2(y + 2) - 12 = 0 \implies 3x + 4y - 18 + 4 - 12 = 0 \implies 3x + 4y - 26 = 0. \quad \checkmark$$

SOLUTION Exercise 7 — Tangency of a line to a circle

Circle $C: x^2 + y^2 = 25$ (centre $O(0, 0)$, radius $r = 5$). Line $L: 3x + 4y - 25 = 0$. Perpendicular distance from centre O to line L :

$$d = \frac{|3(0) + 4(0) - 25|}{\sqrt{3^2 + 4^2}} = \frac{25}{5} = 5.$$

Since $d = r = 5$, the line is a tangent to the circle.

The point of contact T is the foot of the perpendicular from O to L . The normal vector to L is $(3, 4)$. Parametric form of perpendicular through O : $(x, y) = (3t, 4t)$. Substitute into L : $3(3t) + 4(4t) - 25 = 0 \implies 25t = 25 \implies t = 1$. Hence $T = (3, 4)$. Check: T lies on L : $3(3) + 4(4) = 25$. T lies on C : $3^2 + 4^2 = 25$. ✓

SOLUTION Exercise 8 — Length of a tangent from an external point

Circle: $x^2 + y^2 - 4x + 6y - 3 = 0$. External point $P(7, 1)$. Length of tangent $\ell = \sqrt{S(7, 1)}$, where $S(x, y) = x^2 + y^2 - 4x + 6y - 3$.

$$\ell = \sqrt{7^2 + 1^2 - 4(7) + 6(1) - 3} = \sqrt{49 + 1 - 28 + 6 - 3} = \sqrt{25} = \boxed{5}.$$

Geometric check: Centre $C(2, -3)$, radius $r = \sqrt{4 + 9 + 3} = 4$. Distance $CP = \sqrt{(7-2)^2 + (1+3)^2} = \sqrt{25 + 16} = \sqrt{41}$. By Pythagoras in right triangle CPT (where T is point of contact): $\ell^2 = CP^2 - r^2 = 41 - 16 = 25$, so $\ell = 5$. ✓

SOLUTION Exercise 9 — Circle through three points and tangent condition

1. Let the circle be $x^2 + y^2 + 2gx + 2fy + c = 0$. Substituting the three points:

$$A(1, 1): 1 + 1 + 2g + 2f + c = 0 \implies 2g + 2f + c = -2 \quad (1)$$

$$B(2, -1): 4 + 1 + 4g - 2f + c = 0 \implies 4g - 2f + c = -5 \quad (2)$$

$$C(3, 2): 9 + 4 + 6g + 4f + c = 0 \implies 6g + 4f + c = -13 \quad (3)$$

$$(2) - (1): 2g - 4f = -3 \quad (4)$$

$$(3) - (2): 2g + 6f = -8 \quad (5)$$

$$(5) - (4): 10f = -5 \implies f = -\frac{1}{2}. \text{ From (4): } 2g = 4f - 3 = -2 - 3 = -5 \implies g = -\frac{5}{2}. \text{ From (1): } c = -2 - 2g - 2f = -2 + 5 + 1 = 4. \text{ Equation:}$$

$$\boxed{x^2 + y^2 - 5x - y + 4 = 0.}$$

Centre $\left(\frac{5}{2}, \frac{1}{2}\right)$, radius $r = \sqrt{\frac{25}{4} + \frac{1}{4} - 4} = \sqrt{\frac{10}{4}} = \sqrt{\frac{5}{2}}$. Examiner's expectation: Any correct method (general equation or perpendicular bisectors) earns full credit; the three equations must be set up correctly before any method mark for solving is awarded.

2. Tangent at $A(1, 1)$. Centre $O\left(\frac{5}{2}, \frac{1}{2}\right)$. Gradient of radius OA :

$$m_{OA} = \frac{1 - \frac{1}{2}}{1 - \frac{5}{2}} = \frac{\frac{1}{2}}{-\frac{3}{2}} = -\frac{1}{3}.$$

Gradient of tangent (perpendicular to radius): $m_T = 3$. Equation:

$$y - 1 = 3(x - 1) \Rightarrow \boxed{3x - y - 2 = 0.}$$

3. Circle of radius 5 passing through $(2, 0)$ and $(6, 0)$. The centre is equidistant from $(2, 0)$ and $(6, 0)$, so it lies on the perpendicular bisector of this segment. Midpoint is $(4, 0)$, segment is horizontal, so perpendicular bisector is the vertical line $x = 4$. Let centre be $(4, k)$. Then

$$r^2 = (4 - 2)^2 + (k - 0)^2 = 4 + k^2 = 25 \Rightarrow k^2 = 21 \Rightarrow k = \pm\sqrt{21}.$$

Since k takes two distinct values, there are exactly two such circles. Centres: $(4, \sqrt{21})$ and $(4, -\sqrt{21})$. Equations:

$$\boxed{(x - 4)^2 + (y - \sqrt{21})^2 = 25 \quad \text{and} \quad (x - 4)^2 + (y + \sqrt{21})^2 = 25.}$$

Expanded: $x^2 + y^2 - 8x \mp 2\sqrt{21}y + 16 + 21 - 25 = 0$, i.e.

$$\boxed{x^2 + y^2 - 8x \mp 2\sqrt{21}y + 12 = 0.}$$

Examiner's expectation: The explicit statement "two values of k , hence two circles" is required for the justification mark.

SOLUTION Exercise 10 — Intersection of two circles, common chord and family of circles

Given circles:

$$C_1 : x^2 + y^2 - 10 = 0 \quad (\text{Centre } O_1(0, 0), \text{ radius } r_1 = \sqrt{10})$$

$$C_2 : x^2 + y^2 - 6x - 2y - 10 = 0 \quad (\text{Centre } O_2(3, 1), \text{ radius } r_2 = \sqrt{9 + 1 + 10} = \sqrt{20} = 2\sqrt{5})$$

1. The common chord (radical axis) is $C_1 - C_2 = 0$:

$$(x^2 + y^2 - 10) - (x^2 + y^2 - 6x - 2y - 10) = 0 \Rightarrow 6x + 2y = 0 \Rightarrow \boxed{3x + y = 0.}$$

2. On the chord, $y = -3x$. Substitute into C_1 :

$$x^2 + (-3x)^2 = 10 \Rightarrow 10x^2 = 10 \Rightarrow x^2 = 1 \Rightarrow x = \pm 1.$$

Corresponding y : $x = 1 \Rightarrow y = -3$; $x = -1 \Rightarrow y = 3$. Points of intersection:

$$(1, -3) \text{ and } (-1, 3).$$

Check in C_2 : $1 + 9 - 6 + 6 - 10 = 0$ and $1 + 9 + 6 - 6 - 10 = 0$. ✓

3. Family of circles through the intersections of C_1 and C_2 :

$$C_1 + \lambda C_2 = 0 \Rightarrow (x^2 + y^2 - 10) + \lambda(x^2 + y^2 - 6x - 2y - 10) = 0.$$

This circle passes through $(1, 2)$:

$$(1 + 4 - 10) + \lambda(1 + 4 - 6 - 4 - 10) = 0 \Rightarrow -5 + \lambda(-15) = 0 \Rightarrow \lambda = -\frac{1}{3}.$$

Substitute $\lambda = -\frac{1}{3}$ and multiply by 3:

$$3(x^2 + y^2 - 10) - (x^2 + y^2 - 6x - 2y - 10) = 0$$

$$3x^2 + 3y^2 - 30 - x^2 - y^2 + 6x + 2y + 10 = 0$$

$$2x^2 + 2y^2 + 6x + 2y - 20 = 0 \text{ or } x^2 + y^2 + 3x + y - 10 = 0.$$

SOLUTION Exercise 10 (continued) — Conditions for circles to touch

Given circles:

$$C_1 : x^2 + y^2 - 2x - 4y - 20 = 0 \Rightarrow \text{Centre } O_1(1, 2), r_1 = \sqrt{1 + 4 + 20} = 5.$$

$$C_2 : x^2 + y^2 + 6x + 2y - 90 = 0 \Rightarrow \text{Centre } O_2(-3, -1), r_2 = \sqrt{9 + 1 + 90} = 10.$$

Distance between centres:

$$d = O_1O_2 = \sqrt{(1+3)^2 + (2+1)^2} = \sqrt{16+9} = 5.$$

Compare d with radii:

$$r_2 - r_1 = 10 - 5 = 5 = d \text{ and } d \neq 0.$$

Since $d = |r_2 - r_1|$, the circles touch internally.

The point of contact T lies on the line of centres O_1O_2 , on the side of O_1 opposite to O_2 (since $r_2 > r_1$). Vector $\vec{O_2O_1} = (4, 3)$. Unit vector $\left(\frac{4}{5}, \frac{3}{5}\right)$. Starting from O_1 and moving a distance $r_1 = 5$ along this direction:

$$T = O_1 + 5\left(\frac{4}{5}, \frac{3}{5}\right) = (1, 2) + (4, 3) = (5, 5).$$

Check: $O_2T = \sqrt{(5+3)^2 + (5+1)^2} = \sqrt{64+36} = 10 = r_2$. Also T satisfies C_1 : $25+25-10-20-20 = 0$. ✓

SOLUTION Exercise 11 — Locus problems and circles under geometric conditions

1. Let $P(x, y)$ be the moving point. Fixed points $A(-2, 0)$, $B(2, 0)$. Condition: $PA^2 + PB^2 = 20$.

$$(x + 2)^2 + y^2 + (x - 2)^2 + y^2 = 20.$$

Expand:

$$(x^2 + 4x + 4) + y^2 + (x^2 - 4x + 4) + y^2 = 20$$

$$2x^2 + 2y^2 + 8 = 20 \implies 2x^2 + 2y^2 = 12 \implies x^2 + y^2 = 6.$$

This is a circle with centre $(0, 0)$ — the midpoint of AB — and radius $\sqrt{6}$.

2. A circle in the first quadrant touching both coordinate axes has centre (a, a) and radius a ($a > 0$). It passes through $(2, 1)$:

$$(2 - a)^2 + (1 - a)^2 = a^2$$


$$4 - 4a + a^2 + 1 - 2a + a^2 = a^2$$

$$a^2 - 6a + 5 = 0 \implies (a - 1)(a - 5) = 0 \implies a = 1 \text{ or } a = 5.$$

Two circles satisfy the conditions:

$$(x - 1)^2 + (y - 1)^2 = 1 \quad \text{and} \quad (x - 5)^2 + (y - 5)^2 = 25.$$

Check for $a = 5$: $(2 - 5)^2 + (1 - 5)^2 = 9 + 16 = 25$. ✓



Exercise 11.2: Two circles touching both axes and passing through $(2, 1)$.

SOLUTION Exercise 11 (continued) — Circle touching a line at a given point

- Circle touches the line $L : x + y = 1$ at $T(2, -1)$ and passes through $P(4, 1)$. Since the circle touches L at T , the centre C lies on the perpendicular to L at T . Line L has gradient -1 , so the perpendicular has gradient 1 . Parametric form of perpendicular through $T(2, -1)$: $C = (2 + t, -1 + t)$. Radius squared is the squared distance CT^2 :

$$r^2 = t^2 + t^2 = 2t^2.$$

The circle passes through $P(4, 1)$, so $CP^2 = r^2$:


$$(4 - (2 + t))^2 + (1 - (-1 + t))^2 = 2t^2$$

$$(2 - t)^2 + (2 - t)^2 = 2t^2 \implies 2(2 - t)^2 = 2t^2 \implies (2 - t)^2 = t^2.$$

Hence $2 - t = \pm t$. Case $2 - t = -t \Rightarrow 2 = 0$ (impossible). Case $2 - t = t \Rightarrow 2t = 2 \Rightarrow t = 1$. Centre $C = (3, 0)$, $r^2 = 2(1)^2 = 2$. Equation:


$$(x - 3)^2 + y^2 = 2 \quad \text{i.e.} \quad x^2 + y^2 - 6x + 7 = 0.$$

Verification: Distance from $C(3, 0)$ to line $x + y - 1 = 0$: $\frac{|3+0-1|}{\sqrt{2}} = \frac{2}{\sqrt{2}} = \sqrt{2} = r$. ✓ Foot of perpendicular from C to L : $C - \frac{3+0-1}{2}(1, 1) = (3, 0) - (1, 1) = (2, -1) = T$. ✓ $P(4, 1)$: $(4 - 3)^2 + 1^2 = 2 = r^2$. ✓



Exercise 11.3: Circle touching $x + y = 1$ at $(2, -1)$ and passing through $(4, 1)$.

24 SUMMARY SHEET



Mind map of the chapter Circle Geometry.

KEY POINTS Circle Geometry — Summary Sheet

Key definitions.

- A **circle** is the locus of a point P whose distance from a fixed point (the **centre**) is constant (the **radius**).
- A **tangent** is a line touching the circle at exactly one point; it is perpendicular to the radius at the point of contact.
- The **radical axis** of two circles is the locus of points with equal tangent lengths (equal powers) to both circles.

Equations of a circle (know them all!).

- Centre $O(0, 0)$, radius r : $x^2 + y^2 = r^2$.
- Centre $C(a, b)$, radius r : $(x - a)^2 + (y - b)^2 = r^2$.
- General form: $x^2 + y^2 + 2gx + 2fy + c = 0$, with centre $(-g, -f)$ and radius $r = \sqrt{g^2 + f^2 - c}$ (real circle iff $g^2 + f^2 - c > 0$).

- Diameter form, ends $A(x_1, y_1), B(x_2, y_2)$: $(x - x_1)(x - x_2) + (y - y_1)(y - y_2) = 0$ (from $\vec{PA} \cdot \vec{PB} = 0$).
- Parametric form: $x = a + r \cos \theta, y = b + r \sin \theta$; convert to Cartesian by $\cos^2 \theta + \sin^2 \theta = 1$.
- Circle touching both axes: centre $(\pm r, \pm r)$, equation $(x \pm r)^2 + (y \pm r)^2 = r^2$.

Tangents.

- Tangent at (x_1, y_1) on $x^2 + y^2 = r^2$: $xx_1 + yy_1 = r^2$; on the general circle: $xx_1 + yy_1 + g(x + x_1) + f(y + y_1) + c = 0$.
- Line $y = mx + c$ touches $x^2 + y^2 = r^2$ iff $c^2 = r^2(1 + m^2)$; generally: perpendicular distance from centre to line equals r .
- Length of tangent from external point $P(x_1, y_1)$: $\ell = \sqrt{x_1^2 + y_1^2 + 2gx_1 + 2fy_1 + c} = \sqrt{CP^2 - r^2}$.

Intersections and families.

- Line $\cap$ circle: substitute, get a quadratic; $\Delta > 0$ two points, $\Delta = 0$ tangent, $\Delta < 0$ no intersection.
- Family through intersection of circle $S = 0$ and line $L = 0$: $S + \lambda L = 0$.
- Family through intersections of circles $S_1 = 0, S_2 = 0$: $S_1 + \lambda S_2 = 0$; radical axis: $S_1 - S_2 = 0$ (a straight line, the common chord if they intersect).
- Two circles touch externally iff $d = r_1 + r_2$; internally iff $d = |r_1 - r_2|$; intersect iff $|r_1 - r_2| < d < r_1 + r_2$; separate iff $d > r_1 + r_2$.

Methods.

- Circle through 3 points: substitute into the general equation, solve the 3×3 system for g, f, c — or use perpendicular bisectors of two chords.
- Centre on a line: write the centre as (h, k) with the line's relation, then equate radii to the given points.
- Locus: let $P(x, y)$, translate the geometric condition into an equation, simplify.

Common mistakes.

- Sign errors: centre is $(-g, -f)$, not (g, f) ; radius is $\sqrt{g^2 + f^2 - c}$, not $\sqrt{g^2 + f^2 + c}$.
- Forgetting to divide by the coefficient of x^2 before reading off g, f, c .
- Using $d = r_1 + r_2$ for internal touching.
- Mixing up point of tangency and external point in tangent formulas.

GCE tips.

- Always state centre and radius first — most questions build on them.
- Check your final circle passes through all given points.
- Show the discriminant or distance condition clearly: method marks are awarded for the condition, not just the answer.



Signed • OnBuch+
ONBUCH+ TEACHING TEAM
FOLLOWS THE OFFICIAL MINESEC SYLLABUS

OnBuch+