

PURE MATHEMATICS WITH MECHANICS • UPPER SIXTH

UPPER SIXTH — PURE MATHEMATICS WITH MECHANICS

CHAPTER 9

KINEMATICS OF PARTICLES MOVING IN A STRAIGHT LINE

SUGGESTED TIME: 10 PERIODS

IN THIS CHAPTER

This chapter introduces the mathematical description of motion in one dimension: displacement, velocity, acceleration and their graphical representations. Learners master the equations of uniformly accelerated motion and apply them to free fall and real-life situations, in full preparation for the GCE Advanced Level mechanics paper.

- By the end of this chapter I can distinguish between distance and displacement, speed and velocity, and average and instantaneous quantities.
- By the end of this chapter I can state and use the five equations of motion under uniform acceleration (suvat equations).
- By the end of this chapter I can select the appropriate suvat equation for a given problem and solve it systematically.
- By the end of this chapter I can plot, read and interpret displacement–time, velocity–time and acceleration–time graphs.
- By the end of this chapter I can deduce velocity from the gradient of an x – t graph and displacement from the area under a v – t graph.
- By the end of this chapter I can connect the three types of motion graphs and sketch any one of them from another.



DOWNLOAD ON- BUCH

Free • Google Play
AI tutor, past papers,
results



GO ONBUCH+

All signed courses, unlimited AI
tutor, PDF sheets — OnBuch+
tab in the app

THIS SIGNED COURSE CONTAINS

COURSE

complete, illustrated,
step by step

METHODS

and worked examples

EXERCISES

exam style + solutions

SUMMARY SHEET

the essentials to
remember



Signed • OnBuch+

ONBUCH+ TEACHING TEAM
FOLLOWS THE OFFICIAL MINESEC SYLLABUS

OnBuch+

CONTENTS

- | | |
|---|--|
| 1. Section 1: Concept of Motion in One Dimension (Lesson 92) | Gravity (Lessons 99, 100) |
| 2. Section 2: Equations of Motion under Uniform Acceleration (Lesson 93) | 6. Section 6: Real-Life Applications and Synthesis (Lessons 94 & 101, consolidation) |
| 3. Section 3: Displacement–Time and Velocity–Time Graphs (Lessons 94, 95, 96) | 7. Integration activity |
| 4. Section 4: Acceleration–Time Graphs and Connecting the Three Graphs (Lessons 97, 98) | 8. Methods and worked examples |
| 5. Section 5: Free Fall and Motion under | 9. Exercises |
| | 10. Solutions to the exercises |
| | 11. Summary sheet |

CHAPTER OBJECTIVES

- By the end of this chapter I can distinguish between distance and displacement, speed and velocity, and average and instantaneous quantities.
- By the end of this chapter I can state and use the five equations of motion under uniform acceleration (suvat equations).
- By the end of this chapter I can select the appropriate suvat equation for a given problem and solve it systematically.
- By the end of this chapter I can plot, read and interpret displacement–time, velocity–time and acceleration–time graphs.
- By the end of this chapter I can deduce velocity from the gradient of an x – t graph and displacement from the area under a v – t graph.
- By the end of this chapter I can connect the three types of motion graphs and sketch any one of them from another.
- By the end of this chapter I can model free fall as motion under uniform acceleration $g \approx 9.8 \text{ m/s}^2$ and solve vertical motion problems.
- By the end of this chapter I can apply kinematics to real-life situations such as braking distances, road safety and falling objects.
- By the end of this chapter I can solve GCE-style problems involving two moving particles, meeting points and overtaking.

PREREQUISITES

- Algebraic manipulation: solving linear and quadratic equations, rearranging formulae (Lower Sixth algebra).
- Gradient of a straight line and area of basic shapes (rectangle, triangle, trapezium).
- Plotting and reading Cartesian graphs; equation of a line $y = mx + c$.
- Basic SI units (metre, second) and unit conversions ($\text{km/h} \leftrightarrow \text{m/s}$).
- Simple differentiation of polynomials is helpful but NOT required (flagged as enrichment only).

1 SECTION 1: CONCEPT OF MOTION IN ONE DIMENSION (LESSON 92)

Every morning, the bend-skin motorbike rider in Bamenda must judge in a split second whether he can stop before the junction when the traffic light turns orange. A driver on the Yaoundé-Douala highway who follows too closely at 100 km h^{-1} may never stop in time, while a mango falling from a tree in Buea accelerates in exactly the same way as a stone dropped from a storey building. How far does the bike travel before stopping? How long does the mango take to hit the ground? These everyday questions are all answered by one powerful tool: the **kinematics** of a particle moving in a straight line.

In this first section we build the vocabulary of kinematics: what a particle is, how we describe where it is (position, displacement), how fast it moves (speed, velocity) and how its motion changes (acceleration). Everything in the rest of the chapter rests on these ideas, so we shall define them carefully.

1.1 The Particle Model

Look again at the bend-skin. It has a length, a width, wheels that spin, a rider who leans into corners. If we tried to describe the motion of every part of it, the problem would be impossibly complicated. But if all we want to know is how far along the road the bike is at each instant, then its size and shape do not matter: we can pretend the whole bike, rider included, is a single point carrying the total mass.

DEFINITION The particle model

A **particle** is a body whose dimensions are negligible compared with the distances involved in its motion. We model it as a single point at which the whole mass of the body is concentrated.

WARNING The particle model is an idealisation

No real object is a point. The model is valid when the size of the body is small compared with the distances travelled and when we do not care about rotation or internal motion. A bend-skin over a 500 m stretch of road is a particle; the same bend-skin parking in a 2 m gap is not. Always state the modelling assumptions you use in an examination answer.

DID YOU KNOW? Why physicists love simple models

The particle model is the first of a family of idealisations you will meet in mechanics: the light inextensible string, the smooth pulley, the smooth plane. Each one removes a complication (mass of the string, friction, ...) so that the essential mathematics appears clearly. Master-

ing mechanics is largely learning which idealisation fits which situation.

1.2 Position and Displacement

To describe motion along a straight line we turn the line into a number line: we choose an **origin** O (a reference point) and a **positive direction**. The **position** of a particle P at time t is then a single number x , its directed distance from O , measured in metres (m).

DEFINITION Displacement

The **displacement** of a particle between two instants is the change of position

$$s = \Delta x = x_{\text{final}} - x_{\text{initial}}.$$

Displacement is a vector quantity: it has a magnitude (in metres) and a direction (given by its sign along the chosen axis). A negative displacement simply means “in the negative direction”.

The **distance** travelled, by contrast, is the total length of the path actually covered. It is a **scalar**: always positive, with no direction attached.

PROPERTY Distance versus displacement

For any motion,

$$\text{distance travelled} \geq |\text{displacement}|,$$

with equality if and only if the particle moves in one fixed direction throughout (it never reverses). Whenever the particle turns back, the distance travelled strictly exceeds the magnitude of the displacement.


The reason is clear: displacement only “sees” the start and the finish, while distance accumulates every metre covered, including any backtracking. A round trip back to the start gives zero displacement but a large distance.

EXAMPLE Distance and displacement on a number line

A particle starts at $x = -2$ m, moves to $x = 5$ m, then returns to $x = 1$ m.

- Displacement: $\Delta x = 1 - (-2) = +3$ m.
- Distance travelled: $|-2 \rightarrow 5| + |5 \rightarrow 1| = 7 + 4 = 11$ m.

Note that $11 > |3|$, exactly as the property predicts, because the particle reversed at $x = 5$.



A particle goes from $x = -2$ m to $x = 5$ m, then back to $x = 1$ m. Distance travelled = 11 m; displacement = +3 m.

1.3 Average Speed and Average Velocity

Knowing where a particle is at two instants, the natural next question is: how fast did it get there?

DEFINITION Average speed and average velocity

$$\text{average speed} = \frac{\text{distance travelled}}{\text{time taken}}, \quad \text{average velocity} = \frac{\text{displacement}}{\text{time taken}} = \frac{\Delta x}{\Delta t}.$$

Average speed is a scalar; average velocity is a vector (its sign gives the direction). The SI unit of both is the metre per second, m s^{-1} ; in everyday life we also use kilometres per hour, km h^{-1} .

PROPERTY Conversion of units

$$1 \text{ m s}^{-1} = 3.6 \text{ km h}^{-1}, \quad 1 \text{ km h}^{-1} = \frac{1}{3.6} \text{ m s}^{-1}.$$

Derivation (not examinable but instructive): 1 m s^{-1} means 1 m in 1 s, i.e. 3600 m in 3600 s, i.e. 3.6 km in 1 hour.

WARNING Mixing units

The most frequent source of wrong answers in kinematics is mixing km h^{-1} with m s^{-1} , or hours with seconds. Before any calculation, convert everything to SI units (metres, seconds, m s^{-1}). For example, $90 \text{ km/h} = 90/3.6 = 25 \text{ m/s}$.

WORKED EXAMPLE 1 Average speed and average velocity

A trader walks from her stall, taken as origin, 120 m east to the market in 100 s, then walks 40 m back west in 40 s to meet a customer. Take east as positive. Find (a) the total distance, (b) the displacement, (c) her average speed, (d) her average velocity for the whole journey.

SOLUTION

Solution.

- (a) Distance = $120 + 40 = 160$ m.
- (b) Displacement $\Delta x = 120 - 40 = +80$ m (i.e. 80 m east of the stall).
- (c) Total time = $100 + 40 = 140$ s. Average speed = $\frac{160}{140} \approx 1.14 \text{ m s}^{-1}$.
- (d) Average velocity = $\frac{\Delta x}{\Delta t} = \frac{80}{140} \approx +0.57 \text{ m s}^{-1}$, directed east.

Notice the average speed is about twice the magnitude of the average velocity, because of the backtracking.

1.4 Instantaneous Velocity

Average velocity over a whole journey can hide a lot: the bend-skin may average 30 km/h yet reach 60 km/h on a clear stretch. To describe the motion at one instant, we shrink the time interval.

Imagine computing the average velocity $\frac{\Delta x}{\Delta t}$ over the interval $[t, t + \Delta t]$, and letting Δt become smaller and smaller. The ratio settles down to a definite value: the **instantaneous velocity** at time t .

DEFINITION Instantaneous velocity

$$v = \lim_{\Delta t \rightarrow 0} \frac{\Delta x}{\Delta t} = \frac{dx}{dt}.$$

The instantaneous velocity is the gradient of the position–time graph at time t (we exploit this fully in Section 3). Its magnitude is the instantaneous **speed**.

WARNING Speed is not velocity

Speed is a scalar (how fast); velocity is a vector (how fast and in which direction). Two cars moving at 50 km/h in opposite directions have the same speed but opposite velocities. In a problem with a chosen positive direction, a velocity of -8 m/s means a speed of 8 m/s .

in the negative direction. Never drop the sign, and never report a negative speed.

1.5 Acceleration

Vehicles rarely keep a constant velocity: they speed up, slow down, stop. **Acceleration** measures how quickly the velocity changes.

DEFINITION Acceleration

The **average acceleration** over a time interval is

$$a = \frac{\Delta v}{\Delta t} = \frac{v_{\text{final}} - v_{\text{initial}}}{\Delta t},$$

and the instantaneous acceleration is $a = \frac{dv}{dt} = \frac{d^2x}{dt^2}$. Acceleration is a vector; its SI unit is the metre per second squared, m s^{-2} .

The sign of a is read along the chosen positive axis:

- $a > 0$: the velocity is increasing in the positive direction (speeding up if moving positively, slowing down if moving negatively);
- $a < 0$: the velocity is decreasing in the positive direction.

DEFINITION Deceleration (retardation)

When a particle is slowing down, its acceleration is opposite to its velocity. We then speak of **deceleration** or **retardation**: a deceleration of 3 m s^{-2} means an acceleration $a = -3 \text{ m s}^{-2}$ (with the direction of motion taken as positive).

WARNING Negative acceleration is not always slowing down

A negative acceleration only means the acceleration vector points in the negative direction. If the particle is already moving in the negative direction ($v < 0$), a negative acceleration makes it go faster. Deceleration occurs exactly when a and v have opposite signs.

EXAMPLE Reading acceleration

A car moving in the positive direction increases its velocity from 10 m/s to 22 m/s in 4 s:

$$a = \frac{22 - 10}{4} = +3 \text{ m s}^{-2}.$$

If instead it brakes from 22 m/s to 10 m/s in 4 s, then $a = \frac{10 - 22}{4} = -3 \text{ m s}^{-2}$: a deceleration of 3 m s^{-2} .

1.6 Uniform and Non-Uniform Acceleration

Acceleration itself may change with time (a car in town traffic) or stay constant (a freely falling stone, neglecting air resistance).


DEFINITION Uniform acceleration

Acceleration is **uniform** (constant) when the velocity changes by the same amount in every equal time interval: a is a constant. Otherwise the acceleration is **non-uniform** (variable).

This chapter focuses almost entirely on **uniform acceleration**, for two reasons: it describes many real situations well (free fall, a vehicle braking firmly), and it leads to the simple, powerful constant-acceleration formulae of Section 2. Variable acceleration requires calculus and is treated later in the course.

METHOD Choosing signs and keeping them

1. Draw a quick sketch of the situation and mark the straight line of motion.
2. Choose an origin O and a positive direction; write this choice down explicitly.
3. Express every given quantity (position, velocity, acceleration) as a signed number along this axis.
4. Carry the signs through every formula; never “fix” a sign at the end by guesswork.
5. Interpret the sign of each answer in words (“the particle is 3 m to the left of O ”, “it is moving in the negative direction”).



A bend-skin braking on a straight road: positions at $t = 0, 2, 4$ s with velocity arrows. The arrows shrink steadily — the signature of uniform (constant) deceleration.

WORKED EXAMPLE 2 Velocity and acceleration with signs

A particle moves along the x -axis (positive to the right). At $t = 1$ s its velocity is $+6$ m/s; at $t = 5$ s its velocity is -2 m/s. (a) Find its average acceleration over this interval. (b) Explain in words what happened to the particle.

SOLUTION

Solution.

- (a) $a = \frac{\Delta v}{\Delta t} = \frac{-2 - 6}{5 - 1} = \frac{-8}{4} = -2 \text{ m s}^{-2}$.
- (b) The particle was first moving to the right at 6 m/s. The constant negative acceleration slowed it down, brought it momentarily to rest, then made it move to the left, reaching 2 m/s leftwards by $t = 5$ s. Note the trap: the acceleration is negative throughout, yet the particle only decelerates while $v > 0$; once $v < 0$, the same negative acceleration speeds it up in the negative direction.

EXERCISE 1

A student jogs 300 m due north in 150 s, then 100 m due south in 50 s. Taking north as positive, find (a) the distance travelled, (b) the displacement, (c) the average speed, (d) the average velocity. Convert the average speed to km h^{-1} .

SOLUTION Exercise 1

(a) Distance = $300 + 100 = 400$ m. (b) Displacement = $300 - 100 = +200$ m (north). (c) Average speed = $\frac{400}{200} = 2 \text{ m s}^{-1} = 2 \times 3.6 = 7.2 \text{ km h}^{-1}$. (d) Average velocity = $\frac{200}{200} = +1 \text{ m s}^{-1}$, directed north.

KEY POINTS Key points of Section 1

- A **particle** is a point mass: a valid model when size is negligible compared with distances.
- **Displacement** $\Delta x = x_f - x_i$ is a vector; **distance** is a scalar, and distance $\geq |\text{displacement}|$.
- Average velocity $= \frac{\Delta x}{\Delta t}$ (vector); average speed $= \frac{\text{distance}}{\text{time}}$ (scalar). Instantaneous velocity $v = \frac{dx}{dt}$.
- Acceleration $a = \frac{\Delta v}{\Delta t}$, in m s^{-2} ; deceleration means a and v have opposite signs.
- $1 \text{ m s}^{-1} = 3.6 \text{ km h}^{-1}$: always convert to SI units before calculating.
- Choose a positive direction once, and keep every sign consistent to the end.

2 SECTION 2: EQUATIONS OF MOTION UNDER UNIFORM ACCELERATION (LESSON 93)

In Section 1 we learned to describe motion in one dimension using **displacement** s , **velocity** v and **acceleration** a , and we saw that acceleration is the rate of change of velocity. We now specialise to the most important case in the whole of GCE kinematics: motion in a straight line with **constant acceleration**. This case is important for two reasons. First, it occurs very often in practice: a car pulling away from a junction in Bamenda, a taxi braking at a checkpoint, a stone falling freely near the ground (Section 5). Second, when a is constant, the definitions of Section 1 can be turned into a small set of simple algebraic equations — the famous suvat equations — which allow us to solve almost any straight-line problem at Advanced Level.

2.1 The Uniform Acceleration Assumption and the suvat Variables

DEFINITION Uniformly accelerated motion

A particle moves with **uniform acceleration** (or constant acceleration) when its acceleration a keeps the same value — in magnitude and in sign — throughout the motion. Its velocity then changes by equal amounts in equal intervals of time.

Throughout this chapter we use the five standard letters, collectively called the **suvat variables**:

KEY POINTS The five suvat variables

- s : displacement from the starting point (in metres, m);
- u : initial velocity, i.e. the velocity at time $t = 0$ (in m s^{-1});
- v : final velocity, i.e. the velocity at time t (in m s^{-1});
- a : constant acceleration (in m s^{-2});
- t : time elapsed (in seconds, s).

All five are signed quantities: each may be positive or negative depending on the chosen positive direction.

2.2 First Equation: $v = u + at$

Start from the definition of constant acceleration: acceleration is the change in velocity divided by the time taken,

$$a = \frac{v - u}{t}.$$

Multiplying both sides by t and adding u gives the first equation.

PROPERTY First equation of motion

For motion with constant acceleration,

$$v = u + at$$

This links u , v , a and t ; the variable s is absent. The derivation from the definition of acceleration is examinable.

Intuitively: the velocity gained in time t is at (acceleration $\times$ time), so the final velocity is the initial velocity plus this gain.

2.3 Second and Third Equations: the Average Velocity Idea

Here is the key intuition. If the velocity changes steadily from u to v , then the average velocity over the journey is simply the mean of the two:

$$\bar{v} = \frac{u + v}{2}.$$

But displacement is average velocity multiplied by time, so

$$s = \frac{u + v}{2} t.$$

This is the second equation. To obtain the third, substitute $v = u + at$ into it:

$$s = \frac{u + (u + at)}{2} t = \frac{2u + at}{2} t = ut + \frac{1}{2}at^2.$$

PROPERTY Second and third equations of motion

$$s = \frac{1}{2}(u + v)t$$

(no a)

$$s = ut + \frac{1}{2}at^2$$

(no v)

The term ut is the displacement the particle would have covered at constant velocity u ; the term $\frac{1}{2}at^2$ is the extra displacement due to the acceleration.

2.4 Fourth Equation: Eliminating the Time

Sometimes the time is neither given nor required. From $v = u + at$ we get $t = \frac{v - u}{a}$ (for $a \neq 0$).

Substituting into $s = \frac{1}{2}(u + v)t$:

$$s = \frac{u + v}{2} \cdot \frac{v - u}{a} = \frac{v^2 - u^2}{2a},$$

since $(u + v)(v - u) = v^2 - u^2$. Multiplying by $2a$ and rearranging:

PROPERTY Fourth equation of motion

$$v^2 = u^2 + 2as$$

(no t)

This is the equation to use whenever time does not appear in the problem.

2.5 The Fifth Equation and the Summary

Finally, write $u = v - at$ (rearranging the first equation) and substitute into $s = \frac{1}{2}(u + v)t$:

$$s = \frac{(v - at) + v}{2} t = vt - \frac{1}{2}at^2.$$

PROPERTY The five equations of uniformly accelerated motion

$$v = u + at \quad \text{missing } s$$

$$s = \frac{1}{2}(u + v)t \quad \text{missing } a$$

$$s = ut + \frac{1}{2}at^2 \quad \text{missing } v$$

$$v^2 = u^2 + 2as \quad \text{missing } t$$

$$s = vt - \frac{1}{2}at^2 \quad \text{missing } u$$

Each equation involves exactly four of the five variables; each omits a different one. These formulae are valid only when a is constant.

Equation	Absent variable
$v = u + at$	no s
$s = \frac{1}{2}(u + v)t$	no a
$s = ut + \frac{1}{2}at^2$	no v
$v^2 = u^2 + 2as$	no t
$s = vt - \frac{1}{2}at^2$	no u

The five suvat equations: each one leaves out exactly one variable. To choose an equation, look at which variable is neither given nor wanted.

2.6 Strategy: Choosing the Right Equation

METHOD The four-step suvat strategy

1. List. State the positive direction, then write down the values of the suvat variables that are known, converting all units to metres and seconds.
2. Choose. Identify the unknown you want and the variable that is neither given nor wanted; pick the equation that omits that variable.
3. Substitute. Insert the known values, keeping signs carefully (a deceleration is a negative acceleration).
4. Solve and give the answer with correct units and a sensible number of significant figures.

WARNING Exam tip: direction first

Before substituting anything, always state the positive direction and write down the known values with their signs. Most suvat errors at GCE come from sign mistakes made in the first line, not from algebra.

WORKED EXAMPLE 3 A car accelerating from rest

A car starts from rest at a junction and accelerates uniformly at 2.5 m s^{-2} for 8 s. Find (a) its

velocity after 8 s; (b) the distance covered in this time.

SOLUTION

Solution. Take the direction of motion as positive. Known: $u = 0$, $a = 2.5$, $t = 8$.

(a) Want v ; s is absent, so use $v = u + at$:

$$v = 0 + 2.5 \times 8 = 20 \text{ m s}^{-1}.$$

(b) Want s ; use $s = ut + \frac{1}{2}at^2$:

$$s = 0 \times 8 + \frac{1}{2} \times 2.5 \times 8^2 = 1.25 \times 64 = 80 \text{ m}.$$

Check with $v^2 = u^2 + 2as$: $v^2 = 2 \times 2.5 \times 80 = 400$, so $v = 20 \text{ m s}^{-1}$. ✓

WORKED EXAMPLE 4 A taxi braking to a stop

A taxi travelling at 25 m s^{-1} brakes uniformly and stops in a distance of 50 m. Find (a) the acceleration; (b) the time taken to stop.

SOLUTION

Solution. Positive direction = direction of motion. Known: $u = 25$, $v = 0$, $s = 50$.

(a) Want a ; t is absent, so use $v^2 = u^2 + 2as$:

$$0 = 25^2 + 2a(50) \implies 100a = -625 \implies a = -6.25 \text{ m s}^{-2}.$$

The negative sign confirms a deceleration: the taxi slows at 6.25 m s^{-2} .

(b) Want t ; use $v = u + at$:

$$0 = 25 - 6.25t \implies t = \frac{25}{6.25} = 4 \text{ s}.$$

WARNING Sign errors with deceleration

When a particle slows down while moving in the positive direction, its acceleration is negative. Never write $v^2 = u^2 - 2as$ “by hand” to force a deceleration: keep the standard formula $v^2 = u^2 + 2as$ and let $a < 0$ do the work. Mixing the two conventions is the commonest source of wrong braking distances.

WARNING Using suvat when a is not constant

The five equations are valid only for constant acceleration. If a problem states that acceleration varies (for example a depends on t or on v), suvat does not apply: you must return to calculus, $v = \frac{ds}{dt}$ and $a = \frac{dv}{dt}$. Always check the “uniform acceleration” hypothesis before reaching for a suvat formula.

2.7 Two-Phase and Multi-Phase Journeys

Real journeys rarely have a single acceleration. A typical trip consists of phases: accelerate, cruise at constant velocity, then brake. The method is always the same: **treat each phase separately**, using the final velocity of one phase as the initial velocity of the next.

WORKED EXAMPLE 5 Accelerate, cruise, brake

A bus accelerates uniformly from rest at 1.5 m s^{-2} for 10 s, travels at constant velocity for 20 s, then brakes uniformly to rest in 5 s. Find the total distance travelled.

SOLUTION

Solution. Positive direction = direction of motion.

Phase 1 ($u = 0$, $a = 1.5$, $t = 10$):

$$v_1 = 0 + 1.5 \times 10 = 15 \text{ m s}^{-1}, \quad s_1 = \frac{1}{2}(0 + 15) \times 10 = 75 \text{ m}.$$


Phase 2 (constant velocity 15 m s^{-1} , $t = 20$, so $a = 0$):

$$s_2 = 15 \times 20 = 300 \text{ m}.$$

Phase 3 ($u = 15$, $v = 0$, $t = 5$): use $s = \frac{1}{2}(u + v)t$:

$$s_3 = \frac{1}{2}(15 + 0) \times 5 = 37.5 \text{ m}.$$

Total: $s = 75 + 300 + 37.5 = 412.5 \text{ m}$.



A three-phase journey. The final velocity of each phase becomes the initial velocity of the next; distances add up.

2.8 Two-Particle Problems (Enrichment for GCE)

Some GCE questions involve two particles moving along the same line, starting at the same time. Write a suvat expression for the displacement of each particle as a function of the same time t , then impose the condition of the problem — typically a **meeting condition**: the particles meet when their positions are equal.

WORKED EXAMPLE 6 When does the second car catch the first?

Car A passes a point O at 20 m s^{-1} and continues at constant velocity. At the same instant, car B starts from rest at O and accelerates uniformly at 4 m s^{-2} in the same direction. Find the time at which B catches up with A, and the distance from O at that moment.

SOLUTION

Solution. Measure displacement from O , in the direction of motion, with t the time since B started.

Car A (constant velocity): $s_A = 20t$.

Car B ($u = 0$, $a = 4$): $s_B = \frac{1}{2}(4)t^2 = 2t^2$.

Meeting condition $s_A = s_B$:

$$20t = 2t^2 \implies 2t(t - 10) = 0.$$

$t = 0$ is the start; the catching-up occurs at $t = 10 \text{ s}$, at distance $s = 20 \times 10 = 200 \text{ m}$ from O .

EXERCISE 2 Braking distance

A car travelling at 30 m s^{-1} brakes with uniform deceleration 5 m s^{-2} . Find (a) the braking distance; (b) the time to stop; (c) the distance covered during the first 2 s of braking.

SOLUTION Exercise 1

Positive direction = direction of motion: $u = 30$, $a = -5$.

(a) $v^2 = u^2 + 2as$: $0 = 900 - 10s$, so $s = 90 \text{ m}$.

(b) $v = u + at$: $0 = 30 - 5t$, so $t = 6 \text{ s}$.

(c) $s = ut + \frac{1}{2}at^2 = 30(2) - \frac{1}{2}(5)(2^2) = 60 - 10 = 50 \text{ m}$.

EXERCISE 3 Two phases

A motorcyclist accelerates uniformly from 5 m s^{-1} to 15 m s^{-1} over 100 m , then rides at constant velocity for 30 s . Find (a) the acceleration and the duration of phase 1; (b) the total distance travelled.

SOLUTION Exercise 2

(a) $v^2 = u^2 + 2as$: $225 = 25 + 200a$, so $a = 1 \text{ m s}^{-2}$. Then $v = u + at$: $15 = 5 + t$, so $t = 10 \text{ s}$.

(b) Phase 2: $s_2 = 15 \times 30 = 450 \text{ m}$. Total: $100 + 450 = 550 \text{ m}$.

KEY POINTS Key points of Section 2

- The five suvat equations apply only to straight-line motion with constant acceleration.
- Each equation omits one variable: choose the equation that omits the variable which is neither given nor wanted.
- Always fix a positive direction first; deceleration means $a < 0$.
- Multi-phase journeys: treat each phase separately; the final velocity of one phase is the initial velocity of the next.
- Two-particle problems: same clock for both particles; meeting means equal positions.

In the next section we give these motions a face: we draw and interpret displacement–time and velocity–time graphs, and we shall see that the suvat equations reappear as gradients and areas.

3 SECTION 3: DISPLACEMENT–TIME AND VELOCITY–TIME GRAPHS (LESSONS 94, 95, 96)

In Section 2 we derived the equations of motion for uniform acceleration using algebra and calculus. Those equations are powerful, but they assume constant acceleration. In many real situations — a bus leaving a station in Yaoundé, a ball thrown upwards during a physics practical, a car braking on the highway to Douala — the acceleration changes. Graphs give us a universal language to describe motion, whether acceleration is constant or not. They also provide a visual check on our calculations: the area under a velocity–time graph, for instance, always equals the displacement, no matter how the acceleration varies. In this section we master the two most important graphs in kinematics: displacement–time and velocity–time.

3.1 Displacement–Time Graphs (x – t graphs)


A displacement–time graph plots the position x of a particle (measured from a fixed origin along a straight line) against time t . The horizontal axis represents time, the vertical axis represents displacement. The graph tells us where the particle is at any instant.

PROPERTY Gradient of a Displacement–Time Graph

The gradient (slope) of the tangent to an x – t graph at any point gives the instantaneous velocity v at that time:

$$v = \frac{dx}{dt} = \text{gradient of } x\text{--}t \text{ graph.}$$

- A horizontal line (gradient = 0) means the particle is at rest.
- A straight line with constant positive gradient means constant velocity in the positive direction.
- A straight line with constant negative gradient means constant velocity in the negative direction.
- A curve means the velocity is changing; the steeper the curve, the greater the speed.



Displacement–time graph with three distinct regions: (1) constant positive velocity, (2) at rest, (3) constant negative velocity (return journey). Tangents show the velocity in each region.

EXAMPLE Reading an x – t Graph

The graph above shows a particle that:

1. moves from the origin with constant velocity $v = 2 \text{ m s}^{-1}$ for 3 s (reaching $x = 6 \text{ m}$),
2. remains at rest at $x = 6 \text{ m}$ for 3 s,
3. then returns towards the origin with constant velocity $v = -1 \text{ m s}^{-1}$ for 4 s, ending at $x = 2 \text{ m}$.

Questions:

1. What is the displacement at $t = 5 \text{ s}$? Answer: $x = 6 \text{ m}$ (read directly from graph).
2. What is the velocity at $t = 5 \text{ s}$? Answer: $v = 0$ (horizontal segment).

3. What is the displacement between $t = 2$ s and $t = 8$ s? Answer: $\Delta x = x(8) - x(2) = 4 - 4 = 0$ m.

4. What is the total distance travelled in the first 10 s? Answer: $6 \text{ m} + 0 \text{ m} + 4 \text{ m} = 10 \text{ m}$.

KEY POINTS Average Velocity from an $x-t$ Graph

The average velocity between times t_1 and t_2 is the gradient of the chord joining the points (t_1, x_1) and (t_2, x_2) :

$$v_{\text{avg}} = \frac{x_2 - x_1}{t_2 - t_1} = \frac{\Delta x}{\Delta t}.$$

This is the slope of the straight line connecting the two points on the curve, not the slope of the tangent.

WARNING Common Mistake: Confusing Position with Velocity

On an $x-t$ graph, the height of the curve gives the position x , while the slope gives the velocity v . A common error is to read the value on the vertical axis at a given time and call it the velocity. Always remember: gradient = velocity.

3.2 Velocity–Time Graphs ($v-t$ graphs)

A velocity–time graph plots velocity v against time t . It tells us how fast and in which direction the particle is moving at each instant.

PROPERTY Gradient and Area of a Velocity–Time Graph

For a $v-t$ graph:

1. The gradient of the graph at any point equals the acceleration a :

$$a = \frac{dv}{dt} = \text{gradient of } v-t \text{ graph.}$$

2. The area under the graph between two times t_1 and t_2 equals the displacement Δx during that interval (areas above the t -axis are positive, areas below are negative):

$$\Delta x = \int_{t_1}^{t_2} v \, dt = \text{signed area under } v-t \text{ graph.}$$


KEY POINTS**Standard Shapes of v - t Graphs for Uniform Acceleration**

- Constant velocity: horizontal line ($a = 0$). Area = rectangle $\Rightarrow \Delta x = v \times \Delta t$.
- Uniform acceleration from rest: straight line through origin with positive slope. Area = triangle $\Rightarrow \Delta x = \frac{1}{2}v_{\text{final}} \times t$.
- Uniform deceleration to rest: straight line sloping down to the t -axis. Area = triangle.
- Combined journey (accelerate – cruise – brake): trapezoid. Area = area of rectangle + two triangles.

METHOD**Computing Displacement from a v - t Graph**

To find the displacement between t_1 and t_2 :

1. Identify the geometric shapes formed by the graph and the t -axis (triangles, rectangles, trapezia).
2. Compute the area of each shape using standard formulae:
 - Rectangle: base $\times$ height.
 - Triangle: $\frac{1}{2} \times$ base $\times$ height.
 - Trapezium: $\frac{1}{2} \times$ (sum of parallel sides) $\times$ height.
3. Assign a positive sign to areas above the t -axis, negative to areas below.
4. Sum the signed areas to get the displacement Δx .
5. For distance travelled, sum the absolute values of the areas (ignore signs).



Velocity-time graph of a bus journey between two stops: uniform acceleration (0–5 s), constant velocity (5–20 s), uniform deceleration (20–30 s). The total shaded area equals the distance between the stops.

EXAMPLE Bus Journey – Graphical Analysis

A bus starts from rest at a bus stop, accelerates uniformly at 3 m s^{-2} for 5 s, travels at constant velocity for 15 s, then decelerates uniformly to rest at the next stop in 10 s.

1. Sketch the v – t graph (as above).
2. Find the maximum velocity reached. $v_{\text{max}} = at = 3 \times 5 = 15 \text{ m s}^{-1}$.
3. Find the distance between the two stops (total area under graph).

$$\Delta x_1 = \frac{1}{2} \times 5 \times 15 = 37.5 \text{ m (triangle)}$$

$$\Delta x_2 = 15 \times 15 = 225 \text{ m (rectangle)}$$

$$\Delta x_3 = \frac{1}{2} \times 10 \times 15 = 75 \text{ m (triangle)}$$

$$\text{Total distance} = 37.5 + 225 + 75 = 337.5 \text{ m.}$$


4. Find the deceleration during braking. $a = \frac{\Delta v}{\Delta t} = \frac{0-15}{10} = -1.5 \text{ m s}^{-2}$.

DID YOU KNOW? Recovering the suvat Equations Graphically

The area of the trapezium in a v – t graph for uniform acceleration gives $\Delta x = \frac{1}{2}(u + v)t$. Since $v = u + at$, substituting yields $\Delta x = ut + \frac{1}{2}at^2$. The other suvat equations follow similarly. This shows that the algebraic equations are just the geometric areas in disguise!

3.3 Displacement vs Distance from v - t Graphs

When the velocity changes sign (the graph crosses the t -axis), the particle reverses direction. The **displacement** is the algebraic sum of the areas (positive above axis, negative below). The **distance travelled** is the sum of the absolute areas.



Velocity-time graph crossing the time axis at $t = 3$ s. The pink region (below axis) contributes negative displacement; the green region (above axis) contributes positive displacement.

EXAMPLE Displacement and Distance from a v - t Graph Crossing the Axis

The graph shows $v = 2t - 6$ for $0 \leq t \leq 10$.

1. Displacement from $t = 0$ to $t = 10$:

$$\Delta x = \int_0^{10} (2t - 6) dt = [t^2 - 6t]_0^{10} = 100 - 60 = 40 \text{ m.}$$

Geometrically: $A_- = -\frac{1}{2} \times 3 \times 6 = -9$ m (triangle below axis), $A_+ = \frac{1}{2} \times 7 \times 14 = 49$ m (triangle above axis). Sum = 40 m.

2. Distance travelled from $t = 0$ to $t = 10$:

$$\text{Distance} = |A_-| + |A_+| = 9 + 49 = 58 \text{ m.}$$

WARNING Common Mistake: Forgetting the Sign of Area Below the Axis

When a v - t graph goes below the t -axis, the area is negative for displacement but positive for distance. Many candidates lose marks by giving the distance as the net area (displacement) or by forgetting to take the absolute value of the negative part. Always ask yourself: Am I asked for displacement or distance?

WARNING Common Mistake: Reading v as Displacement

Another frequent error: at a given time t , reading the value v on the vertical axis and calling it the displacement. Remember: on a v - t graph, the height is velocity, the area is displacement.

3.4 Real-Life Applications and GCE-Style Questions

In the GCE examination, you will often be given a worded description of a journey and asked to:

- sketch the v - t graph (with correct shape, labels, and units on axes),
- calculate the total distance or displacement,
- find the acceleration at a given stage,
- find the average speed or average velocity.

KEY POINTS Exam Tips for Graph Questions

- Always label axes with the quantity and its unit (e.g., v (m s^{-1}), t (s)). Marks are awarded for this.
- Use a sharp pencil and ruler for straight lines; draw curves smoothly.
- Show key coordinates (intercepts, maximum/minimum points) on the graph.
- When computing area, show the geometric breakdown (e.g., $\frac{1}{2} \times 4 \times 10$) — this earns method marks even if the final arithmetic is wrong.
- For average speed: $\frac{\text{total distance}}{\text{total time}}$. For average velocity: $\frac{\text{total displacement}}{\text{total time}}$.

WORKED EXAMPLE 7 GCE-Style Question

A car starts from rest at a traffic light and accelerates uniformly at 2 m s^{-2} for 10 s. It then travels at constant velocity for 20 s before decelerating uniformly to rest in 5 s at the next traffic light.

1. Sketch the velocity-time graph for the whole journey.
2. Calculate the distance between the two traffic lights.
3. Find the deceleration during the final stage.
4. Calculate the average speed of the car for the whole journey.

SOLUTION

Solution:

1. Graph:

- $0 \leq t \leq 10$: $v = 2t$ (straight line from $(0, 0)$ to $(10, 20)$).
- $10 \leq t \leq 30$: $v = 20$ (horizontal line).
- $30 \leq t \leq 35$: $v = -4(t - 30) + 20 = -4t + 140$ (straight line from $(30, 20)$ to $(35, 0)$).

Axes labelled v (m s^{-1}) and t (s).

2. Distance = total area under graph:

$$A_1 = \frac{1}{2} \times 10 \times 20 = 100 \text{ m}$$

$$A_2 = 20 \times 20 = 400 \text{ m}$$

$$A_3 = \frac{1}{2} \times 5 \times 20 = 50 \text{ m}$$

$$\text{Total distance} = 100 + 400 + 50 = 550 \text{ m.}$$

3. Deceleration = gradient of last segment:

$$a = \frac{0 - 20}{35 - 30} = \frac{-20}{5} = -4 \text{ m s}^{-2}.$$

$$4. \text{ Average speed} = \frac{\text{total distance}}{\text{total time}} = \frac{550}{35} = 15.7 \text{ m s}^{-1} \text{ (to 3 s.f.)}.$$

EXERCISE 4 Practice

A particle moves along a straight line. Its velocity–time graph consists of three straight-line segments:

- From $t = 0$ to $t = 4$ s: velocity increases uniformly from 0 to 8 m s^{-1} .
- From $t = 4$ s to $t = 10$ s: velocity decreases uniformly to -4 m s^{-1} .
- From $t = 10$ s to $t = 14$ s: velocity increases uniformly to 0.

1. Sketch the v – t graph.
2. Find the acceleration during each stage.
3. Calculate the displacement of the particle at $t = 14$ s relative to its starting point.
4. Calculate the total distance travelled in the 14 s.

SOLUTION Exercise

- Graph: three straight lines connecting $(0, 0)$, $(4, 8)$, $(10, -4)$, $(14, 0)$.
- Accelerations: Stage 1: $a_1 = \frac{8-0}{4} = 2 \text{ m s}^{-2}$. Stage 2: $a_2 = \frac{-4-8}{6} = -2 \text{ m s}^{-2}$. Stage 3: $a_3 = \frac{0-(-4)}{4} = 1 \text{ m s}^{-2}$.
- Displacement = signed area: $A_1 = \frac{1}{2} \times 4 \times 8 = 16 \text{ m}$ (positive). A_2 : trapezium from $t = 4$ to 10 with heights 8 and -4 . Area $= \frac{1}{2}(8 + (-4)) \times 6 = 12 \text{ m}$ (positive). A_3 : triangle below axis from $t = 10$ to 14 with height -4 . Area $= \frac{1}{2} \times 4 \times (-4) = -8 \text{ m}$. Total displacement $= 16 + 12 - 8 = 20 \text{ m}$.
- Distance $= |16| + |12| + |-8| = 36 \text{ m}$.

4 SECTION 4: ACCELERATION-TIME GRAPHS AND CONNECTING THE THREE GRAPHS (LESSONS 97, 98)

In Section 3 we learned to read motion from **displacement-time** and **velocity-time** graphs: the gradient of an $x-t$ graph gives velocity, and the area under a $v-t$ graph gives displacement. We now complete the picture by adding the third member of the family, the **acceleration-time graph**, and we learn to move freely between all three. This is one of the most frequently examined skills in the GCE Advanced Level mechanics paper: an examiner gives you one graph and asks you to sketch or interpret the other two.

4.1 The Acceleration-Time Graph

DEFINITION Acceleration-time graph

An **acceleration-time graph** (or $a-t$ graph) is a graph of the acceleration of a particle plotted on the vertical axis against time on the horizontal axis. Since $a = \frac{dv}{dt}$, the graph shows how quickly the velocity itself is changing at each instant.

Because acceleration is the gradient of velocity, the $a-t$ graph plays for the $v-t$ graph exactly the role that the $v-t$ graph plays for the $x-t$ graph. Everything we learned in Section 3 transfers directly, one step further down the chain:

Pair of graphs	Going down (differentiate)	Going up (integrate)
$x-t$ and $v-t$	gradient of $x-t$ = velocity	area under $v-t$ = displacement
$v-t$ and $a-t$	gradient of $v-t$ = acceleration	area under $a-t$ = change in velocity

Reading the basic shapes.

- A horizontal line above the time axis means **constant positive acceleration**: velocity increases uniformly.
- A horizontal line below the time axis means constant negative acceleration (deceleration if the particle is moving forward): velocity decreases uniformly.
- The line $a = 0$ (the graph lies on the time axis) means zero acceleration, i.e. the velocity is **constant** — not necessarily zero.
- A **sloping line** means the acceleration itself is changing (non-uniform acceleration); at this level we mostly meet piecewise-constant accelerations.

WARNING Zero acceleration does not mean zero velocity

The most common error at this point is to read “ $a = 0$ ” as “the particle is at rest”. Wrong. Zero acceleration means the velocity is not changing: the particle moves with constant velocity (which may be large!). A bus cruising steadily at 60 km h^{-1} along a straight stretch of the Douala–Yaoundé road has zero acceleration. Only when the v – t graph touches the time axis is the particle instantaneously at rest.

4.2 Area Under the Acceleration–Time Graph

PROPERTY Area under the a – t graph

The (signed) area between the a – t graph and the time axis, between times t_1 and t_2 , equals the change in velocity over that interval:

$$\text{area under } a\text{--}t \text{ graph} = \Delta v = v(t_2) - v(t_1).$$

Areas below the time axis count negatively. The proof of this result (it is the Fundamental Theorem of Calculus applied to $a = \frac{dv}{dt}$) is not examinable, but the result itself must be known and used fluently.

Why this is true (instructive derivation).

For constant acceleration a over a time interval Δt , the defining equation $a = \frac{\Delta v}{\Delta t}$ gives $\Delta v = a \Delta t$, which is precisely the area of the rectangle of height a and width Δt under the graph. If the acceleration varies, split the interval into very small pieces of duration δt ; on each piece the acceleration is nearly constant, so the velocity changes by approximately $a \delta t$, the area of a thin strip. Adding all the strips, the total change of velocity equals the total area. Making $\delta t \rightarrow 0$ turns this approximation into an exact equality: $\Delta v = \int_{t_1}^{t_2} a \, dt$.

WARNING Area gives the change in velocity, not the velocity

The area under an $a-t$ graph is Δv , not v . To obtain the velocity at time t you must add the accumulated area to the initial velocity:

$$v(t) = v(0) + (\text{signed area under the } a-t \text{ graph from } 0 \text{ to } t).$$

Forgetting $v(0)$ is the second most common mistake in this topic. A particle can have zero area under its $a-t$ graph yet be moving fast, simply because it started fast.

4.3 Reconstructing $v-t$ from $a-t$: Accumulating Areas**METHOD** From the $a-t$ graph to the $v-t$ graph

1. Write down the initial velocity $v(0)$ (given in the question).
2. Work phase by phase. On each phase where a is constant, compute the signed area $a \times \Delta t$ and add it to the running velocity: $v_{\text{end}} = v_{\text{start}} + a \Delta t$.
3. Plot the velocity values at each phase boundary, then join them: constant acceleration means the $v-t$ graph is a straight line on each phase (rising if $a > 0$, falling if $a < 0$, horizontal if $a = 0$).
4. Check: the gradient of each segment of your $v-t$ sketch must equal the value of a on that phase.

EXAMPLE Accumulating areas

A particle starts with velocity $v(0) = 2 \text{ m s}^{-1}$. Its acceleration is 3 m s^{-2} for $0 \leq t \leq 4$, then 0 for $4 \leq t \leq 7$, then -2 m s^{-2} for $7 \leq t \leq 12$.

Phase 1: $\Delta v = 3 \times 4 = 12$, so $v(4) = 2 + 12 = 14 \text{ m s}^{-1}$.

Phase 2: $\Delta v = 0$, so $v(7) = 14 \text{ m s}^{-1}$ (constant velocity).

Phase 3: $\Delta v = (-2) \times 5 = -10$, so $v(12) = 14 - 10 = 4 \text{ m s}^{-1}$.

The $v-t$ graph is therefore: straight line from $(0, 2)$ to $(4, 14)$, horizontal from $(4, 14)$ to $(7, 14)$, straight line down from $(7, 14)$ to $(12, 4)$. Note that the velocity stays positive throughout: the particle never stops, it only slows down.

4.4 From $v-t$ to $x-t$: Accumulating Areas Again

Exactly the same philosophy, one level up: the area under the $v-t$ graph is the change in displacement, so

$$x(t) = x(0) + (\text{signed area under the } v-t \text{ graph from } 0 \text{ to } t).$$

There is one new subtlety concerning shape. On a phase of constant acceleration, the v - t graph is a straight line, so the displacement accumulates like the area under a sloping line: the x - t graph is a **curve (parabola)**, not a straight line. Concretely:

- velocity increasing $\Rightarrow$ x - t curve bends **upwards** (concave up, getting steeper);
- velocity decreasing (but positive) $\Rightarrow$ x - t curve bends **downwards** (concave down, flattening);
- constant velocity $\Rightarrow$ x - t graph is a straight line whose gradient is that velocity.

4.5 From x - t to v - t to a - t : Taking Gradients

Going the other way down the chain, we differentiate graphically: at each instant, the velocity is the gradient of the x - t graph, and the acceleration is the gradient of the v - t graph. For the piecewise-linear graphs met at this level:

- Each **straight segment** of the x - t graph has constant gradient, so it produces a **horizontal segment** on the v - t graph.
- A **horizontal segment** of the v - t graph has zero gradient, so it produces $a = 0$ on the a - t graph.
- A **corner** (sudden change of gradient) in the x - t graph produces a **jump** in the v - t graph; a corner in the v - t graph produces a jump in the a - t graph.
- A **curved** part of the x - t graph (parabola) produces a **sloping straight line** on the v - t graph and a **non-zero horizontal line** on the a - t graph.

METHOD The “gradient going down, area going up” rule

To connect the three graphs, remember one sentence:

Going down the chain ($x \rightarrow v \rightarrow a$), take gradients.

Going up the chain ($a \rightarrow v \rightarrow x$), accumulate areas (and add the initial value).

Concretely:

1. Down: value of v at time t = gradient of x - t at t ; value of a at time t = gradient of v - t at t .
2. Up: $v(t) = v(0) + \text{area under } a$ - t ; $x(t) = x(0) + \text{area under } v$ - t .
3. Always mark the phase-change instants on all three graphs before sketching any curve.

4.6 Complete Worked Chain: A Lift Journey

WORKED EXAMPLE 8 The lift journey

A lift (elevator) in a Douala office block starts from rest at the ground floor. Its acceleration is modelled in three phases: $a = 1.2 \text{ m s}^{-2}$ for $0 \leq t \leq 5$ (speeding up), $a = 0$ for $5 \leq t \leq 11$ (steady motion), $a = -1.5 \text{ m s}^{-2}$ for $11 \leq t \leq 15$ (slowing down to a stop). Take $x(0) = 0$. (a) Find the velocity at the end of each phase. (b) Sketch the a - t , v - t and x - t graphs. (c) Find the total height risen.

SOLUTION

(a) Velocities by accumulating areas. The lift starts from rest: $v(0) = 0$.

Phase 1 (0 to 5 s): $\Delta v = 1.2 \times 5 = 6$, so $v(5) = 0 + 6 = 6 \text{ m s}^{-1}$.

Phase 2 (5 to 11 s): $\Delta v = 0 \times 6 = 0$, so $v(11) = 6 \text{ m s}^{-1}$.

Phase 3 (11 to 15 s): $\Delta v = (-1.5) \times 4 = -6$, so $v(15) = 6 - 6 = 0$. The lift stops exactly at $t = 15$ s, as expected.

(b) The three graphs.

- a - t : horizontal segments at 1.2, then 0, then -1.5.
- v - t : straight line from (0, 0) to (5, 6); horizontal from (5, 6) to (11, 6); straight line from (11, 6) down to (15, 0). The gradient of each piece (1.2, 0, -1.5) matches the acceleration.
- x - t : parabola bending upwards from (0, 0) to (5, 15); straight line of gradient 6 from (5, 15) to (11, 51); parabola bending downwards, flattening to a horizontal tangent, from (11, 51) to (15, 63).


(c) Height risen = area under the v - t graph.

Phase 1 (triangle): $\frac{1}{2} \times 5 \times 6 = 15 \text{ m}$, so $x(5) = 15 \text{ m}$.

Phase 2 (rectangle): $6 \times 6 = 36 \text{ m}$, so $x(11) = 15 + 36 = 51 \text{ m}$.

Phase 3 (triangle): $\frac{1}{2} \times 4 \times 6 = 12 \text{ m}$, so $x(15) = 51 + 12 = 63 \text{ m}$.

The lift rises a total of 63 m. Consistency check: the total signed area under the a - t graph is $1.2 \times 5 + 0 - 1.5 \times 4 = 0$, matching the fact that the velocity returns to its initial value 0.



The lift journey on three aligned graphs. The dashed vertical lines mark the phase changes at $t = 5$ and $t = 11$ s. Going down: gradients. Going up: areas (the shaded areas on the a - t graph, $+6$ and -6 , are the velocity changes).

4.7 Consistency Checks Between the Three Graphs

Before handing in any sketch, run through this checklist. Each row is a direct consequence of “gradient going down, area going up”.

Feature on a graph	Consequence on the graph immediately below	Consequence on the graph two steps below
$x-t$ has a maximum or minimum (horizontal tangent)	$v-t$ crosses zero (sign change of v)	$a-t$... (no fixed rule)
$v-t$ has a maximum or minimum	$a-t$ crosses zero (sign change of a)	$x-t$ has an inflection point (change of concavity)
$v-t$ is horizontal	$a-t$ lies on the time axis ($a = 0$)	$x-t$ is a straight line
$a-t$ is positive	$v-t$ is increasing	$x-t$ bends upwards (concave up)
$a-t$ is negative	$v-t$ is decreasing	$x-t$ bends downwards (concave down)
$v-t$ crosses the time axis (changes sign)	particle reverses direction	$x-t$ has a turning point (maximum or minimum)

WARNING Deceleration is not “negative velocity”

A negative acceleration does not mean the particle moves backwards. In the lift example, $a = -1.5 \text{ m s}^{-2}$ during the last phase, yet the lift keeps moving upwards (velocity still positive) — it simply slows down. Negative acceleration only means the velocity is decreasing. The particle reverses direction only when the velocity changes sign.

EXAMPLE Quick consistency test

A $v-t$ graph reaches a maximum of 10 m s^{-1} at $t = 4 \text{ s}$ and crosses the time axis at $t = 9 \text{ s}$. What must the other graphs show? Answer: at $t = 4$ the $a-t$ graph crosses zero (from positive to negative, since the velocity stops increasing and starts decreasing); at $t = 9$ the particle reverses direction, so the $x-t$ graph has a turning point (a maximum) there.

4.8 Enrichment: The Link with Calculus (Beyond the Syllabus)

DID YOU KNOW? A preview of Further Mathematics

Everything in this section is secretly calculus. If $x(t)$ is the displacement, then

$$v = \frac{dx}{dt} = \dot{x}, \quad a = \frac{dv}{dt} = \frac{d^2x}{dt^2} = \ddot{x},$$

and conversely $v(t) = v(0) + \int_0^t a \, ds$ and $x(t) = x(0) + \int_0^t v \, ds$. “Gradient going down” is differentiation; “area going up” is integration. For example, constant acceleration a integrates to $v = u + at$ and again to $x = ut + \frac{1}{2}at^2$ — the very equations of Lesson 93. You will meet this formally in Further Mathematics, where the same ideas handle non-uniform acceleration such as $a = 3t^2 - 2t$.

EXERCISE 5 Reconstructing the graphs

A motorbike taxi (bendskin) in Bamenda starts from rest. Its acceleration is 2 m s^{-2} for $0 \leq t \leq 6$, then -1 m s^{-2} for $6 \leq t \leq 10$, then 0 for $10 \leq t \leq 16$.

1. Find the velocity at $t = 6$, $t = 10$ and $t = 16$.
2. Sketch the three graphs $a-t$, $v-t$ and $x-t$, aligned vertically.
3. Find the total distance travelled in the 16 s.

SOLUTION Exercise 1

1. $v(0) = 0$. Phase 1: $\Delta v = 2 \times 6 = 12$, so $v(6) = 12 \text{ m s}^{-1}$. Phase 2: $\Delta v = (-1) \times 4 = -4$, so $v(10) = 12 - 4 = 8 \text{ m s}^{-1}$. Phase 3: $\Delta v = 0$, so $v(16) = 8 \text{ m s}^{-1}$.
2. $a-t$: horizontal segments at 2, -1 , 0. $v-t$: line from $(0, 0)$ to $(6, 12)$, line down to $(10, 8)$, horizontal to $(16, 8)$. $x-t$: parabola bending up to $(6, 36)$, parabola bending down to $(10, 76)$, straight line of gradient 8 to $(16, 124)$.
3. Distance = area under $v-t$: triangle $\frac{1}{2}(6)(12) = 36$; trapezium $\frac{1}{2}(12 + 8)(4) = 40$; rectangle $8 \times 6 = 48$. Total = $36 + 40 + 48 = 124 \text{ m}$.

KEY POINTS Key points of Section 4

- The $a-t$ graph shows how velocity changes; a horizontal line means constant acceleration, and $a = 0$ means constant velocity (not rest).
- Area under the $a-t$ graph = change in velocity Δv ; always add the initial velocity: $v(t) = v(0) + \Delta v$.
- Gradient going down the chain $x \rightarrow v \rightarrow a$; area going up the chain $a \rightarrow v \rightarrow x$, never forgetting the initial value.

- Constant acceleration phases give: horizontal $a-t$ segments, straight $v-t$ segments, parabolic $x-t$ curves.
- Consistency: a turning point on one graph corresponds to a zero crossing on the graph below it.
- Exam tip: mark every phase-change instant on all three graphs first, then sketch one phase at a time, checking gradients against areas as you go.

5 SECTION 5: FREE FALL AND MOTION UNDER GRAVITY (LESSONS 99, 100)

In Section 4 we saw how the three graphs of motion — displacement–time, velocity–time and acceleration–time — are linked: the gradient of the $x-t$ graph gives velocity, the gradient of the $v-t$ graph gives acceleration, and areas under the $v-t$ and $a-t$ graphs give displacement and change of velocity. We now apply this whole machinery to the most important natural motion of all: the motion of a body moving under **gravity** alone. Every stone dropped from a bridge over the River Wouri, every mango falling from a tree in Bamenda, every ball tossed into the air on a playground in Buea obeys the same simple laws — and these laws are a favourite of the GCE examiners.

5.1 The Concept of Free Fall (Lesson 99)

An observation to start with.

Drop a coin and a crumpled piece of paper from the same height at the same instant. They hit the ground together. Now drop the coin and an uncrumpled sheet of paper: the paper flutters down much later. What changed? Not the mass of the paper — it is the same paper. What changed is the effect of air resistance. This simple classroom experiment reveals a deep truth: when air resistance is removed or neglected, **all bodies fall with the same acceleration**, whatever their mass.

This was Galileo Galilei's great insight in the early seventeenth century, contradicting the ancient belief (due to Aristotle) that heavier bodies fall faster. Legend says Galileo dropped different masses from the Leaning Tower of Pisa; whether or not the story is literally true, his conclusion has been confirmed by countless experiments, most dramatically when astronaut David Scott dropped a hammer and a feather on the airless Moon in 1971 — they struck the ground simultaneously.

DEFINITION Free fall

A body is in **free fall** when the only force acting on it is its weight, so that its only acceleration is the acceleration due to gravity $\vec{g}$, directed vertically downward. In free fall, air resistance and all other forces (such as buoyancy or tension) are neglected. Near the

Earth's surface, the magnitude of $\vec{g}$ is approximately constant.

PROPERTY Acceleration due to gravity

Near the surface of the Earth, every freely falling body has a constant downward acceleration of magnitude

$$g \approx 9.8 \text{ m s}^{-2} \quad (\text{often taken as } 10 \text{ m s}^{-2} \text{ in GCE exercises}).$$

This acceleration is the same for all bodies, whatever their mass, shape or material. Its direction is always vertically downward, even when the body is moving upward.

WARNING A heavier object does NOT fall faster

A very common misconception is that a stone falls faster than a piece of chalk because it is heavier. In free fall, mass does not appear anywhere in the kinematics: the acceleration is g for every body. The only reason everyday objects seem to fall at different rates is air resistance, which we neglect in this chapter. In your answers, never write “the heavier object lands first”.

WARNING Exam tip — which value of g ?

GCE questions frequently state “take $g = 10 \text{ m s}^{-2}$ ”. If the question says so and you use 9.8, you will obtain different numerical answers and lose accuracy marks. If nothing is said, $g = 9.8 \text{ m s}^{-2}$ is the safe choice. Always read the question to the end before starting.

5.2 Applying the suvat Equations to Vertical Motion

Free fall is motion with uniform acceleration, so all the equations of Section 2 apply unchanged. They can be derived from calculus: $a = \frac{dv}{dt} = -g$ (with up positive) $\Rightarrow v = u - gt$; $v = \frac{dx}{dt} \Rightarrow x = ut - \frac{1}{2}gt^2$; eliminating t gives $v^2 = u^2 - 2gx$. The standard set is:

$$v = u + at, \quad s = ut + \frac{1}{2}at^2, \quad v^2 = u^2 + 2as, \quad s = \frac{1}{2}(u + v)t.$$

The only new ingredient is that the acceleration is known in advance: it is g , downward. Because the motion is vertical, we must choose a positive direction and stick to it.

METHOD Vertical suvat checklist

1. Draw a diagram. Sketch the motion, indicate the chosen positive direction (up or down), and mark the initial velocity $\vec{u}$ and the acceleration $\vec{g}$.
2. Fix the positive direction. Decide whether “up” or “down” is positive, and write it clearly on your script, e.g. “take upward as positive”.
3. Set the acceleration. If up is positive, $a = -g$; if down is positive, $a = +g$. This is the single most common source of error.
4. List the known quantities u, v, s, a, t with their correct signs, and identify the unknown.
5. Choose the suvat equation containing the three knowns and the unknown, exactly as in Section 2.
6. Solve and interpret the sign of the answer (a negative velocity with “up positive” means the body is moving downward).

WARNING Sign of g — the classic trap

If you choose upward as positive, then the acceleration is $a = -9.8 \text{ m s}^{-2}$, because gravity points down. Writing $a = +9.8$ while taking up as positive is the most frequent mistake in this topic and it destroys the whole question. Remember: $g = 9.8 \text{ m s}^{-2}$ is a magnitude (always positive); the acceleration is $\pm g$ depending on your chosen direction.

5.3 Body Dropped from Rest

Suppose a stone is released (not thrown) from a height h above the ground. Take **downward as positive**. Then

$$u = 0, \quad a = +g, \quad s = h.$$

From $v^2 = u^2 + 2as$ we get $v^2 = 2gh$, and from $s = ut + \frac{1}{2}at^2$ we get $h = \frac{1}{2}gt^2$. Hence:

PROPERTY Body dropped from rest — proof examinable

For a body dropped from rest at height h :

$$t = \sqrt{\frac{2h}{g}} \quad (\text{time of fall}), \quad v = \sqrt{2gh} \quad (\text{velocity on impact}).$$

Neither formula contains the mass of the body — Galileo’s observation in algebraic form.

EXAMPLE Stone dropped from a bridge

A stone is dropped from a bridge and hits the water 3 s later. Taking $g = 10 \text{ m s}^{-2}$ and down as positive: $h = \frac{1}{2}gt^2 = \frac{1}{2}(10)(3^2) = 45 \text{ m}$, and the impact velocity is $v = gt = 10 \times 3 = 30 \text{ m s}^{-1}$.

5.4 Body Projected Vertically Upwards

Now throw a ball straight up with initial speed u . Take **upward as positive**, so $a = -g$. The ball slows down on the way up, stops instantaneously at the top, then falls back. The displacement s is measured from the launch point (positive upward).

Time to maximum height. At the top, $v = 0$. Using $v = u + at$:

$$0 = u - gt \quad \Rightarrow \quad t_{\text{up}} = \frac{u}{g}.$$

Maximum height. Using $v^2 = u^2 + 2as$ with $v = 0$, $a = -g$:

$$0 = u^2 - 2gH \quad \Rightarrow \quad H = \frac{u^2}{2g}.$$

PROPERTY At the top: $v = 0$ but $a \neq 0$

At the highest point of the motion, the velocity is instantaneously zero, but the acceleration is still g directed downward ($a = -g$ if up is positive). The body is not in equilibrium at the top — gravity never switches off.

WARNING “Acceleration is zero at the top” — wrong!

Many candidates write that at maximum height “ $v = 0$ so $a = 0$ ”. This is false. If the acceleration were zero at the top, the ball would stay there forever! The velocity changes sign at the top precisely because the acceleration is still $-g$ throughout the whole flight.


Total time of flight (back to launch level). The displacement when the ball returns to its launch point is $s = 0$. Using $s = ut + \frac{1}{2}at^2$ with $a = -g$:

$$0 = ut - \frac{1}{2}gt^2 = t(u - \frac{1}{2}gt) \quad \Rightarrow \quad t = 0 \text{ (launch) or } t = \frac{2u}{g}.$$

Hence the total time of flight is $T = \frac{2u}{g}$.

Symmetry of the motion. Consider the ball passing a given height on the way up and on the way down. From $v^2 = u^2 - 2gs$, the value of v^2 at a given height s is fixed, so the speed is the same

at the same height — only the direction differs. It follows that the time to go up to the highest point equals the time to come back down from it, and the ball returns to its launch point with speed u (moving downward).



A ball thrown vertically upward from a balcony in Buea, shown at equal time intervals. The spacing decreases on the way up (deceleration) and increases on the way down (acceleration); the acceleration $\vec{g}$ is downward throughout.

WORKED EXAMPLE 9 Ball thrown upward from a balcony

A ball is thrown vertically upward with speed 14 m s^{-1} from a balcony 15 m above the ground. Take $g = 10 \text{ m s}^{-2}$ and upward as positive. Find: (a) the maximum height above the ground; (b) the total time until the ball hits the ground; (c) the speed of impact.

SOLUTION

(a) Maximum height above launch point:

$$H = \frac{u^2}{2g} = \frac{14^2}{2 \times 10} = \frac{196}{20} = 9.8 \text{ m.}$$

Maximum height above the ground: $15 + 9.8 = 24.8 \text{ m.}$

(b) Total time of flight. Use $s = ut + \frac{1}{2}at^2$ with $s = -15 \text{ m}$ (the ball finishes 15 m below the launch point, and up is positive), $u = 14$, $a = -10$:

$$-15 = 14t - 5t^2 \implies 5t^2 - 14t - 15 = 0.$$

By the quadratic formula:

$$t = \frac{14 \pm \sqrt{196 + 300}}{10} = \frac{14 \pm \sqrt{496}}{10} \approx \frac{14 \pm 22.27}{10}.$$

Taking the positive root: $t \approx 3.63$ s. (The negative root $t \approx -0.83$ s is rejected: time cannot be negative.)


(c) Impact velocity. $v = u + at = 14 - 10(3.63) \approx -22.3 \text{ m s}^{-1}$. The negative sign means the ball is moving downward; the impact speed is about 22.3 m s^{-1} .

5.5 Graphical Representation of Motion under Gravity (Lesson 100)

Everything we learned in Sections 3 and 4 about graphs now pays off. For a ball projected vertically upward with initial speed u , taking **upward as positive**:

$$a(t) = -g, \quad v(t) = u - gt, \quad x(t) = ut - \frac{1}{2}gt^2.$$

- The **acceleration–time graph** is a horizontal straight line at $-g$: the acceleration is constant and downward for the whole flight, including at the top.
- The **velocity–time graph** is a straight line of gradient $-g$, starting at $v = u$ and crossing the time axis at $t = u/g$ (the instant the ball is at the top). The area between the graph and the time axis (counted negative below the axis) gives the displacement.
- The **displacement–time graph** is a parabola opening downward, with maximum $H = u^2/(2g)$ at $t = u/g$. Its gradient at any point is the velocity: positive on the way up, zero at the top, negative on the way down.



The three graphs for a ball projected upward at 10 m s^{-1} with $g = 10 \text{ m s}^{-2}$, upward positive. The v - t line has gradient -10 and crosses the axis at the top of the flight; the a - t line stays at $-g$ even at the highest point. The x - t parabola crosses the time axis at $t = 2$ s (return to launch level).

Note how the three graphs are perfectly consistent with Section 4: the gradient of the parabola at $t = 0$ is u (the initial slope of the x - t curve equals the initial velocity), the gradient of the v - t line equals the constant value of the a - t graph, and the area under the v - t graph from 0 to $2u/g$ is zero — the ball is back where it started.

5.6 Dropped, Thrown Down, Thrown Up: a Comparison

A classic GCE question compares three bodies released from the same height h : one dropped, one thrown downward with speed u_0 , one thrown upward with speed u_0 . Which lands first? Which lands fastest?

WORKED EXAMPLE 10 Three balls from the same height

From a point 20 m above the ground, ball A is dropped, ball B is thrown vertically downward at 5 m s^{-1} , and ball C is thrown vertically upward at 5 m s^{-1} . Take $g = 10 \text{ m s}^{-2}$. Find the time of flight and impact speed of each ball.

SOLUTION

Take downward as positive, so $a = +10$ and $s = +20$ for all three balls.

Ball A (dropped): $u = 0$.

$$t_A = \sqrt{\frac{2h}{g}} = \sqrt{\frac{2 \times 20}{10}} = 2 \text{ s}, \quad v_A = \sqrt{2gh} = \sqrt{400} = 20 \text{ m s}^{-1}.$$

Ball B (thrown down): $u = +5$. From $s = ut + \frac{1}{2}at^2$:

$$20 = 5t + 5t^2 \implies t^2 + t - 4 = 0 \implies t_B = \frac{-1 + \sqrt{17}}{2} \approx 1.56 \text{ s}.$$

$$v_B^2 = u^2 + 2gh = 25 + 400 = 425 \implies v_B \approx 20.6 \text{ m s}^{-1}.$$

Ball C (thrown up): $u = -5$ (upward is negative here). The same equation gives

$$20 = -5t + 5t^2 \implies t^2 - t - 4 = 0 \implies t_C = \frac{1 + \sqrt{17}}{2} \approx 2.56 \text{ s}.$$

$$v_C^2 = u^2 + 2gh = 25 + 400 = 425 \implies v_C \approx 20.6 \text{ m s}^{-1}.$$

Comparison. B lands first (1.56 s), then A (2 s), then C (2.56 s). But B and C hit the ground with the same speed (20.6 m s^{-1}): by symmetry, C passes its launch point moving downward at exactly 5 m s^{-1} — the same state as B at launch — so from that instant on, B and C have identical motion. Note also $t_C - t_B = 1 \text{ s} = \frac{2u_0}{g}$, exactly the time C spends going up and coming back to the launch level.

KEY POINTS Key points of Section 5

- Free fall: the only acceleration is $g \approx 9.8 \text{ m s}^{-2}$, always directed vertically downward; it is the same for all bodies regardless of mass.
- Fix a positive direction first: up positive $\implies a = -g$; down positive $\implies a = +g$.

- Dropped from rest: $t = \sqrt{2h/g}$, $v = \sqrt{2gh}$.
- Projected upward: $t_{\text{up}} = u/g$, $H = u^2/(2g)$; at the top $v = 0$ but $a = -g \neq 0$; total flight time back to launch level is $2u/g$.
- Up–down symmetry: same speed at the same height; time up = time down.
- Graphs (up positive): $x-t$ is a downward parabola, $v-t$ a line of gradient $-g$, $a-t$ a horizontal line at $-g$.
- Check the question: use $g = 10 \text{ m s}^{-2}$ if told to.

EXERCISE 6 Practice — free fall

Take $g = 10 \text{ m s}^{-2}$ unless stated otherwise.

1. A mango falls from a branch 5 m above the ground. Find the time of fall and the speed on impact.
2. A stone is thrown vertically upward at 20 m s^{-1} from ground level. Find (a) the maximum height, (b) the total time of flight, (c) its velocity after 3 s, and interpret the sign.
3. A ball is dropped from a tower and takes 4 s to reach the ground. Find the height of the tower and the impact speed.
4. From the top of a 45 m cliff, a stone is thrown vertically downward at 10 m s^{-1} . Find the time to reach the foot of the cliff and the impact speed.
5. A ball is projected vertically upward from the ground with speed u . Show that the times at which it is at height h (with $h < u^2/2g$) are the roots of $gt^2 - 2ut + 2h = 0$, and deduce that the sum of these two times is the total time of flight.

SOLUTION Exercise 5

1. $t = \sqrt{2h/g} = \sqrt{10/10} = 1 \text{ s}$; $v = \sqrt{2gh} = \sqrt{100} = 10 \text{ m s}^{-1}$.
2. Up positive, $a = -10$. (a) $H = u^2/2g = 400/20 = 20 \text{ m}$. (b) $T = 2u/g = 4 \text{ s}$. (c) $v = u - gt = 20 - 30 = -10 \text{ m s}^{-1}$: the stone is moving downward at 10 m s^{-1} (it is on its way back down).
3. $h = \frac{1}{2}gt^2 = \frac{1}{2}(10)(16) = 80 \text{ m}$; $v = gt = 40 \text{ m s}^{-1}$.
4. Down positive: $45 = 10t + 5t^2 \Rightarrow t^2 + 2t - 9 = 0 \Rightarrow t = \frac{-2 + \sqrt{40}}{2} \approx 2.16 \text{ s}$. $v^2 = 100 + 2(10)(45) = 1000 \Rightarrow v \approx 31.6 \text{ m s}^{-1}$.
5. With up positive, $h = ut - \frac{1}{2}gt^2$, i.e. $gt^2 - 2ut + 2h = 0$ after multiplying by 2 and rearranging. The two roots t_1, t_2 are the times of passing height h (up and down). By Viète's formula,

$t_1 + t_2 = \frac{2u}{g}$, which is exactly the total time of flight — a beautiful confirmation of the up-down symmetry.

DID YOU KNOW? Did you know?

The value of g is not exactly the same everywhere on Earth: it is slightly larger at the poles (about 9.83 m s^{-2}) and slightly smaller at the equator (about 9.78 m s^{-2}), because the Earth is not a perfect sphere and because of its rotation. Mount Cameroon, rising about 4100 m above sea level, experiences a value of g marginally smaller than at the coast of Limbe below it — the difference is tiny, but modern instruments called gravimeters can detect it, and geophysicists use such variations to prospect for oil and minerals.

6 SECTION 6: REAL-LIFE APPLICATIONS AND SYNTHESIS (LESSONS 94 & 101, CONSOLIDATION)

In Section 5 we studied free fall and motion under gravity: we saw that every object near the Earth's surface, when air resistance is neglected, accelerates downward at $g \approx 9.81 \text{ m s}^{-2}$, and we learned to read this motion on displacement–time and velocity–time graphs. We now close the chapter by putting all of our kinematics to work on real situations: stopping a car on the Douala–Yaoundé highway, estimating the height of a building with a stopwatch, predicting where two moving objects meet, and reading a full journey from a verbal description. This is exactly the style of the long, multi-part questions set at the GCE Advanced Level.

6.1 Road Safety: Thinking Distance, Braking Distance, Stopping Distance

Imagine you are driving at a steady speed v when a goat suddenly crosses the road. Two things happen before the car stops. First, there is a reaction time t_r (typically 0.5 to 1 s) during which you see the danger, decide to brake, and press the pedal. During this time the car is still moving at full speed v . Second, once the brakes bite, the car decelerates uniformly with acceleration a (negative) until it stops.

DEFINITION Thinking, braking and stopping distances

The **thinking distance** is the distance travelled at constant speed during the reaction time: $d_{\text{think}} = vt_r$. The **braking distance** is the distance travelled while decelerating uniformly from speed v to rest. The **stopping distance** is the sum:

$$d_{\text{stop}} = d_{\text{think}} + d_{\text{brake}}.$$

PROPERTY Stopping–distance formula


For a vehicle moving at speed v , with reaction time t_r and braking deceleration of magnitude $|a|$,

$$d_{\text{stop}} = vt_r + \frac{v^2}{2|a|}.$$

In particular the braking distance grows with the square of the speed: doubling the speed multiplies the braking distance by four.

Derivation (examinable). During the reaction time the velocity is constant, so $d_{\text{think}} = vt_r$ immediately. During braking the acceleration is constant, so we may use $v_f^2 = v_i^2 + 2as$ with final velocity $v_f = 0$, initial velocity $v_i = v$, and acceleration $a = -|a|$:

$$0 = v^2 - 2|a|s \implies s = \frac{v^2}{2|a|}. \quad \blacksquare$$




Stopping distance = thinking distance + braking distance.

EXAMPLE Stopping distances on the Douala–Yaoundé highway

Take a reaction time $t_r = 0.7$ s and a braking deceleration $|a| = 6 \text{ m s}^{-2}$ (good tyres, dry road). Convert speeds: $50 \text{ km/h} = 13.9 \text{ m s}^{-1}$, $80 \text{ km/h} = 22.2 \text{ m s}^{-1}$, $110 \text{ km/h} = 30.6 \text{ m s}^{-1}$.

Speed	Thinking vt_r	Braking $v^2/(2 a)$	Stopping
50 km/h	9.7 m	16.1 m	25.8 m
80 km/h	15.6 m	41.1 m	56.7 m
110 km/h	21.4 m	78.0 m	99.4 m

Going from 50 to 100 km/h (doubling the speed) would multiply the braking part by 4; here, from 50 to 110 km/h the braking distance is multiplied by about 4.8. Speed kills through the square in the formula.



Braking distance at 50, 80 and 110 km/h for $|a| = 6 \text{ ms}^{-2}$.

WARNING Forgetting the thinking distance

A very common GCE error is to compute only $v^2/(2|a|)$ and call it the stopping distance. Unless the question says the driver reacts instantly, you must add $v t_r$. Read the question twice and underline “reaction time”.

6.2 Falling Objects in Daily Life: Measuring Heights with a Stopwatch

Drop a stone into a well and time its fall. Since the stone starts from rest and falls freely, $h = \frac{1}{2}gt^2$, so measuring t gives the depth h . This is a genuine field technique.

WORKED EXAMPLE 11 Depth of a well

A stone dropped from rest into a well is heard to splash after 2.4 s. Estimate the depth of the well, taking $g = 9.81 \text{ m s}^{-2}$ and neglecting the time sound takes to come back up.

SOLUTION

Solution. The stone starts from rest ($u = 0$) with acceleration g :

$$h = \frac{1}{2}gt^2 = \frac{1}{2}(9.81)(2.4)^2 = \frac{1}{2}(9.81)(5.76) \approx 28.3 \text{ m.}$$

The well is about 28 m deep. (Including the sound travel time would reduce the answer slightly.)

The same idea estimates the height of a storey building: an object taking 1.0 s to fall covers $\frac{1}{2}(9.81)(1)^2 \approx 4.9 \text{ m}$, roughly one and a half storeys.

6.3 Vertical Motion in Sport and Construction

A ball kicked vertically upward, or a spanner slipped from scaffolding, is motion under uniform acceleration $-g$ (taking upward as positive). The full toolkit of Section 5 applies: time to the top is u/g , maximum height is $u^2/(2g)$, and the speed of return to the launch level equals the launch speed.

WORKED EXAMPLE 12 A tool dropped from scaffolding

A mason drops a hammer from scaffolding 15 m above the ground. A colleague stands below. (a) How long does the hammer take to reach the ground? (b) With what speed does it arrive? (c) The colleague needs 0.8 s to step aside after a warning shout; is a shout at the moment of release sufficient? Take $g = 9.81 \text{ m s}^{-2}$.

SOLUTION

Solution. Take downward positive, $u = 0$, $a = g$, $s = 15 \text{ m}$. (a) $s = \frac{1}{2}gt^2 \Rightarrow t = \sqrt{2s/g} = \sqrt{30/9.81} \approx 1.75 \text{ s}$. (b) $v = gt = 9.81 \times 1.75 \approx 17.2 \text{ m s}^{-1}$ (about 62 km/h!). (c) $1.75 \text{ s} > 0.8 \text{ s}$, so the colleague has just under a second to spare — but a shout even 0.5 s late would leave no safety margin. This is why hard hats and exclusion zones matter on construction sites in Douala and Yaoundé.

6.4 Two-Particle Problems: The Meeting Condition

Many GCE questions involve two objects moving together: one dropped from a height, another projected upward from the ground. The whole secret is organisation.

METHOD Solving meeting problems

1. Choose one origin and one positive direction for both particles.
2. Start one clock: $t = 0$ at the same instant for both particles (if one starts later, shift its time, e.g. replace t by $t - 2$).
3. Write the position of each particle: $x_1(t)$ and $x_2(t)$, using $x = x_0 + ut + \frac{1}{2}at^2$.
4. The meeting condition is $x_1(t) = x_2(t)$. Solve for t , then substitute back to find where they meet.
5. Check that the answer is physically sensible (positive time, position within the allowed region).

WARNING Different origins or different clocks

The classic trap is to measure the dropped stone's position from the top and the thrown ball's position from the ground, then equate them — this compares distances measured from different origins and gives nonsense. Similarly, using $t = 0$ at the drop for one particle and at the throw for the other corrupts everything. One origin, one clock.

WORKED EXAMPLE 13 Drop and throw

A stone is dropped from rest from the top of a tower 45 m high. At the same instant, a ball is thrown vertically upward from the foot of the tower with speed 30 m s^{-1} . Taking $g = 10 \text{ m s}^{-2}$, find when and where they meet.

SOLUTION


Solution. Origin at the ground, upward positive, $t = 0$ at the instant of release. Stone: $x_0 = 45, u = 0, a = -10$:

$$x_1(t) = 45 - 5t^2.$$

Ball: $x_0 = 0, u = 30, a = -10$:

$$x_2(t) = 30t - 5t^2.$$

Meeting: $x_1 = x_2 \Rightarrow 45 - 5t^2 = 30t - 5t^2 \Rightarrow 45 = 30t \Rightarrow t = 1.5 \text{ s}$. Position: $x_2(1.5) = 30(1.5) - 5(1.5)^2 = 45 - 11.25 = 33.75 \text{ m}$ above the ground. Check: the ball is still rising at $t = 1.5 \text{ s}$ (its top is at $t = u/g = 3 \text{ s}$), and the stone has fallen only $5(1.5)^2 = 11.25 \text{ m}$: consistent.



The drop-and-throw meeting problem: one origin (ground), one clock.

6.5 Graphical Synthesis: From a Verbal Description to All Three Graphs

Long GCE questions often describe a journey in words and ask for the displacement–time, velocity–time and acceleration–time graphs, plus numerical answers. The strategy: cut the journey into phases of constant acceleration; compute the velocity at the end of each phase; remember

that the area under the v - t graph gives displacement, and the gradient of the v - t graph gives acceleration.

WORKED EXAMPLE 14 A complete journey

A bus starts from rest at a stop, accelerates uniformly at 1.5 m s^{-2} for 8 s, travels at constant velocity for 20 s, then brakes uniformly to rest in 6 s. Find (a) the maximum velocity, (b) the total distance travelled, (c) the deceleration during braking, and describe the three graphs.

SOLUTION

Solution. (a) $v_{\max} = u + at = 0 + 1.5 \times 8 = 12 \text{ m s}^{-1}$. (b) Use areas under the v - t graph (triangle + rectangle + triangle):

$$s = \frac{1}{2}(8)(12) + (20)(12) + \frac{1}{2}(6)(12) = 48 + 240 + 36 = 324 \text{ m}.$$

(c) $a = \frac{0 - 12}{6} = -2 \text{ m s}^{-2}$, a deceleration of 2 m s^{-2} . Graphs: the v - t graph is a trapezium rising from 0 to 12 over $[0, 8]$, flat over $[8, 28]$, falling to 0 over $[28, 34]$. The a - t graph is $+1.5$, then 0, then -2 m s^{-2} on the same intervals. The x - t graph is a curve bending upward (parabola) for 8 s, then a straight line of slope 12, then a curve flattening to a horizontal tangent at $t = 34$ s.

6.6 Chapter Synthesis and GCE Strategy

KEY POINTS Chapter summary map

- Language: displacement s , velocity $v = \dot{s}$, acceleration $a = \dot{v}$; scalars vs vectors.
- Uniform acceleration: $v = u + at$, $s = ut + \frac{1}{2}at^2$, $s = \frac{1}{2}(u + v)t$, $v^2 = u^2 + 2as$.
- Graphs: gradient of x - t = velocity; gradient of v - t = acceleration; area under v - t = displacement; area under a - t = change of velocity.
- Gravity: $a = -g \approx -9.81 \text{ m s}^{-2}$ (upward positive); time to top u/g ; max height $u^2/(2g)$.
- Applications: stopping distance $vt_r + v^2/(2|a|)$; meeting problems via $x_1(t) = x_2(t)$ with one origin and one clock.

Common GCE question patterns. (i) A two-phase journey with graph sketching and area calculations. (ii) A vertical projection with “greatest height” and “time of flight” parts. (iii) Two particles (drop-and-throw, or overtaking) with a meeting condition. (iv) A stopping-distance question with reaction time. In all of these, later parts reuse earlier results.

WARNING Exam tip: keep intermediate values exact

In long questions, part (b) almost always uses the answer of part (a). Keep intermediate values exact (fractions, surds, or unrounded decimals in your calculator memory) and only round at the final line. Rounding $v_{\max} = 12$ early is safe, but rounding $\sqrt{30/9.81}$ to 1.7 before squaring it again can cost accuracy marks.

DID YOU KNOW? Why real falling objects do not accelerate forever

Our model ignores air resistance. In reality, air drag grows with speed until it balances weight; the object then falls at a constant **terminal velocity**. A skydiver in free fall reaches roughly 190–200 km/h before opening the parachute; a raindrop only a few metres per second — which is why rain does not hurt us. At the GCE level you only need the concept: drag explains why the uniform-acceleration model is an approximation, excellent for dense objects over short falls.

EXERCISE 7 Consolidation

1. A car travels at 90 km/h on a wet road where the braking deceleration is only 4 m s^{-2} ; the driver's reaction time is 0.9 s. Find the stopping distance.
2. A mango falls from a tree and takes 1.1 s to reach the ground. How high was it? With what speed does it land? ($g = 9.81 \text{ m s}^{-2}$)
3. A ball is thrown vertically upward at 20 m s^{-1} from the ground at the same instant as a stone is dropped from a 60 m tower beside it. Taking $g = 10 \text{ m s}^{-2}$, find when and where they meet.
4. A lorry accelerates from rest at 0.8 m s^{-2} for 15 s, keeps constant speed for 40 s, then decelerates uniformly to rest over 100 m. Find the total distance and the total time.

SOLUTION Exercise 1

1. $v = 90/3.6 = 25 \text{ m s}^{-1}$. Thinking: $25 \times 0.9 = 22.5 \text{ m}$. Braking: $25^2/(2 \times 4) = 78.1 \text{ m}$. Stopping $\approx 100.6 \text{ m}$. 2. $h = \frac{1}{2}(9.81)(1.1)^2 \approx 5.9 \text{ m}$; $v = 9.81 \times 1.1 \approx 10.8 \text{ m s}^{-1}$. 3. $x_1 = 60 - 5t^2$, $x_2 = 20t - 5t^2$; $x_1 = x_2 \Rightarrow t = 3 \text{ s}$, $x = 20(3) - 5(9) = 15 \text{ m}$ above the ground. 4. $v = 0.8 \times 15 = 12 \text{ m s}^{-1}$. Distances: $\frac{1}{2}(15)(12) = 90$; $40 \times 12 = 480$; braking 100 m given. Total = 670 m. Braking time: $100 = \frac{1}{2}(12 + 0)t \Rightarrow t = 16.7 \text{ s}$. Total time $\approx 71.7 \text{ s}$.

7 INTEGRATION ACTIVITY

8 REAL-LIFE APPLICATIONS AND SYNTHESIS

8.1 Aims of the Activity

This integration activity brings together everything you have learnt in this chapter: the **suvat equations**, the interpretation and construction of **displacement–time**, **velocity–time** and **acceleration–time** graphs, and **free fall** under gravity. Working in groups, you will act as road-safety engineers studying a real emergency stop in Bamenda, and you will communicate your findings on a poster, exactly as an engineer presents a technical report.

By the end of the activity you should be able to:

- convert speeds between km h^{-1} and m s^{-1} fluently;
- decompose a stopping distance into **thinking distance** and **braking distance**;
- use $v^2 = u^2 + 2as$ and the area under a velocity–time graph to find distances, and check that both methods agree;
- sketch coherent $v-t$, $x-t$ and $a-t$ graphs for the same motion;
- model a falling object with $s = \frac{1}{2}gt^2$ and combine it with horizontal motion at constant velocity;
- compare stopping distances at different speeds and write a justified safety recommendation.

8.2 Situation

The Bend–Skin Safety Report.

In Bamenda, bend-skins (motorbike taxis) are one of the most common means of transport between Commercial Avenue and Nkwen. The Students' Road Safety Club of your school has been asked by the local council to prepare a short technical report on emergency braking.

A bend-skin is travelling at a steady 60 km h^{-1} along a straight, level, tarred road when a goat suddenly crosses 40 m ahead. The rider's **reaction time** (the time between seeing the danger and the brakes starting to act) is 0.8 s . Once applied, the brakes produce a constant deceleration of 4 m s^{-2} until the bike stops. Take $g = 10 \text{ m s}^{-2}$.

During the braking, the rider's phone slips from his pocket at a height of 1.2 m above the ground. The phone's vertical motion may be treated as free fall from rest vertically, while the bike continues horizontally.

Your group must produce a poster containing: the numerical results, the three motion graphs ($v-t$, $x-t$, $a-t$), and a one-paragraph safety recommendation comparing stopping distances at 40 , 60 and 80 km h^{-1} .

8.3 Tasks

1. **Unit conversion.** Express 60 km h^{-1} in m s^{-1} . Show the conversion factor you use.
2. **Thinking distance.** Compute the distance travelled during the reaction time (before braking). Why is the speed constant during this phase?
3. **Braking distance and time.** Using a suitable suvat equation, compute the braking distance and the braking time. Hence find the **total stopping distance**. Does the bike stop before reaching the goat 40 m ahead?
4. **Velocity–time graph.** Draw the velocity–time graph of the whole emergency stop (reaction phase + braking phase). Compute the total stopping distance as the area under the graph and verify that it agrees with Task 3.
5. **The other two graphs.** On the same time axis, sketch the displacement–time graph and the acceleration–time graph. Indicate clearly the gradient of the x – t curve at $t = 0$ and at the stopping instant, and the value of a in each phase.
6. **The falling phone.** Compute the time the phone takes to fall 1.2 m vertically. Assuming the phone slips at the instant braking begins, compute the horizontal distance travelled by the bike during the fall. (Bonus: what horizontal distance does the phone cover if it keeps the horizontal velocity it had when it slipped?)
7. **Speed comparison.** Repeat the stopping–distance calculation for 40 km h^{-1} and 80 km h^{-1} (same reaction time, same deceleration). Present the three results in a table.
8. **Recommendation.** In at most six lines, write a safety recommendation for bend–skin riders, using your table to justify a speed limit near schools and markets.

8.4 Model Answer

Task 1. Unit conversion. Since $1 \text{ km h}^{-1} = \frac{1000}{3600} \text{ m s}^{-1} = \frac{1}{3.6} \text{ m s}^{-1}$,

$$u = \frac{60}{3.6} = \frac{50}{3} \approx 16.7 \text{ m s}^{-1}.$$

Task 2. Thinking distance. During the reaction time the brakes are not yet applied, so the velocity is constant (no horizontal force acts on the bike):

$$s_{\text{think}} = u t_r = \frac{50}{3} \times 0.8 = \frac{40}{3} \approx 13.3 \text{ m}.$$

Task 3. Braking distance and time. Braking phase: $u = \frac{50}{3} \text{ m s}^{-1}$, $v = 0$, $a = -4 \text{ m s}^{-2}$. From $v^2 = u^2 + 2as$:

$$0 = \left(\frac{50}{3}\right)^2 + 2(-4)s_{\text{brake}} \implies s_{\text{brake}} = \frac{2500/9}{8} = \frac{625}{18} \approx 34.7 \text{ m}.$$

Braking time from $v = u + at$:


$$t_b = \frac{u}{|a|} = \frac{50/3}{4} = \frac{25}{6} \approx 4.17 \text{ s}.$$

Total stopping distance:

$$s_{\text{stop}} = \frac{40}{3} + \frac{625}{18} = \frac{865}{18} \approx 48.1 \text{ m}$$

Since $48.1 > 40$, the bike hits the goat: it cannot stop within 40 m.

Task 4. Velocity–time graph and area check. The v – t graph consists of a horizontal segment at $v = 50/3 \text{ m s}^{-1}$ from $t = 0$ to $t = 0.8 \text{ s}$, followed by a straight line of gradient -4 m s^{-2} reaching $v = 0$ at $t = 0.8 + 25/6 = 149/30 \approx 4.97 \text{ s}$.




Velocity–time graph for the emergency stop.


Area under the graph = rectangle + triangle:

$$A = \underbrace{\left(\frac{50}{3} \times 0.8\right)}_{\text{rectangle}} + \underbrace{\frac{1}{2} \left(\frac{50}{3}\right) \left(\frac{25}{6}\right)}_{\text{triangle}} = \frac{40}{3} + \frac{625}{18} = \frac{865}{18} \approx 48.1 \text{ m.} \quad \checkmark$$

Task 5. Displacement–time and acceleration–time graphs.



Displacement–time graph. Gradient at $t = 0$ is $u = 16.7 \text{ m s}^{-1}$; gradient at stop is 0.



Acceleration–time graph. $a = 0$ during reaction, $a = -4 \text{ m s}^{-2}$ during braking, $a = 0$ after stop.

Task 6. The falling phone. Vertical free fall from rest: $h = \frac{1}{2}gt^2$.

$$t_{\text{fall}} = \sqrt{\frac{2h}{g}} = \sqrt{\frac{2 \times 1.2}{10}} = \sqrt{0.24} \approx 0.49 \text{ s.}$$

During this time the bike is still braking (since $0.49 < 4.17$). Distance travelled by bike in 0.49 s after brakes applied:

$$s_{\text{bike}} = ut - \frac{1}{2}at^2 = \frac{50}{3}(0.49) - 2(0.24) \approx 8.17 - 0.48 = 7.69 \text{ m.}$$

(Bonus: the phone retains the horizontal velocity $u = 50/3 \text{ m s}^{-1}$ at the moment it slips, so its horizontal distance is $s_{\text{phone}} = u t_{\text{fall}} = \frac{50}{3} \times 0.49 \approx 8.17 \text{ m.}$)

Task 7. Speed comparison table.

Speed	Thinking distance	Braking distance	Total stopping distance
40 km h ⁻¹ (11.1 m s ⁻¹)	8.9 m	15.4 m	24.3 m
60 km h ⁻¹ (16.7 m s ⁻¹)	13.3 m	34.7 m	48.1 m
80 km h ⁻¹ (22.2 m s ⁻¹)	17.8 m	61.7 m	79.5 m

Task 8. Safety recommendation. “Doubling the speed from 40 to 80 km h⁻¹ more than triples the stopping distance (24.3 m → 79.5 m), because braking distance grows like u^2 . At 60 km h⁻¹ a bend-skin needs about 48 m to stop — more than the length of a football pitch’s half. We recommend a 40 km h⁻¹ limit near schools and markets, and that riders keep at least 50 m of clear road ahead.”

WARNING Marking Points

- Do not forget to convert km h⁻¹ to m s⁻¹ before using suvat equations.
- Do not mix the reaction phase (constant velocity, $a = 0$) with the braking phase (uniform deceleration, $a = -4 \text{ m s}^{-2}$).
- The phone’s fall time depends only on the vertical height and g , not on the bike’s speed.
- When sketching graphs, ensure the time axis is consistent across v - t , x - t and a - t plots.
- The gradient of the x - t graph gives velocity; the gradient of the v - t graph gives acceleration; the area under the v - t graph gives displacement.

9 METHODS AND WORKED EXAMPLES

10 KINEMATICS OF PARTICLES MOVING IN A STRAIGHT LINE

10.1 Concept of Motion in One Dimension

DEFINITION Displacement, velocity and acceleration

A particle moves along a straight line. Choose an origin O and a positive direction.

- Displacement s (or x) is the signed distance from O ; unit: metre (m).
- Velocity v is the rate of change of displacement: $v = \frac{ds}{dt}$; unit: m s^{-1} .
- Acceleration a is the rate of change of velocity: $a = \frac{dv}{dt} = \frac{d^2s}{dt^2}$; unit: m s^{-2} .
- Speed is the magnitude of velocity: $|v|$.
- Retardation (deceleration) means acceleration opposite to the velocity; its magnitude is positive.

PROPERTY Uniform acceleration

If acceleration a is constant, the motion is uniformly accelerated. The five quantities s (displacement from start), u (initial velocity), v (final velocity), a , t (time elapsed) are linked by the **suvat equations**:

$$v = u + at \quad (1)$$

$$s = ut + \frac{1}{2}at^2 \quad (2)$$

$$v^2 = u^2 + 2as \quad (3)$$

$$s = \frac{u+v}{2}t \quad (4)$$

$$s = vt - \frac{1}{2}at^2 \quad (5)$$

Derivation (examinable): From $a = \frac{dv}{dt} \Rightarrow v = u + at$. Then $v = \frac{ds}{dt} \Rightarrow s = \int_0^t (u + at) dt = ut + \frac{1}{2}at^2$. Eliminate t from (1) and (2) to get (3). Equation (4) follows from average velocity $\frac{u+v}{2}$ for constant a . Equation (5) is obtained by substituting $u = v - at$ into (2).

KEY POINTS Sign convention

Always fix a positive direction at the start. All vector quantities (s, u, v, a) carry signs relative to that direction. A negative velocity means motion in the negative direction; a negative acceleration opposite to the velocity means retardation.

WARNING Common pitfalls

- Forgetting to convert km h^{-1} to m s^{-1} (multiply by $\frac{5}{18}$).
- Using $g = 9.8 \text{ m s}^{-2}$ unless the question specifies $g = 10 \text{ m s}^{-2}$.
- Confusing distance (scalar, sum of path lengths) with displacement (vector, final position minus initial position).
- Taking the wrong root when solving a quadratic for time; always reject negative or unphysical times.

10.2 Equations of Motion under Uniform Acceleration**METHOD** Solving a uniform-acceleration problem with the suvat equations

1. Draw a diagram and fix a positive direction (usually the direction of motion). Write down the five quantities: s (displacement), u (initial velocity), v (final velocity), a (acceleration), t (time). Give each known quantity its correct sign.
2. Identify the three known and the one unknown you need. Note which quantity is not mentioned.
3. Select the equation that omits the missing quantity:

Equation	Contains	Omits
$v = u + at$	v, u, a, t	s
$s = ut + \frac{1}{2}at^2$	s, u, a, t	v
$v^2 = u^2 + 2as$	v, u, a, s	t
$s = \frac{(u+v)}{2}t$	s, u, v, t	a
$s = vt - \frac{1}{2}at^2$	s, v, a, t	u

4. Substitute with units, solve, and check the sign and the magnitude of the answer against common sense.

5. Interpret: a negative velocity means motion in the negative direction; a negative acceleration opposite to motion means a **retardation** (deceleration).

Reflexes: “starts from rest” $\Rightarrow u = 0$; “comes to rest” $\Rightarrow v = 0$; convert km h^{-1} to m s^{-1} by multiplying by $\frac{1000}{3600} = \frac{5}{18}$.

WORKED EXAMPLE 15 Braking of a taxi in Douala

A taxi travelling at 72 km h^{-1} along a straight road in Douala sees an obstacle 50 m ahead. The driver applies the brakes, producing a uniform retardation. The taxi just stops at the obstacle. Find (a) the retardation, (b) the time taken to stop.

SOLUTION

Solution. Take the direction of motion as positive.

Step 1: convert units.

$$u = 72 \times \frac{5}{18} = 20 \text{ m s}^{-1}, \quad v = 0, \quad s = 50 \text{ m}.$$

- (a) Time is not given and not required, so use $v^2 = u^2 + 2as$:

$$0 = 20^2 + 2a(50) \Rightarrow 100a = -400 \Rightarrow a = -4 \text{ m s}^{-2}.$$

The retardation is therefore 4 m s^{-2} (the negative sign indicates acceleration opposing the motion).

- (b) Now use $v = u + at$:

$$0 = 20 - 4t \Rightarrow t = 5 \text{ s}.$$

Check: average velocity $= \frac{u+v}{2} = 10 \text{ m s}^{-1}$, so $s = 10 \times 5 = 50 \text{ m}$. ✓

10.3 Displacement–Time and Velocity–Time Graphs

METHOD Interpreting a displacement–time (s – t) graph

1. The gradient of the s – t graph at any instant gives the **velocity**: $v = \frac{ds}{dt}$.
2. A horizontal section means the particle is at rest; a straight inclined section means constant velocity; a curve means the velocity is changing (acceleration).
3. A negative gradient means motion back towards the origin (velocity in the negative direction).

- The total distance travelled is found by adding the magnitudes of all changes of position; the displacement is the final position minus the initial position (signs included).
- To sketch an s - t graph from a description: mark the key times, join them with straight segments (constant velocity) or smooth curves (acceleration), making the gradient steeper where the speed is greater.


WORKED EXAMPLE 16 A walk to the market

A trader leaves home and walks along a straight road. Her displacement s (in metres) from home at time t (in seconds) is recorded: she walks at constant velocity and reaches the market 400 m away after 200 s; she stays there for 100 s; she then walks back home at constant speed, arriving at $t = 500$ s. (a) Sketch the displacement–time graph. (b) Find her velocity on each stage of the journey. (c) Find her total distance travelled and her average speed for the whole journey.

SOLUTION

Solution.

(a) The graph has three straight sections: rising from $(0, 0)$ to $(200, 400)$, horizontal from $(200, 400)$ to $(300, 400)$, and falling from $(300, 400)$ to $(500, 0)$.



Displacement–time graph for the trader's journey.

(b) Velocity = gradient of the s - t graph.

$$\text{Outward: } v_1 = \frac{400 - 0}{200 - 0} = 2 \text{ m s}^{-1}.$$

$$\text{At market: } v_2 = \frac{400 - 400}{300 - 200} = 0 \text{ m s}^{-1} \quad (\text{at rest}).$$

$$\text{Return: } v_3 = \frac{0 - 400}{500 - 300} = -2 \text{ m s}^{-1}.$$

The negative sign shows she is moving back towards home.

(c) Total distance = $400 + 0 + 400 = 800$ m. Total time = 500 s.

$$\text{Average speed} = \frac{800}{500} = 1.6 \text{ m s}^{-1}.$$

Note: her average velocity is $\frac{0 - 0}{500} = 0 \text{ m s}^{-1}$, since the total displacement is zero.

METHOD Extracting information from a velocity–time (v – t) graph

1. The gradient of the v – t graph gives the **acceleration**: $a = \frac{dv}{dt}$.
2. The area between the graph and the time axis gives the **displacement**. Areas below the axis count as negative.
3. Split the area into triangles, rectangles and trapezia: triangle = $\frac{1}{2}bh$, rectangle = bh , trapezium = $\frac{1}{2}(a + b)h$.
4. Total distance = sum of the magnitudes of the areas; displacement = algebraic sum.
5. To sketch a v – t graph from a description: constant acceleration $\Rightarrow$ straight sloping line; constant velocity $\Rightarrow$ horizontal line; rest $\Rightarrow$ line on the time axis.


WORKED EXAMPLE 17 Journey of a bus between two stops

A bus starts from rest at a stop and accelerates uniformly for 10 s until it reaches 15 m s^{-1} . It maintains this velocity for 30 s, then decelerates uniformly, coming to rest at the next stop after a further 15 s. (a) Sketch the velocity–time graph. (b) Find the acceleration during each stage. (c) Find the distance between the two stops.

SOLUTION

Solution.

(a) Three straight sections: rising from (0, 0) to (10, 15), horizontal from (10, 15) to (40, 15), falling from (40, 15) to (55, 0).



Velocity–time graph of the bus; the shaded area gives the distance.

(b) Acceleration = gradient.

$$\text{Stage 1: } a_1 = \frac{15 - 0}{10 - 0} = 1.5 \text{ m s}^{-2}.$$

$$\text{Stage 2: } a_2 = 0 \quad (\text{constant velocity}).$$

$$\text{Stage 3: } a_3 = \frac{0 - 15}{55 - 40} = -1 \text{ m s}^{-2} \quad (\text{retardation } 1 \text{ m s}^{-2}).$$

(c) Distance = total area under the graph = triangle + rectangle + triangle:

$$s = \frac{1}{2}(10)(15) + (30)(15) + \frac{1}{2}(15)(15) = 75 + 450 + 112.5 = \boxed{637.5 \text{ m}}.$$

Check (trapezium formula): the whole area is a trapezium of parallel sides 30 and 55, height 15: $s = \frac{1}{2}(30 + 55)(15) = 637.5 \text{ m}$. ✓

10.4 Acceleration–Time Graphs and Connecting the Three Graphs

METHOD Moving between a - t , v - t and s - t graphs

1. From a - t to v - t : the change in velocity over an interval equals the area under the a - t graph: $\Delta v = \int a \, dt$. Add the initial velocity u .
2. From v - t to s - t : the change in displacement equals the area under the v - t graph: $\Delta s = \int v \, dt$.
3. From s - t to v - t to a - t : differentiate: $v = \frac{ds}{dt}$, $a = \frac{dv}{dt} = \frac{d^2s}{dt^2}$.
4. Key correspondences to memorise:

s - t graph	v - t graph	a - t graph
Straight, sloping	Horizontal (constant v)	On the axis ($a = 0$)
Curve, steepening	Straight, sloping	Horizontal ($a \neq 0$)
Curve, changing curvature	Curve	Sloping line

5. Always check consistency: where the v - t graph crosses the axis, the s - t graph has a turning point; where the a - t graph crosses the axis, the v - t graph has a turning point.


WORKED EXAMPLE 18 From an acceleration-time graph to velocity and displacement

A particle starts from rest at O and moves in a straight line. Its acceleration is 4 m s^{-2} for the first 5 s, then zero for the next 5 s, then -2 m s^{-2} for the final 10 s. (a) Sketch the acceleration-time and velocity-time graphs. (b) Find the velocity at $t = 5$, $t = 10$ and $t = 20$. (c) Find the total displacement after 20 s.

SOLUTION

Solution.

- (a) The a - t graph consists of three horizontal segments at heights 4, 0, -2.



Acceleration-time graph of the particle.

- (b) Change in velocity = area under the a - t graph, with $u = 0$.


$$v(5) = 0 + (4)(5) = 20 \text{ m s}^{-1}.$$

$$v(10) = 20 + (0)(5) = 20 \text{ m s}^{-1}.$$

$$v(20) = 20 + (-2)(10) = 0 \text{ m s}^{-1}.$$

The particle ends at rest — a useful check.

The v - t graph is therefore: rising line from $(0, 0)$ to $(5, 20)$, horizontal from $(5, 20)$ to $(10, 20)$, falling from $(10, 20)$ to $(20, 0)$.



Velocity-time graph obtained from the areas under the a - t graph.

(c) Displacement = area under the v - t graph:

$$s = \frac{1}{2}(5)(20) + (5)(20) + \frac{1}{2}(10)(20) = 50 + 100 + 100 = \boxed{250 \text{ m}}.$$

10.5 Free Fall and Motion under Gravity

DEFINITION Free fall

A body moving under the influence of gravity alone (air resistance neglected) is in **free fall**. Near the Earth's surface the acceleration has constant magnitude $g = 9.8 \text{ ms}^{-2}$ (use $g = 10 \text{ ms}^{-2}$ only if the question instructs so) directed vertically downwards.

METHOD Solving free-fall problems

1. Choose a positive direction (usually upwards) and keep it for the whole solution. With upwards positive, $a = -g = -9.8 \text{ ms}^{-2}$.
2. List s, u, v, a, t with signs. Remember: at the greatest height, $v = 0$; when the body returns to the launch level, $s = 0$.
3. Apply the suvat equations exactly as for any uniform acceleration — gravity is a uniform acceleration.
4. For a body dropped from rest: $u = 0$; for a body projected downwards, $u > 0$ downwards.
5. Time of flight for a body projected upwards from ground level: $T = \frac{2u}{g}$; greatest

height: $H = \frac{u^2}{2g}$. (Derive these in the exam rather than quoting blindly.)

6. Ignore air resistance unless the question states otherwise — this is the standard GCE modelling assumption.

WORKED EXAMPLE 19 Stone thrown upwards from a cliff

A stone is thrown vertically upwards with velocity 14 m s^{-1} from the top of a cliff 30 m above the sea. Taking $g = 9.8 \text{ m s}^{-2}$ and neglecting air resistance, find (a) the greatest height above the cliff top reached by the stone, (b) the time at which the stone passes the cliff top on its way down, (c) the total time until the stone hits the sea, (d) the velocity with which it hits the sea.

SOLUTION

Solution. Take upwards as positive, origin at the cliff top. Then $u = 14 \text{ m s}^{-1}$, $a = -9.8 \text{ m s}^{-2}$.

(a) At greatest height, $v = 0$. Using $v^2 = u^2 + 2as$:

$$0 = 14^2 + 2(-9.8)s \implies s = \frac{196}{19.6} = \boxed{10 \text{ m}}.$$

(b) Passing the cliff top again means $s = 0$. Using $s = ut + \frac{1}{2}at^2$:

$$0 = 14t - 4.9t^2 = t(14 - 4.9t).$$

The solutions are $t = 0$ (the throw) and $t = \frac{14}{4.9} = \frac{20}{7} \approx \boxed{2.86 \text{ s}}$.

(c) Hitting the sea means $s = -30 \text{ m}$ (below the origin):

$$-30 = 14t - 4.9t^2 \implies 4.9t^2 - 14t - 30 = 0.$$

By the quadratic formula:

$$t = \frac{14 \pm \sqrt{14^2 + 4(4.9)(30)}}{2(4.9)} = \frac{14 \pm \sqrt{196 + 588}}{9.8} = \frac{14 \pm 28}{9.8}.$$

Taking the positive root: $t = \frac{42}{9.8} = \frac{30}{7} \approx \boxed{4.29 \text{ s}}$. (The negative root $t = -\frac{12}{9.8}$ is rejected.)

(d) Using $v = u + at$:

$$v = 14 - 9.8\left(\frac{30}{7}\right) = 14 - 42 = -28 \text{ m s}^{-1}.$$

The stone hits the sea with speed $\boxed{28 \text{ m s}^{-1}}$ directed downwards.

Check: $v^2 = u^2 + 2as = 196 + 2(-9.8)(-30) = 196 + 588 = 784$, so $v = \pm 28$. ✓ The negative root is chosen since the stone is moving downwards.

10.6 Two-Particle Problems (Meeting and Overtaking)

METHOD Solving problems with two moving particles

1. Write the displacement of each particle as a function of time, measured from the same origin in the same positive direction: $s_A(t)$ and $s_B(t)$.
2. If one particle starts later, use $t - t_0$ for its time of motion.
3. Meeting / overtaking condition: set $s_A(t) = s_B(t)$ and solve for t . Reject negative or otherwise impossible roots.
4. Substitute back to find the position of the meeting and the velocities at that instant if required.
5. Alternatively, read the meeting point off a pair of $s-t$ graphs drawn on the same axes: the intersection of the two curves.

WORKED EXAMPLE 20 The police car and the speeding lorry

A lorry travels along a straight highway at a constant velocity of 20 m s^{-1} . As it passes a police checkpoint, a police car starts from rest and accelerates uniformly at 2.5 m s^{-2} in pursuit. Find (a) the time at which the police car catches the lorry, (b) the distance from the checkpoint at which this happens, (c) the velocity of the police car at that instant.

SOLUTION

Solution. Take the checkpoint as origin, direction of travel positive, and $t = 0$ when the lorry passes.

Displacements:

$$s_L = 20t \quad (\text{lorry, constant velocity}),$$

$$s_P = 0 \cdot t + \frac{1}{2}(2.5)t^2 = 1.25t^2 \quad (\text{police car, from rest}).$$

(a) Catching up: $s_P = s_L$:

$$1.25t^2 = 20t \implies t(1.25t - 20) = 0.$$

$t = 0$ corresponds to the moment the lorry passes the checkpoint. The overtaking occurs at

$$t = \frac{20}{1.25} = 16 \text{ s}.$$

(b) Distance: $s_L = 20 \times 16 = 320 \text{ m}$ from the checkpoint.

(c) Police car's velocity: $v = u + at = 0 + 2.5 \times 16 = 40 \text{ m s}^{-1}$.

Remark: on the same $s-t$ axes, the lorry's graph is the straight line $s = 20t$ and the police car's graph is the parabola $s = 1.25t^2$; they intersect at $t = 0$ and $t = 16$, confirming the answer graphically.

10.7 Graphical Treatment of Motion under Gravity

METHOD Sketching and using graphs for vertical motion

- For a body projected upwards (upwards positive):
 - the $v-t$ graph is a straight line of gradient $-g$, starting at u and crossing the axis at the time of greatest height;
 - the $s-t$ graph is a parabola opening downwards, with its vertex at the greatest height;
 - the $a-t$ graph is a horizontal line at $-g$ for the entire flight (including at the top!).
- The area under the $v-t$ graph up to the crossing time gives the greatest height; the signed area up to any later time gives the displacement.
- The speed-time graph differs from the velocity-time graph: speed is never negative, so the part below the axis is reflected upwards (a “V” shape).
- Use graph areas to verify algebraic answers — a powerful checking reflex in the examination.


WORKED EXAMPLE 21 Ball dropped from a water tower

A ball is released from rest from the top of a water tower in Bamenda, 45 m above the ground. Taking $g = 10 \text{ m s}^{-2}$, (a) sketch the velocity-time and displacement-time graphs (taking downwards as positive) for the whole fall; (b) use the $v-t$ graph to find the time of fall and the velocity of impact; (c) verify your answers using the equations of motion.

SOLUTION

Solution. Downwards positive: $u = 0$, $a = g = 10 \text{ m s}^{-2}$.

(a) The $v-t$ graph is a straight line through the origin of gradient 10: $v = 10t$. The $s-t$ graph is the parabola $s = 5t^2$, ending at $s = 45$.



Velocity–time graph of the falling ball; the shaded area equals 45 m.

(b) The fall ends when the area under the v – t graph equals 45 m. If T is the time of fall, the area is a triangle:

$$\frac{1}{2} \times T \times 10T = 45 \Rightarrow 5T^2 = 45 \Rightarrow T^2 = 9 \Rightarrow T = \boxed{3 \text{ s}}.$$

Impact velocity: $v = 10T = \boxed{30 \text{ m s}^{-1}}$.

(c) Verification with equations. Using $s = ut + \frac{1}{2}at^2$:

$$45 = 0 + \frac{1}{2}(10)t^2 \Rightarrow t^2 = 9 \Rightarrow t = 3 \text{ s.} \checkmark$$

Using $v^2 = u^2 + 2as$: $v^2 = 0 + 2(10)(45) = 900$, so $v = 30 \text{ m s}^{-1}$. $\checkmark$

The graphical and algebraic methods agree, as they must: the suvat equations are exactly the “area and gradient” statements for a straight-line v – t graph.

10.8 Real-Life Applications and Synthesis

EXPERIMENT Reaction time and stopping distance

A driver travelling at 30 m s^{-1} ($\approx 108 \text{ km h}^{-1}$) sees a hazard. The typical human reaction time is 0.7 s. During this time the car travels at constant speed. The driver then brakes with a uniform retardation of 6 m s^{-2} .

1. Find the thinking distance (distance during reaction time).
2. Find the braking distance.
3. Find the total stopping distance.
4. Sketch the v – t graph for the whole event.

Solution.

1. Thinking distance = $u \times t_{\text{react}} = 30 \times 0.7 = 21 \text{ m}$.
2. Braking: $u = 30$, $v = 0$, $a = -6$. Use $v^2 = u^2 + 2as$: $0 = 900 - 12s \Rightarrow s = 75 \text{ m}$.
3. Total stopping distance = $21 + 75 = \boxed{96 \text{ m}}$.
4. v - t graph: horizontal line at $v = 30$ from $t = 0$ to $t = 0.7$, then straight line down to $v = 0$ at $t = 0.7 + \frac{30}{6} = 5.7 \text{ s}$. The area under the graph is $21 + 75 = 96 \text{ m}$.

DID YOU KNOW?

Did you know?

The Highway Code in Cameroon (and many countries) quotes stopping distances based on a reaction time of 0.7 s and a deceleration of about 6 m s^{-2} on dry roads. At 50 km h^{-1} ($\approx 13.9 \text{ m s}^{-1}$) the total stopping distance is roughly 23 m; at 100 km h^{-1} it quadruples to about 96 m because braking distance is proportional to u^2 . This is why speed limits are critical for safety.

EXERCISE 8

Consolidation exercises

1. A particle moves with uniform acceleration. It passes point A with velocity 10 m s^{-1} and point B with velocity 20 m s^{-1} . The distance AB is 45 m. Find the acceleration and the time taken to travel from A to B.
2. A ball is thrown vertically upwards from ground level with speed 24.5 m s^{-1} . Taking $g = 9.8 \text{ m s}^{-2}$, find (a) the greatest height, (b) the total time of flight, (c) the velocity when it is 14.7 m above the ground on its way down.
3. Two particles P and Q start simultaneously from the same point O on a straight line. P moves with constant velocity 5 m s^{-1} . Q starts from rest with constant acceleration 2 m s^{-2} . Find (a) the time when Q overtakes P, (b) the distance from O at that moment, (c) the velocity of Q then.
4. A car accelerates uniformly from rest to 20 m s^{-1} in 10 s, travels at constant speed for 20 s, then decelerates uniformly to rest in 5 s. Sketch the v - t graph and find the total distance travelled.
5. A stone is dropped from a bridge 78.4 m above a river. At the same instant, another stone is thrown vertically downwards from the bridge with speed 19.6 m s^{-1} . Taking $g = 9.8 \text{ m s}^{-2}$, find the time interval between the two stones hitting the water.

SOLUTION Exercise 1

$$a = 3.33 \text{ m s}^{-2} \text{ (or } \frac{10}{3}), t = 3 \text{ s.}$$

SOLUTION Exercise 2

(a) $H = \frac{u^2}{2g} = \frac{24.5^2}{19.6} = 30.625 \text{ m.}$ (b) $T = \frac{2u}{g} = \frac{49}{9.8} = 5 \text{ s.}$ (c) Downwards positive from max height: $s = 30.625 - 14.7 = 15.925 \text{ m, } u = 0, a = 9.8, v^2 = 2 \times 9.8 \times 15.925 = 312.13, v \approx 17.67 \text{ m s}^{-1}$ downwards.

SOLUTION Exercise 3

(a) $5t = t^2 \Rightarrow t = 5 \text{ s}$ (excluding $t = 0$). (b) $s = 5 \times 5 = 25 \text{ m.}$ (c) $v = 2 \times 5 = 10 \text{ m s}^{-1}.$

SOLUTION Exercise 4

$$\text{Area} = \frac{1}{2}(10)(20) + (20)(20) + \frac{1}{2}(5)(20) = 100 + 400 + 50 = 550 \text{ m.}$$

SOLUTION Exercise 5

Dropped stone: $78.4 = \frac{1}{2}(9.8)t_1^2 \Rightarrow t_1 = 4 \text{ s.}$ Thrown stone: $78.4 = 19.6t_2 + 4.9t_2^2 \Rightarrow t_2^2 + 4t_2 - 16 = 0 \Rightarrow t_2 = -2 + 2\sqrt{5} \approx 2.47 \text{ s.}$ Interval $\approx 4 - 2.47 = 1.53 \text{ s.}$

11 EXERCISES

11.1 I check my knowledge

EXERCISE 9 MCQ: Uniform acceleration

A car starts from rest and accelerates uniformly at 2 m s^{-2} for 10 s. What is its final velocity?

1. 5 m s^{-1}
2. 10 m s^{-1}
3. 20 m s^{-1}
4. 40 m s^{-1}

EXERCISE 10 True/False with justification

State whether each statement is true or false. Justify your answer.


1. The gradient of a displacement–time graph gives the acceleration.
2. The area under a velocity–time graph gives the displacement.
3. For a particle in free fall, the acceleration–time graph is a horizontal line.
4. If the velocity–time graph is a straight line with negative slope, the particle is decelerating.

EXERCISE 11 Definitions and direct calculations

1. Define displacement, velocity and acceleration for motion in a straight line.
2. A particle moves with constant velocity 15 m s^{-1} for 8 s. Calculate its displacement.
3. A body accelerates uniformly from 5 m s^{-1} to 25 m s^{-1} in 4 s. Find its acceleration and the distance covered.

EXERCISE 12 Graph interpretation

The velocity–time graph below shows the motion of a particle for 12 s.



1. Describe the motion in each of the three phases.
2. Calculate the acceleration during the first 4 s.
3. Find the total displacement after 12 s.

11.2 I practise

EXERCISE 13 Car journey with three phases

A car accelerates uniformly from rest to 25 m s^{-1} in 10 s, cruises at this speed for 20 s, then brakes uniformly to rest in 5 s.

1. Sketch the velocity–time graph for the whole journey.
2. Calculate the total distance travelled.
3. Find the average velocity for the journey.

EXERCISE 14 Free fall from a building

A stone is dropped from the top of a storey building and hits the ground after 3 s. Taking $g = 10 \text{ m s}^{-2}$,

1. find the height of the building,
2. find the impact velocity.

EXERCISE 15 Vertical projection

A ball is thrown vertically upward at 20 m s^{-1} from ground level. Taking $g = 10 \text{ m s}^{-2}$,

1. find the maximum height reached,
2. find the total time of flight,
3. find the velocity at $t = 3 \text{ s}$.


EXERCISE 16 Connecting graphs

A particle starts from rest at $x = 0$. Its acceleration–time graph consists of two rectangles: $a = 2 \text{ m s}^{-2}$ for 4 s , then $a = -3 \text{ m s}^{-2}$ until the particle comes to rest.

1. Sketch the velocity–time and displacement–time graphs.
2. Find the total displacement when the particle stops.

11.3 I prepare for the GCE**EXERCISE 17** GCE style: Trapezium velocity–time graph

(12 marks) A particle moves along a straight line. Its velocity v (in m/s) at time t (in s) is given by the trapezium-shaped graph below:



1. Find the acceleration during each of the three phases. (3 marks)
2. Calculate the total distance travelled in 10 s . (4 marks)
3. Determine the average velocity for the whole journey. (2 marks)
4. Sketch the corresponding acceleration–time graph. (3 marks)

EXERCISE 18 GCE style: Two objects meeting

(14 marks) A coconut falls from the top of a 20 m palm tree at the same instant a stone is thrown vertically upward from the ground with an initial velocity of 15 m s^{-1} . Take $g = 10 \text{ m s}^{-2}$.

1. Write down the equations of motion for the coconut and the stone, choosing the upward direction as positive. (4 marks)
2. Find the time at which they cross each other. (4 marks)
3. Find the height above the ground at which they cross. (3 marks)
4. Determine the velocities of both objects at the crossing instant. (3 marks)

EXERCISE 19 GCE style: Overtaking and stopping distance

(14 marks) Two taxis leave the same park. Taxi A moves at a constant velocity of 15 m s^{-1} . Taxi B starts from rest 5 s later with a uniform acceleration of 2 m s^{-2} .

1. Find the time (measured from the departure of taxi A) when taxi B overtakes taxi A. (5 marks)
2. Find the distance from the park where the overtaking occurs. (3 marks)

A driver is travelling at 90 km h^{-1} and sees an obstacle 60 m ahead. His reaction time is 0.7 s and the maximum braking deceleration is 5 m s^{-2} .

3. Calculate the distance travelled during the reaction time. (2 marks)
4. Calculate the braking distance. (2 marks)
5. Does the car stop before hitting the obstacle? Justify. (2 marks)

12 SOLUTIONS TO THE EXERCISES

SOLUTION Exercise 1 — MCQ: Uniform acceleration

The car starts from rest, so $u = 0$, with $a = 2 \text{ m s}^{-2}$ and $t = 10 \text{ s}$. Using the first equation of motion:

$$v = u + at = 0 + 2 \times 10 = 20 \text{ m s}^{-1}.$$

Correct answer: (c) 20 m s^{-1} .

SOLUTION Exercise 2 — True/False with justification

1. False. The gradient of a displacement–time graph gives the velocity, since $v = \frac{dx}{dt}$. Acceleration is the gradient of the velocity–time graph.
2. True. Since $v = \frac{dx}{dt}$, we have $x = \int v dt$, which is exactly the (signed) area under the velocity–time graph.
3. True. In free fall (neglecting air resistance) the acceleration is constant and equal to g , so the acceleration–time graph is a horizontal line $a = -g$ (taking upward as positive).
4. False in general. A straight velocity–time line with negative slope means the acceleration is negative. The particle decelerates (speed decreases) only if velocity and acceleration have opposite signs. If the velocity is itself negative, a negative acceleration makes the particle speed up in the negative direction.

SOLUTION Exercise 3 — Definitions and direct calculations

1. Displacement is the change in position of the particle, measured from a fixed origin on the line; it is a vector quantity (in 1-D, a signed scalar), unit: metre (m). Velocity is the rate of change of displacement with respect to time, $v = \frac{dx}{dt}$; unit: m s^{-1} . Acceleration is the rate of change of velocity with respect to time, $a = \frac{dv}{dt}$; unit: m s^{-2} .
2. Constant velocity: $x = vt = 15 \times 8 = 120 \text{ m}$.
3. Acceleration: $a = \frac{v - u}{t} = \frac{25 - 5}{4} = 5 \text{ m s}^{-2}$.
Distance: $s = \frac{u + v}{2} t = \frac{5 + 25}{2} \times 4 = 60 \text{ m}$. (Check: $s = ut + \frac{1}{2}at^2 = 20 + 40 = 60 \text{ m}$.)

SOLUTION Exercise 4 — Graph interpretation

1. Phase 1 ($0 \leq t \leq 4$): velocity increases uniformly from 0 to 8 m s^{-1} — uniform acceleration. Phase 2 ($4 \leq t \leq 8$): velocity constant at 8 m s^{-1} — zero acceleration. Phase 3 ($8 \leq t \leq 12$): velocity decreases uniformly from 8 m s^{-1} to 0 — uniform deceleration (retardation).

$$2. a = \frac{\Delta v}{\Delta t} = \frac{8 - 0}{4 - 0} = 2 \text{ m s}^{-2}.$$


3. The displacement is the area of the trapezium under the graph (bases 12 s and 4 s , height 8 m s^{-1}):

$$s = \frac{1}{2}(12 + 4) \times 8 = 64 \text{ m}.$$

(Equivalently: triangle $\frac{1}{2}(4)(8) = 16$, rectangle $4 \times 8 = 32$, triangle $\frac{1}{2}(4)(8) = 16$; total $16 + 32 + 16 = 64 \text{ m}$.)

SOLUTION Exercise 5 — Car journey with three phases

1. The graph consists of: a straight line from $(0, 0)$ to $(10, 25)$; a horizontal segment from $(10, 25)$ to $(30, 25)$; a straight line from $(30, 25)$ to $(35, 0)$.



2. Total distance = area under the graph (trapezium with parallel sides 35 s and 20 s , height 25 m s^{-1}):

$$s = \frac{1}{2}(35 + 20) \times 25 = \frac{1}{2} \times 55 \times 25 = 687.5 \text{ m}.$$

(Phase by phase: $\frac{1}{2}(10)(25) = 125$; $20 \times 25 = 500$; $\frac{1}{2}(5)(25) = 62.5$; total 687.5 m .)

$$3. \text{ Average velocity} = \frac{\text{total displacement}}{\text{total time}} = \frac{687.5}{35} \approx 19.6 \text{ m s}^{-1}.$$

SOLUTION Exercise 6 — Free fall from a building

Take the downward direction as positive. Then $u = 0$ (dropped), $a = g = 10 \text{ m s}^{-2}$, $t = 3 \text{ s}$.

1. Height of the building:

$$h = ut + \frac{1}{2}gt^2 = 0 + \frac{1}{2} \times 10 \times 3^2 = 45 \text{ m}.$$

2. Impact velocity:

$$v = u + gt = 0 + 10 \times 3 = 30 \text{ m s}^{-1} \quad (\text{downwards}).$$

SOLUTION Exercise 7 — Vertical projection

Take upward as positive: $u = 20 \text{ m s}^{-1}$, $a = -g = -10 \text{ m s}^{-2}$.

1. At maximum height, $v = 0$. Using $v^2 = u^2 + 2as$:

$$0 = 20^2 - 2(10)h \implies h = \frac{400}{20} = 20 \text{ m}.$$


2. Time to reach the top: $0 = 20 - 10t \implies t = 2 \text{ s}$. By symmetry, the descent takes the same time, so the total time of flight is $T = 2 \times 2 = 4 \text{ s}$. (Alternatively, solve $s = 20t - 5t^2 = 0$, giving $t = 0$ or $t = 4$.)

3. At $t = 3 \text{ s}$: $v = u - gt = 20 - 10(3) = -10 \text{ m s}^{-1}$. The negative sign means the ball is moving downwards at 10 m s^{-1} (it is on its way down).

SOLUTION Exercise 8 — Connecting graphs

1. Phase 1 ($0 \leq t \leq 4$): $a = 2 \text{ m s}^{-2}$, so $v = 2t$ (straight line through the origin); at $t = 4$, $v_{\max} = 8 \text{ m s}^{-1}$. Phase 2: $a = -3 \text{ m s}^{-2}$, so $v = 8 - 3(t - 4)$; the particle stops when $v = 0$, i.e. $t - 4 = \frac{8}{3}$, so $t = \frac{20}{3} \approx 6.67 \text{ s}$.

The velocity–time graph is a triangle with vertices $(0, 0)$, $(4, 8)$, $(20/3, 0)$. The displacement–time graph is a parabola opening upwards for $0 \leq t \leq 4$ ($x = t^2$), then a parabola opening downwards from $t = 4$ to $t = 20/3$, with horizontal tangent at the end (since $v = 0$).



2. Total displacement = area of the triangle:

$$s = \frac{1}{2} \times \frac{20}{3} \times 8 = \frac{80}{3} \approx 26.7 \text{ m.}$$

(Phase by phase: $s_1 = \frac{1}{2}(4)(8) = 16 \text{ m}$; $s_2 = \frac{1}{2}\left(\frac{8}{3}\right)(8) = \frac{32}{3} \text{ m}$; total $16 + \frac{32}{3} = \frac{80}{3} \text{ m.}$)

SOLUTION
Exercise 9 — GCE style: Trapezium velocity–time graph

(a) Accelerations (gradient of each segment):


- Phase 1 ($0 \leq t \leq 2$): $a = \frac{6-0}{2-0} = 3 \text{ m s}^{-2}$.
- Phase 2 ($2 \leq t \leq 6$): $a = \frac{6-6}{6-2} = 0 \text{ m s}^{-2}$.
- Phase 3 ($6 \leq t \leq 10$): $a = \frac{0-6}{10-6} = -1.5 \text{ m s}^{-2}$ (retardation of 1.5 m s^{-2}).

(b) Total distance = area of the trapezium (parallel sides 10 s and 4 s, height 6 m s^{-1}):

$$s = \frac{1}{2}(10 + 4) \times 6 = 42 \text{ m.}$$

(c) Average velocity = $\frac{s}{t} = \frac{42}{10} = 4.2 \text{ m s}^{-1}$.

(d) Acceleration–time graph: three horizontal segments: $a = 3$ on $[0, 2]$, $a = 0$ on $[2, 6]$, $a = -1.5$ on $[6, 10]$.



Indicative mark scheme (12 marks): (a) M1 for gradient method, A1 for 3 and 0, A1 for -1.5 (3 marks). (b) M1 area method, A1 split into three correct parts, M1 summing, A1 for 42 m (4 marks). (c) M1 formula, A1 for 4.2 m s^{-1} (2 marks). (d) B1 horizontal segments, B1 correct values, B1 correct time intervals with discontinuities shown (3 marks). Examiner's expectations: units must be quoted; a negative acceleration must be interpreted as deceleration only because velocity is positive; the area must be identified with displacement.

SOLUTION Exercise 10 — GCE style: Two objects meeting

Take upward as positive, origin at the ground, $g = 10 \text{ m s}^{-2}$.

1. Coconut (dropped from rest at $x = 20$): $u = 0$, $a = -10$:

$$x_C = 20 - 5t^2, \quad v_C = -10t.$$

Stone (projected from the ground at 15 m s^{-1}): $u = 15$, $a = -10$:

$$x_S = 15t - 5t^2, \quad v_S = 15 - 10t.$$

2. They cross when $x_C = x_S$:

$$20 - 5t^2 = 15t - 5t^2 \implies 20 = 15t \implies t = \frac{4}{3} \text{ s} \approx 1.33 \text{ s}.$$

3. Height of crossing: $x_C = 20 - 5\left(\frac{4}{3}\right)^2 = 20 - \frac{80}{9} = \frac{100}{9} \approx 11.1 \text{ m}$ above the ground.

4. Velocities at $t = \frac{4}{3} \text{ s}$:

$$v_C = -10 \times \frac{4}{3} = -\frac{40}{3} \approx -13.3 \text{ m s}^{-1} \text{ (coconut moving down),}$$

$$v_S = 15 - 10 \times \frac{4}{3} = \frac{5}{3} \approx 1.67 \text{ m s}^{-1} \text{ (stone still moving up).}$$

Indicative mark scheme (14 marks): (a) B1 sign convention stated, B1 coconut equation, B1 stone equation, B1 both velocity equations (4 marks). (b) M1 equating positions, A1 simplification (the t^2 terms cancel), M1 solving, A1 $t = 4/3 \text{ s}$ (4 marks). (c) M1 substitution, A1

exact value, A1 decimal with unit (3 marks). (d) M1 use of $v = u + at$, A1 v_C , A1 v_S with directions interpreted (3 marks). Examiner's expectations: a clearly stated sign convention; the physical meaning of the signs of the velocities; exact fractions accepted.

SOLUTION Exercise 11 — GCE style: Overtaking and stopping distance

(a) Overtaking time. Let t be the time measured from the departure of taxi A. Taxi A (constant velocity): $x_A = 15t$. Taxi B starts at $t = 5$ from rest with $a = 2 \text{ m s}^{-2}$, so for $t \geq 5$:

$$x_B = \frac{1}{2}(2)(t - 5)^2 = (t - 5)^2.$$

B overtakes A when $x_B = x_A$:

$$(t - 5)^2 = 15t \implies t^2 - 10t + 25 = 15t \implies t^2 - 25t + 25 = 0.$$

$$t = \frac{25 \pm \sqrt{625 - 100}}{2} = \frac{25 \pm \sqrt{525}}{2} = \frac{25 \pm 5\sqrt{21}}{2}.$$

Numerically $t \approx \frac{25 \pm 22.91}{2}$, giving $t \approx 1.04 \text{ s}$ (rejected, since B has not started yet, $t < 5$) or

$$t \approx 23.96 \text{ s} \approx 24.0 \text{ s} \quad \text{after A's departure.}$$

(b) Distance from the park:

$$x_A = 15 \times 23.96 \approx 359 \text{ m.}$$

(c) Reaction distance. Convert the speed: $90 \text{ km h}^{-1} = \frac{90}{3.6} = 25 \text{ m s}^{-1}$.

$$d_{\text{reaction}} = vt = 25 \times 0.7 = 17.5 \text{ m.}$$

(d) Braking distance. Using $v^2 = u^2 + 2as$ with final velocity 0 and $a = -5 \text{ m s}^{-2}$:

$$0 = 25^2 - 2(5)d \implies d = \frac{625}{10} = 62.5 \text{ m.}$$


(e) Total stopping distance:

$$d_{\text{total}} = 17.5 + 62.5 = 80 \text{ m} > 60 \text{ m.}$$

No, the car does not stop in time: it needs 80 m but the obstacle is only 60 m ahead, so it hits the obstacle (with a residual speed $v = \sqrt{625 - 10(60 - 17.5)} = \sqrt{200} \approx 14.1 \text{ m s}^{-1}$ — optional comment).

Indicative mark scheme (14 marks): (a) B1 expression for x_A , B1 expression for x_B with the time shift $(t - 5)$, M1 equating, M1 solving the quadratic, A1 selecting the valid root $t \approx 24.0 \text{ s}$ with rejection of the other root justified (5 marks). (b) M1 substitution, A1 value, A1 unit (3 marks). (c) M1 conversion of km/h to m/s, A1 17.5 m (2 marks). (d) M1 use of $v^2 = u^2 + 2as$, A1 62.5 m (2 marks). (e) M1 adding both distances, A1 correct conclusion with comparison (2 marks). Examiner's expectations: the time delay of taxi B must appear as $(t - 5)$; the extraneous root must be rejected with a reason; the final answer to (e) must be an explicit justified statement, not just a number.

13 SUMMARY SHEET



Mind map of the chapter: kinematics of particles moving in a straight line.

KEY POINTS Summary Sheet — Kinematics of Particles Moving in a Straight Line

Key definitions

- **Position** x : location of a particle relative to a fixed origin O on the line; **displacement** s : change of position, a vector quantity (its sign gives the direction).
- **Distance**: total length of the path travelled, a scalar (always positive or zero).
- **Velocity**: $v = \frac{ds}{dt}$, the rate of change of displacement; **speed** = $|v|$, the magnitude of velocity.
- **Acceleration**: $a = \frac{dv}{dt} = \frac{d^2s}{dt^2}$; **deceleration** (retardation) means the acceleration is in the direction opposite to the motion.
- **Uniform acceleration**: the acceleration a remains constant throughout the motion.
- **Free fall**: motion under gravity alone (air resistance neglected), with $a = g \approx 9.81 \text{ m s}^{-2}$ directed vertically downwards (take $g = 10 \text{ m s}^{-2}$ only when the question instructs it).

Formulas to know (uniform acceleration, "suvat")

$$v = u + at, \quad s = ut + \frac{1}{2}at^2, \quad s = vt - \frac{1}{2}at^2,$$

$$v^2 = u^2 + 2as, \quad s = \frac{1}{2}(u + v)t,$$

where u = initial velocity, v = final velocity, a = acceleration, s = displacement, t = time. Each equation involves exactly four of the five quantities. These are valid only when a is constant.

Graphical facts

- x - t graph: gradient = velocity; a horizontal part means the particle is at rest; a curve bending upwards means it is speeding up.
- v - t graph: gradient = acceleration; signed area between the graph and the time axis = displacement; area below the time axis counts negative.
- a - t graph: area between the graph and the time axis = change in velocity $v - u$; for uniform acceleration the graph is a horizontal line.
- The three graphs are linked by calculus: the gradient of the x - t graph gives the v - t graph; the gradient of the v - t graph gives the a - t graph; going the other way uses areas (integration).

Free fall essentials

- Choose a positive direction (usually upwards) and keep it throughout: then $a = -g$ for the whole motion, going up and coming down.
- At the highest point, $v = 0$ but $a = -g$: the acceleration is never zero during free fall.
- For a particle projected vertically upwards with speed u : time to the highest point $t_{\text{up}} = \frac{u}{g}$, maximum height $H = \frac{u^2}{2g}$; the time up equals the time down to the same level, and the particle returns with speed u .
- An object dropped from rest at height h : $h = \frac{1}{2}gt^2$, impact speed $v = \sqrt{2gh}$.

Methods

1. Draw a diagram, mark the origin and the positive direction.
2. List the known suvat quantities and the one required; pick the equation containing exactly those four.
3. Substitute with signs, solve, then interpret the result (a negative time, or a negative speed for a car still moving forwards, may have to be rejected).
4. For multi-stage motion, treat each stage separately: the final velocity of one stage is the initial velocity of the next.
5. For graph questions: compute gradients and areas stage by stage; use the total signed area for displacement, and the sum of the absolute areas for distance.

Common mistakes

- Using the suvat equations when the acceleration is not constant.

- Mixing signs: forgetting that $a = -g$ when “up” is positive, or giving a falling object $a = +g$ while taking upwards as positive.
- Confusing distance with displacement on a $v-t$ graph (ignoring the area below the axis).
- Thinking that $v = 0$ at maximum height means $a = 0$ — false, $a = -g$ throughout the flight.
- Reading the gradient of a $v-t$ graph as velocity instead of acceleration.
- Forgetting units, or mixing km/h with m/s (divide by 3.6 to convert km/h to m/s).

GCE tips

- Always state the equation you use before substituting — method marks are awarded for this.
- Use $g = 9.81 \text{ m s}^{-2}$ unless the paper says $g = 10 \text{ m s}^{-2}$; give final answers to 3 significant figures.
- Sketch the $v-t$ graph even when not asked: it often reveals the quickest solution (the area method).
- Check your answer physically: a braking car must have $a < 0$ and $v \geq 0$ until it stops.
- In “two particles” problems (one thrown up, one dropped from height h), write $s_1 + s_2 = h$ at the meeting instant and solve for t .



Signed • OnBuch+

ONBUCH+ TEACHING TEAM
FOLLOWS THE OFFICIAL MINESEC SYLLABUS

OnBuch+