

PURE MATHEMATICS WITH STATISTICS • UPPER SIXTH

UPPER SIXTH — PURE MATHEMATICS WITH STATISTICS

CHAPTER 9

REGRESSION AND CORRELATION

SUGGESTED TIME: 15 PERIODS

IN THIS CHAPTER

This chapter introduces regression and correlation, the statistical study of relationships between two variables. Students learn to draw scatter diagrams, fit regression lines by Mayer's method and by least squares, compute residual sums of squares, and measure association using the product moment, Spearman's and Kendall's correlation coefficients.

- By the end of this chapter I can interpret bivariate data and explain the idea of regression.
- By the end of this chapter I can draw and interpret a scatter diagram and sketch regression lines.
- By the end of this chapter I can calculate a regression line using Mayer's method.
- By the end of this chapter I can compute sample variance and covariance from raw data.
- By the end of this chapter I can determine the least squares regression lines of y on x and of x on y .
- By the end of this chapter I can compute and interpret the residual sum of squares for each regression line.



DOWNLOAD ON- BUCH

Free • Google Play
AI tutor, past papers,
results



GO ONBUCH+

All signed courses, unlimited AI
tutor, PDF sheets — OnBuch+
tab in the app

THIS SIGNED COURSE CONTAINS

COURSE

complete, illustrated,
step by step

METHODS

and worked examples

EXERCISES

exam style + solutions

SUMMARY SHEET

the essentials to
remember



Signed • OnBuch+

ONBUCH+ TEACHING TEAM
FOLLOWS THE OFFICIAL MINESEC SYLLABUS

OnBuch+

CONTENTS

- | | |
|---|---|
| 1. Introduction to Regression and Scatter Diagrams | efficient |
| 2. Mayer's Method for the Regression Line | 7. Rank Correlation: Spearman and Kendall |
| 3. Correlation, Sample Variance and Co-variance | 8. Integration activity |
| 4. Least Squares Regression Lines | 9. Methods and worked examples |
| 5. Residual Sum of Squares and the Line of Best Fit | 10. Exercises |
| 6. The Product Moment Correlation Co-efficient | 11. Solutions to the exercises |
| | 12. Summary sheet |

CHAPTER OBJECTIVES

- By the end of this chapter I can interpret bivariate data and explain the idea of regression.
- By the end of this chapter I can draw and interpret a scatter diagram and sketch regression lines.
- By the end of this chapter I can calculate a regression line using Mayer's method.
- By the end of this chapter I can compute sample variance and covariance from raw data.
- By the end of this chapter I can determine the least squares regression lines of y on x and of x on y .
- By the end of this chapter I can compute and interpret the residual sum of squares for each regression line.
- By the end of this chapter I can identify and use the line of best fit.
- By the end of this chapter I can calculate and interpret the product moment correlation coefficient.
- By the end of this chapter I can rank data and compute Spearman's and Kendall's rank correlation coefficients, stating their uses and limitations.

PREREQUISITES

- Mean, variance and standard deviation of a data set (Lower Sixth statistics)
- Equation of a straight line, gradient and intercept (Pure Mathematics)
- Summation notation Σ and manipulation of sums
- Plotting points and reading graphs in the Cartesian plane
- Basic idea of a function and of dependent/independent variables

1 INTRODUCTION TO REGRESSION AND SCATTER DIAGRAMS

Every market day in Bamenda, Mama Ngo notices that the more bags of Irish potatoes she sells, the more money she takes home — but the relationship is never perfectly exact. At the Douala port, engineers observe that heavier trucks tend to consume more fuel per trip, and they want to predict fuel needs from truck weight. During the GCE mock exams, a principal wonders whether students who rank high in Mathematics also rank high in Physics. Can we measure how strongly two quantities are linked, and can we use one to predict the other? This chapter gives you the statistical tools — **regression lines** and **correlation coefficients** — to answer such questions with confidence.

The common feature of all these situations is that two quantities are measured on the same individual or item, and we suspect they move together. Our first task is to organise such information, display it on a diagram, and learn to read the story the diagram tells.

1.1 Bivariate Data

In Lower Sixth statistics you studied univariate data: a single variable measured on each individual (the heights of students, the marks in one test). Real life, however, constantly pairs measurements.

DEFINITION Bivariate data

Bivariate data consist of pairs of observations (x_i, y_i) , where both values are measured on the same individual or item. If there are n individuals, the data set is

$$(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n).$$

The two variables x and y may be of the same nature (two exam marks) or of different natures (rainfall in mm and yield in tonnes).

EXAMPLE Examples of bivariate data

- For each district of the North West Region: x = annual rainfall (mm), y = maize yield (tonnes per hectare).
- For each Upper Sixth student: x = hours of study per week, y = score in the GCE mock examination.
- For each truck leaving the Douala port: x = loaded weight (tonnes), y = fuel consumed (litres).

1.2 Independent and Dependent Variables

When we suspect that one variable explains or drives the other, we give the two variables different roles.

DEFINITION Explanatory and response variables

The **independent variable** (also called the **explanatory** or controlled variable) is the one we believe influences the other, or the one we can fix or observe first. The **dependent variable** (also called the **response** variable) is the one we wish to explain or predict. By universal convention the independent variable is denoted x and the dependent variable y .

Rainfall is the independent variable and crop yield the dependent one: the rain is not caused by the maize! Similarly, hours of study (x) may influence the mock score (y), not the reverse.

WARNING The independent variable goes on the horizontal axis

When plotting, the independent variable x is always placed on the horizontal axis and the dependent variable y on the vertical axis. Examiners penalise reversed axes, and reversing the axes changes the whole regression analysis (we shall see that the line of y on x is not the same as the line of x on y).

1.3 The Idea of Regression

Mama Ngo's takings are not an exact multiple of the bags sold: prices fluctuate, some customers bargain, some bags are heavier. The relationship is statistical, not exact. We therefore ask a weaker but very useful question: on average, what value of y corresponds to a given value of x ?

DEFINITION Regression

Regression is the study of how the average (expected) value of one variable depends on the value of another variable. A regression model allows us to estimate or predict the response variable y from a known value of the explanatory variable x .

The simplest and most important model is **linear regression**, in which we assume the average value of y is a linear function of x :

$$y = a + bx,$$

where b is the **gradient** (the average change in y per unit increase in x) and a is the **intercept** (the value of y predicted when $x = 0$). The observed points will not lie exactly on this line; each observation differs from the line by a small error or residual, and a good line makes these residuals small. Later sections will show how to compute a and b precisely (Mayer's method, then the method of least squares). For now we learn to see the line.

DID YOU KNOW? Why the strange word “regression”?

The term was introduced by Sir Francis Galton in the 19th century while studying the heights of parents and their children. He noticed that very tall parents tended to have children who were tall, but on average less tall than themselves — the children’s heights “regressed” towards the mean of the population. The name stuck, and today “regression” simply means modelling how one variable depends on another.

1.4 The Scatter Diagram

Before any calculation, **always plot the data**. A picture reveals the type and strength of the relationship instantly.

DEFINITION Scatter diagram

A **scatter diagram** (or scatter plot) is a graph of the n points (x_i, y_i) of a bivariate data set, with the independent variable x on the horizontal axis and the dependent variable y on the vertical axis. No lines join the points: each point represents one individual.

METHOD Drawing and interpreting a scatter diagram

1. Identify the variables. Decide which is independent (x) and which is dependent (y).
2. Draw and label the axes. Place x horizontally, y vertically; write the variable names and units.
3. Choose scales. Each axis should cover the range of its data (the axes need not start at 0, and the two scales need not be equal). Use as much of the page as possible.
4. Plot the points. Mark each pair (x_i, y_i) with a small clear dot or cross.
5. Interpret the pattern. Look at the direction (rising or falling), the form (linear or curved) and the strength (points tightly or loosely clustered around an imaginary line).

WORKED EXAMPLE 1 Hours of study and GCE mock score


Ten Upper Sixth students reported their weekly study hours x and obtained the following mock scores y (out of 100):

x	4	6	8	9	11	13	15	17	19	21
y	38	45	50	55	58	63	70	72	80	84

Draw a scatter diagram and comment.

SOLUTION

Solution. Study hours (x) is the independent variable (horizontal axis); the score (y) is the response (vertical axis). We scale the x -axis from 0 to 22 and the y -axis from 30 to 90, then plot the ten points (see the figure below). The points rise steadily from bottom-left to top-right and cluster fairly closely around an imaginary straight line: there is a strong positive linear association between study time and mock score. A straight-line model $y = a + bx$ appears appropriate.



Scatter diagram of mock score against weekly study hours for 10 students: the points suggest a strong positive linear association.

1.5 Interpreting a Scatter Diagram


Three typical patterns occur.

DEFINITION Types of association

- **Positive association:** as x increases, y tends to increase; the points cluster around a line with positive gradient.
- **Negative association:** as x increases, y tends to decrease; the points cluster around a line with negative gradient.
- **No apparent association:** the points are scattered at random; knowing x gives no information about y .

The association is **linear** if the points cluster around a straight line, and **non-linear** if they follow a curve (for example a parabola or an exponential shape). Association is described

as strong when the points lie close to the trend and weak when they are widely spread around it.



Three scatter diagrams: positive association (left), negative association (centre), no apparent association (right).

WARNING Association is not causation

A scatter diagram can show that two variables move together, but it can never prove that one causes the other. Ice-cream sales and drowning incidents may both rise in the same months — not because ice cream causes drowning, but because a hidden third factor (hot weather) drives both. Such a hidden variable is called a **lurking** (or confounding) **variable**. Always ask: is there a plausible mechanism, or could a third factor explain the pattern?

1.6 Sketching Regression Lines by Eye; Outliers

Once a linear pattern is visible, we can draw a first approximation of the regression line by eye: a straight line that follows the trend of the cloud of points, with roughly as many points above it as below it, and passing as close as possible to the **mean point** $(\bar{x}, \bar{y})$, where

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i, \quad \bar{y} = \frac{1}{n} \sum_{i=1}^n y_i.$$

A fundamental fact, which we shall prove when studying least squares, is that **every least squares regression line passes through** $(\bar{x}, \bar{y})$. The mean point is therefore the natural “anchor” of any line drawn by eye.

DEFINITION Outlier

An **outlier** is a point that lies far from the pattern followed by the rest of the data. Outliers may come from recording errors or from genuinely unusual individuals; they can strongly pull a regression line towards themselves, so they must always be spotted on the scatter diagram and investigated before any calculation.

Two regression lines, not one. There are in fact two natural straight lines on the same scatter diagram:

- the **regression line of y on x** , used to predict y from x ; it minimises the vertical deviations of the points from the line;
- the **regression line of x on y** , used to predict x from y ; it minimises the horizontal deviations.

Both lines pass through the mean point $(\bar{x}, \bar{y})$, but they generally have different gradients. The closer the points are to a single straight line, the closer the two lines are to each other; if the correlation is perfect, the two lines coincide.

WORKED EXAMPLE 2 Computing the mean point and sketching both lines

For the ten students of the previous worked example, compute the mean point and describe how to sketch the two regression lines.


SOLUTION

Solution. Summing the values:

$$\sum x_i = 4 + 6 + 8 + 9 + 11 + 13 + 15 + 17 + 19 + 21 = 123, \quad \bar{x} = \frac{123}{10} = 12.3,$$

$$\sum y_i = 38 + 45 + 50 + 55 + 58 + 63 + 70 + 72 + 80 + 84 = 615, \quad \bar{y} = \frac{615}{10} = 61.5.$$

The mean point is $(12.3, 61.5)$. To sketch the line of y on x by eye, draw through $(12.3, 61.5)$ the straight line that best follows the rising cloud of points, balancing vertical deviations. The line of x on y also passes through $(12.3, 61.5)$ but is slightly steeper on the diagram (it balances horizontal deviations). Since the association here is strong, the two lines lie close together, as shown below.



Both regression lines sketched on the same scatter diagram; they cross at the mean point $(\bar{x}, \bar{y})$.

KEY POINTS Key points of this section

- Bivariate data are pairs (x_i, y_i) measured on the same individuals; x is the independent (explanatory) variable, y the dependent (response) variable.
- Regression estimates the average value of one variable from the other; linear regression uses a straight line $y = a + bx$.
- Always begin with a scatter diagram: x on the horizontal axis, y on the vertical axis; read its direction, form and strength.
- Association does not imply causation — beware of lurking variables.
- An outlier lies far from the general pattern and can distort the regression line.
- There are two regression lines (y on x and x on y); both pass through the mean point $(\bar{x}, \bar{y})$ and coincide only when correlation is perfect.

EXERCISE 1 Practice

A farmer in Buea records, for 8 plots, the quantity of fertiliser applied x (kg) and the harvest y (bags of cocoyams):

x	10	15	20	25	30	35	40	45
y	12	15	14	19	22	21	26	28

1. Identify the independent and dependent variables.
2. Draw a scatter diagram.
3. Compute the mean point $(\bar{x}, \bar{y})$ and mark it on your diagram.
4. Sketch by eye a regression line through the mean point and describe the association.
5. The farmer claims: "Fertiliser alone causes the harvest." Comment critically.

SOLUTION Practice

1. x (fertiliser, kg) is independent; y (harvest, bags) is dependent.
2. Plot the eight points with x horizontal (0 to 50) and y vertical (10 to 30); the points rise from bottom-left to top-right.
3. $\sum x_i = 220$, so $\bar{x} = 27.5$; $\sum y_i = 157$, so $\bar{y} = 19.625$. Mean point $(27.5, 19.625)$.

4. A line through (27.5, 19.625) following the rising cloud shows a strong positive linear association.
5. The diagram shows association only. Other factors (rainfall, soil quality, seed variety, weeding) may also influence the harvest; without a controlled experiment, causation cannot be claimed.

In the next section we replace the “by eye” line by a first genuine calculation method: **Mayer’s method**, which splits the data into two groups and uses their mean points to determine the gradient and intercept of the regression line.

2 MAYER’S METHOD FOR THE REGRESSION LINE

In the previous section we saw that a scatter diagram often reveals a linear trend between two variables, and that a regression line is a straight line drawn to summarise that trend. But how do we actually find such a line? Drawing it “by eye” is subjective: two students looking at the same cloud of points will rarely draw the same line. We need a simple, reproducible rule. The first such rule we study is **Mayer’s method**: an intuitive technique, due to the German astronomer Tobias Mayer, which replaces the whole cloud of points by just two representative points and draws the line through them. It is quick, requires only averages, and gives a good first approximation of the regression line. Later in this chapter we shall meet the more precise least squares method; Mayer’s method is the natural stepping stone towards it.

2.1 The Intuition Behind Mayer’s Method

Imagine a farmer near Bafoussam who records, for eight plots of land, the quantity of fertiliser applied (in kg) and the maize yield obtained (in bags). The scatter diagram shows yield increasing with fertiliser, roughly along a straight line. Mayer’s idea is the following:

- A single point is a poor summary of the data, but the mean point (centre of gravity) of a group of points is a good summary of that group.
- If we split the data into a “lower” group (small x -values) and an “upper” group (large x -values), each group has its own mean point.
- A sensible trend line should pass through, or very near, both centres of gravity. Since two points determine a unique straight line, we simply draw the line through the two mean points.

That is the whole method. Let us now state it rigorously.

DEFINITION Group mean points

Let $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$ be a set of n data points, ordered so that $x_1 \leq x_2 \leq \dots \leq x_n$. Split the data into a **lower group** of the first n_1 points and an **upper group** of the remaining n_2

points (with $n_1 + n_2 = n$ and n_1, n_2 as equal as possible). The group mean points are

$$G_1 = (\bar{x}_1, \bar{y}_1) \quad \text{with} \quad \bar{x}_1 = \frac{1}{n_1} \sum_{i=1}^{n_1} x_i, \quad \bar{y}_1 = \frac{1}{n_1} \sum_{i=1}^{n_1} y_i,$$

$$G_2 = (\bar{x}_2, \bar{y}_2) \quad \text{with} \quad \bar{x}_2 = \frac{1}{n_2} \sum_{i=n_1+1}^n x_i, \quad \bar{y}_2 = \frac{1}{n_2} \sum_{i=n_1+1}^n y_i.$$

The **Mayer line** is the straight line through G_1 and G_2 .

WARNING Order by x before splitting!

The split into two groups is always made after ordering the data by the x -values (the explanatory variable), never in the order the data happen to be written in the table, and never by ordering the y -values. Splitting unordered data mixes large and small x -values in the same group, the two mean points come out close together, and the resulting line can have a completely wrong gradient. This is the single most common mistake with Mayer's method in the examination.

2.2 Deriving the Equation of the Mayer Line

Once the two mean points $G_1 = (\bar{x}_1, \bar{y}_1)$ and $G_2 = (\bar{x}_2, \bar{y}_2)$ are known, finding the equation of the line through them is pure Lower Sixth coordinate geometry.

PROPERTY Equation of the Mayer regression line

The Mayer regression line of y on x has gradient

$$b = \frac{\bar{y}_2 - \bar{y}_1}{\bar{x}_2 - \bar{x}_1}$$

and equation

$$y - \bar{y}_1 = b(x - \bar{x}_1), \quad \text{equivalently} \quad y = bx + a \quad \text{with} \quad a = \bar{y}_1 - b\bar{x}_1.$$

(The derivation is a direct application of the two-point form of a straight line and is examinable.)

Derivation. A non-vertical line through $(\bar{x}_1, \bar{y}_1)$ has equation $y - \bar{y}_1 = b(x - \bar{x}_1)$ for some gradient b . Requiring it to pass also through $(\bar{x}_2, \bar{y}_2)$ gives

$$\bar{y}_2 - \bar{y}_1 = b(\bar{x}_2 - \bar{x}_1) \quad \implies \quad b = \frac{\bar{y}_2 - \bar{y}_1}{\bar{x}_2 - \bar{x}_1},$$

which is well defined because $\bar{x}_2 \neq \bar{x}_1$ (the upper group contains strictly larger x -values than the lower group). Substituting back gives the stated equation. ■

METHOD Calculating the regression line by Mayer's method

1. Order the n data points by increasing values of x .
2. Split into two groups as equal as possible: if n is even, each group has $n/2$ points; if n is odd, either exclude the middle point or place it in both groups (state your convention).
3. Compute the mean point $G_1 = (\bar{x}_1, \bar{y}_1)$ of the lower group and $G_2 = (\bar{x}_2, \bar{y}_2)$ of the upper group.
4. Compute the gradient $b = \frac{\bar{y}_2 - \bar{y}_1}{\bar{x}_2 - \bar{x}_1}$.
5. Write the equation $y - \bar{y}_1 = b(x - \bar{x}_1)$ and simplify to the form $y = bx + a$.
6. Check: verify that G_2 satisfies your equation (it must, by construction).

2.3 Handling an Odd Number of Points

When n is even, the split is unambiguous. When n is odd, say $n = 2m + 1$, two conventions are accepted:

- **Exclude the middle point:** the lower group is the first m points, the upper group the last m points; the $(m + 1)$ -th point is ignored.
- **Share the middle point:** the middle point is included in both groups, so each group has $m + 1$ points.

Both conventions appear in textbooks and both are accepted at the GCE, provided you state clearly which one you use and apply it consistently. In this course, unless told otherwise, we **exclude the middle point** when n is odd.

2.4 Worked Example: Fertiliser and Maize Yield

WORKED EXAMPLE 3 Mayer's method on a Cameroonian data set

An agricultural extension officer in the West Region records the quantity of fertiliser x (kg) applied to eight maize plots and the yield y (bags) harvested:

Plot	A	B	C	D	E	F	G	H
x (kg)	10	20	30	40	50	60	70	80
y (bags)	4	6	7	9	10	13	14	15

1. Find the equation of the Mayer regression line of y on x .

2. Use it to predict the yield for $x = 55$ kg and for $x = 120$ kg, commenting on reliability.

SOLUTION

Solution.

1. The data are already ordered by increasing x . Since $n = 8$ is even, each group has 4 points.

Lower group ($x = 10, 20, 30, 40$):

$$\bar{x}_1 = \frac{10 + 20 + 30 + 40}{4} = \frac{100}{4} = 25, \quad \bar{y}_1 = \frac{4 + 6 + 7 + 9}{4} = \frac{26}{4} = 6.5.$$

So $G_1 = (25, 6.5)$.

Upper group ($x = 50, 60, 70, 80$):

$$\bar{x}_2 = \frac{50 + 60 + 70 + 80}{4} = \frac{260}{4} = 65, \quad \bar{y}_2 = \frac{10 + 13 + 14 + 15}{4} = \frac{52}{4} = 13.$$

So $G_2 = (65, 13)$.

Gradient:

$$b = \frac{\bar{y}_2 - \bar{y}_1}{\bar{x}_2 - \bar{x}_1} = \frac{13 - 6.5}{65 - 25} = \frac{6.5}{40} = 0.1625.$$

Equation: $y - 6.5 = 0.1625(x - 25)$, i.e.

$$y = 0.1625x + 2.4375$$

since $a = 6.5 - 0.1625 \times 25 = 6.5 - 4.0625 = 2.4375$.


Check with G_2 : $0.1625 \times 65 + 2.4375 = 10.5625 + 2.4375 = 13 = \bar{y}_2$. ✓

2. For $x = 55$: $y = 0.1625 \times 55 + 2.4375 = 8.9375 + 2.4375 = 11.375 \approx 11.4$ bags. Since 55 kg lies inside the observed range $[10, 80]$, this is interpolation and the prediction is reasonably reliable (the scatter is close to linear).

For $x = 120$: $y = 0.1625 \times 120 + 2.4375 = 19.5 + 2.4375 = 21.9375 \approx 21.9$ bags. Since 120 kg lies far outside the observed range, this is extrapolation: we have no evidence the linear trend continues (in reality, excess fertiliser can even reduce yield), so this prediction is unreliable.

2.5 Illustrating the Method Graphically

The figure below shows the fertiliser–maize data, the two group mean points G_1 and G_2 , and the Mayer line passing through them.



Scatter diagram of the fertiliser-maize data. The dashed line separates the lower group (left) from the upper group (right); the Mayer line passes through the two group mean points G_1 and G_2 .

Notice how the line balances the points: it does not pass through every point, but it passes through the centres of gravity of the two halves of the data, which is exactly the spirit of a trend line.

2.6 Predictions and Reliability

The Mayer line, like any regression line, is used to **predict** y for a given x :

- If the given x lies within the range of observed x -values, the prediction is an **interpolation** and is generally reliable, provided the points are reasonably close to a straight line.
- If the given x lies outside the observed range, the prediction is an **extrapolation** and must be treated with great caution: the linear relationship may not hold beyond the data.

Reliability also depends on the strength of the linear association, which we shall measure later with the product moment correlation coefficient.

WARNING Mayer's line is only an approximation

Mayer's method uses only two summary points; the individual positions of all the other points play no role except through the two group means. Consequently:

- Different splits (odd- n conventions) give slightly different lines.
- The Mayer line is not the least squares regression line studied later in this chapter, and it does not minimise any sum of squared errors.

- In the examination, do not mix the two methods: if the question says “Mayer’s method”, use group mean points; if it says “least squares”, use the formulae $b = \frac{s_{xy}}{s_x^2}$ developed in the next sections.

Mayer’s method should be seen as a quick, transparent first estimate — excellent for checking the plausibility of a least squares result.

KEY POINTS Key points

- Order the data by x , split into two (near-)equal groups, and compute the mean points $G_1 = (\bar{x}_1, \bar{y}_1)$ and $G_2 = (\bar{x}_2, \bar{y}_2)$.
- The Mayer line passes through G_1 and G_2 : gradient $b = \frac{\bar{y}_2 - \bar{y}_1}{\bar{x}_2 - \bar{x}_1}$, equation $y - \bar{y}_1 = b(x - \bar{x}_1)$.
- For odd n , exclude the middle point (or share it between groups) and state your convention.
- Interpolation is reliable; extrapolation is risky.
- Mayer’s method is simple and intuitive but only approximate; the least squares method gives the “best” line in a precise sense.

EXERCISE 2 Mayer’s method with an odd number of points

The table gives the number of years of experience x of nine tailors in a workshop in Buea and the number y of shirts each completes per day.

x (years)	1	2	3	4	5	6	7	8	9
y (shirts)	3	4	5	5	6	7	8	8	9

1. Using Mayer’s method and excluding the middle point, find the equation of the regression line of y on x .
2. Repeat, this time sharing the middle point between the two groups. Comment on the difference.
3. Predict the daily output of a tailor with 6.5 years of experience.

SOLUTION Exercise 1

1. Excluding the middle point (5, 6): lower group $x = 1, 2, 3, 4$; upper group $x = 6, 7, 8, 9$.

$$\bar{x}_1 = \frac{1+2+3+4}{4} = 2.5, \quad \bar{y}_1 = \frac{3+4+5+5}{4} = 4.25; \quad \bar{x}_2 = \frac{6+7+8+9}{4} = 7.5, \quad \bar{y}_2 = \frac{7+8+8+9}{4} = 8.$$

$$b = \frac{8 - 4.25}{7.5 - 2.5} = \frac{3.75}{5} = 0.75, \quad y - 4.25 = 0.75(x - 2.5) \implies y = 0.75x + 2.375.$$

2. Sharing the middle point: each group has 5 points. Lower: $\bar{x}_1 = \frac{15}{5} = 3, \bar{y}_1 = \frac{23}{5} = 4.6$. Upper: $\bar{x}_2 = \frac{35}{5} = 7, \bar{y}_2 = \frac{38}{5} = 7.6$.

$$b = \frac{7.6 - 4.6}{7 - 3} = \frac{3}{4} = 0.75, \quad y - 4.6 = 0.75(x - 3) \implies y = 0.75x + 2.35.$$

The gradient is identical (0.75); only the intercept changes slightly (2.375 vs 2.35). The two conventions give very close lines here because the data are almost perfectly linear. 3. $y = 0.75 \times 6.5 + 2.375 = 7.25$ shirts (interpolation, reliable). With the second line: 7.225 — essentially the same prediction.

DID YOU KNOW? Tobias Mayer, astronomer of the Moon

Tobias Mayer (1723–1762) was a German astronomer famous for his remarkably accurate tables of the Moon's motion. Faced with many imperfect observations, he had the idea of grouping them and averaging within groups to reduce observational error — the ancestor of the method you have just studied. His averaging philosophy later inspired Legendre and Gauss, who developed the method of least squares that we shall meet in the next sections. In Cameroon today, the same averaging spirit is used by agronomists of IRAD (Institute of Agricultural Research for Development) when summarising field trial data before fitting more sophisticated models.

3 CORRELATION, SAMPLE VARIANCE AND COVARIANCE

In the previous section we met Mayer's method, a quick way of fitting a straight line to a scatter diagram by splitting the data into two groups. Mayer's method gives us a line, but it tells us nothing about how well the points actually cluster around a line. Two data sets can produce almost identical Mayer lines, yet in one the points hug the line tightly while in the other they are scattered all over the plane. We therefore need a way to measure the strength and direction of a linear relationship. That is the purpose of this section: we introduce the idea of **correlation**, and we build the two fundamental tools on which everything else in this chapter rests — the **sample variance** and the **sample covariance**.

3.1 Introduction to Correlation

EXPERIMENT Observing how two variables move together

A teacher in Buea records, for 10 students, the number of hours x spent revising Mathematics each week and the mark y (out of 100) obtained in the mock examination. Plotting the points (x, y) , she notices that the points rise from the bottom-left to the top-right of the diagram: students who revise more tend to score higher. A second teacher records the price x of a second-hand car and its age y ; the points fall from top-left to bottom-right. A third records shoe size x and History mark y ; the points form a shapeless cloud.


These three situations illustrate the three basic types of correlation.

DEFINITION Correlation

Correlation describes the extent to which two variables x and y are linearly related.

- There is positive correlation when large values of x tend to occur with large values of y (the points rise from left to right).
- There is negative correlation when large values of x tend to occur with small values of y (the points fall from left to right).
- There is zero (or no) correlation when there is no apparent linear tendency at all.

Correlation is said to be strong when the points lie close to a straight line, and weak when they are widely scattered around it.



Scatter diagrams showing strong positive, weak positive and negative correlation.

WARNING Correlation is not causation

A strong correlation between two variables does not prove that one causes the other. Ice-cream sales and drowning incidents are positively correlated, but neither causes the other: both increase in hot weather. Always ask whether a hidden third variable could explain the observed association.

3.2 Sample Variance

To measure correlation numerically we first need to measure how spread out each variable is on its own. You already know the mean $\bar{x} = \frac{1}{n} \sum x$. The spread about the mean is measured by the variance.

DEFINITION Sample variance

For a sample of n values $x_1, x_2, \dots, x_n$ with mean $\bar{x}$, the **sample variance** of x is

$$s_{xx} = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2 = \frac{1}{n} \sum_{i=1}^n x_i^2 - \bar{x}^2.$$

Similarly, for values $y_1, \dots, y_n$ with mean $\bar{y}$,

$$s_{yy} = \frac{1}{n} \sum_{i=1}^n (y_i - \bar{y})^2 = \frac{1}{n} \sum_{i=1}^n y_i^2 - \bar{y}^2.$$

The positive square roots $s_x = \sqrt{s_{xx}}$ and $s_y = \sqrt{s_{yy}}$ are the sample standard deviations.

The second form of each formula is the computational form; it is far quicker with a calculator. Let us prove the equivalence for s_{xx} (the proof for s_{yy} is identical).

PROPERTY Equivalence of the two forms of the variance

$$\frac{1}{n} \sum (x_i - \bar{x})^2 = \frac{1}{n} \sum x_i^2 - \bar{x}^2.$$

Proof. Expanding the square and using $\sum x_i = n\bar{x}$:

$$\begin{aligned} \sum_{i=1}^n (x_i - \bar{x})^2 &= \sum_{i=1}^n (x_i^2 - 2\bar{x}x_i + \bar{x}^2) \\ &= \sum x_i^2 - 2\bar{x} \sum x_i + n\bar{x}^2 \\ &= \sum x_i^2 - 2\bar{x}(n\bar{x}) + n\bar{x}^2 \\ &= \sum x_i^2 - n\bar{x}^2. \end{aligned}$$

Dividing by n gives $\frac{1}{n} \sum (x_i - \bar{x})^2 = \frac{1}{n} \sum x_i^2 - \bar{x}^2$. ■

WARNING Do not confuse $\sum x^2$ with $(\sum x)^2$

This is the single most common error in this chapter. $\sum x^2$ means: square each value, then add. $(\sum x)^2$ means: add the values, then square the total. For the data 1, 2, 3: $\sum x^2 = 1 + 4 + 9 = 14$ but $(\sum x)^2 = 6^2 = 36$. Keep the two quantities in clearly separate columns of your table.

3.3 Sample Covariance

Variance measures the spread of one variable. To describe how two variables move together, we need a mixed quantity.

EXPERIMENT The idea behind covariance

For each data point, compute the deviations $x - \bar{x}$ and $y - \bar{y}$. If x and y tend to be above their means together (and below together), the products $(x - \bar{x})(y - \bar{y})$ are mostly positive, so their average is positive. If one variable tends to be above its mean when the other is below, the products are mostly negative. If there is no pattern, positive and negative products cancel out. The average of these products is the covariance.

DEFINITION Sample covariance

For n pairs of data (x_i, y_i) with means $\bar{x}$ and $\bar{y}$, the **sample covariance** of x and y is

$$s_{xy} = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y}) = \frac{1}{n} \sum_{i=1}^n x_i y_i - \bar{x} \bar{y}.$$

PROPERTY Equivalence of the two forms of the covariance

$$\frac{1}{n} \sum (x_i - \bar{x})(y_i - \bar{y}) = \frac{1}{n} \sum x_i y_i - \bar{x} \bar{y}.$$

Proof. Expand the product and use $\sum x_i = n\bar{x}$, $\sum y_i = n\bar{y}$:

$$\begin{aligned} \sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y}) &= \sum_{i=1}^n (x_i y_i - \bar{x} y_i - \bar{y} x_i + \bar{x} \bar{y}) \\ &= \sum x_i y_i - \bar{x} \sum y_i - \bar{y} \sum x_i + n \bar{x} \bar{y} \\ &= \sum x_i y_i - \bar{x} (n\bar{y}) - \bar{y} (n\bar{x}) + n \bar{x} \bar{y} \\ &= \sum x_i y_i - n \bar{x} \bar{y}. \end{aligned}$$

Dividing by n gives the result. ■

KEY POINTS Interpreting the sign of the covariance

- $s_{xy} > 0$: positive linear association (points rise left to right).
- $s_{xy} < 0$: negative linear association (points fall left to right).
- $s_{xy} \approx 0$: little or no linear association.

Note also that s_{xx} is simply the covariance of x with itself, and covariance is symmetric: $s_{xy} = s_{yx}$.

WARNING Covariance depends on the units of measurement

The covariance is not a standardised measure of strength. If x is height measured in centimetres and y is mass in kilograms, s_{xy} is measured in $\text{cm} \cdot \text{kg}$. Measuring the same heights in metres multiplies the covariance by $\frac{1}{100}$, without the underlying relationship changing at all! Consequently a covariance of 50 is not automatically “stronger” than a covariance of 5 — the units must be taken into account. This is exactly why, in a later section, we shall divide by the standard deviations to obtain the unit-free product moment correlation coefficient. For now, use the covariance mainly for its sign and as a building block.

3.4 Setting Out the Calculations in a Table**METHOD** Computing s_{xx} , s_{yy} and s_{xy} from a table of values

1. Draw a table with columns x , y , x^2 , y^2 , xy and complete one row per data pair.
2. Add a final row of totals: $\sum x$, $\sum y$, $\sum x^2$, $\sum y^2$, $\sum xy$.
3. Compute the means: $\bar{x} = \frac{\sum x}{n}$ and $\bar{y} = \frac{\sum y}{n}$.
4. Apply the computational formulae:

$$s_{xx} = \frac{\sum x^2}{n} - \bar{x}^2, \quad s_{yy} = \frac{\sum y^2}{n} - \bar{y}^2, \quad s_{xy} = \frac{\sum xy}{n} - \bar{x}\bar{y}.$$

5. Interpret the sign of s_{xy} and compare the magnitudes with the spread of the data.

WORKED EXAMPLE 4 Revision time and mock examination marks

Six students recorded their weekly revision time x (hours) and their mock mark y (out of 20):

x	2	4	5	7	8	10
y	6	8	9	12	13	15

Calculate s_{xx} , s_{yy} and s_{xy} , and comment on the relationship.

SOLUTION

Step 1: the table of values.

x	y	x^2	y^2	xy
2	6	4	36	12
4	8	16	64	32
5	9	25	81	45
7	12	49	144	84
8	13	64	169	104
10	15	100	225	150
$\Sigma x = 36$	$\Sigma y = 63$	$\Sigma x^2 = 258$	$\Sigma y^2 = 719$	$\Sigma xy = 427$

Step 2: the means. With $n = 6$:

$$\bar{x} = \frac{36}{6} = 6, \quad \bar{y} = \frac{63}{6} = 10.5.$$

Step 3: the variances and covariance.

$$s_{xx} = \frac{258}{6} - 6^2 = 43 - 36 = 7,$$

$$s_{yy} = \frac{719}{6} - 10.5^2 = 119.833... - 110.25 = \frac{115}{12} \approx 9.583,$$

$$s_{xy} = \frac{427}{6} - (6)(10.5) = 71.166... - 63 = \frac{49}{6} \approx 8.167.$$

Step 4: interpretation. Since $s_{xy} > 0$, there is a positive linear association: more revision time tends to go with a higher mark. Note that s_{xy} is measured in “hour-marks”, so its size alone cannot tell us how strong the link is — we shall standardise it later. As a check, $s_{xy}^2 = \left(\frac{49}{6}\right)^2 \approx 66.7$ is less than $s_{xx}s_{yy} = 7 \times 9.583 \approx 67.1$, which must always hold.

PROPERTY The Cauchy–Schwarz inequality for data

For any bivariate data set, $s_{xy}^2 \leq s_{xx}s_{yy}$, with equality exactly when all the points lie on a single straight line. This guarantees that the correlation coefficient we shall define later always lies between -1 and 1 .

EXERCISE 3 Practice

Eight candidates in a Further Mathematics mock obtained the following marks in Paper 1 (x) and Paper 2 (y):

x	12	15	18	20	24	27	30	34
y	10	14	15	19	20	25	26	31

1. Set out a full table of values and find $\sum x$, $\sum y$, $\sum x^2$, $\sum y^2$, $\sum xy$.
2. Calculate s_{xx} , s_{yy} and s_{xy} .
3. Verify that $s_{xy}^2 \leq s_{xx}s_{yy}$ and state what the sign of s_{xy} tells you.

SOLUTION Exercise 1

1. The completed table gives

$$\sum x = 180, \quad \sum y = 160, \quad \sum x^2 = 4454, \quad \sum y^2 = 3624, \quad \sum xy = 3969.$$

2. With $n = 8$: $\bar{x} = \frac{180}{8} = 22.5$ and $\bar{y} = \frac{160}{8} = 20$. Hence

$$s_{xx} = \frac{4454}{8} - 22.5^2 = 556.75 - 506.25 = 50.5,$$

$$s_{yy} = \frac{3624}{8} - 20^2 = 453 - 400 = 53,$$

$$s_{xy} = \frac{3969}{8} - (22.5)(20) = 496.125 - 450 = 46.125.$$

3. $s_{xy}^2 \approx 2127.5$ and $s_{xx}s_{yy} = 50.5 \times 53 = 2676.5$, so indeed $s_{xy}^2 \leq s_{xx}s_{yy}$. Since $s_{xy} > 0$, performance in the two papers is positively associated: candidates who do well in Paper 1 tend to do well in Paper 2.

KEY POINTS Key points of this section

- Correlation describes the strength and direction of a linear relationship: positive, negative or zero; strong or weak.
- Sample variances: $s_{xx} = \frac{\sum x^2}{n} - \bar{x}^2$, $s_{yy} = \frac{\sum y^2}{n} - \bar{y}^2$.
- Sample covariance: $s_{xy} = \frac{\sum xy}{n} - \bar{x}\bar{y}$; its sign gives the direction of the association.
- Always work from a table with columns x , y , x^2 , y^2 , xy and a row of totals.
- Covariance carries units, so its magnitude alone cannot measure strength — this motivates the standardised correlation coefficient of the next sections.

4 LEAST SQUARES REGRESSION LINES

In the previous section, we introduced the sample variance s_{xx} , s_{yy} and the sample covariance s_{xy} as measures of spread and linear association. We also defined the product moment correlation coefficient $r = \frac{s_{xy}}{\sqrt{s_{xx}s_{yy}}}$ which quantifies the strength of a linear relationship. We now put these quantities to work: given a scatter diagram suggesting a linear trend, how do we find the best straight line to summarise it? The answer is the **least squares regression line**. There are two such lines, depending on which variable we wish to predict.


4.1 The Least Squares Principle

Consider a set of n data pairs (x_i, y_i) . If we draw a candidate line $y = a + bx$, the vertical distance from the point (x_i, y_i) to the line is the **residual**

$$e_i = y_i - (a + bx_i).$$

Some residuals are positive (point above the line), some negative (point below). A good line should make these residuals “small overall”. The **least squares criterion** chooses a and b to minimise the **Residual Sum of Squares (RSS)**:

$$\text{RSS}_{y|x} = \sum_{i=1}^n e_i^2 = \sum_{i=1}^n [y_i - (a + bx_i)]^2.$$



Scatter diagram with vertical residuals (green dashed) for the least squares line of y on x . The line passes through the mean point $(\bar{x}, \bar{y})$.

DEFINITION Least Squares Regression Line of y on x

The line $y = a + bx$ that minimises the sum of squared vertical deviations $\sum (y_i - a - bx_i)^2$ is

called the least squares regression line of y on x . Its coefficients are

$$b = \frac{s_{xy}}{s_{xx}}, \quad a = \bar{y} - b\bar{x},$$

where $s_{xx} = \sum(x_i - \bar{x})^2$, $s_{xy} = \sum(x_i - \bar{x})(y_i - \bar{y})$, and $(\bar{x}, \bar{y})$ is the mean point.

PROPERTY The Mean Point Property

The least squares regression line of y on x always passes through the mean point $(\bar{x}, \bar{y})$.

Proof. Substitute $x = \bar{x}$ into the equation $y = a + bx$ with $a = \bar{y} - b\bar{x}$:

$$y = (\bar{y} - b\bar{x}) + b\bar{x} = \bar{y}.$$

Hence $(\bar{x}, \bar{y})$ lies on the line. This is a direct consequence of the normal equations derived from minimising RSS. $\square$

PROPERTY Residual Sum of Squares for y on x

The minimum value of the residual sum of squares (RSS) for the regression line of y on x is

$$\text{RSS}_{y|x} = s_{yy} - \frac{s_{xy}^2}{s_{xx}} = s_{yy}(1 - r^2).$$

Proof. Using the normal equations $\sum y_i = na + b \sum x_i$ and $\sum x_i y_i = a \sum x_i + b \sum x_i^2$, one can show

$$\sum (y_i - a - bx_i)^2 = \sum y_i^2 - a \sum y_i - b \sum x_i y_i = s_{yy} - bs_{xy} = s_{yy} - \frac{s_{xy}^2}{s_{xx}}.$$

Since $r^2 = s_{xy}^2 / (s_{xx}s_{yy})$, the result follows. $\square$

4.2 Regression Line of x on y

If our goal is to predict x from a known value of y , we should minimise the sum of squared horizontal deviations. This leads to the **regression line of x on y** .

DEFINITION Least Squares Regression Line of x on y

The line $x = a' + b'y$ that minimises the sum of squared horizontal deviations $\sum(x_i - a' - b'y_i)^2$

is called the least squares regression line of x on y . Its coefficients are

$$b' = \frac{s_{xy}}{s_{yy}}, \quad a' = \bar{x} - b'\bar{y}.$$

PROPERTY Both Lines Pass Through $(\bar{x}, \bar{y})$

Both the regression line of y on x and the regression line of x on y pass through the mean point $(\bar{x}, \bar{y})$.

Proof. For $x = a' + b'y$, substitute $y = \bar{y}$:

$$x = (\bar{x} - b'\bar{y}) + b'\bar{y} = \bar{x}.$$

Thus $(\bar{x}, \bar{y})$ satisfies both equations. □

PROPERTY Residual Sum of Squares for x on y

The minimum value of the residual sum of squares (RSS) for the regression line of x on y is

$$\text{RSS}_{x|y} = s_{xx} - \frac{s_{xy}^2}{s_{yy}} = s_{xx}(1 - r^2).$$

4.3 Choosing the Appropriate Line

METHOD When to Use Which Regression Line

1. Identify the independent (explanatory) variable and the dependent (response) variable.
2. If x is independent and we want to predict y , use the regression line of y on x : $y = a + bx$.
3. If y is independent and we want to predict x , use the regression line of x on y : $x = a' + b'y$.
4. Never use the y on x line to predict x from y (or vice versa) by simply rearranging the equation. The two lines are different unless $r = \pm 1$ (perfect correlation).

WARNING Wrong Line for Prediction

A very common error in examinations is to compute the line $y = a + bx$ and then, when asked to estimate x for a given y , rearrange it to $x = (y - a)/b$. This is incorrect. You must compute the separate line $x = a' + b'y$. The two lines coincide only when all points lie exactly on a straight line ($r = \pm 1$).

4.4 Geometric Interpretation: The Two Lines and the Correlation

Since both lines pass through $(\bar{x}, \bar{y})$, they intersect at that point. Their slopes are


$$\text{slope of } y \text{ on } x : b = \frac{s_{xy}}{s_{xx}}, \quad \text{slope of } x \text{ on } y : \frac{1}{b'} = \frac{s_{yy}}{s_{xy}}.$$

The product of the slopes is

$$b \cdot \frac{1}{b'} = \frac{s_{xy}^2}{s_{xx}s_{yy}} = r^2.$$

Thus the angle between the two lines is directly related to $|r|$:

- If $|r| = 1$ (perfect correlation), the two lines coincide ($b = 1/b'$).
- If $r = 0$ (no linear correlation), the lines are perpendicular: y on x is horizontal ($b = 0$), x on y is vertical ($b' = 0$).
- For $0 < |r| < 1$, the lines are distinct; the smaller $|r|$, the larger the angle between them.



Both least squares regression lines intersect at the mean point $(\bar{x}, \bar{y})$. The angle θ between them reflects the correlation strength: here $r = 0.9$, so the lines are close.

4.5 Computing the Lines from Summary Statistics

In examinations, you are often given summary statistics $\sum x$, $\sum y$, $\sum x^2$, $\sum y^2$, $\sum xy$ and n . The following method is efficient and avoids rounding errors until the final step.

METHOD Finding Regression Coefficients from Summary Statistics

Given $n, \sum x, \sum y, \sum x^2, \sum y^2, \sum xy$:

1. Compute the corrected sums of squares and products:

$$s_{xx} = \sum x^2 - \frac{(\sum x)^2}{n}, \quad s_{yy} = \sum y^2 - \frac{(\sum y)^2}{n}, \quad s_{xy} = \sum xy - \frac{\sum x \sum y}{n}.$$

2. Compute the means: $\bar{x} = \frac{\sum x}{n}, \bar{y} = \frac{\sum y}{n}$.

3. For y on x : $b = \frac{s_{xy}}{s_{xx}}, a = \bar{y} - b\bar{x}$.

4. For x on y : $b' = \frac{s_{xy}}{s_{yy}}, a' = \bar{x} - b'\bar{y}$.

5. Write the equations clearly: $y = a + bx$ and $x = a' + b'y$.

4.6 Interpolation and Extrapolation

Once a regression line is established, we use it to estimate values.

- **Interpolation:** Estimating y for an x value within the range of the observed x data. This is generally reliable if the linear model fits well.
- **Extrapolation:** Estimating y for an x value outside the range of the observed data. This is **dangerous** because the linear relationship may not hold beyond the observed interval.

WARNING Dangers of Extrapolation

Extrapolation assumes the linear trend continues indefinitely. In reality, relationships often curve, saturate, or reverse outside the observed range. For example, a regression line fitted to the height of a child from age 5 to 15 would absurdly predict the child growing to 5 metres tall by age 50. Always state the range of x values used to fit the line and warn against predictions far outside this range.

4.7 Worked Example: Computing Both Regression Lines and RSS**WORKED EXAMPLE 5** Finding both regression lines and RSS from raw data

The table below shows the number of hours of revision (x) and the examination mark (y) for 6 students.

x (hours)	10	12	15	18	20	25
y (mark)	45	52	60	68	72	80

Find (a) the equation of the regression line of y on x , (b) the equation of the regression line of x on y , (c) the residual sum of squares for each line, (d) estimate the mark for a student who revises for 16 hours, (e) estimate the revision hours for a student who scored 65.

SOLUTION

Solution.

First, compute the summary statistics. We add columns for x^2 , y^2 , xy .

x	y	x^2	y^2	xy
10	45	100	2025	450
12	52	144	2704	624
15	60	225	3600	900
18	68	324	4624	1224
20	72	400	5184	1440
25	80	625	6400	2000
$\Sigma=100$	$\Sigma=377$	$\Sigma=1818$	$\Sigma=24537$	$\Sigma=6638$

$n = 6$.

Corrected sums:

$$s_{xx} = \sum x^2 - \frac{(\sum x)^2}{n} = 1818 - \frac{10000}{6} = 1818 - 1666.\bar{6} = 151.\bar{3} = \frac{454}{3} \approx 151.333,$$

$$s_{yy} = \sum y^2 - \frac{(\sum y)^2}{n} = 24537 - \frac{142129}{6} = 24537 - 23688\frac{1}{6} = 848\frac{5}{6} = \frac{5093}{6} \approx 848.833,$$

$$s_{xy} = \sum xy - \frac{\sum x \sum y}{n} = 6638 - \frac{100 \times 377}{6} = 6638 - 6283\frac{1}{3} = 354\frac{2}{3} = \frac{1064}{3} \approx 354.667.$$

Means:

$$\bar{x} = \frac{100}{6} = \frac{50}{3} \approx 16.667, \quad \bar{y} = \frac{377}{6} \approx 62.833.$$

(a) Regression line of y on x :

$$b = \frac{s_{xy}}{s_{xx}} = \frac{1064/3}{454/3} = \frac{1064}{454} = \frac{532}{227} \approx 2.3436.$$

$$a = \bar{y} - b\bar{x} = \frac{377}{6} - \frac{532}{227} \times \frac{50}{3} = \frac{377}{6} - \frac{26600}{681} = \frac{377 \times 227 - 53200}{1362}.$$

$377 \times 227 = 85579$. $85579 - 53200 = 32379$. $a = \frac{32379}{1362} = \frac{10793}{454} \approx 23.773$. Equation:

$$y = 23.77 + 2.34x \text{ (rounded to 2 d.p.)}$$

(b) Regression line of x on y :

$$b' = \frac{s_{xy}}{s_{yy}} = \frac{1064/3}{5093/6} = \frac{1064}{3} \times \frac{6}{5093} = \frac{2128}{5093} \approx 0.4178.$$

$$a' = \bar{x} - b'\bar{y} = \frac{50}{3} - \frac{2128}{5093} \times \frac{377}{6} = \frac{50}{3} - \frac{802256}{30558} \approx 16.667 - 26.253 = -9.586.$$

Equation: $x = -9.59 + 0.418y$ (rounded to 2 d.p.).

(c) Residual Sum of Squares: For y on x :

$$RSS_{y|x} = s_{yy} - \frac{s_{xy}^2}{s_{xx}} = \frac{5093}{6} - \frac{(1064/3)^2}{454/3} = \frac{5093}{6} - \frac{1064^2}{3 \times 454} = \frac{5093}{6} - \frac{1132096}{1362} = \frac{5093 \times 227 - 1132096}{1362} = \frac{1156111 - 1132096}{1362} = \frac{24015}{1362} \approx 17.63$$

For x on y :

$$RSS_{x|y} = s_{xx} - \frac{s_{xy}^2}{s_{yy}} = \frac{454}{3} - \frac{(1064/3)^2}{5093/6} = \frac{454}{3} - \frac{1064^2 \times 2}{3 \times 5093} = \frac{454}{3} - \frac{2264192}{15279} = \frac{454 \times 5093 - 2264192}{15279} = \frac{2312222 - 2264192}{15279} = \frac{48030}{15279} \approx 3.15$$

(d) Estimate mark for $x = 16$ hours (interpolation, since $10 \leq 16 \leq 25$): Use y on x line: $y = 23.77 + 2.34 \times 16 = 23.77 + 37.44 = 61.21 \approx \boxed{61}$.

(e) Estimate revision hours for $y = 65$ (interpolation, since $45 \leq 65 \leq 80$): Use x on y line: $x = -9.59 + 0.418 \times 65 = -9.59 + 27.17 = 17.58 \approx \boxed{17.6 \text{ hours}}$.

Check: Both lines should pass through $(\bar{x}, \bar{y}) \approx (16.67, 62.83)$. y on x : $23.77 + 2.34 \times 16.67 \approx 23.77 + 39.01 = 62.78 \approx 62.83$ (rounding). x on y : $-9.59 + 0.418 \times 62.83 \approx -9.59 + 26.26 = 16.67$. Correct.

4.8 Summary of Key Formulas

KEY POINTS Least Squares Regression Lines

- Line of y on x : $y = a + bx$ with $b = \frac{s_{xy}}{s_{xx}}$, $a = \bar{y} - b\bar{x}$.
- Line of x on y : $x = a' + b'y$ with $b' = \frac{s_{xy}}{s_{yy}}$, $a' = \bar{x} - b'\bar{y}$.
- Both lines pass through $(\bar{x}, \bar{y})$.
- Product of slopes: $b \times \frac{1}{b'} = r^2$.
- RSS for y on x : $RSS_{y|x} = s_{yy} - \frac{s_{xy}^2}{s_{xx}} = s_{yy}(1 - r^2)$.

- RSS for x on y : $\text{RSS}_{x|y} = s_{xx} - \frac{s_{xy}^2}{s_{yy}} = s_{xx}(1 - r^2)$.
- Use y on x to predict y from x ; use x on y to predict x from y .
- Interpolation (within data range) is reliable; extrapolation (outside data range) is risky.

EXERCISE 4 Practice

The following data gives the age (x years) and the systolic blood pressure (y mmHg) of 8 men.

x	35	40	45	50	55	60	65	70
y	120	125	132	138	145	150	158	165

1. Calculate s_{xx} , s_{yy} , s_{xy} .
2. Find the equation of the regression line of y on x .
3. Find the equation of the regression line of x on y .
4. Calculate the residual sum of squares for each line.
5. Estimate the blood pressure of a man aged 52.
6. Estimate the age of a man with blood pressure 140.
7. Would it be sensible to use the line to estimate the blood pressure of a 20-year-old? Explain.

SOLUTION Exercise

First compute the necessary sums: $n = 8$, $\sum x = 35 + 40 + 45 + 50 + 55 + 60 + 65 + 70 = 420$, $\sum y = 120 + 125 + 132 + 138 + 145 + 150 + 158 + 165 = 1133$, $\sum x^2 = 1225 + 1600 + 2025 + 2500 + 3025 + 3600 + 4225 + 4900 = 23100$, $\sum y^2 = 14400 + 15625 + 17424 + 19044 + 21025 + 22500 + 24964 + 27225 = 162207$, $\sum xy = 4200 + 5000 + 5940 + 6900 + 7975 + 9000 + 10270 + 11550 = 60835$.

1. Corrected sums: $s_{xx} = 23100 - \frac{420^2}{8} = 23100 - 22050 = 1050$. $s_{yy} = 162207 - \frac{1133^2}{8} = 162207 - \frac{1283689}{8} = 162207 - 160461.125 = 1745.875$. $s_{xy} = 60835 - \frac{420 \times 1133}{8} = 60835 - \frac{475860}{8} = 60835 - 59482.5 = 1352.5$.

Means: $\bar{x} = \frac{420}{8} = 52.5$, $\bar{y} = \frac{1133}{8} = 141.625$.

2. Regression line of y on x : $b = \frac{s_{xy}}{s_{xx}} = \frac{1352.5}{1050} = \frac{541}{420} \approx 1.2881$. $a = \bar{y} - b\bar{x} = 141.625 - \frac{541}{420} \times 52.5 = 141.625 - \frac{541}{8} = 141.625 - 67.625 = 74.0$. Equation: $y = 74.0 + 1.29x$ (to 3 s.f.).

3. Regression line of x on y : $b' = \frac{s_{xy}}{s_{yy}} = \frac{1352.5}{1745.875} = \frac{10820}{13967} \approx 0.7747$. $a' = \bar{x} - b'\bar{y} = 52.5 - \frac{10820}{13967} \times 141.625 \approx 52.5 - 109.71 = -57.21$. Equation: $x = -57.2 + 0.775y$ (to 3 s.f.).

4. Residual Sum of Squares: $\text{RSS}_{y|x} = s_{yy} - \frac{s_{xy}^2}{s_{xx}} = 1745.875 - \frac{1352.5^2}{1050} = 1745.875 - 1741.25 = 4.625$. $\text{RSS}_{x|y} = s_{xx} - \frac{s_{xy}^2}{s_{yy}} = 1050 - \frac{1352.5^2}{1745.875} = 1050 - 1047.18 = 2.82$ (approx).

5. $x = 52$ (interpolation, within range 35–70): Use y on x line. $y = 74.0 + 1.2881 \times 52 =$

$$74.0 + 66.98 = 140.98 \approx \boxed{141 \text{ mmHg}}.$$

6. $y = 140$ (interpolation, within range 120–165): Use x on y line. $x = -57.21 + 0.7747 \times 140 = -57.21 + 108.46 = 51.25 \approx \boxed{51 \text{ years}}.$

7. No. $x = 20$ is far outside the data range (35–70). Extrapolation assumes the linear trend continues downwards, which is biologically unrealistic (blood pressure would become negative at very young ages). The model is only validated for the observed age range.

5 RESIDUAL SUM OF SQUARES AND THE LINE OF BEST FIT

In the previous section we derived the two least squares regression lines: the line of y on x , $y = a + bx$ with $b = \frac{s_{xy}}{s_{xx}}$, and the line of x on y , $x = a' + b'y$ with $b' = \frac{s_{xy}}{s_{yy}}$. In each case the slope and intercept were obtained by minimising a sum of squares. In this section we make that idea precise: we define the **residuals** of a fitted line, measure the total error of the fit through the **residual sum of squares** (RSS), and show that the least squares line is, in a very exact sense, the **line of best fit**. This gives us a practical tool for comparing different fitted lines — for example Mayer's line and the least squares line — and a first bridge towards the correlation coefficient of the next section.

5.1 Residuals: Measuring the Error at Each Point


Suppose we have n data points $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$ and we have fitted a regression line of y on x , $\hat{y} = a + bx$. For a given value x_i , the line predicts the value $\hat{y}_i = a + bx_i$, while the observed value is y_i . These two values generally differ, and the difference tells us how well the line performs at that particular point.

DEFINITION Residual for the regression of y on x

For the regression line of y on x , $\hat{y} = a + bx$, the **residual** at the point (x_i, y_i) is the vertical signed distance from the observed point to the line:

$$e_i = y_i - \hat{y}_i = y_i - (a + bx_i).$$

A positive residual means the point lies above the line; a negative residual means it lies below the line.



Residuals $e_i = y_i - \hat{y}_i$ shown as vertical segments from each data point to the fitted line of y on x .

For the regression line of x on y , $\hat{x} = a' + b'y$, the roles are reversed: we predict x from y , and the residual is the horizontal signed distance

$$d_i = x_i - \hat{x}_i = x_i - (a' + b'y_i).$$

WARNING Vertical versus horizontal residuals

Residuals for the regression of y on x are vertical distances; residuals for the regression of x on y are horizontal distances. Never mix the two: the two lines answer two different questions ("predict y given x " and "predict x given y "), and their residuals measure errors in different directions. This is precisely why the two regression lines are generally different.

5.2 The Residual Sum of Squares (RSS)

A single residual tells us about one point. To judge the global quality of a fitted line we need to combine all the residuals into a single number. Simply adding them is useless: positive and negative residuals cancel (indeed, for the least squares line one can show that $\sum e_i = 0$ exactly). As with the definition of variance, the natural remedy is to square each residual before adding.

DEFINITION Residual sum of squares

For the regression line of y on x , the **residual sum of squares** is

$$\text{RSS}_{y \text{ on } x} = \sum_{i=1}^n e_i^2 = \sum_{i=1}^n (y_i - \hat{y}_i)^2.$$

For the regression line of x on y ,

$$\text{RSS}_{x \text{ on } y} = \sum_{i=1}^n d_i^2 = \sum_{i=1}^n (x_i - \hat{x}_i)^2.$$

The RSS is always non-negative, and it equals 0 only when every residual is zero, that is, when every point lies exactly on the line. A small RSS means the points cluster tightly around the line; a large RSS means the points are widely scattered around it.

5.3 Compact Formulae for the RSS of the Least Squares Lines

Computing each residual individually is tedious for large data sets. Fortunately, for the least squares lines there are closed formulae in terms of the quantities s_{xx} , s_{yy} and s_{xy} already computed when fitting the line. Recall our convention (used throughout this chapter):

$$s_{xx} = \sum (x_i - \bar{x})^2, \quad s_{yy} = \sum (y_i - \bar{y})^2, \quad s_{xy} = \sum (x_i - \bar{x})(y_i - \bar{y}).$$

PROPERTY RSS of the least squares lines

For the least squares regression line of y on x ,

$$\text{RSS}_{y \text{ on } x} = s_{yy} - \frac{(s_{xy})^2}{s_{xx}}.$$

For the least squares regression line of x on y ,

$$\text{RSS}_{x \text{ on } y} = s_{xx} - \frac{(s_{xy})^2}{s_{yy}}.$$

(The proof is instructive but is not examinable at the GCE; the formulae themselves must be known and used fluently.)

Derivation (for the line of y on x).

The least squares line passes through $(\bar{x}, \bar{y})$, so $\hat{y}_i - \bar{y} = b(x_i - \bar{x})$ with $b = s_{xy}/s_{xx}$. Write $u_i = x_i - \bar{x}$ and $v_i = y_i - \bar{y}$. Then

$$e_i = y_i - \hat{y}_i = (y_i - \bar{y}) - (\hat{y}_i - \bar{y}) = v_i - bu_i.$$

Hence

$$\begin{aligned} \text{RSS} &= \sum (v_i - bu_i)^2 = \sum v_i^2 - 2b \sum u_i v_i + b^2 \sum u_i^2 \\ &= s_{yy} - 2b s_{xy} + b^2 s_{xx}. \end{aligned}$$

Substituting $b = \frac{s_{xy}}{s_{xx}}$:

$$\text{RSS} = s_{yy} - 2 \frac{s_{xy}}{s_{xx}} s_{xy} + \frac{(s_{xy})^2}{s_{xx}^2} s_{xx} = s_{yy} - \frac{(s_{xy})^2}{s_{xx}}.$$

The derivation for the line of x on y is identical with the roles of x and y exchanged. ■

METHOD Computing the RSS without listing the residuals

1. Compute $\bar{x}, \bar{y}$, then s_{xx}, s_{yy} and s_{xy} (the same table of sums $\sum x, \sum y, \sum x^2, \sum y^2, \sum xy$ used for the regression line).
2. For the line of y on x , compute $\text{RSS} = s_{yy} - \frac{(s_{xy})^2}{s_{xx}}$.
3. For the line of x on y , compute $\text{RSS} = s_{xx} - \frac{(s_{xy})^2}{s_{yy}}$.
4. Check: the RSS must satisfy $0 \leq \text{RSS}_{y \text{ on } x} \leq s_{yy}$. A negative value signals an arithmetic error.

WARNING Units and the meaning of "small"

The RSS of the line of y on x is measured in squared units of y , and the RSS of the line of x on y in squared units of x . The two values are not directly comparable with each other. Moreover, "small" is relative: an RSS of 12 is excellent if $s_{yy} = 4000$ but poor if $s_{yy} = 15$. Always judge the RSS against the total variation s_{yy} (or s_{xx}) — this idea is made precise by the product moment correlation coefficient in the next section, since $\frac{\text{RSS}_{y \text{ on } x}}{s_{yy}} = 1 - r^2$.

5.4 The Line of Best Fit

We now come to the central result of this section, which justifies the name "least squares".

PROPERTY The least squares line is the line of best fit

Among all straight lines $y = a + bx$, the one that makes $\text{RSS} = \sum (y_i - a - bx_i)^2$ as small as possible is exactly the least squares regression line of y on x , with $b = \frac{s_{xy}}{s_{xx}}$ and $a = \bar{y} - b\bar{x}$. For this reason the least squares line is called the **line of best fit** (in the least squares sense). Similarly, the least squares line of x on y minimises $\sum (x_i - a' - b'y_i)^2$ among all lines $x = a' + b'y$.

Why this is true.

For fixed b , the quantity $S(a, b) = \sum (y_i - a - bx_i)^2$ is minimised with respect to a when $a = \bar{y} - b\bar{x}$ (this is the same "sum of squared deviations is minimised about the mean" fact used for the variance). Substituting and writing $u_i = x_i - \bar{x}$, $v_i = y_i - \bar{y}$ gives

$$S = \sum (v_i - bu_i)^2 = s_{xx} b^2 - 2s_{xy} b + s_{yy},$$

a quadratic in b with positive leading coefficient $s_{xx} > 0$. Its minimum occurs at $b = \frac{s_{xy}}{s_{xx}}$, and the minimum value is $s_{yy} - \frac{(s_{xy})^2}{s_{xx}}$, which is precisely the RSS formula above. ■

KEY POINTS Key facts about RSS and the line of best fit

- Residual: $e_i = y_i - \hat{y}_i$ (vertical, y on x); $d_i = x_i - \hat{x}_i$ (horizontal, x on y).
- $\text{RSS}_{y \text{ on } x} = s_{yy} - \frac{(s_{xy})^2}{s_{xx}}$ and $\text{RSS}_{x \text{ on } y} = s_{xx} - \frac{(s_{xy})^2}{s_{yy}}$.
- The least squares line minimises the RSS: it is the line of best fit.
- $\text{RSS} = 0 \iff$ all points lie exactly on the line (perfect linear fit, $|r| = 1$).
- Any other line (e.g. Mayer's line) has a larger RSS than the least squares line.

5.5 Worked Example: Residuals and RSS for a Small Data Set

WORKED EXAMPLE 6 Computing residuals and RSS directly

A small trader in Buea records the number of crates of plantain sold, y , against the midday temperature in $^{\circ}\text{C}$, x , on five market days:

x	20	22	25	27	30
y	14	15	18	17	21

The least squares line of y on x is $\hat{y} = 1.1 + 0.7x$. Compute the five residuals, verify that they sum to zero, and deduce the RSS. Check the answer with the compact formula.

SOLUTION

Solution. Compute $\hat{y}_i = 1.1 + 0.7x_i$ and $e_i = y_i - \hat{y}_i$:

x_i	20	22	25	27	30	
y_i	14	15	18	17	21	
$\hat{y}_i$	15.1	16.5	18.6	20.0	22.1	
e_i	-1.1	-1.5	-0.6	-3.0	-1.1	

Wait — let us recompute carefully: $\hat{y}_1 = 1.1 + 0.7(20) = 15.1$, so $e_1 = 14 - 15.1 = -1.1$; $\hat{y}_2 = 1.1 + 15.4 = 16.5$, $e_2 = -1.5$; $\hat{y}_3 = 1.1 + 17.5 = 18.6$, $e_3 = -0.6$; $\hat{y}_4 = 1.1 + 18.9 = 20.0$, $e_4 = -3.0$; $\hat{y}_5 = 1.1 + 21.0 = 22.1$, $e_5 = -1.1$.

Hmm, these do not sum to zero, which means the stated line is not exactly the least

squares line for this data. Let us instead derive the line ourselves. We have $\bar{x} = \frac{124}{5} = 24.8$, $\bar{y} = \frac{85}{5} = 17$.

$$s_{xx} = \sum x^2 - n\bar{x}^2 = (400 + 484 + 625 + 729 + 900) - 5(24.8)^2 = 3138 - 3075.2 = 62.8,$$

$$s_{xy} = \sum xy - n\bar{x}\bar{y} = (280 + 330 + 450 + 459 + 630) - 5(24.8)(17) = 2149 - 2108 = 41,$$

$$s_{yy} = \sum y^2 - n\bar{y}^2 = (196 + 225 + 324 + 289 + 441) - 5(289) = 1475 - 1445 = 30.$$

Hence $b = \frac{41}{62.8} \approx 0.6529$ and $a = 17 - 0.6529(24.8) \approx 0.8089$. The least squares line is $\hat{y} = 0.809 + 0.653x$.

The residuals are:

x_i	20	22	25	27	30
$\hat{y}_i$	13.87	15.17	17.13	18.44	20.40
e_i	0.13	-0.17	0.87	-1.44	0.60

Sum of residuals: $0.13 - 0.17 + 0.87 - 1.44 + 0.60 \approx 0$ (up to rounding), as expected for a least squares line. Then

$$RSS = (0.13)^2 + (-0.17)^2 + (0.87)^2 + (-1.44)^2 + (0.60)^2 \approx 0.017 + 0.029 + 0.757 + 2.074 + 0.360 \approx 3.24.$$

Check with the compact formula:

$$RSS = s_{yy} - \frac{(s_{xy})^2}{s_{xx}} = 30 - \frac{41^2}{62.8} = 30 - \frac{1681}{62.8} \approx 30 - 26.77 = 3.23.$$

The two answers agree (the tiny difference comes from rounding the residuals). Conclusion: $RSS \approx 3.23$, small compared with $s_{yy} = 30$, so the line fits the data well.

5.6 Comparing Two Candidate Lines by their RSS

Because the least squares line minimises the RSS, any other line fitted to the same data — for instance Mayer's line from Section 2 — must have an RSS at least as large. This gives a concrete way to compare two candidate lines: compute the RSS of each; the smaller RSS indicates the better fit (in the least squares sense).

WORKED EXAMPLE 7 Mayer's line versus the least squares line


For the data of the previous worked example, Mayer's method (splitting into the groups $\{20, 22, 25\}$ and $\{27, 30\}$) gives the line $y = 0.8x - 2.73$ approximately; take $\hat{y} = 0.8x - 2.7$ for simplicity. Compute its RSS and compare with the least squares RSS of 3.23.

SOLUTION

Solution. Predictions from Mayer's line: $\hat{y}_1 = 13.3, \hat{y}_2 = 14.9, \hat{y}_3 = 17.3, \hat{y}_4 = 18.9, \hat{y}_5 = 21.3$. Residuals: 0.7, 0.1, 0.7, -1.9, -0.3. Hence

$$\text{RSS}_{\text{Mayer}} = 0.49 + 0.01 + 0.49 + 3.61 + 0.09 = 4.69.$$

Since $4.69 > 3.23$, Mayer's line fits less well than the least squares line — exactly as the theory guarantees. The difference is modest here because the points are nearly collinear; for more scattered data the gap can be much larger.



Two candidate lines through the same data. The least squares line (solid blue) has the smaller RSS; Mayer's line (dashed orange) is close but not optimal.

5.7 RSS and the Quality of a Linear Model

The RSS connects directly with correlation. Dividing the RSS formula by s_{yy} :

$$\frac{\text{RSS}_{y \text{ on } x}}{s_{yy}} = 1 - \frac{(s_{xy})^2}{s_{xx}s_{yy}} = 1 - r^2,$$

where r is the product moment correlation coefficient studied in the next section. Consequently:

- $\text{RSS} = 0 \iff r^2 = 1$: perfect linear fit, all points exactly on the line;
- the closer r^2 is to 1, the smaller the RSS relative to the total variation s_{yy} , and the better the line explains the data;
- if $r = 0$ then $\text{RSS} = s_{yy}$: the line explains nothing, and predicting $\hat{y} = \bar{y}$ (a horizontal line) is just as good.

WARNING A small RSS does not prove the model is appropriate

The least squares line is the best straight line, but the true relationship may not be linear at all. A curved pattern can still yield a moderately small RSS while the residuals show a clear systematic pattern (all positive in the middle, negative at the ends, for example). Always look at the scatter diagram first, and if possible at the pattern of the residuals, before trusting a linear model. RSS measures the quality of the fit within the family of straight lines — nothing more.

DID YOU KNOW? Why squares and not absolute values?

Minimising $\sum |e_i|$ would seem just as natural, and such "least absolute deviations" methods do exist in modern statistics. But squared residuals lead to clean algebra (a quadratic to minimise), explicit formulae for a and b , and the elegant identity $\text{RSS} = s_{yy}(1 - r^2)$. This is why Legendre and Gauss, who introduced the least squares method around 1800 to predict the orbit of the asteroid Ceres, chose squares — and why your calculator and the GCE syllabus still use them today.

EXERCISE 5 RSS for the line of x on y

Using the plantain data of the worked example ($s_{xx} = 62.8$, $s_{yy} = 30$, $s_{xy} = 41$):

1. Find the equation of the least squares regression line of x on y .
2. Compute $\text{RSS}_{x \text{ on } y}$ using the compact formula.
3. Explain why this value cannot be compared directly with $\text{RSS}_{y \text{ on } x} = 3.23$.

SOLUTION Exercise 1

1. $b' = \frac{s_{xy}}{s_{yy}} = \frac{41}{30} \approx 1.3667$; $a' = \bar{x} - b'\bar{y} = 24.8 - 1.3667(17) \approx 1.567$. So $\hat{x} = 1.567 + 1.367y$.
2. $\text{RSS}_{x \text{ on } y} = s_{xx} - \frac{(s_{xy})^2}{s_{yy}} = 62.8 - \frac{1681}{30} = 62.8 - 56.03 \approx 6.77$.
3. $\text{RSS}_{x \text{ on } y}$ is measured in squared units of x (here $(^\circ\text{C})^2$) while $\text{RSS}_{y \text{ on } x}$ is in squared crates; quantities with different units cannot be compared. Each RSS must be judged against its own total variation, s_{xx} or s_{yy} respectively.

EXERCISE 6 Comparing two fitted lines

A student fits two lines to a data set with $s_{yy} = 120$: line L_1 gives $\text{RSS} = 30$ and line L_2 gives $\text{RSS} = 24$.

1. Which line is the better fit in the least squares sense? Justify.
2. Can L_1 be the least squares regression line of y on x ? Explain.
3. Compute r^2 if L_2 is the least squares line.

SOLUTION Exercise 2

1. L_2 , because it has the smaller residual sum of squares ($24 < 30$): on average its points lie closer to the line.
2. No. The least squares line minimises the RSS over all lines, so no other line can have a strictly smaller RSS. Since L_2 achieves $24 < 30$, L_1 cannot be the least squares line.
3. $r^2 = 1 - \frac{\text{RSS}}{s_{yy}} = 1 - \frac{24}{120} = 0.8$. Hence $|r| = \sqrt{0.8} \approx 0.894$, indicating a strong linear relationship.

KEY POINTS Summary

The residual $e_i = y_i - \hat{y}_i$ measures the vertical error of the line of y on x at each point; the RSS, $\sum e_i^2$, measures the global error. The least squares line is the unique line minimising the RSS — the **line of best fit** — and its RSS is given by $s_{yy} - \frac{(s_{xy})^2}{s_{xx}}$. Comparing RSS values ranks competing lines, and $\text{RSS} = 0$ characterises a perfect linear fit. The ratio $\text{RSS}/s_{yy} = 1 - r^2$ leads naturally to the product moment correlation coefficient, the subject of the next section.

6 THE PRODUCT MOMENT CORRELATION COEFFICIENT

In the previous section we used the **residual sum of squares** to decide which of the two least squares regression lines, y on x or x on y , fits a given scatter of points better. The RSS measures the quality of fit of a chosen line. But it does not answer a more basic question that every GCE examiner loves to ask: how strong is the linear relationship between the two variables? A small RSS tells us the line fits well, but the RSS is measured in squared units of y , so its size alone does not allow us to compare different data sets. What we need is a dimensionless number, lying on

a fixed scale, that summarises the strength and direction of the linear association. That number is the **product moment correlation coefficient**, denoted r , and it is the subject of this section.

6.1 Definition of the Product Moment Correlation Coefficient

Think of a scatter diagram. If the points cluster tightly around an upward sloping line, we say x and y are strongly positively correlated. If they cluster around a downward sloping line, the correlation is strongly negative. If the points form a shapeless cloud, there is little or no linear correlation. We already possess all the ingredients needed to measure this: the covariance s_{xy} tells us the direction of the association (positive when large x goes with large y , negative otherwise), while s_{xx} and s_{yy} measure the spreads of x and y separately. Dividing the covariance by the geometric mean of the two spreads removes the units and bounds the result between -1 and 1 .

DEFINITION Product moment correlation coefficient

For a sample of n pairs $(x_1, y_1), \dots, (x_n, y_n)$, the **product moment correlation coefficient** (PMCC) is

$$r = \frac{s_{xy}}{\sqrt{s_{xx} s_{yy}}} = \frac{\sum (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum (x_i - \bar{x})^2 \sum (y_i - \bar{y})^2}}.$$

It is also called Pearson's correlation coefficient.

For hand computation at the GCE, we never work with deviations directly; we use the summary sums $\sum x$, $\sum y$, $\sum x^2$, $\sum y^2$, $\sum xy$.

PROPERTY Computational formula for r

Using $S_{xy} = \sum xy - n\bar{x}\bar{y}$, $S_{xx} = \sum x^2 - n\bar{x}^2$ and $S_{yy} = \sum y^2 - n\bar{y}^2$,

$$r = \frac{\sum xy - n\bar{x}\bar{y}}{\sqrt{(\sum x^2 - n\bar{x}^2)(\sum y^2 - n\bar{y}^2)}}.$$

This formula is examinable: candidates must be able to derive the numerator and denominator forms from the definitions of S_{xy} , S_{xx} and S_{yy} (expanding $\sum (x_i - \bar{x})(y_i - \bar{y})$ and using $\sum x_i = n\bar{x}$).

6.2 Properties and Interpretation of r

PROPERTY Range and boundary values of r

For every bivariate data set,

$$-1 \leq r \leq 1.$$

Moreover:


- $r = 1$ if and only if all the points lie exactly on a straight line of positive gradient (perfect positive linear correlation);
- $r = -1$ if and only if all the points lie exactly on a straight line of negative gradient (perfect negative linear correlation);
- $r = 0$ when $S_{xy} = 0$: there is no linear correlation.

The inequality $-1 \leq r \leq 1$ follows from the Cauchy–Schwarz inequality applied to the deviations $x_i - \bar{x}$ and $y_i - \bar{y}$; its proof is not required at the GCE, but the result and the meaning of the boundary values must be known.

Intermediate values are interpreted by their sign (direction) and their absolute value (strength). A commonly used guide is:

Value of r	Interpretation
$0.8 \leq r \leq 1$	Strong linear correlation
$0.5 \leq r < 0.8$	Moderate linear correlation
$0.2 \leq r < 0.5$	Weak linear correlation
$ r < 0.2$	Very weak or no linear correlation

These bands are conventions, not laws: always interpret r in the context of the question (for example, “there is a strong positive linear correlation between revision hours and mock examination marks”).



Four scatter diagrams illustrating perfect positive, moderate positive, zero, and strong negative linear correlation.

6.3 Link with the Two Regression Gradients

Recall from the least squares section that the regression line of y on x has gradient $b = \frac{S_{xy}}{S_{xx}}$, while the regression line of x on y has gradient $b' = \frac{S_{xy}}{S_{yy}}$.

PROPERTY r^2 is the product of the two regression gradients

$$b \cdot b' = \frac{S_{xy}}{S_{xx}} \cdot \frac{S_{xy}}{S_{yy}} = \frac{S_{xy}^2}{S_{xx} S_{yy}} = r^2.$$

Consequently $r = \pm\sqrt{b \cdot b'}$, the sign being the common sign of b and b' (which is the sign of S_{xy}). This identity is frequently examined: given the two regression lines, one can recover r without touching the raw data.

This result also explains why the two regression lines coincide exactly when $r = \pm 1$: then $|b \cdot b'| = 1$, the two lines are mutual inverses with the same slope through $(\bar{x}, \bar{y})$, i.e. the same line.

EXAMPLE Recovering r from the regression lines

The least squares regression lines for a data set are $y = 0.8x + 2$ (y on x) and $x = 0.45y + 1$ (x on y). Rewriting the second as x on y gives gradient $b' = 0.45$. Hence

$$r^2 = 0.8 \times 0.45 = 0.36, \quad r = +0.6$$

(positive because both gradients are positive). There is a moderate positive linear correlation.

6.4 Effect of a Linear Change of Scale (GCE Extra)**PROPERTY** Invariance of r under coding

If the data are coded by linear transformations $u = \frac{x-a}{p}$ and $v = \frac{y-c}{q}$ with $p, q > 0$, then the correlation coefficient of the coded data equals that of the original data:

$$r_{uv} = r_{xy}.$$

Indeed, deviations satisfy $u_i - \bar{u} = (x_i - \bar{x})/p$, so $S_{uu} = S_{xx}/p^2$, $S_{vv} = S_{yy}/q^2$, $S_{uv} = S_{xy}/(pq)$, and the factors p, q cancel in the ratio. This is a useful GCE extra: coding large or awkward values (e.g. years, or prices in thousands of FCFA) does not change r , and can greatly simplify the arithmetic.

6.5 Computing r from a Table of Sums**METHOD** Computing and interpreting r

1. From the data (or the question), obtain the five sums $\sum x$, $\sum y$, $\sum x^2$, $\sum y^2$, $\sum xy$ and the sample size n .
2. Compute the means $\bar{x} = \sum x/n$ and $\bar{y} = \sum y/n$.
3. Compute $S_{xy} = \sum xy - n\bar{x}\bar{y}$, $S_{xx} = \sum x^2 - n\bar{x}^2$, $S_{yy} = \sum y^2 - n\bar{y}^2$.
4. Substitute into $r = \frac{S_{xy}}{\sqrt{S_{xx}S_{yy}}}$ and give the value to 3 or 4 significant figures.
5. Interpret the sign and magnitude of r in the context of the problem.

WORKED EXAMPLE 8 GCE-style worked example

A teacher in Buea records, for 8 candidates, the number of hours x spent revising Pure Mathematics during the week before a mock examination and the mark y (out of 100) obtained. The results are summarised by

$$\sum x = 96, \quad \sum y = 424, \quad \sum x^2 = 1392, \quad \sum y^2 = 23938, \quad \sum xy = 5802.$$

Calculate the product moment correlation coefficient between revision time and mark, and interpret your result.

SOLUTION

Solution. Here $n = 8$, so

$$\bar{x} = \frac{96}{8} = 12, \quad \bar{y} = \frac{424}{8} = 53.$$

Compute the three summary quantities:

$$S_{xy} = \sum xy - n\bar{x}\bar{y} = 5802 - 8(12)(53) = 5802 - 5088 = 714,$$

$$S_{xx} = \sum x^2 - n\bar{x}^2 = 1392 - 8(144) = 1392 - 1152 = 240,$$

$$S_{yy} = \sum y^2 - n\bar{y}^2 = 23938 - 8(2809) = 23938 - 22472 = 1466.$$

Therefore

$$r = \frac{714}{\sqrt{240 \times 1466}} = \frac{714}{\sqrt{351840}} = \frac{714}{593.16...} \approx 0.904.$$

Interpretation. Since $r \approx 0.90$ is close to 1, there is a strong positive linear correlation between revision time and mock mark: candidates who revised more tended to obtain higher marks.

6.6 Limitations of the Product Moment Correlation Coefficient

WARNING $r \approx 0$ does not mean “no relationship”

The PMCC measures linear association only. A data set can have $r \approx 0$ while x and y are perfectly related by a curve. For example, points lying exactly on the parabola $y = x^2$ for x symmetric about 0 give $S_{xy} = 0$, hence $r = 0$, yet y is completely determined by x . Always look at the scatter diagram before concluding that two variables are unrelated.

WARNING Correlation is not causation

A value of r close to ± 1 shows that x and y vary together linearly; it does not prove that changes in x cause changes in y . Both variables may be driven by a hidden third factor

(a confounding variable). For instance, ice cream sales and cases of sunburn are strongly correlated in many towns, not because one causes the other, but because both increase in hot sunny weather. In GCE answers, never write “ x causes y ” on the basis of r alone.

Two further cautions deserve mention. First, r is sensitive to outliers: a single point far from the main cloud can drag r towards $+1$ or -1 and completely distort the conclusion. Second, r computed on a restricted range of x may badly understate the true association. Both issues are visible only on the scatter diagram, which is why examiners reward candidates who sketch or refer to it.

KEY POINTS Key points on the PMCC

- $r = \frac{S_{xy}}{\sqrt{S_{xx}S_{yy}}}$ with $S_{xy} = \sum xy - n\bar{x}\bar{y}$, etc.; always compute the five sums first.
- $-1 \leq r \leq 1$; $r = \pm 1$ means a perfect linear relationship, $r = 0$ means no linear relationship.
- $r^2 = b b'$, the product of the two least squares gradients; the sign of r is the common sign of b and b' .
- r is unchanged by linear coding of the data.
- r measures linear association only, is sensitive to outliers, and never by itself proves causation.

EXERCISE 7 Computing and using r

For six shops in Douala, the daily number x of customers and the daily revenue y (in thousands of FCFA) give

$$\sum x = 330, \quad \sum y = 612, \quad \sum x^2 = 20100, \quad \sum y^2 = 64452, \quad \sum xy = 36420.$$

1. Calculate r and interpret it.
2. Given that the regression line of y on x has gradient $b = 1.5$, find the gradient b' of the regression line of x on y .

SOLUTION Exercise 1

$$1. n = 6, \bar{x} = 55, \bar{y} = 102.$$

$$S_{xy} = 36420 - 6(55)(102) = 36420 - 33660 = 2760,$$

$$S_{xx} = 20100 - 6(55)^2 = 20100 - 18150 = 1950,$$

$$S_{yy} = 64452 - 6(102)^2 = 64452 - 62424 = 2028.$$

$$r = \frac{2760}{\sqrt{1950 \times 2028}} = \frac{2760}{\sqrt{3954600}} \approx \frac{2760}{1988.6} \approx 0.888.$$

There is a strong positive linear correlation between the number of customers and daily revenue. 2. From $r^2 = b b'$: $b' = \frac{r^2}{b} = \frac{0.888^2}{1.5} \approx \frac{0.788}{1.5} \approx 0.526$.

7 RANK CORRELATION: SPEARMAN AND KENDALL

In the previous section we studied the **product moment correlation coefficient** r , which measures the strength of a linear relationship between two quantitative variables. However, real data often do not satisfy the assumptions required for Pearson's r : the variables may be ordinal (only ranks are available), the relationship may be non-linear but monotonic, or the data may contain outliers that distort r . In such situations we turn to **rank correlation** coefficients, which are based solely on the order of the observations. This section introduces the two most widely used rank correlation measures: **Spearman's rank correlation coefficient** (R_s) and **Kendall's rank correlation coefficient** (τ).

7.1 Ranking Data and Tied Ranks

Before we can compute any rank correlation, we must assign **ranks** to the observations. The smallest value receives rank 1, the next smallest rank 2, and so on up to n . Alternatively, one may rank in descending order (largest = 1) provided the same convention is used for both variables.

DEFINITION Ranks

Given a sample of n observations $x_1, x_2, \dots, x_n$, the rank of x_i is its position when the data are sorted in ascending order. If several observations have the same value (a tie), each of them receives the average of the ranks they would have occupied had they been slightly different.

WARNING Tied Ranks

When ties occur, always assign the mean rank to each tied item. For example, if the 3rd and 4th smallest values are equal, they both receive rank $\frac{3+4}{2} = 3.5$. The next distinct value

then receives rank 5. Failing to do this systematically is a common error that invalidates both Spearman's and Kendall's formulas.

EXAMPLE Ranking with Ties

Consider the marks of 7 students in a Mathematics test:

52, 58, 58, 61, 65, 65, 65.

Sorting gives the same order. The ranks are assigned as follows:

- 52 is smallest $\rightarrow$ rank 1.
- The two 58's occupy positions 2 and 3 $\rightarrow$ each gets rank $\frac{2+3}{2} = 2.5$.
- 61 is next $\rightarrow$ rank 4.
- The three 65's occupy positions 5, 6, 7 $\rightarrow$ each gets rank $\frac{5+6+7}{3} = 6$.

The rank vector is (1, 2.5, 2.5, 4, 6, 6, 6).

Ranking with Tied Values

Value	52	58	58	61	65	65	65
Rank	1	2.5	2.5	4	6	6	6

Illustration of tied ranks: the two 58's share rank 2.5 (orange), the three 65's share rank 6 (green).

7.2 Spearman's Rank Correlation Coefficient

Spearman's coefficient, denoted R_s (or sometimes ρ), is simply the Pearson correlation coefficient computed on the ranks of the two variables. When there are no ties, it reduces to a very simple formula.

DEFINITION Spearman's Rank Correlation Coefficient

Let (x_i, y_i) , $i = 1, \dots, n$ be paired observations. Let $R(x_i)$ and $R(y_i)$ be their ranks. Spearman's rank correlation coefficient is

$$R_s = \frac{\sum_{i=1}^n (R(x_i) - \overline{R_x})(R(y_i) - \overline{R_y})}{\sqrt{\sum_{i=1}^n (R(x_i) - \overline{R_x})^2 \sum_{i=1}^n (R(y_i) - \overline{R_y})^2}}.$$

If there are no tied ranks, this simplifies to

$$R_s = 1 - \frac{6 \sum_{i=1}^n d_i^2}{n(n^2 - 1)}, \quad d_i = R(x_i) - R(y_i).$$

If ties are present, one must compute the Pearson correlation coefficient on the ranked data (using the averaged ranks). The simplified formula $1 - \frac{6 \sum d_i^2}{n(n^2 - 1)}$ is not valid when ties exist. In GCE examinations, the simplified formula is used only when explicitly stated or clearly evident that there are no ties; otherwise compute the Pearson correlation on the ranked data.

PROPERTY Properties of R_s

- $-1 \leq R_s \leq 1$.
- $R_s = 1 \iff$ the two rankings are in perfect agreement (monotonically increasing).
- $R_s = -1 \iff$ the two rankings are in perfect disagreement (one is the reverse of the other).
- $R_s = 0$ indicates no monotonic association.
- R_s measures the strength of a monotonic relationship, not necessarily linear.

METHOD Computing Spearman's R_s (Step-by-Step)

1. List the paired observations (x_i, y_i) .
2. Assign ranks $R(x_i)$ to the x 's and $R(y_i)$ to the y 's. Handle ties by averaging ranks.
3. Check for ties:
 - No ties: compute $d_i = R(x_i) - R(y_i)$ for each pair, then $\sum d_i^2$ and apply $R_s = 1 - \frac{6 \sum d_i^2}{n(n^2 - 1)}$.
 - Ties present: compute the Pearson correlation coefficient on the ranked data (using the averaged ranks).
4. Interpret the value: close to 1 $\rightarrow$ strong positive monotonic agreement; close to -1 $\rightarrow$ strong negative monotonic agreement; near 0 $\rightarrow$ little or no monotonic association.

WORKED EXAMPLE 9 Spearman's R_s with No Ties

Two judges rank 8 candidates in a singing competition. The ranks given by Judge A and Judge B are:

Candidate	1	2	3	4	5	6	7	8
Judge A	1	2	3	4	5	6	7	8
Judge B	2	1	4	3	6	5	8	7

Compute Spearman's rank correlation coefficient.

SOLUTION

Solution. Here the ranks are already given and there are no ties. We compute $d_i = R_A - R_B$:

Candidate	1	2	3	4	5	6	7	8
R_A	1	2	3	4	5	6	7	8
R_B	2	1	4	3	6	5	8	7
d_i	-1	1	-1	1	-1	1	-1	1
d_i^2	1	1	1	1	1	1	1	1

$\sum d_i^2 = 8$. With $n = 8$:

$$R_s = 1 - \frac{6 \times 8}{8(8^2 - 1)} = 1 - \frac{48}{8 \times 63} = 1 - \frac{48}{504} = 1 - 0.09524 = 0.90476.$$

The judges show a very strong positive agreement in their rankings.

WORKED EXAMPLE 10 Spearman's R_s with Tied Ranks

Eight students are ranked by their Mathematics teacher (Mr. T) and their Physics teacher (Ms. K) based on test scores. The raw scores and ranks are:

Student	A	B	C	D	E	F	G	H
Maths score	72	65	65	58	58	58	50	45
Physics score	80	75	70	70	65	60	60	55

Compute Spearman's rank correlation coefficient.

SOLUTION

Solution.

Step 1: Assign ranks to Maths scores (with ties). Sorted Maths: 45, 50, 58, 58, 58, 65, 65, 72.

- 45 (H) → rank 1
- 50 (G) → rank 2
- 58 (D,E,F) occupy positions 3,4,5 → each gets rank $\frac{3+4+5}{3} = 4$
- 65 (B,C) occupy positions 6,7 → each gets rank $\frac{6+7}{2} = 6.5$
- 72 (A) → rank 8

Maths ranks R_M : A=8, B=6.5, C=6.5, D=4, E=4, F=4, G=2, H=1.

Step 2: Assign ranks to Physics scores (with ties). Sorted Physics: 55, 60, 60, 65, 70, 70, 75, 80.

- 55 (H) → rank 1
- 60 (F,G) occupy positions 2,3 → each gets rank $\frac{2+3}{2} = 2.5$
- 65 (E) → rank 4
- 70 (C,D) occupy positions 5,6 → each gets rank $\frac{5+6}{2} = 5.5$
- 75 (B) → rank 7
- 80 (A) → rank 8

Physics ranks R_P : A=8, B=7, C=5.5, D=5.5, E=4, F=2.5, G=2.5, H=1.

Step 3: Compute Pearson correlation on the ranks.

Student	A	B	C	D	E	F	G	H
R_M	8	6.5	6.5	4	4	4	2	1
R_P	8	7	5.5	5.5	4	2.5	2.5	1

$\overline{R_M} = \overline{R_P} = \frac{8+7+6+5+4+3+2+1}{8} = 4.5$ (sum of ranks 1 to 8 is always 36). Compute deviations and products:

Student	A	B	C	D	E	F	G	H
$R_M - 4.5$	3.5	2.0	2.0	-0.5	-0.5	-0.5	-2.5	-3.5
$R_P - 4.5$	3.5	2.5	1.0	1.0	-0.5	-2.0	-2.0	-3.5
Product	12.25	5.00	2.00	-0.50	0.25	1.00	5.00	12.25

$\sum(R_M - \overline{R_M})(R_P - \overline{R_P}) = 37.25$. $\sum(R_M - \overline{R_M})^2 = 12.25 + 4 + 4 + 0.25 + 0.25 + 0.25 + 6.25 + 12.25 = 39.5$.
 $\sum(R_P - \overline{R_P})^2 = 12.25 + 6.25 + 1 + 1 + 0.25 + 4 + 4 + 12.25 = 41$.

$$R_s = \frac{37.25}{\sqrt{39.5 \times 41}} = \frac{37.25}{\sqrt{1619.5}} = \frac{37.25}{40.243} \approx 0.9256.$$

There is a very strong positive monotonic association between the two teachers' rankings.

7.3 Kendall's Rank Correlation Coefficient

Kendall's τ (tau) takes a different approach: instead of comparing rank differences, it compares the order of every pair of observations. It counts how many pairs are concordant (same order in both rankings) and how many are discordant (opposite order).

DEFINITION Concordant and Discordant Pairs

Consider two distinct observations i and j ($i < j$). Let their ranks in the two variables be $(R_x(i), R_y(i))$ and $(R_x(j), R_y(j))$.

- The pair is concordant if $(R_x(i) - R_x(j))(R_y(i) - R_y(j)) > 0$ (both differences have the same sign).
- The pair is discordant if $(R_x(i) - R_x(j))(R_y(i) - R_y(j)) < 0$ (differences have opposite signs).
- If either difference is zero (a tie in one variable), the pair is neither concordant nor discordant and is excluded from the count in the basic τ formula. (Tie-corrected versions τ_b and τ_c exist but are beyond the GCE syllabus.)

Let C be the number of concordant pairs and D the number of discordant pairs. The total number of pairs is $\binom{n}{2} = \frac{n(n-1)}{2}$. Kendall's τ is defined as

$$\tau = \frac{C - D}{\frac{1}{2}n(n-1)} = \frac{2(C - D)}{n(n-1)}.$$

Often the score $S = C - D$ is used, giving $\tau = \frac{2S}{n(n-1)}$.

PROPERTY Properties of Kendall's τ

- $-1 \leq \tau \leq 1$.
- $\tau = 1 \iff$ perfect agreement (all pairs concordant).
- $\tau = -1 \iff$ perfect disagreement (all pairs discordant).
- $\tau = 0 \iff$ number of concordant pairs equals number of discordant pairs (no monotonic association).
- τ is more robust to outliers than R_s because it only uses the sign of the difference, not its magnitude.

METHOD Counting Concordant and Discordant Pairs (Kendall's τ)

1. Arrange the data so that one variable (say X) is in natural order $1, 2, \dots, n$. Reorder the Y

ranks accordingly.

2. For each i from 1 to $n - 1$, compare Y_i with all Y_j for $j > i$.

3. Count:

- Concordant if $Y_j > Y_i$ (since $X_j > X_i$ by construction).
- Discordant if $Y_j < Y_i$.
- Ignore if $Y_j = Y_i$ (tie in Y).

4. Sum C and D over all i . Compute $S = C - D$ and $\tau = \frac{2S}{n(n-1)}$.




Illustration of concordant (green) and discordant (orange) pairs for Kendall's τ . The left column is in natural order; arrows connect to the right column. Pair (1,2) is discordant because item 1 is above item 2 in X but below in Y .

WORKED EXAMPLE 11 Kendall's τ on the Same Data

Using the same judges' rankings as in the Spearman example (8 candidates, no ties), compute Kendall's τ .

SOLUTION

Solution. We have $n = 8$. Judge A's ranks are already in natural order 1, 2, ..., 8. Judge B's ranks in that order are:

$$Y = (2, 1, 4, 3, 6, 5, 8, 7).$$

We count concordant and discordant pairs by scanning each i and comparing with later j .

- $i = 1, Y_1 = 2$: later values 1, 4, 3, 6, 5, 8, 7. Values < 2 : only 1 $\rightarrow$ 1 discordant. Values > 2 : 4, 3, 6, 5, 8, 7 $\rightarrow$ 6 concordant.
- $i = 2, Y_2 = 1$: later values 4, 3, 6, 5, 8, 7. All are $> 1 \rightarrow$ 6 concordant, 0 discordant.
- $i = 3, Y_3 = 4$: later values 3, 6, 5, 8, 7. < 4 : 3 $\rightarrow$ 1 discordant. > 4 : 6, 5, 8, 7 $\rightarrow$ 4 concordant.
- $i = 4, Y_4 = 3$: later values 6, 5, 8, 7. All $> 3 \rightarrow$ 4 concordant.
- $i = 5, Y_5 = 6$: later values 5, 8, 7. < 6 : 5 $\rightarrow$ 1 discordant. > 6 : 8, 7 $\rightarrow$ 2 concordant.
- $i = 6, Y_6 = 5$: later values 8, 7. Both $> 5 \rightarrow$ 2 concordant.
- $i = 7, Y_7 = 8$: later value 7. $< 8 \rightarrow$ 1 discordant.

Total concordant $C = 6 + 6 + 4 + 4 + 2 + 2 = 24$. Total discordant $D = 1 + 1 + 1 + 1 = 4$. Check: total pairs $= \binom{8}{2} = 28$, $C + D = 28 \equiv$. $S = C - D = 20$.

$$\tau = \frac{2S}{n(n-1)} = \frac{40}{8 \times 7} = \frac{40}{56} = \frac{5}{7} \approx 0.7143.$$

7.4 Uses and Limitations of Kendall's Coefficient

KEY POINTS Uses of Kendall's τ

- Small samples: The exact distribution of τ under the null hypothesis of independence is known for small n , making it ideal for exact hypothesis tests.
- Ordinal data: When data are only available as ranks (e.g., preference rankings, competition results), τ is a natural choice.
- Agreement between judges/raters: τ directly measures the degree of concordance between two rankings, which is exactly what is needed in inter-rater reliability studies.
- Robustness: Because it uses only the sign of rank differences, τ is less affected by outliers than Spearman's R_s or Pearson's r .

WARNING Limitations of Kendall's τ

- Computational burden for large n : Counting concordant/discordant pairs requires $O(n^2)$ comparisons. For $n > 30$ it becomes tedious by hand (though software handles it easily).

- Ties complicate the count: Ties in either variable produce pairs that are neither concordant nor discordant. The standard formula $\tau = \frac{2(C-D)}{n(n-1)}$ assumes no ties. With ties, one must use Kendall's τ_b or τ_c which adjust the denominator. This is beyond the GCE syllabus but you should be aware that ties reduce the maximum possible $|\tau|$.
- Only measures monotonic agreement: Like R_s , τ captures only monotonic relationships. A perfect U-shaped relationship would give $\tau \approx 0$.

7.5 Comparison of r , R_s and Kendall's τ

Feature	Pearson's r	Spearman's R_s	Kendall's τ
Data type	Quantitative (interval/ratio)	Quantitative or ordinal	Ordinal or quantitative
Relationship measured	Linear	Monotonic	Monotonic (concordance)
Sensitivity to outliers	High	Moderate (uses rank differences)	Low (uses only signs)
Assumptions	Bivariate normality, linearity, homoscedasticity	None (non-parametric)	None (non-parametric)
Interpretation	Strength of linear fit	Strength of monotonic trend	Probability concordance minus discordance
Computation	Simple formula	Simple if no ties; otherwise Pearson on ranks	Pair counting ($O(n^2)$)
When to use	Linear relationship, quantitative data, no outliers	Monotonic trend, ordinal data, outliers present	Small samples, ordinal data, judge agreement, many ties

7.6 Worked Example: Two Teachers Ranking Six Candidates

WORKED EXAMPLE 12 Comparing R_s and τ on a Small Sample

Two teachers, Mr. Ngwa and Ms. Fon, independently rank 6 students (A–F) based on their overall performance. The rankings are:

Student	A	B	C	D	E	F
Mr. Ngwa	1	2	3	4	5	6
Ms. Fon	2	1	4	3	6	5

1. Compute Spearman's rank correlation coefficient R_s .
2. Compute Kendall's rank correlation coefficient τ .
3. Compare the two values and comment on the agreement between the teachers.

SOLUTION

Solution.

1. Spearman's R_s . There are no ties. Differences $d_i = R_{\text{Ngwa}} - R_{\text{Fon}}$:

Student	A	B	C	D	E	F
R_N	1	2	3	4	5	6
R_F	2	1	4	3	6	5
d_i	-1	1	-1	1	-1	1
d_i^2	1	1	1	1	1	1

$\sum d_i^2 = 6$. With $n = 6$:

$$R_s = 1 - \frac{6 \times 6}{6(6^2 - 1)} = 1 - \frac{36}{6 \times 35} = 1 - \frac{36}{210} = 1 - 0.17143 = 0.82857.$$

2. Kendall's τ . Arrange in Mr. Ngwa's order (already natural). Ms. Fon's ranks in that order: $Y = (2, 1, 4, 3, 6, 5)$. Count concordant/discordant pairs:

- $i = 1, Y_1 = 2$: later 1, 4, 3, 6, 5 $\rightarrow$ discordant: 1 (value 1); concordant: 4 (4, 3, 6, 5).
- $i = 2, Y_2 = 1$: later 4, 3, 6, 5 $\rightarrow$ all concordant (4).
- $i = 3, Y_3 = 4$: later 3, 6, 5 $\rightarrow$ discordant: 1 (3); concordant: 2 (6, 5).
- $i = 4, Y_4 = 3$: later 6, 5 $\rightarrow$ both concordant (2).
- $i = 5, Y_5 = 6$: later 5 $\rightarrow$ discordant: 1 (5).

$C = 4 + 4 + 2 + 2 = 12$, $D = 1 + 1 + 1 = 3$. Total pairs $= \binom{6}{2} = 15$, $C + D = 15 \equiv$. $S = C - D = 9$.

$$\tau = \frac{2S}{n(n-1)} = \frac{18}{6 \times 5} = \frac{18}{30} = 0.6.$$

3. Comparison and Comment. Both coefficients indicate strong positive agreement between the two teachers. $R_s \approx 0.83$ is higher than $\tau = 0.60$. This is typical: Spearman's coefficient tends to be larger in magnitude than Kendall's for the same data because R_s uses the squared rank differences, while τ uses only the sign. Kendall's τ has a more direct probabilistic interpretation: the probability that a randomly chosen pair is concordant minus the probability it is discordant. Here, $\tau = 0.6$ means concordant pairs exceed discordant pairs by 60% of all pairs. The teachers' rankings are very similar; the only disagreements are the swaps of adjacent pairs ($A \leftrightarrow B$, $C \leftrightarrow D$, $E \leftrightarrow F$).

DID YOU KNOW? Historical Note

Charles Spearman (1863–1945) developed his rank correlation while studying intelligence testing. Maurice Kendall (1907–1983) introduced τ in 1938 as a measure of rank correlation with a simpler sampling distribution. Both are British statisticians. In Cameroon, rank correlation is often used in educational research to compare rankings of schools or students by different examiners, and in agricultural trials where farmers rank crop varieties by preference.

EXERCISE 8 Practice: Rank Correlation

1. Two coaches rank 7 athletes in a 100m sprint trial. The ranks are:

Athlete	1	2	3	4	5	6	7
Coach X	1	2	3	4	5	6	7
Coach Y	3	1	2	5	4	7	6

Calculate (a) Spearman's R_s , (b) Kendall's τ . Interpret.

2. In a beauty contest, 5 judges rank 8 contestants. Explain why Kendall's τ might be preferred over Spearman's R_s for measuring agreement between two specific judges.
3. A researcher collects data on the ranking of 10 villages by access to healthcare and literacy rate. The data contain several tied ranks. Describe the correct procedure to compute Spearman's coefficient in this case.

SOLUTION Exercise 1

(a) $d_i = R_X - R_Y$: $(-2, 1, 1, -1, 1, -1, 1)$; $\sum d_i^2 = 4 + 1 + 1 + 1 + 1 + 1 + 1 = 10$. $R_s = 1 - \frac{6 \times 10}{7(49-1)} = 1 - \frac{60}{336} = 1 - 0.1786 = 0.8214$.

(b) $Y = (3, 1, 2, 5, 4, 7, 6)$. Count pairs: $C = 15$, $D = 6$ (details omitted). $\tau = \frac{2(15-6)}{7 \times 6} = \frac{18}{42} = 0.4286$. Interpretation: Strong positive agreement by Spearman; moderate agreement by Kendall. The coaches generally agree on the top athletes but differ on exact ordering.

SOLUTION Exercise 2

Kendall's τ is based on pairwise concordance, which directly reflects whether two judges agree on the relative order of each pair of contestants. It is less affected by the magnitude of rank differences and handles small samples well. With 5 judges, one might compute τ for each pair of judges to assess inter-rater reliability.

SOLUTION Exercise 3

When ties are present, assign the average rank to each tied observation. Then compute the Pearson correlation coefficient on these averaged ranks (i.e., use the definition formula for R_s with the ranked data). Do not use the simplified formula $1 - \frac{6 \sum d^2}{n(n^2-1)}$ because it assumes no ties.

8 INTEGRATION ACTIVITY

9 REGRESSION AND CORRELATION

9.1 Introduction to Regression and Scatter Diagrams

In many scientific, economic, and educational contexts, we observe two variables simultaneously and wish to understand how they relate. For instance, a teacher may suspect that the time a student spends revising influences their examination score. Regression analysis provides tools to model the relationship between a dependent variable (response) and one or more independent variables (explanatory). Correlation measures the strength and direction of the association.

DEFINITION Bivariate Data

A set of paired observations (x_i, y_i) , $i = 1, 2, \dots, n$, where x is the independent (explanatory) variable and y is the dependent (response) variable.


DEFINITION Scatter Diagram

A graphical display of bivariate data where each pair (x_i, y_i) is plotted as a point on a Cartesian plane with x on the horizontal axis and y on the vertical axis. It gives a visual impression of the relationship: linear, curved, clustered, or no pattern.

EXAMPLE Scatter Diagram for Revision Hours vs Mock Scores

Consider the data collected by Group A in a Yaoundé bilingual high school:

Student	Hours of revision per week (x)	Mock score (y)
1	2	42
2	3	48
3	4	55
4	5	52
5	6	65
6	7	62
7	8	75
8	9	70
9	10	85
10	11	82
11	12	90



Scatter diagram of mock scores against revision hours.

Comment: The points show a clear upward trend. As revision hours increase, mock scores tend to increase. The points are roughly clustered around a straight line, suggesting a strong positive linear relationship.

9.2 Mayer's Method for the Regression Line

Mayer's method (also called the double mean or median–median method) is a simple graphical technique to estimate a regression line by using the mean points of two halves of the data.

METHOD Mayer's Method – Steps

1. Order the data by increasing x (if not already ordered).
2. Split the n observations into two groups of equal size (or as equal as possible). If n is odd, the middle observation is usually omitted or the groups are split as $\lfloor n/2 \rfloor$ and $\lceil n/2 \rceil$.
3. Calculate the mean point $(\bar{x}_1, \bar{y}_1)$ of the first group and $(\bar{x}_2, \bar{y}_2)$ of the second group.
4. Draw the line passing through these two mean points. Its equation is $y = a_M + b_M x$ where

$$b_M = \frac{\bar{y}_2 - \bar{y}_1}{\bar{x}_2 - \bar{x}_1}, \quad a_M = \bar{y}_1 - b_M \bar{x}_1 \quad (\text{or using } \bar{y}_2, \bar{x}_2).$$

WARNING Limitations of Mayer's Method

Mayer's line depends on the arbitrary split of the data and does not use all points optimally. It is not the "line of best fit" in the least squares sense. It is mainly a pedagogical tool or a quick approximation.

9.3 Correlation, Sample Variance and Covariance

To quantify the linear relationship, we need measures of spread and joint variability.

DEFINITION Sample Variance and Covariance

For a sample $(x_i, y_i), i = 1, \dots, n$, with means $\bar{x}, \bar{y}$:

- Corrected sum of squares for x : $S_{xx} = \sum (x_i - \bar{x})^2 = \sum x_i^2 - \frac{(\sum x_i)^2}{n}$.
- Corrected sum of squares for y : $S_{yy} = \sum (y_i - \bar{y})^2 = \sum y_i^2 - \frac{(\sum y_i)^2}{n}$.
- Corrected sum of cross-products: $S_{xy} = \sum (x_i - \bar{x})(y_i - \bar{y}) = \sum x_i y_i - \frac{\sum x_i \sum y_i}{n}$.

The sample covariance is $\frac{S_{xy}}{n-1}$ (or $\frac{S_{xy}}{n}$ for the population analogue). The sample variances are $\frac{S_{xx}}{n-1}$ and $\frac{S_{yy}}{n-1}$.

PROPERTY Properties of S_{xx}, S_{yy}, S_{xy}

- $S_{xx} \geq 0, S_{yy} \geq 0$; equality holds iff all x (resp. y) are equal.
- S_{xy} can be positive, negative, or zero.
- By the Cauchy–Schwarz inequality: $S_{xy}^2 \leq S_{xx}S_{yy}$.

9.4 Least Squares Regression Lines

The least squares criterion chooses the line that minimises the sum of squared vertical distances (residuals) between the observed y_i and the fitted values $\hat{y}_i$.

PROPERTY Least Squares Regression Line of y on x

The line $y = a + bx$ that minimises $\sum (y_i - a - bx_i)^2$ has slope and intercept:

$$b = \frac{S_{xy}}{S_{xx}}, \quad a = \bar{y} - b\bar{x}.$$

The line always passes through the mean point $(\bar{x}, \bar{y})$.

PROPERTY Least Squares Regression Line of x on y

If we instead minimise horizontal distances (treat x as response), the line $x = a' + b'y$ has:

$$b' = \frac{S_{xy}}{S_{yy}}, \quad a' = \bar{x} - b'\bar{y}.$$

This line also passes through $(\bar{x}, \bar{y})$. In general, the two lines are different unless $|r| = 1$.

KEY POINTS Interpretation of Slopes

- $b = \frac{S_{xy}}{S_{xx}}$: estimated change in y for a one-unit increase in x .
- $b' = \frac{S_{xy}}{S_{yy}}$: estimated change in x for a one-unit increase in y .
- $b \cdot b' = r^2$ (the coefficient of determination).

9.5 Residual Sum of Squares and the Line of Best Fit

DEFINITION Residual

For the regression line of y on x , the residual for observation i is $e_i = y_i - \hat{y}_i = y_i - (a + bx_i)$.

PROPERTY Residual Sum of Squares (RSS)

- For y on x : $\text{RSS}_{y|x} = \sum e_i^2 = S_{yy} - bS_{xy} = S_{yy} - \frac{S_{xy}^2}{S_{xx}}$.
- For x on y : $\text{RSS}_{x|y} = \sum (x_i - \hat{x}_i)^2 = S_{xx} - b'S_{xy} = S_{xx} - \frac{S_{xy}^2}{S_{yy}}$.

The least squares line is the line of best fit because it minimises the respective RSS among all straight lines.

WARNING Residuals vs Errors

Residuals are computed from the sample; errors are the unobservable true deviations. Residuals sum to zero ($\sum e_i = 0$) and are uncorrelated with x ($\sum x_i e_i = 0$) for the least squares line.

9.6 The Product Moment Correlation Coefficient

DEFINITION Pearson's Product Moment Correlation Coefficient (r)

$$r = \frac{S_{xy}}{\sqrt{S_{xx}S_{yy}}}.$$

It measures the strength and direction of the linear relationship between x and y .

PROPERTY Properties of r

- $-1 \leq r \leq 1$.
- $r = 1$: perfect positive linear relationship (all points on a line with positive slope).
- $r = -1$: perfect negative linear relationship.
- $r = 0$: no linear relationship (but there could be a non-linear one).

- r is invariant under linear changes of scale: $r(ax + b, cy + d) = \text{sign}(ac) \cdot r(x, y)$.
- $r^2 = \frac{\text{Explained variation}}{\text{Total variation}} = 1 - \frac{\text{RSS}}{S_{yy}}$ is the coefficient of determination.

WARNING Correlation $\neq$ Causation

A high $|r|$ does not imply that x causes y . There may be confounding variables, reverse causality, or coincidence. In our example, revision hours may correlate with scores, but motivation, prior ability, and teaching quality also play roles.

9.7 Rank Correlation: Spearman and Kendall

When data are not interval-scaled, contain outliers, or the relationship is monotonic but not linear, rank correlation coefficients are preferred.

DEFINITION Spearman's Rank Correlation Coefficient (R_s)

Replace each x_i and y_i by their ranks $R(x_i), R(y_i)$ (average ranks for ties). Then

$$R_s = 1 - \frac{6 \sum d_i^2}{n(n^2 - 1)}, \quad \text{where } d_i = R(x_i) - R(y_i),$$

provided there are no ties. With ties, use the Pearson formula on the ranks.

DEFINITION Kendall's Rank Correlation Coefficient (τ)

Consider all $\binom{n}{2}$ pairs of observations (i, j) with $i < j$. A pair is concordant if $(x_i - x_j)(y_i - y_j) > 0$ and discordant if < 0 . Let C and D be the counts. Then

$$\tau = \frac{C - D}{C + D} = \frac{C - D}{\binom{n}{2}}.$$

(If ties exist, a modified denominator is used: τ_b or τ_c .)

KEY POINTS Comparison of Correlation Measures

Measure	Assesses	Assumptions	Robustness
Pearson r	Linear relationship	Bivariate normality, linearity, interval data	Sensitive to outliers
Spearman R_s	Monotonic relationship	Ordinal or continuous data; ranks used	Less sensitive to outliers
Kendall τ	Concordance (pair-wise agreement)	Ordinal data; ranks used	Most robust; smaller magnitude typically

WARNING Uses and Limitations of Kendall's τ

- Uses: Small samples, ordinal data, many tied ranks, robust alternative to Spearman.
- Limitations: Computationally heavier for large n (though $O(n \log n)$ algorithms exist); τ tends to be smaller in magnitude than R_s for the same data; interpretation as “probability of concordance minus probability of discordance” is less intuitive for non-statisticians.

9.8 Integration Activity: Revision Hours and Mock Scores

9.9 Aims of the Activity

- Consolidate the understanding of regression and correlation techniques covered in the chapter.
- Apply scatter diagrams, Mayer's method, least squares regression, residual analysis, product moment correlation, Spearman's rank correlation, and Kendall's rank correlation to a real dataset.
- Develop skills in data collection, organisation, computation, interpretation, and report writing.
- Critically evaluate the reliability of predictions and compare different correlation measures.
- Foster collaborative work and communication of statistical findings in a Cameroonian educational context.

9.10 Situation

In many Cameroonian secondary schools, Upper Sixth students prepare intensely for the GCE Advanced Level Mathematics examination. A common belief among students and teachers is that the number of hours devoted to personal revision each week strongly influences the mock examination scores. To investigate this claim, the Mathematics teacher of a bilingual high school in Yaoundé proposes an integration activity. The class is divided into groups of four to five stu-

dents. Each group must select 10–12 volunteer classmates (different from the group members) and record two variables for each volunteer:

- x : average number of hours of personal revision per week (over the last month),
- y : score obtained in the recent Mathematics mock examination (out of 100).

The data collected by one group (Group A) is shown in the table below. This data will be used throughout the activity to illustrate the required computations. Each group will use their own data, but the steps are identical.

Student	Hours of revision per week (x)	Mock score (y)
1	2	42
2	3	48
3	4	55
4	5	52
5	6	65
6	7	62
7	8	75
8	9	70
9	10	85
10	11	82
11	12	90

The teacher asks each group to carry out a complete statistical analysis and write a short report. The activity culminates in a plenary session where groups compare their results and discuss the strengths and limitations of each method.

9.11 Tasks

1. **Scatter diagram:** Plot the data on a scatter diagram with x on the horizontal axis and y on the vertical axis. Comment on the pattern observed.
2. **Mayer's method:**
 - (a) Order the data by x (already done). Split the 11 observations into two groups: the first 5 and the last 6 (since $n = 11$ is odd).
 - (b) Calculate the mean point $(\bar{x}_1, \bar{y}_1)$ of the first group and $(\bar{x}_2, \bar{y}_2)$ of the second group.
 - (c) Draw the line passing through these two mean points on the scatter diagram. Find its equation in the form $y = a + bx$.
3. **Least squares regression lines:**
 - (a) Compute the corrected sums of squares and cross-products: S_{xx}, S_{yy}, S_{xy} .
 - (b) Determine the least squares estimates of the slope b and intercept a for the regression line $y = a + bx$ (y on x).

- (c) Determine the least squares estimates for the regression line $x = a' + b'y$ (x on y).
- (d) Write both equations and plot the y on x line on the scatter diagram.

4. Residual Sum of Squares (RSS) and Product Moment Correlation Coefficient:

- (a) Calculate the Residual Sum of Squares (RSS) for the least squares line of y on x and for x on y .
- (b) Compute the product moment correlation coefficient r .
- (c) Interpret the value of r and r^2 in the context of the problem.

5. Rank correlation:

- (a) Assign ranks to the x values (already 1 to 11) and to the y values (handle ties if any).
- (b) Compute Spearman's rank correlation coefficient R_s .
- (c) Compute Kendall's rank correlation coefficient τ .
- (d) Compare the three correlation coefficients (r , R_s , τ) and comment on their similarities and differences.

6. Prediction and report:


- (a) Use the least squares regression line of y on x to predict the mock score of a student who revises 9 hours per week.
- (b) Discuss the reliability of this prediction, considering the value of r , the scatter diagram, the sample size, and whether 9 hours lies within the range of the data.
- (c) Write a concise report (about 200 words) summarising the findings, the prediction, and a comparison of the three correlation measures. State any limitations of the study.

9.12 Model Answer for Group A's Data

WORKED EXAMPLE 13 Model Answer

SOLUTION

1. Scatter Diagram The scatter diagram below shows the 11 data points. There is a clear upward trend: as revision hours increase, mock scores tend to increase. The points are roughly clustered around a straight line, suggesting a strong positive linear relationship.



Scatter diagram with least squares regression lines and Mayer's line.

2. Mayer's Method The data is already ordered by x . Split into two groups:

- Group 1 (first 5): (2, 42), (3, 48), (4, 55), (5, 52), (6, 65)
- Group 2 (last 6): (7, 62), (8, 75), (9, 70), (10, 85), (11, 82), (12, 90)

Mean points:

$$\bar{x}_1 = \frac{2+3+4+5+6}{5} = 4, \quad \bar{y}_1 = \frac{42+48+55+52+65}{5} = \frac{262}{5} = 52.4$$

$$\bar{x}_2 = \frac{7+8+9+10+11+12}{6} = \frac{57}{6} = 9.5, \quad \bar{y}_2 = \frac{62+75+70+85+82+90}{6} = \frac{464}{6} \approx 77.333$$

Slope of Mayer's line:

$$b_M = \frac{\bar{y}_2 - \bar{y}_1}{\bar{x}_2 - \bar{x}_1} = \frac{77.333 - 52.4}{9.5 - 4} = \frac{24.933}{5.5} \approx 4.533$$

Intercept:

$$a_M = \bar{y}_1 - b_M \bar{x}_1 = 52.4 - 4.533 \times 4 = 52.4 - 18.132 = 34.268$$

Equation of Mayer's line: $y = 34.27 + 4.53x$ (rounded to two decimals).

3. Least Squares Regression Lines We compute the necessary sums:

$$\begin{aligned} n &= 11 \\ \sum x &= 77, \quad \sum y = 726 \\ \sum x^2 &= 649, \quad \sum y^2 = 50440 \\ \sum xy &= 5594 \end{aligned}$$

Means:

$$\bar{x} = \frac{77}{11} = 7, \quad \bar{y} = \frac{726}{11} = 66$$

Corrected sums:

$$S_{xx} = \sum x^2 - \frac{(\sum x)^2}{n} = 649 - \frac{5929}{11} = 649 - 539 = 110$$

$$S_{yy} = \sum y^2 - \frac{(\sum y)^2}{n} = 50440 - \frac{527076}{11} = 50440 - 47916 = 2524$$

$$S_{xy} = \sum xy - \frac{\sum x \sum y}{n} = 5594 - \frac{77 \times 726}{11} = 5594 - 5082 = 512$$

Line of y on x:

$$b = \frac{S_{xy}}{S_{xx}} = \frac{512}{110} = \frac{256}{55} \approx 4.6545$$

$$a = \bar{y} - b\bar{x} = 66 - \frac{256}{55} \times 7 = 66 - \frac{1792}{55} = \frac{3630 - 1792}{55} = \frac{1838}{55} \approx 33.4182$$

Equation: $y = 33.42 + 4.65x$ (rounded to two decimals).

Line of x on y:

$$b' = \frac{S_{xy}}{S_{yy}} = \frac{512}{2524} = \frac{128}{631} \approx 0.2029$$

$$a' = \bar{x} - b'\bar{y} = 7 - \frac{128}{631} \times 66 = 7 - \frac{8448}{631} = \frac{4417 - 8448}{631} = -\frac{4031}{631} \approx -6.388$$

Equation: $x = -6.39 + 0.203y$ (or $y = 31.48 + 4.93x$ when rearranged for plotting).

Both lines pass through $(\bar{x}, \bar{y}) = (7, 66)$. The y on x line is plotted in solid orange; the x on y line (rearranged) in dotted purple. They differ because they minimise different distances (vertical vs horizontal).

4. Residual Sum of Squares (RSS) and Product Moment Correlation Coefficient

$$RSS_{y|x} = S_{yy} - bS_{xy} = 2524 - \frac{512}{110} \times 512 = 2524 - \frac{262144}{110} = 2524 - 2383.127 = 140.873$$

$$\text{Alternatively, } RSS_{y|x} = S_{yy} - \frac{S_{xy}^2}{S_{xx}} = 2524 - \frac{512^2}{110} = 140.873.$$

$$RSS_{x|y} = S_{xx} - b'S_{xy} = 110 - \frac{512}{2524} \times 512 = 110 - \frac{262144}{2524} = 110 - 103.861 = 6.139$$

Product moment correlation coefficient:

$$r = \frac{S_{xy}}{\sqrt{S_{xx}S_{yy}}} = \frac{512}{\sqrt{110 \times 2524}} = \frac{512}{\sqrt{277640}} = \frac{512}{526.915} \approx 0.9717$$

$$r^2 = (0.9717)^2 \approx 0.9442$$

Interpretation: $r = 0.972$ indicates a very strong positive linear correlation between hours of revision and mock scores. About 94.4% of the variation in mock scores is explained by the linear relationship with revision hours (coefficient of determination r^2). The remaining 5.6% is due to other factors or random variation.

5. Rank Correlation Ranks of x are simply $1, 2, \dots, 11$ because the x values are distinct and already sorted. Ranks of y : sort the y values and assign ranks 1 to 11.

Student	1	2	3	4	5	6	7	8	9	10	11
y	42	48	55	52	65	62	75	70	85	82	90
Rank of y	1	2	4	3	6	5	8	7	10	9	11

Differences $d = \text{rank}_x - \text{rank}_y$:

$$d : 0, 0, -1, 1, -1, 1, -1, 1, -1, 1, 0$$

$$\sum d^2 = 0 + 0 + 1 + 1 + 1 + 1 + 1 + 1 + 1 + 1 + 0 = 8$$

Spearman's rank correlation coefficient (no ties):

$$R_s = 1 - \frac{6 \sum d^2}{n(n^2 - 1)} = 1 - \frac{6 \times 8}{11 \times 120} = 1 - \frac{48}{1320} = 1 - 0.03636 = 0.9636$$

Kendall's τ : Count concordant (C) and discordant (D) pairs. Since x ranks are in natural order, discordant pairs correspond to inversions in the sequence of y ranks: $[1, 2, 4, 3, 6, 5, 8, 7, 10, 9, 11]$. Inversions: $(4, 3), (6, 5), (8, 7), (10, 9) \rightarrow 4$ discordant pairs. Total pairs $= \binom{11}{2} = 55$. So $C = 51, D = 4$.

$$\tau = \frac{C - D}{C + D} = \frac{51 - 4}{55} = \frac{47}{55} \approx 0.8545$$

Comparison:

- $r = 0.972$ (Pearson) measures linear correlation; assumes bivariate normality and linearity.
- $R_s = 0.964$ (Spearman) measures monotonic correlation; based on ranks, less sensitive to outliers.
- $\tau = 0.855$ (Kendall) measures concordance; more robust but generally smaller in magnitude.

All three indicate a very strong positive association. R_s is closer to r because the relationship is nearly linear. τ is lower because it counts pairwise concordances, giving less weight to the overall trend.

6. Prediction and Report Prediction for $x = 9$ hours using the least squares line of y on x :

$$\hat{y} = 33.418 + 4.655 \times 9 = 33.418 + 41.895 = 75.313 \approx 75.3$$

A student revising 9 hours per week is predicted to score about 75 out of 100 in the mock examination.

Reliability Discussion:

- The correlation is very strong ($r = 0.972$), and the scatter diagram shows points tightly clustered around the line.

- The prediction is within the range of the data ($x = 9$ is between 2 and 12), so it is an interpolation, not an extrapolation.
- However, the sample size is only 11, which is small. The RSS is about 141, giving a residual standard deviation of $s = \sqrt{\text{RSS}/(n-2)} \approx \sqrt{141/9} \approx 3.95$. A rough 95% prediction interval would be $\hat{y} \pm 2 \times 3.95 \approx 75.3 \pm 7.9$, i.e., between 67 and 83.
- The relationship might not be perfectly linear for extreme values, and other factors (teaching quality, prior knowledge, health) also affect scores.

Overall, the prediction is reasonably reliable but should be treated as an estimate with a margin of error.

Report

Our investigation on 11 Upper Sixth students reveals a very strong positive relationship between weekly revision hours and Mathematics mock scores. The scatter diagram shows a clear linear trend. Mayer's method gave the line $y = 34.27 + 4.53x$, while the least squares line of y on x is $y = 33.42 + 4.65x$ and the line of x on y is $x = -6.39 + 0.203y$. The product moment correlation coefficient is $r = 0.972$, indicating that 94.4% of the variation in scores is explained by revision hours. Rank correlations confirm the strong association: Spearman's $R_s = 0.964$ and Kendall's $\tau = 0.855$. Using the least squares line, a student revising 9 hours per week is predicted to score about 75. This prediction is fairly reliable because the correlation is strong and 9 hours lies within our data range. However, the small sample size (11) limits the precision; a larger study across multiple schools would improve generalisability. All three correlation measures agree on a strong positive link, though Kendall's τ is more conservative. We recommend that students aim for at least 9 hours of weekly revision to target scores above 70, but they should also focus on effective study techniques.

9.13 Exercises

EXERCISE 9 Exercise 1: Least Squares Calculations

For the data set: (1, 2), (2, 3), (3, 5), (4, 4), (5, 6).

1. Compute $\bar{x}, \bar{y}, S_{xx}, S_{yy}, S_{xy}$.
2. Find the least squares regression line of y on x and of x on y .
3. Calculate $r, r^2, \text{RSS}_{y|x}, \text{RSS}_{x|y}$.
4. Plot the data and both lines. Comment on their difference.

EXERCISE 10 Exercise 2: Rank Correlation with Ties

The following data show the ranks of 8 students in Mathematics (x) and Physics (y):

Student	A	B	C	D	E	F	G	H
Math rank	1	2	3	4	5	6	7	8
Physics rank	2	1	4	3	6	5	8	7

1. Compute Spearman's R_s using the d^2 formula.
2. Compute Kendall's τ by counting concordant/discordant pairs.
3. If the Physics ranks had a tie (e.g., students C and D both ranked 3.5), how would you compute R_s ?

EXERCISE 11 Exercise 3: Interpretation and Limitations

A study finds $r = 0.85$ between the number of ice creams sold daily and the number of drowning incidents at a beach.

1. Does this mean eating ice cream causes drowning? Explain.
2. Suggest a confounding variable.
3. Why might Spearman's R_s be preferred over Pearson's r if the relationship is curved but monotonic?

SOLUTION Exercise 1

1. $\sum x = 15, \sum y = 20, \sum x^2 = 55, \sum y^2 = 90, \sum xy = 69, n = 5, \bar{x} = 3, \bar{y} = 4, S_{xx} = 55 - 225/5 = 10, S_{yy} = 90 - 400/5 = 10, S_{xy} = 69 - 300/5 = 9.$
2. $b = 9/10 = 0.9, a = 4 - 0.9 \times 3 = 1.3 \rightarrow y = 1.3 + 0.9x. b' = 9/10 = 0.9, a' = 3 - 0.9 \times 4 = -0.6 \rightarrow x = -0.6 + 0.9y.$
3. $r = 9/\sqrt{100} = 0.9, r^2 = 0.81. RSS_{y|x} = 10 - 0.9 \times 9 = 1.9. RSS_{x|y} = 10 - 0.9 \times 9 = 1.9$ (symmetric because $S_{xx} = S_{yy}$).
4. The lines cross at $(3, 4)$. They differ because one minimises vertical, the other horizontal distances.

SOLUTION Exercise 2

1. $d: -1, 1, -1, 1, -1, 1, -1, 1 \rightarrow \sum d^2 = 8$. $R_s = 1 - \frac{6 \times 8}{8 \times 63} = 1 - \frac{48}{504} = 1 - 0.0952 = 0.9048$.
2. Physics ranks in Math order: $[2, 1, 4, 3, 6, 5, 8, 7]$. Inversions: $(2, 1), (4, 3), (6, 5), (8, 7) \rightarrow D = 4$. Total pairs = 28. $C = 24$. $\tau = (24 - 4)/28 = 20/28 = 0.714$.
3. With ties, use the Pearson formula on the ranks (or the adjusted formula $R_s = 1 - \frac{6 \sum d^2 + \frac{1}{2} \sum (t^3 - t)}{n(n^2 - 1)}$ where t are tie sizes).

SOLUTION Exercise 3

1. No. Correlation does not imply causation. It is a spurious correlation.
2. Temperature (or season): hot weather increases both ice cream sales and swimming (hence drowning risk).
3. Pearson's r measures only linear association; a curved monotonic relationship would give $|r| < 1$ even if perfectly monotonic. Spearman's R_s uses ranks and captures any monotonic trend.

10 METHODS AND WORKED EXAMPLES

10.1 Introduction to Regression and Scatter Diagrams

DEFINITION Bivariate Data and Variables

When we collect data on two variables for each individual or observation, we have **bivariate data**. The variable we think might explain or influence the other is called the **independent variable** (or **explanatory variable**), usually denoted x . The variable we think might respond to changes in x is the **dependent variable** (or **response variable**), usually denoted y . In the GCE Advanced Level examination, x is always plotted on the horizontal axis and y on the vertical axis.

METHOD Drawing and interpreting a scatter diagram

1. Identify the variables: Decide which variable is x (independent) and which is y (dependent). Label axes clearly with variable names and units.
2. Choose appropriate scales: Select scales so that the data points occupy most of the graph paper. The origin $(0, 0)$ need not be included unless it is meaningful for the context. Use a broken axis symbol $(\sim)$ if the scale does not start at zero.
3. Plot the points: For each pair (x_i, y_i) , mark a point using a sharp pencil and small crosses ($\times$) or dots ($\bullet$). Do not join the points.
4. Calculate and plot the mean point: Compute $\bar{x} = \frac{1}{n} \sum x_i$ and $\bar{y} = \frac{1}{n} \sum y_i$. Plot $(\bar{x}, \bar{y})$ clearly (e.g., with a ringed dot $\odot$). Key reflex: Any reasonable regression line must pass through $(\bar{x}, \bar{y})$.
5. Observe the pattern: Look for a general trend:
 - **Positive linear correlation**: points tend to rise from left to right, roughly following a straight line.
 - **Negative linear correlation**: points tend to fall from left to right, roughly following a straight line.
 - **No linear correlation**: points appear randomly scattered with no discernible linear trend.
 - **Non-linear pattern**: points follow a curve (e.g., quadratic, exponential). In this case, linear regression is not appropriate unless the data is transformed.
6. Draw a line of best fit by eye (optional): If the pattern appears roughly linear, draw a straight line that passes as close as possible to all points, with roughly equal numbers

of points above and below the line, and passing through $(\bar{x}, \bar{y})$.

- Interpret the line: The slope (gradient) indicates the rate of change of y with respect to x ; the intercept gives the value of y when $x = 0$ (only meaningful if $x = 0$ is within or near the data range).

WARNING Common mistakes with scatter diagrams

- Joining the points with line segments (this is a time-series plot, not a scatter diagram).
- Choosing scales that squash all points into a tiny corner.
- Forgetting to label axes with variable names and units.
- Drawing a line of best fit that does not pass through $(\bar{x}, \bar{y})$.
- Interpreting the y -intercept when $x = 0$ is far outside the data range (extrapolation).

WORKED EXAMPLE 14 Scatter diagram for height and weight of students

The table below shows the heights (in cm) and weights (in kg) of 10 Upper Sixth students in a college in Yaoundé.


Height x	160	165	170	175	180	185	190	195	200	205
Weight y	55	58	62	65	70	73	78	82	85	90

- Draw a scatter diagram for this data.
- Comment on the correlation.
- Draw a line of best fit by eye and estimate the weight of a student of height 178 cm.

SOLUTION

Solution.

- Scatter diagram: We choose the horizontal axis for height (160–205 cm) and vertical axis for weight (55–90 kg). Plot each pair (x_i, y_i) . The points are: (160, 55), (165, 58), (170, 62), (175, 65), (180, 70), (185, 73), (190, 78), (195, 82), (200, 85), (205, 90).



Scatter diagram of height vs weight for 10 students

2. Comment: The points lie very close to a straight line sloping upwards. There is a strong **positive linear correlation** between height and weight. As height increases, weight tends to increase in an approximately linear fashion.
3. Line of best fit by eye: The mean height $\bar{x} = \frac{160+165+\dots+205}{10} = 182.5$ cm. The mean weight $\bar{y} = \frac{55+58+\dots+90}{10} = 71.8$ kg. The mean point is $(182.5, 71.8)$. Draw a straight line through this point that follows the trend of the points. Using the graph, at height 178 cm (which corresponds to $x = 178$), the line gives a weight of approximately 67 kg. (Reading from the graph: between 175 cm (65 kg) and 180 cm (70 kg), 178 cm is about 67 kg.)

10.2 Mayer's Method for the Regression Line (Double Mean Method)

DEFINITION Mayer's Method

Mayer's method (also called the **double mean method**) is a simple arithmetic technique to find an approximate regression line when the scatter diagram suggests a linear relationship. It uses the mean points of two halves of the data ordered by x . It is less accurate than the least squares method but is useful for quick estimation and is examinable in the GCE Advanced Level syllabus.

METHOD Mayer's method (step by step)

1. Order the data by the independent variable x in ascending order.

2. Split the data into two groups of equal size.

- If n is even: first $n/2$ points form Group 1 (lower x), last $n/2$ points form Group 2 (higher x).
- If n is odd: the usual GCE approach is to omit the middle observation, or include it in both groups. We will adopt: omit the middle observation, so Group 1 has the first $(n-1)/2$ points, Group 2 has the last $(n-1)/2$ points.

3. Compute the mean point of each group: $(\bar{x}_1, \bar{y}_1)$ for Group 1 and $(\bar{x}_2, \bar{y}_2)$ for Group 2.

4. Calculate the gradient: $m = \frac{\bar{y}_2 - \bar{y}_1}{\bar{x}_2 - \bar{x}_1}$. (If $\bar{x}_2 = \bar{x}_1$, the method fails — vertical line.)

5. Find the equation: Using point-slope form: $y - \bar{y}_1 = m(x - \bar{x}_1)$, or $y = mx + c$ with $c = \bar{y}_1 - m\bar{x}_1$.

6. Verify: The line should pass through (or very close to) the overall mean point $(\bar{x}, \bar{y})$.

Key formula: $m = \frac{\bar{y}_2 - \bar{y}_1}{\bar{x}_2 - \bar{x}_1}$.

WARNING When Mayer's method is inappropriate

- If the scatter diagram shows a clear non-linear pattern.
- If there are extreme outliers that distort the group means.
- If the two group mean points are very close together (small denominator), making the gradient unstable.
- For precise work, the least squares method (next section) is always preferred.

WORKED EXAMPLE 15 Mayer's method for advertising and sales

A company records its monthly advertising expenditure (in thousands of CFA francs) and the corresponding sales (in millions of CFA francs) for 8 months:

Advertising x	2	3	4	5	6	7	8	9
Sales y	10	12	14	15	18	20	22	24

Use Mayer's method to find the regression line of sales on advertising expenditure. Hence estimate the sales when advertising expenditure is 6.5 thousand CFA francs.

SOLUTION

Solution.

1. The data is already ordered by x . $n = 8$ (even), so we split into two groups of 4. Group 1 (first 4): (2, 10), (3, 12), (4, 14), (5, 15). Group 2 (last 4): (6, 18), (7, 20), (8, 22), (9, 24).
2. Group means: Group 1: $\bar{x}_1 = \frac{2+3+4+5}{4} = \frac{14}{4} = 3.5$, $\bar{y}_1 = \frac{10+12+14+15}{4} = \frac{51}{4} = 12.75$. Group 2: $\bar{x}_2 = \frac{6+7+8+9}{4} = \frac{30}{4} = 7.5$, $\bar{y}_2 = \frac{18+20+22+24}{4} = \frac{84}{4} = 21$.
3. Gradient: $m = \frac{\bar{y}_2 - \bar{y}_1}{\bar{x}_2 - \bar{x}_1} = \frac{21 - 12.75}{7.5 - 3.5} = \frac{8.25}{4} = 2.0625$.
4. Equation: Using point $(\bar{x}_1, \bar{y}_1) = (3.5, 12.75)$: $y - 12.75 = 2.0625(x - 3.5)$ $y = 2.0625x - 7.21875 + 12.75 = 2.0625x + 5.53125$. So the Mayer regression line is $y = 2.0625x + 5.53125$. (Rounded to 3 significant figures: $y = 2.06x + 5.53$.)
5. Check overall mean: $\bar{x} = \frac{2+3+\dots+9}{8} = \frac{44}{8} = 5.5$, $\bar{y} = \frac{10+12+\dots+24}{8} = \frac{135}{8} = 16.875$. Plug $x = 5.5$ into line: $y = 2.0625(5.5) + 5.53125 = 11.34375 + 5.53125 = 16.875$. Exactly the overall mean. Good.
6. Estimate for $x = 6.5$: $y = 2.0625(6.5) + 5.53125 = 13.40625 + 5.53125 = 18.9375$ million CFA francs. (Rounded: 18.9 million CFA francs.)

10.3 Sample Variance, Covariance and Product Moment Correlation Coefficient

DEFINITION Corrected Sums of Squares and Products

For a bivariate sample $(x_i, y_i), i = 1, \dots, n$, with means $\bar{x}, \bar{y}$, we define the **corrected sums of squares and products**:

$$S_{xx} = \sum_{i=1}^n (x_i - \bar{x})^2 = \sum x_i^2 - \frac{(\sum x_i)^2}{n},$$

$$S_{yy} = \sum_{i=1}^n (y_i - \bar{y})^2 = \sum y_i^2 - \frac{(\sum y_i)^2}{n},$$

$$S_{xy} = \sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y}) = \sum x_i y_i - \frac{(\sum x_i)(\sum y_i)}{n}.$$

These quantities measure the spread of x , spread of y , and joint variability of x and y respectively. They are the building blocks for all regression and correlation calculations.

DEFINITION Sample Variance and Covariance

The **sample variances** and **sample covariance** are:

$$s_x^2 = \frac{S_{xx}}{n-1}, \quad s_y^2 = \frac{S_{yy}}{n-1}, \quad s_{xy} = \frac{S_{xy}}{n-1}.$$

The denominator $n - 1$ makes them unbiased estimators of the population variance and covariance.

DEFINITION Product Moment Correlation Coefficient (Pearson's r)

The **product moment correlation coefficient** (Pearson's r) measures the strength and direction of the **linear** relationship between x and y :

$$r = \frac{S_{xy}}{\sqrt{S_{xx}S_{yy}}} = \frac{s_{xy}}{s_x s_y}.$$

Properties:

- $-1 \leq r \leq 1$ always.
- $r = 1$: perfect positive linear correlation (all points lie exactly on a line with positive slope).
- $r = -1$: perfect negative linear correlation (all points lie exactly on a line with negative slope).
- $r = 0$: no linear correlation (but there could be a non-linear relationship).
- r is invariant under linear changes of scale: $r(ax + b, cy + d) = \text{sign}(ac) \cdot r(x, y)$.

METHOD Calculating S_{xx}, S_{yy}, S_{xy} , variances, covariance and r

1. Compute the sums: $\sum x_i, \sum y_i, \sum x_i^2, \sum y_i^2, \sum x_i y_i$.
2. Compute the corrected sums using the shortcut formulas: $S_{xx} = \sum x^2 - \frac{(\sum x)^2}{n}$, $S_{yy} = \sum y^2 - \frac{(\sum y)^2}{n}$, $S_{xy} = \sum xy - \frac{(\sum x)(\sum y)}{n}$.
3. Sample variances: $s_x^2 = \frac{S_{xx}}{n-1}$, $s_y^2 = \frac{S_{yy}}{n-1}$.
4. Sample covariance: $s_{xy} = \frac{S_{xy}}{n-1}$.
5. Correlation coefficient: $r = \frac{s_{xy}}{\sqrt{s_x^2 s_y^2}}$.

6. Interpret r in context.

Reflex: Always compute S_{xx}, S_{yy}, S_{xy} first; they are used in regression lines as well. Keep extra decimal places during calculation; round only the final answer.

WORKED EXAMPLE 16 Variance, covariance and correlation for a small dataset

For the following data:

x	2	4	6	8	10
y	3	5	7	9	11

Calculate:

1. S_{xx}, S_{yy}, S_{xy} .
2. The sample variances s_x^2, s_y^2 and covariance s_{xy} .
3. The product moment correlation coefficient r .
4. Comment on the result.

SOLUTION

Solution.

$n = 5$.

1. Sums: $\sum x = 2+4+6+8+10 = 30$, $\sum y = 3+5+7+9+11 = 35$. $\sum x^2 = 4+16+36+64+100 = 220$. $\sum y^2 = 9+25+49+81+121 = 285$. $\sum xy = 2 \cdot 3 + 4 \cdot 5 + 6 \cdot 7 + 8 \cdot 9 + 10 \cdot 11 = 6+20+42+72+110 = 250$.
2. Corrected sums: $S_{xx} = \sum x^2 - \frac{(\sum x)^2}{n} = 220 - \frac{900}{5} = 220 - 180 = 40$. $S_{yy} = \sum y^2 - \frac{(\sum y)^2}{n} = 285 - \frac{1225}{5} = 285 - 245 = 40$. $S_{xy} = \sum xy - \frac{(\sum x)(\sum y)}{n} = 250 - \frac{30 \cdot 35}{5} = 250 - 210 = 40$.
3. Sample variances and covariance: $s_x^2 = \frac{S_{xx}}{n-1} = \frac{40}{4} = 10$. $s_y^2 = \frac{S_{yy}}{n-1} = \frac{40}{4} = 10$. $s_{xy} = \frac{S_{xy}}{n-1} = \frac{40}{4} = 10$.
4. Correlation coefficient: $r = \frac{s_{xy}}{\sqrt{s_{xx}s_{yy}}} = \frac{40}{\sqrt{40 \cdot 40}} = \frac{40}{40} = 1$.
5. Comment: $r = 1$ indicates a perfect positive linear correlation. Indeed, the data satisfies $y = x + 1$ exactly; all points lie on a straight line with gradient 1.

DID YOU KNOW? Correlation vs Causation

A high correlation (close to ± 1) does not imply causation. For example, ice cream sales and drowning incidents are positively correlated, but eating ice cream does not cause drown-

ing; both are influenced by a third variable: hot weather. In the GCE exam, always state "there is evidence of a linear association" rather than "x causes y".

10.4 Least Squares Regression Line of y on x

DEFINITION Least Squares Regression Line of y on x

The **least squares regression line of y on x** is the line $y = a + bx$ that minimises the sum of squared vertical distances (residuals) between the observed y_i and the predicted values $\hat{y}_i = a + bx_i$:

$$\text{RSS} = \sum_{i=1}^n (y_i - \hat{y}_i)^2 = \sum_{i=1}^n (y_i - a - bx_i)^2.$$

This line is also called the **line of best fit** for predicting y from x.

PROPERTY Equation of the least squares line of y on x

The coefficients are given by:

$$b = \frac{S_{xy}}{S_{xx}}, \quad a = \bar{y} - b\bar{x}.$$

The equation can be written as $y = a + bx$ or equivalently $y - \bar{y} = b(x - \bar{x})$. Examinable proof: Derivation via the normal equations:

$$\begin{aligned} \sum y &= na + b \sum x, \\ \sum xy &= a \sum x + b \sum x^2. \end{aligned}$$

Solving these simultaneously yields the formulas above. You must be able to reproduce this derivation.

METHOD Finding the least squares regression line of y on x

1. Calculate $\sum x, \sum y, \sum x^2, \sum xy$ (and $\sum y^2$ if you also need r or S_{yy}).
2. Compute $S_{xx} = \sum x^2 - \frac{(\sum x)^2}{n}$ and $S_{xy} = \sum xy - \frac{(\sum x)(\sum y)}{n}$.
3. Compute the slope: $b = \frac{S_{xy}}{S_{xx}}$.

4. Compute the intercept: $a = \bar{y} - b\bar{x}$.
5. Write the equation: $y = a + bx$ (round a and b to 3 significant figures unless instructed otherwise).
6. Check: Verify that the line passes through $(\bar{x}, \bar{y})$.

Reflex: Never rearrange the y on x line to get the x on y line; they are different (unless $r = \pm 1$).

WORKED EXAMPLE 17 Least squares line of y on x for agricultural data

An agricultural researcher collects data on the amount of fertilizer applied (in kg/ha, x) and the resulting crop yield (in tonnes/ha, y) for 6 plots:

x	10	20	30	40	50	60
y	2.1	2.9	3.8	4.5	5.2	6.0

Find the equation of the least squares regression line of y on x . Use it to estimate the yield when 35 kg/ha of fertilizer is applied.

SOLUTION

Solution.

$$n = 6.$$

1. Sums: $\sum x = 10 + 20 + 30 + 40 + 50 + 60 = 210$. $\sum y = 2.1 + 2.9 + 3.8 + 4.5 + 5.2 + 6.0 = 24.5$.
 $\sum x^2 = 100 + 400 + 900 + 1600 + 2500 + 3600 = 9100$. $\sum y^2 = 4.41 + 8.41 + 14.44 + 20.25 + 27.04 + 36.00 = 110.55$. $\sum xy = 10(2.1) + 20(2.9) + 30(3.8) + 40(4.5) + 50(5.2) + 60(6.0) = 21 + 58 + 114 + 180 + 260 + 360 = 993$.
2. Means: $\bar{x} = 210/6 = 35$. $\bar{y} = 24.5/6 = 4.08333 \dots$
3. Corrected sums: $S_{xx} = \sum x^2 - \frac{(\sum x)^2}{n} = 9100 - \frac{44100}{6} = 9100 - 7350 = 1750$. $S_{xy} = \sum xy - \frac{(\sum x)(\sum y)}{n} = 993 - \frac{210 \times 24.5}{6} = 993 - \frac{5145}{6} = 993 - 857.5 = 135.5$.
4. Slope and intercept: $b = \frac{S_{xy}}{S_{xx}} = \frac{135.5}{1750} = 0.07742857 \dots \approx 0.0774$ (4 s.f. or 3 s.f. as required). $a = \bar{y} - b\bar{x} = 4.08333 \dots - 0.07742857 \dots \times 35 = 4.08333 \dots - 2.71 = 1.37333 \dots \approx 1.37$ (3 s.f.).
5. Regression line: $y = 1.37 + 0.0774x$ (or $y = 1.373 + 0.0774x$ keeping more precision).
6. Estimate for $x = 35$: Since $\bar{x} = 35$, the estimate is exactly $\bar{y} = 4.08$ tonnes/ha. Using the equation: $y = 1.37 + 0.0774(35) = 1.37 + 2.709 = 4.079 \approx 4.08$ tonnes/ha.

10.5 Least Squares Regression Line of x on y

DEFINITION Least Squares Regression Line of x on y

When we want to predict x from y , we minimise the sum of squared **horizontal** distances. The line is written as $x = c + dy$ (or $x = a' + b'y$). The formulas are symmetric to the y on x case.

PROPERTY Equation of the least squares line of x on y

The coefficients are given by:

$$d = \frac{S_{xy}}{S_{yy}}, \quad c = \bar{x} - d\bar{y}.$$

The equation is $x = c + dy$ or $x - \bar{x} = d(y - \bar{y})$. Relationship between the two lines:

- Both lines pass through the mean point $(\bar{x}, \bar{y})$.
- The product of the slopes is $b \times d = \frac{S_{xy}}{S_{xx}} \cdot \frac{S_{xy}}{S_{yy}} = \frac{S_{xy}^2}{S_{xx}S_{yy}} = r^2$.
- The slopes are reciprocals ($d = 1/b$) only if $r = \pm 1$ (perfect correlation).
- The angle between the two lines decreases as $|r|$ increases; they coincide when $r = \pm 1$.

METHOD Finding the least squares regression line of x on y

1. Compute $S_{yy} = \sum y^2 - \frac{(\sum y)^2}{n}$ and S_{xy} (same as before).
2. Compute the slope: $d = \frac{S_{xy}}{S_{yy}}$.
3. Compute the intercept: $c = \bar{x} - d\bar{y}$.
4. Write the equation: $x = c + dy$.
5. Check: Verify that the line passes through $(\bar{x}, \bar{y})$.

Reflex: Do not simply rearrange the y on x line; that gives a different line (unless $r = \pm 1$). Always compute d using S_{yy} .

WORKED EXAMPLE 18 Least squares line of x on y for the same agricultural data

Using the data from the previous example (fertilizer x , yield y), find the least squares regression line of x on y . Hence estimate the amount of fertilizer needed to achieve a yield of 4.5 tonnes/ha.

SOLUTION

Solution.

We already have: $n = 6$, $\sum x = 210$, $\sum y = 24.5$, $\sum x^2 = 9100$, $\sum y^2 = 110.55$, $\sum xy = 993$, $\bar{x} = 35$, $\bar{y} = 4.08333 \dots$, $S_{xy} = 135.5$.

1. Compute S_{yy} : $S_{yy} = \sum y^2 - \frac{(\sum y)^2}{n} = 110.55 - \frac{600.25}{6} = 110.55 - 100.04166 \dots = 10.50833 \dots$
2. Slope d : $d = \frac{S_{xy}}{S_{yy}} = \frac{135.5}{10.50833 \dots} = 12.894 \dots \approx 12.9$ (3 s.f.).
3. Intercept c : $c = \bar{x} - d\bar{y} = 35 - 12.894 \dots \times 4.08333 \dots = 35 - 52.65 \dots = -17.65 \dots \approx -17.7$ (3 s.f.).
4. Regression line of x on y : $x = -17.7 + 12.9y$ (or $x = 12.9y - 17.7$).
5. Estimate for $y = 4.5$: $x = -17.65 + 12.894(4.5) = -17.65 + 58.023 = 40.373 \approx 40.4$ kg/ha.
6. Check: The two lines intersect at $(\bar{x}, \bar{y}) = (35, 4.0833)$. For $y = 4.0833$, $x = -17.65 + 12.894(4.0833) \approx 35$. Correct.

10.6 Residual Sum of Squares (RSS) and the Line of Best Fit**DEFINITION** Residuals and Residual Sum of Squares

For a fitted regression line:

- For the line $y = a + bx$ (y on x), the **residual** for point i is the vertical distance: $e_i = y_i - \hat{y}_i = y_i - (a + bx_i)$.
- The **Residual Sum of Squares** for y on x is $\text{RSS}_{y|x} = \sum e_i^2 = \sum (y_i - a - bx_i)^2$.
- For the line $x = c + dy$ (x on y), the residual is the horizontal distance: $f_i = x_i - \hat{x}_i = x_i - (c + dy_i)$.
- The RSS for x on y is $\text{RSS}_{x|y} = \sum f_i^2 = \sum (x_i - c - dy_i)^2$.

PROPERTY Formulas for RSS

The RSS can be computed without calculating individual residuals:

$$\text{RSS}_{y|x} = S_{yy} - bS_{xy} = S_{yy} - \frac{S_{xy}^2}{S_{xx}} = S_{yy}(1 - r^2),$$

$$\text{RSS}_{x|y} = S_{xx} - dS_{xy} = S_{xx} - \frac{S_{xy}^2}{S_{yy}} = S_{xx}(1 - r^2).$$

The **coefficient of determination** is $R^2 = r^2 = 1 - \frac{\text{RSS}_{y|x}}{S_{yy}}$. It represents the proportion of the variation in y explained by the linear regression on x .

METHOD Calculating RSS and R^2

1. Compute S_{xx} , S_{yy} , S_{xy} and r (or b and d).
2. For y on x : $\text{RSS}_{y|x} = S_{yy} - \frac{S_{xy}^2}{S_{xx}}$ or $S_{yy}(1 - r^2)$.
3. For x on y : $\text{RSS}_{x|y} = S_{xx} - \frac{S_{xy}^2}{S_{yy}}$ or $S_{xx}(1 - r^2)$.
4. Compute $R^2 = r^2 = 1 - \frac{\text{RSS}_{y|x}}{S_{yy}}$.
5. Interpret R^2 as a percentage: " $R^2 \times 100\%$ of the variation in y is explained by the linear relationship with x ."

Reflex: RSS is always non-negative. A smaller RSS indicates a better fit. R^2 close to 1 means a good linear fit.

WORKED EXAMPLE 19 RSS for both lines and interpretation

For the agricultural data (fertilizer x , yield y) used in the previous two examples, calculate:

1. The Residual Sum of Squares for the regression line of y on x .
2. The Residual Sum of Squares for the regression line of x on y .
3. The coefficient of determination R^2 and interpret it.

SOLUTION

Solution.

We have: $S_{xx} = 1750$, $S_{yy} = 10.50833 \dots$, $S_{xy} = 135.5$. $r = \frac{S_{xy}}{\sqrt{S_{xx}S_{yy}}} = \frac{135.5}{\sqrt{1750 \times 10.50833 \dots}} = \frac{135.5}{\sqrt{18389.58 \dots}} = \frac{135.5}{135.608 \dots} \approx 0.9992$. $r^2 \approx 0.9984$.

1. RSS for y on x : $RSS_{y|x} = S_{yy} - \frac{S_{xy}^2}{S_{xx}} = 10.50833 \dots - \frac{135.5^2}{1750} = 10.50833 \dots - \frac{18360.25}{1750} = 10.50833 \dots - 10.4915714 \dots = 0.01676 \dots \approx 0.0168$.
2. RSS for x on y : $RSS_{x|y} = S_{xx} - \frac{S_{xy}^2}{S_{yy}} = 1750 - \frac{18360.25}{10.50833 \dots} = 1750 - 1747.18 \dots = 2.82 \dots \approx 2.82$.
3. Coefficient of determination: $R^2 = r^2 = 0.9984$ (or $1 - \frac{RSS_{y|x}}{S_{yy}} = 1 - \frac{0.01676}{10.50833} = 0.9984$).
Interpretation: 99.84% of the variation in crop yield (y) is explained by the linear relationship with fertilizer amount (x). The fit is extremely good; the linear model is highly appropriate for this data.

10.7 Rank Correlation: Spearman and Kendall

DEFINITION Rank Correlation

When data is ordinal (ranks) or when the relationship is monotonic but not necessarily linear, or when outliers make Pearson's r unreliable, we use **rank correlation coefficients**. The two main ones are **Spearman's rank correlation coefficient** (r_s or ρ) and **Kendall's rank correlation coefficient** (τ).

METHOD Spearman's rank correlation coefficient (r_s)

1. Rank the data: Assign ranks to x values (1 for smallest, 2 for next, etc.) and to y values. If there are ties (equal values), assign the **average rank** to each tied value. Example: values 10, 12, 12, 15 get ranks 1, 2.5, 2.5, 4.
2. Compute the difference in ranks: $d_i = \text{rank}(x_i) - \text{rank}(y_i)$.
3. Calculate r_s :
 - If there are no ties (or very few ties), use the shortcut formula: $r_s = 1 - \frac{6 \sum d_i^2}{n(n^2-1)}$.
 - If there are many ties, the shortcut formula is biased. Use the Pearson formula on the ranks: $r_s = \frac{\sum (R_{x_i} - \bar{R}_x)(R_{y_i} - \bar{R}_y)}{\sqrt{\sum (R_{x_i} - \bar{R}_x)^2 \sum (R_{y_i} - \bar{R}_y)^2}}$, where $\bar{R}_x = \bar{R}_y = \frac{n+1}{2}$.
4. Interpret: Same as Pearson's r : +1 perfect positive monotonic association, -1 perfect negative monotonic association, 0 no monotonic association.
5. Hypothesis test: For $n \geq 10$, r_s is approximately normally distributed; for small n , use critical value tables provided in the exam.

Key formula (no ties): $r_s = 1 - \frac{6 \sum d_i^2}{n(n^2-1)}$. Reflex: Always check for ties. If ties exist, mention

them and decide which formula to use. In GCE exams, if ties are present, the Pearson-on-ranks formula is safer.

METHOD Kendall's rank correlation coefficient (τ)

1. Rank both variables (same as Spearman, handling ties by average ranks).
2. Arrange the pairs in order of one variable (say x ranks ascending). Then look at the sequence of y ranks.
3. Count concordant and discordant pairs: For each pair of observations (i, j) with $i < j$ (in the ordered x list):
 - The pair is **concordant** if $(R_{y_i} - R_{y_j})$ has the same sign as $(R_{x_i} - R_{x_j})$ (which is positive since x ranks are in ascending order). So concordant means $R_{y_i} < R_{y_j}$.
 - The pair is **discordant** if $R_{y_i} > R_{y_j}$ (an inversion).
 - If $R_{y_i} = R_{y_j}$ (tie in y), the pair is neither concordant nor discordant (handled in τ_b or τ_c).
4. Compute τ (simple version, no ties or few ties): $\tau = \frac{N_c - N_d}{N_0}$, where N_c = number of concordant pairs, N_d = number of discordant pairs, $N_0 = \frac{1}{2}n(n-1)$ = total pairs.
5. With ties (Kendall's τ_b): $\tau_b = \frac{N_c - N_d}{\sqrt{(N_0 - N_1)(N_0 - N_2)}}$, where $N_1 = \sum \frac{t_k(t_k-1)}{2}$ (ties in x), $N_2 = \sum \frac{u_k(u_k-1)}{2}$ (ties in y), t_k, u_k are sizes of tie groups. For GCE, the simple version $\tau = \frac{N_c - N_d}{N_0}$ is often used when ties are few or absent.

KEY POINTS Uses and Limitations of Kendall's τ

- Uses: More robust to outliers than Spearman; better for small samples; directly interpretable as the difference between the probability of concordance and discordance; τ estimates a population parameter (Kendall's τ) naturally.
- Limitations: Computationally heavier for large n (though $O(n \log n)$ algorithms exist); less intuitive than Spearman's r_s ; not as widely tabulated for hypothesis testing in exam tables; the simple formula ignores ties, so τ_b is needed for tied data.

WORKED EXAMPLE 20 Spearman and Kendall for examination ranks

Two examiners rank 8 candidates in an oral examination:

Candidate	A	B	C	D	E	F	G	H
Examiner 1 (x)	1	2	3	4	5	6	7	8
Examiner 2 (y)	2	1	4	3	6	5	8	7

1. Calculate Spearman's rank correlation coefficient r_s .
2. Calculate Kendall's τ .
3. Comment on the agreement between the examiners.

SOLUTION

Solution.

1. Spearman's r_s : The data are already ranks (1 to 8). No ties. Compute $d_i = R_{x_i} - R_{y_i}$:

Candidate	R_x	R_y	d	d^2
A	1	2	-1	1
B	2	1	1	1
C	3	4	-1	1
D	4	3	1	1
E	5	6	-1	1
F	6	5	1	1
G	7	8	-1	1
H	8	7	1	1
Total				8

$$n = 8, \sum d^2 = 8. r_s = 1 - \frac{6 \times 8}{8(8^2 - 1)} = 1 - \frac{48}{8 \times 63} = 1 - \frac{48}{504} = 1 - \frac{2}{21} = \frac{19}{21} \approx 0.9048.$$

2. Kendall's τ : Arrange in order of Examiner 1 (already ordered). Look at Examiner 2's ranks: 2, 1, 4, 3, 6, 5, 8, 7. Count concordant and discordant pairs among the y ranks. Total pairs $N_0 = \frac{1}{2} \times 8 \times 7 = 28$. We count inversions (discordant pairs) in the sequence 2, 1, 4, 3, 6, 5, 8, 7: - (2,1): inversion $\rightarrow$ discordant - (2,4): concordant - (2,3): concordant - (2,6): concordant - (2,5): concordant - (2,8): concordant - (2,7): concordant - (1,4): concordant - (1,3): concordant - (1,6): concordant - (1,5): concordant - (1,8): concordant - (1,7): concordant - (4,3): inversion $\rightarrow$ discordant - (4,6): concordant - (4,5): concordant - (4,8): concordant - (4,7): concordant - (3,6): concordant - (3,5): concordant - (3,8): concordant - (3,7): concordant - (6,5): inversion $\rightarrow$ discordant - (6,8): concordant - (6,7): concordant - (5,8): concordant - (5,7): concordant - (8,7):

inversion $\rightarrow$ discordant Total discordant $N_d = 4$ (the inversions: (2,1), (4,3), (6,5), (8,7)).

Concordant $N_c = 28 - 4 = 24$. $\tau = \frac{N_c - N_d}{N_0} = \frac{24 - 4}{28} = \frac{20}{28} = \frac{5}{7} \approx 0.7143$.

3. Comment: Both coefficients indicate strong positive agreement between the examiners. Spearman's $r_s = 0.905$ is higher than Kendall's $\tau = 0.714$; this is typical because τ is more conservative (it counts pairwise agreements/disagreements directly). The examiners consistently rank candidates in a very similar order, with only minor swaps of adjacent ranks (each adjacent pair is swapped). At the 5% significance level, both coefficients would indicate statistically significant agreement (critical values for $n = 8$: $r_s \approx 0.738$, $\tau \approx 0.619$).

EXERCISE 12 Practice: Complete Regression and Correlation Analysis

The table below shows the number of hours of revision (x) and the examination mark (y) for 8 students.

Hours x	10	12	14	16	18	20	22	24
Mark y	45	50	55	60	65	70	75	80

1. Draw a scatter diagram and comment.
2. Calculate S_{xx} , S_{yy} , S_{xy} and the product moment correlation coefficient r .
3. Find the least squares regression line of y on x .
4. Find the least squares regression line of x on y .
5. Calculate $RSS_{y|x}$ and $RSS_{x|y}$ and the coefficient of determination R^2 .
6. Use the appropriate line to estimate the mark for a student who revises for 15 hours, and the hours needed to achieve a mark of 72.
7. Calculate Spearman's r_s and Kendall's τ for this data (treating the raw data as ranks after verifying they are already in rank order). Compare with r .

SOLUTION Exercise: Complete Regression and Correlation Analysis

1. Scatter diagram: Points lie exactly on a straight line. Perfect positive linear correlation.
2. Sums: $n = 8$, $\sum x = 136$, $\sum y = 500$, $\sum x^2 = 2520$, $\sum y^2 = 32400$, $\sum xy = 9160$. $\bar{x} = 17$, $\bar{y} = 62.5$. $S_{xx} = 2520 - 136^2/8 = 2520 - 2312 = 208$. $S_{yy} = 32400 - 500^2/8 = 32400 -$

$$31250 = 1150. S_{xy} = 9160 - (136 \times 500)/8 = 9160 - 8500 = 660. r = 660/\sqrt{208 \times 1150} = 660/\sqrt{239200} = 660/489.08 = 1 \text{ (exactly, since } y = 2.5x + 20 \text{)}.$$

3. y on x : $b = 660/208 = 3.17308 \dots$? Wait: $y = 2.5x + 20$ exactly. Let's recalc: $S_{xy} = 660$, $S_{xx} = 208$, $b = 660/208 = 3.173$? But $y = 2.5x + 20$ gives $b = 2.5$. Check sums: x : 10,12,14,16,18,20,22,24. y : 45,50,55,60,65,70,75,80. $\sum x = 136$, $\sum y = 500$. $\sum xy = 10 \times 45 + 12 \times 50 + 14 \times 55 + 16 \times 60 + 18 \times 65 + 20 \times 70 + 22 \times 75 + 24 \times 80 = 450 + 600 + 770 + 960 + 1170 + 1400 + 1650 + 1920 = 8920$. Not 9160. My previous sum was wrong. Let's recompute properly. $\sum xy = 450 + 600 + 770 + 960 + 1170 + 1400 + 1650 + 1920 = 8920$. $S_{xy} = 8920 - (136 \times 500)/8 = 8920 - 8500 = 420$. $S_{xx} = 2520 - 2312 = 208$. $b = 420/208 = 2.019$? No, $y = 2.5x + 20$ exactly. $\sum x^2 = 100 + 144 + 196 + 256 + 324 + 400 + 484 + 576 = 2480$. Not 2520. Let's do it cleanly: x : 10,12,14,16,18,20,22,24. $\sum x = 136$. $\sum x^2 = 100 + 144 + 196 + 256 + 324 + 400 + 484 + 576 = 2480$. y : 45,50,55,60,65,70,75,80. $\sum y = 500$. $\sum y^2 = 2025 + 2500 + 3025 + 3600 + 4225 + 4900 + 5625 + 6400 = 32300$. $\sum xy = 450 + 600 + 770 + 960 + 1170 + 1400 + 1650 + 1920 = 8920$. $S_{xx} = 2480 - 136^2/8 = 2480 - 2312 = 168$. $S_{yy} = 32300 - 500^2/8 = 32300 - 31250 = 1050$. $S_{xy} = 8920 - 136 \times 500/8 = 8920 - 8500 = 420$. $b = 420/168 = 2.5$. $a = 62.5 - 2.5 \times 17 = 62.5 - 42.5 = 20$. $y = 20 + 2.5x$. Correct. $r = 420/\sqrt{168 \times 1050} = 420/\sqrt{176400} = 420/420 = 1$.
4. x on y : $d = 420/1050 = 0.4$. $c = 17 - 0.4 \times 62.5 = 17 - 25 = -8$. $x = -8 + 0.4y$. (Equivalent to $y = 20 + 2.5x$).
5. RSS: $r = 1 \Rightarrow \text{RSS}_{y|x} = 0, \text{RSS}_{x|y} = 0, R^2 = 1$.
6. Estimates: For $x = 15$, $y = 20 + 2.5(15) = 57.5$. For $y = 72$, $x = -8 + 0.4(72) = 20.8$ hours.
7. Rank correlation: Data is perfectly linear, so ranks are perfectly correlated. $r_s = 1, \tau = 1$. Same as r .

11 EXERCISES

11.1 I check my knowledge

EXERCISE 13 MCQ: Regression Line Properties

Which of the following statements is always true for the least squares regression line of y on x ?

1. It passes through the origin $(0, 0)$.
2. It passes through the mean point $(\bar{x}, \bar{y})$.
3. Its slope is equal to the product moment correlation coefficient r .
4. The sum of the residuals is equal to the sum of the y -values.

EXERCISE 14 True/False: Correlation vs Causation

State whether the following statements are True or False. Justify your answer briefly.

1. If the product moment correlation coefficient $r = 0$, then there is no relationship whatsoever between the variables x and y .
2. The regression line of x on y can be obtained by simply rearranging the equation of the regression line of y on x to make x the subject.
3. Spearman's rank correlation coefficient R_s can only be calculated if the data are quantitative (numerical).
4. For a given data set, the residual sum of squares (RSS) for the least squares line of y on x is always less than or equal to the RSS for any other line.

EXERCISE 15 Definitions and Direct Formulas

1. Define the residual for a data point (x_i, y_i) with respect to the regression line $y = a + bx$.
2. Write down the formula for the slope b of the least squares regression line of y on x in terms of S_{xy} and S_{xx} .
3. State the relationship between the product moment correlation coefficient r , the slope b of the regression line of y on x , and the slope b' of the regression line of x on y .

4. Given the ranks of 5 items by two judges, explain how to handle tied ranks when calculating Spearman's rank correlation coefficient.

EXERCISE 16 Direct Calculation: Covariance and Correlation

For a bivariate data set of $n = 6$ pairs, the following summary statistics are given:

$$\sum x = 42, \quad \sum y = 60, \quad \sum x^2 = 320, \quad \sum y^2 = 650, \quad \sum xy = 440.$$

Calculate:

1. The sample covariance S_{xy} .
2. The sample variances S_{xx} and S_{yy} .
3. The product moment correlation coefficient r .
4. Comment on the strength and direction of the linear relationship.

11.2 I practise

EXERCISE 17 Least Squares Regression Line: Rainfall and Crop Yield

The table below shows the annual rainfall (in cm) and the corresponding maize yield (in tonnes per hectare) for a farm in Buea over 8 years.

Rainfall x (cm)	120	145	130	160	155	140	170	150
Yield y (t/ha)	2.1	3.0	2.5	3.8	3.5	2.8	4.0	3.2

1. Plot a scatter diagram of yield against rainfall. (Graph paper not required; describe the trend).
2. Calculate the equation of the least squares regression line of y on x in the form $y = a + bx$.
3. Use your equation to estimate the maize yield when the rainfall is 158 cm.
4. Calculate the residual for the year when rainfall was 145 cm and yield was 3.0 t/ha. Interpret its sign.

EXERCISE 18 Regression Line of x on y and Comparison

Using the same data from the previous exercise (Rainfall x , Yield y):

1. Calculate the equation of the least squares regression line of x on y in the form $x = c + dy$.
2. Using the line of x on y , estimate the rainfall required to achieve a yield of 3.6 t/ha.
3. Rearrange the regression line of y on x (from the previous exercise) to make x the subject. Compare this equation with the line of x on y obtained in (i). Explain why they are different.

EXERCISE 19 Mayer's Method vs Least Squares

A small business in Douala records the advertising expenditure (in 1000 FCFA) and the resulting sales (in 1000 FCFA) for 9 months.

Advertising x	2	4	5	7	8	10	11	12	14
Sales y	15	22	25	30	33	38	40	44	48

1. Divide the data into two groups of equal size (or as equal as possible) to apply Mayer's method. Find the mean points $(\bar{x}_1, \bar{y}_1)$ and $(\bar{x}_2, \bar{y}_2)$.
2. Determine the equation of the regression line by Mayer's method (line joining the two mean points).
3. Calculate the equation of the least squares regression line of y on x .
4. Compute the Residual Sum of Squares (RSS) for both lines. Which line provides a better fit? Justify.

EXERCISE 20 Rank Correlation: Singing Competition

Two judges rank 7 singers in a "Yaoundé Talent Show" as follows:

Singer	A	B	C	D	E	F	G
Judge 1 Rank	1	2	3	4	5	6	7
Judge 2 Rank	2	1	4	3	6	7	5

1. Calculate Spearman's rank correlation coefficient R_s .
2. Interpret the value of R_s in the context of the competition.

3. If Singer H is added and both judges rank H as 8th, explain how R_s would change without recalculating fully.

11.3 I prepare for the GCE

EXERCISE 21 GCE Style: Mixed Regression and Correlation Problem – 18 marks

The ages (x years) and prices (y thousand FCFA) of 10 second-hand taxis in Douala are summarised below:

$$n = 10, \quad \sum x = 65, \quad \sum y = 820, \quad \sum x^2 = 485, \quad \sum y^2 = 72500, \quad \sum xy = 5020.$$

1. Calculate the product moment correlation coefficient r between age and price. [3 marks]
2. Interpret your value of r in the context of the question. [2 marks]
3. Find the equation of the least squares regression line of y on x in the form $y = a + bx$. [4 marks]
4. A buyer wants to purchase a 6-year-old taxi. Use your regression line to estimate a fair price. [2 marks]
5. The oldest taxi in the sample is 9 years old. Explain why it would be unreliable to use your regression line to estimate the price of a 12-year-old taxi. [2 marks]
6. Calculate the Residual Sum of Squares (RSS) for the regression line of y on x . [3 marks]
7. Given that the regression line of x on y is $x = 7.2 - 0.0045y$, verify that $r^2 = b \times b'$, where b and b' are the slopes of the two regression lines. [2 marks]

EXERCISE 22 GCE Style: Kendall's Rank Correlation and Interpretation – 14 marks

Five students are ranked by their Mathematics teacher and their Physics teacher according to their performance in the mock exams.

Student	P	Q	R	S	T
Maths Rank	1	2	3	4	5
Physics Rank	2	1	5	3	4

1. List all $\binom{5}{2} = 10$ pairs of students. For each pair, determine whether the rankings are concordant or discordant. [5 marks]

2. Calculate Kendall's rank correlation coefficient τ . [3 marks]
3. Calculate Spearman's rank correlation coefficient R_s for the same data. [3 marks]
4. Compare the values of τ and R_s . State one use and one limitation of Kendall's τ compared to Spearman's R_s . [3 marks]

EXERCISE 23 GCE Style: Regression Lines and Mean Point – 12 marks

For a certain bivariate data set, the least squares regression line of y on x is $y = 0.8x + 3$ and the regression line of x on y is $x = 0.45y + 1$.

1. Find the coordinates of the mean point $(\bar{x}, \bar{y})$. [3 marks]
2. Calculate the product moment correlation coefficient r . [3 marks]
3. Given that $S_{xx} = 50$, find the value of S_{yy} . [3 marks]
4. If a new data point $(\bar{x}, \bar{y} + 2)$ is added to the data set, describe qualitatively what would happen to the slope of the regression line of y on x and to the value of r . [3 marks]

12 SOLUTIONS TO THE EXERCISES

SOLUTION Exercise 1 — MCQ: Regression Line Properties

The correct answer is (b): the least squares regression line of y on x always passes through the mean point $(\bar{x}, \bar{y})$.

Justification. The least squares line is $y = a + bx$ with $b = \frac{S_{xy}}{S_{xx}}$ and $a = \bar{y} - b\bar{x}$. Substituting $x = \bar{x}$:

$$a + b\bar{x} = (\bar{y} - b\bar{x}) + b\bar{x} = \bar{y},$$

so the point $(\bar{x}, \bar{y})$ lies on the line, for any data set.

Why the others are false:

- (a) The line passes through $(0, 0)$ only if $a = \bar{y} - b\bar{x} = 0$, which is not true in general.
- (c) The slope is $b = \frac{S_{xy}}{S_{xx}} = r \frac{s_y}{s_x}$, which equals r only when $s_x = s_y$.
- (d) The sum of the residuals is always 0 (a property of least squares), not the sum of the y -values.

SOLUTION Exercise 2 — True/False: Correlation vs Causation

1. False. $r = 0$ only means there is no linear relationship. The variables may be strongly related in a non-linear way (e.g. $y = x^2$ on symmetric data gives $r = 0$).
2. False. The line of x on y minimises the sum of squares of horizontal residuals, whereas the line of y on x minimises vertical residuals. They are two different lines (they coincide only when $r = \pm 1$). Rearranging one equation does not give the other.
3. False. Spearman's coefficient only requires ranks. Qualitative data that can be ordered (e.g. beauty, taste, quality of a performance) can be ranked and R_s computed, even with no numerical measurements at all.
4. True. By definition, the least squares line is the line that minimises the residual sum of squares. Hence its RSS is less than or equal to the RSS of any other line fitted to the same data.

SOLUTION Exercise 3 — Definitions and Direct Formulas

1. The residual of the point (x_i, y_i) is the vertical signed distance between the observed

value and the value predicted by the line:

$$e_i = y_i - \hat{y}_i = y_i - (a + bx_i).$$

2. $b = \frac{S_{xy}}{S_{xx}}$, where $S_{xy} = \sum x_i y_i - \frac{\sum x_i \sum y_i}{n}$ and $S_{xx} = \sum x_i^2 - \frac{(\sum x_i)^2}{n}$.
3. The slopes satisfy $b \times b' = r^2$, i.e. $r = \pm\sqrt{bb'}$, the sign of r being the common sign of b and b' . (Indeed $bb' = \frac{S_{xy}}{S_{xx}} \cdot \frac{S_{xy}}{S_{yy}} = \frac{S_{xy}^2}{S_{xx}S_{yy}} = r^2$.)
4. When two or more items are tied, assign to each of them the mean of the ranks they occupy. For example, if two items tie for positions 3 and 4, each receives rank $\frac{3+4}{2} = 3.5$, and the next item receives rank 5. One then computes d and $\sum d^2$ as usual in $R_s = 1 - \frac{6 \sum d^2}{n(n^2 - 1)}$ (a correction factor may be applied if ties are numerous).

SOLUTION Exercise 4 — Direct Calculation: Covariance and Correlation

Given $n = 6$, $\sum x = 42$, $\sum y = 60$, $\sum x^2 = 320$, $\sum y^2 = 650$, $\sum xy = 440$.

1. Covariance:

$$S_{xy} = \sum xy - \frac{\sum x \sum y}{n} = 440 - \frac{42 \times 60}{6} = 440 - 420 = \boxed{20}.$$

2. Variances:

$$S_{xx} = 320 - \frac{42^2}{6} = 320 - 294 = \boxed{26}, \quad S_{yy} = 650 - \frac{60^2}{6} = 650 - 600 = \boxed{50}.$$

3. Product moment correlation coefficient:

$$r = \frac{S_{xy}}{\sqrt{S_{xx}S_{yy}}} = \frac{20}{\sqrt{26 \times 50}} = \frac{20}{\sqrt{1300}} = \frac{20}{36.0555...} \approx \boxed{0.555}.$$

4. Comment: $r \approx 0.56 > 0$, so there is a moderate positive linear relationship: as x increases, y tends to increase, but the points are fairly scattered around any straight line.

SOLUTION Exercise 5 — Least Squares Regression Line: Rainfall and Crop Yield

Preliminary sums ($n = 8$):

$$\sum x = 1170, \quad \sum y = 24.9, \quad \sum x^2 = 172950, \quad \sum xy = 3714.5, \quad \sum y^2 = 80.43.$$

$$\bar{x} = \frac{1170}{8} = 146.25, \quad \bar{y} = \frac{24.9}{8} = 3.1125.$$

1. Scatter diagram: plotting yield y against rainfall x , the points rise from the lower left to the upper right in a roughly linear band: the trend is positive and approximately linear (more rainfall $\Rightarrow$ higher yield).
2. Regression line of y on x :

$$S_{xx} = 172950 - \frac{1170^2}{8} = 172950 - 171112.5 = 1837.5,$$

$$S_{xy} = 3714.5 - \frac{1170 \times 24.9}{8} = 3714.5 - 3641.625 = 72.875.$$

$$b = \frac{72.875}{1837.5} = 0.03966 \dots \approx 0.0397, \quad a = 3.1125 - 0.03966 \dots \times 146.25 \approx -2.6875.$$

$$y = -2.688 + 0.0397x$$

3. Estimate for $x = 158$:

$$y = -2.6875 + 0.03966 \dots \times 158 \approx 3.58 \text{ t/ha.}$$

4. Residual at (145, 3.0): predicted value $\hat{y} = -2.6875 + 0.03966 \dots \times 145 \approx 3.063$.

$$e = y - \hat{y} = 3.0 - 3.063 = -0.063 \text{ t/ha.}$$

The residual is negative: the actual yield that year was about 0.06 t/ha below what the model predicts — the line slightly overestimates the yield for that year.

SOLUTION Exercise 6 — Regression Line of x on y and Comparison

Using the sums of Exercise 5: $S_{xy} = 72.875$, and

$$S_{yy} = 80.43 - \frac{24.9^2}{8} = 80.43 - 77.50125 = 2.92875.$$

1. Line of x on y : $x = c + dy$ with

$$d = \frac{S_{xy}}{S_{yy}} = \frac{72.875}{2.92875} \approx 24.88, \quad c = \bar{x} - d\bar{y} = 146.25 - 24.88 \times 3.1125 \approx 68.81.$$

$$x = 68.81 + 24.88y$$

2. Rainfall for a yield of 3.6 t/ha:

$$x = 68.81 + 24.88 \times 3.6 \approx 158.4 \text{ cm.}$$

3. Rearranging the line of y on x : from $y = -2.688 + 0.0397x$,

$$x = \frac{y + 2.688}{0.0397} \approx 67.7 + 25.2y.$$

This is close to but different from $x = 68.81 + 24.88y$. The two lines are different because they answer different minimisation problems: the line of y on x minimises the sum of squared vertical deviations, while the line of x on y minimises the sum of squared horizontal deviations. Both pass through $(\bar{x}, \bar{y}) = (146.25, 3.1125)$ but have different slopes; they would coincide only if $r = \pm 1$ (here $r = \sqrt{0.0397 \times 24.88} \approx 0.994$, which is why they are nearly identical).

SOLUTION Exercise 7 — Mayer's Method vs Least Squares

Preliminary sums ($n = 9$): $\sum x = 73$, $\sum y = 295$, $\sum x^2 = 719$, $\sum xy = 2737$, $\sum y^2 = 10607$.

1. Mayer's groups. Order the data by x and split into a group of 4 and a group of 5:

$$G_1 = \{(2, 15), (4, 22), (5, 25), (7, 30)\}, \quad G_2 = \{(8, 33), (10, 38), (11, 40), (12, 44), (14, 48)\}.$$

$$\bar{x}_1 = \frac{18}{4} = 4.5, \quad \bar{y}_1 = \frac{92}{4} = 23; \quad \bar{x}_2 = \frac{55}{5} = 11, \quad \bar{y}_2 = \frac{203}{5} = 40.6.$$

2. Mayer's line through $(4.5, 23)$ and $(11, 40.6)$:

$$b_M = \frac{40.6 - 23}{11 - 4.5} = \frac{17.6}{6.5} \approx 2.708, \quad a_M = 23 - 2.708 \times 4.5 \approx 10.82.$$

$$y = 10.82 + 2.708x$$

3. Least squares line:

$$S_{xx} = 719 - \frac{73^2}{9} = 126.89, \quad S_{xy} = 2737 - \frac{73 \times 295}{9} = 344.22.$$

$$b = \frac{344.22}{126.89} \approx 2.713, \quad a = \frac{295}{9} - 2.713 \times \frac{73}{9} \approx 32.78 - 22.00 = 10.77.$$

$$y = 10.77 + 2.713x$$

4. Residual sums of squares. For the least squares line, use $RSS = S_{yy} - \frac{S_{xy}^2}{S_{xx}}$:

$$S_{yy} = 10607 - \frac{295^2}{9} = 937.56, \quad RSS_{LS} = 937.56 - \frac{344.22^2}{126.89} \approx 937.56 - 933.80 \approx 3.76.$$

For Mayer's line, computing each residual $e_i = y_i - (10.82 + 2.708x_i)$ and summing the squares:

(x_i, y_i)	$\hat{y}_i$ (Mayer)	e_i	e_i^2
(2,15)	16.24	-1.24	1.54
(4,22)	21.65	0.35	0.12
(5,25)	24.36	0.64	0.41
(7,30)	29.78	0.22	0.05
(8,33)	32.48	0.52	0.27
(10,38)	37.90	0.10	0.01
(11,40)	40.61	-0.61	0.37
(12,44)	43.32	0.68	0.46
(14,48)	48.73	-0.73	0.53

$$\text{RSS}_M \approx 1.54 + 0.12 + 0.41 + 0.05 + 0.27 + 0.01 + 0.37 + 0.46 + 0.53 \approx 3.76.$$

The two fits are almost identical here, but $\text{RSS}_{LS} \leq \text{RSS}_M$: the least squares line provides the better fit, as guaranteed by the minimisation property (Mayer's method only uses the two group means, whereas least squares uses every data point).

SOLUTION Exercise 8 — Rank Correlation: Singing Competition

1. Differences $d = R_1 - R_2$:

Singer	A	B	C	D	E	F	G
d	-1	1	-1	1	-1	-1	2
d^2	1	1	1	1	1	1	4

$$\sum d^2 = 10, n = 7:$$

$$R_s = 1 - \frac{6 \times 10}{7(7^2 - 1)} = 1 - \frac{60}{336} = 1 - 0.1786 \approx \boxed{0.821}.$$

2. Interpretation: $R_s \approx 0.82$ shows a strong positive agreement between the two judges: singers ranked highly by Judge 1 tend to be ranked highly by Judge 2. The judges assess the performances in a very similar way.
3. Adding Singer H ranked 8th by both judges: the new difference is $d_H = 0$, so $\sum d^2$ stays equal to 10, but n increases from 7 to 8:

$$R_s = 1 - \frac{6 \times 10}{8(64 - 1)} = 1 - \frac{60}{504} \approx 0.881.$$

Since the denominator $n(n^2 - 1)$ grows while $\sum d^2$ is unchanged, the subtracted term shrinks and R_s increases towards 1: a new pair in perfect agreement strengthens the measured agreement.

SOLUTION Exercise 9 — GCE Style: Mixed Regression and Correlation Problem

Mark scheme and full solution. First compute the summary quantities:

$$\bar{x} = 6.5, \quad \bar{y} = 82, \quad S_{xx} = 485 - \frac{65^2}{10} = 62.5,$$

$$S_{yy} = 72500 - \frac{820^2}{10} = 5260, \quad S_{xy} = 5020 - \frac{65 \times 820}{10} = -310.$$

1. [3 marks] $r = \frac{-310}{\sqrt{62.5 \times 5260}} = \frac{-310}{573.4} \approx \boxed{-0.541}$. (M1 correct formula; A1 correct S_{xy} , S_{xx} , S_{yy} ; A1 answer to 3 s.f.)

2. [2 marks] $r \approx -0.54$: a moderate negative linear correlation — as the age of a taxi increases, its price tends to decrease, with considerable scatter. (B1 direction + strength; B1 contextual interpretation.)

3. [4 marks]

$$b = \frac{S_{xy}}{S_{xx}} = \frac{-310}{62.5} = -4.96, \quad a = 82 - (-4.96)(6.5) = 114.24.$$

$$\boxed{y = 114.24 - 4.96x} \quad (\text{M1 for } b, \text{ A1; M1 for } a \text{ via } (\bar{x}, \bar{y}), \text{ A1.})$$

4. [2 marks] $x = 6$: $y = 114.24 - 4.96 \times 6 = 84.48$. Fair price $\approx \boxed{84\,500 \text{ FCFA}}$. (M1 substitution; A1 with units.)

5. [2 marks] 12 years lies outside the range of the data (maximum age 9 years): this is extrapolation. There is no evidence the linear trend continues (prices cannot fall below zero, for instance), so the estimate would be unreliable. (B1 extrapolation identified; B1 reason.)

6. [3 marks]

$$\text{RSS} = S_{yy} - \frac{S_{xy}^2}{S_{xx}} = 5260 - \frac{(-310)^2}{62.5} = 5260 - 1537.6 = \boxed{3722.4}.$$

(M1 correct formula; A1 substitution; A1 answer.)

7. [2 marks] With $b = -4.96$ and the given $b' = -0.0045$: $b \times b' = 0.0223$, whereas $r^2 = 0.292$. These are not equal, which reveals that the line given in the question is inconsistent with the data (a misprint). The correct line of x on y has slope $b' = \frac{S_{xy}}{S_{yy}} =$

$$\frac{-310}{5260} \approx -0.0589, \text{ and then}$$

$$b \times b' = (-4.96)(-0.0589) \approx 0.292 = r^2. \checkmark$$

Examiner expectation: M1 for computing both bb' and r^2 ; A1 for the comparison and conclusion. A candidate who spots and corrects the inconsistency with justification earns full credit.

SOLUTION Exercise 10 — GCE Style: Kendall's Rank Correlation and Interpretation

1. [5 marks] A pair is concordant if the two judges rank the two students in the same order, discordant otherwise. Maths order: $P < Q < R < S < T$.

Pair	PQ	PR	PS	PT	QR	QS	QT	RS	RT	ST
Type	D	C	C	C	C	C	C	D	D	C

(PQ: Maths $P < Q$ but Physics $P > Q$ — discordant; RS: Maths $R < S$ but Physics $R > S$ — discordant; RT: similarly discordant.) Hence $C = 7$ concordant, $D = 3$ discordant. (B1 per two correct pairs, rounded to 5 marks.)

2. [3 marks]

$$\tau = \frac{C - D}{\binom{5}{2}} = \frac{7 - 3}{10} = \boxed{0.4}.$$

(M1 formula; A1 correct counts; A1 answer.)

3. [3 marks] Differences d : P: -1, Q: 1, R: -2, S: 1, T: 1; $\sum d^2 = 1 + 1 + 4 + 1 + 1 = 8$.

$$R_s = 1 - \frac{6 \times 8}{5(25 - 1)} = 1 - \frac{48}{120} = \boxed{0.6}.$$

(M1 differences; A1 $\sum d^2$; A1 answer.)

4. [3 marks] Both coefficients are positive and indicate moderate agreement, but $R_s = 0.6 > \tau = 0.4$: the two coefficients measure agreement differently and need not be equal.

- Use of Kendall's τ : it has a direct probabilistic interpretation — τ measures (probability of concordance) - (probability of discordance) — and it behaves better than R_s for small samples or when there are many ties.
- Limitation: it is more tedious to compute for large n (all $\binom{n}{2}$ pairs must be examined), and its values are typically smaller in magnitude than R_s , so the two scales cannot be compared directly.

(B1 comparison; B1 valid use; B1 valid limitation.)

SOLUTION Exercise 11 — GCE Style: Regression Lines and Mean Point

1. [3 marks] Both regression lines pass through $(\bar{x}, \bar{y})$, so solve

$$\bar{y} = 0.8\bar{x} + 3, \quad \bar{x} = 0.45\bar{y} + 1.$$

Substituting: $\bar{y} = 0.8(0.45\bar{y} + 1) + 3 = 0.36\bar{y} + 3.8$, so $0.64\bar{y} = 3.8$, giving $\bar{y} = 5.9375$ and

$$\bar{x} = 0.45 \times 5.9375 + 1 = 3.671875.$$

$$(\bar{x}, \bar{y}) \approx (3.67, 5.94) \quad (\text{M1 using the mean-point property; A1 A1.})$$

2. [3 marks] $r^2 = b \times b' = 0.8 \times 0.45 = 0.36$. Since both slopes are positive,

$$r = +\sqrt{0.36} = \boxed{0.6}. \quad (\text{M1 } r^2 = bb'; \text{ A1 value; B1 correct sign justified.})$$

3. [3 marks] $b = \frac{S_{xy}}{S_{xx}} = 0.8$ and $S_{xx} = 50$ give $S_{xy} = 0.8 \times 50 = 40$. Then

$$b' = \frac{S_{xy}}{S_{yy}} = 0.45 \implies S_{yy} = \frac{40}{0.45} \approx \boxed{88.9}.$$


(M1 finding S_{xy} ; M1 using b' ; A1 answer.)

4. [3 marks] The new point $(\bar{x}, \bar{y} + 2)$ lies vertically above the mean point. Its x -deviation from the mean is exactly zero, so it adds nothing to S_{xy} and S_{xx} , but it adds a positive amount to S_{yy} . Consequently:

- the slope $b = S_{xy}/S_{xx}$ is unchanged;
- $r = \frac{S_{xy}}{\sqrt{S_{xx}S_{yy}}}$ decreases in magnitude (the denominator grows while the numerator stays fixed).

Intuitively, the new point does not follow the linear trend, so it weakens the measured linear correlation without tilting the line. (B1 slope unchanged with reason; B1 r decreases; B1 quality of explanation.)

13 SUMMARY SHEET



Mind map of the chapter: from scatter diagrams to rank correlation.

FICHE BILAN L'essentiel à connaître pour l'examen

Key definitions

- **Bivariate data**: pairs (x_i, y_i) observed on the same individuals.
- **Scatter diagram**: plot of the points (x_i, y_i) ; it suggests the type of association (positive, negative, none, linear or not).
- **Regression line of y on x**: line used to predict y from x; **regression line of x on y**: line used to predict x from y.
- **Residual**: $e_i = y_i - \hat{y}_i$, vertical gap between observation and fitted line.
- **Covariance**: measures how x and y vary together.
- **Correlation coefficient**: a number in $[-1, 1]$ measuring the strength and direction of a linear (or rank) association.

Formulas to know

- Means: $\bar{x} = \frac{\sum x_i}{n}$, $\bar{y} = \frac{\sum y_i}{n}$.

- Sample variance and covariance:

$$s_x^2 = \frac{1}{n} \sum (x_i - \bar{x})^2 = \frac{\sum x_i^2}{n} - \bar{x}^2, \quad s_{xy} = \frac{1}{n} \sum (x_i - \bar{x})(y_i - \bar{y}) = \frac{\sum x_i y_i}{n} - \bar{x}\bar{y}.$$

- Least squares line of y on x : $y - \bar{y} = b(x - \bar{x})$ with $b = \frac{s_{xy}}{s_x^2}$.
- Least squares line of x on y : $x - \bar{x} = b'(y - \bar{y})$ with $b' = \frac{s_{xy}}{s_y^2}$.
- Both lines pass through the **mean point** $(\bar{x}, \bar{y})$.
- Residual sums of squares:

$$\text{RSS}_{y \text{ on } x} = \sum (y_i - \hat{y}_i)^2 = \sum y_i^2 - n\bar{y}^2 - \frac{(\sum x_i y_i - n\bar{x}\bar{y})^2}{\sum x_i^2 - n\bar{x}^2},$$

and symmetrically for $\text{RSS}_{x \text{ on } y}$.

- **Line of best fit**: the line with the smaller RSS (equivalently the larger $|r|$ choice); the y -on- x line is best for predicting y .
- Product moment correlation coefficient:

$$r = \frac{s_{xy}}{s_x s_y} = \frac{\sum x_i y_i - n\bar{x}\bar{y}}{\sqrt{(\sum x_i^2 - n\bar{x}^2)(\sum y_i^2 - n\bar{y}^2)}}, \quad -1 \leq r \leq 1.$$

Also $bb' = r^2$.

- Spearman's rank correlation (no tied ranks):

$$r_s = 1 - \frac{6 \sum d_i^2}{n(n^2 - 1)}, \quad d_i = \text{rank}(x_i) - \text{rank}(y_i).$$

- Kendall's rank correlation:

$$\tau = \frac{2(P - Q)}{n(n - 1)} = \frac{S}{\frac{1}{2}n(n - 1)},$$

where P = number of concordant pairs, Q = number of discordant pairs.

Methods

- **Mayer's method**: order the data by x , split into two equal (or nearly equal) groups, compute the mean point of each group, and draw the line through these two points.
- **Least squares**: compute $\sum x$, $\sum y$, $\sum x^2$, $\sum y^2$, $\sum xy$ in a table; then b , then the line through $(\bar{x}, \bar{y})$.
- **Ranking**: rank from highest (or lowest) consistently; give tied values the mean of the ranks they occupy.

- **Kendall:** order one variable's ranks naturally, then count, for each rank of the second variable, how many later ranks are larger (P) and smaller (Q).

Common mistakes

- Confusing y on x with x on y : they are different lines (unless $|r| = 1$).
- Forgetting that both regression lines pass through $(\bar{x}, \bar{y})$ — a useful check.
- Using r to claim causation: correlation is not causation.
- Predicting outside the data range (extrapolation) without caution.
- Sign errors in $\sum d_i^2$; always check $\sum d_i = 0$.
- Mishandling tied ranks in Spearman or Kendall.
- Interpreting $r \approx 0$ as "no relation": there may be a strong non-linear relation.

GCE tips

- Show the table of sums explicitly; examiners award method marks for $\sum x^2$, $\sum xy$.
- Quote the formula before substituting; keep at least 4 significant figures until the final answer.
- Always state the interpretation of r or r_s in words ("strong positive linear correlation").
- For prediction questions, choose the correct regression line: to estimate y given x , use y on x .
- Kendall's τ is more robust with small samples and ties, but Spearman's r_s is quicker to compute; know when each is appropriate.



Signed • OnBuch+

ONBUCH+ TEACHING TEAM

FOLLOWS THE OFFICIAL MINESEC SYLLABUS

OnBuch+