

BIOLOGY • UPPER SIXTH

UPPER SIXTH — BIOLOGY

CHAPTER 1

REPRODUCTION

SUGGESTED TIME: 21 PERIODS

IN THIS CHAPTER

This chapter explores how living organisms perpetuate their species: asexual and sexual reproduction in plants and animals, the human reproductive systems and their hormonal control, fertilization, embryonic development, birth and birth control. It ends with the study of life cycles (alternation of generations, metagenesis) and the patterns and hormonal regulation of growth.

- By the end of this chapter I can define reproduction and distinguish asexual from sexual reproduction with Cameroonian examples.
- By the end of this chapter I can describe the mechanisms of asexual reproduction in plants and animals.
- By the end of this chapter I can explain pollen and ovule formation, pollination and double fertilization in flowering plants.
- By the end of this chapter I can describe seed formation, dispersal and the conditions and stages of germination.
- By the end of this chapter I can label and state the functions of the parts of the male and female human reproductive systems.
- By the end of this chapter I can explain the hormonal control of the testes and ovaries, spermatogenesis, oogenesis and the menstrual cycle.



DOWNLOAD ON- BUCH

Free • Google Play
AI tutor, past papers,
results



GO ONBUCH+

All signed courses, unlimited AI
tutor, PDF sheets — OnBuch+
tab in the app

THIS SIGNED COURSE CONTAINS

COURSE

complete, illustrated,
step by step

METHODS

and worked examples

EXERCISES

exam style + solutions

SUMMARY SHEET

the essentials to
remember



Signed • OnBuch+

ONBUCH+ TEACHING TEAM
FOLLOWS THE OFFICIAL MINESEC SYLLABUS

OnBuch+

CONTENTS

1. Asexual Reproduction: Concepts and Mechanisms
2. Sexual Reproduction in Flowering Plants: Gamete Formation to Fertilization
3. Seeds, Dispersal and Germination
4. The Male Reproductive System and Spermatogenesis
5. The Female Reproductive System, Oogenesis and the Menstrual Cycle
6. Fertilization, Embryonic Development and Birth
7. Birth Control Methods
8. Life Cycles: Alternation of Generations and Metagenesis
9. Growth: Curves, Patterns and Hormonal Control
10. Integration activity
11. Methods and worked examples
12. Exercises
13. Solutions to the exercises
14. Summary sheet

CHAPTER OBJECTIVES

- By the end of this chapter I can define reproduction and distinguish asexual from sexual reproduction with Cameroonian examples.
- By the end of this chapter I can describe the mechanisms of asexual reproduction in plants and animals.
- By the end of this chapter I can explain pollen and ovule formation, pollination and double fertilization in flowering plants.
- By the end of this chapter I can describe seed formation, dispersal and the conditions and stages of germination.
- By the end of this chapter I can label and state the functions of the parts of the male and female human reproductive systems.
- By the end of this chapter I can explain the hormonal control of the testes and ovaries, spermatogenesis, oogenesis and the menstrual cycle.
- By the end of this chapter I can describe fertilization, embryonic development, the birth process and methods of birth control.
- By the end of this chapter I can compare alternation of generations and metagenesis with labelled life-cycle diagrams.

- By the end of this chapter I can interpret growth curves, distinguish growth patterns and explain the hormonal control of growth.

PREREQUISITES

- Cell structure and the differences between mitosis and meiosis (Lower Sixth).
- Structure of a flower and its floral parts (Lower Sixth / Form 5).
- Basic notions of hormones and endocrine glands (Form 5 / Lower Sixth).
- Chromosomes, genes and the haploid/diploid distinction (Lower Sixth).
- General plant and animal tissue organisation (Lower Sixth).

1 ASEXUAL REPRODUCTION: CONCEPTS AND MECHANISMS

In a village near Bafoussam, Ma Ngo plants cassava stem cuttings that grow into whole new plants without any seed, while her neighbour's maize field must be sown with seeds every season. At the Bamenda regional hospital, a midwife explains to a young couple why a pregnancy test turned positive only two weeks after fertilization. Meanwhile, a Form Five student wonders why her younger brother suddenly grew taller and developed a deeper voice at age fourteen. All these everyday Cameroonian observations are governed by one great biological theme: reproduction and growth. This chapter unravels the cellular, hormonal and developmental mechanisms behind them — knowledge that is heavily examined at the GCE Advanced Level.

We begin with the simplest strategy of all: reproduction without sex, without a partner, and without gametes.

1.1 Reproduction and Its Two Great Strategies

From observation to concept.

Drop a single Amoeba into a drop of pond water and return a few days later: you may find several. No mate was involved, no egg, no sperm. Yet the population increased. Reproduction is clearly something more general than the familiar union of male and female.

DEFINITION Reproduction

Reproduction is the biological process by which living organisms produce new individuals of their own kind, thereby ensuring the continuity of the species from one generation to the next.

Biologists recognise two fundamental strategies:

- **Asexual reproduction**: a single parent produces offspring without the formation or fusion of gametes. Because all the offspring inherit their genetic material (by mitosis) from one parent only, they are **genetically identical** to that parent and to one another.
- **Sexual reproduction**: two parents (or two gamete-producing structures) contribute gametes which fuse (fertilization) to form a zygote. The offspring receive a new combination of genes and are therefore genetically different from each parent.

DEFINITION Clone

A **clone** is a group of genetically identical individuals derived from a single parent by asexual reproduction (or, artificially, by cloning techniques).

EXAMPLE Clones around us

All the plantain suckers taken from one mother plant in a plantation near Kumba are clones of that plant. Likewise, all the bacteria in a colony grown from one cell on an agar plate form a clone.

1.2 Advantages and Disadvantages of Asexual Reproduction

Why should an organism give up the genetic mixing of sexual reproduction? The answer lies in a balance of costs and benefits:

	Advantages	Disadvantages
Speed	Very rapid population increase (a bacterium can divide every 20 minutes)	—
No mate needed	A single isolated individual can reproduce; useful in sparse populations or sessile organisms	—
Energy	No energy wasted producing gametes, courtship or finding partners	—
Fidelity	Well-adapted genotypes are preserved exactly	No genetic variation: offspring cannot easily adapt if the environment changes
—	—	Accumulation of harmful mutations and uniform susceptibility: one disease can wipe out an entire clone

WARNING The variation trap

A very common exam error is to write that asexual reproduction produces “weak” offspring. This is wrong: clones can be perfectly vigorous. The correct point is that they show no genetic variation (apart from rare mutations), so the whole population is equally vulnerable to a new parasite, drought or pesticide.

1.3 Mechanisms of Asexual Reproduction in Animals and Micro-organisms

Binary fission.

In **binary fission** the parent cell grows, duplicates its genetic material, and then divides into two approximately equal daughter cells, each of which becomes an independent organism. This is the mode of reproduction of Amoeba, Paramecium, and of bacteria.

In *Amoeba*, the nucleus first divides by mitosis, then the cytoplasm constricts in the middle until two daughter amoebae separate. In bacteria, the circular chromosome is replicated, the two copies move apart, and a new transverse cell wall grows across the middle of the cell.

Budding.

In **budding** a small outgrowth (bud) forms on the parent body, grows, and eventually detaches as a new individual. The offspring is initially much smaller than the parent — this distinguishes budding from binary fission, where the two products are roughly equal. Budding occurs in *Hydra* (a freshwater cnidarian) and in yeast (*Saccharomyces*), the fungus used in baking and in brewing palm wine and “sha’a”.

Fragmentation.

In **fragmentation** the body of the parent breaks into two or more pieces, and each piece regenerates the missing parts to become a complete organism. The filamentous alga *Spirogyra* reproduces this way when its filaments break; each fragment grows by cell division into a new filament. Among animals, flatworms such as planarians show remarkable fragmentation: a worm cut into pieces can regenerate a complete individual from each piece.


Sporulation (spore formation).

Many organisms produce **spores**: small, usually unicellular reproductive bodies, often protected by a resistant wall, each capable of developing directly into a new individual without fertilization. Spore formation occurs in fungi (e.g. *Mucor* and *Rhizopus*, the black bread mould, which produces sporangia full of spores), in mosses and in ferns (sporangia on the underside of fern fronds). Spores are light and are dispersed by wind, which spreads the organism widely.

Binary fission (*Amoeba*)



Fragmentation (*Spirogyra*)



Budding (*Hydra*)



each fragment grows into a new filament

Three mechanisms of asexual reproduction. Top: binary fission in *Amoeba* — nucleus divides, then cytoplasm splits into two equal daughters. Bottom left: budding in *Hydra* — a small bud grows on the parent and detaches. Right: fragmentation in *Spirogyra* — the filament breaks and each fragment grows into a complete filament.

KEY POINTS Key distinctions


- Binary fission: parent splits into two equal daughters (*Amoeba*, bacteria).
- Budding: unequal division — a small bud on a larger parent (*Hydra*, yeast).

- **Fragmentation:** body breaks into pieces, each regenerates a whole organism (Spirogyra, planaria).
- **Sporulation:** resistant spores develop directly into new individuals (fungi, mosses, ferns).

1.4 Vegetative Propagation in Plants

Natural vegetative propagation.

Flowering plants possess specialised vegetative organs — modified stems, roots or leaves — that can detach or spread and give rise to new plants. This is **vegetative propagation**: asexual reproduction using vegetative (non-reproductive) organs.



Classification of natural vegetative propagules with a familiar Cameroonian example for each. Runners are horizontal aerial stems; rhizomes are horizontal underground stems; tubers are swollen storage stems or roots; bulbs are compressed stems with fleshy leaves; suckers are shoots arising from the base of the mother plant.

- **Runners:** horizontal stems growing along the soil surface; at each node, roots and a new shoot develop (grasses, strawberry).
- **Rhizomes:** horizontal underground stems with nodes and scale leaves; buds on the rhizome produce new shoots (ginger, canna lily).
- **Tubers:** swollen underground storage organs. Stem tubers such as yam and Irish potato bear “eyes” (buds) that sprout into new plants.
- **Bulbs:** a very short stem surrounded by fleshy storage leaves (onion); lateral buds develop into new bulbs.
- **Suckers:** shoots arising from buds at the base of the mother plant or on its roots (plantain, banana, pineapple).

Artificial vegetative propagation.

Farmers and horticulturists exploit the same principle deliberately:

- **Cuttings:** a piece of stem (cassava, hibiscus, sugar cane) or root is planted and develops adventitious roots and shoots.
- **Grafting:** a shoot or bud (the scion) of a desirable variety is joined to the rooted stem (the stock) of another plant. Widely used for mango, citrus and improved cocoa varieties in the

South West and Centre Regions.

- **Layering:** a low branch is bent to the ground and covered with soil; it roots while still attached to the parent, then is separated.
- **Tissue culture (micropropagation):** tiny pieces of plant tissue are grown on sterile nutrient medium to produce thousands of identical plantlets — used for disease-free plantain and banana planting material.

METHOD Demonstrating vegetative propagation with a cassava cutting

1. Select a healthy mature cassava stem and cut portions about 20 cm to 25 cm long, each bearing at least 3–4 nodes (buds). Cut the lower end obliquely.
2. Plant the cuttings upright (or slightly slanting) in moist, well-drained soil or in a trough of moist sawdust, burying about two-thirds of the length. Keep some cuttings upside-down as a control.
3. Water regularly and observe weekly for 4–6 weeks. Record the appearance of shoots (from nodes above ground) and adventitious roots (from buried nodes).
4. Conclusion: a complete new plant has formed from a stem fragment, without seed — evidence of vegetative propagation. Note that polarity matters: inverted cuttings root poorly.

WARNING A tuber is not a seed!

A classic GCE trap: candidates write that yam or Irish potato “seeds” are planted. Wrong. A seed is a fertilized ovule containing an embryo — the product of sexual reproduction. A tuber is a swollen stem or root used for vegetative propagation. Saying “yam seeds” or “cassava reproduces by seeds” in an essay costs marks. Similarly, do not confuse rhizome (underground stem, with nodes) with root tuber.

1.5 Parthenogenesis: A Special Case

In a few animals, an **unfertilized egg** can develop into a new individual. This is **parthenogenesis** (from Greek parthenos, virgin). Although an egg cell is involved, there is no fertilization, so the process is functionally asexual.

- In **aphids**, females produce generation after generation of daughters from unfertilized eggs during the favourable season — a spectacularly rapid multiplication on crop plants.
- In **honey bees**, unfertilized eggs develop into males (drones), while fertilized eggs develop into females (workers and queens). This is exploited by beekeepers in the North West Region.

DID YOU KNOW? Vegetative propagation and the Cameroonian economy

Cameroon's food security rests heavily on vegetative propagation. Cassava, the staple of the forest zone, is multiplied almost exclusively by stem cuttings; plantain and banana, the backbone of the Littoral and South West food economy, are propagated by suckers. Improved cocoa and robusta coffee planting material distributed to farmers is produced by grafting and rooted cuttings, so that the desirable traits of high-yielding, disease-tolerant clones are preserved exactly — something seeds could not guarantee.

WORKED EXAMPLE 1 Worked exercise — GCE style

Statement. (a) Define asexual reproduction. (b) A farmer near Bafoussam multiplies her plantain plantation using suckers, while her neighbour propagates improved cocoa by grafting. State the type of reproduction involved in each case, and give two advantages and one disadvantage of multiplying crops this way.

SOLUTION

Solution.

(a) Asexual reproduction is the production of offspring from a single parent without the formation or fusion of gametes, so that the offspring are genetically identical to the parent (clones).

(b) Both suckers and grafting are vegetative propagation, a form of asexual reproduction.

Advantages (any two):

- Rapid multiplication — many plantlets obtained in one season;
- No need for a mate or pollinator, and no waiting for seed production;
- The desirable qualities of the parent (yield, taste, disease tolerance) are preserved exactly in the clones.

Disadvantage (any one):

- No genetic variation: all plants are equally susceptible to a new disease or pest;
- Harmful mutations or viral infections of the parent are transmitted to all offspring.

EXERCISE 1 Check your understanding

1. Distinguish clearly between binary fission and budding, naming one organism for each.
2. Explain why a field of cassava planted from cuttings of a single parent plant could be

devastated by one viral disease, whereas a maize field grown from seed might survive.

3. Classify the following as natural or artificial vegetative propagation, and name the organ involved: (i) ginger spreading in a garden; (ii) a farmer planting cassava stems; (iii) onion bulbs multiplying; (iv) a cocoa scion joined to a stock.
4. A planarian is cut into three pieces and three complete worms are obtained. Name and define the process.

SOLUTION Exercise — Check your understanding

1. Binary fission: the parent divides into two equal daughter cells (e.g. Amoeba, bacteria). Budding: a small outgrowth forms on the parent and detaches as a smaller new individual (e.g. Hydra, yeast).
2. The cassava plants are clones: they share identical genes, so none possesses resistance to the virus and all are infected. The maize plants, produced sexually, are genetically varied, so some individuals may carry resistance and survive.
3. (i) Natural — rhizome (underground stem); (ii) artificial — stem cutting; (iii) natural — bulb; (iv) artificial — grafting.
4. Fragmentation: asexual reproduction in which the parent's body breaks into pieces, each of which regenerates the missing parts to form a complete new organism.

KEY POINTS Exam tip — GCE Advanced Level

Whenever you are asked to describe a type of asexual reproduction, always: (1) define it precisely, (2) describe the mechanism step by step, and (3) give a named example (Amoeba, Hydra, Spirogyra, Mucor, cassava, plantain...). An answer without a named example rarely earns full marks. And remember: tubers, rhizomes and cuttings are not seeds.

2 SEXUAL REPRODUCTION IN FLOWERING PLANTS: GAMETE FORMATION TO FERTILIZATION

In the previous section, we saw that asexual reproduction is fast and economical, but it produces offspring that are genetically identical to the parent. In a changing environment — a new parasite, a drought, a shift in climate — such uniformity can be fatal. Nature's answer is **sexual reproduction**: the mixing of genetic material from two parents so that every offspring is

a unique combination. In flowering plants (angiosperms), this process is centred on a remarkable organ: the **flower**. In this section we follow the whole journey, from the formation of pollen and ovules, through pollination, to the unique event of double fertilization.

2.1 The Concept of Sexual Reproduction

DEFINITION Sexual reproduction

Sexual reproduction is the type of reproduction involving the fusion of two haploid gametes — a male gamete and a female gamete — during **fertilization**, to form a diploid **zygote** which develops into a new individual.

Because each gamete carries a different shuffled set of genes (thanks to meiosis and random fertilization), the zygote — and therefore the offspring — is genetically different from both parents and from its siblings. This **genetic variation** is the raw material on which natural selection acts; it is the great advantage of sexual reproduction over asexual reproduction.


PROPERTY Advantage of sexual reproduction

Sexual reproduction produces genetic variation among offspring, increasing the chance that some individuals survive environmental changes, diseases or new predators. Its costs are that it is slower, requires two parents (or both sex organs), and uses more energy than asexual reproduction.

2.2 Structure of a Typical Flower

A flower is essentially a shoot modified for reproduction. Its parts are arranged in four **whorls** on a swollen tip of the stalk called the **receptacle**:

- **Sepals** (collectively the calyx): usually green, they protect the flower in bud.
- **Petals** (collectively the corolla): often large, coloured and scented to attract pollinators.
- **Stamens** (the androecium, male part): each consists of an **anther**, which produces pollen, borne on a stalk called the **filament**.
- **Carpels** (the gynoecium, female part): each consists of a sticky **stigma** (receives pollen), a **style** (stalk supporting the stigma) and an **ovary** containing one or more **ovules**.



Longitudinal section through a typical insect-pollinated flower, showing the four whorls: sepals, petals, stamens (anther + filament) and carpel (stigma, style, ovary with ovule).

2.3 Pollen Formation: Microsporogenesis and Microgametogenesis

Pollen grains are the male gametophytes of flowering plants. They are produced inside the **anther**, which typically has four pollen sacs (microsporangia). The process occurs in two phases: **microsporogenesis** (meiosis) and **microgametogenesis** (mitotic divisions).

METHOD Microsporogenesis and pollen maturation



1. Inside each pollen sac, diploid **pollen mother cells** (microsporocytes, $2n$) are formed from the sporogenous tissue.
2. Each pollen mother cell undergoes meiosis, producing a tetrad of four haploid microspores (n).
3. Each microspore matures into a **pollen grain** (the male gametophyte): its nucleus divides by mitosis to give a large **tube nucleus** (vegetative nucleus) and a smaller **generative nucleus**. The wall thickens into a resistant outer exine (sporopollenin) and an inner intine (cellulose/pectin).
4. In many species, the generative nucleus divides by a second mitosis after pollination, inside the growing pollen tube, to form two male gamete nuclei (sperm cells). In some species this division occurs before pollen release (bicellular vs tricellular pollen).

2.4 Ovule Formation: Megasporogenesis and Megagametogenesis

Inside the ovary, each **ovule** is attached to the placenta by a stalk (the **funicle**) and is protected by one or two **integuments**, leaving a small pore, the **micropyle**. The female gametophyte (embryo sac) develops as follows:

METHOD Megasporogenesis and embryo sac development (Polygonum type)

1. A diploid **megaspore mother cell** (megasporeocyte, $2n$) forms in the ovule tissue (nucellus).
2. It undergoes meiosis to give four haploid megaspores (n); three degenerate and only one functional megaspore survives (usually the chalazal one).
3. The nucleus of the functional megaspore divides by three successive mitotic divisions (without cytokinesis at first) to give eight haploid nuclei.
4. These organise into the **embryo sac**, which is 7-celled and 8-nucleate:
 - At the micropylar end: one **egg cell** flanked by two **synergids** (together forming the egg apparatus).
 - At the chalazal end: three **antipodal cells**.
 - In the centre: a large **central cell** containing two polar nuclei (which often fuse before fertilization to form a diploid secondary nucleus, $2n$).

Microsporogenesis (anther)**Megasporogenesis (ovule)**

Microsporogenesis (left) and megasporogenesis (right). Meiosis produces haploid spores; mitosis then builds the pollen grain (tube + generative nuclei) and the 7-celled, 8-nucleate embryo sac.

WARNING Ploidy trap

All eight nuclei of the embryo sac are haploid (n) because they come from mitotic divi-

sions of one haploid megaspore. The pollen mother cell and megaspore mother cell are diploid ($2n$). Do not confuse the two polar nuclei ($n + n$) with a diploid nucleus — they only become part of a triploid ($3n$) nucleus after fusion with a male gamete at fertilization. Note: the two polar nuclei often fuse before fertilization to form a diploid ($2n$) central cell nucleus; the endosperm is then formed by fusion of this $2n$ nucleus with a male n nucleus, still giving $3n$.

2.5 Pollination

DEFINITION Pollination

Pollination is the transfer of pollen grains from the anther to the stigma of a flower.

Two situations are possible:

- **Self-pollination:** pollen lands on the stigma of the same flower (autogamy), or of another flower on the same plant (geitonogamy).
- **Cross-pollination:** pollen is carried to the stigma of a flower on a different plant of the same species (xenogamy).

PROPERTY Self- versus cross-pollination

Cross-pollination mixes genes from two different parents and therefore increases genetic variation, giving offspring a better chance of adapting to change. **Self-pollination** produces more uniform offspring and preserves desirable characters (useful to farmers who want a stable crop variety), and it can occur even when pollinators are scarce.

Agents of pollination. Pollen may be carried by **insects** (bees, butterflies), **wind**, **water** (in some aquatic plants e.g. *Vallisneria*) or **animals** such as birds (sunbirds) and bats. Each agent corresponds to a set of floral adaptations.

METHOD Comparing insect- and wind-pollinated flowers

To distinguish the two types at the GCE, examine the following features:

Feature	Insect-pollinated	Wind-pollinated
Petals	Large, brightly coloured	Small, dull or absent
Scent and nectar	Scented; nectar produced	No scent, no nectar
Pollen grains	Few, large, sticky or spiny	Many, small, light, smooth
Stigma	Small, sticky, inside flower	Large, feathery, hangs outside
Stamens	Enclosed in flower	Long filaments, anthers hang out

Examples: hibiscus, flamboyant, cocoa (insect-pollinated); maize, grasses, oil palm (wind-pollinated).

Mechanisms favouring cross-pollination. Many plants reduce self-pollination by:

- **Dichogamy:** anthers and stigma of the same flower mature at different times (anthers first = **protandry**; stigma first = **protogyny**).
- **Self-incompatibility:** pollen landing on the stigma of the same plant fails to germinate or grows too slowly to fertilize (genetic recognition).
- **Unisexual flowers:** male and female flowers are separate, either on the same plant (**monoecious**, e.g. maize, oil palm) or on different plants (**dioecious**, e.g. pawpaw, date palm).
- **Herkogamy:** physical separation of anthers and stigma within the same flower.

WARNING Pollination is NOT fertilization

A very common GCE mistake: pollination is only the transfer of pollen to the stigma; fertilization is the fusion of gamete nuclei which happens later, after the pollen tube has grown down the style. Pollination always precedes fertilization.

2.6 Double Fertilization in Angiosperms


Once a compatible pollen grain lands on the stigma, it germinates: the tube nucleus directs the growth of a **pollen tube** down through the style towards an ovule. This growth is guided by chemical signals (chemotropism) from the synergids. The generative nucleus divides by mitosis into **two male gamete nuclei** (if not already divided). The pollen tube enters the ovule through the **micropyle**, penetrates one synergid (which degenerates), and releases the two male nuclei into the embryo sac.

DEFINITION Double fertilization

Double fertilization is the unique event in angiosperms in which one male nucleus (n) fuses

with the egg cell (n) to form a diploid zygote ($2n$), while the second male nucleus (n) fuses with the two polar nuclei ($n + n$) or the fused secondary nucleus ($2n$) to form a triploid endosperm nucleus ($3n$).

The zygote develops into the **embryo**; the triploid endosperm develops into the **food store** that nourishes it; the ovule becomes the **seed** and the ovary becomes the **fruit** — the subject of the next section. The synergids and antipodal cells degenerate after fertilization.



Double fertilization. The pollen tube enters the ovule through the micropyle and discharges two male nuclei: one fuses with the egg to form the diploid zygote, the other with the two polar nuclei to form the triploid endosperm.

WORKED EXAMPLE 2 Ploidy of embryo sac cells and fertilization products

A flowering plant has a diploid number $2n = 24$. State the number of chromosomes in: (a) a pollen mother cell; (b) a microspore; (c) the egg cell; (d) a polar nucleus; (e) the zygote; (f) the endosperm nucleus.

SOLUTION

Since $2n = 24$, the haploid number is $n = 12$.

- (a) Pollen mother cell: diploid, 24.
- (b) Microspore (product of meiosis): haploid, 12.
- (c) Egg cell: haploid, 12.
- (d) Each polar nucleus: haploid, 12.

- (e) Zygote ($n + n$): 24.
- (f) Endosperm nucleus ($n + n + n$): triploid, 36.

WARNING GCE exam tip: know your ploidy

Examiners frequently ask for the ploidy of every cell in the embryo sac and its products. Memorise this summary: pollen mother cell and megaspore mother cell = $2n$; microspores, megaspores, pollen nuclei, egg, synergids, antipodals and polar nuclei = n ; zygote = $2n$; endosperm = $3n$. Note that the endosperm is triploid, never diploid — this is the classic trap.

EXERCISE 2 Check your understanding

1. Define pollination and distinguish clearly between pollination and fertilization.
2. A maize plant and a hibiscus plant are grown side by side. For each, state the likely pollinating agent and give three floral features that support your answer.
3. Draw and label the embryo sac of a flowering plant just before fertilization, indicating the ploidy of each nucleus.
4. Explain two mechanisms by which a flower may favour cross-pollination, and state the advantage this confers.
5. Describe the fate of the two male nuclei after the pollen tube enters the embryo sac.

SOLUTION Exercise 1

1. Pollination is the transfer of pollen grains from anther to stigma. Fertilization is the fusion of male and female gamete nuclei; it occurs after pollination, once the pollen tube has reached the embryo sac.
2. Maize: wind-pollinated — dull flowers without petals or nectar, large feathery stigmas hanging outside, abundant small light pollen. Hibiscus: insect-pollinated — large colourful petals, nectar and scent, sticky stigma and large sticky pollen grains.
3. The drawing should show an oval sac with: egg cell (n) and two synergids (n) at the micropylar end, three antipodals (n) at the opposite end, and two polar nuclei (n each) in the central cell (often fused as a $2n$ secondary nucleus).

4. Any two of: dichogamy (anthers and stigma mature at different times, preventing self-pollination within a flower); self-incompatibility (own pollen fails to grow on the stigma); unisexual flowers (male and female flowers separated); herkogamy (physical separation). Advantage: cross-pollination mixes genes from two parents, increasing genetic variation and adaptability of the offspring.
5. The generative nucleus (if not already divided) divides by mitosis to form two male nuclei. One fuses with the egg cell to form the diploid zygote ($2n$); the other fuses with the two polar nuclei (or the fused secondary nucleus) to form the triploid endosperm nucleus ($3n$).

KEY POINTS Key points

- Sexual reproduction is the fusion of haploid gametes to form a diploid zygote; it generates genetic variation.
- The stamen (anther + filament) is male; the carpel (stigma, style, ovary) is female.
- Microsporogenesis: pollen mother cell ($2n$) $\xrightarrow{\text{meiosis}}$ 4 microspores (n) → pollen grain with tube and generative nuclei (microgametogenesis).
- Megasporogenesis: megaspore mother cell ($2n$) $\xrightarrow{\text{meiosis}}$ 1 functional megaspore (n) → 7-celled, 8-nucleate embryo sac (megagametogenesis).
- Pollination = transfer of pollen to the stigma; it precedes fertilization. Cross-pollination increases variation; self-pollination preserves characters.
- Double fertilization: male (n) + egg (n) → zygote ($2n$); male (n) + polar nuclei ($n + n$) → endosperm ($3n$).

3**SEEDS, DISPERSAL AND GERMINATION**

Having explored how pollen grains reach the stigma and how double fertilization gives rise to a diploid zygote and a triploid endosperm, we now follow the remarkable transformation of the fertilized ovule into a seed and the ovary into a fruit. This section connects the reproductive act to the next generation: how the embryo is packaged with food reserves, how the fruit protects and disperses it, and how the dormant embryo springs back to life during germination.

3.1 From Fertilized Ovule to Seed and Fruit

Immediately after double fertilization, the flower undergoes dramatic changes. The sepals, petals, stamens, style and stigma usually wither and fall off (they are deciduous), having completed their roles. The fate of the remaining parts is summarized in the table below.

Flower part	Fate after fertilization
Ovule	Becomes the seed
Ovary	Becomes the fruit (pericarp develops from ovary wall)
Zygote (2n)	Develops into the embryo (radicle, plumule, cotyledon(s))
Primary endosperm nucleus (3n)	Divides to form endosperm (nutritive tissue)
Integuments of ovule	Become the testa (seed coat)
Micropyle	Remains as a small pore in the testa
Funiculus	Becomes the stalk attaching seed to fruit; leaves hilum (scar) when detached

DEFINITION Seed


A seed is a mature, fertilized ovule containing an embryonic plant, a food store (endosperm or cotyledons), and a protective seed coat (testa), capable of developing into a new plant under favourable conditions.

DEFINITION Fruit

A fruit is a mature, ripened ovary (often with accessory tissues) that encloses the seeds and aids in their protection and dispersal.

3.2 Structure of Dicot and Monocot Seeds

Seeds vary greatly in size, shape and internal organization. The fundamental distinction lies in the number of cotyledons: **dicotyledonous** (two cotyledons) and **monocotyledonous** (one cotyledon). We examine two classic examples: the bean (*Phaseolus vulgaris*) and the maize grain (*Zea mays*).



Longitudinal sections of a dicot seed (bean) and a monocot grain (maize). Note the large cotyledons storing food in bean, versus the single scutellum absorbing food from the massive endosperm in maize.

KEY POINTS Key structural differences

- **Bean (dicot):** Food stored in two large cotyledons; endosperm absorbed during seed development. Embryo consists of radicle, plumule and two cotyledons attached to a short hypocotyl.
- **Maize (monocot):** Food stored mainly in endosperm; single cotyledon modified into scutellum (absorptive shield). Embryo protected by coleoptile (sheath covering plumule) and coleorhiza (sheath covering radicle). Pericarp (fruit wall) fused to testa.

3.3 Fruit Types (Brief Overview)

Fruits are classified primarily by the texture of the pericarp (ovary wall) and whether they split open at maturity.

Category	Type	Description and Cameroonian examples
Dry	Dehiscent (split open)	Legume/Pod: splits along two sutures (bean, groundnut, flamboyant). Capsule: splits in various ways (cotton, okra).
Dry	Indehiscent (do not split)	Caryopsis (grain): pericarp fused to testa (maize, rice, millet, guinea corn). Achene: single seed, free pericarp (sunflower). Nut: hard woody pericarp (cashew).
Fleshy	Berry: many seeds, fleshy throughout (tomato, guava, papaya). Drupe: one seed enclosed in hard stone (mango, plum, coconut). Pome: fleshy swollen receptacle, true fruit is core (apple — less common in Cameroon).	

DID YOU KNOW? Cameroon context

The Flamboyant (*Delonix regia*) pods explode violently on hot dry days, scattering seeds metres away — a classic example of explosive dispersal. The mango (*Mangifera indica*), a drupe, relies on bats, birds and humans eating the fleshy mesocarp and discarding the stone. Maize (*Zea mays*), a caryopsis, is primarily dispersed by humans (agriculture) but its light chaff can be wind-blown.

3.4 Seed and Fruit Dispersal

Dispersal is the scattering of seeds and fruits away from the parent plant. It is crucial for reducing competition among siblings and with the parent for light, water and minerals, and for colonizing new habitats.

DEFINITION Dispersal


Dispersal is the movement, spread or transport of seeds/fruits away from the parent plant, achieved by structural adaptations that exploit abiotic (wind, water) or biotic (animals) agents, or by the plant's own mechanical forces.

PROPERTY Importance of dispersal

1. Reduces intraspecific competition (overcrowding leads to weak, disease-prone seedlings).
2. Enables colonization of new, potentially favourable habitats (range expansion).
3. Reduces predation and pathogen buildup near parent plant.
4. Promotes genetic mixing by separating related individuals.

The table below links dispersal agents to structural adaptations. Memorize these pairings for the GCE.

Agent	Adaptation	Examples (Cameroon)
Wind (Anemochory)	Winged (samara), plumed/hairy (pappus), light/dust-like , balloon-like inflated calyx	Flamboyant (winged), Ceiba pentandra (kapok — hairy), Combretum (winged), orchids (dust seeds)
Water (Hydrochory)	Buoyant , waterproof , fibrous/spongy mesocarp/husk, air spaces	Coconut (fibrous husk), water lily (spongy fruit), mangrove propagules (buoyant)
Animals (Zoochory)	Edible fleshy fruit (endozoochory), hooks/spines (epizoochory), sticky mucilage	Mango, papaya, bush mango (Irvingia) — fleshy; Desmodium (beggar's lice), Xanthium (cocklebur) — hooks
Explosive / Self (Autochory)	Tension-building fruit walls that snap, coiling pods, explosive dehiscence	Flamboyant, Impatiens (balsam), Hura crepitans (sandbox tree)



Structural adaptations for dispersal by wind, water, animals and explosive mechanisms.

WARNING GCE Exam Tip

When asked to "state the agent of dispersal and the adaptation", give a single sentence linking them explicitly. Example: "The fruit is dispersed by wind because it has a winged pericarp (samaras) that increases air resistance and allows it to be carried far from the parent plant." Do not just list "wind" and "wing" separately.

3.5 Germination: Definition, Conditions and Biochemistry

DEFINITION Germination

Germination is the resumption of active growth of the embryo in a seed after a period of dormancy, resulting in the emergence of the radicle and plumule and the establishment of a seedling, provided favourable environmental conditions are met.

PROPERTY Conditions necessary for germination

Three external factors are essential:

1. **Water:** Rehydrates cells (imbibition), activates hydrolytic enzymes, provides medium for metabolic reactions, and ruptures the testa.
2. **Oxygen:** Required for aerobic respiration to release energy (ATP) from food reserves. Most seeds cannot germinate anaerobically (rice is an exception).
3. **Suitable temperature:** Enzymes have an optimum temperature range (typically 25 °C–35 °C for tropical crops). Too low: enzymes inactive; too high: enzymes denature.

Internal factors: Viable embryo, non-dormant state (dormancy broken), sufficient food reserves.

METHOD Mobilisation of food reserves during germination

1. Imbibition: Water enters via micropyle; cells swell, testa ruptures.
2. Hormonal signal: Embryo produces gibberellins (GA) which diffuse to the aleurone layer (in cereals) or cotyledons (in dicots).
3. Enzyme synthesis: GA stimulates synthesis of hydrolytic enzymes: amylases (starch → maltose/glucose), proteases (proteins → amino acids), lipases (lipids → fatty acids + glycerol).
4. Translocation: Soluble products transported via vascular tissue to growing points (radicle, plumule).
5. Utilization: Respiration (glycolysis, Krebs cycle, oxidative phosphorylation) provides ATP and carbon skeletons for new cell synthesis (cell division in meristems, cell elongation).


WARNING Common mistake

Do not confuse imbibition (physical water uptake) with germination (metabolic activation and growth). A dead seed can imbibe water but will not germinate. Also, light is not a universal requirement; some seeds need light (lettuce), some need darkness, most are indifferent.

3.6 Epigeal vs Hypogeal Germination

The behaviour of the cotyledons during germination defines two major patterns.

Feature	Epigeal germination	Hypogeal germination
Cotyledons	Pulled above ground by elongating hypocotyl ; become photosynthetic (green).	Remain below ground ; epicotyl elongates instead. Cotyledons stay in soil, non-photosynthetic.
Hypocotyl	Elongates vigorously, forms a hook that pulls cotyledons up.	Remains short; does not elongate significantly.
Epicotyl	Short initially; elongates later after cotyledons emerge.	Elongates vigorously, pushing plumule above ground.
Protection of plumule	Cotyledons enclose plumule; may be damaged during emergence.	Plumule protected by coleoptile (monocots) or by remaining inside cotyledons (dicots like pea).
Examples	Bean, castor oil, sunflower, cotton, flamboyant.	Maize, rice, millet, guinea corn, pea, groundnut, coconut.



Comparison of epigeal (bean) and hypogeal (maize) germination stages. Note the elongating hypocotyl pulling cotyledons up in bean, versus the elongating epicotyl/coleoptile pushing plumule up while cotyledon (scutellum) stays below in maize.

WARNING GCE Trap

In hypogeal germination, the cotyledon(s) remain underground. Do not draw them above ground! In maize, the scutellum never emerges; in pea (dicot but hypogeal), the cotyledons stay below soil. The key is: which axis elongates? Hypocotyl → epigeal; Epicotyl → hypogeal.

3.7 Experimental Investigation: Conditions for Germination

The classic controlled experiment uses four test tubes (or boiling tubes) to isolate each condi-

tion.

METHOD Four-tube experiment to demonstrate requirements for germination

1. Set up four tubes with cotton wool at the bottom and 10–20 viable seeds of the same species (e.g., bean) on top.
2. Tube A (Control): Cotton wool moistened with water; tube open to air; kept at room temperature (~ 25 °C). All conditions present.
3. Tube B (No water): Cotton wool dry; tube open; room temperature. Water absent.
4. Tube C (No oxygen): Cotton wool moistened with boiled and cooled water (drives out dissolved O₂); add a layer of liquid paraffin or pyrogallol + NaOH to prevent atmospheric O₂ diffusion; room temperature. Oxygen absent.
5. Tube D (Unsuitable temperature): Cotton wool moistened; tube open; placed in a refrigerator (~ 4 °C) or an incubator at 45 °C. Temperature unsuitable.
6. Observe daily for 5–7 days. Record which tubes show germination (radicle emergence).

WORKED EXAMPLE 3 Interpreting germination experiment results

Statement: In the four-tube experiment described above, after 5 days at room temperature, seeds in Tube A have germinated (radicles 10 mm long). Tubes B, C and D show no radicle emergence. Seeds from Tube B are transferred to moist cotton wool at room temperature; they germinate within 2 days. Seeds from Tube C are transferred to moist cotton wool open to air; they germinate within 2 days. Seeds from Tube D are transferred to room temperature; they germinate within 2 days.

Questions:

1. State the condition being tested in each of Tubes B, C and D.
2. Explain why boiled water cooled and covered with paraffin is used in Tube C.
3. What conclusion can be drawn about the viability of seeds in Tubes B, C and D?
4. Why is it important to use seeds from the same batch and same species?

SOLUTION

Detailed solution:

1. Tube B: Absence of water (dry cotton wool). Tube C: Absence of oxygen (boiled water + paraffin excludes O₂). Tube D: Unsuitable (low) temperature (refrigerator).
2. Boiling water removes dissolved oxygen. The layer of liquid paraffin (or oil) creates a

physical barrier preventing atmospheric oxygen from dissolving back into the water. This ensures anaerobic conditions.

3. The seeds in Tubes B, C and D are viable (alive) because they germinated once the missing condition was supplied. They were dormant due to the lack of that specific factor, not dead.
4. Using the same batch and species controls genetic variability and seed age/size, ensuring that any difference in germination is due only to the manipulated environmental factor (validity of the controlled experiment).

3.8 Worked Example: Linking Structure to Function

WORKED EXAMPLE 4 Seed structure and dispersal adaptation

Statement: The diagram below shows a longitudinal section of a fruit of *Xanthium* (cocklebur). The fruit is a dry, indehiscent achene covered in stiff, hooked spines. The seed inside has a thin testa and a large, oil-rich embryo.

1. Name the agent of dispersal for this fruit.
2. Explain how the structure of the fruit adapts it to this agent.
3. The seed germinates epigeally. Describe the role of the hypocotyl during germination.
4. State two environmental conditions necessary for the germination of this seed.

SOLUTION

Detailed solution:

1. Agent: Animal (epizoochory / external attachment to fur/feathers/clothing).
2. Adaptation: The fruit wall (pericarp) is modified into a hard, woody achene covered with stiff, hooked spines. These hooks catch onto the fur of passing mammals (or feathers/clothing of humans), carrying the fruit away from the parent plant. The fruit is indehiscent, so the seed remains protected inside until it falls off or the fruit decays.
3. Role of hypocotyl: In epigeal germination, the hypocotyl elongates rapidly, forming a hook that pushes through the soil. The hook straightens in response to light (phototropism), pulling the cotyledons above ground. The cotyledons then expand, become green and photosynthetic, providing energy for the seedling until true leaves form.
4. Conditions: (1) Water (for imbibition and enzyme activation). (2) Oxygen (for aéro-

bic respiration). (3) Suitable temperature (approx. 25 °C–30 °C). (Any two correctly stated.)

3.9 Summary of Key Points

KEY POINTS Seeds, Dispersal and Germination — Essentials for GCE

- Seed formation: Ovule → seed; Ovary → fruit; Zygote → embryo; Endosperm nucleus → endosperm; Integuments → testa; Funiculus → stalk/hilum.
- Dicot seed (bean): Testa, hilum, micropyle, two large cotyledons (food store), embryo (radicle, plumule, hypocotyl). No endosperm.
- Monocot grain (maize): Pericarp+testa fused, scutellum (single cotyledon, absorptive), large starchy endosperm, embryo with coleoptile & coleorhiza.
- Fruit types: Dry dehiscent (pod), dry indehiscent (caryopsis, nut), fleshy (berry, drupe). Local examples: bean pod, maize grain, mango, flamboyant.
- Dispersal agents & adaptations: Wind — wing/hair; Water — fibrous husk/buoyancy; Animal — fleshy edible / hooks; Explosive — tension-splitting pods.
- Importance of dispersal: Reduces competition, colonizes new habitats, reduces predation/disease.
- Germination conditions: Water, Oxygen, Suitable Temperature. (Light not universal).
- Biochemistry: Imbibition → GA synthesis → enzyme production (amylase, protease, lipase) → hydrolysis of reserves → translocation → respiration/growth.
- Epigeal: Hypocotyl elongates, cotyledons pulled above ground, become photosynthetic (bean).
- Hypogeal: Epicotyl elongates, cotyledons remain underground (maize, pea).
- Experiment: Four tubes (Control, –Water, –Oxygen, –Temp) prove all three factors essential. Viability confirmed by recovery.

EXERCISE 3 Practice Questions

1. Draw a large, labelled diagram of the longitudinal section of a maize grain. Indicate the positions of the scutellum, coleoptile, coleorhiza, endosperm and pericarp.
2. A student sets up the four-tube germination experiment but forgets to boil the water

in Tube C. What would be the likely result in Tube C, and why?

3. Distinguish clearly between the terms imbibition, germination and seedling establishment.
4. The fruit of *Triumfetta* (a common weed in Cameroon) has numerous hooked bristles. Name the dispersal agent and explain the advantage of this dispersal method for the species.
5. In an experiment, bean seedlings grown in the dark have long, pale hypocotyls and small, unexpanded cotyledons. Seedlings grown in light have short, green hypocotyls and large, expanded green cotyledons. Explain these observations.

SOLUTION Exercise 1

Maize grain LS: Oval outline (pericarp+testa fused). Large central endosperm (starchy). Embryo at one side: shield-shaped scutellum pressed against endosperm. Plumule covered by coleoptile (sheath). Radicle covered by coleorhiza (sheath). Labels: Pericarp/Testa, Endosperm, Scutellum, Coleoptile, Plumule, Coleorhiza, Radicle.

SOLUTION Exercise 2

Tube C would likely show germination because unboiled water contains dissolved oxygen. The paraffin layer alone does not remove oxygen already dissolved; it only prevents further diffusion. Boiling is essential to drive out dissolved O_2 .

SOLUTION Exercise 3

Imbibition: Physical absorption of water by dry seed (colloidal matrix swells), passive, occurs in dead seeds too. Germination: Metabolic activation of embryo, enzyme synthesis, hydrolysis of reserves, cell division/elongation, radicle emergence — requires viable seed. Seedling establishment: Post-germination phase where seedling becomes autotrophic (photosynthetic), true leaves form, root system develops, reliance on seed reserves ends.

SOLUTION Exercise 4

Agent: Animal (epizoochory). Advantage: Hooked bristles attach firmly to fur of passing animals (antelopes, cattle, humans), transporting fruits far from parent. Reduces com-

petition among dense seedlings under parent plant; colonizes new areas like paths, forest edges, farmland.

SOLUTION Exercise 5

Dark-grown seedlings show etiolation: hypocotyl elongates rapidly (skotomorphogenesis) to reach light, but lacks chlorophyll (pale/yellow) because chlorophyll synthesis requires light (protochlorophyllide → chlorophyllide needs photons). Cotyledons remain small and unexpanded (no photosynthetic demand). Light-grown seedlings: light inhibits hypocotyl elongation (photomorphogenesis), promotes chlorophyll synthesis (green), and stimulates cotyledon expansion for photosynthesis.


4 THE MALE REPRODUCTIVE SYSTEM AND SPERMATOGENESIS

In the previous sections we followed the flowering plant from pollination to the germination of a new seedling. We now turn to reproduction in humans, and we begin with the male partner. Every day, from puberty until old age, a healthy man produces several hundred million spermatozoa — yet each ejaculate releases cells so small that about two thousand of them could fit on the head of a pin. How is such a prodigious, continuous output organised? The answer lies in a beautifully coordinated system of organs, tubes, glands and hormones, which we now study in detail.

4.1 Overview of the Male Reproductive System

DEFINITION Male reproductive system

The **male reproductive system** is the set of organs that produce male gametes (**spermatozoa**), nourish and store them, and deliver them into the female reproductive tract during copulation. It comprises the testes (held in the scrotum), a system of ducts (epididymis, vas deferens, urethra), accessory glands (seminal vesicles, prostate gland, Cowper's glands) and the penis.



Simplified sagittal section of the male reproductive system, showing the testis in the scrotum, the sperm ducts and the accessory glands.

4.2 Structure and Functions of the Organs

The testes and the scrotum.

The two **testes** are the male gonads: they produce spermatozoa and secrete the male sex hormone **testosterone**. Each testis is an oval organ about 4 cm long and 2.5 cm wide, suspended outside the abdominal cavity in a skin pouch, the **scrotum**. This external position is not accidental. Spermatogenesis is extremely temperature-sensitive and proceeds normally only at about 2 °C–3 °C below core body temperature (i.e. around 34 °C–35 °C instead of 37 °C). The scrotum acts as a thermostat: in cold weather its muscles (dartos and cremaster) contract and draw the testes closer to the warm body; in hot weather it relaxes so the testes hang further away and lose heat. This is why tight clothing that keeps the testes pressed against the body can temporarily reduce sperm production — a fact relevant to reproductive health advice in our climate.

The epididymis.

A long, highly coiled tube (about 6 m if uncoiled) lying on the posterior surface of each testis. Newly formed spermatozoa are immature and cannot swim; during their journey through the epididymis, which takes 10 --14 days, they **mature**, gain motility and are **stored** until ejaculation. The epididymis also absorbs excess fluid and phagocytoses defective sperm.

The vas deferens (sperm duct).

A muscular tube, about 45 cm long, that carries spermatozoa from the epididymis up into the abdomen, looping over the ureter and the bladder to join the urethra at the ejaculatory duct. During ejaculation, waves of **peristalsis** in its thick muscular wall propel the sperm forward. Cutting and tying the vas deferens is the principle of male sterilisation (vasectomy), studied in the section on birth control.

The accessory glands.

Spermatozoa alone would form a tiny, sticky mass. Three sets of glands add fluids that together with sperm form **semen** (typical volume 2 --5 ml per ejaculate):

- the **seminal vesicles** (paired, 5 cm long) secrete an alkaline fluid (~60 % of semen volume) rich in **fructose** (the main energy source for sperm swimming), prostaglandins (which stimulate uterine contractions to help sperm transport), and clotting proteins;
- the **prostate gland** (single, chestnut-sized) secretes a milky, slightly acidic fluid (~30 % of volume) containing citrate, enzymes (including prostate-specific antigen, PSA) and zinc; it helps **neutralise the acidity** of the female vagina, which would otherwise kill the spermatozoa;
- the **Cowper's (bulbo-urethral) glands** (paired, pea-sized) secrete a clear, viscous, alkaline mucus before ejaculation, which lubricates the urethra and neutralises traces of acidic urine.

The urethra and the penis.

The **urethra** (18 --20 cm long) is the common passage for urine and semen (never at the same time: the internal urethral sphincter closes the bladder during ejaculation). The **penis** contains three columns of **erectile tissue**: two **corpora cavernosa** dorsally and one **corpus spongiosum** ventrally (which expands to form the glans penis). During sexual arousal, parasympathetic stimulation causes arterial dilation and venous compression, filling these spaces with blood and making the organ erect so it can be inserted into the vagina to deposit semen near the cervix.

KEY POINTS Functions at a glance

Organ	Function
Testis	Produces spermatozoa and testosterone
Scrotum	Keeps testes 2 °C–3 °C below body temperature
Epididymis	Maturation, storage and transport of spermatozoa
Vas deferens	Transports sperm by peristalsis during ejaculation
Seminal vesicles	Secrete fructose-rich fluid (energy), prostaglandins, clotting proteins
Prostate gland	Secretes fluid with enzymes, citrate, zinc; neutralises vaginal acidity
Cowper's glands	Lubrication; neutralise urine traces in urethra
Penis / urethra	Copulation and delivery of semen


4.3 Microscopic Structure of the Testis

If we cut a thin section of a testis and observe it under the microscope, we see that it is packed with hundreds of tightly coiled tubes, the **seminiferous tubules** (total length ~250 m per testis). It is inside these tubules that spermatozoa are manufactured. Between the tubules lie clusters of **interstitial cells** (also called **Leydig cells**), which secrete **testosterone**. Blood capillaries and lymphatic vessels are also found in this interstitial tissue.

The wall of each seminiferous tubule contains two important cell types:

- the **germinal epithelial cells** (spermatogonia, primary/secondary spermatocytes, spermatids) at different stages of division, which give rise to spermatozoa (the youngest cells lie at the outer edge near the basal lamina, the most mature near the central cavity or lumen);
- the large **Sertoli cells** (sustentacular cells), which stretch from the basal lamina to the lumen. They **nourish and support** the developing sperm cells, phagocytose excess cytoplasm, form the **blood–testis barrier** (protecting developing sperm from immune attack), and secrete the hormone **inhibin** and **androgen–binding protein (ABP)** which concentrates testosterone in the tubule.

Mature spermatozoa are released into the lumen of the tubule (spermiation) and carried by fluid towards the **rete testis**, then to the epididymis.



Transverse section of a seminiferous tubule showing the arrangement of germ cells and Sertoli cells.

4.4 Spermatogenesis: The Production of Spermatozoa

DEFINITION Spermatogenesis


Spermatogenesis is the production of spermatozoa inside the seminiferous tubules of the testes, from the germinal epithelium. It begins at puberty (around 12 --14 years in Cameroon) and continues continuously throughout adult life, unlike oogenesis in the female, which is cyclic and stops at the menopause. The entire process takes approximately 64 --74 days in humans.

The process combines **growth**, **meiosis** (which halves the chromosome number from diploid $2n$ to haploid n) and **differentiation (spermiogenesis)**. In humans $2n = 46$ and $n = 23$.

METHOD Following spermatogenesis step by step

1. Multiplication (mitotic phase): diploid spermatogonia ($2n$) at the tubule edge (near basal lamina) divide repeatedly by mitosis. Some remain as stem cells (type A spermatogonia) to maintain the germ line; others (type B spermatogonia) commit to differentiation.
2. Growth phase: type B spermatogonia enlarge to become primary spermatocytes ($2n$), entering prophase I of meiosis (which lasts ~22 days).
3. Meiosis I (reduction division): each primary spermatocyte completes meiosis I to form two haploid secondary spermatocytes (n). This division separates homologous chromosomes.
4. Meiosis II (equational division): each secondary spermatocyte rapidly divides (without DNA replication) to form two spermatids (n) — four spermatids per primary spermatocyte. This division separates sister chromatids.
5. Spermiogenesis (differentiation): each spermatid, still round and immobile, transforms into a streamlined spermatozoon: the nucleus condenses and elongates, the acrosome forms from the Golgi apparatus over the anterior nucleus, mitochondria gather into a tight helix in the middle piece, the flagellum (tail) grows from the centriole, and excess cytoplasm is shed as residual bodies (phagocytosed by Sertoli cells).

The whole sequence takes roughly 74 days and occurs continuously in waves along the tubule, so millions of spermatozoa are released every day.



Flow diagram of spermatogenesis. Meiosis halves the chromosome number from $2n$ to n ; one primary spermatocyte yields four functional spermatozoa.


WARNING Spermatogenesis versus oogenesis

Do not confuse the two processes. In spermatogenesis, one primary spermatocyte gives

four functional spermatozoa, and production is continuous from puberty. In oogenesis (next section), one primary oocyte gives only one functional ovum plus polar bodies, and the process is cyclic. Examiners love this comparison.

4.5 Structure of the Mature Spermatozoon

A human spermatozoon is about $60\ \mu\text{m}$ long and has three distinct regions, each adapted to its function of reaching and fertilising the ovum:



Structure of a mature human spermatozoon.

- the **head** (flattened, $5\ \mu\text{m}$ long), containing the highly condensed haploid **nucleus** (paternal genetic material) capped by the **acrosome**, a modified lysosome derived from the Golgi apparatus containing hydrolytic enzymes (hyaluronidase, acrosin) that digest the corona radiata and zona pellucida of the ovum at fertilisation;
- the **neck** (connecting piece) containing the proximal and distal centrioles;
- the **middle piece**, packed with a helical sheath of **mitochondria** which produce the ATP that powers flagellar movement;
- the **tail (flagellum)**, the longest part ($50\ \mu\text{m}$), whose axoneme (9+2 microtubule arrangement) beats in a whip-like motion to propel the cell through the female tract.

METHOD Drawing and labelling a spermatozoon (GCE practical skill)

1. Draw an oval head ($5\ \mu\text{m}$); shade a large nucleus inside it and add a cap-like acrosome over the anterior half.
2. Attach a short neck and a thicker middle piece ($5\ \mu\text{m}$) and fill it with small dots or a helical line representing mitochondria.
3. Draw a long, thin, tapering tail ($50\ \mu\text{m}$), at least ten times the head length.
4. Label each region and state its function in one line: nucleus — paternal chromosomes; acrosome — enzymes to penetrate ovum coats; mitochondria — ATP production; tail — propulsion.


4.6 Hormonal Control of the Testes

Why does sperm production start at puberty and not before? Because the testes are switched on by hormones arriving through the blood, in a chain of command involving three levels: the **hypothalamus** (brain), the **anterior pituitary gland**, and the **testes** themselves. This is the **hypothalamus–pituitary–testis axis**.

PROPERTY The hypothalamus–pituitary–testis axis

- At puberty the hypothalamus secretes **GnRH** (gonadotrophin-releasing hormone) in pulsatile fashion, which stimulates the anterior pituitary.
- The anterior pituitary releases two gonadotrophins: **FSH** (follicle-stimulating hormone), which acts on the Sertoli cells and, together with testosterone, drives spermatogenesis (stimulates ABP production, supports spermatogonial differentiation); and **LH** (luteinising hormone, also called ICSH — interstitial cell stimulating hormone — in the male), which stimulates the Leydig cells to secrete testosterone.
- Testosterone acts locally on Sertoli cells (via ABP) to support spermatogenesis, and systemically to maintain secondary sexual characteristics and libido. It also exerts negative feedback on the hypothalamus (reducing GnRH pulse frequency) and anterior pituitary (reducing LH response to GnRH).
- The Sertoli cells secrete inhibin, which specifically inhibits FSH secretion from the anterior pituitary (negative feedback on FSH only).

(The names of the hormones, their cellular targets, and the logic of the feedback loops are examinable; learn to reproduce the diagram from memory.)



Hormonal control of the testes. GnRH stimulates FSH and LH release; testosterone and inhibin exert negative feedback (orange/pink arrows) on the hypothalamus and pituitary.

Roles of testosterone.

Testosterone is far more than a sperm-production hormone. It is responsible for:

- the development of the **secondary sexual characteristics** at puberty: deepening of the voice, growth of beard and body hair, broadening of shoulders, muscular development, growth of the penis and testes, increased bone density;
- the maintenance of **libido** (sexual drive) and erectile function;
- the **maintenance of spermatogenesis**, acting together with FSH on the seminiferous tubules (via ABP);
- anabolic effects on muscle and bone, erythropoiesis stimulation, and metabolic effects.

WARNING Common mistakes in hormonal control

- FSH and LH are secreted by the anterior pituitary, not by the testes. The testes are the target, not the source.
- Testosterone comes from the Leydig (interstitial) cells, not from the Sertoli cells. Sertoli cells nourish sperm and secrete inhibin and ABP.
- LH in the male is the same hormone as in the female; in the male it is often called ICSH (interstitial cell stimulating hormone) — a name that tells you exactly where it acts.
- GnRH is secreted in pulses; continuous GnRH (as from an agonist implant) down-regulates pituitary receptors and suppresses gonadotrophin release — the basis of medical castration.

WORKED EXAMPLE 5 Explaining negative feedback (typical GCE question)

Statement. A man receives regular injections of testosterone as part of a medical treatment. After several months, his sperm count falls dramatically. Explain this observation using your knowledge of the hormonal control of the testes.

SOLUTION

Solution.

1. The injected testosterone raises the blood testosterone level far above normal.
2. By negative feedback, the high testosterone level inhibits the hypothalamus (reduced GnRH pulse frequency) and the anterior pituitary (reduced sensitivity to GnRH).
3. The pituitary therefore secretes less LH and less FSH.
4. FSH, together with locally high intratesticular testosterone (maintained by LH), is required to drive spermatogenesis in the seminiferous tubules; with FSH and LH lacking, sperm production slows or stops, so the sperm count falls.

This is exactly the principle explored for a possible male contraceptive: suppressing FSH/LH by external hormones switches off sperm production reversibly.

DID YOU KNOW? Why so many sperm?

A single ejaculate contains 200 – 500 million spermatozoa, yet only one can fertilise the ovum. Most die on the way: the vagina is acidic (pH ~4), many swim in the wrong direction (only ~1% enter the cervix), and the journey to the oviduct is enormous relative to their size — comparable to a human swimming 100 km. Producing vast numbers is the male strategy that maximises the chance that at least one spermatozoon reaches the egg. In Cameroon, as elsewhere, a sperm count below 15 million per ml (oligospermia) is considered subfertile.

EXERCISE 4 Check your understanding

1. List, in order, the structures through which a spermatozoon passes from its production site to the outside of the body.
2. State precisely which cells secrete (a) testosterone, (b) inhibin, (c) FSH, (d) GnRH.
3. Explain why the testes are located outside the abdominal cavity.
4. A tumour destroys a man's anterior pituitary. Predict the effect on his testosterone level and sperm production, and justify your answer.

5. Describe the role of Sertoli cells in spermatogenesis.
6. What is the significance of the blood–testis barrier?

SOLUTION Exercise 1

1. Seminiferous tubules → rete testis → epididymis → vas deferens → ejaculatory duct → urethra → outside (via the penis).
2. (a) Leydig (interstitial) cells; (b) Sertoli cells; (c) anterior pituitary gonadotrophs; (d) hypothalamic neurosecretory cells.
3. Spermatogenesis requires a temperature 2°C – 3°C below core body temperature; the scrotum keeps the testes cooler than the abdomen, and adjusts their position via dartos and cremaster muscles to regulate temperature. Cryptorchidism (undescended testes) leads to infertility and increased cancer risk.
4. Without the anterior pituitary there is no LH and no FSH. Without LH, the Leydig cells are not stimulated, so testosterone secretion collapses; without FSH (and testosterone), spermatogenesis stops. The man becomes sterile and loses testosterone-dependent characteristics unless given hormone replacement therapy.
5. Sertoli cells: nourish developing sperm (provide nutrients, growth factors); phagocytose residual bodies; form blood–testis barrier (tight junctions) isolating adluminal compartment from immune system; secrete inhibin (FSH feedback), ABP (concentrates testosterone), and anti-Müllerian hormone (in fetal life).
6. The blood–testis barrier prevents immune cells from recognising haploid sperm antigens as foreign (since they arise after immune tolerance is established), and creates a specialised microenvironment for meiosis and spermiogenesis.

KEY POINTS Key points of this section

- The testes produce spermatozoa (seminiferous tubules) and testosterone (Leydig cells); the scrotum keeps them 2°C – 3°C below body temperature.
- Spermatogenesis: spermatogonia ($2n$) $\xrightarrow{\text{mitosis}}$ primary spermatocytes ($2n$) $\xrightarrow{\text{meiosis I}}$ secondary spermatocytes (n) $\xrightarrow{\text{meiosis II}}$ spermatids (n) $\xrightarrow{\text{spermiogenesis}}$ spermatozoa (n); continuous from puberty; 64–74 days per cycle; four sperm per primary spermatocyte.
- A spermatozoon = head (nucleus + acrosome) + neck (centrioles) + middle piece (mi-

tochondria) + tail (flagellum, 9+2 microtubules).

- Control: hypothalamic GnRH (pulsatile) → pituitary FSH (Sertoli cells, spermatogenesis support) and LH/ICSH (Leydig cells, testosterone); testosterone and inhibin exert negative feedback on hypothalamus/pituitary.

Having seen how the male gametes are produced and controlled, we now turn to the female partner: the structure of the female reproductive system, the production of ova, and the remarkable monthly cycle that prepares the body for a possible pregnancy.

5 THE FEMALE REPRODUCTIVE SYSTEM, OOGENESIS AND THE MENSTRUAL CYCLE


In the previous section we studied the male reproductive system and saw that spermatogenesis is a continuous process producing millions of small, motile gametes every day from puberty until old age. The female system tells a very different story. A baby girl is born with all the potential egg cells she will ever have, already paused part-way through meiosis; from puberty onwards, roughly one of them matures and is released each month in a beautifully orchestrated hormonal cycle. Understanding this cycle — the interplay between the brain, the pituitary gland and the ovaries — is one of the most frequently examined topics at GCE Advanced Level. We shall proceed from anatomy, to gamete production, to the ovarian and uterine cycles and their hormonal control, ending with the hormonal changes of pregnancy.

5.1 Organs of the Female Reproductive System

Observation. During a caesarean delivery in a Cameroonian hospital, the surgeon works through the abdomen to reach the uterus, a muscular organ normally the size of a pear. That such a small organ can shelter a baby of 3 kg or more tells us immediately that the female reproductive organs are specialised, elastic and under powerful hormonal control.

DEFINITION The female reproductive system

The **female reproductive system** is the set of organs that produce female gametes (ova), receive sperm, provide the site of fertilisation, and nourish and protect the developing embryo and foetus until birth.



Front view of the female reproductive system. The vulva (external genitalia: labia majora, labia minora, clitoris and the vaginal opening) is not shown in this internal view.

Structure and functions of each organ.

- **Ovaries** (one pair, almond-shaped, about 3 cm long): the female gonads. They contain thousands of follicles and have two functions: production of oocytes (oogenesis) and secretion of the hormones oestrogen and progesterone.
- **Fallopian tubes (oviducts)** (one pair, about 10 cm): muscular tubes lined with ciliated cells. Their funnel-shaped ends bear finger-like projections, the **fimbriae**, which sweep over the ovary at ovulation to capture the released oocyte. Fertilisation normally occurs in the upper third of the Fallopian tube.
- **Uterus**: a hollow, thick-walled muscular organ. Its wall has two layers of interest: the **myometrium** (smooth muscle which contracts during birth) and the **endometrium** (a glandular lining which thickens each month and into which the embryo implants).
- **Cervix**: the narrow, muscular neck of the uterus. It produces mucus whose consistency varies with the cycle (thin and watery at ovulation to allow sperm passage, thick at other times).
- **Vagina**: a muscular, elastic tube that receives the penis and semen during intercourse, serves as the passage for menstrual flow, and forms the birth canal.
- **Vulva**: the external genital organs (labia majora and minora, clitoris), which protect the vaginal opening and play a role in sexual sensation.

WARNING Common mistake

Do not confuse the urethra (part of the urinary system) with the vagina. In females, unlike males, the urinary and reproductive openings are separate. Also note: fertilisation occurs in the Fallopian tube, not in the uterus; the uterus is the site of implantation and development.

5.2 Oogenesis: Production of the Female Gamete

Observation. A woman releases roughly 400 oocytes in her whole reproductive life, while a man produces hundreds of millions of sperm per day. This enormous difference in numbers is matched by a difference in strategy: the female gamete is huge, packed with cytoplasm and food reserves, and produced in a slow, discontinuous, precisely timed process.

DEFINITION Oogenesis

Oogenesis is the process by which diploid germ cells in the ovary (oogonia) develop into a haploid female gamete. It begins before birth, is arrested for years, and is completed only if fertilisation occurs.


The stages of oogenesis, step by step.

1. **Before birth (foetal life).** Oogonia multiply by mitosis, then enlarge to become **primary oocytes**. Each primary oocyte begins meiosis I but is arrested in prophase I. At birth, the ovary already contains its full stock of primary oocytes, each surrounded by a layer of cells forming a **primordial follicle**.
2. **From puberty, each month.** Under FSH stimulation, several follicles start to grow, but usually only one becomes a mature **Graafian follicle**. Inside it, the primary oocyte completes **meiosis I** just before ovulation, producing a large **secondary oocyte** and a tiny **first polar body** (the cytoplasm is divided very unequally so that the future gamete keeps nearly all the reserves).
3. **Ovulation.** The cell released is the **secondary oocyte**, which has begun meiosis II but is arrested in metaphase II.
4. **Only if fertilisation occurs.** The entry of a sperm triggers completion of **meiosis II**, producing the mature **ovum** (whose nucleus fuses with the sperm nucleus) and a **second polar body**. The polar bodies degenerate.

WARNING The classic GCE trap

The cell released at ovulation is a secondary oocyte arrested in metaphase II, not a fully mature ovum. Meiosis II is only completed after a sperm penetrates it. Writing “a mature ovum is released at ovulation” costs marks.

SPERMATOGENESIS



OOGENESIS



Comparison of spermatogenesis and oogenesis. Note the unequal cytoplasmic divisions in oogenesis: one large ovum plus small polar bodies, versus four equal sperm.

Feature	Spermatogenesis	Oogenesis
Timing	Continuous from puberty to old age	Begins before birth; one oocyte per month from puberty to menopause
Meiosis	Completed without interruption	Arrested twice (prophase I, then metaphase II); completed only after fertilisation
Gametes per meiosis	4 functional spermatozoa	1 functional ovum + 2–3 polar bodies
Cytoplasm division	Equal	Very unequal (ovum keeps the reserves)
Gamete size/motility	Small, motile	Large, non-motile

5.3 The Ovarian and Uterine (Menstrual) Cycles

Observation. Most women of reproductive age experience menstrual bleeding roughly every 28 days. This visible event is only the tip of the iceberg: it is the external sign of two coordinated cycles — one in the ovary (preparing and releasing an oocyte) and one in the uterus (preparing a bed for a possible embryo).

DEFINITION Menstruation

Menstruation is the monthly shedding of the thickened endometrium (with blood and mucus) through the vagina, occurring when fertilisation has not taken place. It marks day 1 of a new cycle.

The **ovarian cycle** (average 28 days) has three key events:

1. **Follicular phase (days 1–13)**. Under FSH, several follicles grow; one becomes dominant and matures into a **Graafian follicle**, which secretes increasing amounts of **oestrogen**.
2. **Ovulation (about day 14)**. The Graafian follicle ruptures and releases the secondary oocyte.
3. **Luteal phase (days 15–28)**. The remains of the follicle transform into the **corpus luteum** (“yellow body”), which secretes **progesterone** (and some oestrogen). If there is no pregnancy, the corpus luteum degenerates after about 10–12 days.

The **uterine cycle** follows in parallel:

1. **Menstruation (days 1–5)**: the old endometrium breaks down and is shed.
2. **Proliferative (repair) phase**: under oestrogen from the growing follicles, the endometrium is rebuilt and its glands and blood vessels develop.
3. **Secretory phase**: under progesterone from the corpus luteum, the endometrium thickens further, becomes spongy and richly vascularised — ready for implantation.
4. **Breakdown**: if no embryo implants, the corpus luteum degenerates, progesterone falls, and the endometrium breaks down: a new menstruation begins.

DEFINITION Ovulation

Ovulation is the release of a secondary oocyte from a mature Graafian follicle at the surface of the ovary, triggered by the mid-cycle surge of LH. It occurs at about day 14 of a 28-day cycle.

5.4 Hormonal Control of the Cycle


The cycle is directed by a chain of command: **hypothalamus** → **pituitary** → **ovary** → **uterus**, with feedback loops returning information at every level.

1. The hypothalamus secretes **GnRH** (gonadotrophin-releasing hormone), which stimulates the anterior pituitary.
2. The pituitary releases **FSH**, which stimulates follicle growth. The growing follicles secrete **oestrogen**.
3. At low/moderate concentration, oestrogen inhibits FSH and LH (negative feedback), so only the dominant follicle survives.

- When oestrogen reaches a high, sustained concentration (mature Graafian follicle), it exerts **positive feedback**: a massive **LH surge** (with a smaller FSH peak) occurs, triggering **ovulation** about 24–36 hours later.
- LH converts the ruptured follicle into the **corpus luteum**, which secretes **progesterone** (plus oestrogen). These hormones inhibit FSH and LH (negative feedback), preventing new follicles from maturing.
- Without pregnancy, the corpus luteum degenerates; progesterone and oestrogen fall; the endometrium breaks down (menstruation), and the removal of inhibition allows FSH to rise again — a new cycle begins.

PROPERTY Positive feedback of oestrogen — the key exception

At **high, sustained concentration**, oestrogen switches from negative to positive feedback on the pituitary (and hypothalamus), provoking the LH surge that causes ovulation. This is the classic exception to the rule that reproductive hormones exert negative feedback, and examiners love to test it. (No proof is required; you must be able to explain it and locate it on a graph.)



Hormone levels across a 28-day cycle. The oestrogen peak (green) immediately precedes and triggers the LH surge (orange), which causes ovulation (dashed line). Progesterone (purple) dominates the luteal phase; its fall at day 26–28 causes menstruation.

METHOD Identifying the day of ovulation on a hormone graph

1. Locate the sharpest, tallest peak: this is the LH surge.
2. Check that it is immediately preceded by an oestrogen peak (positive feedback) and followed by a rise in progesterone (corpus luteum forming).
3. Ovulation occurs at, or about 24–36 hours after, the LH peak — typically day 14 of a 28-day cycle.
4. Justify in words: “the LH surge, triggered by high oestrogen, causes the Graafian follicle to rupture.”

WORKED EXAMPLE 6 Interpreting a cycle graph

On a graph of a 28-day cycle, LH rises sharply from day 12 to a maximum at day 13, then falls; progesterone begins to rise from day 15. (a) On which day does ovulation most probably occur? (b) Explain the hormonal cause of the LH peak. (c) Predict the consequence for the endometrium if progesterone falls sharply at day 26.

SOLUTION

(a) Ovulation occurs about 24–36 hours after the LH peak, so most probably on day 14. (b) The mature Graafian follicle has been secreting increasing amounts of oestrogen. When oestrogen reaches a high, sustained level, it exerts positive feedback on the pituitary (via the hypothalamus), provoking the sudden LH surge. (c) Progesterone maintains the thickened endometrium. A sharp fall of progesterone (due to degeneration of the corpus luteum in the absence of pregnancy) removes this support, so the endometrium breaks down and is shed: menstruation begins, marking day 1 of the next cycle.

5.5 Hormonal Changes During Pregnancy

Observation. If the cycle is a monthly rehearsal for pregnancy, what happens when the rehearsal succeeds? A woman who conceives does not menstruate — and the reason is entirely hormonal.

If fertilisation occurs, the young embryo implants in the endometrium around day 20–22 and its outer layer (the future placenta) secretes **hCG** (human chorionic gonadotrophin). hCG acts like LH: it **maintains the corpus luteum**, which therefore continues to secrete **progesterone and oestrogen**. These hormones keep the endometrium intact (no menstruation) and keep FSH and LH inhibited (no new ovulation). From about the third month, the **placenta takes over** the production of progesterone and oestrogen, and the corpus luteum is no longer needed. Progesterone thus remains high throughout pregnancy — this is the hormone detected indirectly by the absence of menstruation, and hCG itself is the hormone detected by pregnancy test kits.

WARNING GCE exam tip

A very frequent question: “Explain why progesterone levels remain high during pregnancy.” The full-mark answer has two stages: (1) the embryo secretes hCG, which maintains the corpus luteum so it keeps producing progesterone; (2) later, the placenta takes over progesterone (and oestrogen) production. Mentioning only one stage loses marks.

KEY POINTS Key points of this section

- The ovary produces oocytes and secretes oestrogen and progesterone; fertilisation occurs in the Fallopian tube; implantation occurs in the endometrium of the uterus.
- Oogenesis is discontinuous: primary oocytes are formed before birth; meiosis I is completed at ovulation (secondary oocyte + first polar body); meiosis II is completed only after fertilisation. One ovum per cycle.
- The cell released at ovulation is a secondary oocyte arrested in metaphase II.
- Cycle logic: FSH → follicle growth → oestrogen → (positive feedback) LH surge → ovulation → corpus luteum → progesterone → negative feedback; fall of progesterone → menstruation.
- In pregnancy, hCG maintains the corpus luteum, then the placenta takes over hormone production.

EXERCISE 5 Check your understanding

1. Name the organ where fertilisation normally occurs and the organ where implantation occurs.
2. State precisely the stage of meiosis of the cell released at ovulation, and explain when meiosis is completed.
3. Explain why only one functional ovum results from oogenesis while spermatogenesis yields four spermatozoa.
4. On a hormone graph, which peak signals that ovulation is imminent, and what causes it?
5. A woman's progesterone level remains high three weeks after ovulation. Suggest a biological explanation.

SOLUTION Exercise 1

1. Fertilisation: the Fallopian tube (oviduct). Implantation: the endometrium of the uterus.
2. A secondary oocyte arrested in metaphase II. Meiosis II is completed only if a sperm penetrates the oocyte (after fertilisation), producing the ovum and the second polar body.
3. In oogenesis the cytoplasm divides unequally: almost all cytoplasm and reserves go to one cell (the ovum), the rest forming tiny polar bodies that degenerate. In spermatogenesis divisions are equal, giving four functional sperm.
4. The LH surge. It is caused by positive feedback of the high, sustained oestrogen secreted by the mature Graafian follicle.
5. She is probably pregnant: the implanted embryo secretes hCG, which maintains the corpus luteum, so progesterone secretion continues instead of falling.

DID YOU KNOW? Did you know?

The hormone hCG is the basis of the pregnancy test kits sold in pharmacies across Cameroon: it is detectable in urine from about two weeks after fertilisation. Also, the “fertile window” used in natural family planning corresponds to the few days around ovulation, because the secondary oocyte survives only about 24 hours while sperm can survive up to about 3–5 days in the female tract.

We are now equipped to follow the gametes to their meeting point: in the next section we shall study fertilisation itself, the development of the embryo, and the process of birth.

6 FERTILIZATION, EMBRYONIC DEVELOPMENT AND BIRTH

In the previous section we followed the female gamete: we saw how oogenesis produces a secondary oocyte, how ovulation releases it into the Fallopian tube, and how the menstrual cycle prepares the endometrium to receive a possible embryo. All of this elaborate preparation has a single purpose: the meeting of one sperm and one oocyte. In this section we follow that meeting and its extraordinary consequences — fertilization, the nine months of development that transform a single cell into a baby, the drama of birth, and the production of milk that nourishes the newborn. This is one of the most frequently examined topics in the GCE Advanced Level paper, so master every stage.

6.1 From Copulation to the Meeting of Gametes

Copulation and ejaculation. During sexual intercourse (copulation), the erect penis is inserted into the vagina. Rhythmic stimulation triggers the **ejaculation** reflex: coordinated contractions of the epididymis, vas deferens, seminal vesicles and prostate expel semen into the urethra and out into the vagina, near the cervix. A single ejaculate (about $3\text{--}4\text{ cm}^3$) contains roughly 300 million spermatozoa, yet only a few hundred will ever reach the egg. Why this enormous wastage? The female tract is a hostile obstacle course.

The journey of sperm through the female tract. From the vagina, sperm must:

1. **Cross the cervix.** Around ovulation, under the influence of oestrogen, the cervical mucus becomes thin, watery and alkaline — channels form that guide sperm through. At other times of the cycle the mucus is thick and almost impassable. Millions of sperm are stopped here.
2. **Swim through the uterus,** helped by muscular contractions of the uterine wall and by the tail (flagellum) beat powered by mitochondria in the mid-piece.
3. **Enter a Fallopian tube.** Only the tube on the side that ovulated contains an egg. Sperm reaching the wrong tube are lost.
4. **Reach the ampulla** (the wider region of the tube), where the secondary oocyte, surrounded by follicle cells (the corona radiata), awaits — fertilization usually occurs here, within about 24 hours of ovulation.

Of the hundreds of millions deposited, fewer than 1 000 typically arrive near the egg, and only one will fertilize it.

Capacitation. Freshly ejaculated sperm are not yet able to fertilize. During their passage through the female tract (over several hours), secretions of the uterus and tube remove cholesterol and glycoproteins from the sperm surface membrane. This change, called **capacitation**, destabilises the membrane of the acrosome and makes the sperm capable of undergoing the acrosome reaction. You only need to know it briefly: capacitation is the final maturation of sperm inside the female tract.

6.2 Fertilization in the Fallopian Tube

DEFINITION Fertilization

Fertilization is the fusion of a haploid sperm nucleus with a haploid egg nucleus to form a diploid **zygote**. In humans it normally takes place in the ampulla of the Fallopian tube.

It proceeds in a precise sequence:

1. **Penetration of the corona radiata.** The capacitated sperm pushes between the follicle cells surrounding the oocyte, aided by enzymes.
2. **The acrosome reaction.** On contact with the **zona pellucida** (the glycoprotein coat of the egg), the acrosome — a cap-like vesicle at the sperm head — releases hydrolytic enzymes (e.g. hyaluronidase, acrosin). These digest a path through the zona pellucida.

3. **Fusion of membranes.** The membrane of the sperm head fuses with the egg membrane, and the sperm nucleus (and usually the whole sperm, minus the tail) enters the cytoplasm.
4. **The block to polyspermy.** Entry of the first sperm triggers the egg to release cortical granules whose contents alter the zona pellucida, hardening it (zona reaction). This **block to polyspermy** prevents any other sperm from entering — essential, because a zygote with three sets of chromosomes cannot develop normally.
5. **Completion of meiosis and fusion of nuclei.** The secondary oocyte, which was arrested in metaphase II, now completes meiosis II, producing the ovum and a second polar body. The haploid sperm nucleus ($n = 23$) and the haploid egg nucleus ($n = 23$) — now called pronuclei — fuse. **Diploidy is restored:** the zygote has $2n = 46$ chromosomes, half from each parent. Sex is determined at this instant: an X-bearing sperm gives a girl (XX), a Y-bearing sperm a boy (XY).

KEY POINTS Fertilization — the essentials

Site: ampulla of the Fallopian tube. Key events: acrosome reaction → penetration of zona pellucida → block to polyspermy → fusion of pronuclei → restoration of the diploid number ($2n = 46$).


6.3 Early Development: From Zygote to Implanted Blastocyst

The zygote does not wait. As it is swept along the Fallopian tube by cilia and muscular contractions (a journey of about 4–5 days), it begins to divide.

- **Cleavage** (days 1–3): repeated mitotic divisions without growth — the cells (blastomeres) become smaller at each division, so the overall size stays constant. The embryo passes through 2-cell, 4-cell, 8-cell stages.
- **Morula** (day 3–4): a solid ball of about 16–32 cells ("mulberry"), still enclosed in the zona pellucida. It enters the uterus.
- **Blastocyst** (day 5–6): fluid accumulates inside, forming a hollow ball. Two parts can now be distinguished: the **inner cell mass** (which will form the embryo proper) and the outer **trophoblast** (which will form the chorion and the fetal part of the placenta). The zona pellucida breaks down ("hatching").
- **Implantation** (day 6–7 after fertilization): the trophoblast secretes enzymes that digest a small area of endometrium, and the blastocyst burrows into it.

DEFINITION Implantation

Implantation is the embedding of the blastocyst in the endometrium (the vascularised lining) of the uterus. It occurs about 6–7 days after fertilization and marks the true beginning of pregnancy.



From zygote to implanted blastocyst. Cleavage divisions occur without growth; implantation takes place about day 6–7.

METHOD Sequencing exercise — from ejaculation to implantation

A classic GCE task: arrange scrambled events in order. Reason as follows.

1. Identify the starting point: ejaculation into the vagina.
2. Follow the sperm: cervix → uterus → Fallopian tube (capacitation occurs during this journey).
3. Fertilization events: acrosome reaction → penetration of zona pellucida → fusion of pronuclei (zygote).
4. Follow the embryo: cleavage → morula → blastocyst → implantation (day 6–7).
5. Check the logic: anything involving the uterus (morula arrival, implantation) must come after events in the tube.

6.4 Extra-embryonic Membranes and the Placenta

After implantation, the embryo develops four **extra-embryonic membranes** which protect and sustain it but form no part of the baby's body:

- **Amnion:** a fluid-filled sac enclosing the embryo. The amniotic fluid cushions the fetus against shocks, allows movement, and keeps a constant temperature. ("The waters" that break before birth.)
- **Chorion:** the outermost membrane, derived from the trophoblast. It develops finger-like chorionic villi that grow into the endometrium and form the fetal part of the placenta. It also secretes **hCG** (human chorionic gonadotrophin), the hormone detected by pregnancy tests, which maintains the corpus luteum during the first weeks.
- **Allantois:** a small outgrowth whose blood vessels become the blood vessels of the umbilical cord.
- **Yolk sac:** in humans it contains no yolk; it produces the first blood cells and the primordial germ cells.

The placenta. By about week 10–12, the **placenta** is fully formed from the chorionic villi (fetal contribution) and the endometrium (maternal contribution). The fetus is connected to it by the **umbilical cord**, containing two umbilical arteries (carrying deoxygenated blood and wastes from fetus to placenta) and one umbilical vein (carrying oxygenated, nutrient-rich blood back to the fetus). The placenta has three great functions:


1. **Exchange organ:** oxygen and nutrients (glucose, amino acids, fatty acids, vitamins, minerals) diffuse from maternal blood to fetal blood; carbon dioxide, urea and other wastes diffuse in the opposite direction. Some antibodies (IgG) also cross, giving the newborn passive immunity.
2. **Endocrine gland:** the placenta secretes hCG, then **progesterone** and **oestrogens** (taking over from the corpus luteum after about 3 months), maintaining the endometrium and suppressing further ovulation and menstruation; it also produces human placental lactogen.
3. **Selective barrier:** it blocks many (but not all) harmful agents — some viruses, the rubella virus, alcohol, nicotine and certain drugs can cross and damage the fetus.

PROPERTY The placental barrier — no mixing of bloods

Exchange across the placenta occurs by diffusion (and active transport for some substances) through the thin walls of the chorionic villi. Maternal and fetal blood do NOT mix: they remain separated by the villus epithelium, connective tissue and capillary endothelium. This is essential because mother and fetus may have different, incompatible blood groups (e.g. Rhesus). (Statement examinable; no proof required.)

WARNING Common mistake

Never write "the mother's blood flows into the baby" or "the two bloods mix in the placenta". Examiners penalise this heavily. Correct phrasing: substances diffuse between two separate circulations.



The fetus in utero: the amnion encloses the fetus in amniotic fluid; the umbilical cord links the fetus to the placenta attached to the uterus wall.

6.5 Gestation: Nine Months of Development

Gestation (pregnancy) lasts about **38–40 weeks** (roughly 9 months) from fertilization, conventionally divided into three trimesters.

- **First trimester (weeks 1–12):** the period of **organogenesis** — all the major organs are laid down. The heart begins beating around week 4; the neural tube, limbs, eyes and digestive system form. This is the most delicate period: the embryo is highly sensitive to alcohol, drugs and infections (e.g. rubella), which can cause malformations. By week 12 the fetus is recognisably human, about 7–8 cm long.
- **Second trimester (weeks 13–26):** mainly growth and maturation. Movements are felt by the mother ("quickening"); the fetus can suck its thumb; hair and nails appear.
- **Third trimester (weeks 27–40):** rapid growth and fat deposition; the lungs mature (surfactant production); the fetus usually turns head-down in preparation for birth.

DID YOU KNOW?

Pregnancy care in Cameroon

Because the first trimester is the period of organogenesis, health workers in Cameroonian antenatal clinics insist that expectant mothers avoid alcohol and self-medication, attend consultations early, and sleep under insecticide-treated mosquito nets: malaria infection during pregnancy reduces placental exchange and is a major cause of low birth weight. Protecting the placenta means protecting the baby.

6.6 Birth (Parturition)


DEFINITION Parturition

Parturition is the process of giving birth: the expulsion of the fetus and then of the placenta from the uterus, driven by powerful uterine contractions under the control of **oxytocin** and **prostaglandins**.

Hormonal trigger. As pregnancy reaches term, the fetal pituitary and the placenta shift the hormonal balance: oestrogen rises relative to progesterone, making the myometrium (uterine muscle) sensitive to oxytocin. The fetal head pressing on the cervix sends nerve impulses to the mother's hypothalamus, causing the posterior pituitary to release **oxytocin**. Oxytocin stimulates uterine contractions; the uterus and fetal membranes also release **prostaglandins**, which intensify contractions and soften (ripen) the cervix.

WARNING Oxytocin: a **POSITIVE** feedback — the classic trap

Most hormonal controls you have studied (e.g. thyroxine, blood glucose) are negative feedback. Oxytocin in labour is the famous exception: contractions push the baby's head against the cervix → more stretching → more oxytocin released → stronger contractions → and so on, until the baby is born and the stimulus stops. Do not describe it as negative feedback.



Positive feedback of oxytocin during labour: each contraction reinforces the next until delivery breaks the loop.

The three stages of labour:

1. **Dilation of the cervix** (longest stage, up to several hours): regular contractions dilate the cervix to about 10 cm; the amnion usually ruptures ("breaking of the waters").
2. **Expulsion of the baby**: powerful contractions, aided by the mother's abdominal muscles, push the baby head-first through the cervix and vagina. The umbilical cord is clamped and cut.
3. **Delivery of the placenta (afterbirth)**: about 10–30 minutes later, further contractions detach and expel the placenta and membranes. The uterus then contracts firmly, limiting bleeding.

6.7 Lactation: Feeding the Newborn

During pregnancy, high levels of oestrogen and progesterone develop the mammary glands but inhibit milk secretion. After birth, the fall in these hormones lifts the inhibition, and two pituitary hormones take over:

- **Prolactin** (anterior pituitary): stimulates the alveoli of the mammary glands to produce milk. Suckling maintains prolactin secretion — the more the baby suckles, the more milk is made.
- **Oxytocin** (posterior pituitary): released by the suckling reflex, it causes the myoepithelial cells around the alveoli to contract, ejecting milk into the ducts (the “let-down” reflex). Even the baby’s cry can trigger it.

The first secretion, **colostrum**, is a yellowish fluid rich in proteins and maternal antibodies but low in fat; it gives the newborn crucial passive immunity during its first days. Breastfeeding is strongly recommended in Cameroon and worldwide: breast milk is sterile, at the right temperature, perfectly balanced nutritionally, transmits antibodies, and is free — a decisive advantage where clean water for formula preparation is not guaranteed.

WORKED EXAMPLE 7 Worked exercise — functions of the placenta

Statement (GCE style). (a) Define implantation. (b) State four functions of the placenta, explaining briefly how exchange occurs. (c) Explain why a Rhesus-negative mother and a Rhesus-positive fetus can coexist during a first pregnancy.

SOLUTION

Solution. (a) Implantation is the embedding of the blastocyst in the endometrium of the uterus, about 6–7 days after fertilization. (b) Any four of: (i) diffusion of oxygen from maternal to fetal blood; (ii) diffusion of nutrients (glucose, amino acids) to the fetus; (iii) removal of carbon dioxide and urea from fetal blood; (iv) transfer of maternal antibodies (passive immunity); (v) secretion of hormones — hCG, progesterone and oestrogens — which maintain pregnancy. Exchange occurs by diffusion (and active transport for some substances) across the thin walls of the chorionic villi, which present a large surface area bathed in maternal blood. (c) Maternal and fetal blood do not mix in the placenta; they are separated by the villus tissues. During a first pregnancy the mother’s immune system therefore does not meet the fetal Rhesus antigens, so no anti-Rhesus antibodies are produced and the fetus is unharmed. (Sensitisation may occur at birth, when bloods can mix — a risk for later pregnancies.)

WARNING Exam tip (GCE)

When asked for the functions of the placenta, always give both categories: the nutritive/respiratory/excretory exchanges and the endocrine role (secretion of progesterone, oestrogens, hCG). Answers giving only “exchange of gases and food” earn only half the marks.

EXERCISE 6 Check your understanding

1. Place in correct order: morula; ejaculation; implantation; acrosome reaction; blastocyst; capacitation; fusion of pronuclei.
2. Explain why the block to polyspermy is essential for normal development.
3. Distinguish clearly between the roles of prolactin and oxytocin in lactation.
4. State the three stages of parturition and the hormone(s) responsible for the contractions.

SOLUTION Exercise n°1

1. Ejaculation → capacitation → acrosome reaction → fusion of pronuclei → morula → blastocyst → implantation.
2. If several sperm entered, the zygote would receive extra sets of chromosomes (triploidy or more); such abnormal chromosome numbers prevent normal development, so the embryo would die.
3. Prolactin (anterior pituitary) stimulates milk production by the alveoli; oxytocin (posterior pituitary) causes contraction of myoepithelial cells and milk ejection (let-down) in response to suckling.
4. (i) Dilation of the cervix; (ii) expulsion of the baby; (iii) delivery of the placenta (after-birth). Contractions are driven by oxytocin (positive feedback) and prostaglandins.

KEY POINTS Section summary

Fertilization occurs in the Fallopian tube: acrosome reaction, block to polyspermy, fusion of pronuclei restoring $2n = 46$. The zygote undergoes cleavage → morula → blastocyst, which implants in the endometrium around day 6–7. The extra-embryonic membranes (amnion, chorion, allantois, yolk sac) protect and nourish the embryo; the placenta exchanges gases, nutrients and wastes by diffusion — the two bloods never mix — and secretes hCG, progesterone and oestrogens. Gestation lasts 38–40 weeks, with organogenesis in the first trimester. Birth is driven by oxytocin (positive feedback) and prostaglandins through three stages: dilation, expulsion, afterbirth. Lactation depends on prolactin (production) and oxytocin (ejection); colostrum gives the newborn its first antibodies.

7 BIRTH CONTROL METHODS

Having studied the intricate processes of fertilisation, embryonic development and birth, we now turn to a crucial application of reproductive biology: the deliberate regulation of fertility. Understanding **contraception** is not merely a medical topic; it is a cornerstone of public health, family welfare and socio-economic planning, particularly in our Cameroonian context where maternal health and population dynamics are national priorities.

7.1 Concept and Rationale for Birth Control

DEFINITION Contraception

Contraception (or birth control) is the deliberate use of artificial or natural methods to prevent **fertilisation** (the fusion of sperm and ovum) or, in some cases, to prevent **implantation** of the blastocyst in the uterine endometrium. It allows individuals and couples to anticipate and attain their desired number of children and the spacing and timing of their births.

DID YOU KNOW? Why Birth Control Matters in Cameroon

In Cameroon, the rationale for birth control is multi-dimensional:


- **Maternal and Child Health:** Spacing pregnancies (at least 24 months) reduces the risk of maternal anaemia, haemorrhage, and low birth weight babies. It lowers maternal mortality (MMR) and infant mortality (IMR).
- **Family Welfare:** Planned families can invest more in nutrition, education (school fees, books) and healthcare per child.
- **Socio-economic Development:** A controlled population growth rate eases pressure on infrastructure (schools, hospitals, roads), food security and youth employment.
- **Women's Empowerment:** Control over reproduction enables girls to stay in school longer and women to participate fully in the workforce.

The National Population Policy of Cameroon explicitly promotes family planning as a right and a development tool.

7.2 Classification of Contraceptive Methods

Contraceptive methods are classified by their **mechanism of action** and **permanence**. The following chart summarises the landscape.

Classification of Contraceptive Methods



Classification of contraceptive methods by mechanism, showing site of action.

7.3 Natural Methods

These methods rely on observing physiological signs or behavioural discipline. They involve no devices or drugs.

EXPERIMENT

The Menstrual Cycle and the Fertile Window

Principle: Fertilisation can only occur if intercourse happens during the **fertile window** — approximately 5 days before ovulation (sperm survival) to 24 hours after ovulation (ovum lifespan). **Calendar (Rhythm) Method:** The woman records cycle lengths for 6–12 months.

- Shortest cycle - 18 = First fertile day.
- Longest cycle - 11 = Last fertile day.
- Abstain or use barrier during this window.

Basal Body Temperature (BBT): Progesterone raises body temperature by $\approx 0.3^{\circ}\text{C}$ after ovulation. Daily morning temperature charting confirms ovulation has occurred (retrospective). **Cervical Mucus (Billings/ovulation method):** Oestrogen → clear, stretchy, slippery mucus (spinnbarkeit) → fertile. Progesterone → thick, sticky, opaque → infertile. **Symptothermal Method:** Combines BBT + Mucus + Calendar calculation → higher reliability.

KEY POINTS Natural Methods — Key Points

- Abstinence: 100% effective if total; requires high motivation.
- Withdrawal (Coitus Interruptus): Penis withdrawn before ejaculation. Unreliable: Pre-ejaculatory fluid (Cowper's gland secretion) may contain viable sperm. High failure rate ($\approx 20\%$ typical use).
- Lactational Amenorrhea Method (LAM): Exclusive breastfeeding (on demand, day & night) $\rightarrow$ high prolactin $\rightarrow$ suppresses GnRH $\rightarrow$ no LH surge $\rightarrow$ no ovulation. Effective ($> 98\%$) only if: (1) Baby < 6 months, (2) Amenorrhoeic, (3) Exclusive breastfeeding.
- Advantages: No cost, no side effects, religiously acceptable.
- Disadvantages: High user failure rate; requires training, discipline, regular cycles; NO protection against STIs/HIV.

7.4 Barrier Methods

These physically block sperm from reaching the ovum.

DEFINITION Barrier Contraception

A **barrier method** is a device that creates a physical block preventing spermatozoa deposited in the vagina from entering the uterus and reaching the fallopian tubes.

Method	Mechanism / Site	Effectiveness (Typical Use)	STI Protection
Male Condom (Latex/Polyurethane)	Sheath over erect penis; collects semen.	$\approx 85\%$	YES (Best)
Female Condom (Polyurethane/Nitrile)	Loose sheath lining vagina; inner ring at cervix, outer ring at vulva.	$\approx 79\%$	YES
Diaphragm / Cervical Cap (+ Spermicide)	Dome/cap fits over cervix; spermicide immobilises sperm. Requires fitting by clinician.	$\approx 83\%$	Partial (Herpes/HPV limited)

WARNING Common GCE Trap

Only condoms (male and female) offer significant protection against Sexually Transmitted Infections (STIs) including HIV/AIDS. The Pill, IUD, Implants, Injectables, Diaphragm, Withdrawal DO NOT protect against STIs. This is a favourite distinction in GCE Paper 2 essay questions.


7.5 Hormonal Methods

These exploit the **negative feedback** control of the hypothalamic-pituitary-ovarian axis.

PROPERTY Mechanism of Hormonal Contraception

Exogenous **oestrogen** and/or **progestin** (synthetic progesterone) suppress the secretion of **GnRH** from the hypothalamus and **FSH/LH** from the anterior pituitary.

- Oestrogen component: Inhibits FSH → prevents follicular development → no ovulation.
- Progestin component: Inhibits LH surge → no ovulation; thickens cervical mucus → blocks sperm penetration; thins endometrium → unfavourable for implantation.

Negative Feedback Mechanism of the Combined Oral Contraceptive Pill

The pill maintains high steroid levels, mimicking the luteal phase, suppressing GnRH and the LH surge, thereby preventing ovulation.

Method	Composition / Regimen	Key Features / Side Effects
Combined Oral Contraceptive Pill (COC)	Oestrogen (Ethinylestradiol) + Progestin. 21 days on, 7 days off (withdrawal bleed).	High efficacy ($\approx 91\%$ typical). Regulates cycles, reduces dysmenorrhoea, acne. Risks: Thromboembolism (DVT/PE), hypertension, contraindicated in smokers > 35 yrs, breastfeeding mums (reduces milk).
Progestin-Only Pill (POP / Mini-Pill)	Progestin only (e.g. Desogestrel). Taken continuously, same time daily (± 3 hrs).	Safe for breastfeeding, no oestrogen risks. Mainly thickens mucus; may suppress ovulation. Irregular bleeding common.
Injectables (e.g. Depo-Provera / Sayana Press)	DMPA (Depot Medroxyprogesterone Acetate) IM/SC every 12–13 weeks.	High efficacy ($\approx 94\%$). Amenorrhoea common. Delay in return to fertility (avg 9–10 months). Bone density loss with long use.
Implants (e.g. Jadelle 2-rods, Implanon 1-rod)	Levonorgestrel / Etonogestrel rods under skin of upper arm. 3–5 years.	Highest efficacy ($\approx 99.9\%$). Long-acting, reversible (LARC). Insertion/removal minor surgery. Irregular bleeding.
Emergency Contraceptive Pill (ECP)	Levonorgestrel 1.5mg (single dose) or Ulipristal Acetate. Within 72–120h post coitus.	Delays/blocks ovulation. NOT an abortifacient (does not disrupt implanted pregnancy). Not for regular use.

7.6 Intrauterine Devices (IUD / IUCD)

DEFINITION Intrauterine Device

An **IUD** is a small, T-shaped device inserted into the uterine cavity by a trained provider. It provides long-acting reversible contraception (LARC).

Type	Mechanism	Duration / Notes
Copper IUD (Cu-T 380A)	Copper ions → toxic to sperm (motility/viability) and ova; sterile inflammatory reaction in endometrium → prevents fertilisation & implantation.	10–12 years. Non-hormonal. Can be used as emergency contraception (inserted ≤ 5 days post coitus). Heavier, more painful periods initially.
Hormonal IUS (Levonorgestrel-IUS, e.g. Mirena)	Releases levonorgestrel locally → thickens cervical mucus, thins endometrium (often amenorrhoea), may suppress ovulation.	5–7 years. Treats heavy menstrual bleeding (menorrhagia). Lower systemic hormones.

WARNING IUD Misconceptions

- **Mechanism:** Primary action is prevention of fertilisation (spermicidal/ovicidal), not abortion. It does not disrupt an established pregnancy.
- **Infection Risk:** Slightly increased risk of PID (Pelvic Inflammatory Disease) only in first 20 days post-insertion (aseptic technique crucial). Not contraindicated in nulliparous women (modern guidelines).
- **Expulsion:** Check strings monthly after menses.

7.7 Surgical (Permanent) Methods

These are **sterilisation** procedures intended to be irreversible. Counselling is mandatory.

METHOD Vasectomy (Male Sterilisation)

1. Under local anaesthesia, a small incision or puncture is made in the scrotum.
2. The **vas deferens** (sperm duct) is isolated, a segment excised, and ends ligated/sealed (cautery/clips).
3. Not immediately effective: Residual sperm proximal to cut. Must use backup contraception until azoospermia confirmed by semen analysis (usually $\approx$ 20 ejaculates / 3 months).
4. Testosterone production (Leydig cells) and sexual function unchanged.

METHOD Tubal Ligation (Female Sterilisation)

1. Usually laparoscopic (minilaparotomy post-partum).
2. **Fallopian tubes** are occluded: clips (Filshie), rings, coagulation, or excision (Pomeroy).
3. Effective immediately.
4. Ovarian function (hormones, ovulation) preserved; ovum degenerates in peritoneal cavity.
5. Higher surgical risk than vasectomy (anaesthesia, bowel/bladder injury).

KEY POINTS Sterilisation — Key Exam Points

- Considered permanent. Reversal (vasovasostomy, tubal reanastomosis) is microsurgery, expensive, not always successful, and not guaranteed by health systems.
- No protection against STIs.
- Ethical/Legal: In Cameroon, requires informed written consent, minimum age (often 21 or 2 children), spousal consent often requested (though legally debated), cooling-off period.

7.8 Comparative Evaluation and Exam Strategy

METHOD Evaluating a Contraceptive Method — Three Criteria

For GCE essays/tables, evaluate every method against these three pillars:

1. Effectiveness (Pearl Index): **Perfect Use** (method efficacy) vs **Typical Use** (real world). LARC (Implants, IUD, Sterilisation) $\approx$ Perfect Use. User-dependent methods (Pills, Condoms, Natural) have wide gap.
2. Reversibility: How quickly does fertility return?
 - Immediate: Condoms, Diaphragm, IUD removal, Implant removal, Pills (usually 1–3 cycles).
 - Delayed: Injectables (months).
 - Permanent/Difficult: Sterilisation.
3. STI/HIV Protection: Only Male/Female Condoms.

WORKED EXAMPLE 8 GCE Style Question

Question: A newly married couple in Buea wants to delay their first pregnancy for 3 years. The husband travels frequently. The wife has a history of migraine with aura. They want a method that is highly effective, reversible, and does not require daily action. Recommend the most suitable method and justify your choice by comparing it with two other methods.

Solution: Recommendation: Subdermal Implant (e.g. Jadelle / Implanon NXT).

Justification (Comparison Table):

Criterion	Subdermal Implant	Combined Oral Pill (COC)	Copper IUD
Effectiveness	> 99.9% (LARC, user-independent)	≈ 91% (User-dependent, missed pills)	> 99% (LARC)
Duration / Action	3–5 years; insert & forget	Daily pill at same time	10–12 years; insert & forget
Migraine with Aura	SAFE (Progestin only)	CONTRAINDICATED (Oestrogen ↑ stroke risk)	SAFE (Non-hormonal)
Husband Travel	No adherence needed during absence	Requires daily discipline; missed pills on travel	No adherence needed
Reversibility	Rapid return (weeks)	Rapid return (1–3 cycles)	Rapid return (immediate)
STI Protection	No	No	No
Side Effects	Irregular bleeding (manageable)	Nausea, weight, thromboembolism risk	Heavier/painful periods

Conclusion: The Implant is optimal: safe for her medical history, unaffected by husband's travel, highly effective for the 3-year span, and easily removable when they desire pregnancy. The IUD is a strong second choice but may worsen dysmenorrhoea; the COC is medically unsafe.

7.9 Ethical, Religious and Health Considerations

DID YOU KNOW? Discussion Points for GCE Paper 2 Essays

- **Religious Views:** Catholic Church teaches against artificial contraception (*Humanae Vitae*), promoting Natural Family Planning (NFP). Many Protestant and Muslim leaders support spacing for maternal health but may oppose sterilisation/abortifacients.
- **Adolescent Access:** Cameroon law allows adolescents access to reproductive health services. Barrier: provider bias, stigma, cost. School-based programmes vs parental consent debates.
- **Gender Burden:** Most methods (Pill, Injectables, Implants, IUD, Tubal ligation) are female-controlled. Vasectomy and Male Condom are only male methods. Promoting shared responsibility is a public health goal.

- **Abortion vs Contraception:** Emergency Contraception (ECP) prevents ovulation; it is not medical abortion (Mifepristone/Misoprostol). Clarifying this distinction reduces stigma and increases ECP uptake.
- **Health System Integration:** Contraception integrated into ANC (post-partum), PNC, HIV clinics (dual protection), youth corners. Task-shifting (nurses inserting implants/IUDs) expands access.

EXERCISE 7 Practice Questions

- Define contraception and distinguish between contraception and contragestion.
- Explain the physiological basis of the Lactational Amenorrhea Method (LAM). State the three criteria for its effectiveness.
- Describe the mechanism of action of the Combined Oral Contraceptive Pill with reference to negative feedback.
- Compare and contrast the Copper IUD and the Levonorgestrel IUS (Hormonal IUD) in terms of mechanism, duration, and non-contraceptive benefits.
- A 35-year-old woman, smoker (15 cigarettes/day), requests the Combined Pill. Advise her with reasons.
- Outline the procedure for Vasectomy. Why is it not immediately effective?
- Discuss the statement: "The most effective contraceptive methods are those that do not depend on user compliance."

SOLUTION Exercise 1

Contraception is the deliberate prevention of fertilisation or implantation. Contragestion (interception) refers to agents acting after fertilisation but before implantation (e.g. high-dose progestins, IUD insertion post-coitus) to prevent implantation. The distinction is subtle; most modern definitions classify IUD/ECP as contraception (primary action pre-fertilisation).

SOLUTION Exercise 2

LAM: Suckling → Prolactin ↑ → GnRH pulse frequency ↓ → LH surge blocked → No ovulation. Criteria: (1) Infant < 6 months, (2) Amenorrhoea (no menses), (3) Exclusive/near-exclusive breastfeeding on demand (day & night).

SOLUTION Exercise 3

COC provides constant exogenous oestrogen/progestin → Negative feedback on Hypothalamus (GnRH ↓) and Ant. Pituitary (FSH/LH ↓) → No follicular recruitment (FSH low) → No LH surge → Anovulation. Progestin also thickens cervical mucus (sperm block) and atrophies endometrium (implantation block).

SOLUTION Exercise 4

Cu-IUD: Copper toxic to gametes, sterile inflammation → prevents fertilisation. 10+ yrs. Non-hormonal. Emergency contraception. Side effect: heavier menses. LNG-IUS: Local progestin → thick mucus, thin endometrium, maybe anovulation. 5–7 yrs. Treats menorrhagia, dysmenorrhoea, endometriosis. Side effect: irregular spotting → amenorrhoea.

SOLUTION Exercise 5

Advise against COC. Oestrogen + Smoking + Age > 35 = High risk of cardiovascular events (Venous Thromboembolism VTE, Stroke, MI). Absolute contraindication (UKMEC Category 4). Offer Progestin-only methods (POP, Implant, Injectables, LNG-IUS) or Copper IUD.

SOLUTION Exercise 6

Local anaesthesia → access vas deferens via scrotal incision/puncture → excise segment → ligate/cauterise ends. Not immediately effective because sperm remain in distal vas deferens/ampulla. Requires ≈ 20 ejaculates / 3 months + semen analysis confirming azoospermia before relying on it.

SOLUTION Exercise 7

Agree. LARC methods (Implants, IUDs, Sterilisation) have Pearl Indices ≈ 0.1 – 0.8 (Perfect ≈ Typical). User-dependent methods (Pills, Condoms, Natural) have Typical Use failure rates 7–24% due to human error (forgetting, incorrect use, breakage). LARC removes "user failure". However, "best" method includes acceptability, side effects, STI protection (condoms), and medical eligibility.

8 LIFE CYCLES: ALTERNATION OF GENERATIONS AND METAGENESIS

In the previous section we examined how human beings control their own reproduction. But reproduction is not always a simple matter of two parents producing offspring of the same kind. Walk through the humid forest on the slopes of Mount Cameroon after the rains: the green carpet of mosses and the delicate fronds of ferns are reproducing too — yet they do it through a remarkable strategy in which two different kinds of plant, one haploid and one diploid, take turns to produce each other. Even some animals, such as jellyfish, alternate between two body forms across their life cycle. Understanding these strategies is essential for the GCE, where life-cycle diagrams with n and $2n$ labels are a favourite examination topic.

8.1 The Concept of a Life Cycle: Haploid and Diploid Phases

DEFINITION Life cycle

The **life cycle** of an organism is the complete sequence of stages through which an individual passes, from its formation (by fertilization or by asexual reproduction) until it in turn produces the next generation.

Every sexually reproducing organism must solve a chromosome problem. If gametes contained the full diploid number of chromosomes ($2n$), the chromosome number would double at every generation. The solution is a rhythm of two key events:

- **Meiosis** halves the chromosome number: a diploid cell ($2n$) produces haploid cells (n).
- **Fertilization** (syngamy) restores the diploid number: two haploid gametes ($n + n$) fuse to form a diploid zygote ($2n$).

In animals like ourselves, meiosis directly produces the gametes, and the only haploid cells in the whole life cycle are the sperm and the egg. Plants (and some algae) do something different and far more elaborate: meiosis produces not gametes but **spores**, and each spore grows by mitosis into a whole haploid organism which then produces gametes by mitosis. This creates two alternating multicellular generations.

KEY POINTS The three key rules of any life cycle

- Rule 1: Meiosis is the only event that changes $2n \rightarrow n$.
- Rule 2: Fertilization is the only event that changes $n \rightarrow 2n$.
- Rule 3: Everything else (growth of gametophyte or sporophyte, gamete formation, budding, larval development) occurs by mitosis, which conserves the chromosome number.

8.2 Alternation of Generations in Plants

DEFINITION Alternation of generations

Alternation of generations is the regular alternation, in the life cycle of a plant, between a haploid (n) gamete-producing generation called the **gametophyte** and a diploid ($2n$) spore-producing generation called the **sporophyte**. The sporophyte produces haploid spores by meiosis; each spore germinates into a gametophyte, which produces gametes by mitosis; fusion of gametes restores the diploid sporophyte.

Notice the logic: the sporophyte is named after what it makes (spores, by meiosis), and the gametophyte after what it makes (gametes, by mitosis). The two generations differ in ploidy, in appearance, and often in which one is dominant.

The life cycle of a moss (e.g. *Funaria*)


The green, leafy moss plant you see on damp walls and stones in Buea during the rainy season is the **gametophyte** (n) — it is the dominant, photosynthetic, independent generation. It develops from a filamentous **protonema** that germinates from the spore. At its tip it bears the sex organs:

- **Antheridia** (male organs) release many motile, biflagellate sperm cells.
- **Archegonia** (female organs), flask-shaped, each contain a single egg cell.

Fertilization **requires liquid water**: the sperm must swim through a film of rain or dew to reach the egg inside the archegonium. This dependence on water is why mosses are restricted to moist habitats. The resulting zygote ($2n$) is not released; it grows in place, on top of the gametophyte, into the **sporophyte**: a thin stalk (seta) carrying a capsule. The sporophyte is **dependent** — it has no roots of its own and draws water and nutrients from the gametophyte. Inside the capsule, spore mother cells undergo **meiosis** to produce haploid spores, which are scattered by wind and germinate into a new protonema and then a green gametophyte.

The life cycle of a fern (e.g. *Dryopteris*)

In ferns the roles are reversed. The familiar fern plant with large divided fronds is the **sporophyte** ($2n$) — dominant, independent, with true roots, stems and leaves. On the underside of the fronds you can see brown dots called **sori** (singular: sorus); each sorus is a cluster of **sporangia** in which meiosis produces haploid **spores**. A spore that lands on moist soil germinates into the gametophyte: a small, flat, heart-shaped green plate called the **prothallus**, only a few millimetres across. The prothallus is independent and photosynthetic, but short-lived and delicate. It bears antheridia and archegonia on its lower surface; sperm swim through a water film to the egg; the zygote ($2n$) grows on the prothallus into a young fern, and the prothallus withers away.



Life cycle of a fern (*Dryopteris*). The diploid sporophyte is the dominant, conspicuous plant. Meiosis in sporangia (within sori) produces haploid spores. These germinate into the small, independent haploid prothallus (gametophyte) which produces gametes by mitosis. Fertilization requires water for sperm to swim to the egg.

PROPERTY Dominance of the generations — GCE comparison

In mosses (bryophytes), the **gametophyte** (n) is the dominant, independent, photosynthetic generation, while the sporophyte is small, short-lived and nutritionally dependent on the gametophyte. In ferns (pteridophytes), the **sporophyte** ($2n$) is dominant, independent and long-lived, while the gametophyte (prothallus) is small, short-lived but still independent and free-living. (This paired comparison is frequently examined; learn it as a set of contrasting statements.)

8.3 The Evolutionary Trend: From Mosses to Flowering Plants

Comparing mosses, ferns and flowering plants reveals one of the clearest trends in plant evolution:

Group	Dominant generation	Gametophyte	Water needed for fertilization?
Mosses (Bryophytes)	Gametophyte (n)	Large, independent (the leafy plant + protonema)	Yes — flagellated sperm swim
Ferns (Pteridophytes)	Sporophyte ($2n$)	Small prothallus, independent, free-living	Yes — flagellated sperm swim
Flowering plants (Angiosperms)	Sporophyte ($2n$)	Extremely reduced: pollen grain (male) and embryo sac (female), both dependent on the sporophyte	No — pollen tube delivers male nuclei

KEY POINTS The trend to remember for GCE

From mosses to flowering plants, the sporophyte becomes increasingly dominant while the gametophyte becomes increasingly reduced and dependent. In angiosperms, the male gametophyte is reduced to the **pollen grain** (and the pollen tube it produces), and the female gametophyte to the **embryo sac** inside the ovule. The conquest of dry land was possible because pollination by wind or insects replaced the need for liquid water at fertilization.

DID YOU KNOW? Why this matters in Cameroon

The mist forests of Mount Cameroon and the Bakossi Mountains are famous for their richness in ferns and mosses. These plants thrive there precisely because the permanent humidity provides the water film their swimming sperm need — a direct, visible consequence of alternation of generations. Flowering plants, freed from this constraint by the pollen tube, dominate the drier savanna regions of the North.

8.4 Metagenesis in Animals

Some animals also alternate between two generations, but of a very different nature. Consider the colonial cnidarian *Obelia*, a relative of jellyfish.


DEFINITION Metagenesis

Metagenesis is the alternation, in the life cycle of certain animals (especially cnidarians),

between an asexual generation (typically a sessile **polyp**) and a sexual generation (typically a free-swimming **medusa**). Both generations are diploid; there is no change in ploidy between them.

In *Obelia*, the life cycle runs as follows:

1. The **polyp colony** ($2n$), fixed to rocks or seaweed, grows and produces new individuals by **budding** — asexual reproduction (mitosis). Specialised reproductive buds (blastostyles) produce **medusae** by budding, which detach and swim away.
2. The free-swimming **medusa** ($2n$) is the sexual stage: it produces gametes (n) by meiosis in its gonads (located on the manubrium).
3. **Fertilization** of the egg produces a zygote ($2n$), which develops into a ciliated, free-swimming larva called the **planula**.
4. The planula settles on a solid surface and grows, by budding (mitosis), into a new polyp colony — completing the cycle.



Life cycle of *Obelia*, illustrating metagenesis: a sessile asexual polyp colony buds off sexual medusae; gametes fuse to give a zygote, which develops into a planula larva that settles and forms a new colony. Note that polyp, medusa, zygote and planula are all diploid ($2n$). Only the gametes are haploid (n).

WARNING Do not confuse metagenesis with alternation of generations — GCE TRAP

This is one of the most frequent errors at the GCE. In alternation of generations (plants), the two generations differ in ploidy: haploid gametophyte versus diploid sporophyte. In metagenesis (animals), the two generations (polyp and medusa) are both diploid; they differ only in body form and in the mode of reproduction (asexual budding versus sexual reproduction). The only haploid cells in metagenesis are the gametes, as in any animal. Writing “the medusa is haploid” or “meiosis produces the polyp” will cost you marks.

Feature	Alternation of Generations (Plants)	Metagenesis (Animals, e.g. Cnidarians)
Ploidy of the two generations	Haploid (n) gametophyte ↔ Diploid ($2n$) sporophyte	Both generations diploid ($2n$)
Chromosome number change	Yes: Meiosis ($2n \rightarrow n$) and Fertilization ($n \rightarrow 2n$)	Only at gamete formation (meiosis) and fertilization
Nature of generations	Morphologically distinct, often one dominant	Morphologically distinct (polyp vs medusa)
Reproductive mode of each	Gametophyte: sexual (gametes); Sporophyte: asexual (spores)	Polyp: asexual (budding); Medusa: sexual (gametes)
Examples	Mosses, ferns, flowering plants	Obelia, Aurelia (jellyfish), some flatworms

8.5 Biological Significance of These Strategies

Why should organisms bother with two generations at all? Both strategies confer real advantages:

- **Dispersal.** In plants, light spores (and later pollen and seeds) scatter the species far from the parent. In Obelia, the sessile colony cannot move, but the swimming medusa and planula larva carry the species to new substrates.
- **Multiplication.** The asexual phase (sporophyte producing many spores by meiosis; polyp budding many medusae by mitosis) multiplies the output of a single successful fertilization many times over.
- **Survival in varying conditions.** Each generation is adapted to a different way of life: the resistant sporophyte of the fern withstands dry air, while the delicate prothallus exploits moist microhabitats; the polyp resists and colonises, the medusa disperses.
- **Genetic mixing.** The sexual phase maintains genetic recombination, providing the variation on which natural selection acts.

WORKED EXAMPLE 9 Analysing a life-cycle diagram (GCE style)

A diagram of the fern life cycle shows the following structures: (A) frond with sori, (B) spore, (C) prothallus, (D) egg and sperm, (E) zygote. (1) State the ploidy (n or $2n$) of each structure. (2) Indicate between which structures meiosis and fertilization occur. (3) Name the dominant generation.

SOLUTION

Solution.

1. (A) Frond with sori: $2n$ (it belongs to the sporophyte). (B) Spore: n (product of meiosis). (C) Prothallus: n (gametophyte, grown by mitosis from the spore). (D) Egg and sperm: n (gametes, produced by mitosis on the gametophyte). (E) Zygote: $2n$ (product of fertilization).
2. Meiosis occurs in the sporangia of the sori, i.e. in the transition from (A) to (B) ($2n \rightarrow n$). Fertilization occurs when sperm and egg fuse, i.e. from (D) to (E) ($n + n \rightarrow 2n$).
3. The dominant generation is the sporophyte (the fern plant bearing the fronds), which is independent and photosynthetic; the prothallus is small and short-lived.

Marking tip: one mark per correct ploidy, one for each correctly placed event, one for the dominant generation — an easy six marks if the three key rules are applied methodically.

METHOD How to analyse any life-cycle diagram in the exam

1. Locate meiosis and fertilization first — they are the only two points where ploidy changes.
2. Label everything between fertilization and meiosis as $2n$ (sporophyte side in plants; polyp/medusa/zygote/planula side in metagenesis); label everything between meiosis and fertilization as n (gametophyte side in plants; gametes only in metagenesis).
3. Identify which generation is dominant (largest, longest-lived, independent) — for plants only.
4. For animals, check: if both generations are diploid and differ only in reproductive mode, it is metagenesis, not alternation of generations.

EXERCISE 8 Checking your understanding

1. Define alternation of generations and state how it differs from metagenesis.

2. In the moss *Funaria*, state the ploidy of: the leafy plant, the capsule, the spore, the sperm, the protonema.
3. Explain why mosses and ferns are restricted to moist habitats while flowering plants are not.
4. Draw and fully label the life cycle of *Obelia*, marking the points of meiosis and fertilization, and indicating the ploidy of each stage.
5. A student claims: "In the fern life cycle, the prothallus is diploid because it grows from the zygote." Explain why this statement is incorrect.

SOLUTION Exercise 1

1. Alternation of generations: regular alternation of a haploid gametophyte (produces gametes by mitosis) and a diploid sporophyte (produces spores by meiosis) in a plant life cycle. Metagenesis (animals) alternates an asexual polyp and a sexual medusa, both diploid — no change of ploidy between the two generations; meiosis occurs only during gamete formation in the medusa.
2. Leafy plant (gametophyte): n ; capsule (part of the sporophyte): $2n$; spore: n ; sperm: n ; protonema: n (grows from spore by mitosis).
3. In mosses and ferns the sperm are flagellated and must swim through liquid water to reach the egg, so reproduction is only possible in moist conditions. In flowering plants the pollen tube carries the male gametes to the embryo sac, making fertilization independent of external water.
4. The diagram must show: polyp colony ($2n$) $\xrightarrow{\text{budding (mitosis)}}$ medusa ($2n$) $\xrightarrow{\text{meiosis}}$ gametes (n) $\xrightarrow{\text{fertilization}}$ zygote ($2n$) $\xrightarrow{\text{cleavage}}$ planula ($2n$) $\xrightarrow{\text{settling \& growth}}$ new polyp colony. Meiosis is placed at gamete formation in the medusa; fertilization at the fusion of gametes.
5. The prothallus grows from the spore (which is n , produced by meiosis), not from the zygote. The zygote grows into the young sporophyte ($2n$). The student has confused the direction of the cycle.

KEY POINTS Summary of the section — GCE Essentials

- A life cycle alternates haploid (n) and diploid ($2n$) phases; only meiosis and fertilization change the ploidy; all other growth is by mitosis.
- Alternation of generations (plants): haploid gametophyte $\rightleftharpoons$ diploid sporophyte.

Mosses: gametophyte dominant, sporophyte dependent. Ferns: sporophyte dominant, prothallus small but independent.

- Evolutionary trend: increasing dominance of the sporophyte and reduction of the gametophyte, ending in the pollen grain and embryo sac of flowering plants.
- Metagenesis (animals, e.g. Obelia): asexual polyp $\rightleftharpoons$ sexual medusa, both diploid — do not confuse with alternation of generations.
- Both strategies improve dispersal, multiplication and survival.
- Exam technique: On any life-cycle diagram, first mark meiosis ($2n \rightarrow n$) and fertilization ($n \rightarrow 2n$), then deduce ploidy of all other stages.

9 GROWTH: CURVES, PATTERNS AND HORMONAL CONTROL

In the previous section we followed organisms through their **life cycles**, seeing how a fern alternates between sporophyte and gametophyte, and how a jellyfish passes through a polyp stage. Every stage of every life cycle has one thing in common: the organism must grow to reach it. A maize seedling in a field near Bafoussam becomes a two-metre plant in a few months; a newborn baby in Yaoundé becomes an adult of perhaps 170 cm in about eighteen years; a caterpillar becomes a butterfly only after a series of moults. How do we measure this growth, what shape does it follow, and what controls it? These are the questions of this final section.

9.1 What is Growth?

Intuition. If you weigh a baby every month, the mass increases. But if you soak a dry sponge in water, its mass also increases — yet the sponge has not grown. True growth is not just “getting bigger”; it is a permanent increase in the amount of living material, built from new cells and new cytoplasm.

DEFINITION Growth

Growth is the permanent and irreversible increase in size and dry mass of an organism (or of one of its parts), resulting from three cellular processes:

- **cell division** (mitosis), which increases the number of cells;
- **cell enlargement** (expansion), which increases the size of cells — in plants mainly by uptake of water into the vacuole (vacuolation);
- **cell differentiation**, by which cells become specialised in structure for particular functions.

WARNING Growth is not just swelling

A seed soaked in water increases in mass and volume, but this is imbibition, a reversible physical uptake of water — not growth. Growth implies the synthesis of new protoplasm and is irreversible: an adult who loses water through sweating has not “un-grown”.

9.2 Measuring Growth

To study growth scientifically we must measure a chosen parameter repeatedly over time. Three measures are commonly used.

Measure	Advantages	Limitations
Length / height	Quick, cheap, non-destructive; ideal for seedlings and children	One dimension only; ignores changes in width and mass
Fresh mass	Easy (weigh the organism); non-destructive	Fluctuates with water content (meals, rain, transpiration); unreliable
Dry mass	The true measure of growth: mass of organic matter actually synthesised	Destructive (the organism must be killed and dried in an oven at about 100 °C to constant mass); needs several identical samples

WARNING Fresh mass versus dry mass — the classic trap

A very common examination error is to use fresh mass as evidence of growth. Fresh mass rises after drinking or rainfall and falls during drought, without any change in the amount of living tissue. Dry mass is the only reliable measure of true growth, because it measures the organic material actually built by the organism.

METHOD Plotting and interpreting a growth curve


To obtain and analyse a growth curve (e.g. height of maize seedlings):

1. Germinate a large batch of genetically similar seeds under identical conditions.
2. At regular intervals (e.g. every 3 days), measure the chosen parameter (height, or dry mass of a sample of seedlings).
3. Plot the parameter (y-axis) against time (x-axis).
4. Identify the phases: slow start, rapid middle, levelling off.
5. Calculate the **growth rate** at any time as the gradient of the curve: $\text{rate} = \frac{\Delta \text{size}}{\Delta t}$. The

steepest part of the curve corresponds to the maximum growth rate.

9.3 The Sigmoid Growth Curve

When the size of an organism (a maize plant, a yeast population, a child measured over years) is plotted against time, the curve obtained is usually **S-shaped**, called a **sigmoid curve**. It shows four phases.



The sigmoid (S-shaped) growth curve with its four phases.

KEY POINTS Phases of the sigmoid curve

- Lag phase: slow growth; the organism is adjusting, cells are small and few, enzymes and organelles are being built up.
- Exponential (log) phase: growth rate is maximal; cells divide rapidly and resources are abundant. Size increases by a constant factor per unit time.
- Decelerating phase: growth slows as limiting factors appear (nutrients, space, accumulation of waste; in animals, the approach of adult size).
- Stationary phase: growth stops; in plants the organs are mature, in animals adult size is reached and cell production only balances cell loss.

The whole period from the start of growth to the stationary phase is called the **grand period of growth**.


9.4 Intermittent Growth: the Stepped Curve of Arthropods

Not all organisms grow smoothly. A grasshopper or a cockroach is enclosed in a rigid **exoskeleton** (cuticle) made of chitin, which cannot stretch. Such an animal can only increase in size by shedding its cuticle at intervals.

DEFINITION Ecdysis

Ecdysis (moulting) is the periodic shedding of the rigid exoskeleton of an arthropod, allowing a rapid increase in body size before the new, larger cuticle hardens. The stage between two successive moults is called an **instar**.

Consequently, the growth curve of an insect is **stepped (intermittent)**: long plateaux where length hardly changes (the cuticle restricts growth), separated by sudden vertical jumps at each moult, when the soft new cuticle allows rapid expansion by uptake of air or water.



Stepped growth curve of an insect: each vertical jump corresponds to a moult (ecdysis); the horizontal plateaux are the instars.


WARNING Exam tip — stepped versus smooth curves

A frequent GCE question asks: “Explain why the growth curve of an insect is stepped while that of a human is smooth.” The key idea: the insect’s rigid, inextensible exoskeleton prevents continuous expansion, so size can only increase suddenly at each moult; the human skeleton is internal and grows continuously with the body, so growth is smooth and continuous.

9.5 The Human Growth Curve

Human growth, plotted as total height against age, is roughly sigmoid but with a special fea-

ture: **two periods of rapid growth** — in infancy (the first two years) and at puberty (the **adolescent growth spurt**). This is seen most clearly on a **height-velocity curve**, which plots the rate of growth (cm per year) against age.



Height-velocity curves for boys and girls. Note the high rate in infancy and the adolescent growth spurt, which occurs about two years earlier in girls.

KEY POINTS Features of human growth

- Growth is fastest in infancy, then slows during childhood.
- A marked adolescent growth spurt occurs at puberty: earlier in girls (about 11–13 years) than in boys (about 13–15 years), which is why girls of 12 are often taller than boys of the same age.
- Growth stops when the epiphyseal (growth) plates of the long bones fuse, under the action of sex hormones.
- Body parts grow at different rates — **allometric growth**: the head is proportionally large in a newborn (about a quarter of body length) but grows relatively little afterwards, while the legs grow greatly, changing the body proportions.

9.6 Growth Patterns: Determinate and Indeterminate

DEFINITION Determinate and indeterminate growth

Determinate growth is growth that stops at a characteristic adult size (or when the organ matures): it is shown by most animals, by leaves, flowers and fruits, and by annual plants

such as maize and cowpea. **Indeterminate growth** continues throughout life, because permanent growth zones (meristems, growth plates) remain active: it is shown by perennial plants (trees), fish, and reptiles.

In plants, two kinds of growth are distinguished:

- **Primary growth:** elongation of roots and shoots from **apical meristems** (at the tips); it produces the sigmoid curve of a seedling.
- **Secondary growth:** increase in girth (thickening) of stems and roots in woody dicotyledons, due to the **vascular cambium** producing secondary xylem (wood) and secondary phloem, and the cork cambium producing bark. This is why an old iroko or baobab tree keeps thickening year after year.

9.7 Hormonal Control of Growth in Humans

Growth is coordinated by **hormones**, chemical messengers secreted by endocrine glands and carried in the blood to their target organs.

Hormone	Gland	Role in growth
Growth hormone (GH)	Anterior pituitary	Stimulates protein synthesis, cell division and elongation of long bones (via the epiphyseal plates) throughout childhood
Thyroxine	Thyroid	Controls the basal metabolic rate; essential for normal growth and for development of the brain and skeleton
Testosterone / Oestrogen	Testes / Ovaries	Trigger the pubertal growth spurt and development of secondary sexual characteristics; later cause fusion of the epiphyseal plates, ending growth
Insulin	Pancreas (islets of Langerhans)	Promotes uptake of glucose and protein synthesis; needed for normal tissue growth

PROPERTY Disorders of growth hormone

The effect of growth hormone depends on the age at which the imbalance occurs (this is examinable as applied knowledge, not as a formal proof):

- Deficiency before puberty → **dwarfism** (pituitary dwarf): very short stature but normal body proportions and normal intelligence.
- Excess before puberty → **gigantism**: abnormal overall tallness, because the epiphyseal plates are still open.
- Excess in adults → **acromegaly**: the long bones can no longer lengthen, so only the bones of the hands, feet, jaw and brow thicken, giving coarse features.

DID YOU KNOW? Thyroxine and cretinism

Severe thyroxine deficiency in childhood (for example where dietary iodine is lacking) causes cretinism: stunted growth together with mental retardation. Iodised salt, distributed in Cameroon precisely to combat iodine-deficiency disorders, prevents this condition.

9.8 Hormonal Control of Growth in Plants (Brief)

Plant growth is controlled by **plant growth substances** (phytohormones), produced in small quantities and transported to their sites of action.

- **Auxins** (e.g. indole-acetic acid, IAA), produced in shoot and root tips, promote **cell elongation** by softening cell walls; they also mediate phototropism and apical dominance.
- **Gibberellins** promote **stem elongation** (internode lengthening), stimulate seed germination and can induce flowering; dwarf varieties of maize and rice often lack active gibberellin.
- **Cytokinins**, produced mainly in roots, promote **cell division** (cytokinesis) and delay ageing (senescence) of leaves.

Auxins and cytokinins act together: the balance between them determines whether cultured plant cells form shoots or roots — the principle behind in vitro propagation of plantain and banana plantlets used by Cameroonian farmers.

WORKED EXAMPLE 10 Interpreting a growth experiment

The dry mass of samples of maize seedlings was measured every week after germination:

Week	1	2	3	4	5	6	7
Mean dry mass (g)	0.4	0.6	1.1	2.0	3.0	3.6	3.8

Questions: (a) Why was dry mass, not fresh mass, used? (b) Between which weeks was the growth rate greatest? (c) Identify the phases of growth shown and explain the slowing after week 5.

SOLUTION

Solution.

- (a) Dry mass measures the organic matter actually synthesised by the seedlings; fresh mass would fluctuate with water uptake and loss and would not prove true growth.
- (b) Growth rate = $\frac{\Delta m}{\Delta t}$. Weekly increases: 0.2, 0.5, 0.9, 1.0, 0.6, 0.2 g. The greatest increase is between weeks 4 and 5 (1.0 g/week).
- (c) Weeks 1–2: lag phase (slow growth, seedlings establishing). Weeks 2–5: exponential (log) phase (rapid, accelerating growth). Weeks 5–7: decelerating then stationary phase — growth slows and nearly stops as the plants approach maturity and limiting factors (nutrients, light, space) take effect. The curve is therefore sigmoid.

EXERCISE 9 Growth curves and hormones

1. Define growth and state the three cellular processes that bring it about.
2. Sketch and label the sigmoid growth curve, naming its four phases.
3. Explain, with a labelled sketch, why the growth curve of a locust is stepped.
4. A boy receives injections of growth hormone from age 8. Predict the consequence if the treatment is continued in excessive doses into adulthood, and name the resulting condition.
5. State one role of each of the following in plant growth: auxin, gibberellin, cytokinin.

SOLUTION Exercise — Growth curves and hormones

1. Growth is the permanent, irreversible increase in size and dry mass of an organism, brought about by cell division, cell enlargement and cell differentiation.
2. S-shaped curve with: lag phase (slow start), exponential/log phase (steepest), decelerating phase, stationary phase (plateau at adult size).
3. The locust's rigid chitinous exoskeleton cannot stretch, so length stays constant during each instar (plateau); at each moult (ecdysis) the old cuticle is shed and the body expands rapidly before the new cuticle hardens (vertical jump). The curve is therefore a series of steps.
4. Excess GH before puberty causes gigantism; if excess continues after the epiphyseal plates have fused, the long bones cannot lengthen but the bones of the hands, feet

and face thicken — this is acromegaly.

5. Auxin: cell elongation (and phototropism); gibberellin: stem/internode elongation and germination; cytokinin: cell division and delay of senescence.

KEY POINTS Chapter summary — Growth

- Growth = irreversible increase in size and dry mass (division, enlargement, differentiation).
- The sigmoid curve (lag, log, decelerating, stationary) describes most organisms; the grand period of growth covers all four phases.
- Arthropods grow in steps because of moulting (ecdysis); humans grow smoothly with spurts in infancy and at puberty.
- Determinate growth stops at maturity; indeterminate growth continues for life (trees, fish).
- Human growth is controlled by GH, thyroxine, sex hormones and insulin; plant growth by auxins, gibberellins and cytokinins.

10 INTEGRATION ACTIVITY

10.1 Aims of the Activity

KEY POINTS Learning Objectives

This integration activity consolidates knowledge from the entire Reproduction chapter. By completing it, you will:

- Demonstrate understanding of **sexual reproduction** in humans and flowering plants, including hormonal control, gametogenesis, fertilisation and early development.
- Apply concepts of **asexual reproduction** and **vegetative propagation** to agricultural practice.
- Design a controlled **germination** experiment and interpret **growth curves** (sigmoid and pubertal spurt).
- Evaluate **contraceptive methods** in relation to the menstrual cycle and make evidence-based recommendations.
- Communicate scientific reasoning clearly in a written advisory report and an oral presentation.

10.2 The Mfoundi Family Scenario

EXPERIMENT Case Study Context

The Mfoundi family lives on the outskirts of Yaoundé, Cameroon. Mr. and Mrs. Mfoundi have three children aged 14, 10 and 6. They wish to space their next child by at least three years for health and economic reasons. Mrs. Mfoundi is 32 years old, has a regular 28-day menstrual cycle, and has no medical contraindications to hormonal contraception. The family also runs a small diversified farm of 0.5 ha where they cultivate:

- Plantain (*Musa* spp.) – they want to multiply an improved high-yielding cultivar using suckers.
- Mango (*Mangifera indica*) – they plan to graft scions of a late-fruited variety onto established rootstocks.
- Maize (*Zea mays*) – they have just harvested seeds from their best plants and want to test germination before the next planting season.

Their 14-year-old son, Pierre, is in the middle of puberty. The family has recorded his height every six months since age 10. They also measured the height of their maize crop

weekly from sowing to harvest (12 weeks). The data are summarised below.

Week after sowing	Maize height (cm)	Notes
1	2.1	Emergence
2	4.8	
3	9.5	
4	16.2	
5	24.7	
6	34.1	
7	43.0	
8	50.5	
9	55.8	
10	58.9	
11	60.2	
12	60.5	Harvest

Age (years)	Pierre's height (cm)	Notes
10.0	138.2	Pre-pubertal
10.5	140.5	
11.0	143.0	
11.5	146.2	
12.0	150.8	Onset of growth spurt
12.5	156.5	
13.0	163.0	Peak height velocity
13.5	168.2	
14.0	171.5	Deceleration

The family asks you, as a team of Upper Sixth biology students, to act as their advisers. You must produce a one-page advisory report (maximum 300 words) and defend it in a 5-minute class presentation.

10.3 Tasks

- Contraceptive choice:** Select the most appropriate contraceptive method for Mrs. Mfoundi. Justify your choice by linking its mode of action to the hormonal events of the menstrual cycle (FSH, LH, oestrogen, progesterone). Discuss effectiveness, reversibility, side-effects and cultural acceptability in the Cameroonian context.
- Vegetative propagation plan:** Design a step-by-step plan to multiply the improved plan-

tain cultivar using suckers and to graft the mango scions onto rootstocks. Include timing, selection criteria, sanitation, and after-care. Explain why these methods are considered asexual reproduction and how they preserve the genotype.

3. **Germination experiment:** Propose a controlled experiment to test the germination capacity of the maize seeds. State the hypothesis, variables (independent, dependent, controlled), experimental set-up (number of replicates, substrate, temperature, moisture, light), data to be collected, and statistical analysis (e.g., germination percentage, mean germination time).
4. **Growth curve analysis:**
 - (a) Plot the maize height data and Pierre's height data on separate graphs (time on x-axis, height on y-axis). Identify the type of growth curve for each (sigmoid vs. pubertal spurt).
 - (b) For the maize curve, label the lag, exponential, and stationary phases. Calculate the approximate maximum growth rate (cm week^{-1}) during the exponential phase.
 - (c) For Pierre's curve, determine the age at peak height velocity (PHV) and the peak height velocity value (cm year^{-1}). Relate these to hormonal changes (testosterone, growth hormone, IGF-1).
5. **Integration and report writing:** Synthesize your answers into a concise advisory report (≤ 300 words) addressing all four points. Prepare a 5-minute presentation highlighting the scientific reasoning and practical recommendations.

10.4 Model Answers

EXAMPLE Task 1: Contraceptive Choice for Mrs. Mfoundi

Recommended method: Combined Oral Contraceptive Pill (COCP) containing ethinylestradiol (oestrogen) and levonorgestrel (progestogen).

Mode of action linked to menstrual cycle:

- The COCP maintains high circulating levels of synthetic oestrogen and progesterone.
- These exert negative feedback on the hypothalamus and anterior pituitary, suppressing the secretion of **GnRH**, **FSH** and **LH**.
- Crucially, the mid-cycle LH surge is prevented; without it, **ovulation** does not occur.
- Additional effects: progestogen thickens **cervical mucus** (blocking sperm penetration) and alters the **endometrium** (making it unsuitable for implantation).

Effectiveness: Perfect use $\approx 99.7\%$ (Pearl Index 0.3); typical use $\approx 91\%$ (Pearl Index 9).

Reversibility: Full fertility returns within 1–3 cycles after stopping. Side-effects: Nausea, breast tenderness, weight change, mood changes (usually transient); rare but serious risk of thromboembolism (counselling required). Cameroonian context: Widely available in public health centres and pharmacies at subsidised cost; culturally accepted among educated urban couples; daily adherence counselling is essential. Alternative: If daily adherence is a concern, Long-Acting Reversible Contraceptives (LARCs) such as the sub-

dermal implant (etonogestrel) or hormonal IUD (levonorgestrel) offer >99% effectiveness with user-independent action.

EXAMPLE Task 2: Vegetative Propagation Plan

A. Plantain (*Musa* spp.) – Multiplication by Sword Suckers

METHOD Step-by-step

1. Timing: Start of rainy season (March–April in Yaoundé area) for optimal establishment.
2. Mother plant selection: Healthy, disease-free (no Banana Bunchy Top Virus, no nematode damage), 6–8 months after planting, bearing the desired high-yielding cultivar traits.
3. Sucker selection: Choose sword suckers (narrow, lanceolate leaves, strong conical corm) 30–50 cm tall. Avoid water suckers (broad leaves, weak corm).
4. Detachment: Use a clean, sharp machete. Cut the sucker from the mother corm, ensuring a portion of corm (5–10 cm) and roots remain attached.
5. Sanitation: Trim leaves to 2–3 cm from the corm to reduce transpiration. Dip corm in 1% sodium hypochlorite (bleach) solution for 5 minutes, then rinse in clean water. Optional: hot water treatment (50 °C for 20 min) against nematodes.
6. Planting: Prepare holes 30 cm × 30 cm × 30 cm at 3 m × 3 m spacing. Add 5–10 kg compost or well-decomposed manure per hole. Plant sucker upright, cover corm with soil, firm gently.
7. After-care: Mulch heavily with dry grass or banana leaves. Water regularly until established. Apply mulch and organic fertiliser every 3–4 months. Monitor for pests (weevils, nematodes) and diseases.

Genetic basis: Suckers arise from meristematic tissue of the mother corm → **mitotic divisions** only → genetically identical clones. The improved cultivar's genotype is preserved exactly.

B. Mango (*Mangifera indica*) – Grafting (Cleft Graft)

METHOD Step-by-step

1. Rootstock: 6–12 month-old seedlings of a local, hardy, well-adapted variety (e.g., 'Kensington' or local 'Mangoro'), pencil-thickness (≈ 1 cm diameter), actively growing.
2. Scion collection: During dry season dormancy (December–January). Select 15–20 cm mature shoots with 3–4 plump buds from the desired late-fruiting variety. Wrap in damp cloth, keep cool.
3. Grafting operation: Under shade. Decapitate rootstock 20–30 cm above soil. Make a vertical cleft (3–4 cm deep) in the rootstock stem. Cut scion base into a wedge. Insert scion into cleft, aligning **cambium** layers on at least one side. Secure tightly with grafting tape (polythene).
4. Sealing: Apply grafting wax or petroleum jelly over the union and cut surfaces to prevent desiccation.
5. After-care: Place in high-humidity propagator (or cover with clear polythene bag) for 3–4 weeks. Ventilate gradually. Remove any rootstock sprouts below the graft. Harden off by increasing exposure to ambient conditions over 2 weeks.
6. Field planting: Transplant at onset of rains (March), 10 m $\times$ 10 m spacing.

Genetic basis: Grafting joins two genotypes: the scion (fruiting variety) and the rootstock (adaptability). The union is **asexual** – no meiosis, no gamete fusion. The scion's genotype is preserved, ensuring true-to-type fruit quality and late-fruiting trait.

EXAMPLE Task 3: Controlled Germination Experiment

METHOD Experimental Design

1. Hypothesis: Maize seeds selected from the family's best plants have a germination percentage $\geq 90\%$ and a mean germination time ≤ 4 days under optimal conditions.
2. Variables:
 - Independent: Seed source (Family's saved seeds vs. Certified commercial seeds of same variety – control).
 - Dependent: Germination percentage (%), Mean Germination Time (MGT, days).
 - Controlled: Substrate (sterile river sand or moist filter paper), Temperature ($25^{\circ}\text{C} \pm 1^{\circ}\text{C}$), Moisture (field capacity / constant moisture), Light (continuous darkness – maize germinates in dark), Seed depth (3 cm in sand / between papers), Seed viability (use seeds from same harvest year).
3. Set-up: Randomised Complete Block Design. 4 replicates (blocks) $\times$ 2 treatments = 8 experimental units. Each unit: 50 seeds. Seeds placed in Petri dishes (filter paper) or sand trays. Incubate in a germination cabinet or dark room at 25°C .
4. Data collection: Count germinated seeds (radicle $\geq 2\text{mm}$) daily for 10 days. Record number germinated per day (n_i) for each replicate.
5. Statistical analysis:
 - Germination percentage: $GP = \frac{\text{total germinated}}{\text{total seeds}} \times 100$. Compare treatments using Chi-square test (χ^2) on contingency table (germinated vs. non-germinated).
 - Mean Germination Time (MGT): $MGT = \frac{\sum(n_i \times d_i)}{\sum n_i}$ where d_i = day of count. Compare MGT between treatments using unpaired t-test (check normality, equal variance) or Mann-Whitney U test. Significance level $\alpha = 0.05$.

EXAMPLE Task 4: Growth Curve Analysis

A. Maize Height Growth

KEY POINTS Sigmoid Growth Curve

The maize data produce a classic sigmoid (S-shaped) curve typical of annual plants.

- Lag phase (Weeks 1–2): Slow growth; establishment of root system, limited photosynthetic area.
- Exponential (log) phase (Weeks 3–7): Rapid cell division and elongation; maximum photosynthetic rate; steepest slope.
- Stationary phase (Weeks 8–12): Growth slows and stops; resources diverted to grain filling; senescence begins.

Maximum Growth Rate Calculation: Use the steepest interval (Weeks 3–7):

$$\text{Rate} = \frac{\Delta \text{height}}{\Delta \text{time}} = \frac{43.0 - 9.5}{7 - 3} = \frac{33.5}{4} = 8.375 \approx 8.4 \text{ cm week}^{-1}$$

B. Pierre's Height Growth

KEY POINTS Pubertal Growth Spurt

The data show a pubertal growth spurt – a rapid increase in height velocity during adolescence.

- Peak Height Velocity (PHV) age: 13.0 years (height = 163.0 cm).
- PHV value: Calculate over the interval centred on PHV (12.5–13.5 years):

$$\text{PHV} = \frac{168.2 - 156.5}{13.5 - 12.5} = \frac{11.7}{1.0} = 11.7 \text{ cm year}^{-1}$$

Hormonal basis: Rising **testosterone** (Leydig cells) stimulates **GH** secretion from anterior pituitary → GH stimulates liver to produce **IGF-1** (Insulin-like Growth Factor 1). GH and IGF-1 act synergistically on **epiphyseal plates** (growth plates) to increase chondrocyte proliferation and matrix synthesis → longitudinal bone growth. Testosterone also directly stimulates growth plate maturation (eventually causing epiphyseal closure).

EXAMPLE Task 5: Advisory Report (Model – 298 words)

To the Mfoundi Family,

We recommend the combined oral contraceptive pill (COCP) for child spacing. It

maintains high oestrogen/progestogen levels, suppressing the hypothalamic-pituitary axis and preventing the mid-cycle LH surge; thus ovulation is blocked. Additional effects: thickened cervical mucus and altered endometrium. Perfect-use effectiveness is 99.7%; typical use 91%. It is fully reversible (fertility returns in 1–3 cycles), affordable, and widely available in Yaoundé health centres. Cultural acceptability is high; we advise daily intake at the same time and counselling on transient side-effects (nausea, weight change) and rare thromboembolism risk. If adherence is difficult, a subdermal implant or hormonal IUD are excellent alternatives.

For the farm: Plantain – multiply using sword suckers (30–50 cm) from healthy mother plants at the start of the rains. Sanitise corms in 1% bleach, plant at 3 m × 3 m with compost and mulch. This **clonal propagation** preserves the improved genotype exactly. Mango – cleft-graft scions of the late-fruited variety onto 6–12 month local rootstocks during the dry season. Align cambium, seal with wax, maintain high humidity until union forms (3–4 weeks). Grafting combines the scion's fruited genotype with the rootstock's hardiness – an asexual method ensuring true-to-type fruit.

Maize germination test: Four replicates of 50 seeds each on sterile moist sand at 25 °C in darkness. Count daily for 10 days. Calculate germination percentage and Mean Germination Time $MGT = \frac{\sum n_i d_i}{\sum n_i}$. Compare with certified seeds using χ^2 and t-tests ($\alpha = 0.05$). Expect ≥90% germination if seeds are viable.

Growth analysis: Maize follows a sigmoid curve; maximum growth rate $\approx 8.4 \text{ cm week}^{-1}$ (weeks 3–7). Pierre shows a pubertal spurt; PHV $\approx 11.7 \text{ cm year}^{-1}$ at age 13, driven by testosterone, GH and IGF-1 acting on epiphyseal plates. Both curves illustrate distinct biological growth patterns: deterministic in annual plants, hormone-mediated in adolescents.

Implementing these recommendations will support family health, farm productivity, and informed monitoring of Pierre's development. We are available for follow-up guidance. Upper Sixth Biology Advisory Team

WARNING Common Pitfalls in the Integration Activity

- Do not confuse the contraceptive mechanism: the COCP prevents the LH surge (not the FSH surge alone) via negative feedback.
- Vegetative propagation must emphasise genetic identity (clones) – do not describe it as sexual reproduction or imply genetic recombination.
- In the germination experiment, control all abiotic factors (temperature, moisture, light, substrate); omitting any invalidates the test.
- When calculating growth rates, use the steepest part of the curve: exponential phase for maize (weeks 3–7), PHV interval (12.5–13.5 years) for Pierre.
- The advisory report must be concise (≤ 300 words) and address all four tasks; marks

are awarded for integration and application, not just isolated facts.

- For mango grafting, specify cambial alignment – this is the key to a successful vascular union.
- In growth analysis, distinguish between absolute growth rate (cm/time) and relative growth rate; the question asks for absolute rate.

11 METHODS AND WORKED EXAMPLES

11.1 Skill 1: Identifying the Type of Asexual Reproduction

METHOD Recognising asexual reproduction mechanisms

When faced with a description, a diagram or a micrograph of an organism reproducing asexually, proceed as follows.

1. Count the parents. If only **one parent** is involved and no gametes are formed or fused, the reproduction is asexual.
2. Look at what is produced. Ask: is the offspring a whole new individual, an outgrowth, or a resistant cell?
3. Match the observation to the mechanism:
 - **Binary fission**: one cell splits into two equal daughter cells (Amoeba, Paramecium, bacteria).
 - **Budding**: a small outgrowth (bud) forms on the parent, then detaches (Hydra, yeast).
 - **Fragmentation / regeneration**: the body breaks into pieces, each regenerating a whole organism (planaria, starfish, Spirogyra).
 - **Sporulation**: spores produced in a sporangium germinate into new individuals (Mucor, ferns, mosses).
 - **Vegetative propagation**: new plants from vegetative organs — stems (runners, rhizomes, tubers, cuttings), roots or leaves — e.g. cassava and sugarcane cuttings, Irish potato tubers, ginger rhizomes, onion bulbs.
 - **Parthenogenesis**: an unfertilised egg develops into an adult (aphids, drones in bees).
4. State the genetic consequence: offspring are **genetically identical** (clones) to the parent, except for rare mutations.

Reflex: “One parent + no gamete fusion = asexual; offspring = clone.”

WORKED EXAMPLE 11 Identifying modes of asexual reproduction

An Upper Sixth student observes the following in the school biology garden and laboratory: (a) a cassava stem cutting planted in the soil develops roots and shoots; (b) a slide of Mucor shows black sporangia releasing spores; (c) a Hydra with a small lateral outgrowth; (d) an Amoeba constricting into two cells. For each case, name the type of asexual repro-

duction and justify your answer. State one advantage and one disadvantage of asexual reproduction.

SOLUTION

Solution.

1. (a) Cassava cutting: **Vegetative propagation** (propagation by stem cuttings). Justification: a new plant develops from a vegetative organ (stem), not from a seed or a gamete.
2. (b) Mucor: **Sporulation**. Justification: spores are produced inside sporangia; each spore can germinate into a new mycelium.
3. (c) Hydra: **Budding**. Justification: a small bud grows from the parent's body wall and will detach as an independent individual.
4. (d) Amoeba: **Binary fission**. Justification: the single parent cell divides by mitosis into two genetically identical daughter cells.

Advantage: rapid reproduction and colonisation (no mate needed, no energy spent on gametes); all offspring are well adapted to the parent's environment.

Disadvantage: no genetic variation, so the whole population may be wiped out by a single disease or environmental change (e.g. cassava mosaic disease spreading through infected cuttings).


11.2 Skill 2: Accounting for Chromosome Numbers in Gamete Formation and Double Fertilization

METHOD Tracking chromosome and DNA numbers in flowering plants

1. Fix the diploid number $2n$ given in the question. Write it down before any calculation.
2. Apply the key rule: **meiosis** halves the chromosome number ($2n \rightarrow n$); **fertilization** restores it ($n + n \rightarrow 2n$).
3. In the flower, remember the ploidy of each structure:
 - Pollen grain (male gametophyte): n ; it contains a **generative nucleus** (n) which divides by mitosis into two male gamete nuclei (n each), plus a tube nucleus (n).
 - Embryo sac (female gametophyte): n ; it contains the **egg cell** (n) and two **polar nuclei** (n each, i.e. $2n$ when fused).
4. Double fertilization: one male nucleus (n) + egg (n) $\rightarrow$ **zygote** ($2n$); the other male nucleus (n) + the two polar nuclei ($2n$) $\rightarrow$ **endosperm nucleus** ($3n$, triploid).

5. Check: the zygote must be $2n$; the endosperm must be $3n$. If your answer differs, re-examine step 4.

Reflex: "Egg n , zygote $2n$, endosperm $3n$."



Double fertilization in a flowering plant: one male nucleus fuses with the egg cell (zygote, $2n$); the second fuses with the two polar nuclei (endosperm, $3n$).

WORKED EXAMPLE 12 Chromosome numbers in maize reproduction

Maize (*Zea mays*) has a diploid number $2n = 20$. (a) State the number of chromosomes in: (i) a pollen mother cell; (ii) a pollen grain nucleus; (iii) the egg cell; (iv) the zygote; (v) the endosperm. (b) Explain how the endosperm comes to have its chromosome number. (c) How many chromosomes would the embryo of the seed contain?

SOLUTION

Solution.

(a)

- (i) Pollen mother cell: it is a diploid cell of the anther, so $2n = 20$ chromosomes.
- (ii) Pollen grain nucleus: produced by meiosis of the pollen mother cell, so $n = 10$ chromosomes.
- (iii) Egg cell: a haploid gamete, $n = 10$ chromosomes.

4. (iv) Zygote: egg (n) + male nucleus (n) = $2n = 20$ chromosomes.

5. (v) Endosperm: triploid, $3n = 30$ chromosomes.

(b) Origin of the endosperm: during **double fertilization**, the second male gamete nucleus ($n = 10$) fuses with the two polar nuclei ($n + n = 20$) in the embryo sac. The fusion product therefore contains $10 + 20 = 30$ chromosomes ($3n$). It divides mitotically to form the endosperm, the nutritive tissue of the seed.

(c) Embryo: the embryo develops by repeated mitosis of the zygote, so every embryonic cell contains $2n = 20$ chromosomes.

Check: zygote $2n = 20$ ✓; endosperm $3n = 30$ ✓.

11.3 Skill 3: Interpreting Germination Experiments and Calculating Germination Percentage

METHOD Analysing a germination experiment

1. Identify the variables: independent variable (the factor varied: water, oxygen, temperature, light), dependent variable (germination: number or percentage of seeds germinated), controlled variables (same seed batch, same number of seeds, same duration).
2. Compute the germination percentage:

$$\text{Germination \%} = \frac{\text{number of seeds germinated}}{\text{total number of seeds sown}} \times 100$$

3. Compare set-ups: a seed germinates only if **water**, **oxygen** and a **suitable temperature** are all present. Light is needed only by some species (e.g. lettuce); most seeds germinate equally well in darkness.
4. Draw the conclusion by linking each result to the missing or present condition, and state the role of each factor (water: imbibition and enzyme activation; oxygen: aerobic respiration for energy; temperature: enzyme activity).
5. Evaluate reliability: large sample, replicates, control of other variables.

WORKED EXAMPLE 13 Germination of bean seeds

A student in Buea sets up four flasks, each with 50 bean seeds: Flask A — moist cotton wool, open to air, at 25°C ; Flask B — dry cotton wool, open to air, 25°C ; Flask C — seeds completely covered with boiled, cooled water, 25°C ; Flask D — moist cotton wool, open to air, at 4°C in a refrigerator. After 5 days: A = 45 germinated, B = 0, C = 0, D = 2. (a)

Calculate the germination percentage in each flask. (b) Explain the result of each flask. (c) State the purpose of boiling and cooling the water in flask C.

SOLUTION

Solution.

(a) Germination % = $\frac{\text{germinated}}{50} \times 100$:

- A: $\frac{45}{50} \times 100 = 90\%$
- B: $\frac{0}{50} \times 100 = 0\%$
- C: 0%
- D: $\frac{2}{50} \times 100 = 4\%$

(b) Flask A (control): water, oxygen and a suitable temperature are all present, so enzymes are activated, food reserves are mobilised and respiration supplies energy — high germination (90%). Flask B: no water, so no imbibition and no enzyme activation — no germination. Flask C: boiled water contains almost no dissolved oxygen; without oxygen, aerobic respiration cannot release the energy needed for growth — no germination. Flask D: the low temperature slows enzyme activity drastically, so germination is very poor (4%).

(c) Boiling expels dissolved air (oxygen) from the water; cooling prevents the heat from killing the seeds, so that the only missing factor is oxygen. This makes flask C a valid test of the need for oxygen.

Conclusion: bean seeds need water, oxygen and a suitable temperature to germinate.

11.4 Skill 4: Reading Hormone Graphs of the Menstrual Cycle


METHOD Interpreting the menstrual cycle graph

1. Identify each curve by its profile over the 28 days: **FSH** rises early (follicle growth); **oestrogen** rises to a peak just before mid-cycle; **LH** shows a sharp peak at about day 14 (this triggers ovulation); **progesterone** dominates the second half (secreted by the corpus luteum).
2. Locate ovulation: it occurs just after the LH peak, around **day 14**.
3. Link hormones to events: FSH → follicle development; oestrogen → repair and thickening of the endometrium + positive feedback causing the LH surge; LH → ovulation and formation of the corpus luteum; progesterone → maintenance of the endometrium and negative feedback inhibiting FSH and LH.
4. Explain menstruation: if no fertilization occurs, the corpus luteum degenerates, pro-

gesterone and oestrogen fall, the endometrium breaks down and is shed (days 1–5).

5. Answer “what if pregnant?”: the corpus luteum is maintained (by HCG from the embryo), progesterone stays high, no menstruation.

Reflex: “LH peak = ovulation; progesterone fall = menstruation.”



Typical profiles of FSH, LH, oestrogen and progesterone during a 28-day menstrual cycle (schematic).

WORKED EXAMPLE 14 Analysing a hormone graph

The graph of a woman's cycle shows: FSH rising from day 1 to day 6; oestrogen peaking on day 12; a sharp LH peak on day 13–14; progesterone rising from day 15 to day 22 then falling. (a) On which day did ovulation most likely occur? Justify. (b) Which structure secretes progesterone after day 15, and from what does it develop? (c) Explain why progesterone falls after day 22. (d) Explain how a pregnancy would change the progesterone curve. (e) Explain how the combined contraceptive pill exploits this cycle.

SOLUTION

Solution.

(a) Ovulation: about day 14. The LH surge (peak on day 13–14) is the direct trigger of ovulation; the egg is released roughly a day after the LH peak, i.e. around day 14.

(b) Progesterone is secreted by the **corpus luteum**, which develops from the ruptured Graafian follicle left in the ovary after ovulation, under the influence of LH.

(c) Fall of progesterone: in the absence of fertilization, the corpus luteum degenerates after about 10–12 days (it becomes the corpus albicans). With no corpus luteum, progesterone (and oestrogen) secretion collapses; the endometrium is no longer maintained and breaks down — menstruation begins.

(d) If pregnant: the embryo implants and secretes **HCG** (human chorionic go-

nadotrophin), which maintains the corpus luteum. Progesterone therefore remains high instead of falling; the endometrium is preserved and menstruation does not occur. Later, the placenta takes over progesterone production.

(e) The combined pill contains synthetic oestrogen and progesterone. These hormones exert negative feedback on the hypothalamus and pituitary, inhibiting FSH and LH secretion. Without FSH, follicles do not mature; without the LH surge, ovulation does not occur, so no egg is available for fertilization.

11.5 Skill 5: Comparing Spermatogenesis and Oogenesis

METHOD Constructing a comparison of gametogenesis

1. Fix the framework: both processes involve **mitosis** of germ cells, growth, **meiosis**, and maturation/differentiation.
2. Compare point by point (never mix the two columns of your reasoning): site, timing, number of functional gametes per meiosis, size of gametes, continuity of the process.
3. Key figures to remember:
 - Spermatogenesis: 1 primary spermatocyte → 4 functional spermatozoa; continuous from puberty; occurs in the seminiferous tubules; controlled by FSH and testosterone (LH acts on Leydig cells).
 - Oogenesis: 1 primary oocyte → 1 functional ovum + polar bodies; begins before birth, meiosis I completed only just before ovulation, meiosis II completed only at fertilization; controlled by FSH, LH, oestrogen, progesterone.
4. Hormonal control: name the gland (hypothalamus/pituitary), the hormone, the target and the effect. GnRH → FSH + LH.

Feature	Spermatogenesis	Oogenesis
Site	Seminiferous tubules of testes	Ovaries
Timing	Continuous from puberty	Begins before birth; meiosis completed only at ovulation/fertilization
Functional gametes per meiosis	4 spermatozoa	1 ovum + polar bodies
Cytokinesis	Equal	Unequal (cytoplasm kept in one large cell)
Gamete	Small, motile, no reserves	Large, non-motile, rich in yolk
Main hormones	FSH, LH, testosterone	FSH, LH, oestrogen, progesterone

WORKED EXAMPLE 15 Gamete yield and hormonal control

(a) A man produces sperm continuously from puberty. Starting from 100 primary spermatocytes, how many functional spermatozoa are produced? Starting from 100 primary oocytes, how many functional ova are produced? Explain the difference. (b) Describe the hormonal control of the testes. (c) A tumour destroys a man's Leydig (interstitial) cells. Predict and explain the consequences.

SOLUTION

Solution.

(a) Each primary spermatocyte undergoes meiosis I (two secondary spermatocytes) then meiosis II, giving 4 spermatids, each of which differentiates into a functional spermatozoon. So 100 primary spermatocytes $\rightarrow 100 \times 4 = 400$ spermatozoa.

Each primary oocyte undergoes meiosis with unequal cytokinesis: meiosis I gives one large secondary oocyte and one small polar body; meiosis II gives one large ovum and further polar bodies. Only 1 functional ovum results per primary oocyte. So 100 primary oocytes $\rightarrow 100$ ova (plus polar bodies that degenerate).

Explanation: in oogenesis the cytoplasm is conserved in a single large cell so that the ovum contains abundant food reserves for the early embryo; in spermatogenesis all four products are small, motile gametes, maximising the chance that one reaches the egg.

(b) Hormonal control of the testes: the hypothalamus secretes **GnRH**, which stimulates the anterior pituitary to release **FSH** and **LH**. FSH acts on the Sertoli cells of the seminiferous tubules, promoting spermatogenesis and the nourishment of developing sperm. LH stimulates the Leydig (interstitial) cells to secrete **testosterone**, which is essential for sperm maturation, the development of secondary sexual characteristics and libido. Testosterone exerts negative feedback on the hypothalamus and pituitary, keeping its level stable.

(c) Destruction of Leydig cells: testosterone secretion falls drastically. Consequences: spermatogenesis is impaired or stops (infertility), because testosterone is required for

sperm maturation; secondary sexual characteristics may regress; libido decreases. Because the negative feedback is removed, blood levels of LH (and FSH) rise — a diagnostic sign that the failure is in the testis, not the pituitary.

11.6 Skill 6: Analysing Growth Curves and Growth Patterns

METHOD Interpreting a growth curve


1. Identify the type of curve:

- **Sigmoid (S-shaped)** curve: slow growth (lag phase) → rapid exponential growth → stationary phase — typical of annual plants and many organs.
- **Intermittent/stepped** curve: growth in steps after each moult (ecdysis) — typical of insects and arthropods with an exoskeleton.
- Human growth curve: rapid in infancy, steady in childhood, a pubertal growth spurt, then a plateau at adulthood.

2. Compute the growth rate between two times: $\text{rate} = \frac{\text{size at } t_2 - \text{size at } t_1}{t_2 - t_1}$.

3. Explain each phase biologically (cell division, cell elongation, differentiation; moulting; hormonal spurts).

4. Link to hormones when asked: in plants, auxins, gibberellins, cytokinins; in humans, growth hormone (pituitary) and thyroid hormones, plus sex hormones at puberty.



Growth of a maize seedling: lag phase, exponential phase and stationary phase form a sigmoid (S-shaped) curve.

WORKED EXAMPLE 16 Growth of a maize seedling and of a locust

The height of a maize seedling was measured weekly: week 1: 4 cm; week 2: 8 cm; week 3: 18 cm; week 4: 34 cm; week 5: 50 cm; week 6: 58 cm; week 7: 60 cm; week 8: 60 cm. (a) Calculate the growth rate between weeks 3 and 5, and between weeks 6 and 8. (b) Describe and explain the shape of the curve. (c) A locust's growth curve is stepped, not smooth. Explain. (d) Name one hormone that promotes stem elongation in maize and state its action.

SOLUTION

Solution.

(a) Growth rate, weeks 3–5: $\frac{50 - 18}{5 - 3} = \frac{32}{2} = 16 \text{ cm/week}$.

Growth rate, weeks 6–8: $\frac{60 - 58}{8 - 6} = \frac{2}{2} = 1 \text{ cm/week}$.

(b) Shape: the curve is **sigmoid** (S-shaped). Lag phase (weeks 1–2): growth is slow because the seedling has few cells and is still establishing leaves and roots. Exponential (log) phase (weeks 2–5): rapid growth by active cell division in meristems and cell elongation, with photosynthesis supplying abundant food — the rate peaks at 16 cm/week. Stationary phase (weeks 6–8): growth slows and stops as the plant reaches maturity; cells differentiate and resources are diverted to flowering and fruiting.


(c) Stepped curve of the locust: insects have a rigid **exoskeleton** (cuticle) that cannot expand. Growth in size is only possible when the old cuticle is shed (**ecdysis**/moulting) and the new, soft cuticle expands before hardening. Between moults, size is constant, so the curve rises in steps at each moult. Moulting is controlled by the hormone ecdysone.

(d) Gibberellins promote stem elongation: they stimulate both cell division and especially cell elongation in the internodes. (Auxins also promote cell elongation in shoots by stimulating cell wall loosening.)

11.7 Skill 7: Evaluating Birth Control Methods and Life Cycles**METHOD** Answering questions on contraception and life cycles

1. Classify the method before discussing it: natural (abstinence, rhythm/calendar method), barrier (condom, diaphragm), hormonal (pill, implant, injectable), intrauterine device (IUD), surgical/permanent (vasectomy, tubal ligation).
2. State the mechanism precisely: e.g. pill = negative feedback preventing ovulation; condom = physical barrier preventing sperm meeting egg (and reduces sexually transmitted infections); IUD = prevents implantation; vasectomy = cutting and tying the vas deferens.
3. Evaluate: reliability, advantages, disadvantages, protection (or not) against STIs such as HIV.

- For life cycles: in **alternation of generations** (ferns, mosses), identify the sporophyte ($2n$, produces spores by meiosis) and the gametophyte (n , produces gametes by mitosis); fertilization restores $2n$. In **metagenesis** (e.g. Obelia, jellyfish), an asexual phase (polyp) alternates with a sexual phase (medusa).
- Always place meiosis correctly: meiosis occurs at spore formation in plants, at gamete formation in animals.



Alternation of generations in a fern: a diploid sporophyte alternates with a haploid gametophyte; meiosis occurs at spore formation.

WORKED EXAMPLE 17 Contraception and the fern life cycle

(a) A couple in Yaoundé asks a health worker for advice. Compare the condom and the combined pill under the headings: mechanism, reliability, and protection against sexually transmitted infections. (b) Explain why vasectomy is considered a permanent method and why it does not affect a man's sexual characteristics. (c) In a fern, state the ploidy of the sporophyte, the spores, the gametophyte (prothallus) and the zygote, and state where meiosis occurs.

SOLUTION

Solution.

(a) Condom: Mechanism — a thin sheath worn over the penis acts as a physical barrier, preventing sperm from entering the vagina. Reliability — high when used correctly every time, but it can fail through tearing or incorrect use. STI protection — yes: it is the only common method that greatly reduces transmission of HIV and other sexually transmitted infections.

Combined pill: Mechanism — synthetic oestrogen and progesterone cause negative feedback on the pituitary, inhibiting FSH and LH, so follicles do not mature and ovulation is prevented; the cervical mucus is also thickened. Reliability — very high when taken daily

without omission. STI protection — none: it offers no barrier against infections.

Advice: for protection against both pregnancy and STIs, the condom (or both methods combined) is preferable.

(b) Vasectomy: the **vas deferens** (sperm duct) on each side is cut and tied, so sperm can no longer leave the testes and appear in the semen. It is permanent because reconnection is difficult and usually not attempted. Sexual characteristics are unaffected because the Leydig cells still secrete testosterone directly into the blood (not through the vas deferens); erection, libido and ejaculation of seminal fluid (from the prostate and seminal vesicles) continue normally — only sperm are absent from the ejaculate.

(c) Fern life cycle (alternation of generations):

- Sporophyte (the familiar fern plant): diploid ($2n$).
- Spores: produced in sporangia on the fronds by meiosis, so haploid (n).
- Gametophyte (prothallus): develops from a spore by mitosis, so haploid (n); it bears antheridia and archegonia producing gametes (n) by mitosis.
- Zygote: formed by fusion of sperm and egg, so diploid ($2n$); it grows into a new sporophyte.

Meiosis occurs at spore formation in the sporangia of the sporophyte — this is the key event that halves the chromosome number and switches the cycle from the diploid to the haploid generation.

WARNING Frequent errors in GCE answers

- Confusing pollination (transfer of pollen to the stigma) with fertilization (fusion of gamete nuclei) — they are distinct events.
- Forgetting that the endosperm is triploid ($3n$), not diploid.
- Saying “LH causes menstruation” — in fact it is the fall of progesterone that triggers menstruation; the LH surge triggers ovulation.
- Claiming oogenesis produces four ova per meiosis — it produces one ovum plus polar bodies.
- Placing meiosis at gamete formation in plants — in plants meiosis produces spores, not gametes.
- Writing percentages without showing the formula and substitution — always show $\frac{\text{germinated}}{\text{total}} \times 100$.

12 EXERCISES

12.1 I check my knowledge

EXERCISE 10 MCQ: Asexual reproduction mechanisms

Which of the following is not a form of asexual reproduction in animals?

1. Budding in Hydra
2. Binary fission in Amoeba
3. Parthenogenesis in aphids
4. Conjugation in Paramecium

EXERCISE 11 True/False with justification

State whether each statement is true or false. Justify your answer briefly.

1. The pollen grain of a flowering plant is the male gamete.
2. Double fertilization occurs in all vascular plants.
3. The corpus luteum secretes mainly oestrogen during the luteal phase.
4. In human spermatogenesis, one primary spermatocyte produces four functional spermatozoa.

EXERCISE 12 Definitions

Define the following terms precisely:

1. Alternation of generations
2. Metagenesis
3. Epigeal germination
4. Menarche

EXERCISE 13 Direct calculation: Growth rate

A locust nymph measures 8 mm at the beginning of an instar and 12 mm at the end of the same instar, which lasts 5 days. Calculate the average daily growth rate in mm/day. If the growth follows a sigmoid curve, explain why the rate would not be constant throughout the instar.

12.2 I practise

EXERCISE 14 Labelling and hormonal events

Draw a large, labelled diagram of the human female reproductive system (anterior view). On your diagram, indicate the site of fertilization and the site of implantation. Then, describe the hormonal events that occur from day 1 to day 14 of a typical 28-day menstrual cycle, naming the hormones, their sources and their main actions.

EXERCISE 15 Data analysis: Hormone profiles

The table below shows the concentrations of four hormones in the blood of a woman over a 28-day menstrual cycle.

Day	FSH (mIU/mL)	LH (mIU/mL)	Oestrogen (pg/mL)	Progesterone (ng/mL)
1	5.2	4.8	30	0.5
4	6.1	5.0	45	0.4
7	7.5	6.2	80	0.3
10	8.0	7.5	150	0.3
13	9.5	25.0	280	0.4
14	10.0	45.0	300	0.5
17	4.0	8.0	180	12.0
20	3.5	5.5	150	25.0
23	3.0	4.0	120	30.0
26	2.8	3.5	80	20.0
28	4.5	4.0	40	2.0

1. Plot the four hormone curves on the same axes (use a suitable scale).
2. Deduce the most probable day of ovulation. Justify your answer using the data.
3. Explain the role of the LH surge in ovulation and the subsequent formation of the corpus luteum.

EXERCISE 16 Comparison: Spermatogenesis vs Oogenesis

Copy and complete the table below to compare spermatogenesis and oogenesis in humans.

Feature	Spermatogenesis	Oogenesis
Site		
Timing (when does it start/end?)		
Number of functional gametes per meiosis		
Hormonal control (key hormones)		
Presence of a resting phase		

EXERCISE 17 Experiment design: Oxygen and germination

Design an investigation to show that oxygen is necessary for the germination of bean seeds (*Phaseolus vulgaris*). Your answer must include:

1. A clear hypothesis.
2. The independent, dependent and two controlled variables.
3. A description of the experimental set-up (including a control).
4. The expected results and how they support the hypothesis.
5. One precaution to ensure validity.

12.3 I prepare for the GCE**EXERCISE 18 Structured essay: Female reproductive system and menstrual cycle**

(20 marks)

1. Draw a fully labelled diagram of the human female reproductive system (anterior view). [6 marks]
2. Describe the hormonal changes that occur during the follicular phase (days 1–14) of the menstrual cycle. Explain how these changes lead to ovulation. [8 marks]
3. Outline the role of progesterone in preparing the uterus for implantation. [6 marks]

EXERCISE 19 Life cycle interpretation: Fern

(15 marks) The diagram below shows the life cycle of a fern (*Dryopteris*) with stages labelled A to F. (Imagine a typical alternation of generations diagram: A = mature sporophyte, B = sporangium, C = spores, D = gametophyte/prothallus, E = antheridia/archegonia, F = zygote/young sporophyte.)

1. Identify which stages are haploid (n) and which are diploid ($2n$). [4 marks]
2. State where meiosis occurs and where fertilization occurs. [3 marks]
3. Explain the significance of the alternation of generations in ferns, particularly in relation to genetic variation and dispersal. [8 marks]

EXERCISE 20 Problem question: Growth disorders and birth control

(20 marks)

1. A 10-year-old child shows stunted growth (height well below the 3rd percentile). Using your knowledge of growth hormone (GH) and thyroxine (T_4), suggest three possible endocrine causes for this condition. For each cause, describe a diagnostic test that could confirm it. [12 marks]
2. A couple wishes to delay pregnancy for two years. They are considering the following methods: combined oral contraceptive pill, copper intrauterine device (IUD), male condom, and subdermal implant. For each method, state:
 - The precise stage of reproduction it blocks.
 - One advantage and one disadvantage.

[8 marks]

13 SOLUTIONS TO THE EXERCISES

13.1 I check my knowledge

SOLUTION Exercise 1 — MCQ: Asexual reproduction mechanisms

The correct answer is d) Conjugation in Paramecium.

Justification:

- Budding in Hydra is asexual: a small outgrowth of the body wall develops tentacles, then detaches as a new individual.
- Binary fission in Amoeba is asexual: the nucleus divides by mitosis and the cytoplasm splits into two genetically identical daughter cells.
- Parthenogenesis in aphids is asexual: unfertilized eggs develop into new females (a form of reproduction without fertilization).
- Conjugation in Paramecium is not asexual reproduction: two individuals exchange micronuclear material (a sexual process of genetic recombination). It does not by itself increase the number of individuals, and it involves meiosis and exchange of gametic nuclei.

SOLUTION Exercise 2 — True/False with justification

1. False. The pollen grain is the male gametophyte; it contains the male gametes (two sperm nuclei formed from the generative nucleus). The male gamete itself is the sperm nucleus, not the whole grain.
2. False. Double fertilization (one sperm nucleus fusing with the egg cell to form the zygote, the other with the polar nuclei to form the endosperm) is characteristic of angiosperms (flowering plants) only. Gymnosperms and pteridophytes (ferns) are vascular plants but do not show double fertilization.
3. False. The corpus luteum secretes mainly progesterone during the luteal phase (with some oestrogen). Oestrogen is secreted mainly by the developing Graafian follicles during the follicular phase.
4. True. In spermatogenesis, one primary spermatocyte undergoes meiosis I to give two secondary spermatocytes, then meiosis II to give four spermatids, all of which differentiate into four functional spermatozoa (no polar bodies are formed).

SOLUTION Exercise 3 — Definitions

1. Alternation of generations: a life cycle in which a multicellular **diploid** generation (the **sporophyte**, which produces spores by meiosis) alternates regularly with a multicellular **haploid** generation (the **gametophyte**, which produces gametes by mitosis). Example: ferns, mosses.
2. Metagenesis: a particular alternation of generations in which a sexual generation alternates with an asexual generation that reproduces by budding, as in Obelia (the polyp colony buds off medusae; the medusae reproduce sexually).
3. Epigeal germination: germination in which the **hypocotyl** elongates and curves, carrying the cotyledons above the soil surface (e.g. bean, *Phaseolus vulgaris*).
4. Menarche: the first menstrual period (first menstruation) of a girl, marking the beginning of reproductive life at puberty (typically around 11–14 years).

SOLUTION Exercise 4 — Direct calculation: Growth rate

Calculation of the average daily growth rate:

$$\text{Growth rate} = \frac{\text{final length} - \text{initial length}}{\text{time}} = \frac{12 \text{ mm} - 8 \text{ mm}}{5 \text{ days}} = \frac{4}{5} = 0.8 \text{ mm/day.}$$

Why the rate is not constant over a sigmoid curve: A sigmoid (S-shaped) growth curve has three phases:

- a lag phase at the start, when growth is slow (cells are few, the nymph is adapting and feeding little);
- a log (exponential) phase, when growth is rapid and the rate is maximal;
- a stationary/deceleration phase, when growth slows as the instar approaches its end (the cuticle hardens before ecdysis, food intake falls).

Because the instantaneous rate (slope of the curve) rises then falls, the value 0.8 mm/day is only an average; the actual daily rate is lower than this at the beginning and end of the instar and higher in the middle.

13.2 I practise**SOLUTION** Exercise 5 — Labelling and hormonal events

Diagram (anterior view) — the candidate's drawing must show and label: the two ovaries,

the Fallopian tubes (oviducts) with fimbriae, the uterus (with endometrium and myometrium), the cervix, the vagina. The site of fertilization is the ampulla of the Fallopian tube (outer third of the oviduct); the site of implantation is the endometrium of the uterus (usually the upper posterior wall).

Hormonal events, day 1 to day 14 (follicular phase):

1. Days 1–5 (menstruation): the previous corpus luteum has degenerated, so progesterone and oestrogen levels are low. The endometrium breaks down and is shed. Low oestrogen/progesterone removes negative feedback on the hypothalamus and pituitary.
2. From day 1: the hypothalamus secretes GnRH, stimulating the anterior pituitary to release FSH (and some LH).
3. FSH (source: anterior pituitary) stimulates the growth of several ovarian follicles; the developing follicles secrete oestrogen.
4. Oestrogen (source: growing follicles) causes repair and thickening (proliferation) of the endometrium; at moderate levels it exerts negative feedback on FSH (so only one dominant follicle survives).
5. Around days 12–13: oestrogen reaches a high, sustained peak and switches to positive feedback on the pituitary, triggering a surge of LH (and a smaller FSH surge).
6. Day 14: the LH surge causes ovulation — rupture of the mature Graafian follicle and release of the secondary oocyte.

SOLUTION Exercise 6 — Data analysis: Hormone profiles

1. Plotting. Use days 1–28 on the x-axis. Because the units and magnitudes differ, plot LH and FSH on one scale (0–50 mIU/mL) and use a secondary y-axis (or a separate scaled plot) for oestrogen (0–300 pg/mL) and progesterone (0–30 ng/mL). Key features the graph must show: a sharp LH peak at day 14 (45 mIU/mL), a smaller FSH peak around days 13–14, an oestrogen peak at day 14 (300 pg/mL) just before/at ovulation, and a progesterone peak around day 23 (30 ng/mL) in the luteal phase.

2. Day of ovulation: day 14. Justification from the data:

- LH rises sharply from 7.5 (day 10) to 45.0 mIU/mL (day 14) — the **LH surge** directly triggers ovulation.
- Oestrogen peaks at 300 pg/mL on day 14, then falls (180 by day 17) — the oestrogen peak precedes and triggers the LH surge.
- Progesterone is low (≤ 0.5 ng/mL) until day 14, then rises steeply (12.0 by day 17) — progesterone is secreted by the corpus luteum, which can only form after ovulation.

3. Role of the LH surge: the LH surge (a) completes the maturation of the Graafian follicle

and the oocyte (resumption of meiosis I), (b) weakens the follicle wall (enzymatic digestion) causing its rupture and release of the oocyte — ovulation; (c) transforms the remaining granulosa and theca cells of the ruptured follicle into the **corpus luteum** (luteinization), which then secretes progesterone (and oestrogen) to maintain the endometrium during the luteal phase.

SOLUTION Exercise 7 — Comparison: Spermatogenesis vs Oogenesis

Feature	Spermatogenesis	Oogenesis
Site	Seminiferous tubules of the testes	Ovaries (follicles)
Timing	Starts at puberty; continuous throughout life	Begins before birth (oocytes arrest in prophase I); resumes at puberty; ends at menopause
Functional gametes per meiosis	4 functional spermatozoa	1 functional ovum + 2–3 polar bodies (which degenerate)
Hormonal control	FSH (Sertoli cells), LH (testosterone from Leydig cells), testosterone	FSH, LH, oestrogen, progesterone (cyclic)
Resting phase	Absent — continuous process	Present — long arrests in prophase I and metaphase II

SOLUTION Exercise 8 — Experiment design: Oxygen and germination

1. Hypothesis: oxygen is necessary for the germination of bean seeds; seeds deprived of oxygen will not germinate.

2. Variables:

- Independent variable: presence/absence of oxygen.
- Dependent variable: germination (percentage of seeds germinated, or radicle length after a fixed time).
- Controlled variables (any two): same species/age/number of seeds; same temperature; same water availability; same light conditions; same duration of the experiment.

3. Set-up:

- Control (Set-up A): soak bean seeds, place them on moist cotton wool in an open flask exposed to air at room temperature.
- Experiment (Set-up B): place identical soaked seeds in a flask of freshly boiled and cooled water (boiling expels dissolved air), and cover the water surface with a layer of oil to prevent oxygen diffusing back in. (Alternatively, use alkaline pyrogallol in a sealed flask to absorb oxygen.)
- Keep both flasks in the same place (same temperature and light) for 4–5 days.

4. Expected results: seeds in Set-up A (with oxygen) germinate — radicles emerge and grow; seeds in Set-up B (without oxygen) do not germinate. Since the only difference is the availability of oxygen, the result supports the hypothesis that oxygen is necessary for germination (needed for aerobic respiration, which releases the energy for growth of the embryo).

5. Precaution: use a large number of seeds (e.g. 20 per set-up) and repeat the experiment, to avoid the effect of dead or non-viable seeds; also check that seeds in B are still alive by transferring them to air afterwards — they should then germinate, proving lack of oxygen (not death) prevented germination.

13.3 I prepare for the GCE

SOLUTION Exercise 9 — Structured essay: Female reproductive system and menstrual cycle (20 marks)

(a) Diagram [6 marks]. Examiner expectations: large clear drawing in pencil, ruled label lines, no shading. Award 1 mark for each correctly drawn and labelled structure, up to 6: ovary, Fallopian tube (oviduct) with fimbriae, uterus, endometrium/myometrium, cervix, vagina. Deduct for labels touching the wrong structure or for a diagram too small.

(b) Hormonal changes in the follicular phase [8 marks] — indicative mark scheme (1 mark each):

1. At the start of the cycle, progesterone and oestrogen levels are low because the previous corpus luteum has degenerated.
2. Low hormone levels remove negative feedback, so the hypothalamus secretes GnRH.
3. GnRH stimulates the anterior pituitary to secrete FSH (and LH).
4. FSH stimulates the growth and maturation of ovarian follicles.
5. The developing follicles secrete increasing amounts of oestrogen.
6. Oestrogen repairs and thickens the endometrium (proliferation) after menstruation.

7. High, sustained oestrogen exerts positive feedback, causing a surge of LH (and FSH) around days 13–14.
 8. The LH surge triggers ovulation: rupture of the Graafian follicle and release of the secondary oocyte (day 14).
- (c) Role of progesterone in preparing the uterus [6 marks] — 1 mark each:
1. Progesterone is secreted by the corpus luteum after ovulation.
 2. It maintains and further thickens the endometrium (secretory phase).
 3. It stimulates the endometrial glands to secrete nutrients (uterine milk) for the early embryo.
 4. It increases the blood supply/vascularization of the endometrium.
 5. It inhibits uterine (myometrial) contractions, preventing expulsion of the embryo.
 6. It inhibits FSH/LH release (negative feedback), preventing development of new follicles during a possible pregnancy; it also thickens cervical mucus, blocking sperm entry.

SOLUTION Exercise 10 — Life cycle interpretation: Fern (15 marks)

(a) Ploidy of the stages [4 marks] — 1 mark each:

- Diploid ($2n$): A (mature sporophyte), B (sporangium — its tissues are part of the sporophyte), F (zygote and young sporophyte).
- Haploid (n): C (spores), D (gametophyte/prothallus), E (antheridia and archegonia, and the gametes they produce).

(Award 4 marks for all six correct; 2–3 marks for partial correct classification.)

(b) Sites of meiosis and fertilization [3 marks]:

- Meiosis occurs in the sporangium (B): spore mother cells divide by meiosis to produce haploid spores (C). [2 marks]
- Fertilization occurs on/in the gametophyte (D), when a sperm swims from an antheridium to the egg in an archegonium (E), forming the diploid zygote (F). [1 mark]

(c) Significance of alternation of generations [8 marks] — indicative points:

- Genetic variation (4 marks): meiosis in the sporangium produces genetically varied spores (independent assortment and crossing over); the gametophyte generation produces gametes, and sexual fusion of gametes from different prothalli combines genes from two parents, increasing variation; variation provides raw material for natural selection and adaptation.
- Dispersal (3 marks): the sporophyte produces large numbers of tiny, light spores that

are dispersed by wind over long distances, allowing ferns to colonize new, moist habitats (e.g. the wet slopes of Mount Cameroon); spores are resistant and can survive unfavourable conditions.

- Life-cycle efficiency (1 mark): the dominant sporophyte is adapted to land (vascular tissue, roots, leaves) while the small gametophyte ensures sexual reproduction where water is available for sperm to swim — each generation exploits the conditions suited to it.

SOLUTION Exercise 11 — Problem question: Growth disorders and birth control (20 marks)

(a) Three possible endocrine causes of stunted growth [12 marks — 4 per cause: 2 for the cause, 2 for the test]:

1. Growth hormone (GH) deficiency (hyposecretion by the anterior pituitary, e.g. pituitary dwarfism). Diagnostic test: measure blood GH level, which will be abnormally low; since GH is secreted in pulses, use a stimulation test (e.g. insulin or arginine provocation) — a deficient child shows no rise in GH; low IGF-1 level also confirms it.
2. Primary hypothyroidism (thyroxine/T4 deficiency) — thyroxine is needed for normal growth and for the action of GH; deficiency in childhood causes stunted growth (cretinism if congenital). Diagnostic test: measure blood T4 (low) and TSH (high in primary hypothyroidism, since negative feedback is removed).
3. Secondary hypothyroidism (TSH deficiency) — the pituitary fails to stimulate the thyroid, leading to low thyroxine and stunted growth. Diagnostic test: low T4 with low or normal TSH (distinguishing it from primary hypothyroidism); a TRH stimulation test shows whether the pituitary can respond.

(Any three coherent endocrine causes with a matching test are accepted; e.g. GH-receptor insensitivity (Laron syndrome) with a GH/IGF-1 assay showing high GH but low IGF-1.)

(b) Birth control methods [8 marks — 2 per method]:

- Combined oral contraceptive pill: blocks ovulation (oestrogen + progesterone inhibit FSH/LH by negative feedback, so no follicle matures); also thickens cervical mucus. Advantage: very effective when taken correctly and reversible. Disadvantage: must be taken daily; possible side effects (nausea, weight gain, increased risk of blood clots); no protection against STIs.
- Copper IUD: acts mainly before fertilization — copper ions are spermicidal and the device causes a uterine reaction hostile to sperm and to implantation. Advantage: long-acting (several years), no daily action needed, hormone-free. Disadvantage: can cause heavier/more painful periods; requires insertion by trained personnel; no protection against STIs.

- Male condom: blocks the deposition of sperm in the vagina (physical barrier preventing sperm from reaching the egg). Advantage: cheap, available without prescription, and protects against STIs including HIV. Disadvantage: can tear or slip if misused; must be used correctly at every intercourse.
- Subdermal implant: releases progestogen which blocks ovulation and thickens cervical mucus (blocking sperm passage). Advantage: very effective and long-acting (3–5 years), reversible on removal. Disadvantage: requires minor surgery for insertion/removal; may cause irregular bleeding; no protection against STIs.

Examiner expectations: for full marks the candidate must state the precise stage blocked (ovulation, sperm transport, fertilization or implantation), not merely "prevents pregnancy", and give one realistic advantage and disadvantage per method.

14 SUMMARY SHEET



Mind map of the chapter "Reproduction": the nine major blocks of the GCE A-Level syllabus.

FICHE BILAN L'essentiel à connaître pour l'examen

Key definitions

- **Reproduction**: biological process by which organisms produce new individuals, ensuring the continuity of the species.
- **Asexual reproduction**: production of offspring from a single parent, without gametes or fertilization; offspring are genetically identical to the parent (**clones**). Mechanisms: binary fission (Amoeba, bacteria), budding (Hydra, yeast), fragmentation (Spirogyra, planarians), sporulation (ferns, fungi such as Mucor), vegetative

propagation (runners, tubers, rhizomes, bulbs, cuttings – e.g. cassava, cocoyam and plantain in Cameroon).

- **Parthenogenesis**: development of an unfertilized egg into a new individual (aphids, honeybees – drones develop from unfertilized eggs).
- **Sexual reproduction**: fusion of two haploid gametes (**fertilization**) to form a diploid **zygote**; meiosis and fertilization together produce genetic variation.
- **Pollination**: transfer of pollen grains from the anther to the stigma (self-pollination vs cross-pollination; agents: wind, insects, water, birds).
- **Double fertilization** (flowering plants): one male nucleus fuses with the egg cell → diploid zygote ($2n$); the second male nucleus fuses with the two polar nuclei → triploid **endosperm** nucleus ($3n$), which develops into the nutritive tissue of the seed.
- **Seed dispersal**: scattering of seeds away from the parent plant, reducing competition; agents: wind, water, animals, explosive (self-dispersal) mechanisms.
- **Germination**: resumption of growth of the embryo when water, oxygen and a suitable temperature are present; **epigeal** germination (cotyledons carried above ground, e.g. bean) vs **hypogeal** germination (cotyledons remain below ground, e.g. maize).
- **Spermatogenesis**: continuous production of spermatozoa in the seminiferous tubules of the testes from puberty onwards: spermatogonia ($2n$) → primary spermatocytes → (meiosis I) secondary spermatocytes (n) → (meiosis II) spermatids → spermatozoa.
- **Oogenesis**: production of ova in the ovaries; begins before birth (oocytes arrested in meiosis I), resumes at puberty; usually one secondary oocyte is released per cycle at **ovulation**, and meiosis II is completed only if fertilization occurs.
- **Menstrual cycle**: approximately 28 days; phases: menstrual (days 1–5), follicular/proliferative (days 6–13), ovulation (about day 14), luteal/secretory (days 15–28).
- **Fertilization** (human): fusion of sperm and ovum nuclei, normally in the oviduct (Fallopian tube), restoring the diploid number.
- **Implantation**: embedding of the blastocyst in the endometrium of the uterus, about 6–7 days after fertilization.
- **Alternation of generations**: alternation of a haploid **gametophyte** generation (produces gametes by mitosis) and a diploid **sporophyte** generation (produces spores by meiosis) in the life cycle of a plant (mosses, ferns).
- **Metagenesis**: alternation of a sexual generation and an asexual generation in the life cycle of an animal (e.g. Obelia: polyp colony reproduces asexually to release medusae, which reproduce sexually).

- **Growth**: irreversible increase in size, cell number and dry mass, accompanied by cell division, cell enlargement and differentiation.

Hormonal control – the essentials

- Male: hypothalamus secretes **GnRH** → anterior pituitary: **FSH** stimulates Sertoli cells (support of spermatogenesis); **LH** (= ICSH) stimulates the interstitial (Leydig) cells → **testosterone** (development of secondary sexual characteristics, maintenance of spermatogenesis; negative feedback on GnRH, FSH and LH).
- Female: **FSH** → growth of Graafian follicles → **oestrogen** (repair and thickening of the endometrium; high level triggers the LH surge); **LH** surge → **ovulation** and formation of the corpus luteum → **progesterone** (maintains the endometrium; negative feedback on FSH and LH). If there is no pregnancy, the corpus luteum degenerates → fall of oestrogen and progesterone → breakdown of the endometrium → menstruation.
- Pregnancy: **hCG** (secreted by the embryo, then the placenta) maintains the corpus luteum; later the placenta itself secretes oestrogen and progesterone, which maintain the uterus and inhibit FSH/LH (no further ovulation). **Oxytocin** triggers uterine contractions at birth (positive feedback); **prolactin** stimulates milk production after birth.
- Growth: **GH** (growth hormone, anterior pituitary) stimulates protein synthesis and growth of bones and soft tissues; **thyroxine** (thyroid) is essential for normal growth, development and metamorphosis; sex hormones (testosterone, oestrogen) cause the pubertal growth spurt and later the closure of the epiphyses (end of growth in height).

Development sequence (learn the order!)

Zygote → cleavage → morula → blastocyst → implantation (day 6–7) → embryo → foetus (from week

Birth control methods

- Natural: abstinence, rhythm/calendar method (avoiding intercourse around ovulation), withdrawal – all relatively unreliable.
- Barrier: condom (also protects against STIs including HIV), diaphragm.
- Hormonal: contraceptive pill (oestrogen + progesterone inhibit FSH/LH and block ovulation), implants, injections.
- Intra-uterine device (IUD): prevents implantation/fertilization.
- Surgical (permanent): vasectomy (cutting of the sperm ducts) in the male, tubal ligation (cutting of the oviducts) in the female.

Growth curves and patterns

- Sigmoid (S-shaped) curve in annual plants and many organs: lag phase → exponential (log) phase → stationary phase.

- Human growth curve: rapid growth in infancy, slow steady growth in childhood, a spurt at puberty, then growth stops in adulthood.
- Intermittent (stepwise) growth in arthropods due to **ecdysis** (moulting of the rigid exoskeleton).
- Measurements: length/height, fresh mass, and **dry mass** (most accurate – it excludes variations in water content).

Common mistakes to avoid

- Confusing pollination (transfer of pollen to the stigma) with fertilization (fusion of gamete nuclei).
- Forgetting that the endosperm is triploid ($3n$), not diploid.
- Saying FSH/LH act on the uterus: they act on the ovary; oestrogen and progesterone act on the uterus.
- Placing ovulation at the start of the cycle: it occurs around day 14 of a 28-day cycle.
- Confusing oestrogen (follicular phase) with progesterone (luteal phase and pregnancy).
- Confusing gametophyte (produces gametes, n) with sporophyte (produces spores, $2n$).
- Mixing up epigeal (cotyledons above ground) and hypogeal (cotyledons below ground) germination.
- Writing that the sperm fertilizes the ovum “in the uterus”: fertilization normally occurs in the oviduct.

GCE tips

- Draw and label diagrams: flower, seed, testis/seminiferous tubule, ovary, menstrual cycle hormone graph – marks are awarded for correct labels, proportions and a title.
- In hormonal essays, always state: gland – hormone – target organ – effect – feedback.
- For graph questions (hormone levels, growth curves), describe each phase and link it to a precise biological event (e.g. LH peak → ovulation).
- Use precise vocabulary: “implantation” not “attachment”; “ovulation” not “release of the ovum into the uterus”.
- Manage time: structured questions on reproduction carry many small marks – answer every sub-question.





Signed • OnBuch+

ONBUCH+ TEACHING TEAM

FOLLOWS THE OFFICIAL MINESEC SYLLABUS

OnBuch+