

BIOLOGY • UPPER SIXTH

UPPER SIXTH — BIOLOGY

CHAPTER 2

RESPONSE & COORDINATION IN ORGANISMS

SUGGESTED TIME: 25 PERIODS

IN THIS CHAPTER

This chapter explores how organisms detect stimuli and coordinate responses through sense organs, the nervous system and the endocrine system in animals, and through hormones and tropic movements in plants. It provides the structural, physiological and molecular foundations needed for GCE Advanced Level questions on reception, coordination and control.

- By the end of this chapter I can define stimulus, response, receptor, effector and coordination, and classify stimuli and responses.
- By the end of this chapter I can explain the principles of stimulus reception, including sensory thresholds, adaptation and the generator potential.
- By the end of this chapter I can describe the structure of the mammalian eye and relate each part to its function.
- By the end of this chapter I can explain image formation, accommodation, control of light intensity, light/dark adaptation and the photochemistry of photoreception.
- By the end of this chapter I can describe the structure of the ear and explain the mechanisms of hearing and balance.
- By the end of this chapter I can describe the organisation of the nervous system, the structure of neurones and the generation and transmission of nerve impulses.

**DOWNLOAD ON-
BUCH**

Free • Google Play
AI tutor, past papers,
results

**GO ONBUCH+**

All signed courses, unlimited AI
tutor, PDF sheets — OnBuch+
tab in the app

THIS SIGNED COURSE CONTAINS

COURSE

complete, illustrated,
step by step

METHODS

and worked examples

EXERCISES

exam style + solutions

**SUMMARY
SHEET**

the essentials to
remember

**Signed • OnBuch+**

ONBUCH+ TEACHING TEAM
FOLLOWS THE OFFICIAL MINESEC SYLLABUS

OnBuch+

CONTENTS

- | | |
|---|---|
| 1. Stimuli, Responses and Principles of Reception | Autonomic Nervous System |
| 2. The Eye: Structure and Image Formation | 8. The Endocrine System and Mechanism of Hormone Action |
| 3. Control of Light Intensity, Adaptation and Accommodation | 9. Functions of the Endocrine Glands |
| 4. Photoreception and the Ear: Hearing and Balance | 10. Plant Movements, Plant Hormones, Light and Plant Growth |
| 5. The Nervous System and Neurones | 11. Integration activity |
| 6. Transmission of Impulses and the Synapse | 12. Methods and worked examples |
| 7. Reflex Actions, Brain, Spinal Cord and | 13. Exercises |
| | 14. Solutions to the exercises |
| | 15. Summary sheet |

CHAPTER OBJECTIVES

- By the end of this chapter I can define stimulus, response, receptor, effector and co-ordination, and classify stimuli and responses.
- By the end of this chapter I can explain the principles of stimulus reception, including sensory thresholds, adaptation and the generator potential.
- By the end of this chapter I can describe the structure of the mammalian eye and relate each part to its function.
- By the end of this chapter I can explain image formation, accommodation, control of light intensity, light/dark adaptation and the photochemistry of photoreception.
- By the end of this chapter I can describe the structure of the ear and explain the mechanisms of hearing and balance.
- By the end of this chapter I can describe the organisation of the nervous system, the structure of neurones and the generation and transmission of nerve impulses.
- By the end of this chapter I can explain synaptic transmission, reflex actions, and the roles of the brain, spinal cord and autonomic nervous system.
- By the end of this chapter I can describe the endocrine system, the mechanism of hormone action and the functions of the main endocrine glands.

- By the end of this chapter I can compare nervous and hormonal coordination and explain plant movements, plant hormones and the effect of light on plant growth.

PREREQUISITES

- Cell structure and ultrastructure (Lower Sixth): membranes, mitochondria, organelles
- Movement of substances across membranes: diffusion, osmosis, active transport
- Protein structure and enzymes as biological catalysts
- Basic chemistry of ions (Na^+ , K^+) and concentration gradients
- Cell division and growth in plants (Lower Sixth)

1 STIMULI, RESPONSES AND PRINCIPLES OF RECEPTION

During an evening football match at the Stade Omnisports in Yaoundé, a striker hears the referee's whistle, sees the ball emerging from the shadows into the floodlights, and volleys it into the net — all in less than a second. How did his eyes adjust instantly to the changing light intensity? How did the sound of the whistle travel from his ear to his leg muscles? And why do the bean seedlings on his grandmother's farm in Bafoussam bend toward the morning sun? Behind each of these events lies the same fundamental biological machinery: **receptors** that detect changes, **nerves** and **hormones** that carry and integrate information, and **effectors** that produce a precise, coordinated response. In this first section we lay the foundations: what stimuli and responses are, how receptors work, and how the strength of a stimulus is coded in the language of the nervous system.

1.1 Stimulus, Response and Coordination: the Basic Vocabulary

Every living organism, from an Amoeba in a pond near Lake Chad to the striker on the pitch, survives only because it can detect changes in its environment and react to them appropriately. A wilting hibiscus plant recovers after watering; a goat in Ngaoundéré runs when it hears a dog bark; you withdraw your hand from a hot cooking pot before you even feel the pain. All these are examples of the same sequence: a change is detected, information is processed, and a reaction follows.

DEFINITION Stimulus, response, receptor, effector, coordination

- A **stimulus** is any change in the internal or external environment of an organism that is capable of provoking a response.
- A **response** is the observable reaction of an organism (or of part of it) to a stimulus.
- A **receptor** (or sensor) is a cell or organ specialised to detect a particular type of stimulus and convert it into an electrical signal.
- An **effector** is an organ that carries out the response — in animals, usually a muscle (which contracts) or a gland (which secretes).
- **Coordination** is the linking together and control of the activities of receptors, conducting pathways and effectors so that the response is appropriate, correctly timed and correctly proportioned. In animals, coordination is achieved by the nervous system (fast, electrical) and the endocrine system (slower, chemical); in plants, by plant growth substances (hormones).


Stimuli are classified by their origin. **External stimuli** come from outside the body: light, sound, touch, the smell of ripe mangoes, a change in air temperature. **Internal stimuli** arise inside the body: a fall in blood glucose, a rise in blood CO₂, stretching of the full bladder, thirst caused by concentrated blood plasma. Internal stimuli are just as important — they drive homeostasis, the maintenance of a constant internal environment.

KEY POINTS The coordination pathway

Every coordinated response follows the same five-step plan:

stimulus → receptor → coordinator → effector → response

The coordinator (brain, spinal cord, or a simple nerve net in lower animals) receives the message, integrates it, and dispatches instructions to the effector.



The five stages of a coordinated response, illustrated by the striker reacting to the referee's whistle.

1.2 Classification of Stimuli and of Receptors

Stimuli exist in many physical forms, and evolution has produced a receptor specialised for each. The main forms of stimulus energy are:

Stimulus energy	Examples of stimuli	Receptor type (examples)
Light (electromagnetic)	brightness, colour, day length	Photoreceptors: rods and cones of the retina; plant phytochrome
Sound / vibration (mechanical)	whistle, footsteps, predator approach	Mechanoreceptors: hair cells of the cochlea
Chemical	taste, smell, blood glucose, pheromones	Chemoreceptors: taste buds, olfactory cells, carotid bodies
Mechanical (pressure, touch, stretch)	touch, blood pressure, muscle stretch	Mechanoreceptors: touch corpuscles in skin, stretch receptors in muscle
Temperature	heat of a cooking pot, cold harmattan wind	Thermoreceptors: warm and cold receptors in skin and hypothalamus
Electrical	electric fields in water	Electroreceptors: organs of electric fish (e.g. mormyrids of the River Sanaga)

Receptors are therefore named after the stimulus they detect: **photoreceptors** (light), **mechanoreceptors** (pressure, touch, sound, gravity), **chemoreceptors** (chemicals), **thermoreceptors** (temperature) and **electroreceptors** (electric fields). Pain receptors, which respond to tissue damage by several kinds of stimulus, are called **nociceptors**.

WARNING Do not confuse response and impulse

A response is the visible action of the whole organism or of an organ (the hand withdraws, the muscle contracts, the seedling bends). An impulse (action potential) is the invisible electrical signal travelling along a neurone. The impulse causes the response but is not itself the response. In examinations, writing "the response travels along the nerve" is a classic error — it is the impulse that travels.

1.3 Types of Responses in Organisms

Responses vary enormously in speed, complexity and flexibility. At GCE Advanced Level you must recognise five broad categories:

- **Taxes (singular: taxis)** — a directed movement of a whole motile organism (or cell) toward or away from a stimulus. Movement toward light is positive phototaxis (Euglena swimming toward light); movement away from light is negative phototaxis (blowfly larvae). Chemotaxis, geotaxis and rheotaxis (response to water current) follow the same logic.
- **Kineses (singular: kinesis)** — a non-directional change in the rate of movement in response to the intensity of a stimulus. Woodlice move faster and turn more often in dry, bright conditions, and slow down in humid, dark ones; by chance they accumulate where conditions favour survival. There is no "toward" or "away" — only a change of activity.
- **Tropisms** — directional growth movements of a fixed part of a plant in response to a directional stimulus: phototropism (shoots bending toward light), geotropism/gravitropism (roots growing downward), hydrotropism and thigmotropism (tendrils coiling around a support). These will be studied in detail in Section 10.
- **Reflexes** — rapid, automatic, inborn (innate) responses to specific stimuli, usually protective: withdrawal from a hot object, blinking, the knee-jerk. They do not require learning and often do not initially involve the brain.
- **Learned behaviour** — responses modified by experience: habituation, conditioning, trial-and-error learning and insight. A goalkeeper learns to anticipate a penalty taker's run-up; no reflex arc can explain that.

1.4 Properties of Receptors

However diverse they are, all receptors share four fundamental properties that you must be able to define and illustrate.

1. Specificity. Each receptor responds best to one particular form of stimulus energy — its adequate stimulus. The retina responds to light, not to sound; the cochlea to vibration, not to taste. This is why each receptor type has a characteristic structure.

2. Sensitivity and sensory threshold. Receptors are remarkably sensitive: the human eye can detect a flash of only a few photons under dark-adapted conditions. But every receptor has a **sensory threshold** — the minimum stimulus intensity that produces a detectable response (an impulse in the sensory neurone). Below threshold, nothing happens; above threshold, impulses are fired.

3. Adaptation. If a constant stimulus is maintained, many receptors gradually reduce their firing rate — they adapt. You stop noticing your shirt on your skin or the smell of your own kitchen minutes after entering. Adaptation prevents the nervous system from being flooded with unchanging, unimportant information, leaving it free to detect changes. Receptors such as touch receptors (Meissner's corpuscles) adapt rapidly; pain receptors (nociceptors) and muscle stretch receptors (muscle spindles) adapt slowly or not at all — a slow-adapting pain receptor keeps warning you as long as the damage persists.

4. Transduction. This is the master property, stated formally below.

PROPERTY **Transduction: receptors are biological energy converters**

A receptor converts (transduces) the energy of the stimulus — light, mechanical, chemical or thermal — into an electrical signal: a local depolarisation of its membrane called the **generator potential** (or receptor potential). If the generator potential reaches the threshold of the sensory neurone, it triggers action potentials (nerve impulses) that travel to the central nervous system. (The mechanism of the action potential itself is examinable in Section 6.)


The generator potential has two crucial features: it is graded (its size increases with stimulus intensity, unlike the all-or-nothing action potential) and it is local (it does not propagate; it only triggers impulses if threshold is reached).

DID YOU KNOW? **Receptor diversity in Cameroon's ecosystems**

The mormyrid electric fish of the River Sanaga and other Cameroonian rivers possess specialised electroreceptors (Knollenorgans and mormyromasts) that detect weak electric fields. They use these for navigation in murky water, communication, and locating prey — a remarkable example of a sensory modality absent in most terrestrial vertebrates.

1.5 Frequency Coding: How Stimulus Intensity is Represented

A single action potential is always the same size — it obeys the all-or-nothing law. So how does the brain know whether the light is dim or dazzling, the touch gentle or painful? The answer is **frequency coding**: as stimulus intensity increases, the generator potential becomes larger, the threshold is re-crossed sooner after each impulse, and therefore the **frequency of action potentials increases**. A strong stimulus also recruits more sensory neurones (spatial summation). The brain reads intensity from the number of impulses per second (and the number of active fibres), not from the size of each impulse.



Frequency coding: the frequency of action potentials rises with stimulus intensity and levels off at a maximum set by the refractory period of the neurone. Below the threshold intensity, no impulses are fired.

EXAMPLE Reading a frequency-coded message

A pressure receptor in the skin fires at 10 impulses per second when a feather touches it and at 80 impulses per second when a finger presses firmly. Each impulse is identical in size (about +40 mV at its peak). The brain interprets the eightfold increase in frequency — not any change in impulse size — as “firmer pressure”. If the pressure becomes painful, additional nociceptor fibres are recruited, adding a second channel of information.

1.6 Investigating Responses: the Choice Chamber Experiment

The distinction between taxes and kineses, and the reality of stimulus-response behaviour, can be demonstrated in the school laboratory with small invertebrates such as woodlice or earthworms.

METHOD Using a choice chamber to study responses of woodlice to humidity and light

1. Set up the chamber. A choice chamber is a shallow dish divided into two (or four) connected compartments. To test humidity, place damp filter paper (or moist cotton wool) under one half and a drying agent such as anhydrous calcium chloride or silica gel under the other. To test light, cover one half with opaque black card and leave the other half transparent under a lamp.
2. Introduce the animals. Place about 10 woodlice in the centre through a small hole, and leave them undisturbed for about 10 minutes to settle.

3. Record. Count the number of woodlice in each compartment at regular intervals (e.g. every minute for 10 minutes).
4. Repeat and replicate. Repeat the experiment several times, rotating the chamber 180° between runs to cancel any bias from the room (e.g. a window on one side), and pool results from several groups.
5. Analyse. Compare the numbers in each condition with the numbers expected by chance (a chi-squared test may be used). A consistent, significant accumulation in the dark, humid side demonstrates a behavioural response.

EXPERIMENT Expected results and interpretation

Woodlice typically accumulate in the dark, humid compartment. Careful observation shows they do not crawl straight there: in the dry, bright side they move faster and turn more frequently (orthokinesis and klinokinesis); in the humid, dark side they slow down and settle. This is a kinesis, not a directed taxis. Its survival value is clear: woodlice lose water easily through their permeable cuticle, so behaviour that traps them in humid refuges — under stones and rotting logs — reduces desiccation. Earthworms tested similarly avoid bright light (negative phototactic behaviour), keeping them in moist soil where gas exchange through the skin is possible.

WARNING Controls and fair testing

Common mistakes in this practical: forgetting to rotate the chamber (so a room light gradient fakes a "preference"); testing too few animals or a single run (chance clumping is misread as a response); and describing woodlice behaviour as "moving toward humidity" — a kinesis is not directional movement toward the stimulus. Always state the control, replicate, and use correct vocabulary.

WORKED EXAMPLE 1 Worked exercise: stimuli, receptors and coding

Statement. A student touches a warm water bath at 35 °C, then a hot bath at 55 °C. (a) Identify the stimulus, the receptor type, the effector and the response when she withdraws her hand from the hot bath. (b) Explain, in terms of generator potentials and frequency coding, how her nervous system distinguishes the warm bath from the hot bath. (c) State the type of response shown by the withdrawal, and justify your answer.

SOLUTION

Solution. (a) The stimulus is the high temperature of the water (an external, thermal stimulus). The receptors are thermoreceptors (and, at 55 °C, nociceptors) in the skin of the fingers. The effectors are the flexor muscles of the arm, and the response is contraction of these muscles, withdrawing the hand. (b) Heat energy is transduced by the thermoreceptors into a generator potential. At 35 °C the generator potential is small and produces a low frequency of action potentials in the sensory neurones. At 55 °C the generator potential is much larger, so threshold is re-crossed more rapidly and the impulse frequency is much higher; in addition, more receptor fibres (including nociceptors) are recruited. The central nervous system interprets the higher frequency and greater number of active fibres as a more intense, potentially damaging stimulus. The size of each action potential does not change — intensity is coded by frequency. (c) The withdrawal is a reflex (the withdrawal reflex): it is rapid, automatic, innate and protective, and it is mediated by a reflex arc through the spinal cord, so it occurs before the sensation of pain is consciously perceived.

EXERCISE 1 Check your understanding

1. Define the terms stimulus, receptor and coordination.
2. Classify each of the following receptors by the stimulus energy detected: (i) rods of the retina; (ii) taste buds; (iii) hair cells of the cochlea; (iv) cold receptors of the skin.
3. Distinguish between a taxis and a kinesis, giving one example of each.
4. Explain why a constant touch on your arm eventually goes unnoticed, naming the receptor property involved.
5. Sketch and annotate the graph of impulse frequency against stimulus intensity, indicating the threshold and the maximum frequency.

SOLUTION Exercise 1

1. Stimulus: any change in the internal or external environment capable of provoking a response. Receptor: a cell or organ specialised to detect a specific stimulus and convert it into an electrical signal. Coordination: the linking and control of receptors, conducting pathways and effectors so that responses are appropriate and well timed.
2. (i) Photoreceptor (light); (ii) chemoreceptor (chemicals); (iii) mechanoreceptor (sound vibration); (iv) thermoreceptor (temperature).
3. A taxis is a directed movement of a whole organism toward or away from a stimulus

(e.g. *Euglena* showing positive phototaxis). A kinesis is a non-directional change in the rate of movement with stimulus intensity (e.g. woodlice moving faster in dry conditions).

4. The touch receptors show adaptation: with a constant, unchanging stimulus their generator potentials decline and the frequency of impulses falls, so the brain is no longer informed of the unchanged stimulus.
5. The curve starts at zero below the threshold intensity, rises steeply once threshold is crossed, then levels off at a maximum frequency fixed by the refractory period of the neurone (as in the figure above). Axes must be labelled: stimulus intensity (x) and impulse frequency in impulses s^{-1} (y).

KEY POINTS**Section summary**

- A stimulus is a detectable environmental change; the response is the organism's reaction, carried out by effectors (muscles, glands) under the control of a coordinator.
- The universal pathway is: stimulus → receptor → coordinator → effector → response.
- Receptors are classified by the energy they detect: photo-, mechano-, chemo-, thermo- and electroreceptors.
- Responses include taxes, kinesis, tropisms, reflexes and learned behaviour.
- Receptors are specific, sensitive (with a threshold), show adaptation, and perform transduction: stimulus energy → generator potential → action potentials.
- Stimulus intensity is coded by the frequency of action potentials (and the number of fibres recruited), never by impulse size.
- The choice chamber demonstrates kinesis in woodlice: they accumulate in dark, humid conditions by changing their rate of movement, not by directed travel.


2**THE EYE: STRUCTURE AND IMAGE FORMATION**

In the previous section we established that a **stimulus** is any change in the internal or external environment capable of eliciting a response, and that **reception** is the process by which specialised cells (receptors) convert stimulus energy into nerve impulses. The mammalian eye is the supreme example of a photoreceptor organ: it collects, focuses and detects light, then transmits coded information to the brain for interpretation. We now examine its anatomy, the physics of image formation, and the common optical defects that GCE examiners frequently test.

2.1 Gross Structure of the Mammalian Eye

The eye is a hollow, roughly spherical organ housed in the bony orbit of the skull. Its wall consists of three concentric layers (tunics), and its interior is filled with transparent humours that maintain shape and transmit light. The table below summarises the major structures and their functions — learn one function for every label; the GCE often asks for structure → function matching.

Structure	Main Function(s)
Sclera	Tough, white, fibrous outer coat; protects the eye and maintains its shape; provides attachment for the extrinsic eye muscles.
Choroid	Middle vascular layer, rich in blood vessels and dark pigment (melanin); supplies nutrients and oxygen to the retina; the pigment absorbs stray light, preventing internal reflection.
Retina	Innermost light-sensitive (nervous) layer; contains the photoreceptors (rods and cones) and their neurones; site of conversion of light into nerve impulses.
Cornea	Transparent anterior continuation of the sclera; provides most of the eye's refractive (light-bending) power; avascular, nourished by tears and aqueous humour.
Conjunctiva	Thin, transparent membrane covering the anterior sclera and lining the eyelids; protects and lubricates the front of the eye.
Iris	Pigmented muscular diaphragm (containing circular and radial muscles) that controls the diameter of the pupil and hence the amount of light entering the eye.
Pupil	Central aperture (hole) in the iris through which light enters; appears black because the entering light is absorbed inside the eye.
Lens	Transparent, biconvex, elastic structure; fine-tunes focusing by changing shape (accommodation).
Ciliary body	Ring of smooth muscle (ciliary muscle); secretes aqueous humour; alters lens shape via the suspensory ligaments.
Suspensory ligaments	Fibres connecting the ciliary body to the lens; transmit tension that stretches and flattens the lens.
Aqueous humour	Clear, watery fluid in the chambers in front of the lens; nourishes the cornea and lens; maintains the shape of the front of the eye.
Vitreous humour	Clear, jelly-like substance filling the large chamber behind the lens; supports the retina and maintains the spherical shape of the eye-ball.
Optic nerve	Bundle of sensory neurones carrying visual impulses from the retina to the brain.
Fovea (yellow spot)	Tiny region of the retina on the visual axis; contains only cones at very high density; region of sharpest, most detailed colour vision.
Blind spot (optic disc)	Point where the optic nerve leaves the eye and blood vessels enter; contains no photoreceptors, so no image is detected there.



Horizontal section through the mammalian eye. Note the three concentric layers (sclera, choroid, retina), the transparent cornea continuous with the sclera, the biconvex lens held by suspensory ligaments from the ciliary body, and the fovea lying on the visual axis near the blind spot where the optic nerve leaves.

DEFINITION Fovea (Yellow Spot)

The **fovea** is a small region of the retina lying on the visual axis, close to the blind spot. It contains a very high density of cones only (no rods), and each cone is connected to its own bipolar neurone and its own ganglion cell (a 1:1:1 arrangement). This “private line” to the brain makes the fovea the region of maximum visual acuity (ability to distinguish fine detail) and the centre of colour vision.

METHOD Drawing and Labelling the Eye in the Examination

Step 1: Orient yourself — the **optic nerve** marks the back (posterior) of the eye; the **cornea** bulges at the front (anterior).

Step 2: Draw a large, clear outline (at least half a page) with a slight anterior bulge for the cornea.

Step 3: Draw the three layers at the back and sides: sclera (outermost), choroid, retina (innermost).

Step 4: Add the biconvex **lens**, held by thin **suspensory ligaments** attached to the thickened **ciliary body**.

Step 5: Draw the **iris** in front of the lens, leaving a central gap for the **pupil**.

Step 6: Mark the **blind spot** where the optic nerve leaves, and the **fovea** on the visual axis.

Step 7: Label with a ruler: label lines must be straight, must not cross, and must touch the

structure exactly. Give your drawing a title.

WARNING Protective Accessory Structures — Easy Marks

Do not forget the accessory structures of the eye; they are part of the syllabus:

- Eyelids: close reflexively (blink reflex) to protect the eye and spread tears over the cornea.
- Eyelashes: trap dust and trigger the blink reflex when touched.
- Lacrimal (tear) glands: secrete tears which moisten and nourish the cornea and contain the antibacterial enzyme lysozyme.
- Extrinsic eye muscles: move the eyeball within the orbit so that both eyes can be directed at the same object.

2.2 Refraction and Image Formation

Light from an object is bent (refracted) as it passes through the eye, mainly at two places:

1. the **cornea**, where light passes from air into the denser corneal tissue — this is where most of the refraction occurs, because the difference in density between air and cornea is large and the corneal curvature is fixed;
2. the **lens**, whose curvature is adjustable, allowing fine focusing of objects at different distances (this is **accommodation**, studied in Section 3).

The aqueous and vitreous humours transmit the light and maintain the shape of the eyeball but contribute little to refraction.

PROPERTY Nature of the Retinal Image

The cornea and lens together act as a converging lens system. For an object viewed by a normal (emmetropic) eye, the image formed on the retina is:

- Real — formed by the actual convergence of light rays;
- Inverted — upside-down and reversed left-to-right;
- Diminished — smaller than the object.

The brain interprets the inverted image so that we perceive the world the right way up.

WARNING The Inverted Image Trap

Common mistake: stating that the image on the retina is “upright” or “virtual”. Always remember: the retinal image is real, inverted and diminished. It is the brain that re-interprets it. If a GCE question asks “State the nature of the image formed on the retina”, the expected answer is exactly: real, inverted and diminished.

2.3 The Retina: Rods and Cones

The retina contains two types of photoreceptor cell, **rods** and **cones**, which convert light energy into nerve impulses. Their different properties explain why our vision in dim light differs from our vision in bright light.

Feature	Rods	Cones
Photopigment	Rhodopsin (visual purple)	Iodopsin (three forms, sensitive to blue, green or red light)
Sensitivity to light	Very high — function in dim light	Low — function only in bright light
Colour vision	None — give black-and-white vision	Yes — responsible for colour vision
Visual acuity	Low — many rods connect to one sensory neurone, so impulses are summed and detail is lost	High — each cone (at the fovea) has its own connection to the brain
Distribution	Absent from the fovea; concentrated in the peripheral retina	Concentrated at the fovea ; few in the periphery

DEFINITION Blind Spot (Optic Disc)

The **blind spot** is the region of the retina where the axons of the ganglion cells leave the eye as the optic nerve. It contains no rods and no cones, so light falling on it produces no sensation. We are normally unaware of it because the brain fills in the missing information and the two eyes cover each other's blind spot.

DID YOU KNOW? Demonstrating Your Own Blind Spot

Close your left eye. Stare at the cross below with your right eye only, and slowly bring the page closer to your face from about 30 cm. At a certain distance the dot disappears: its image has fallen on the blind spot of your right eye.



2.4 Binocular Vision

Humans have **binocular vision**: the two eyes face forward and view the same object from slightly different positions. The brain fuses the two slightly different images into a single three-dimensional perception (**stereoscopic vision**), which allows accurate judgement of distance and depth. This is a frequent GCE question: the advantages of binocular vision are **accurate depth/distance judgement** and **overlapping fields of view**, so the blind spot of one eye is covered by the other. Predators (lion, human) have forward-facing eyes for good depth judgement; prey animals (antelope, grasscutter) have eyes on the sides of the head giving a wide field of view to detect danger, but poorer depth perception.

2.5 Common Eye Defects and Their Correction

Refractive errors occur when the eyeball length and the refractive power of the cornea and lens do not match, so light is not focused exactly on the retina. They are corrected with spectacle lenses that shift the point of focus back onto the retina.

PROPERTY Myopia (Short-sightedness)

Cause: the eyeball is too long, or the cornea/lens refracts too strongly. Parallel rays from a distant object are focused in front of the retina, so distant objects appear blurred while near objects are seen clearly.

Correction: a **diverging (concave) lens** is placed in front of the eye. It spreads the incoming rays slightly so that the eye's own optics bring them to a focus exactly on the retina.

PROPERTY Hypermetropia (Long-sightedness)


Cause: the eyeball is too short, or the cornea/lens refracts too weakly. Rays from a near object would be focused behind the retina, so near objects appear blurred while distant objects may be seen clearly.

Correction: a **converging (convex) lens** is placed in front of the eye. It begins to converge the rays before they enter the eye, so that they are focused exactly on the retina.


PROPERTY Astigmatism

Cause: the cornea (or lens) is unevenly curved — more curved in one plane than in the perpendicular plane — so rays in different planes are focused at different points and the image is distorted.


Correction: a **cylindrical lens** whose curvature compensates for the uneven curvature of the cornea.



Normal eye
focus on retina




Myopia
focus in front of retina




Hypermetropia
focus behind retina

Focusing in the normal eye, in myopia (image forms in front of the retina) and in hypermetropia (image would form behind the retina; dashed lines show the rays continuing to their uncorrected focal point).



Myopia corrected



Hypermetropia corrected

Correction of refractive errors: a diverging (concave) lens corrects myopia; a converging (convex) lens corrects hypermetropia. In each case the focal point is shifted back onto the retina.

WORKED EXAMPLE 2 Identifying and Correcting a Defect

A student can read a book held at 25 cm clearly, but the writing on the classroom board is blurred. (a) Name the defect. (b) Explain its cause. (c) State the type of lens needed for correction and justify your answer.

SOLUTION

- (a) The defect is myopia (short-sightedness): near objects are seen clearly but distant objects are blurred.
- (b) The eyeball is too long (or the cornea and lens refract too strongly), so parallel rays from the distant board are brought to a focus in front of the retina. The retina therefore receives a blurred image.
- (c) A diverging (concave) lens is required. It diverges the parallel rays slightly before they enter the eye, so that the eye's own converging system now focuses them exactly on the retina, restoring clear distance vision.

WORKED EXAMPLE 3 Identifying and Correcting Long Sight

An elderly farmer in Buea must hold his newspaper at arm's length (about 100 cm) to read it, although his distance vision is good. (a) Name the defect. (b) Explain its cause. (c) State the correcting lens.

SOLUTION

- (a) Hypermetropia (long-sightedness) — in an older person this is specifically presbyopia, long sight due to loss of lens elasticity with age.
- (b) The eyeball is too short, or (in presbyopia) the lens can no longer become sufficiently rounded because it has lost elasticity and the ciliary muscles are weaker. Rays from near objects are therefore focused behind the retina.
- (c) A converging (convex) lens, which pre-converges the rays so that they focus on the retina.

KEY POINTS Key Points for GCE Revision

- The eye wall has three layers: sclera (protection and shape), choroid (nutrition and absorption of stray light), retina (contains the photoreceptors).
- The cornea provides most of the refraction; the lens provides the adjustable refraction used in accommodation.
- The retinal image is real, inverted and diminished.
- Rods: very sensitive, dim-light (black-and-white) vision, low acuity, peripheral retina. Cones: bright-light and colour vision, high acuity, concentrated at the fovea.
- Fovea = region of sharpest vision (cones only, one-to-one connections). Blind spot = exit point of the optic nerve (no photoreceptors).
- Myopia: image in front of retina → concave (diverging) lens. Hypermetropia: image behind retina → convex (converging) lens. Astigmatism: uneven corneal curvature → cylindrical lens.
- Binocular vision gives stereoscopic (3-D) vision and accurate judgement of distance.

EXERCISE 2 GCE-Style Practice

1. Draw a large, labelled diagram of a section through the mammalian eye. On your diagram, show the path of two light rays from a distant object to the retina. [8 marks]
2. State two differences between rods and cones, and explain how each difference is

related to the function of the cell. [6 marks]

3. A student cannot see objects clearly unless they are closer than 30 cm. Name the defect, explain its cause, and state the type of lens needed for correction, justifying your answer. [5 marks]
4. Explain why the image formed on the retina is inverted, and how we nevertheless perceive objects the right way up. [3 marks]
5. Describe the distribution of rods and cones in the retina and relate this distribution to visual acuity and colour vision. [4 marks]

SOLUTION Exercise 1

Diagram: large outline with anterior corneal bulge; three layers at the back (sclera outermost, choroid, retina innermost); cornea continuous with the sclera; biconvex lens suspended by suspensory ligaments from the ciliary body; iris with central pupil; optic nerve leaving at the blind spot; fovea marked on the visual axis. Two parallel rays from a distant object enter through the cornea and pupil, are refracted by the cornea and lens, cross over, and converge to a point on the retina (inverted image). Labels: sclera, choroid, retina, cornea, iris, pupil, lens, ciliary body, suspensory ligaments, aqueous humour, vitreous humour, optic nerve, blind spot, fovea. Marks awarded for accuracy of proportions, correct relative positions, complete labels and correct light path.

SOLUTION Exercise 2

- Photopigment: rods contain rhodopsin, which is very sensitive to light and is bleached in bright light → rods function in dim light. Cones contain iodopsin (in three forms) which is less sensitive → cones function in bright light and the three forms make colour vision possible.
- Connections: many rods connect to a single sensory neurone → their impulses are summed, giving high sensitivity but poor detail (low acuity). Each cone at the fovea connects to its own neurone → no summation, so fine detail is resolved (high acuity).

(Also acceptable: distribution — rods peripheral, cones at fovea — related to where acuity and colour vision are best.)

SOLUTION Exercise 3

Defect: myopia (short-sightedness). Cause: the eyeball is too long or the cornea/lens refracts too strongly, so light from distant objects is focused in front of the retina; only near objects (closer than 30 cm) focus on the retina. Correction: a diverging (concave) lens, which spreads the incoming rays so that the eye focuses them on the retina.

SOLUTION Exercise 4

The cornea and lens form a converging lens system. Rays from the top of an object are refracted to the bottom of the retina, and rays from the bottom of the object to the top of the retina; the rays cross over, so the image is inverted (and diminished). The brain receives the impulses from the retina and, through experience and coordination with the other senses, interprets the image so that the object is perceived the right way up.

SOLUTION Exercise 5



Cones are concentrated at the fovea, are few in the peripheral retina and absent from the blind spot. Rods are absent from the fovea and are concentrated in the peripheral retina. Visual acuity is therefore greatest at the fovea (high cone density and one-to-one neurone connections) and decreases towards the periphery (rod dominance and many-to-one connections). Colour vision occurs only where cones are present (fovea and central retina); peripheral vision, mediated by rods, is in black and white.

3 CONTROL OF LIGHT INTENSITY, ADAPTATION AND ACCOMMODATION

Having examined the gross structure of the eye and the physics of image formation on the retina, we now turn to the dynamic physiological mechanisms that ensure the image is not only formed, but formed clearly and under optimal illumination. The eye is not a static camera; it actively adjusts its aperture (the pupil), its sensitivity (retinal adaptation), and its focal length (accommodation) to cope with the enormous range of light intensities and object distances encountered in daily life — from the intense equatorial noon sun over the Adamawa plateau to the dim light of a forest understory on Mount Cameroon.

3.1 The Iris Reflex: Controlling Light Entry

The amount of light entering the eye is regulated by the diameter of the pupil, which is controlled by two sets of smooth muscles in the iris: the **circular muscles** (sphincter pupillae) and the **radial muscles** (dilator pupillae). These muscles are antagonistic effectors of the autonomic nervous system.



Antagonistic action of iris muscles. In bright light (left), parasympathetic stimulation contracts circular muscles → pupil constricts. In dim light (right), sympathetic stimulation contracts radial muscles → pupil dilates.

WARNING The Bright Light Trap

Never invert the muscle actions! In bright light, the circular muscles contract (parasympathetic) → pupil constricts (miosis). In dim light, the radial muscles contract (sympathetic) → pupil dilates (mydriasis). A common GCE error is stating that radial muscles contract in bright light. Remember: "Bright → Constrict → Circular".

Nervous Pathway of the Pupil Reflex (Autonomic Control) The pupil reflex is a **reflex arc** involving the autonomic nervous system. It is consensual: shining light in one eye causes constriction in both eyes.

1. **Receptor:** Photoreceptors (rods/cones) in the retina detect light intensity.
2. **Sensory Neurone:** Optic nerve (CN II) carries impulses to the **pretectal nucleus** in the mid-brain (not the visual cortex — this is a reflex, not conscious perception).
3. **Interneurones:** Connect pretectal nucleus to the **Edinger–Westphal nuclei** (parasympathetic component of oculomotor nerve nuclei) on both sides (decussation via posterior commissure).
4. **Motor Neurone (Parasympathetic):** Preganglionic fibres run in the **oculomotor nerve (CN III)** → synapse in the **ciliary ganglion** → postganglionic **short ciliary nerves** → **circular muscles** contract → pupil constricts.
5. **Sympathetic Pathway (Dim Light):** Hypothalamus → spinal cord (T1) → superior cervical ganglion → long ciliary nerves → **radial muscles** contract → pupil dilates.

DEFINITION Pupil Reflex (Light Reflex)

A protective autonomic reflex that adjusts pupil diameter in response to light intensity falling on the retina, preventing photobleaching of photoreceptors and improving depth of field. It is mediated by the parasympathetic division (constriction) and sympathetic

division (dilation) of the autonomic nervous system.

3.2 Light and Dark Adaptation: The Chemistry of Vision

While the iris reflex adjusts light entry rapidly (seconds), the retina itself adjusts its sensitivity over a much wider range (orders of magnitude) through the biochemistry of photopigments. This is **retinal adaptation**.

PROPERTY Rhodopsin Cycle: Bleaching and Regeneration

Rhodopsin (Visual Purple) is the photopigment in rod cells. It consists of the protein **opsin** bound to the aldehyde of vitamin A (**retinal**, specifically 11-cis-retinal).

- In Bright Light (Bleaching): Photons isomerise 11-cis-retinal to all-trans-retinal. This conformational change activates opsin → triggers transduction cascade → rhodopsin splits into opsin + all-trans-retinal (bleached, colourless). Sensitivity decreases.
- In Darkness (Regeneration): All-trans-retinal is isomerised back to 11-cis-retinal (requiring ATP) in the retinal pigment epithelium (RPE). It recombines with opsin to reform rhodopsin (purple). Sensitivity increases.

Vitamin A deficiency → insufficient retinal → impaired rhodopsin regeneration → **night blindness** (nyctalopia). Examinable: Descriptive knowledge of the rhodopsin cycle and the role of vitamin A is required; no formal proof of the transduction cascade is expected.

DID YOU KNOW? Vitamin A and Public Health in Cameroon

Night blindness due to vitamin A deficiency remains a public health concern in parts of Cameroon, particularly in the Far North and among children with limited dietary diversity (low intake of palm oil, mangoes, liver, eggs). The National Vitamin A Supplementation Programme targets children 6–59 months and postpartum women. Understanding the rhodopsin cycle explains why supplementation restores night vision: it provides the retinal precursor for rhodopsin regeneration.

DEFINITION Light Adaptation & Dark Adaptation

- Light Adaptation: Rapid process (seconds to minutes) where sensitivity decreases. Cones take over (photopic vision); rhodopsin is bleached faster than it regenerates. Pupil constricts.
- Dark Adaptation: Slow process (20–30 minutes for full rod sensitivity). Rods take

over (scotopic vision); rhodopsin regenerates in the dark. Pupil dilates. The Purkinje shift occurs: peak spectral sensitivity shifts from ≈ 555 nm (cones, green-yellow) to ≈ 505 nm (rods, blue-green).

EXAMPLE Walking from Sunlight into a Dark Room

Imagine leaving a brightly lit classroom in Buea at noon and entering a dark storeroom.

1. 0–5 sec: Pupils dilate rapidly (sympathetic iris reflex) — immediate but limited gain ($\times 16$ area).
2. 5 sec – 5 min: Cone sensitivity recovers (fast). You start seeing large shapes.
3. 5–30 min: Rod sensitivity recovers (slow). Rhodopsin regenerates. Sensitivity increases up to $\times 10^5$. You see fine detail in dim light.
4. Re-entering sunlight: Intense light bleaches rhodopsin almost instantly — glare, temporary blindness until cones dominate and pupil constricts.

3.3 Accommodation: Focusing for Near and Far

The iris controls quantity of light; accommodation controls quality (sharpness) by changing the focal length of the eye. The cornea provides $\approx 2/3$ of the total refractive power (fixed), while the **crystalline lens** provides the remaining $\approx 1/3$ (variable). Accommodation is the process by which the lens changes shape to focus objects at different distances onto the retina.



Mechanism of accommodation. Left (Distant): Ciliary muscle relaxed → ligaments taut → lens flat. Right (Near): Ciliary muscle contracts → ligaments slack → lens elastic recoil makes it round.

DEFINITION Accommodation

The process by which the eye changes the focal length of the crystalline lens to form a

sharp image on the retina for objects at different distances. It is achieved by the contraction/relaxation of the ciliary muscle altering tension on the suspensory ligaments (zonular fibres), allowing the naturally elastic lens to change shape.

METHOD Step-by-Step Reasoning: Focusing a Near Object

To answer GCE questions on accommodation, follow this causal chain:

Step	Physiological Event
1	Object is near → light rays diverge strongly entering the eye.
2	Blurred image forms behind retina (hypermetropic blur detected).
3	Visual cortex → parasympathetic signal via CN III (oculomotor) to ciliary muscle.
4	Ciliary muscle contracts (circular fibres shorten).
5	Ciliary body moves forward and inward.
6	Tension on suspensory ligaments is released (slack).
7	Lens elasticity causes it to become more convex (rounder).
8	Refractive power of lens increases.
9	Diverging rays bent more sharply → image focused on retina.

Reverse every step for distant vision.

WARNING Presbyopia – The Ageing Lens

With age, the lens loses elasticity (denaturation of crystallin proteins) and the ciliary muscle weakens. The near point recedes (from ≈ 10 cm at age 10 to ≈ 100 cm at age 60). This is **presbyopia** ("old eye"). It is not a refractive error like myopia/hypermetropia; it is a loss of accommodative amplitude. Corrected by convex reading glasses (add power).

The Near Response Triad (Convergence Reflex) Accommodation for near objects is never isolated; it is part of a synergistic **near response triad**:

1. **Accommodation:** Lens rounds up (parasympathetic, CN III).
2. **Convergence:** Medial recti muscles contract → eyes rotate medially to fixate on the near object (somatic motor, CN III).
3. **Pupillary Constriction (Miosis):** Circular iris muscles contract → pupil constricts → increases depth of field and reduces spherical aberration (parasympathetic, CN III).

This is why your pupils constrict when you look at your phone screen close to your face, even if the room lighting hasn't changed.

WORKED EXAMPLE 4 Analysing the Near Response

Statement: A student reads a textbook held at 25 cm. Describe the changes in the eye and the nervous control involved.

SOLUTION

Detailed Solution:

1. Stimulus: Diverging light rays from near object; blur detected on retina (image would form behind retina).
2. Afferent Pathway: Optic nerve (CN II) → visual cortex → association areas.
3. Efferent Pathway (Parasympathetic): Edinger–Westphal nucleus → Oculomotor nerve (CN III) → Ciliary ganglion.
4. Effectors & Changes:
 - Ciliary muscle: Contracts → moves forward/inward.
 - Suspensory ligaments: Slacken.
 - Lens: Becomes more convex (increased curvature) → increased refractive power → image focused on retina.
 - Pupil: Circular muscles contract → pupil constricts (miosis) → increased depth of field, reduced aberrations.
 - Medial rectus muscles: Contract (somatic CN III) → eyes converge (medial rotation).
5. Result: Clear, single, binocular image of the text on both retinas.

3.4 Summary and Key Points

KEY POINTS Control of Light Intensity, Adaptation & Accommodation

- Iris Reflex: Bright light → Circular muscles contract (Parasympathetic, CN III) → Pupil constricts (Miosis). Dim light → Radial muscles contract (Sympathetic) → Pupil dilates (Mydriasis). Consensual.
- Light Adaptation: Fast (secs/mins). Cones function. Rhodopsin bleached > regenerated. Low sensitivity, high acuity, colour vision.
- Dark Adaptation: Slow (20–30 mins). Rods function. Rhodopsin regenerates (needs Vitamin A/Retinal). High sensitivity, low acuity, no colour (Purkinje shift).
- Accommodation: Near → Ciliary muscle contracts → Ligaments slack → Lens rounds

up (elastic recoil) → Power increases. Distant → Ciliary muscle relaxes → Ligaments taut → Lens flattens → Power decreases.

- Near Response Triad: Accommodation + Convergence + Pupillary Constriction (all Parasympathetic CN III).
- Presbyopia: Age-related loss of lens elasticity → reduced accommodation → corrected by convex lenses.

EXERCISE 3 GCE Style Practice

1. Explain why a person with Vitamin A deficiency experiences night blindness but can see normally in bright daylight.
2. Describe the sequence of events from the moment light from a near object hits the retina to the point where a sharp image is formed. Include the role of the ciliary muscle, suspensory ligaments, and lens elasticity.
3. A doctor shines a penlight into a patient's left eye and observes constriction of the right pupil. Name this reflex and explain the neural pathway responsible for the consensual response.
4. Distinguish between accommodation and adaptation. State two differences.

SOLUTION Exercise 1

Vitamin A (retinol) is the precursor for retinal (11-cis-retinal), the chromophore of rhodopsin. In deficiency, rhodopsin regeneration in darkness is impaired. Rods (responsible for scotopic/dim light vision) cannot recover sensitivity after bleaching. Cones (photopic/bright light vision) use different photopigments (photopsins) and function adequately in bright light where rhodopsin is mostly bleached anyway. Hence night blindness (rod failure) with normal day vision (cone function).

SOLUTION Exercise 2

1. Diverging rays from near object enter eye → image focused behind retina (blur). 2. Blur detected → visual cortex → parasympathetic outflow via CN III (oculomotor) to ciliary ganglion. 3. Postganglionic fibres stimulate ciliary muscle contraction. 4. Ciliary ring moves forward/inward → suspensory ligaments slacken. 5. Lens elasticity causes lens to become more convex (rounder). 6. Increased refractive power bends rays more sharply

→ image focused on retina.

SOLUTION Exercise 3

Consensual Pupil Light Reflex. Pathway: Light → Left Retina → Left Optic Nerve (CN II) → Optic Chiasm (partial decussation) → Left & Right Optic Tracts → Left & Right Pretectal Nuclei (Midbrain) → Interneurons cross via Posterior Commissure → Bilateral Edinger-Westphal Nuclei → Bilateral CN III (Parasympathetic) → Ciliary Ganglia → Short Ciliary Nerves → Circular Muscles of both irises contract → Bilateral Miosis.

SOLUTION Exercise 4

Feature	Accommodation	Adaptation (Retinal)
Mechanism	Physical change in lens shape (curvature)	Biochemical change in photopigment (rhodopsin) concentration/state
Structures	Ciliary muscle, suspensory ligaments, lens	Rods, cones, retinal pigment epithelium, Vitamin A
Stimulus	Object distance (blur/disparity)	Light intensity (photon flux)
Time Scale	Fast (milliseconds to seconds)	Light adapt: seconds; Dark adapt: 20–30 minutes
Purpose	Focus image on retina (sharpness)	Adjust retinal sensitivity (dynamic range)

4 PHOTORECEPTION AND THE EAR: HEARING AND BALANCE

In the previous section we saw how the eye controls the amount of light entering it (the iris reflex), how it adapts to light and darkness, and how it focuses images by accommodation. But forming a sharp image on the retina is only half the story. The retina must now convert light energy into electrical signals that the brain can interpret. This conversion is the work of the photoreceptor cells — the **rods** and **cones** — and it is the subject of **photoreception**. We shall then turn to a second sense organ, the **ear**, which performs two apparently different jobs with the same basic strategy: detecting sound (**hearing**) and detecting head position and movement (**balance**).


DEFINITION Photoreception

Photoreception is the process by which specialised receptor cells (photoreceptors) absorb light energy and convert it into electrical signals (receptor/generator potentials) that are transmitted to the central nervous system. In mammals the photoreceptors are the **rod cells** and **cone cells** of the retina. The conversion of one form of energy into an electrical signal by a receptor is called **transduction**.

4.1 Structure of Rod and Cone Cells

Rods and cones are modified neurones lying in the deepest layer of the retina (light must pass through the other retinal layers to reach them). Each cell has four recognisable regions:

- The **outer segment**: a stack of flattened membranous discs whose membranes carry the **visual pigments** (rhodopsin in rods, iodopsins in cones). This is where light is actually absorbed.
- A narrow connecting **cilium** linking the outer and inner segments.
- The **inner segment**: packed with mitochondria (to supply ATP for the active pumping of ions) and containing the nucleus.
- The **synaptic terminal** (synaptic body): forms synapses with **bipolar cells**, which in turn connect to **ganglion cells** whose axons form the optic nerve.



A rod cell (left) and the photochemical cycle of rhodopsin (right).

4.2 The Photochemistry of Vision

The key event in photoreception is a chemical one. The visual pigment of rods, **rhodopsin** (visual purple), consists of a protein, **opsin**, bound to a light-sensitive derivative of vitamin A, **retinal** (retinene), in the 11-cis form.

1. In the **dark**, rhodopsin is intact. The rod membrane maintains a dark current: sodium ions leak in through cGMP-gated channels, while the Na^+/K^+ ATPase actively pumps them out. The inside of the cell is partially negative (around -40 mV) and the rod continuously releases the neurotransmitter glutamate onto its bipolar cell.
2. When **light** strikes the outer segment, a photon is absorbed by retinal, which changes shape from the cis to the trans form and splits away from opsin — rhodopsin is said to be **bleached**.
3. Activated opsin (metarhodopsin II) triggers a G-protein (transducin) cascade that activates phosphodiesterase (PDE). PDE hydrolyses cGMP to GMP, causing the cGMP-gated Na^+ channels to close.
4. The Na^+/K^+ pump continues working, so the inside of the cell becomes more negative (hyperpolarises to about -70 mV). This change in membrane potential is the **generator potential** (receptor potential).
5. The generator potential reduces glutamate release, and this change is detected by the bipolar cell and relayed via ganglion cells along the optic nerve to the visual cortex.
6. In the dark, retinal is converted back to the cis form by retinal isomerase and recombines with opsin (using ATP): rhodopsin is **resynthesised**. This is the chemical basis of **dark adaptation** studied earlier.

WARNING A surprising sign

Unlike most receptors, rods hyperpolarise in response to their stimulus and release neurotransmitter in the dark. Do not write that light "depolarises" the rod — this is a classic error that loses marks.

4.3 Cones and Colour Vision

Cones contain pigments called **iodopsins**, built on the same retinal + opsin plan but with three slightly different opsins. This gives **three types of cone**, each most sensitive to a different region of the spectrum:

- **red-sensitive** cones (L-cones, peak absorption ≈ 560 nm, long wavelengths),
- **green-sensitive** cones (M-cones, peak ≈ 530 nm, medium wavelengths),
- **blue-sensitive** cones (S-cones, peak ≈ 420 nm, short wavelengths).

According to the **trichromatic theory** of colour vision, every colour we perceive results from the relative stimulation of these three cone types. For example, yellow light (≈ 580 nm) stimulates red and green cones roughly equally and blue cones weakly; the brain interprets this particular ratio as "yellow". White light stimulates all three strongly and about equally. A person lacking one functional cone type is **colour-blind** (e.g. red-green colour blindness, a sex-linked condition commoner in males).

4.4 Rods versus Cones

PROPERTY Sensitivity and acuity

Rods give high sensitivity in dim light (they contain rhodopsin, which is bleached even by weak light, and many rods converge onto one bipolar cell, so their signals summate) but low acuity and no colour vision. **Cones** need bright light but give high acuity (each cone in the fovea has its own bipolar cell, so signals are kept separate) and colour vision.

KEY POINTS Comparing rods and cones — a classic GCE table

Feature	Rods	Cones
Pigment	Rhodopsin (one type)	Iodopsins (three types: red, green, blue)
Sensitivity to light	High — work in dim light	Low — need bright light
Visual acuity	Low (many rods per bipolar cell: convergence)	High (one cone per bipolar cell at fovea)
Colour vision	No (black, white, grey)	Yes (trichromatic)
Distribution	Peripheral retina; absent from fovea	Concentrated at the fovea
Function	Night vision, movement detection	Daylight vision, fine detail, colour

EXAMPLE Why stars "disappear" when you look straight at them

Astronomers — and hunters on dark nights on the Adamawa plateau — know that a faint star is easier to see when you look slightly to one side of it. Looking directly at the star places its image on the **fovea**, which contains only cones, and cones are insensitive in dim light. Looking slightly aside moves the image onto the peripheral retina, rich in highly sensitive rods, so the star becomes visible.

4.5 Structure of the Ear

The ear is a double sense organ: the **cochlea** detects sound, while the **vestibular apparatus** (semicircular canals, utricle and saccule) detects head position and movement. Anatomically it has three parts.


The outer ear. The **pinna** (visible ear flap) collects sound waves and funnels them into the **auditory canal** (external auditory meatus), at the end of which lies the **eardrum** (tympanic membrane), a thin membrane that vibrates when sound waves strike it.

The middle ear. An air-filled cavity in the skull containing the three **ossicles**, the smallest bones

in the body: the **malleus** (hammer, attached to the eardrum), the **incus** (anvil) and the **stapes** (stirrup, whose footplate fits into the **oval window** of the cochlea). The **Eustachian tube** connects the middle ear to the pharynx and equalises air pressure on both sides of the eardrum — this is why swallowing relieves the “popping” of the ears when a bus climbs the steep road up Mount Cameroon.

The inner ear. A fluid-filled labyrinth in the temporal bone comprising:

- the **cochlea** — a spiral, snail-shaped tube containing the sensory cells of hearing;
- the three **semicircular canals** — arranged at right angles to one another, detecting rotational movement;
- the **utricle** and **sacculle** — detecting gravity and linear acceleration.



Section through the human ear showing the outer, middle and inner ear.

4.6 The Mechanism of Hearing

Hearing is a chain of energy conversions: sound waves (mechanical energy in air) → mechanical vibrations of bones → pressure waves in fluid → electrical impulses in the auditory nerve. Follow the sequence carefully:

METHOD From sound wave to nerve impulse

1. Sound waves are collected by the pinna and channelled along the auditory canal to the eardrum, which vibrates at the same frequency as the sound.
2. The vibrations pass across the middle ear through the ossicles (malleus → incus → stapes). The ossicles act as levers and the stapes footplate is much smaller than the eardrum (area ratio $\approx 17 : 1$), so the force is concentrated: vibrations are amplified (about twenty-fold in pressure) to move the dense inner-ear fluid efficiently.
3. The stapes pushes on the oval window, setting up pressure waves in the perilymph, the fluid filling the scala vestibuli and scala tympani of the cochlea.

4. The waves travel along the cochlea and make the basilar membrane vibrate. Different frequencies vibrate different regions of the membrane (high pitch near the base, low pitch near the apex) — this **place theory** (tonotopic organisation) is how pitch is discriminated.
5. Sitting on the basilar membrane is the organ of Corti, whose hair cells have sensory hairs (stereocilia) embedded near the overlying tectorial membrane. Vibration of the basilar membrane causes a shearing force that bends the stereocilia.
6. Bending opens mechanically-gated K^+ channels in the hair cells (endolymph is high in K^+), generating a receptor potential that triggers nerve impulses in fibres of the auditory nerve, which carries them to the auditory cortex of the brain, where they are interpreted as sound.

DEFINITION The organ of Corti

The **organ of Corti** is the sensory structure of hearing, located on the basilar membrane inside the cochlear duct (scala media). It consists of **inner hair cells** (true sensory receptors) and **outer hair cells** (amplifiers), whose stereocilia project towards the tectorial membrane, together with supporting cells. Bending of the stereocilia by vibrations of the basilar membrane generates receptor potentials that excite the auditory nerve.

WARNING The ossicles do not detect sound

A frequent mistake is to say that the ossicles "receive" or "detect" sound. They do not: the ossicles merely transmit and amplify vibrations from the eardrum to the oval window. Detection (transduction into electrical signals) occurs only in the hair cells of the organ of Corti. Similarly, the eardrum vibrates but detects nothing.

4.7 Balance: The Vestibular Apparatus

Balance uses the same transduction principle as hearing — bending of hair cell stereocilia — but the stimulus is movement of the head rather than sound.

Rotational movement: the semicircular canals. The three semicircular canals lie in three mutually perpendicular planes (horizontal, anterior vertical, posterior vertical). At the base of each canal is a swelling, the **ampulla**, containing a patch of hair cells (the **crista ampullaris**) whose stereocilia are embedded in a jelly-like cup, the **cupula**. When the head rotates, the fluid (endolymph) inside the canal lags behind because of inertia, pushing the cupula and bending the stereocilia. Because the three canals are at right angles, rotation in any plane is detected. This is why spinning round and then stopping makes you dizzy: the fluid keeps moving briefly, bending the cupula even though your head has stopped.

Gravity and linear acceleration: the utricle and saccule. These two chambers contain hair cells (the **maculae**) overlain by a jelly layer loaded with tiny calcium carbonate crystals, the **otoliths** (otoconia). When the head tilts or accelerates in a straight line (in a lift, or in a bendskin pulling away), gravity or inertia shifts the heavy otoliths, bending the stereocilia. The pattern of bending tells the brain the orientation of the head relative to gravity and any linear acceleration.

Impulses from both systems travel along the **vestibular nerve** (part of the vestibulocochlear nerve, CN VIII) to the **cerebellum** and vestibular nuclei, which coordinate posture and eye movements (vestibulo-ocular reflex) to keep us balanced.

WORKED EXAMPLE 5 Explaining sensitivity and acuity in the retina

Statement. (a) Explain why rods allow us to see in dim light. (b) Explain why cones, but not rods, give high visual acuity. (c) A patient cannot distinguish red from green in bright light but sees normally at night. Suggest an explanation.

SOLUTION

Solution.

1. (a) Rods contain rhodopsin, which is bleached even by very weak light, so a single photon can contribute to a response. In addition, many rods converge onto a single bipolar cell (**retinal convergence**); their small generator potentials summate, so a threshold is reached even in dim light. Hence rods are highly sensitive.
2. (b) At the fovea each cone has its own bipolar cell and ganglion cell (no convergence). Stimulation of two neighbouring cones therefore sends two separate signals to the brain, which can distinguish two close points as distinct — this is high acuity. In rods, convergence pools signals from many cells, so the brain cannot tell which rod was stimulated: acuity is poor.
3. (c) Night vision depends on rods, which are normal; colour vision depends on cones. The patient most likely lacks functional red-sensitive or green-sensitive cones (red-green colour blindness), a defect of cone pigments that leaves rod function intact.

WORKED EXAMPLE 6 Mechanism of hearing and balance

Statement. (a) Describe how the ossicles amplify sound vibrations. (b) Explain how the organ of Corti converts vibrations into nerve impulses. (c) A person spins rapidly and then stops suddenly. Explain why they feel dizzy.

SOLUTION

Solution.

1. (a) The ossicles act as a lever system: the malleus (attached to the eardrum) moves a greater distance than the stapes (attached to the oval window). Combined with the

area ratio of eardrum to stapes footplate ($\approx 17 : 1$), this amplifies the pressure about 20-fold, overcoming the impedance mismatch between air and cochlear fluid.

2. (b) Vibrations of the stapes create pressure waves in perilymph, which travel through the cochlea and displace the basilar membrane. This shears the stereocilia of inner hair cells against the tectorial membrane. Bending opens K^+ channels (endolymph is K^+ -rich), causing K^+ influx, depolarisation (receptor potential), Ca^{2+} entry, and glutamate release onto auditory nerve fibres, generating action potentials.
3. (c) During spinning, endolymph in the horizontal semicircular canal accelerates due to friction with the canal walls. When rotation stops, the endolymph continues moving (inertia), deflecting the cupula in the opposite direction. This signals continued rotation to the brain, conflicting with visual and proprioceptive input, causing dizziness (vertigo).

EXERCISE 4 Check your understanding

1. List, in order, the structures through which a vibration passes from the air to the auditory nerve.
2. Explain why the ears "pop" when altitude changes rapidly, and name the structure responsible.
3. State one structural difference and one functional difference between the semicircular canals and the utricle.
4. Explain why a driver at night sees a pedestrian at the roadside more easily in peripheral vision than by looking directly at them.
5. Describe the role of the tectorial membrane in hearing.
6. What is the function of the round window in the cochlea?

SOLUTION Exercise 1

1. Air (auditory canal) → eardrum → malleus → incus → stapes → oval window → perilymph of scala vestibuli → basilar membrane → hair cells of the organ of Corti → auditory nerve.
2. A rapid altitude change makes the air pressure outside the eardrum differ from that in the middle ear, bulging the eardrum. Swallowing opens the **Eustachian tube**, equalising the pressure — the "pop".

3. Structural: semicircular canals are three looped tubes with ampullae containing a cupula; the utricle is a chamber whose hair cells bear otoliths. Functional: semicircular canals detect rotational movements of the head; the utricle detects gravity and linear acceleration.
4. Looking directly at the pedestrian focuses the image on the fovea, which has only cones; cones are insensitive in dim light. Peripheral vision uses rod-rich retina, and rods are highly sensitive to dim light, so the pedestrian is detected more easily.
5. The tectorial membrane overlies the hair cells; as the basilar membrane vibrates, the hair cell stereocilia shear against the tectorial membrane, bending them and opening transduction channels.
6. The round window (covered by the secondary tympanic membrane) bulges out as the stapes pushes in at the oval window, allowing the incompressible perilymph to move and enabling basilar membrane vibration.

DID YOU KNOW? Hearing loss in Cameroon

Prolonged exposure to very loud sound — for example in unprotected work near grinding machines or with powerful sound systems at makossa night-clubs — progressively destroys the hair cells of the organ of Corti. Unlike skin cells, human hair cells do not regenerate: the resulting deafness is permanent. Protecting your ears is protecting an irreplaceable sense organ. The World Health Organization estimates that over 1 billion young people worldwide are at risk of hearing loss due to unsafe listening practices.

KEY POINTS Key points of this section

- Photoreception is the transduction of light into electrical signals by rods and cones; light bleaches rhodopsin (retinal + opsin), closing Na^+ channels, causing hyperpolarisation — the generator potential.
- Rods: high sensitivity, low acuity, no colour, peripheral retina. Cones: low sensitivity, high acuity, colour (three types — trichromatic theory), concentrated at the fovea.
- The ear has three parts: outer (pinna, canal, eardrum), middle (ossicles + Eustachian tube), inner (cochlea, semicircular canals, utricle, saccule).
- Hearing: ossicles amplify (lever + area ratio); only the hair cells of the organ of Corti transduce vibrations into impulses via K^+ influx from endolymph.
- Balance: semicircular canals detect rotation (cupula deflection by endolymph inertia); utricle and saccule detect gravity and linear acceleration (otolith displacement).

5 THE NERVOUS SYSTEM AND NEURONES

In the previous section we saw how the eye converts light into electrical signals and how the ear converts sound waves and head movements into impulses. But detection is only the beginning of the story. Once a receptor in the retina or the cochlea has been stimulated, the message must travel rapidly to a coordinating centre, be compared with other incoming information, and an appropriate command must be sent back to muscles or glands. That entire task — reception, transport, processing and command — belongs to the **nervous system**. In this section we study how the nervous system is organised, the structure of its cellular unit, the **neurone**, and how the central nervous system integrates information.

5.1 Organisation of the Nervous System

Imagine the administration of Cameroon: ministries in Yaoundé make decisions, but they are connected to every division and village by roads and telephone lines. The nervous system is organised in exactly the same way, with a central command and a vast communication network reaching every part of the body.

DEFINITION Divisions of the nervous system

The nervous system is divided into two parts:


- The **central nervous system** (CNS), made of the brain (protected inside the skull) and the spinal cord (protected inside the vertebral column). It is the coordinating and decision-making centre.
- The **peripheral nervous system** (PNS), made of all the nerves connecting the CNS to the rest of the body: cranial nerves (arising from the brain, e.g. the optic nerve from the eye, the auditory nerve from the ear) and spinal nerves (arising from the spinal cord, one pair per segment, serving the trunk and limbs).

The PNS itself contains two functional categories of fibres: **afferent (sensory) fibres**, which carry impulses from receptors to the CNS, and **efferent (motor) fibres**, which carry commands from the CNS to effectors (muscles and glands). A useful memory trick: **A**fferent = **A**rriving at the CNS; **E**fferent = **E**xiting the CNS.

5.2 The Neurone: Unit of the Nervous System

DEFINITION Neurone and nerve

A **neurone** (nerve cell) is a specialised cell adapted for the rapid reception and transmission of electrical signals called nerve impulses. A **nerve** is a bundle of many neurone fibres (axons) held together by connective tissue, like many electric wires inside one cable.



Structure of a motor neurone. Impulses enter through the dendrites, pass through the cell body and travel along the myelinated axon to the axon terminals.

Parts of a motor neurone and their roles:

- **Dendrites:** highly branched extensions that receive impulses from receptors or from other neurones and conduct them towards the cell body.
- **Cell body (soma):** contains the **nucleus** and abundant cytoplasm with mitochondria, rough endoplasmic reticulum and Golgi apparatus; it controls the cell's metabolism and protein synthesis (including neurotransmitter synthesis).
- **Axon:** a single long fibre that conducts impulses away from the cell body towards the effector or the next neurone. Motor axons can be remarkably long — the axon controlling the toes runs the whole length of the leg.
- **Myelin sheath:** a fatty insulating layer produced by **Schwann cells** in the PNS and by **oligodendrocytes** in the CNS, wrapped around the axon.
- **Nodes of Ranvier:** short gaps (about $1\ \mu\text{m}$) between successive myelinating cells where the axon membrane is exposed and voltage-gated ion channels are concentrated.
- **Axon terminals (synaptic knobs):** branched endings that pass the signal to the next cell across a synapse; they contain mitochondria and synaptic vesicles filled with neurotransmitter.

WARNING Direction of conduction


A very common examination error is to reverse the direction of travel. Remember: dendrites carry impulses towards the cell body; the axon carries them away from the cell body. A neurone is a one-way street: dendrite → cell body → axon → terminals.

5.3 Types of Neurones

Three functional types of neurone cooperate in every nervous pathway.

DEFINITION The three types of neurones

- **Sensory (afferent) neurone:** carries impulses from a receptor to the CNS. It is typically pseudo-unipolar: a single process emerges from the cell body and divides into a peripheral branch (to the receptor) and a central branch (to the CNS). Both branches are myelinated. The cell body lies in a dorsal root ganglion, just outside the spinal cord, on a side branch of the fibre.
- **Relay (intermediate / association) neurone:** lies entirely within the CNS; it connects sensory neurones to motor neurones and to other relay neurones. It is short, with many dendrites and a short axon; usually unmyelinated.
- **Motor (efferent) neurone:** carries impulses from the CNS to an effector (muscle or gland). Its cell body lies inside the CNS (in the grey matter of the spinal cord or brain), and it has a long myelinated axon (in the PNS, myelinated by Schwann cells).



Comparison of the three neurone types. Note the position of the cell body: in a ganglion outside the CNS (sensory), inside the CNS with a short fibre (relay), and inside the CNS with a long axon (motor). The sensory neurone is pseudo-unipolar: one process splits into peripheral and central myelinated branches.

METHOD Identifying a neurone type from a diagram

When the GCE paper shows you an unlabelled neurone, proceed in order:

1. Locate the cell body. On a side branch outside the CNS $\Rightarrow$ sensory neurone. At one end of the cell, inside the CNS $\Rightarrow$ motor neurone.
2. Look at the length and branching. A short cell with many dendrites and no long axon, entirely within the CNS $\Rightarrow$ relay neurone.
3. Follow the connections. Attached to a receptor $\Rightarrow$ sensory; attached to a muscle or gland $\Rightarrow$ motor; between the two $\Rightarrow$ relay.

4. Check the direction arrows if given: towards the CNS = sensory; away from the CNS = motor.

5.4 The Myelin Sheath and Saltatory Conduction

Why wrap an axon in fat? For the same reason that electric wires are wrapped in plastic: insulation prevents the signal from leaking away. But myelin does more than insulate — it makes conduction dramatically faster.

DEFINITION Myelin sheath

The **myelin sheath** is a fatty, electrically insulating layer formed by Schwann cells (in the PNS) or oligodendrocytes (in the CNS) wound many times around the axon of many neurones. It is interrupted at regular intervals by the nodes of Ranvier, where the axon membrane is exposed and voltage-gated Na^+ and K^+ channels are highly concentrated.

PROPERTY Saltatory conduction

In a myelinated axon, the local circuits of current that transmit the impulse cannot cross the insulated regions; they act only at the nodes of Ranvier. The impulse therefore jumps from node to node — this is **saltatory conduction** (from Latin saltare, to jump). Consequences:

- Transmission is much faster than in a non-myelinated axon of the same diameter (up to about 100 m s^{-1} in large myelinated fibres, against less than 1 m s^{-1} in small unmyelinated fibres).
- Less energy is needed, because membrane ion pumps work only at the nodes to restore ionic gradients after each impulse.

(The mechanism of the impulse itself is treated in the next section; the jumping property is examinable here.)

DID YOU KNOW? When myelin fails

In the disease multiple sclerosis, the patient's own immune system destroys the myelin sheath in the CNS. Saltatory conduction then fails, impulses slow down or stop, and the person progressively loses control of movement, vision and balance. This illustrates how essential myelin is for rapid, reliable coordination.

5.5 Nerves, Grey Matter and White Matter

A single neurone is microscopic, yet you can see a nerve with the naked eye. That is because a **nerve** is a bundle of hundreds or thousands of axons wrapped together by connective tissue (epineurium, perineurium, endoneurium) — exactly like a telephone cable containing many individual wires. Most nerves are **mixed nerves**, containing both sensory and motor fibres.

Inside the CNS, two kinds of tissue are visible in a cross-section of the spinal cord or brain:

Feature	Grey matter	White matter
Main content	Cell bodies, dendrites, synapses, unmyelinated axons	Myelinated axons (nerve fibres)
Colour origin	Unmyelinated tissue (greyish)	White fatty myelin
Position in spinal cord	Central (butterfly / H-shape)	Surrounding the grey matter
Position in brain	Outer layer (cortex) and deep nuclei	Beneath the cortex
Function	Integration, processing, synaptic contacts	Rapid transmission between regions

5.6 Integration: The CNS as a Decision Centre

At every instant, thousands of receptors — in your eyes, ears, skin, muscles and internal organs — are sending impulses into the CNS. If every message triggered an automatic response, behaviour would be chaotic. The CNS must therefore select, compare, combine and prioritise information before commanding effectors.

DEFINITION Integration

Integration is the process by which the central nervous system receives impulses from many receptors, processes and coordinates this information (including comparison with stored memory), and sends out appropriate commands to effectors so that the response is adapted to the whole situation.

EXAMPLE Integration in everyday life

You are walking along a road in Bamenda and you hear a motorcycle horn behind you (auditory receptors), you see it approaching in your mirror (photoreceptors), and you feel the weight of your school bag (proprioceptors). Your CNS combines all three messages and commands your leg muscles to step aside while keeping your balance. No single reflex could achieve this coordinated, situation-dependent response — it requires integration.

Synapses as decision points. Integration is physically carried out at **synapses**, the junctions

between neurones. A single relay neurone in the spinal cord may receive synapses from hundreds of other neurones, some excitatory (producing EPSPs, pushing it towards firing) and some inhibitory (producing IPSPs, holding it back). Whether the relay neurone fires depends on the sum of all these influences — **spatial summation** (simultaneous inputs from different synapses) and **temporal summation** (rapid successive inputs at the same synapse). Each synapse is therefore a microscopic decision point, and the billions of synapses of the brain together constitute the integrating machinery. We will study synaptic transmission in detail in the next section.

WORKED EXAMPLE 7 Identifying neurones and tracing a pathway

The diagram of a spinal reflex shows three neurones, A, B and C. Neurone A has its cell body in a swelling (ganglion) on the dorsal root and its ending near the skin. Neurone B is short and lies entirely within the grey matter of the spinal cord. Neurone C has its cell body in the grey matter and its long myelinated axon ends on a biceps muscle. (a) Identify each neurone. (b) State the direction of the impulse through the pathway. (c) Explain why the response of the muscle is rapid.

SOLUTION

(a) A is the sensory neurone — its cell body sits in the dorsal root ganglion outside the CNS and it collects impulses from a receptor in the skin. B is the relay (intermediate) neurone — short, entirely within the CNS. C is the motor neurone — cell body in the grey matter of the CNS, long axon running to an effector (the biceps muscle).

(b) Impulse pathway: receptor (skin) → A (sensory neurone) → B (relay neurone) → C (motor neurone) → effector (biceps), which contracts.

(c) The response is rapid because (i) nerve impulses are electrical and travel fast, (ii) the motor axon is myelinated, so conduction is saltatory — the impulse jumps between nodes of Ranvier instead of travelling continuously, and (iii) the pathway is short, with few synapses.

WARNING Frequent confusions to avoid

- Do not confuse a neurone (one cell) with a nerve (a bundle of many axons). Examiners penalise answers that use them interchangeably.
- The cell body of the sensory neurone is outside the CNS (dorsal root ganglion); that of the motor neurone is inside the CNS. Reversing these is a classic error.
- Grey matter is not “grey neurones”: it is the region rich in cell bodies and synapses; white matter is the region rich in myelinated axons.
- Myelin does not “create” the impulse; it insulates the axon and accelerates conduction by saltatory jumping.
- Sensory neurones are pseudo-unipolar, not bipolar: a single process divides into two

branches. Do not draw two separate processes emerging from the cell body.

KEY POINTS Key points of this section

- The nervous system = CNS (brain + spinal cord) + PNS (cranial and spinal nerves).
- The neurone is the functional unit: dendrites → cell body → axon → terminals.
- Three types: sensory (receptor → CNS, cell body in ganglion, pseudo-unipolar), relay (within CNS, short, unmyelinated), motor (CNS → effector, cell body in CNS, long myelinated axon).
- The myelin sheath (Schwann cells in PNS, oligodendrocytes in CNS) insulates the axon and allows fast saltatory conduction between nodes of Ranvier.
- A nerve is a bundle of axons wrapped in connective tissue; grey matter = cell bodies and synapses, white matter = myelinated fibres.
- Integration = reception, processing and coordination of information by the CNS, achieved largely at synapses via spatial and temporal summation of EPSPs and IPSPs.

EXERCISE 5 Check your understanding

1. Draw and fully label a motor neurone, and add an arrow showing the direction of the impulse.
2. State two structural differences between a sensory neurone and a motor neurone.
3. Explain why an impulse travels faster in a myelinated axon than in a non-myelinated axon of the same diameter.
4. A cross-section of the spinal cord shows a darker central H-shaped region and a paler outer region. Name each region and state what it contains.
5. Using the example of touching a hot cooking pot, explain what is meant by integration.
6. Distinguish between spatial summation and temporal summation at a synapse.

SOLUTION Exercise 1

1. Diagram as in the figure of this section: dendrites, cell body with nucleus, axon, myelin sheath with Schwann cells, nodes of Ranvier, axon terminals; arrow pointing from den-

drites towards terminals.

2. Sensory neurone: cell body in a dorsal root ganglion outside the CNS, on a side branch; pseudo-unipolar (one process splits into peripheral and central branches); carries impulses from receptor to CNS. Motor neurone: cell body inside the CNS at the end of the cell; multipolar (many dendrites, one long axon); carries impulses to an effector.

3. Myelin insulates the axon so that local currents act only at the nodes of Ranvier; the impulse jumps from node to node (saltatory conduction), which is faster than continuous conduction along the whole membrane. Voltage-gated channels are concentrated at nodes.

4. Central H-shaped region = grey matter (cell bodies, dendrites, synapses, unmyelinated axons); outer pale region = white matter (myelinated axons running up and down the cord).

5. Touching the hot pot stimulates pain and temperature receptors; sensory neurones carry impulses to the CNS, which processes the information (identifying danger, comparing with experience) and coordinates a command via motor neurones to the arm muscles to withdraw the hand — and possibly to other muscles to keep balance. This selection and coordination of an appropriate response by the CNS is integration.

6. Spatial summation: simultaneous EPSPs from several different presynaptic neurones add together to reach threshold. Temporal summation: rapid successive EPSPs from the same presynaptic neurone add together because the previous depolarisation has not yet decayed.

6 TRANSMISSION OF IMPULSES AND THE SYNAPSE

In the previous section we examined the structure of neurones and the organisation of the nervous system. We now turn to the fundamental question: how does a neurone actually transmit a signal? The answer lies in rapid, temporary changes in the electrical potential across the neuronal membrane — the **action potential** — and in the specialised junction where one neurone communicates with another: the **synapse**. Understanding these mechanisms is essential not only for the GCE examination but also for appreciating how drugs, toxins and diseases affect nervous function.

6.1 Resting Potential: The Polarised Membrane

Before a neurone can fire, it must establish a **resting potential** — a voltage difference across its membrane with the inside negative relative to the outside. In most mammalian neurones this value is approximately -70 mV.

DEFINITION Resting Potential

The resting potential is the steady membrane potential of an excitable cell (neurone or

muscle fibre) when it is not transmitting an impulse. It is typically around -70 mV (inside negative) and is maintained by the unequal distribution of ions and the selective permeability of the membrane.

Three factors create and maintain this potential:

1. **Concentration gradients:** The extracellular fluid is rich in Na^+ and Cl^- , while the intracellular fluid is rich in K^+ and large organic anions (A^-). These gradients are established by the **sodium–potassium pump** (Na^+/K^+ -ATPase), which actively transports 3 Na^+ out and 2 K^+ in per ATP hydrolysed.
2. **Selective permeability:** At rest, the membrane is far more permeable to K^+ (via **leak channels**) than to Na^+ or Cl^- . K^+ diffuses out down its concentration gradient, leaving behind unbalanced negative charges (mainly A^-) inside.
3. **Electrochemical equilibrium:** K^+ efflux continues until the electrical attraction pulling K^+ back in balances the chemical gradient pushing it out. This equilibrium potential for K^+ (calculated by the Nernst equation) is about -90 mV. The resting potential is slightly less negative because of a small Na^+ leak.

KEY POINTS Key Points on Resting Potential

- The Na^+/K^+ pump is electrogenic (moves 3 positive charges out, 2 in) and contributes directly a few millivolts to the negativity inside.
- The membrane is polarised at rest. Any stimulus that reduces this polarisation (makes the inside less negative) is a **depolarisation**; an increase in negativity is a **hyperpolarisation**.
- The resting potential is a steady state, not a true equilibrium, because it requires continuous ATP expenditure by the pump.

6.2 Action Potential: The Nerve Impulse

When a stimulus depolarises the membrane to a critical **threshold** (typically around -55 mV), a massive, transient reversal of the membrane potential occurs: the **action potential**.


DEFINITION Action Potential

An action potential is a rapid, all-or-nothing reversal of the membrane potential from about -70 mV to about $+30$ mV and back, lasting 1–2 ms in myelinated neurones. It is the electrical signal that propagates along the axon.

PROPERTY Phases of the Action Potential

The action potential results from the sequential opening and closing of **voltage-gated ion channels**:

1. **Depolarisation (rising phase):** Voltage-gated Na^+ channels open rapidly. Na^+ influx drives the membrane potential towards the Na^+ equilibrium potential ($\approx +60 \text{ mV}$). The membrane potential overshoots zero.
2. **Repolarisation (falling phase):** Voltage-gated Na^+ channels inactivate (a distinct conformational state). Voltage-gated K^+ channels open more slowly. K^+ efflux restores the negative interior.
3. **Hyperpolarisation (undershoot):** K^+ channels remain open slightly longer than needed, driving the potential closer to the K^+ equilibrium potential ($\approx -90 \text{ mV}$).
4. **Return to rest:** K^+ channels close; the Na^+/K^+ pump restores the exact ion distributions.



Typical action potential waveform in a myelinated mammalian neurone. The threshold is the voltage at which voltage-gated Na^+ channels open regeneratively.

DEFINITION Refractory Periods

After an action potential, the membrane cannot immediately fire again.

- **Absolute refractory period:** From the start of depolarisation until Na^+ channels recover from inactivation (approx. 1–2 ms). No stimulus, however strong, can trigger a second action potential. This enforces one-way propagation.
- **Relative refractory period:** During hyperpolarisation. A stronger-than-threshold stimulus can elicit a second action potential because the membrane is closer to the K^+ equilibrium potential.

PROPERTY All-or-Nothing Law

The action potential is an **all-or-nothing** event: if the stimulus reaches threshold, the action potential always has the same amplitude and shape for a given neurone; if the stimulus is subthreshold, no action potential occurs. The nervous system encodes stimulus intensity by the frequency of action potentials, not their size.

WARNING Common Mistake: Graded Potentials vs. Action Potentials

Do not confuse graded potentials (local, decremental, amplitude varies with stimulus strength — e.g. generator potentials at receptors, EPSPs/IPSPs at synapses) with action potentials (all-or-nothing, non-decremental, fixed amplitude). Only action potentials propagate over long distances.

6.3 Propagation of the Action Potential

The action potential at one patch of membrane depolarises the adjacent membrane, triggering a new action potential there. This **local current flow** creates a wave of depolarisation that travels along the axon.

PROPERTY Factors Affecting Conduction Speed


1. Myelination: In myelinated axons, voltage-gated channels are concentrated at the **nodes of Ranvier**. The action potential jumps from node to node (**saltatory conduction**), which is much faster and energetically cheaper than continuous conduction in unmyelinated fibres.
2. Axon diameter: Larger diameter → lower axial resistance → faster local current spread → higher speed.
3. Temperature: Higher temperature increases ion channel kinetics and diffusion rates, speeding conduction (up to a physiological limit).

DID YOU KNOW? Cameroon Context: Myelin and Disease

Demyelinating conditions (e.g. multiple sclerosis, though rarer in sub-Saharan Africa) drastically slow conduction, causing neurological deficits. In Cameroon, nutritional neuropathies (e.g. vitamin B12 deficiency) can also affect myelination. Understanding saltatory conduction explains why even partial demyelination has severe functional consequences.

6.4 The Synapse: Structure and Chemical Transmission

When the action potential reaches the axon terminal, it must cross the **synaptic cleft** to excite (or inhibit) the next cell. At a **cholinergic synapse** (using acetylcholine, ACh), the process is as follows.



Structure of a cholinergic synapse. Voltage-gated Ca^{2+} channels open on depolarisation; synaptic vesicles fuse and release ACh; ACh diffuses across the cleft and binds to receptors (AChR) on the postsynaptic membrane; acetylcholinesterase (AChE) rapidly hydrolyses ACh.

METHOD Synaptic Transmission in Six Steps

Learn this sequence for examinations:

1. **Arrival:** An action potential depolarises the presynaptic membrane.
2. **Ca^{2+} entry:** Voltage-gated Ca^{2+} channels open; Ca^{2+} rushes in down its steep electrochemical gradient.
3. **Vesicle fusion:** Ca^{2+} binds to synaptotagmin, triggering SNARE-mediated fusion of synaptic vesicles with the presynaptic membrane.
4. **Release:** Acetylcholine (ACh) is expelled into the synaptic cleft by exocytosis.
5. **Diffusion & binding:** ACh diffuses across the cleft ($\approx 20 \text{ nm}$) and binds to nicotinic ACh receptors (ligand-gated Na^+/K^+ channels) on the postsynaptic membrane.
6. **Postsynaptic response:** Channels open; Na^+ influx (and K^+ efflux) causes a local depolarisation — an **excitatory postsynaptic potential (EPSP)**. If the EPSP reaches threshold, an action potential is initiated in the postsynaptic neurone.

DEFINITION Excitatory and Inhibitory Postsynaptic Potentials

- EPSP: A depolarising graded potential (e.g. via Na^+ influx) that brings the membrane closer to threshold.
- IPSP: A hyperpolarising graded potential (e.g. via Cl^- influx or K^+ efflux) that moves the membrane away from threshold.

6.5 Synaptic Integration, Modulation and Pharmacology

A single EPSP is usually too small to reach threshold. The postsynaptic neurone integrates thousands of inputs.

PROPERTY Summation

- Temporal summation: Rapid successive EPSPs from the same presynaptic neurone add up because the membrane time constant ($\approx 10\text{--}20\text{ ms}$) allows them to overlap.
- Spatial summation: Simultaneous EPSPs from different presynaptic neurones at different synapses add up.

Inhibitory inputs (IPSPs) subtract from this sum. The axon hillock (initial segment) is the **trigger zone** where the net potential is assessed.

WARNING One-Way Transmission

Synaptic transmission is unidirectional (presynaptic $\rightarrow$ postsynaptic) because:

1. Vesicles and release machinery are only in the presynaptic terminal.
2. Receptors are only on the postsynaptic membrane.
3. Acetylcholinesterase (AChE) in the cleft rapidly hydrolyses ACh, preventing back-diffusion and repeated stimulation.

KEY POINTS Role of Acetylcholinesterase (AChE)

AChE hydrolyses ACh into choline and acetate within $< 1\text{ ms}$. This:

- Terminates the signal, allowing high-frequency firing.
- Recycles choline (taken back up by the presynaptic terminal via a high-affinity transporter) for resynthesis of ACh.

Inhibition of AChE (e.g. by organophosphates) causes prolonged depolarisation, depolarisation block, and potentially fatal overstimulation of muscles and glands.

EXAMPLE Drugs and Toxins at the Cholinergic Synapse — Exam Favourites

Agent	Site/Mechanism	Effect
Curare	Competitive antagonist at nicotinic ACh receptors (postsynaptic)	Blocks EPSPs → flaccid paralysis (used historically as arrow poison in South America; principle of neuromuscular blockers in anaesthesia).
Organophosphates (e.g. malathion, sarin)	Irreversible inhibition of AChE	ACh accumulates → sustained depolarisation → depolarisation block, convulsions, respiratory failure.
Neostigmine, physostigmine	Reversible AChE inhibitors	Used to treat myasthenia gravis (autoimmune loss of ACh receptors) and glaucoma.
Botulinum toxin (Botox)	Prevents vesicle fusion (cleaves SNARE proteins)	Blocks ACh release → flaccid paralysis (therapeutic: dystonia, cosmetic).
Nicotine	Agonist at nicotinic receptors	Mimics ACh → stimulation then desensitisation (addiction via dopamine release in reward pathways).

EXPERIMENT Investigating the Effect of Temperature on Conduction Velocity (Core Practical)

Aim: Determine how temperature affects the speed of nerve impulse conduction. **Apparatus:** Isolated sciatic nerve–gastrocnemius muscle preparation (frog), stimulating electrodes, recording electrodes, water bath at controlled temperatures (5°C, 15°C, 25°C, 35°C), kymograph or digital oscilloscope. **Method:**

1. Stimulate the nerve at a fixed point; record the time for the muscle twitch to begin (latency).
2. Measure the distance between stimulation and recording points.
3. Calculate conduction velocity = distance / (latency – synaptic delay – excitation–contraction coupling time).
4. Repeat at each temperature; allow equilibration.

Expected result: Conduction velocity increases with temperature (approx. doubling per 10°C rise, $Q_{10} \approx 2$) until thermal damage occurs. **Examiner's tip:** Be ready to explain why (increased ion channel kinetics, faster diffusion) and to identify sources of error (synaptic delay variability, muscle contraction time).

WORKED EXAMPLE 8 Worked Example: Refractory Period and Maximum Firing Frequency

Question: A myelinated motor neurone has an absolute refractory period of 1.2 ms and a relative refractory period of 3.8 ms. What is the theoretical maximum firing frequency of this neurone? Explain why the actual maximum frequency in vivo is often lower.

Solution: The absolute refractory period sets the hard limit on how quickly a second action potential can be initiated. The minimum interval between two action potentials is the absolute refractory period (ARP).

$$f_{\max} = \frac{1}{\text{ARP}} = \frac{1}{1.2 \times 10^{-3} \text{ s}} \approx 833 \text{ Hz.}$$

Why lower in vivo?

- During the relative refractory period, a stronger stimulus is needed; natural synaptic inputs may not provide sufficient depolarisation.
- Synaptic integration (temporal/spatial summation) takes time.
- Metabolic constraints (ATP for Na^+/K^+ pump) limit sustained high-frequency firing.
- Many neurones exhibit spike-frequency adaptation (slow K^+ currents, Ca^{2+} -activated K^+ channels) that progressively raises the threshold.

Typical sustained firing rates for motor neurones are 50–100 Hz.

EXERCISE 6 Practice Question: Synaptic Transmission

A student applies a toxin that specifically blocks voltage-gated Ca^{2+} channels in the presynaptic terminal of a cholinergic synapse. The presynaptic neurone is stimulated with action potentials at 50 Hz.

1. Predict the effect on the postsynaptic membrane potential. Justify your answer.
2. Would the application of exogenous acetylcholine (via a micropipette) onto the postsynaptic membrane restore the response? Why or why not?
3. Name one natural toxin that acts via this mechanism.

SOLUTION Exercise: Synaptic Transmission

1. Effect: No EPSPs will be generated; the postsynaptic membrane will remain at resting potential. Justification: Voltage-gated Ca^{2+} channels are essential for vesicle fusion and ACh release. Blocking them prevents exocytosis of ACh, so no neurotransmitter reaches the cleft, no binding to receptors occurs, and no ion channels open on the postsynaptic membrane.
2. Yes. Exogenous ACh applied directly to the postsynaptic membrane will bind to the nicotinic receptors (which are unaffected by the presynaptic toxin) and open their ion channels, producing a depolarisation (EPSP). This bypasses the blocked release step.
3. ω -Conotoxin GVIA (from the cone snail *Conus geographus*) specifically blocks N-type voltage-gated Ca^{2+} channels at presynaptic terminals. (Also accept: certain spider toxins, though ω -conotoxin is the classic example.)

KEY POINTS Summary Checklist for Section 6

- Resting potential (≈ -70 mV) is maintained by the Na^+/K^+ pump and high resting K^+ permeability.
- Action potential: threshold $\rightarrow$ Na^+ influx (depolarisation) $\rightarrow$ Na^+ channel inactivation + K^+ efflux (repolarisation) $\rightarrow$ hyperpolarisation $\rightarrow$ return to rest.
- All-or-nothing law; refractory periods enforce one-way propagation and limit frequency.
- Conduction speed: myelination (saltatory) $>$ diameter $>$ temperature.
- Synapse: Ca^{2+} entry $\rightarrow$ vesicle fusion $\rightarrow$ ACh release $\rightarrow$ diffusion $\rightarrow$ receptor binding $\rightarrow$ EPSP/IPSP.

- AChE terminates signal; summation integrates inputs; transmission is one-way.
- Key toxins: curare (receptor block), organophosphates (AChE inhibition), botulinum (release block), ω -conotoxin (Ca^{2+} channel block).

This section provides the mechanistic foundation for the next topic: how the brain, spinal cord and autonomic nervous system organise these signals into coordinated responses.

7 REFLEX ACTIONS, BRAIN, SPINAL CORD AND AUTONOMIC NERVOUS SYSTEM

In the previous section we saw how an impulse travels along a neurone and crosses a synapse. But a single impulse travelling along a single neurone does not yet constitute a coordinated response. Real responses involve chains of neurones organised into pathways. The simplest and fastest of these pathways is the **reflex arc**, and the great coordinating centres that receive, integrate and dispatch impulses are the **brain** and the **spinal cord**. Finally, much of the body's internal housekeeping — heartbeat, digestion, glandular secretion — is controlled without our conscious will by the **autonomic nervous system**. These three topics form the heart of this section.

7.1 Reflex Actions

Observation. Touch a hot cooking pot in the kitchen and your hand is withdrawn before you even feel the pain. Dust blows into your eye and you blink instantly. A doctor taps the tendon below your kneecap and your lower leg kicks forward. In each case the response is so fast that there was no time to “think”. Such responses are called reflex actions.

DEFINITION Reflex action

A **reflex action** is a rapid, automatic, involuntary and stereotyped response to a stimulus, which does not require conscious thought and is usually protective in function.

KEY POINTS Characteristics of reflex actions

- **Rapid:** the pathway is short and involves few synapses (typically one or two).
- **Automatic and involuntary:** they occur without conscious decision; you cannot easily prevent a blink.
- **Stereotyped:** the same stimulus always produces the same response.
- **Protective:** they remove the body (or a part of it) from harm, or maintain posture and

balance.

The reflex arc. The nervous pathway followed by the impulse during a reflex is called the **reflex arc**. It always contains the same five components in the same order.


DEFINITION Reflex arc

A **reflex arc** is the nervous pathway along which an impulse travels from a receptor to an effector during a reflex action. Its components are:

receptor → sensory neurone → relay neurone (interneurone) → motor neurone → effector

In a typical **spinal reflex** (e.g. the withdrawal reflex), the sequence is:

1. A **stimulus** (e.g. heat from a hot object) is detected by a **receptor** (pain/temperature receptors in the skin of the finger).
2. An impulse is generated and travels along the **sensory neurone**, whose cell body lies in the **dorsal root ganglion**, into the spinal cord through the **dorsal root**.
3. Inside the **grey matter** of the spinal cord, the sensory neurone synapses with a **relay neurone** (intermediate neurone / interneurone).
4. The relay neurone synapses with a **motor neurone**, which leaves the cord through the **ventral root**.
5. The motor neurone carries the impulse to an **effector** (the biceps muscle of the arm), which contracts and withdraws the hand.



Flow diagram of a simple spinal reflex arc (withdrawal reflex).

WARNING The brain is not needed first

A very common examination error is to write that the impulse “goes to the brain, which decides to withdraw the hand”. In a simple spinal reflex the impulse does not need to reach the brain before the response occurs: the relay takes place entirely in the spinal cord, which is why the response is so fast. An impulse is also sent up to the brain, so you become aware of the pain — but only after the hand has already been withdrawn.

Common examples of spinal reflexes.

- **Knee jerk (patellar reflex):** tapping the tendon below the kneecap stretches the thigh muscle (quadriceps); stretch receptors (muscle spindles) trigger a reflex contraction of the same muscle. This is a **monosynaptic reflex** (only one synapse between sensory and motor neurone). Used by doctors to test the integrity of the spinal cord segments L2–L4.
- **Withdrawal reflex:** pulling the hand away from a hot or sharp object (protective). This is a **polysynaptic reflex** (involves one or more relay neurones).
- **Blinking (corneal reflex):** protects the eye from foreign bodies and bright light; involves the trigeminal (V) and facial (VII) cranial nerves — a **cranial reflex**.
- **Pupil reflex:** constriction of the pupil in bright light, protecting the retina (studied in Section 3).
- **Coughing and sneezing reflexes:** expel irritants from the respiratory tract.

Conditioned reflexes. All the reflexes above are **inborn (unconditioned)**. The Russian physiologist **Ivan Pavlov** showed that a reflex can be learned: he rang a bell each time he fed dogs; after repeated association, the dogs salivated at the sound of the bell alone. A **conditioned**


reflex is a learned response in which a previously neutral stimulus (the bell) comes to trigger a reflex response (salivation) after repeated association with a natural stimulus (food). Conditioned reflexes involve the **cerebrum** (higher centres) and are the basis of much learning and habit formation.

KEY POINTS Unconditioned vs. conditioned reflexes

- Unconditioned (simple) reflex: inborn, inherited, does not require learning, involves spinal cord or lower brain centres (e.g. knee jerk, withdrawal).
- Conditioned reflex: acquired during life, requires learning and memory, involves the cerebrum (e.g. salivating at the sound of a bell, a student opening a book when the bell rings).

7.2 The Brain

The brain is the anterior, enlarged part of the central nervous system, protected by the skull (cranium) and by three membranes, the **meninges** (dura mater, arachnoid mater, pia mater), and cushioned by **cerebrospinal fluid (CSF)**. A sagittal section reveals several distinct regions, each with precise functions.



Sagittal section of the human brain showing the main regions.

Functions of the main regions.

Region	Function(s)
Cerebrum (cerebral hemispheres)	Centre of intelligence, memory, reasoning, learning, imagination and will ; receives and interprets sensory information (sight, hearing, touch, taste, smell); initiates all voluntary movements . Its folded outer layer is the cerebral cortex (grey matter).
Cerebellum	Coordinates balance, posture and fine muscular movement ; ensures movements are smooth and precise (e.g. walking, writing, riding a bicycle). Receives input from proprioceptors, eyes and vestibular apparatus.
Medulla oblongata	Controls involuntary vital activities : heart rate (cardiac centre), breathing rate (respiratory centre), blood pressure (vasomotor centre), swallowing, coughing, vomiting and peristalsis.
Pons	Relays signals between cerebrum, cerebellum and medulla; contains the pneumotaxic centre which helps regulate breathing rate.
Hypothalamus	Regulates body temperature (thermoregulation), water balance (osmoregulation) , hunger, thirst and sleep; controls the pituitary gland , linking the nervous and endocrine systems.
Thalamus	A relay and sorting station : directs sensory impulses (except smell) to the correct areas of the cerebrum.
Pituitary gland	The “master gland”: secretes hormones controlling other endocrine glands (detailed in Sections 8 and 9).

WARNING Exam tip

Examiners frequently ask: “State one function of each named part of the brain.” Learn one precise function per region — e.g. cerebellum = balance and coordination of movement; medulla = control of breathing and heart rate; hypothalamus = temperature and water balance regulation. Vague answers such as “controls the body” earn no marks.


DID YOU KNOW? The blood–brain barrier

The brain is protected not only by bone and meninges but also by the blood–brain barrier: endothelial cells of brain capillaries are tightly joined, and astrocytes wrap around them, preventing many substances (including most antibiotics and neurotransmitters) from entering the brain tissue. This maintains a stable chemical environment for neurones.

7.3 The Spinal Cord

The spinal cord is a cylinder of nervous tissue running from the medulla oblongata down the vertebral column (to about L1/L2 in adults), protected by the vertebrae, the meninges and cerebrospinal fluid. In transverse section it shows:

- An inner, butterfly-shaped (or H-shaped) region of **grey matter**, containing the **cell bodies** of motor and relay neurones, dendrites, and the synapses of reflex arcs. The grey matter has **dorsal horns** (sensory) and **ventral horns** (motor).
- An outer region of **white matter**, containing **myelinated axons** running up and down the cord in tracts (ascending sensory tracts, descending motor tracts).
- A central canal containing cerebrospinal fluid.
- **Dorsal roots** (carrying sensory impulses into the cord, with a swelling — the **dorsal root ganglion** — containing sensory neurone cell bodies) and **ventral roots** (carrying motor impulses out of the cord). Dorsal and ventral roots unite to form a **spinal nerve** (mixed nerve).



Transverse section of the spinal cord showing grey matter, white matter, central canal, dorsal/ventral roots, and the path of a reflex arc (dashed purple).

KEY POINTS Functions of the spinal cord

1. It is the reflex centre for spinal reflexes (the relay neurones of reflex arcs synapse in its grey matter).
2. It is a pathway conducting sensory impulses up to the brain (ascending tracts) and motor impulses down from the brain to the effectors (descending tracts).

WARNING Spinal cord injury level matters

Damage to the spinal cord in the neck (cervical region) affects all four limbs (tetraple-

gia/quadruplegia) because both ascending and descending tracts for the arms and legs are interrupted. Damage lower down (thoracic/lumbar) affects only the legs (paraplegia). Local spinal reflexes below the injury may persist (e.g. knee jerk) because the reflex arc remains intact, but they are no longer under brain control.

7.4 Voluntary and Involuntary Actions; the Autonomic Nervous System

A **voluntary action** is initiated consciously by the cerebrum (e.g. picking up a pen, walking to school, speaking). An **involuntary action** occurs without conscious control. Involuntary actions include reflexes and the continuous automatic regulation of internal organs — heartbeat, gut movements, pupil size, glandular secretion. This internal regulation is the work of the **autonomic nervous system** (ANS).

The ANS consists of motor neurones running from the brain and spinal cord to internal effectors (smooth muscle, cardiac muscle and glands). It has two divisions with **antagonistic (opposing)** effects.

PROPERTY Antagonism of the two autonomic divisions

The **sympathetic** and **parasympathetic** divisions of the autonomic nervous system act antagonistically on the same organs: the sympathetic division prepares the body for emergency and activity (“fight or flight”), while the parasympathetic division promotes rest, digestion and energy conservation (“rest and digest”). The balance between the two keeps internal conditions stable.

Organ / function	Sympathetic (emergency)	Parasympathetic (rest)
Heart rate	Increases (positive chronotropic)	Decreases (negative chronotropic)
Force of heart contraction	Increases	Decreases slightly
Pupil of the eye	Dilates (mydriasis)	Constricts (miosis)
Gut (peristalsis, digestion)	Slowed / inhibited	Stimulated
Salivary glands	Secretion reduced (thick, mucus-rich; dry mouth)	Secretion increased (watery)
Bronchi of lungs	Dilate (bronchodilation; more air)	Constrict (bronchoconstriction)
Blood vessels of skin/gut	Constrict (blood diverted to muscles)	Dilate (or no direct effect)
Blood vessels of skeletal muscle	Dilate (via adrenaline on β_2 receptors)	No direct effect
Adrenal medulla	Stimulated to release adrenaline (epinephrine)	No effect
Bladder / rectum	Emptying inhibited (sphincters contract)	Emptying promoted (detrusor contracts, sphincters relax)
Liver	Glycogenolysis stimulated (glucose release)	Glycogenesis promoted

Comparison of the antagonistic effects of the sympathetic and parasympathetic divisions.

DID YOU KNOW? Why your mouth goes dry before an oral exam

When you are frightened — for example just before speaking in front of the class — the sympathetic division dominates: your heart beats faster, your pupils dilate, blood is diverted to your muscles and saliva production falls, giving you a dry mouth. This is the same “fight or flight” preparation that helped our ancestors escape danger on the savanna. Afterwards, the parasympathetic division restores calm: the heart slows and digestion resumes.

KEY POINTS Anatomical differences between the two divisions

- Sympathetic: preganglionic neurones have cell bodies in the thoracic and lumbar regions of the spinal cord (thoracolumbar outflow); short preganglionic axons, long postganglionic axons; ganglia form a chain alongside the vertebral column.
- Parasympathetic: preganglionic neurones have cell bodies in the brainstem (cranial nerves III, VII, IX, X) and sacral spinal cord (craniosacral outflow); long preganglionic axons, short postganglionic axons; ganglia lie near or within the target organs.

- **Neurotransmitters:** both divisions release acetylcholine (ACh) at the preganglionic synapse (nicotinic receptors). Parasympathetic postganglionic fibres release ACh (muscarinic receptors). Sympathetic postganglionic fibres mostly release norepinephrine (norepinephrine) (adrenergic receptors), except those to sweat glands and some blood vessels in muscle which release ACh.

The protective role of reflexes in daily life. Reflexes protect us constantly and without effort: blinking shields the cornea from dust; coughing and sneezing expel irritants from the airways; the withdrawal reflex removes limbs from hot or sharp objects; postural reflexes (coordinated by the cerebellum and spinal cord) keep us upright; the pupil reflex prevents damage to the retina by bright light. Because they bypass conscious decision-making, they save the precious fractions of a second that often make the difference between injury and safety.

WORKED EXAMPLE 9 Worked exercise: analysing a reflex

A student accidentally touches a thorn while picking hibiscus flowers and immediately pulls her hand away. (a) Name this type of response and give two of its characteristics. (b) List, in order, the five components of the nervous pathway involved. (c) Explain why the hand is withdrawn before the student feels the pain.

SOLUTION

(a) This is a reflex action (a spinal withdrawal reflex). Any two characteristics: it is rapid, automatic/involuntary (no conscious thought), stereotyped, and protective.

(b) The reflex arc:

receptor (pain receptor in skin) → sensory neurone → relay neurone (grey matter of spinal cord) → motor neurone → effector (muscle)

(c) The impulse passes directly from the sensory neurone to the relay neurone and then to the motor neurone within the spinal cord; this pathway is short and involves few synapses, so the muscle contracts very quickly. The impulse that travels up the spinal cord to the brain — producing the conscious sensation of pain — follows a longer route and arrives later. Hence the hand is withdrawn before the pain is consciously felt.

WORKED EXAMPLE 10 Worked exercise: autonomic control

A man is suddenly confronted by an aggressive dog. Describe the changes in his body brought about by the sympathetic division and explain how they prepare him for “fight or flight”.

SOLUTION

Changes (sympathetic dominance):

- Heart rate and force of contraction increase → more oxygen and glucose delivered to muscles.
- Bronchi dilate → increased air flow and gas exchange.
- Blood vessels in skin and gut constrict; vessels in skeletal muscle dilate → blood redirected to muscles.
- Liver glycogen broken down to glucose → blood glucose rises for energy.
- Pupils dilate → improved vision in low light / wider field of view.
- Gut peristalsis and digestion inhibited → energy diverted from non-essential processes.
- Adrenal medulla releases adrenaline → prolongs and amplifies sympathetic effects.
- Sweating increases → cooling during anticipated exertion.

Preparation: These changes maximise the supply of oxygen and glucose to the brain and muscles, enhance alertness and vision, and suppress non-urgent functions, enabling the man to either fight the dog or run away.

EXERCISE 7 Practice questions

1. Define the term reflex arc and distinguish between a simple (unconditioned) reflex and a conditioned reflex, giving one example of each.
2. State one function of each of the following: cerebellum; medulla oblongata; hypothalamus; cerebrum; pons.
3. Draw a labelled transverse section of the spinal cord and indicate the route taken by an impulse during a spinal reflex.
4. Copy and complete: during sympathetic stimulation, the heart rate ..., the pupil ..., and peristalsis is ...; state the opposite effects produced by the parasympathetic division.
5. Explain why damage to the spinal cord in the neck region can abolish both sensation and movement in the legs.
6. A person eats a large meal. Which division of the autonomic nervous system is dominant? List three effects on the digestive system.
7. Compare the anatomical arrangement (outflow, ganglion position, axon lengths) of the sympathetic and parasympathetic divisions.

SOLUTION Exercise 1

1. A reflex arc is the nervous pathway along which an impulse travels from receptor to effector during a reflex: receptor → sensory neurone → relay neurone → motor neurone → effector. A simple reflex is inborn and automatic (e.g. knee jerk); a conditioned reflex is learned by association (e.g. Pavlov's dogs salivating at a bell).
2. Cerebellum: coordination of balance and muscular movement. Medulla oblongata: control of breathing rate and heart rate. Hypothalamus: regulation of body temperature and water balance (and control of the pituitary). Cerebrum: intelligence, memory, voluntary movement and interpretation of sensation. Pons: relay between brain regions and contains pneumotaxic centre for breathing.
3. The drawing should show: outer white matter, inner butterfly-shaped grey matter (dorsal and ventral horns), central canal, dorsal root with dorsal root ganglion (sensory neurone entering), ventral root (motor neurone leaving). Route: receptor → sensory neurone → dorsal root → relay neurone in grey matter → motor neurone → ventral root → effector.
4. Heart rate increases, the pupil dilates, peristalsis is inhibited. Parasympathetic: heart rate decreases, pupil constricts, peristalsis is stimulated.
5. The spinal cord is the pathway between the brain and the body below the neck. If it is severed, sensory impulses from the legs can no longer ascend to the brain (loss of sensation) and motor impulses from the brain can no longer descend to the leg muscles (paralysis), although local spinal reflexes below the injury may persist.
6. Parasympathetic division dominates ("rest and digest"). Effects: increased salivation, increased gastric secretion, increased peristalsis, relaxation of sphincters, increased blood flow to gut.
7. Sympathetic: thoracolumbar outflow; ganglia in chain alongside vertebral column; short preganglionic, long postganglionic axons; postganglionic neurotransmitter mostly noradrenaline. Parasympathetic: craniosacral outflow; ganglia near/within target organs; long preganglionic, short postganglionic axons; postganglionic neurotransmitter acetylcholine.

KEY POINTS Key points of this section

- A reflex action is rapid, automatic, involuntary, stereotyped and protective; its pathway is the reflex arc (receptor → sensory → relay → motor → effector).
- In a spinal reflex, integration occurs in the spinal cord — the brain is informed afterwards.
- The brain's regions have distinct functions: cerebrum (thought, voluntary action, sensation), cerebellum (balance and coordination), medulla (vital involuntary centres), pons (relay, breathing), hypothalamus (homeostasis and pituitary control), thalamus (sensory relay).

- The spinal cord is both a reflex centre and a two-way conduction pathway (ascending sensory, descending motor tracts).
- The autonomic nervous system controls internal organs through two antagonistic divisions: sympathetic (thoracolumbar, “fight or flight”, noradrenaline) and parasympathetic (craniosacral, “rest and digest”, acetylcholine).

8 THE ENDOCRINE SYSTEM AND MECHANISM OF HORMONE ACTION

In the previous section we explored how the nervous system provides rapid, short-lived coordination through electrical impulses and neurotransmitters. However, not all responses need to be instantaneous. Many vital processes — growth, metabolism, reproduction, water balance — require slower, longer-lasting, and more widespread regulation. This is the domain of the **endocrine system**: a network of ductless glands that secrete chemical messengers called **hormones** directly into the bloodstream. Together, the nervous and endocrine systems form the two pillars of coordination in animals, often working hand in hand.

8.1 Endocrine Glands, Hormones and Target Organs

DEFINITION Hormone and Target Organ

A hormone is a chemical substance, produced in very small quantities by an endocrine gland (ductless gland), secreted directly into the blood plasma, transported by the bloodstream to a specific target organ (or target cells) where it exerts its effect, often at very low concentrations. The target cells possess specific **receptors** for that hormone.

PROPERTY General Properties of Hormones

- Secreted in minute amounts: Concentrations in blood are typically 10^{-12} to 10^{-9} mol dm⁻³.
- Transported by blood: They travel from gland to target via the circulatory system.
- Specificity: Each hormone acts only on cells bearing its specific receptor (lock-and-key principle).
- Action at a distance: Unlike paracrine/autocrine signals, endocrine signals act on distant targets.


- Degradation and excretion: Hormones are inactivated mainly in the liver and kidneys; their half-lives range from minutes (adrenaline) to days (thyroxine).

WARNING Endocrine vs Exocrine Glands — Do Not Confuse!

- Endocrine glands are ductless; they secrete hormones directly into the blood capillaries penetrating the gland. Examples: pituitary, thyroid, adrenal glands.
- Exocrine glands secrete their products (enzymes, mucus, sweat, saliva) into **ducts** that carry them to a body surface or into a lumen (e.g. digestive tract). Examples: salivary glands, sweat glands, pancreas (acinar cells).
- The pancreas is a **mixed gland**: its **acinar cells** (exocrine) secrete digestive enzymes into the pancreatic duct, while its **islets of Langerhans** (endocrine) secrete insulin and glucagon directly into blood. In examination questions, always specify which part you are discussing.

8.2 Location of the Main Endocrine Glands

The major endocrine glands in the human body are positioned strategically. Figure summarises their locations.



Location of the major human endocrine glands. The pancreas is shown for its endocrine portion (islets of Langerhans).

KEY POINTS**Key Endocrine Glands and Their Main Hormones**

Gland	Main Hormones	Primary Functions
Pituitary (Hypophysis)	GH, TSH, ACTH, LH, FSH, Prolactin, ADH, Oxytocin	Master gland; regulates other glands, growth, water balance, reproduction
Thyroid	Thyroxine (T ₄), Triiodothyronine (T ₃), Calcitonin	Basal metabolic rate, growth, calcium homeostasis
Parathyroids	Parathyroid Hormone (PTH)	Raises blood Ca ²⁺
Adrenal Cortex	Cortisol, Aldosterone, Androgens	Stress response, Na ⁺ /K ⁺ balance, secondary sex characteristics
Adrenal Medulla	Adrenaline (Epinephrine), Noradrenaline	Fight-or-flight, rapid metabolic changes
Pancreas (Islets)	Insulin (β -cells), Glucagon (α -cells)	Blood glucose homeostasis
Ovaries	Oestrogen, Progesterone	Female reproductive cycle, secondary sexual characteristics
Testes	Testosterone	Male reproductive function, secondary sexual characteristics

8.3 Chemical Nature of Hormones

Hormones are classified by their chemical structure, which determines their solubility, transport in blood, receptor location, and mechanism of action.

PROPERTY Classification by Chemical Nature

1. Peptide / Protein hormones (hydrophilic): Insulin, glucagon, ADH, GH, TSH, LH, FSH. Synthesised as preprohormones → prohormones → active hormones; stored in secretory vesicles; released by exocytosis.
2. Steroid hormones (lipophilic): Cortisol, aldosterone, testosterone, oestrogen, progesterone. Derived from cholesterol; not stored in vesicles (synthesised on demand); diffuse through plasma membrane; transported in blood bound to carrier proteins.
3. Amine hormones: Derived from tyrosine. Thyroid hormones (T₃, T₄) behave like steroids (lipophilic, nuclear receptors). Catecholamines (adrenaline, noradrenaline) behave like peptides (hydrophilic, membrane receptors).

8.4 Mechanism of Hormone Action

The site of the receptor (cell membrane vs intracellular) dictates the signalling cascade.

8.5 Protein Hormones: Membrane Receptors and Second Messengers

Protein hormones cannot cross the phospholipid bilayer. They bind to specific **transmembrane receptors** on the target cell surface. This binding triggers a conformational change that activates an intracellular signalling cascade, often involving a **second messenger**.

METHOD Mechanism of Protein Hormone Action (cAMP Pathway)

1. First messenger: Hormone (e.g. adrenaline, glucagon) binds to its specific **G-protein-coupled receptor (GPCR)** on the outer surface of the plasma membrane.
2. Receptor activation: The receptor changes shape, activating the associated **G-protein** (G_s) by exchanging GDP for GTP on its α -subunit.
3. Effector activation: The G_α -GTP subunit dissociates and activates **adenylyl cyclase** (membrane enzyme).
4. Second messenger formation: Adenylyl cyclase converts ATP → **cyclic AMP (cAMP)** + PP_i .
5. Amplification: One hormone-receptor complex can generate many cAMP molecules.
6. Protein kinase activation: cAMP activates **Protein Kinase A (PKA)** by binding its regulatory subunits, releasing catalytic subunits.
7. Phosphorylation cascade: PKA phosphorylates specific target proteins (enzymes, ion channels, transcription factors), altering their activity.
8. Termination: **Phosphodiesterase** hydrolyses cAMP → AMP; phosphatases remove phosphate groups; hormone dissociates.

EXAMPLE Adrenaline and Glycogenolysis in Liver

Adrenaline binds to β -adrenergic receptors on hepatocytes → cAMP ↑ → PKA activated → phosphorylates (activates) **phosphorylase kinase** → phosphorylates (activates) **glycogen phosphorylase** → glycogen breakdown → glucose released into blood. Simultaneously, glycogen synthase is phosphorylated (inactivated). This **amplification cascade** ensures a rapid, massive glucose output from a tiny hormone signal.

WARNING Other Second Messenger Pathways

Besides cAMP, two other major pathways exist:

- IP_3 /DAG pathway: Hormone $\rightarrow$ GPCR (G_q) $\rightarrow$ Phospholipase C $\rightarrow$ $PIP_2 \rightarrow IP_3 + DAG$. IP_3 releases Ca^{2+} from ER; DAG activates Protein Kinase C (PKC). Example: ADH (V1 receptor), Oxytocin.
- Receptor Tyrosine Kinase (RTK) pathway: Hormone (e.g. Insulin) binds $\rightarrow$ receptor dimerisation $\rightarrow$ autophosphorylation $\rightarrow$ recruits adaptor proteins (IRS) $\rightarrow$ PI3K/Akt and MAPK cascades. No classic small-molecule second messenger.

8.6 Steroid Hormones: Intracellular Receptors and Gene Transcription


Steroid hormones are lipophilic and diffuse freely through the plasma membrane. Their receptors are located in the **cytoplasm** or **nucleus**.

METHOD Mechanism of Steroid Hormone Action

1. Diffusion: Hormone (e.g. cortisol, testosterone, oestrogen) diffuses through the plasma membrane.
2. Receptor binding: In cytoplasm, receptor is often bound to **heat-shock proteins (HSP90)** keeping it inactive. Hormone binding causes HSP dissociation.
3. Nuclear translocation: Hormone-receptor complex dimerises and translocates into the nucleus (if not already nuclear).
4. DNA binding: Complex binds to specific **Hormone Response Elements (HREs)** — short DNA sequences in promoter/enhancer regions of target genes.
5. Transcription regulation: Recruits co-activators / co-repressors and RNA polymerase $\rightarrow$ increased or decreased **mRNA synthesis**.
6. Translation: New mRNA $\rightarrow$ synthesis of specific proteins (enzymes, structural proteins, receptors) $\rightarrow$ long-term cellular response.

WARNING Time Course Difference

Protein hormone actions (via second messengers) take seconds to minutes (modifying existing proteins). Steroid hormone actions (via gene transcription) take hours to days (synthesising new proteins). This is a classic examination distinction.



Comparison of protein hormone (left, cAMP second messenger) and steroid hormone (right, nuclear receptor / gene transcription) mechanisms of action.

8.7 Negative Feedback Control of Hormone Secretion


Endocrine systems are predominantly regulated by **negative feedback loops**: the output (hormone or its effect) inhibits further secretion. This maintains homeostasis.

METHOD Reading and Interpreting a Negative Feedback Loop

1. Identify the stimulus: What variable changes? (e.g. blood glucose $\uparrow$, blood Ca^{2+} $\downarrow$, metabolic rate $\downarrow$).
2. Identify the sensor: Which gland/cells detect the change? (e.g. pancreatic β -cells, parathyroid chief cells, pituitary thyrotrophs).
3. Identify the hormone secreted: What is released in response? (e.g. insulin, PTH, TSH).
4. Identify the target and effect: Where does the hormone act and what does it do? (e.g. liver $\rightarrow$ glycogenesis; bone/kidney $\rightarrow$ Ca^{2+} release; thyroid $\rightarrow$ thyroxine release).
5. Trace the feedback: Does the effect reverse the original stimulus? (e.g. blood glucose $\downarrow$, blood Ca^{2+} $\uparrow$, metabolic rate $\uparrow$).
6. State the outcome: Homeostasis restored; hormone secretion decreases.

EXPERIMENT Negative Feedback: Thyroxine (T3/T4) Regulation

Scenario: Cold exposure → metabolic rate falls → hypothalamus releases TRH → anterior pituitary releases TSH → thyroid releases T3/T4 → basal metabolic rate rises → body temperature restored → T3/T4 inhibit TRH and TSH secretion (long-loop and short-loop feedback).



Negative feedback control of thyroxine secretion. Long-loop feedback: T3/T4 inhibit hypothalamus. Short-loop feedback: T3/T4 inhibit anterior pituitary.

WORKED EXAMPLE 11 Blood Glucose Negative Feedback

Statement: Describe the negative feedback loop that restores blood glucose concentration after a carbohydrate-rich meal.

Detailed solution:

1. Stimulus: Blood glucose concentration rises above set point ($\sim 5 \text{ mmol dm}^{-3}$).
2. Sensor: β -cells of pancreatic islets of Langerhans detect high glucose via GLUT2 transporters and glucokinase.
3. Hormone secreted: Insulin (protein hormone) released by exocytosis.
4. Target organs and effects:
 - Liver: $\uparrow$ glycogenesis, $\uparrow$ glycolysis, $\downarrow$ gluconeogenesis, $\downarrow$ glycogenolysis.
 - Muscle/Adipose: $\uparrow$ GLUT4 translocation $\rightarrow$ $\uparrow$ glucose uptake.
5. Result: Blood glucose concentration falls.
6. Feedback: Falling glucose removes stimulus $\rightarrow$ insulin secretion decreases. Homeostasis restored.

Examiner's tip: Always mention the sensor, the hormone, the target, the effect, and explicitly state that the effect opposes the original change.

8.8 Comparison of Nervous and Hormonal Coordination

Both systems coordinate body functions, but their modes of operation differ fundamentally.

PROPERTY Nervous vs Hormonal Coordination

Feature	Nervous System	Endocrine System
Nature of signal	Electrical (action potentials) + chemical (neurotransmitters at synapses)	Chemical only (hormones)
Transmission pathway	Along specific neurons (wired)	Via bloodstream (wireless)
Speed of transmission	Very fast (up to 120 m s^{-1})	Slow (seconds to minutes for blood transport)
Speed of response	Milliseconds	Seconds (protein hormones) to hours/days (steroids)
Duration of action	Brief (ms–s); stops when impulses stop	Prolonged (min–days); hormone persists in blood
Specificity	High (specific pathway to specific effector)	High (specific receptor on target cells) but widespread distribution
Adaptation	Rapid adaptation (habituation)	Little adaptation; sustained response
Integration with behaviour	Direct control of voluntary muscle, rapid reflexes	Modulates metabolism, growth, reproduction, mood

DID YOU KNOW? Neuroendocrine Integration in Cameroon's Context

The **hypothalamus** is the master link: it is neural tissue that secretes hormones (releasing/inhibiting factors into the hypophyseal portal system, ADH and oxytocin into posterior pituitary). In stressful situations — like a student facing the GCE Advanced Level exams — the **hypothalamo–pituitary–adrenal (HPA) axis** is activated: neural stress signals → hy-

pothalamus (CRH) → pituitary (ACTH) → adrenal cortex (cortisol). Cortisol then feeds back to the brain to modulate mood and cognition. This neuroendocrine crosstalk explains why chronic stress affects both mental performance and physical health (immune suppression, glucose dysregulation). In Cameroon, rising prevalence of type 2 diabetes in urban areas (Yaoundé, Douala) illustrates the endocrine dimension of lifestyle changes: chronic high-calorie diets → sustained insulin demand → β -cell exhaustion → hyperglycaemia.

8.9 Consolidation Exercises


EXERCISE 8 Exercise 1: Hormone Classification and Mechanism

Classify each hormone below as protein/peptide, steroid, or amine. For each, state: (i) receptor location (membrane / cytoplasmic / nuclear), (ii) second messenger if applicable, (iii) typical time scale of action.

1. Insulin
2. Cortisol
3. Adrenaline
4. Thyroxine (T4)
5. Antidiuretic hormone (ADH)
6. Testosterone

EXERCISE 9 Exercise 2: Negative Feedback Interpretation

The diagram below shows a simplified feedback loop for blood calcium regulation.



1. Identify the stimulus, sensor, hormone, effectors, and response.

2. Explain why this is a negative feedback loop.
3. Predict the effect of a tumour in the parathyroid gland that secretes PTH autonomously (ignoring feedback).

SOLUTION Exercise 1

1. Insulin: Protein. (i) Membrane receptor (RTK). (ii) No classic second messenger; autophosphorylation cascade (PI3K/Akt). (iii) Seconds–minutes.
2. Cortisol: Steroid. (i) Cytoplasmic → nuclear. (ii) None (direct gene regulation). (iii) Hours–days.
3. Adrenaline: Amine (catecholamine). (i) Membrane receptor (GPCR). (ii) cAMP (or IP_3 /DAG via α -receptors). (iii) Seconds.
4. Thyroxine (T4): Amine (iodinated tyrosine) — behaves as steroid. (i) Nuclear receptor. (ii) None. (iii) Hours–days.
5. ADH: Peptide. (i) Membrane receptor (GPCR, V2 receptor in kidney). (ii) cAMP. (iii) Minutes.
6. Testosterone: Steroid. (i) Cytoplasmic/nuclear. (ii) None. (iii) Hours–days.

SOLUTION Exercise 2

1. Stimulus: Low blood Ca^{2+} . Sensor: Parathyroid chief cells (CaSR receptors). Hormone: Parathyroid Hormone (PTH). Effectors: Bone (osteoclasts ↑ resorption), Kidney (↑ Ca^{2+} reabsorption, ↑ 1α -hydroxylase → active vitamin D → ↑ gut Ca^{2+} absorption). Response: Blood Ca^{2+} rises.
2. It is negative feedback because the response (rise in blood Ca^{2+}) inhibits the sensor/gland, reducing further PTH secretion. The output opposes the initial stimulus.
3. Autonomous PTH secretion → persistent high PTH → chronic hypercalcaemia (bone resorption → osteoporosis, kidney stones, abdominal pain, confusion). Feedback inhibition is lost; the gland no longer responds to high Ca^{2+} . This is **primary hyperparathyroidism**.

KEY POINTS Section 8 — Key Points for Revision

- Endocrine glands are ductless; hormones travel in blood to specific target cells with receptors.
- Pancreas = mixed gland (exocrine acini + endocrine islets). Do not confuse roles.
- Protein hormones → membrane receptors → second messengers (cAMP, IP₃/DAG, Ca²⁺) → rapid action via phosphorylation.
- Steroid hormones → intracellular receptors → gene transcription → slow, long-lasting action via new protein synthesis.
- Negative feedback: hormone (or its effect) inhibits its own secretion → homeostasis.
- Nervous = fast, wired, brief. Endocrine = slow, wireless, sustained. They interact (neuroendocrine system).

9 FUNCTIONS OF THE ENDOCRINE GLANDS

In the previous section we saw that hormones are chemical messengers released into the blood by ductless glands, and that they act only on target cells bearing the correct receptors. We now examine each major endocrine gland in turn: the hormones it secretes, the target organs, the effects produced, and the disorders that arise when secretion is too high or too low. As you read, notice a recurring theme: most endocrine secretions are controlled by **negative feedback**, so that the internal environment is kept remarkably constant.

9.1 The Pituitary Gland: the “Master Gland”

The **pituitary gland** is a small pea-sized structure hanging from the hypothalamus at the base of the brain. It is called the **master gland** because many of its hormones control other endocrine glands. It has two lobes with different origins and modes of action.

The **anterior lobe** synthesises and releases several hormones under the control of releasing factors from the hypothalamus:

- **Growth hormone (GH)**: stimulates growth of bones and soft tissues and promotes protein synthesis. **Over-secretion in childhood** causes **gigantism** (abnormal height); **under-secretion in childhood** causes **dwarfism** (stunted growth but normal proportions and intelligence). Over-secretion in adults causes thickening of bones of the hands, feet and jaw (**acromegaly**).
- **Thyroid-stimulating hormone (TSH)**: stimulates the thyroid gland to secrete thyroxine.
- **Adrenocorticotrophic hormone (ACTH)**: stimulates the adrenal cortex to release corticosteroids.
- **Follicle-stimulating hormone (FSH)**: in females stimulates growth of ovarian follicles; in males

stimulates sperm production.

- **Luteinising hormone (LH)**: in females triggers ovulation and formation of the corpus luteum; in males stimulates testosterone secretion.
- **Prolactin**: stimulates milk production by the mammary glands after childbirth.

The **posterior lobe** does not make hormones; it stores and releases **antidiuretic hormone (ADH)** and oxytocin, both produced by the hypothalamus. ADH increases water reabsorption in the kidney collecting ducts, concentrating the urine; deficiency causes **diabetes insipidus** (production of large volumes of dilute urine).

WARNING Anterior versus posterior pituitary

Do not write that the posterior pituitary “secretes” ADH in the sense of manufacturing it. ADH is synthesised in the hypothalamus and only stored and released by the posterior lobe. Examiners penalise this confusion.

9.2 The Thyroid Gland

The **thyroid gland** lies in the neck, on either side of the trachea just below the larynx. It secretes **thyroxine**, an iodine-containing hormone that:

- controls the **metabolic rate** (rate of cellular respiration and heat production);
- is essential for normal **growth and development**, especially of the skeleton and nervous system in children.

Disorders. Iodine is needed to make thyroxine. Where the diet lacks iodine, the thyroid enlarges in an attempt to capture more iodine, producing a visible swelling of the neck called a **goitre** (simple or endemic goitre). Severe thyroxine deficiency in childhood causes **cretinism** (stunted growth and mental retardation); in adults it causes **myxoedema** (low metabolic rate, weight gain, lethargy). Over-secretion (hyperthyroidism) raises metabolic rate, causing weight loss, rapid heartbeat and nervousness.

DID YOU KNOW? Iodine deficiency in Cameroon

Goitre due to iodine deficiency has historically been reported in some hilly and mountainous regions of Cameroon, where soils (and therefore crops) are poor in iodine. The use of iodised table salt is a simple public-health measure that prevents this disorder — a good example of biology applied to everyday life.

9.3 The Adrenal Glands

The two **adrenal glands** sit on top of the kidneys. Each has an inner **medulla** and an outer **cortex**. **Adrenal medulla** — **adrenaline**. **Adrenaline** is released in response to stress, fear or danger and

produces the **fight-or-flight** response:

- increases **heart rate** and the force of contraction, raising blood supply to muscles;
- stimulates **glycogenolysis** (breakdown of glycogen to glucose) in the liver, raising blood glucose for respiration;
- dilates the **pupils** and the bronchioles;
- diverts blood from the gut and skin to skeletal muscles.

Adrenal cortex — corticosteroids. These include glucocorticoids (e.g. cortisol), which raise blood glucose during prolonged stress and suppress inflammation, and mineralocorticoids (e.g. aldosterone), which regulate sodium and potassium balance and hence water reabsorption in the kidney.

WARNING Adrenaline: hormone or nervous effect?

Adrenaline is a hormone: it is secreted into the blood by the adrenal medulla. However, its effects (faster heartbeat, pupil dilation, reduced gut activity) are essentially the same as those of sympathetic nervous stimulation. The distinction to state clearly is this: the sympathetic nervous system acts rapidly and briefly through nerves, while adrenaline acts more slowly but for longer through the bloodstream. The two systems reinforce each other in emergencies.

9.4 The Pancreas and the Control of Blood Glucose

The pancreas is both an exocrine gland (digestive enzymes) and an endocrine gland. Its endocrine tissue, the **islets of Langerhans**, contains two cell types:

- β cells secrete **insulin** when blood glucose is high;
- α cells secrete **glucagon** when blood glucose is low.

Insulin lowers blood glucose by: (i) stimulating the liver and muscles to convert glucose to glycogen (**glycogenesis**); (ii) increasing **glucose uptake** and respiration by cells; (iii) promoting conversion of excess glucose to fat.

Glucagon raises blood glucose by stimulating the liver to break down glycogen into glucose (**glycogenolysis**) and to form glucose from amino acids (**gluconeogenesis**).

PROPERTY Antagonistic action of insulin and glucagon

Insulin and glucagon act **antagonistically**: insulin lowers blood glucose while glucagon raises it. Together, through negative feedback, they keep the blood glucose concentration approximately constant at about 90 mg per 100 cm³ of blood ($\approx 5 \text{ mmol dm}^{-3}$). This is a core example of **homeostasis**.

DEFINITION Negative feedback in blood glucose homeostasis


Negative feedback is a control mechanism in which a change in a regulated variable triggers a response that reverses that change, returning the variable towards its set point. Applied to blood glucose: a rise in blood glucose after a meal is detected by the pancreas, which secretes insulin; insulin lowers glucose, so the stimulus for insulin secretion disappears. A fall in blood glucose triggers glucagon secretion, which raises glucose again. The output thus switches off its own stimulus.

METHOD Describing a negative feedback loop

To explain any negative feedback loop (e.g. blood glucose, thyroxine, ADH, calcium) in an exam answer, follow these four steps:

1. Stimulus: State the change in the internal environment (e.g. blood glucose rises after a meal).
2. Receptor: Identify the sensor (e.g. β cells of the islets of Langerhans detect high glucose).
3. Effector response: Name the hormone secreted and its action (e.g. insulin $\rightarrow$ glycogenesis, glucose uptake).
4. Return to set point: Explain how the response reverses the stimulus and switches off further hormone release.

Always use the phrase “negative feedback” and mention the **set point**.



Blood glucose regulation after a meal. In a normal person, insulin brings the post-meal rise back to the set point within about two hours; if glucose later falls, glucagon restores it. In an untreated diabetic, glucose starts high, rises further and falls only slowly.

Diabetes mellitus. **Diabetes mellitus** is a disorder in which blood glucose is persistently too high because insulin is lacking or ineffective.

- **Causes:** failure of the β cells to secrete insulin (Type 1, usually appearing in youth) or failure of body cells to respond to insulin (Type 2, associated with obesity and age).
- **Symptoms:** high blood glucose (hyperglycaemia), **glucose in the urine**, excessive urine production, persistent thirst, tiredness and weight loss.
- **Control:** Type 1 is managed by regular **insulin injections** (insulin is a protein and would be digested if swallowed); Type 2 is managed by controlling the diet (reducing sugar intake), exercise, and drugs that improve insulin action.

DID YOU KNOW? Diabetes in Cameroon

Diabetes mellitus is a growing public health concern in Cameroon, driven by urbanisation, reduced physical activity and diets richer in refined sugars and fats. The Cameroon Ministry of Public Health runs awareness campaigns on World Diabetes Day (14 November) promoting early screening, balanced nutrition and regular exercise. Traditional foods like ndolé (bitterleaf) and eru are high in fibre and can help regulate blood glucose when prepared with less oil.

9.5 The Ovaries and the Testes

Ovaries. The ovaries secrete **oestrogen** and **progesterone**.

- **Oestrogen** (from the developing follicles) controls the development of female **secondary**

sexual characteristics at puberty (breast development, widening of hips, growth of pubic hair, onset of the menstrual cycle), and repairs and thickens the uterus lining after menstruation.

- **Progesterone** (from the corpus luteum) maintains the thickened uterus lining during the second half of the **menstrual cycle** and throughout **pregnancy**, and inhibits further ovulation during pregnancy.

Testes. The testes secrete **testosterone**, which:

- stimulates the development of male **secondary sexual characteristics** at puberty (deepening of the voice, growth of beard and body hair, muscular development, enlargement of the testes and penis);
- is necessary for the continuous **production of sperm** (spermatogenesis).

9.6 The Parathyroid Glands

The four small **parathyroid glands** are embedded in the back of the thyroid. They secrete **parathormone (PTH)**, which **raises blood calcium** by stimulating calcium release from bones, increasing calcium reabsorption in the kidneys, and promoting calcium absorption from the gut (via vitamin D). Calcium is vital for nerve impulse transmission, muscle contraction and blood clotting; over-secretion weakens bones, while under-secretion causes muscle spasms (**tetany**).

9.7 Summary of the Endocrine Glands

Gland	Hormone(s)	Target	Main effect
Pituitary	GH, TSH, ACTH, FSH, LH, prolactin, ADH	Various endocrine glands, kidney, uterus, breast	Controls other glands; growth; water balance
Thyroid	Thyroxine	All cells	Metabolic rate, growth, development
Key idea: Negative feedback loops maintain homeostasis for glucose, calcium, water, metabolic rate and stress responses.			
Adrenals	Adrenaline, corticosteroids	Heart, liver, pupils, kidney, liver	Fight-or-flight; long-term stress, salt balance
Pancreas	Insulin, glucagon	Liver, muscles, adipose tissue	Blood glucose homeostasis
Ovaries	Oestrogen, progesterone	Uterus, breasts, bones	Female sexual characteristics, menstrual cycle, pregnancy
Testes	Testosterone	Male tissues, testes	Male sexual characteristics, spermatogenesis
Parathyroids	Parathormone (PTH)	Bones, kidneys, gut	Blood calcium homeostasis

Summary map of the major endocrine glands, their hormones, targets and effects.

WORKED EXAMPLE 12 Applying knowledge of endocrine disorders

Statement. A 14-year-old boy in a village in the North-West Region has a visibly swollen neck, feels tired, and is shorter than most boys of his age. Local crops are grown on iodine-poor soils. (a) Name the gland and hormone most likely involved. (b) Name the disorder shown by the swollen neck and explain its cause. (c) Suggest one simple preventive measure for the community. (d) Explain why the boy's growth is affected.

SOLUTION

Solution.

1. The thyroid gland and its hormone thyroxine are involved.
2. The swollen neck is a goitre. Because the diet lacks iodine, the thyroid cannot make enough thyroxine; the gland enlarges as it works harder to trap the little iodine available.
3. Use of iodised table salt (or iodine-rich foods such as sea fish) prevents the deficiency.
4. Thyroxine is needed for normal growth and development; deficiency in childhood slows skeletal growth (and can impair mental development), so the boy is shorter than average.

WARNING Exam tip: one disorder per gland

For the GCE, learn at least one disorder per major gland, with its cause and symptoms: pituitary — gigantism or dwarfism (abnormal GH in childhood); thyroid — goitre (iodine deficiency); pancreas — diabetes mellitus (lack of, or insensitivity to, insulin); parathyroids — tetany (under-secretion of parathormone). Be ready to link cause → hormone level → symptoms in a logical chain.

EXERCISE 10 Checking your understanding

1. Explain why the pituitary is called the master gland, giving two examples of its control over other glands.
2. A patient has a rapid heartbeat, weight loss despite eating well, and bulging eyes. Which gland and hormone are likely involved, and is secretion too high or too low?
3. Describe, step by step, the negative feedback response when blood glucose rises after a meal of ndolé and bobolo.

4. State two differences between the action of adrenaline and the action of insulin.

SOLUTION Exercise 1

1. The pituitary secretes hormones that control other endocrine glands, e.g. TSH stimulates the thyroid to release thyroxine, and ACTH stimulates the adrenal cortex to release corticosteroids; hence “master gland”.
2. The thyroid gland, hormone thyroxine; secretion is too high (hyperthyroidism), raising metabolic rate and heart rate.
3. Rise in blood glucose → detected by β cells of the islets of Langerhans → insulin secreted → liver and muscles convert glucose to glycogen (glycogenesis) and cells take up more glucose → blood glucose falls to the set point → insulin secretion switches off.
4. Adrenaline raises blood glucose (glycogenolysis) and acts quickly in emergencies; insulin lowers blood glucose (glycogenesis, glucose uptake) and acts during normal feeding. Adrenaline also speeds the heart and dilates pupils; insulin does not.

KEY POINTS Key points

- The pituitary is the master gland; GH disorders cause gigantism or dwarfism; ADH (stored in the posterior lobe) controls water reabsorption.
- Thyroxine sets the metabolic rate and drives growth; iodine deficiency causes goitre.
- Adrenaline produces the fight-or-flight response; corticosteroids act in longer-term stress and salt balance.
- Insulin and glucagon act antagonistically, by negative feedback, to hold blood glucose near its set point; failure of this system gives diabetes mellitus.
- Oestrogen and progesterone control female sexual development, the menstrual cycle and pregnancy; testosterone controls male sexual development and sperm production; parathormone raises blood calcium.

10 PLANT MOVEMENTS, PLANT HORMONES, LIGHT AND PLANT GROWTH

In the previous section we saw how animals coordinate their activities through hormones secreted by endocrine glands and transported in the blood. Plants have no blood, no glands and no nervous system, yet a maize seedling in a Cameroonian farm still manages to grow upward out of the soil, bend toward sunlight, and a groundnut plant folds its leaves at night. How is this possible? Plants too produce chemical messengers — the **plant hormones** or **phytohormones** — and they respond to stimuli such as light, gravity, water and touch through slow but precise **growth movements**. In this final section we study these plant responses and show how farmers exploit them in agriculture.

10.1 Plant Movements: Tropisms and Nastic Movements

Observation. Place a potted bean seedling on a window sill in Buea. After a few days, the shoot bends toward the window. Touch the leaflets of *Mimosa pudica* (the “sensitive plant” common along our roadsides) and they fold within seconds. Both are responses to stimuli, but they differ fundamentally: the bending of the shoot is slow, permanent, and directed by the stimulus; the folding of *Mimosa* is rapid, reversible, and occurs in the same direction whatever the direction of touch.

DEFINITION Tropism

A **tropism** is a directional growth movement of a plant organ in response to a directional (unilateral) external stimulus. The direction of growth is related to the direction of the stimulus. A tropism is positive when the organ grows toward the stimulus and negative when it grows away from it.

DEFINITION Nastic movement

A **nastic movement** is a non-directional movement of a plant organ in response to a non-directional (diffuse) stimulus. The direction of the movement is determined by the structure of the organ, not by the direction of the stimulus. Nastic movements are usually rapid, reversible and caused by changes in turgor pressure, not by growth.

The main types of tropisms are named after the stimulus involved:

Tropism	Stimulus	Example
Phototropism	Light	Shoots grow toward light (positive); roots grow away (negative)
Geotropism (gravitropism)	Gravity	Roots grow downward (positive); shoots grow upward (negative)
Hydrotropism	Water	Roots grow toward moist soil (positive)
Thigmotropism (haptotropism)	Touch / contact	Tendrils of passion fruit coil around a support (positive)
Chemotropism	Chemicals	Pollen tube grows toward the ovule (positive)

Examples of nastic movements include:

- **Thigmonasty:** folding of leaflets of *Mimosa pudica* when touched, caused by rapid loss of turgor in swollen motor cells (the **pulvini**) at the leaflet bases;
- **Nyctinasty:** “sleep movements” — leaves of many legumes (e.g. groundnut) fold at night and open in the morning in response to changes in light intensity and temperature;
- **Photonasty:** opening of flowers (e.g. hibiscus) in response to light, and their closing at dusk.

WARNING Tropism or nastic movement?

A very common examination error is to confuse the two. Remember: tropisms are growth movements — they are slow, irreversible (permanent), and their direction depends on the direction of the stimulus. Nastic movements are reversible, often rapid turgor movements, and are not oriented by the stimulus. *Mimosa* folds its leaves whether you touch it from above, below or the side: that is a nastic movement, not a tropism. Also note the spelling geotropism = gravitropism; both are accepted.

10.2 Plant Hormones (Phytohormones)

Plants coordinate growth and responses through small organic molecules produced in one part of the plant and transported (often in the phloem or from cell to cell) to target tissues where they act at very low concentrations.

DEFINITION Plant hormone

A **plant hormone** (phytohormone) is a chemical substance produced naturally in small quantities in one region of a plant and transported to another region, where it regulates growth, development or responses to stimuli.

The five classical groups of phytohormones and their main effects are summarised below.


Hormone	Main site of production	Main effects
Auxins (e.g. IAA)	Shoot tips (apical buds), young leaves, developing seeds	Promote cell elongation in shoots; apical dominance; root initiation; fruit development; mediate phototropism and geotropism
Gibberellins	Young leaves, embryos of seeds, root tips	Promote stem elongation (internode growth); break seed dormancy; stimulate germination; induce flowering and fruit growth
Cytokinins	Root tips, developing fruits and seeds	Promote cell division (cytokinesis); delay leaf senescence (ageing); promote lateral bud growth
Abscisic acid (ABA)	Mature leaves, stems, fruits (especially under stress)	Inhibits growth; promotes seed and bud dormancy; stimulates stomatal closure during water stress ("stress hormone")
Ethene (ethylene)	Ripening fruits, ageing tissues, nodes	Promotes fruit ripening; promotes leaf and fruit abscission (fall); inhibits stem elongation

DEFINITION Auxin

Auxins are plant hormones, the commonest natural one being **indoleacetic acid (IAA)**, synthesised mainly in shoot apical meristems and young leaves. They are transported downwards (basipetally) from cell to cell and promote cell elongation in shoots by loosening cell walls and stimulating proton pumps that acidify the wall space.

PROPERTY Differential sensitivity of shoots and roots to auxin

At the same concentration, auxin promotes cell elongation in shoots but inhibits cell elongation in roots. Roots are far more sensitive to auxin than shoots: the optimum concentration for root elongation is very low (about 10^{-10} to 10^{-8} mol/L), while the optimum for shoots is much higher (about 10^{-5} to 10^{-4} mol/L). Concentrations that stimulate shoot growth therefore inhibit root growth. This property explains the opposite geotropic responses of shoots and roots. (The interpretation of the dose-response curve is examinable.)



Dose-response curves of auxin on shoots and roots. Note the logarithmic concentration axis: roots respond to concentrations about 10^4 times lower than shoots, and are inhibited by concentrations that strongly stimulate shoot elongation.

10.3 The Mechanism of Phototropism

Observation. A coleoptile (the protective sheath of a grass seedling, e.g. maize) illuminated from one side bends toward the light. Classic experiments (Darwin, then Went and others) showed that the tip detects the light, but the bending occurs in the elongation zone below the tip — evidence that a chemical messenger travels downward.

METHOD Demonstrating phototropism with coleoptiles


1. Grow maize or oat seedlings in darkness until the coleoptiles are about 2–3 cm tall.
2. Divide them into four groups and treat as follows: (A) control, left intact; (B) tip covered with an opaque foil cap; (C) tip removed; (D) base of coleoptile covered with foil, tip exposed.
3. Illuminate all groups from one side only (unilateral light) for several hours.
4. Observe: A bends toward the light; B and C do not bend (the tip is needed to perceive light and to produce auxin); D bends normally (the light receptor is at the tip, not the base).
5. A **clinostat** (a slowly rotating drum) can be used as a control to neutralise unilateral stimuli: seedlings rotated on a clinostat grow straight because every side receives the stimulus equally.

Conclusion: the tip perceives light and produces a growth-promoting substance (auxin) that is redistributed and causes differential growth below the tip.

EXPERIMENT Went's agar-block experiment (historical proof that the messenger is chemical)

Went cut off oat coleoptile tips and placed them on agar blocks for some hours; the agar alone, placed asymmetrically on decapitated coleoptiles in darkness, caused curvature away from the side bearing the block — exactly as if a growth-promoting substance had diffused into the agar and then down one side of the shoot. Control blocks of plain agar (never in contact with a tip) caused no curvature. This proved that the tip releases a diffusible chemical — later identified as IAA — and that its asymmetric distribution, not light itself, bends the shoot.

Explanation of the bending. Auxin (IAA) is synthesised at the shoot tip. Under unilateral light, auxin is transported laterally away from the illuminated side toward the shaded side. The shaded side therefore receives more auxin, its cells elongate more, and the shoot curves toward the light.




Auxin redistribution in a shoot tip exposed to unilateral light. Auxin migrates to the shaded side; greater elongation there curves the shoot toward the light.

10.4 The Mechanism of Geotropism (Gravitropism)

Place a germinating seedling horizontally. Within hours the shoot curves upward (negative geotropism) and the root curves downward (positive geotropism). Both responses are caused by the same event — auxin accumulating on the **lower side** — but with opposite outcomes, because of the differential sensitivity stated in the property above.

- **In the shoot:** auxin accumulates on the lower side; the higher concentration promotes elongation there, so the lower side grows faster and the shoot bends upward (away from gravity).
- **In the root:** auxin also accumulates on the lower side, but roots are so sensitive that this

concentration inhibits elongation; the upper side (less auxin) elongates more, so the root bends downward (toward gravity).



Geotropic responses of a horizontally placed seedling. Auxin settles on the lower side of both organs: it stimulates elongation in the shoot but inhibits it in the root.

WORKED EXAMPLE 13 Interpreting a geotropism experiment

A maize seedling is pinned horizontally on a clinostat and another is pinned horizontally on a stationary board, both in darkness. After 24 hours, describe and explain the appearance of each seedling.

SOLUTION

Solution.

- Stationary seedling: the shoot has curved upward and the root downward. Gravity causes auxin to accumulate on the lower side of each organ. In the shoot this higher auxin concentration stimulates cell elongation, so the lower side grows faster and the shoot bends up (negative geotropism). In the root the same concentration inhibits elongation, so the upper side grows faster and the root bends down (positive geotropism).
- Clinostat seedling: both shoot and root continue to grow horizontally, in a straight line. The slow rotation continuously changes the orientation of the seedling relative to gravity, so auxin never accumulates on any one side; distribution remains symmetrical and growth is straight. The clinostat acts as a control showing that gravity, not another factor, caused the curvature.

10.5 Light and Plant Growth: Photomorphogenesis and Etiolation

Light does more than bend shoots: it controls the whole pattern of plant development, a phe-

nomenon called **photomorphogenesis**.

Observation. Potato tubers sprouting in a dark cupboard produce long, pale, weak shoots with tiny leaves. The same tubers sprouting in daylight produce short, sturdy, green shoots. The dark-grown plant is said to be **etiolated**.

DEFINITION Etiolation

Etiolation is the syndrome of abnormal development shown by plants grown in darkness: excessively elongated internodes, pale yellow-white colour (no chlorophyll formed), small unexpanded leaves and a weak stem. On exposure to light, internode elongation slows, chlorophyll forms (greening) and leaves expand.

Etiolation is adaptive: a seedling germinating deep in the soil invests all its reserves in rapid stem elongation to reach the light before its food runs out. Once in the light, energy is redirected to leaf expansion and photosynthesis.

DID YOU KNOW? Phytochrome: the plant's light switch (exam-useful extra)

Plants detect light quality and day length using a blue-green pigment called **phytochrome**. It exists in two interconvertible forms: P_r (absorbs red light, ≈ 660 nm) and P_{fr} (absorbs far-red light, ≈ 730 nm). Red light converts P_r into the active P_{fr} form; far-red light (or darkness, slowly) converts it back. P_{fr} triggers many light responses: greening, leaf expansion, inhibition of etiolation, seed germination in light-sensitive seeds, and the measurement of night length in flowering (photoperiodism). You are not required to know the details, but quoting phytochrome as the receptor of photomorphogenesis earns credit in extended answers.

10.6 Practical Applications in Agriculture

Knowledge of plant hormones is exploited daily, including in Cameroonian agriculture:

- **Rooting powders:** cuttings of cassava, hibiscus or ornamental shrubs dipped in synthetic auxin powder (e.g. IBA, NAA) form adventitious roots faster and more reliably, making vegetative propagation cheap and efficient.
- **Selective weedkillers:** synthetic auxins such as 2,4-D, applied at high concentration, cause uncontrolled, lethal growth in broad-leaved weeds (dicots) while leaving cereal crops like maize and rice (monocots) largely unaffected — widely used in maize fields in the West and North-West Regions.
- **Fruit ripening:** ethene gas is used commercially to ripen harvested bananas, plantains and mangoes uniformly during transport and storage; traders exploit the natural ethene released by ripe fruits placed with unripe ones.
- **Seedless (parthenocarpic) fruits:** spraying unpollinated flowers with auxin or gibberellin induces fruit development without fertilisation, giving seedless tomatoes, cucumbers or

grapes.

- **Breaking dormancy:** gibberellins are used to promote uniform germination of seeds and to induce sprouting of seed yams and potato tubers.
- **Delaying senescence:** cytokinins keep harvested vegetables and cut flowers fresh longer.

WARNING Common mistakes with plant hormones

- Do not write that auxin “moves the shoot” — auxin causes differential cell elongation, and the resulting growth bends the organ. Movement is by growth, not by muscles.
- Do not say auxin is “attracted by darkness” — it is actively transported away from the illuminated side (phototropism) or accumulates on the lower side under gravity (geotropism).
- Remember that the same auxin concentration has opposite effects in shoots and roots; quoting the property precisely is what earns full marks.
- Ethene is a gas; it is the only gaseous plant hormone.

EXERCISE 11 Phototropism and hormones

1. Define the terms tropism and nastic movement, giving one example of each.
2. A coleoptile tip is separated from the rest of the shoot by a thin block of agar, and the shoot is illuminated from one side. Predict, with reasons, whether the shoot will bend toward the light.
3. Explain why a high concentration of synthetic auxin kills dicot weeds but is harmless to maize.
4. State two differences between the etiolated form and the light-grown form of a bean seedling, and name the pigment system responsible for detecting light.

SOLUTION Exercise 1

1. A tropism is a directional growth movement in response to a directional stimulus (e.g. a shoot bending toward light — positive phototropism). A nastic movement is a non-directional, usually reversible movement whose direction is fixed by the organ's structure (e.g. folding of *Mimosa pudica* leaflets on touch).
2. Yes, it will bend. Auxin produced in the tip diffuses down through the agar block (agar is permeable to IAA); unilateral light still causes lateral redistribution of auxin to the

shaded side, which elongates more, curving the shoot toward the light. (If an impermeable mica sheet were inserted on the shaded side instead, bending would be prevented.)

3. Synthetic auxins at high dose cause uncontrolled, disorganised growth (abnormal elongation, disrupted transport tissues) that is lethal to sensitive broad-leaved dicots. Maize, a monocot, is far less sensitive at that concentration, so it survives — the basis of selective weedkillers such as 2,4-D.
4. Etiolated: long internodes, pale yellow (no chlorophyll), small unexpanded leaves, weak stem; light-grown: short internodes, green, expanded leaves, sturdy stem. The light-detecting pigment system is phytochrome (P_r/P_{fr}).

KEY POINTS Key points of the section

- Tropisms are slow, permanent, directional growth movements (photo-, geo-, hydro-, thigmo-, chemotropism); nastic movements are rapid, reversible, non-directional turgor movements.
- Plant hormones: auxins (cell elongation, tropisms), gibberellins (stem elongation, germination), cytokinins (cell division, delay of senescence), abscisic acid (dormancy, stomatal closure), ethene (fruit ripening, abscission).
- Phototropism: auxin made at the tip moves to the shaded side $\Rightarrow$ more elongation there $\Rightarrow$ bending toward light.
- Geotropism: auxin accumulates on the lower side; shoots elongate more there (bend up), roots are inhibited there (bend down) — because roots are far more sensitive to auxin.
- Light controls development (photomorphogenesis) via phytochrome; dark-grown plants are etiolated.
- Applications: rooting powders, 2,4-D weedkillers, ethene ripening of plantain and mango, seedless fruits, breaking dormancy — all relevant to Cameroonian farming.

11 INTEGRATION ACTIVITY

12 RESPONSE & COORDINATION IN ORGANISMS

12.1 Aims of the Activity

KEY POINTS Learning Outcomes

By the end of this integration workshop, you should be able to:


- Consolidate understanding of the fundamental sequence stimulus → reception → coordination → response across animal and plant systems.
- Apply knowledge of taxis, kinesis, tropisms and nastic movements to design and interpret a choice-chamber experiment.
- Analyse clinical endocrine disorders (goitre, diabetes mellitus) by identifying the gland, hormone, feedback mechanism failure and physiological consequences.
- Distinguish between nervous and hormonal coordination in animals, and between hormonal and electrical coordination in plants.
- Develop scientific communication skills by synthesising findings into a structured scientific poster.

12.2 Situation

EXPERIMENT The GBHS Bamenda Integration Workshop

Context: You are Upper Sixth Biology students at Government Bilingual High School (GBHS) Bamenda. The North West Region offers diverse micro-habitats – from the humid forests around Lake Awing to the drier grasslands of the Bamenda Highlands. Your teacher, Mr. Nfor, has organised a Response & Coordination Integration Workshop to prepare you for the GCE Advanced Level practical (Paper 3) and theory (Papers 1 & 2) examinations.

The workshop comprises three linked stations. You will rotate in groups of four, spending 45 minutes at each station. At the end, each group produces an A1 poster titled “Stimulus → Reception → Coordination → Response: Unity in Diversity” linking the three stations.



Station 1: Choice chamber apparatus for investigating wood-louse (*Porcellio scaber*) responses to light and humidity gradients.

Station 1 – The Choice Chamber (Animal Behaviour)

Mr. Nfor provides a choice chamber (see diagram) with four chambers creating combined gradients of light (Light vs Dark) and humidity (Dry vs Humid). Twenty woodlice (*Porcellio scaber*, locally called “kankpe” in Pidgin) are collected from the school compost heap and released at the central junction. After 15 minutes, the distribution is recorded.

Station 2 – The Clinic (Human Endocrine Disorders)

Two case files from the Bamenda Regional Hospital are presented:

Parameter	Case A: Mr. T., 42 years, Farmer from Santa	Case B: Miss N., 16 years, Student, Bamenda Town
Presenting Complaint	Swelling in anterior neck (6 months), fatigue, cold intolerance, weight gain, constipation.	Polyuria, polydipsia, polyphagia, weight loss, fatigue, blurred vision (3 weeks).
Signs	Diffuse, non-tender goitre (Grade 2), bradycardia (56 bpm), dry skin, delayed ankle reflex.	BMI 17.2 kg/m ² , random blood glucose 14.2 mmol/L, glucose++ in urine, ketones+.
Background	Diet: mainly cassava/fufu, low seafood/iodised salt use. No family history.	Sudden onset. No family history of diabetes. Recent severe malaria episode 4 weeks ago.

Station 3 – The School Garden (Plant Responses)

Two observations are made in the school garden behind the science block:

1. **Maize (*Zea mays*) seedlings:** Planted in a row parallel to the western wall of the laboratory. After two weeks, all seedlings show a distinct curvature of the coleoptile and young leaves towards the open field (East), away from the wall.
2. **Mimosa pudica (“Shame-shame plant”):** A potted plant is brought to the lab. When a student touches one leaflet, the entire leaf folds inward and the petiole droops within 1–2 seconds. The response spreads to adjacent leaves. After 15 minutes, the leaves reopen.

12.3 Tasks

Station 1: Choice Chamber Analysis (25 marks)

1. Identify the **stimuli** (adequate and inadequate) and the **receptors** involved in the woodlouse response. Classify the response type (taxis, kinesis, or both) with justification. [4 marks]
2. Predict the expected distribution of the 20 woodlice after 15 minutes in the four chambers (A, B, C, D). Explain your prediction using the concepts of orthokinesis and klinokinesis. [5 marks]
3. The actual results are: A=2, B=3, C=12, D=3. Perform a Chi-squared (χ^2) test to determine if the distribution differs significantly from random expectation (equal distribution). Use $p = 0.05$, critical value $\chi^2_{crit}(3) = 7.82$. State your conclusion. [6 marks]
4. Explain the survival value of this behaviour for woodlice in the leaf litter of the Bamenda Highlands during the dry season (November–March). [3 marks]
5. Propose **two** modifications to the apparatus to investigate the effect of temperature gradient independently of humidity. [2 marks]

Station 2: Clinical Case Analysis (30 marks)

1. For **Case A (Mr. T.)**:
 - (a) Identify the affected endocrine gland and the deficient hormone. [2 marks]
 - (b) Explain the **negative feedback mechanism** involving the hypothalamus, anterior pituitary and thyroid gland that has failed. Use the terms TRH, TSH, Thyroxine (T_4)/Triiodothyronine (T_3). Why does the goitre develop despite low thyroxine? [6 marks]
 - (c) Link the symptoms (cold intolerance, bradycardia, constipation) to the cellular role of thyroid hormone (basal metabolic rate, Na^+/K^+ -ATPase, mitochondrial uncoupling). [4 marks]
2. For **Case B (Miss N.)**:
 - (a) Identify the likely type of diabetes mellitus and the deficient hormone. Explain why ketones appear in urine. [3 marks]
 - (b) Describe the normal **negative feedback loop** regulating blood glucose concentration involving insulin and glucagon. [4 marks]
 - (c) Explain the mechanism of **polyuria** and **polydipsia** in uncontrolled diabetes (renal threshold, osmotic diuresis, ADH). [4 marks]

- (d) Suggest **two** immediate management steps and **one** long-term monitoring test (name the test and target value). [3 marks]

Station 3: Plant Response Analysis (25 marks)

- For the **maize seedlings**:
 - Name the type of response shown. Is it a tropism or a nastic movement? Justify. [3 marks]
 - Identify the stimulus, receptor region, and the hormone responsible. Describe the **asymmetric distribution** mechanism of this hormone causing curvature (Cholodny–Went theory). [6 marks]
- For the **Mimosa pudica**:
 - Name the type of response. Distinguish it clearly from the maize response (directionality, speed, mechanism). [4 marks]
 - Outline the sequence: Mechanical stimulus → Receptor → Electrical signal (Action Potential) → Effector (Pulvinus) → Response. Mention the role of K^+ and Cl^- ion fluxes and water potential (ψ_w) changes in the motor cells. [6 marks]
- Compare and contrast the **coordination systems** in animals (nervous + endocrine) and plants (hormonal + electrical) using examples from Stations 1, 2 and 3. [6 marks]

Synthesis Task: The Poster (20 marks)

- Design the layout of your group's A1 poster. It must contain:
 - A central flow diagram: **Stimulus** → **Reception** → **Coordination** → **Effector** → **Response**.
 - Three branches (Animal Behaviour, Human Endocrine, Plant Responses) feeding into the central flow.
 - For each branch: Specific example, Receptor type, Coordination type (Nervous/Hormonal/Electrical), Effector, Response type.
 - A “Key Concepts” box defining: Taxis, Kinesis, Tropism, Nastic movement, Negative Feedback, Homeostasis.
- Prepare a 3-minute oral defence: “How does the woodlouse finding a humid crack, the thyroxine-deficient patient developing goitre, and the maize seedling bending to light all illustrate the same fundamental biological principle?” [Outline your answer in 150 words.]

12.4 Model Answers

Station 1: Choice Chamber Analysis

- Stimuli & Receptors**:
 - Light**: Adequate stimulus for photoreceptors (compound eyes / ocelli). Woodlice are negatively phototactic (move away from light).
 - Humidity**: Adequate stimulus for hygroreceptors (located on antennae, specifically the flagellum). Woodlice are positively hydrokinetic (orthokinesis: speed decreases in humid air; klinokinesis: turning rate increases in dry air).

Response Classification: Both taxis and kinesis.

- **Negative phototaxis:** Directed movement away from the directional light source (gradient).
 - **Hygrokinesis (Ortho- + Klinokinesis):** Non-directed movement where rate of movement (orthokinesis) and rate of turning (klinokinesis) depend on humidity intensity. In dry air: high speed, high turning → rapid exit. In humid air: low speed, low turning → aggregation.
2. **Predicted Distribution:** Woodlice seek **Dark + Humid** (Chamber D) ideally. However, the chamber creates a conflict: Chamber C (Humid, Light) vs Chamber D (Humid, Dark).
- **Chamber D (Humid, Dark):** Highest aggregation. Both needs met. Low speed, low turning → trap.
 - **Chamber C (Humid, Light):** High humidity retains them (low orthokinesis), but light causes negative phototaxis → movement towards dark partition. Some accumulate near partition.
 - **Chambers A & B (Dry):** High orthokinesis (speed) and klinokinesis (turning) → rapid escape. Very few remain.

Predicted ranking: $D > C \gg B \approx A$.

3. Chi-squared Test:

H_0 : Woodlice are distributed randomly (no preference). Expected (E) = $20/4 = 5$ per chamber.

H_1 : Distribution is not random (preference exists).

$$\begin{aligned}\chi^2 &= \sum \frac{(O - E)^2}{E} \\ &= \frac{(2 - 5)^2}{5} + \frac{(3 - 5)^2}{5} + \frac{(12 - 5)^2}{5} + \frac{(3 - 5)^2}{5} \\ &= \frac{9}{5} + \frac{4}{5} + \frac{49}{5} + \frac{4}{5} = 1.8 + 0.8 + 9.8 + 0.8 = 13.2\end{aligned}$$

Degrees of freedom $df = n - 1 = 3$. Critical value $\chi_{0.05}^2(3) = 7.82$. Since $13.2 > 7.82$, **reject** H_0 . The distribution is highly significant ($p < 0.01$). Woodlice show a strong preference for humid conditions (Chamber C+D = 15 vs A+B = 5) and within humid, a preference for dark (D=3 vs C=12 – see Note below).

WARNING Interpreting C vs D

The data shows C (12) > D (3). This suggests humidity is the dominant drive (kinesis) overriding the light taxis in this specific apparatus, or the light gradient was weak. A good student discusses this: "Humidity kinesis traps them in C before phototaxis moves them to D". This is excellent evaluation.

4. **Survival Value (Dry Season Bamenda):** During the dry season (Harmattan), relative humidity drops below 30%, desiccation risk is extreme. Woodlice lack a waxy cuticle and lose water rapidly via transpiration and excretion.

- **Hygrokinesis** ensures they spend maximum time in moist microhabitats (under logs, deep leaf litter, soil cracks) where ψ_w is high.
- **Negative phototaxis** keeps them under cover (leaf litter, bark) avoiding UV damage and surface heating, which lowers humidity further.
- Combined: Efficient location of the “humid dark” refuge maximises water conservation and survival.


5. Modifications for Temperature Gradient:

- Use a **Peltier element** or water jackets (circulating water baths) at each end of a linear chamber to create a stable thermal gradient, replacing the light/dark covers.
- Control humidity uniformly:** Saturate the entire chamber floor with moist cotton wool / agar gel so humidity is constant (100% RH) and not a confounding variable. Use a data logger with thermistors to verify the gradient.

Station 2: Clinical Case Analysis

1. Case A: Mr. T. (Primary Hypothyroidism / Endemic Goitre)

- Gland:** Thyroid gland (anterior neck). **Hormone:** Thyroxine (T_4) / Triiodothyronine (T_3). Deficiency due to dietary iodine lack (Santa highlands, cassava diet – goitrogens).
- Feedback Failure:**




Low T_3/T_4 → No inhibition of Hypothalamus/Pituitary → **High TRH & High TSH**. High TSH causes **thyroid hyperplasia/hypertrophy** (goitre) attempting to trap more iodide. But without iodine, hormone synthesis fails → colloid accumulation → diffuse goitre.

(c) Symptoms ↔ Cellular Role:

- **Cold intolerance / Low BMR:** T_3 upregulates Na^+/K^+ -ATPase (consumes ATP → heat) and mitochondrial uncoupling proteins (UCP1) → thermogenesis. Deficiency → reduced heat production.
- **Bradycardia:** T_3 increases β -adrenergic receptor density and sino-atrial node automaticity. Deficiency → reduced heart rate/contractility.
- **Constipation:** T_3 maintains gut smooth muscle tone and peristalsis. Deficiency → reduced motility.

2. Case B: Miss N. (Type 1 Diabetes Mellitus – likely autoimmune triggered by viral infection/malaria stress)

- Type 1 DM (Insulin Dependent).** **Deficient hormone:** Insulin (from β -cells of Islets of Langerhans). **Ketones:** Insulin lack → uncontrolled lipolysis in adipose tissue (Hormone-Sensitive Lipase active) → high free fatty acids → liver β -oxidation → acetyl-CoA → ketogenesis (acetoacetate, β -hydroxybutyrate, acetone). Ketones are acidic → metabolic ketoacidosis (DKA risk).

(b) **Glucose Homeostasis Feedback Loop:**(c) **Polyuria & Polydipsia Mechanism:**

- i. Hyperglycaemia ($> 10 \text{ mmol/L}$) exceeds **renal threshold** ($\approx 10 \text{ mmol/L}$).
- ii. Glucose not fully reabsorbed in PCT (SGLT2 saturation) $\rightarrow$ remains in tubular lumen.
- iii. **Osmotic diuresis:** Glucose lowers tubular fluid $\psi_w \rightarrow$ water retained in lumen $\rightarrow$ large volume urine (polyuria).
- iv. Volume depletion $\rightarrow$ increased plasma osmolarity $\rightarrow$ hypothalamic osmoreceptors stimulate **thirst centre** $\rightarrow$ polydipsia. Also ADH released but overwhelmed by osmotic load.

(d) **Management:**

- **Immediate:** 1. IV Fluids (0.9% NaCl) for rehydration + IV Insulin infusion (sliding scale) to lower glucose/ketones. 2. Potassium replacement (insulin drives K^+ into cells $\rightarrow$ hypokalaemia risk). Monitor blood gases (pH, bicarbonate).
- **Long-term monitoring:** HbA_{1c} (**Glycated Haemoglobin**). Target $< 48 \text{ mmol/mol}$ ($< 6.5\%$) for good control; $< 58 \text{ mmol/mol}$ (7.5%) acceptable in adolescents.

Station 3: Plant Response Analysis**1. Maize Seedlings (Phototropism)**

- (a) **Positive Phototropism.** It is a **tropism** (directional growth response where direction of response is determined by direction of stimulus – light from East).
- (b) **Stimulus:** Unilateral blue light (approx. 450 nm). **Receptor:** Phototropins (phot1, phot2) in the **coleoptile tip** (and young leaves). **Hormone:** **Auxin (IAA – Indole-3-Acetic Acid)**.
- (c) **Cholodny–Went Mechanism:**
 - i. Light hits lit side $\rightarrow$ Phototropin activation $\rightarrow$ Phosphorylation of PIN proteins (auxin efflux carriers).
 - ii. PINs relocate to lateral membranes $\rightarrow$ Auxin transported laterally away from light to **shaded side**.
 - iii. **Asymmetric distribution:** Higher [Auxin] on shaded side.
 - iv. In shoots: High [Auxin] $\rightarrow$ stimulates **cell elongation** (via H^+ -ATPase $\rightarrow$ wall acidification $\rightarrow$ expansin activation $\rightarrow$ wall loosening $\rightarrow$ water influx $\rightarrow$ turgor-driven extension).
 - v. Shaded side elongates faster $\rightarrow$ curvature towards light.

2. Mimosa pudica (Thigmonasty / Seismonasty)

(a) **Thigmonasty (Seismonasty)**. **Non-directional** (nastic) response to touch (direction of response – folding – is fixed by plant anatomy, not stimulus direction). **Speed**: Very fast (1–2 s) via **action potentials (APs)**, not growth. **Mechanism**: Turgor loss in pulvinus motor cells (electro-chemical), not differential growth.

(b) **Sequence**:

- i. **Stimulus**: Mechanical deformation of mechanoreceptors (hair cells) on leaflets.
- ii. **Receptor Potential**: Depolarisation → if threshold reached → **Action Potential (AP)** generated.
- iii. **Transmission**: AP propagates via **phloem (sieve tubes)** at $\approx$ few cm/s (electrical + chemical/ H^+ waves) to **pulvini** (swollen leaf bases).
- iv. **Effector (Pulvinus)**: AP reaches extensor/flexor motor cells.
 - **Flexor side (lower)**: AP → Voltage-gated K^+ & Cl^- channels OPEN → massive efflux of K^+ , Cl^- (and malate²⁻) → ψ_s increases (less negative) → ψ_w increases → Water leaves cells (exosmosis) → **Turgor loss** → Cells shrink.
 - **Extensor side (upper)**: Less affected or gains water.
- v. **Response**: Flexor collapse → Leaf folds, petiole droops.
- vi. **Recovery (15 min)**: Active H^+ -pump / K^+ influx → ψ_w drops → Water enters → Turgor restored.

3. Comparison of Coordination Systems

Feature	Animal Nervous (Station 1)	Animal Endocrine (Station 2)	Plant (Station 3)
Signal Type	Electrical (Action Potentials) + Chemical (Neurotransmitters)	Chemical (Hormones in blood)	Chemical (Hormones: Auxin) + Electrical (APs in Mimosa)
Speed	Very fast (ms) – Woodlouse escape	Slow (mins-hours) – Thyroxine/Insulin effects	Slow (hours) – Auxin growth; Fast (s) – Mimosa AP
Target Specificity	High (specific muscle/gland)	Low (all cells with receptor)	Variable (Auxin: specific tissues; AP: whole leaf)
Duration	Short (ms-s)	Long (days-weeks)	Long (growth) / Short (turgor)
Example	Woodlouse hygroreceptor → Interneuron → Leg muscles	Hypothalamus → Pituitary → Thyroid → Body cells	Coleoptile tip → Auxin transport → Shaded cells
Integration	CNS (Brain/Spinal cord)	Feedback loops (Hypothalamo-pituitary axis)	No CNS; Apical dominance, vascular transport

Key Unifying Principle: All systems detect a **change (stimulus)** via a **specific receptor**, transmit information via a **coordination pathway** (neural, hormonal, electrical), and activate an

effector to produce a **response** that restores homeostasis or enhances fitness.

Synthesis Task: The Poster

1. Poster Layout Description:

- **Title Banner:** “Stimulus → Reception → Coordination → Effector → Response: Unity in Diversity” (GBHS Bamenda, Upper Sixth Biology).
- **Central Column (Vertical Flow):** Large arrows connecting 5 boxes: Stimulus (Light, Touch, Hormone level, Humidity) ↓ Reception (Photoreceptor, Mechanoreceptor, Osmoreceptor, Chemoreceptor) ↓ Coordination (Nervous System, Endocrine System, Phloem/Electrical) ↓ Effector (Muscle/Gland, Target Cell, Pulvinus/Meristem) ↓ Response (Taxis/Kinesis, Metabolic change, Growth/Turgor).
- **Left Branch (Animal Behaviour):** Photo of Choice Chamber. Boxes: Stimulus: Light/Humidity → Receptor: Eye/Antenna → Coordination: Ventral Nerve Cord → Effector: Leg Muscles → Response: Kinesis/Taxis.
- **Middle Branch (Human Endocrine):** Diagram of H-P-Thyroid axis & Glucose loop. Boxes: Stimulus: Low T_3 /High Glucose → Receptor: Hypothalamic neurons / β -cells → Coordination: Hormonal (TSH/Insulin) → Effector: Thyroid / Liver/Adipose → Response: Metabolic Rate / Glucose Uptake. **Red Alert Box:** “Feedback Failure = Disease (Goitre, Diabetes)”.
- **Right Branch (Plant Responses):** Split: Top: Maize (Tropism). Stimulus: Unilateral Light → Receptor: Phototropin (Tip) → Coordination: Auxin Transport → Effector: Shaded Cortex → Response: Positive Phototropism. Bottom: Mimosa (Nasty). Stimulus: Touch → Receptor: Mechanoreceptor → Coordination: Action Potential (Phloem) → Effector: Pulvinus Motor Cells → Response: Thigmonasty.
- **Bottom Key Concepts Box:** Definitions of Taxis, Kinesis, Tropism, Nastic Movement, Negative Feedback, Homeostasis, Receptor, Effector.

2. **Oral Defence Outline (150 words):** “All three scenarios illustrate the universal biological principle of **Information Processing for Survival**. In the choice chamber, the woodlouse detects a desiccation threat (low humidity stimulus) via antennal hygroreceptors; its ventral nerve cord coordinates a kinesis/taxis response, moving legs to carry it to a humid refuge – immediate behavioural homeostasis. In the clinic, Mr. T.’s thyroid fails to transduce the metabolic ‘low energy’ signal into thyroxine due to iodine lack; the pituitary over-secretes TSH (failed negative feedback), causing goitre – a chronic failure of hormonal homeostasis. In the garden, the maize seedling detects directional light via tip phototropins; auxin redistribution coordinates differential growth, bending the shoot toward energy – developmental homeostasis. Though the signals (nerves, blood hormones, phloem sap/APs) and timescales (seconds, days, hours) differ, the logic is identical: **Receptor → Transducer → Coordinator → Effector → Adaptive Response**. This unity – from kankpe in the compost to a student in the ward – is the essence of physiology.”

13 METHODS AND WORKED EXAMPLES

14 RESPONSE AND COORDINATION IN ORGANISMS

14.1 Skill 1: Identifying and Interpreting a Reflex Arc

DEFINITION Reflex Action

A **reflex action** is a rapid, automatic, involuntary and stereotyped response to a stimulus, mediated by a **reflex arc** that bypasses conscious brain centres for speed. It is protective and inborn (not learnt).

PROPERTY Components of the Reflex Arc

The reflex arc consists of five essential components in sequence:

1. Receptor: detects the stimulus (e.g. free nerve endings for pain, muscle spindle for stretch).
2. Sensory (afferent) neurone: carries the impulse from receptor to CNS; cell body in **dorsal root ganglion**.
3. Relay (intermediate) neurone: lies entirely within the CNS (spinal cord or brainstem); integrates the signal.
4. Motor (efferent) neurone: carries the impulse from CNS to effector; cell body in **ventral horn** of spinal cord grey matter.
5. Effector: muscle (contracts) or gland (secretes) producing the response.

METHOD Analysing a Reflex Arc Diagram

1. Locate the receptor. Identify the structure in contact with the stimulus (skin, muscle spindle, sense organ). The stimulus arrow points towards the receptor.
2. Trace the afferent pathway. Follow the sensory neurone from receptor to CNS. Note the dorsal root ganglion (swelling on dorsal root).
3. Identify the integration centre. In the spinal cord, the grey matter (butterfly-shaped)

contains the synapse between sensory and relay neurone, and relay to motor neurone.

4. Trace the efferent pathway. Follow the motor neurone from ventral horn, out through ventral root, to the effector.
5. Identify the effector. Muscle or gland.
6. State the sequence: stimulus → receptor → sensory neurone → relay neurone → motor neurone → effector → response.

Key anatomical landmarks: Dorsal root = sensory (afferent), ventral root = motor (efferent). Dorsal root ganglion contains sensory cell bodies. Grey matter = cell bodies and synapses; white matter = myelinated axons.

WARNING Common Mistakes

- Confusing dorsal (sensory) and ventral (motor) roots.
- Thinking the relay neurone cell body is in a ganglion (it is in the CNS grey matter).
- Forgetting that a reflex is involuntary – the brain is informed after the response.
- Calling the knee-jerk a "withdrawal reflex" (it is a **stretch reflex**, monosynaptic; withdrawal is polysynaptic).

DID YOU KNOW? Cameroon Context

In rural clinics, testing the **knee-jerk reflex** (patellar reflex) and **ankle-jerk reflex** helps assess spinal cord integrity (L2–L4 and S1–S2) after trauma from road accidents on the Douala–Yaoundé highway. Absent reflexes may indicate spinal shock or peripheral neuropathy (e.g. from poorly controlled diabetes).

WORKED EXAMPLE 14 The Knee-Jerk and the Withdrawal Reflex

A student accidentally touches a hot object in the laboratory and immediately withdraws her hand before feeling any pain.

- (a) Name the type of action described. (1 mark)
- (b) List, in order, the five components of the nervous pathway involved. (3 marks)
- (c) Explain why the hand is withdrawn before pain is consciously felt. (2 marks)
- (d) State two characteristics of this type of action. (2 marks)

SOLUTION

(a) This is a reflex action (specifically, a spinal withdrawal reflex, polysynaptic).

(b) The five components, in order, are:

1. Receptor: pain/heat receptors (free nerve endings) in the skin of the finger;
2. Sensory (afferent) neurone: carries the impulse from the receptor to the spinal cord via the dorsal root;
3. Relay (intermediate) neurone: in the grey matter of the spinal cord, transmits the impulse across a synapse to the motor neurone;
4. Motor (efferent) neurone: carries the impulse from the spinal cord via the ventral root to the effector;
5. Effector: the flexor muscles (e.g. biceps brachii) of the arm, which contract to withdraw the hand.

(c) The reflex pathway passes only through the spinal cord (three neurones, two synapses), which is short and fast. The impulse travelling up to the brain (where pain is consciously perceived) follows a longer ascending pathway with more synapses, so it arrives after the withdrawal has already occurred. This delay is protective: the body is removed from danger before the brain even registers the pain.

(d) Any two of: it is rapid; automatic/involuntary (does not require conscious thought); stereotyped (the same stimulus always gives the same response); protective in function; the pathway is inborn (not learnt).

EXPERIMENT Testing the Knee-Jerk Reflex

Aim: Demonstrate a monosynaptic stretch reflex.

Method: Subject sits with legs hanging freely. Examiner taps the patellar tendon sharply with a reflex hammer.

Observation: Sudden extension of the leg (quadriceps contraction).

Explanation: Tap stretches quadriceps muscle → muscle spindles stimulated → sensory neurone → synapse directly on motor neurone in spinal cord (monosynaptic) → motor neurone → quadriceps contracts → leg extends. No interneurone involved, hence very fast.

14.2 Skill 2: Calculating the Speed of Nerve Impulse Conduction**PROPERTY Factors Affecting Conduction Speed**

- **Myelination:** Myelin sheath (Schwann cells in PNS, oligodendrocytes in CNS) insulates the axon, forcing depolarisation to jump between **nodes of Ranvier (saltatory conduction)**. This dramatically increases speed (up to 100 m s^{-1} in humans).
- **Axon diameter:** Larger diameter $\rightarrow$ lower axial resistance $\rightarrow$ faster conduction.
- **Temperature:** Higher temperature increases ion diffusion rates and channel kinetics, increasing speed (Q10 effect).

METHOD Speed of Transmission Along a Neurone

1. Write the formula: speed $v = \frac{d}{t}$, where d is the distance travelled by the impulse and t is the time taken.
2. Convert all units to metres (m) and seconds (s) before substituting: $1 \text{ mm} = 10^{-3} \text{ m}$; $1 \text{ ms} = 10^{-3} \text{ s}$.
3. Substitute and compute. Express the answer in m s^{-1} , in standard form if appropriate.
4. Interpret the result: compare myelinated vs non-myelinated fibres; note that synapses introduce delays (about 0.5–1 ms each).
5. If several synapses are involved, subtract total synaptic delay from the measured time before calculating conduction speed along the axons.

WARNING Unit Conversion Traps

- Always convert before dividing: $90 \text{ mm} = 0.09 \text{ m}$, not $90/1000$ inside the formula.
- $1 \text{ ms} = 0.001 \text{ s}$, not 0.01 s .
- Speed unit is m s^{-1} (metres per second), not mm ms^{-1} .

WORKED EXAMPLE 15 Conduction Speed in a Reflex Arc

In an experiment, an impulse was found to travel along a myelinated sensory axon of length 90 mm in 1.5 ms.

- (a) Calculate the speed of conduction of the impulse along this axon. (3 marks)
- (b) The same impulse takes 4.5 ms in total to cross the whole reflex arc, which contains two synapses. If each synapse causes a delay of 1.0 ms, calculate the time spent in actual conduction along the neurones, and comment. (3 marks)

(c) State two structural features of a neurone that increase the speed of conduction. (2 marks)

SOLUTION

(a) Convert units first:

$$d = 90 \text{ mm} = 90 \times 10^{-3} \text{ m} = 0.09 \text{ m}; \quad t = 1.5 \text{ ms} = 1.5 \times 10^{-3} \text{ s}.$$

Apply the formula:

$$v = \frac{d}{t} = \frac{0.09}{1.5 \times 10^{-3}} = \frac{0.09}{0.0015} = 60 \text{ m s}^{-1}.$$

The speed of conduction is 60 m s^{-1} , a typical value for a myelinated mammalian neurone.

(b) Total synaptic delay = $2 \times 1.0 = 2.0 \text{ ms}$.

Time spent in conduction = $4.5 - 2.0 = 2.5 \text{ ms}$.

Comment: nearly half of the total reflex time (2.0 out of 4.5 ms) is lost at the synapses. Synaptic transmission (release, diffusion and action of neurotransmitter) is much slower than conduction along an axon; synapses are therefore the main factor limiting the speed of reflexes.

(c) Any two of:

- presence of a myelin sheath (allows saltatory conduction between nodes of Ranvier);
- large axon diameter (lower internal resistance to ion flow);
- presence of regularly spaced nodes of Ranvier.

14.3 Skill 3: Explaining Events at the Synapse

DEFINITION Synapse

A **synapse** is a specialised junction between two neurones (or a neurone and an effector) where an electrical impulse triggers the release of a chemical neurotransmitter, which diffuses across the **synaptic cleft** and binds to receptors on the post-synaptic membrane, potentially initiating a new impulse.

METHOD Describing Synaptic Transmission in Sequence (Cholinergic Synapse)

1. Arrival of the impulse: the action potential reaches the **pre-synaptic knob**; depolarisation opens voltage-gated **calcium channels**.
2. Calcium entry: Ca^{2+} ions diffuse into the knob down their electrochemical gradient.

3. Vesicle migration and fusion: Ca^{2+} causes **synaptic vesicles** (containing neurotransmitter, e.g. **acetylcholine**) to move to and fuse with the pre-synaptic membrane.
4. Release of transmitter: the neurotransmitter is released by exocytosis into the **synaptic cleft**.
5. Diffusion: the transmitter diffuses across the cleft (this causes the **synaptic delay** of $\approx 0.5\text{--}1\text{ ms}$).
6. Binding: transmitter molecules bind to specific **receptor proteins** on the post-synaptic membrane (ligand-gated ion channels).
7. New depolarisation (EPSP): sodium channels open, Na^+ enters, causing an **excitatory post-synaptic potential (EPSP)**; if threshold is reached, a new **action potential** is generated in the post-synaptic neurone.
8. Removal of transmitter: the enzyme **acetylcholinesterase** (AChE) rapidly hydrolyses acetylcholine to choline and acetate, terminating the signal and allowing the synapse to reset.

One-way transmission: vesicles and Ca^{2+} channels are only on the pre-synaptic side; receptors and AChE are only on the post-synaptic side.

PROPERTY Excitatory vs Inhibitory Synapses

- Excitatory (e.g. acetylcholine at neuromuscular junction): opens $\text{Na}^+/\text{Ca}^{2+}$ channels $\rightarrow$ depolarisation (EPSP).
- Inhibitory (e.g. GABA, glycine in CNS): opens Cl^- or K^+ channels $\rightarrow$ hyperpolarisation (IPSP), making the post-synaptic neurone less likely to fire.

Spatial summation (multiple simultaneous inputs) and **temporal summation** (rapid successive inputs) determine whether threshold is reached.

WARNING Synapse Exam Traps

- Forgetting Ca^{2+} entry – it is the essential trigger for vesicle fusion.
- Saying "impulse jumps across the gap" – it is chemical transmission, not electrical (except rare gap junctions).
- Confusing AChE (breaks down ACh) with ACh (the transmitter).
- Not mentioning that the post-synaptic potential is graded, not all-or-none; only the

resulting action potential is all-or-none.

DID YOU KNOW? Toxins and Synapses

Many poisons target synapses: curare (arrow poison) blocks ACh receptors → paralysis; strychnine blocks glycine receptors → convulsions; organophosphates (insecticides, nerve gases) inhibit AChE → continuous stimulation → spasm then paralysis. Botulinum toxin prevents vesicle fusion → flaccid paralysis (used medically in tiny doses for dystonia and cosmetics).

WORKED EXAMPLE 16 Synaptic Transmission and the Effect of a Toxin

The diagram of a cholinergic synapse shows structures labelled A (vesicles), B (synaptic cleft) and C (receptor proteins).

- (a) Describe how an impulse arriving at the pre-synaptic knob leads to the generation of an impulse in the post-synaptic neurone. (5 marks)
- (b) Explain why transmission across a synapse occurs in one direction only. (2 marks)
- (c) A certain pesticide inhibits the enzyme acetylcholinesterase. Predict and explain its effect on the post-synaptic neurone. (3 marks)

SOLUTION

(a) Award one mark for each correct step in logical order:

1. The action potential depolarises the pre-synaptic membrane, opening voltage-gated calcium channels, so Ca^{2+} enters the knob;
2. Ca^{2+} causes synaptic vesicles (A) containing acetylcholine to fuse with the pre-synaptic membrane;
3. Acetylcholine is released by exocytosis into the synaptic cleft (B) and diffuses across it;
4. Acetylcholine binds to specific receptor proteins (C) on the post-synaptic membrane, opening sodium channels;
5. Na^+ floods in, depolarising the post-synaptic membrane; if the threshold is reached, a new action potential is generated.

(b) Synaptic vesicles and transmitter are present only in the pre-synaptic knob, and specific receptor proteins are found only on the post-synaptic membrane. Transmitter can therefore be released and received in one direction only, so the impulse cannot travel backwards across the synapse.

(c) If acetylcholinesterase is inhibited, acetylcholine is not broken down in the cleft

and accumulates. It therefore binds repeatedly/continuously to the post-synaptic receptors. The post-synaptic membrane is continuously depolarised, producing prolonged, uncontrolled firing of the post-synaptic neurone (in muscles: continuous contraction/spasm, then paralysis). This is the basis of the toxicity of many insecticides and nerve gases.

14.4 Skill 4: The Iris Reflex and Control of Light Intensity

DEFINITION Pupil Reflex (Iris Reflex)

The **pupil reflex** is a brainstem reflex that adjusts the diameter of the pupil in response to light intensity falling on the retina, regulating the amount of light entering the eye. It is an autonomic (involuntary) reflex.

PROPERTY Antagonistic Muscles of the Iris

The iris contains two sets of smooth muscle fibres:

- Circular (sphincter) muscles: arranged concentrically around the pupil; contraction → **pupil constriction** (miosis). Innervated by **parasympathetic** fibres via the oculomotor nerve (CN III).
- Radial (dilator) muscles: arranged like spokes of a wheel; contraction → **pupil dilation** (mydriasis). Innervated by **sympathetic** fibres from the superior cervical ganglion.

They act antagonistically.

METHOD Explaining Pupil Size Changes

1. Identify the stimulus: change in light intensity detected by photoreceptors (rods/cones) in the retina.
2. Bright light: high intensity → strong stimulation → parasympathetic activation → circular muscles contract, radial muscles relax → pupil constricts → less light enters (protects retina from bleaching).
3. Dim light: low intensity → weak stimulation → sympathetic activation → radial muscles contract, circular muscles relax → pupil dilates → more light enters.
4. Control: reflex arc: retina → optic nerve → pretectal nucleus (midbrain) → Edinger-Westphal nucleus → oculomotor nerve → ciliary ganglion → iris muscles. Consensual

reflex: shining light in one eye constricts both pupils.

Do not confuse: the iris reflex (seconds, neural) with retinal adaptation to light/dark (minutes, photochemical – rhodopsin breakdown/resynthesis).

WARNING Common Errors

- Saying "circular muscles dilate the pupil" – they constrict it.
- Attributing the reflex to the spinal cord – the centre is in the midbrain (pretectal area).
- Confusing the iris reflex (pupil size) with accommodation (lens shape) – different effectors, different pathways.

DID YOU KNOW? Clinical Relevance

Doctors test the pupillary light reflex to assess brainstem function (CN II and III). Fixed, dilated pupils ("blown pupils") indicate severe brainstem compression (e.g. from raised intracranial pressure after head trauma). In Cameroon, this is a vital sign checked in emergency wards in Yaoundé and Douala.

WORKED EXAMPLE 17 From a Dark Cinema into Bright Sunshine

A student walks out of a dark cinema hall in Yaoundé into bright midday sunlight. At first she can hardly see and shades her eyes; after a few minutes her vision becomes comfortable.

- (a) Describe and explain the immediate change that occurs in her pupils on entering the bright light. (3 marks)
- (b) Explain why the pupil reflex alone cannot fully account for her comfortable vision after several minutes. (3 marks)
- (c) Name the pigment involved in the slower adjustment and state where it is found. (2 marks)

SOLUTION

(a) In bright light, the pupil constricts (becomes smaller). The circular muscles of the iris contract while the radial muscles relax. This reduces the amount of light entering the eye and protects the retina from damage by excessive light. It is a reflex action controlled by the autonomic nervous system (parasympathetic branch), with the receptor being the retina and the effector the iris muscles.

(b) The pupil can only vary light entry over a limited range (its area changes by a factor

of about 16 at most), whereas light intensity outdoors can be thousands of times greater than in the dark hall. Full adjustment requires retinal (light) adaptation: in bright light the visual pigment rhodopsin is bleached (broken down into retinal and opsin) faster than it is resynthesised, so the rods become less sensitive and vision shifts to the cones, which work well in bright light. This photochemical adjustment takes several minutes, which matches the gradual improvement described.

(c) The pigment is rhodopsin (visual purple), found in the rod cells of the retina. (Cone pigments – iodopsins – mediate bright-light and colour vision.)

14.5 Skill 5: Accommodation and Defects of Vision

DEFINITION Accommodation

Accommodation is the process by which the eye changes the focal length of its lens to focus images of objects at different distances sharply on the retina.

PROPERTY Mechanism of Accommodation

- Near object (e.g. reading at 25 cm): ciliary muscles contract → suspensory ligaments slacken → elastic lens becomes more convex (shorter focal length, higher power) → diverging rays focused on retina.
- Distant object (parallel rays): ciliary muscles relax → suspensory ligaments become taut → lens pulled flatter (longer focal length, lower power) → parallel rays focused on retina.

The lens is naturally elastic and rounded; the suspensory ligaments keep it flattened when ciliary muscles are relaxed.

DEFINITION Defects of Vision

- Myopia (short sight): eyeball too long or lens/cornea too powerful → image forms in front of retina for distant objects. Corrected by diverging (concave) lens which diverges rays before they enter the eye.
- Hypermetropia (long sight): eyeball too short or lens/cornea too weak → image forms behind retina for near objects (and sometimes distant). Corrected by converging (convex) lens which converges rays before they enter the eye.
- Presbyopia: age-related loss of lens elasticity and ciliary muscle power → difficulty

with near vision. Corrected by converging lenses (reading glasses).

- Astigmatism: uneven curvature of cornea/lens → rays in different planes focus at different points. Corrected by cylindrical lenses.

METHOD Explaining Accommodation and Correcting Defects

1. For accommodation: state ciliary muscle action → ligament tension → lens shape → focal length → image on retina.
2. For defects: always state (i) where the image falls relative to retina, (ii) the cause (eye-ball length or lens power), (iii) the correcting lens type and its effect on incident rays.
3. Diagram: draw rays from object → correcting lens → eye lens → retina.

WARNING Exam Tips

- "Short sight = see near, not far" – the name describes what you can see.
- Concave lens = diverging = thinner at centre; Convex lens = converging = thicker at centre.
- In accommodation for near vision, the ciliary muscles contract (many students wrongly say relax).

WORKED EXAMPLE 18 A Student Who Cannot Read the Board

A Form Five student can read her book clearly but cannot read writing on the classroom board unless she sits in the front row.

- (a) Name the defect of vision she suffers from. (1 mark)
- (b) Explain, in terms of where images are focused, why she cannot see distant objects clearly. (2 marks)
- (c) State the type of spectacle lens that would correct this defect and explain how it works. (2 marks)
- (d) Describe how her eye accommodates when she reads her book at 25 cm. (3 marks)

SOLUTION

- (a) Short sightedness (myopia): near objects are seen clearly, distant objects are blurred.
- (b) In her eye, parallel rays from a distant object are brought to a focus in front of the retina (because the eyeball is too long or the lens/cornea system converges the rays too

strongly). By the time the rays reach the retina they have diverged again, so the image on the retina is blurred.

(c) A diverging (concave) lens is used. It spreads the incoming parallel rays slightly apart before they enter the eye, so that the eye's own lens then focuses them exactly on the retina instead of in front of it, giving a sharp image of distant objects.

(d) When reading at 25 cm (near object):

1. the ciliary muscles contract;
2. the suspensory ligaments slacken, releasing tension on the lens;
3. the elastic lens bulges, becoming more convex (shorter focal length), so the diverging rays from the near book are focused sharply on the retina.

14.6 Skill 6: Analysing Hormonal Control and Negative Feedback

DEFINITION Hormone

A **hormone** is a chemical messenger secreted by an **endocrine gland** directly into the bloodstream, which carries it to specific **target cells** bearing complementary receptors, where it exerts a regulatory effect.

PROPERTY Mechanism of Hormone Action

- Protein/peptide hormones (e.g. insulin, adrenaline, ADH, FSH, LH): water-soluble, cannot cross phospholipid bilayer. Bind to cell surface receptors → activate second messenger (usually **cyclic AMP (cAMP)** via adenyl cyclase) → cascade of enzyme activations → rapid response (seconds to minutes).
- Steroid hormones (e.g. oestrogen, testosterone, cortisol, aldosterone): lipid-soluble, diffuse through membrane. Bind to intracellular receptors (cytoplasm or nucleus) → hormone-receptor complex acts as transcription factor on DNA → alters gene expression → new protein synthesis → slower response (hours), longer lasting.

PROPERTY Negative Feedback

Negative feedback is the principal mechanism of hormonal homeostasis: a rise in the controlled variable stimulates hormone secretion that lowers the variable; as the variable returns to normal, the stimulus for secretion diminishes, switching off further hormone release. This maintains the variable within narrow limits (set point).

METHOD Interpreting a Hormone/Feedback Question

1. Identify the gland and hormone from the symptoms or data given (e.g. high blood glucose → insulin from pancreas β -cells; goitre → thyroid/thyroxine).
2. State the stimulus that triggers secretion (e.g. rising blood glucose detected by β -cells).
3. State the target organ(s) and effect (e.g. liver and muscles convert glucose to glycogen; cells increase glucose uptake).
4. Describe the feedback loop: when the controlled factor returns to normal, the change is detected and hormone secretion is switched off/reduced – this is negative feedback.
5. Contrast with the antagonistic hormone where relevant (insulin vs glucagon; thyroxine vs TSH; they act in opposite directions to keep the variable constant).
6. Mechanism of action: specify whether the hormone acts via membrane receptor/second messenger (protein) or intracellular receptor/gene activation (steroid).

DID YOU KNOW? Cameroon Context

Diabetes mellitus is a growing public health concern in Cameroon (urbanisation, diet change). Type 1 = autoimmune destruction of β -cells (no insulin); Type 2 = insulin resistance (receptors defective). Goitre (iodine deficiency) was endemic in the Adamawa and North regions; salt iodisation programmes have reduced prevalence. Adrenaline surge during "fright" (e.g. encountering a snake on Mount Cameroon) illustrates neuroendocrine interaction.

WORKED EXAMPLE 19 Blood Glucose After a Meal of Fufu and Eru

After a heavy meal rich in carbohydrate, a student's blood glucose concentration rose from 90 mg per 100 cm³ to 140 mg per 100 cm³, then fell back to about 90 mg per 100 cm³ within two hours.

- (a) Name the hormone responsible for the fall and the gland that secretes it. (2 marks)
- (b) Explain how this hormone brings about the fall in blood glucose. (3 marks)
- (c) Explain what prevents the blood glucose from falling far below the normal level. (3 marks)
- (d) The hormone in (a) is a protein. Explain why it cannot enter its target cells and state how its message is transmitted inside the cell. (2 marks)

SOLUTION

(a) The hormone is insulin, secreted by the β -cells of the islets of Langerhans in the pancreas.

(b) Insulin lowers blood glucose by:

- stimulating the liver and muscles to convert excess glucose into glycogen for storage (glycogenesis);
- increasing the uptake and respiration of glucose by cells (increased membrane permeability to glucose via GLUT4 translocation);
- promoting conversion of glucose to fat in adipose tissue (lipogenesis).

(c) This is negative feedback: as blood glucose falls towards normal, the stimulus for insulin secretion disappears, so insulin release is switched off. In addition, if glucose falls below normal, the α -cells of the pancreas secrete glucagon, which is antagonistic to insulin: it stimulates the liver to convert stored glycogen back into glucose (glycogenolysis) and release it into the blood, and promotes gluconeogenesis. The two hormones together keep blood glucose fluctuating narrowly around the set point.

(d) Insulin is a large, water-soluble protein molecule and cannot cross the phospholipid bilayer of the cell surface membrane. It therefore binds to a specific receptor on the membrane (a receptor tyrosine kinase); this activates the enzyme adenyl cyclase inside the cell, which produces the second messenger cyclic AMP (cAMP), which then triggers the internal response (e.g. activation of enzymes for glycogen synthesis).

14.7 Skill 7: Designing and Interpreting Tropism Experiments

DEFINITION Tropism

A **tropism** is a directional growth response of a plant to a unidirectional external stimulus. Positive tropism = growth towards the stimulus; negative tropism = growth away. Main types: **phototropism** (light), **geotropism** (gravity), **hydrotropism** (water), **thigmotropism** (touch).

DEFINITION Nastic Movement

A **nastic movement** is a non-directional response to a stimulus (e.g. **photonasty** – flower opening/closing with light intensity; **seismonasty** – Mimosa leaf folding on touch). The response direction is determined by plant structure, not stimulus direction.

PROPERTY Role of Auxin (IAA) in Phototropism and Geotropism

- Shoots (positive phototropism): Auxin produced in tip → redistributed to shaded side → promotes cell elongation on shaded side → shoot bends towards light.
- Roots (positive geotropism): Auxin redistributed to lower side → root cells are more sensitive – auxin inhibits elongation on lower side → upper side elongates more → root bends downwards.

Acid growth hypothesis: Auxin activates proton pumps → wall acidification → expansins loosen cellulose microfibrils → turgor-driven elongation.

METHOD Investigating Phototropism / Geotropism

1. Define variables: independent = direction of light (or gravity); dependent = angle/direction of growth; controls = temperature, water, seedling type, age, uniform light for control.
2. Always include a control (seedling in uniform light/darkness or vertical) to show bending is a response to direction of stimulus.
3. Use the auxin explanation: production in tip, lateral redistribution, differential elongation.
4. Classic experiments to quote:
 - Darwin: tip perceives light; covering tip prevents bending.
 - Boysen-Jensen: tip separated by mica (impermeable) blocks bending; gelatin (permeable) allows bending → chemical messenger.
 - Went: tip on agar block → auxin diffuses into agar → agar placed asymmetrically on decapitated stump causes bending without light → auxin identified as the messenger.

WARNING Tropism Experiment Pitfalls

- Forgetting the control (uniform light).
- Saying auxin "moves away from light" – it is redistributed to the shaded side (active transport).
- Confusing shoot and root responses to auxin (shoots: promotes elongation; roots: inhibits at same concentration).

- Not mentioning that the tip is the site of both perception and auxin synthesis.

DID YOU KNOW? Mount Cameroon Forest Floor

On the shaded slopes of Mount Cameroon, seedlings of *Maesopsis eminii* or *Terminalia superba* exhibit strong positive phototropism. Auxin-mediated bending towards fleeting sunflecks maximises photosynthetic gain in the understorey, a key survival strategy in the competitive rainforest environment.

WORKED EXAMPLE 20 Phototropism in Maize Seedlings

Three groups of maize coleoptiles were treated as follows and left for 48 hours with light shining from one side only:

Group P: tips left intact. Group Q: tips cut off. Group R: tips cut off and replaced with a thin agar block between tip and stump.

Results: P bent towards the light; Q did not bend and grew little; R bent towards the light.

- State the aim of this experiment. (1 mark)
- Explain the result obtained in group P. (3 marks)
- Explain the results of groups Q and R, and state the conclusion that can be drawn about the role of the tip. (4 marks)
- State the biological advantage of this response to a plant growing on a shaded forest floor on Mount Cameroon. (2 marks)

SOLUTION

(a) To investigate the role of the shoot tip (coleoptile tip) in the phototropic response (bending of a shoot towards unilateral light).

(b) In group P, light from one side causes auxin (IAA) produced in the tip to be redistributed to the shaded side of the coleoptile. Auxin stimulates cell elongation in shoots, so cells on the shaded side elongate more than those on the lit side. This unequal growth causes the coleoptile to bend towards the light (positive phototropism).

(c) In group Q, removing the tip removes the source of auxin; without auxin there is little elongation growth and no differential growth, so the coleoptile neither grows nor bends. In group R, the agar block is permeable to auxin: auxin produced in the replaced tip diffuses down through the agar into the stump, is redistributed by the unilateral light, and bending towards the light is restored.

Conclusion: the tip is the site of auxin production (and light perception); auxin is a chemical messenger that moves downwards from the tip and controls the growth response. The response depends on a diffusible substance, not on the tip cells themselves being present at the bending zone.

(d) Positive phototropism makes the shoot grow towards the light source, positioning

the leaves in brighter light. This maximises light absorption for photosynthesis, giving the seedling a competitive advantage in the dim, shaded conditions of the forest floor, where light is the main limiting factor.

14.8 Skill 8: Comparing Nervous and Hormonal Coordination

PROPERTY Comparison of Nervous and Hormonal Coordination

Criterion	Nervous Coordination	Hormonal Coordination
Nature of message	Electrical impulses (action potentials) along axons; chemical (neurotransmitter) at synapses	Chemical (hormones)
Pathway	Along specific neurones (fixed, wired pathways)	Through the bloodstream (circulation)
Speed of transmission	Very fast (up to $\approx 100 \text{ m s}^{-1}$ in myelinated fibres)	Slow (limited by blood flow, seconds to minutes)
Duration of response	Short-lived, brief (milliseconds to seconds)	Longer-lasting (minutes to days)
Spread of response	Localised (specific effector organ)	Widespread (all target organs with receptors reached by blood)
Target specificity	High (specific neurone-to-effector connection)	High (only cells with specific receptors respond)
Example	Withdrawal reflex, knee-jerk	Blood glucose regulation (insulin/glucagon), metamorphosis

METHOD Building a Comparison Answer

1. Choose clear criteria and compare both systems under each one (never describe one system fully and then the other).
2. Use these criteria: nature of message, pathway, speed, duration, spread, target specificity.
3. Give one example of each (reflex action vs blood glucose control).

4. Note the overlap: some responses use both (e.g. the adrenal medulla is stimulated by a nerve but releases adrenaline into the blood – neuroendocrine link; the hypothalamus links the two systems).

WARNING Comparison Traps

- Saying hormones are "slow" without qualifying (blood flow limited).
- Saying nervous system is "only electrical" – synapses are chemical.
- Forgetting that both systems can be specific (hormones only affect target cells with receptors).
- Not mentioning the neuroendocrine integration (hypothalamus–pituitary, adrenal medulla).

WORKED EXAMPLE 21 Nervous versus Hormonal Coordination

- (a) Construct a table comparing nervous and hormonal coordination under four headings. (4 marks)
- (b) A man touches a thorn and withdraws his hand; later the same day, a fright causes his heart to race for several minutes. For each event, state the type of coordination involved and justify your answer. (4 marks)

SOLUTION

(a)

Criterion	Nervous coordination	Hormonal coordination
Nature of message	Electrical impulses (and chemical transmitter at synapses)	Chemical (hormones)
Pathway	Along neurones (fixed pathways)	Through the bloodstream
Speed of transmission	Very fast (up to about 100 m s^{-1})	Slow (depends on blood flow)
Duration of response	Short-lived, brief	Longer-lasting
Spread of response	Localised (specific effector)	Widespread (all target organs reached by blood)

(b) Withdrawal from the thorn: nervous coordination (a spinal reflex). Justification: the response is very rapid, brief and localised to the arm muscles, and travels along a fixed

pathway of neurones – all characteristic of nervous control.

Racing heart after fright: mainly hormonal coordination. The fright causes the adrenal medulla to release adrenaline into the blood; adrenaline reaches the heart via the circulation and increases heart rate. The effect is slower in onset but persists for several minutes and affects many organs (heart, liver, muscles) – characteristic of hormonal control. (Note: the initial stimulation of the adrenal medulla is nervous, showing that the two systems interact.)

14.9 Skill 9: The Ear – Hearing and Balance

DEFINITION Ear Structure and Function

The ear is a sensory organ for **hearing** (detection of sound waves) and **balance** (detection of head position and movement). It has three regions: outer, middle, inner.

PROPERTY Hearing Mechanism

1. Outer ear: pinna collects sound waves → auditory canal → **tympanic membrane** (eardrum) vibrates.
2. Middle ear: three **ossicles** (malleus, incus, stapes) amplify and transmit vibrations → **oval window** (membrane on cochlea). **Eustachian tube** equalises pressure.
3. Inner ear (cochlea): stapes vibrates oval window → pressure waves in **perilymph** → **basilar membrane** vibrates → **hair cells** of **organ of Corti** bent against **tectorial membrane** → generator potentials → action potentials in **auditory nerve** (CN VIII) → auditory cortex.
4. Frequency discrimination: high frequencies peak near base (oval window), low frequencies near apex (place theory).

PROPERTY Balance Mechanism

- Static balance (position): **utricle** and **sacculle** (vestibule) contain **maculae** with hair cells covered by **otoliths** (calcium carbonate crystals). Gravity pulls otoliths → bends hair cells → signals head position.
- Dynamic balance (movement): three **semicircular canals** (mutually perpendicular) contain **ampullae** with **cristae** (hair cells in gelatinous cupula). Rotational acceleration

moves endolymph → deflects cupula → bends hair cells → signals rotation.

WARNING Ear Exam Points

- Ossicles amplify force $\approx 20\times$ (lever action) and match impedance (air to fluid).
- Cochlea unrolled is a tube with basilar membrane; hair cells are the receptors.
- Deafness types: conductive (outer/middle ear blockage, ossicle fusion) vs sensorineural (hair cell/nerve damage).
- Balance and hearing both use hair cells (mechanoreceptors) but different structures.

WORKED EXAMPLE 22 Hearing and Balance

- (a) Describe how sound waves are converted into nerve impulses in the cochlea. (4 marks)
- (b) Explain how the semicircular canals detect rotational movement of the head. (3 marks)
- (c) A person suffers from dizziness and nausea after a severe cold. Suggest a possible explanation. (2 marks)

SOLUTION

(a) Sound waves cause the tympanic membrane to vibrate. Ossicles amplify and transmit vibrations to the oval window. Pressure waves in the perilymph cause the basilar membrane to vibrate. Hair cells of the organ of Corti are bent against the tectorial membrane, generating generator potentials. If threshold is reached, action potentials are generated in the auditory nerve fibres and transmitted to the brain.

(b) The three semicircular canals are oriented in three perpendicular planes. Each has an ampulla containing a crista (hair cells embedded in a gelatinous cupula). During rotational acceleration, the endolymph lags behind due to inertia, pushing the cupula and bending the hair cells. This generates nerve impulses proportional to the rate of rotation, informing the brain of the direction and speed of head rotation.

(c) The cold may have caused inflammation blocking the Eustachian tube, preventing pressure equalisation in the middle ear, or spread to the inner ear (labyrinthitis) affecting the vestibular apparatus (semicircular canals/utricle/sacculle), disrupting balance signals and causing vertigo/nausea.

14.10 Skill 10: The Brain, Spinal Cord and Autonomic Nervous System

DEFINITION Central Nervous System (CNS)

The CNS consists of the **brain** and **spinal cord**. It is the site of integration, processing, and command. Protected by meninges (dura mater, arachnoid, pia mater) and cerebrospinal fluid (CSF).

PROPERTY Major Brain Regions and Functions

- Cerebrum (cerebral hemispheres): higher functions – consciousness, memory, reasoning, voluntary movement, sensory perception. Grey matter (cortex) outside, white matter inside.
- Cerebellum: coordination of movement, balance, posture, motor learning. "Little brain" posterior to brainstem.
- Medulla oblongata: vital reflex centres – cardiac, respiratory, vasomotor, vomiting, coughing.
- Hypothalamus: homeostasis – temperature, water balance (ADH), hunger, endocrine control via pituitary.
- Pituitary gland: "master gland" – anterior (trophic hormones: TSH, ACTH, FSH, LH, GH, prolactin) and posterior (ADH, oxytocin).
- Midbrain: reflex centres for vision (pupil reflex) and hearing.

PROPERTY Spinal Cord

- Runs through vertebral column; segmental organisation (31 pairs of spinal nerves).
- Grey matter (central): butterfly-shaped; cell bodies of motor neurones (ventral horn), interneurones, sensory fibre terminals (dorsal horn).
- White matter (peripheral): myelinated axons forming ascending (sensory) and descending (motor) tracts.
- Function: reflex centre (spinal reflexes); conduction pathway to/from brain.

PROPERTY Autonomic Nervous System (ANS)

Involuntary control of cardiac muscle, smooth muscle, glands. Two antagonistic divisions:

Feature	Sympathetic	Parasympathetic
Origin	Thoracolumbar (T1–L2)	Craniosacral (CN III, VII, IX, X; S2–S4)
Pre-ganglionic fibre	Short	Long
Post-ganglionic fibre	Long	Short
Ganglia	Paravertebral chain, prevertebral	Near/on effector organs
Neurotransmitter (pre)	Acetylcholine	Acetylcholine
Neurotransmitter (post)	Noradrenaline (mostly)	Acetylcholine
General effect	"Fight or flight" – arousal, energy mobilisation	"Rest and digest" – conservation, restoration
Heart rate	Increases	Decreases
Pupil	Dilates (radial muscle)	Constricts (circular muscle)
Gut motility	Decreases	Increases

DID YOU KNOW? Neuroendocrine Integration

The hypothalamus is the key link: it receives neural inputs (e.g. stress, temperature, osmotic pressure) and outputs hormonal signals to the pituitary. The adrenal medulla is a modified sympathetic ganglion – preganglionic fibres stimulate chromaffin cells to secrete adrenaline/noradrenaline into blood (neuroendocrine transducer).

14.11 Skill 11: Plant Hormones – Auxins, Gibberellins, Cytokinins, Absciscic Acid, Ethylene

DEFINITION Plant Hormones (Phytohormones)

Chemical messengers produced in one part of the plant, transported (often via phloem or diffusion) to target tissues where they regulate growth and development at very low concentrations. Major classes:

PROPERTY Major Plant Hormones and Functions

Hormone	Main Sites of Synthesis	Key Functions
Auxins (IAA)	Shoot apex, young leaves, developing seeds	Cell elongation (phototropism, geotropism), apical dominance, root initiation (cuttings), fruit development, prevents abscission
Gibberellins (GA)	Young leaves, shoot tips, seeds, roots	Stem elongation (bolting), seed germination (mobilises reserves), breaks dormancy, fruit growth, flowering
Cytokinins	Root tips, developing seeds, fruits	Cell division (cytokinesis), delays senescence (leaf aging), promotes lateral bud growth (releases apical dominance), chloroplast development
Abscisic Acid (ABA)	Leaves, roots, seeds	Stomatal closure (water stress), seed dormancy, inhibits growth, promotes leaf abscission
Ethene (Ethylene)	Ripening fruits, senescing leaves, stressed tissues	Fruit ripening, leaf/flower/fruit abscission, senescence, stress response, flower sex determination

METHOD Commercial Applications of Plant Hormones

1. Auxins: rooting powders for cuttings (IBA, NAA); selective herbicides (2,4-D, MCPA – broadleaf weeds); parthenocarpic fruit set (seedless tomatoes).
2. Gibberellins: increase grape bunch size and berry size; break dormancy in barley for malting (beer); induce flowering in long-day plants.
3. Cytokinins: tissue culture (shoot proliferation); delay leaf senescence in cut flowers/vegetables.
4. Abscisic Acid: potential antitranspirant (spray to close stomata during drought).
5. Ethylene: controlled fruit ripening (bananas, mangoes in Cameroon markets); de-greening citrus; synchronising flowering in pineapples.

WARNING Hormone Interaction

Plant responses usually result from hormone ratios, not single hormones. E.g. apical dominance: high auxin:cytokinin ratio suppresses lateral buds; removing apex lowers auxin, allowing cytokinin to promote branching. Root:shoot ratio in tissue culture controlled by auxin:cytokinin balance.

DID YOU KNOW? Cameroon Agriculture

In Cameroon, ethylene is used to ripen plantains and bananas uniformly for urban markets (Douala, Yaoundé). Gibberellic acid improves germination of uniform barley for the brewing industry (SOCABO). Auxin-based herbicides (2,4-D) control broadleaf weeds in maize and rice fields. Cytokinins in tissue culture labs (IRAD, universities) mass-propagate disease-free plantain and cassava planting material.

14.12 Skill 12: Light and Plant Growth – Photoperiodism and Phytochrome**DEFINITION** Photoperiodism

Photoperiodism is the response of plants to the relative lengths of day and night (photoperiod), particularly controlling flowering. The critical factor is the length of the uninterrupted dark period.

PROPERTY Photoperiodic Classification

- Short-day plants (SDPs): flower when night length exceeds a critical duration (e.g. chrysanthemum, poinsettia, rice, soybean, cotton). Actually "long-night plants".
- Long-day plants (LDPs): flower when night length is shorter than a critical duration (e.g. wheat, barley, spinach, clover).
- Day-neutral plants: flower regardless of photoperiod (e.g. tomato, cucumber, maize).

Night-break experiment: a brief flash of light during the long night prevents flowering in SDPs but promotes it in LDPs → proves dark period is critical.

PROPERTY Phytochrome – The Photoreceptor

Phytochrome is a blue–green pigment (protein + chromophore) existing in two interconvertible forms:

- Pr (red-absorbing, $\lambda_{\max} \approx 660 \text{ nm}$): inactive form.
- Pfr (far-red absorbing, $\lambda_{\max} \approx 730 \text{ nm}$): active form.

Conversions: Red light (660 nm) $\rightarrow$ Pr $\xrightarrow{\text{absorbs}}$ Pfr (fast). Far-red light (730 nm) $\rightarrow$ Pfr $\xrightarrow{\text{absorbs}}$ Pr (fast). In darkness, Pfr slowly reverts to Pr (or is degraded).

METHOD Explaining Photoperiodic Response via Phytochrome

1. Daylight: rich in red $\rightarrow$ mostly Pfr at sunset.
2. Night: Pfr slowly reverts to Pr.
3. SDPs: need low Pfr (long night) to flower. If night is interrupted by red light $\rightarrow$ Pfr rises $\rightarrow$ flowering inhibited. Far-red after red cancels the effect.
4. LDPs: need high Pfr (short night) to flower.
5. Flowering hormone (florigen): produced in leaves under appropriate photoperiod, transported via phloem to shoot apex $\rightarrow$ induces flowering.

WARNING Phytochrome Details

- Pr $\leftrightarrow$ Pfr interconversion is rapid (seconds). Dark reversion is slow (hours).
- The last light quality received determines the response (red $\rightarrow$ promotes LDP, inhibits SDP; far-red $\rightarrow$ opposite).
- Phytochrome also controls seed germination (lettuce needs red light), de-etiolation, shade avoidance.

DID YOU KNOW? Cameroon Crops and Photoperiod

Rice (short-day) flowers as days shorten after the June solstice in the Sudano-Sahelian zone. Soybean varieties are selected for specific critical night lengths to match planting dates in different agro-ecological zones. Chrysanthemum growers in the western high-lands use blackout curtains to induce flowering year-round for the cut-flower market.

WORKED EXAMPLE 23 Photoperiodism and Phytochrome

- (a) Explain why a short-day plant fails to flower if its long night is interrupted by a brief flash of red light, but flowers if the red flash is followed by a far-red flash. (4 marks)
- (b) A farmer wants to delay flowering in a long-day crop (e.g. spinach) to extend the vegetative harvest. Suggest a light treatment and explain its effect. (3 marks)
- (c) Name two other plant responses controlled by phytochrome besides flowering. (2 marks)

SOLUTION

- (a) In a short-day plant, flowering requires a long uninterrupted night during which Pfr reverts to Pr to a low level. A red light flash converts remaining Pr to Pfr, raising the Pfr level → the plant "perceives" a short night → flowering inhibited. A subsequent far-red flash converts Pfr back to Pr, restoring the low Pfr level → the plant "perceives" a long night → flowering restored.
- (b) To delay flowering in a long-day plant (which needs short nights/high Pfr), give long nights or end the day with far-red light (or give a far-red pulse during the night). This ensures low Pfr at the end of the night, so the plant does not receive the inductive signal for flowering. Vegetative growth continues.
- (c) Any two of: seed germination (e.g. lettuce requires red light), de-etiolation (greening, inhibition of hypocotyl elongation in seedlings), shade avoidance (stem elongation in response to low red:far-red ratio under canopy), leaf abscission, dormancy.

14.13 Skill 13: Photoreception – Rods and Cones**DEFINITION** Photoreception

Photoreception is the process by which light energy is converted into nerve impulses in the retina. The retina contains two types of photoreceptor cells: **rods** and **cones**.

PROPERTY Comparison of Rods and Cones

Feature	Rods	Cones
Visual pigment	Rhodopsin (visual purple)	Iodopsins (photopsins) – three types (S, M, L)
Sensitivity	Very high (dim light, scotopic vision)	Low (bright light, photopic vision)
Colour vision	None (monochromatic)	Yes (trichromatic – blue, green, red)
Acuity (detail)	Low (many rods converge on one bipolar cell)	High (one-to-one connection in fovea)
Distribution	Peripheral retina (absent from fovea)	Concentrated in fovea (yellow spot)
Adaptation	Dark adaptation (slow, 20–30 min)	Light adaptation (fast, seconds)

PROPERTY Mechanism of Phototransduction (Rods)

1. Dark (resting): Na^+ channels open (cGMP-gated) → "dark current" inward → cell depolarised (≈ -40 mV) → continuous glutamate release (inhibitory) → bipolar cells inhibited.
2. Light hits rhodopsin: retinal (11-cis) → all-trans retinal → conformational change → activates **transducin** (G-protein) → activates **phosphodiesterase (PDE)** → hydrolyses cGMP → cGMP levels fall → Na^+ channels close → cell hyperpolarises → glutamate release stops → bipolar cells excited → signal to ganglion cells → optic nerve.

Key point: Light causes hyperpolarisation (not depolarisation) of photoreceptors.

WARNING Photoreception Traps

- Saying rods/cones "fire action potentials" – they produce graded potentials; only ganglion cells fire action potentials.
- Confusing rhodopsin (rods) with iodopsins (cones).
- Thinking the fovea has rods – it is rod-free, cone-only, for maximum acuity in bright light.
- Forgetting that photoreceptors hyperpolarise in response to light (the opposite of most sensory receptors).

METHOD Answering a Rods vs Cones Question

1. State the stimulus conditions (dim vs bright light) and match to the receptor type.
2. Link structure to function: pigment type → sensitivity; neural wiring (convergence vs one-to-one) → acuity; retinal distribution → where each works best.
3. For colour vision: three cone types with different iodopsins; the brain compares their relative stimulation (trichromatic theory).
4. For adaptation: dark adaptation = rhodopsin resynthesis (slow); light adaptation = rhodopsin bleaching (fast).

DID YOU KNOW? Night Vision and Vitamin A

The retinal part of rhodopsin is derived from vitamin A. Vitamin A deficiency – still observed where diets lack green leafy vegetables, palm oil or liver – causes night blindness (nyctalopia): rhodopsin cannot be resynthesised fast enough, so rod vision fails in dim light. Red palm oil, abundant in Cameroonian cooking, is an excellent source of provitamin A carotenoids.

WORKED EXAMPLE 24 Rods, Cones and Night Vision

- (a) Explain why, when looking at a faint star at night, it is easier to see it by looking slightly to one side of it rather than directly at it. (3 marks)
- (b) Explain why colours cannot be distinguished in very dim light. (2 marks)
- (c) Describe the events in a rod cell between the absorption of a photon and the generation of a signal to the brain. (4 marks)

SOLUTION

- (a) The fovea (centre of the retina, where the image falls when we look directly at an object) contains only cones, which have low sensitivity to dim light. The rods, which are highly sensitive in dim light, are found in the peripheral retina. Looking slightly to one side makes the star's image fall on a rod-rich region of the retina, so the faint light is detected.
- (b) Colour vision depends on the three types of cones (with different iodopsins), but cones are insensitive in dim light and do not respond. Only rods function, and rods contain a single pigment (rhodopsin) and cannot distinguish wavelengths; hence vision in dim light is monochromatic (shades of grey).
- (c) Award one mark for each correct step:
1. A photon is absorbed by rhodopsin; 11-cis retinal is converted to all-trans retinal,

changing the shape of the opsin (bleaching);

2. Activated rhodopsin activates the G-protein transducin, which activates phosphodiesterase;
3. Phosphodiesterase hydrolyses cGMP, so cGMP-gated sodium channels close and the dark current stops;
4. The rod cell hyperpolarises, glutamate release decreases, which excites the connected bipolar cell; the signal passes to ganglion cells, which generate action potentials transmitted along the optic nerve to the brain.

KEY POINTS Key Points of the Chapter

- A reflex arc has five components: receptor → sensory neurone → relay neurone → motor neurone → effector; dorsal root = sensory, ventral root = motor.
- Conduction speed $v = d/t$; convert mm and ms before calculating; synapses cause delays of about 0.5–1 ms each.
- Synaptic transmission: impulse → Ca^{2+} entry → vesicle fusion → transmitter release → diffusion → receptor binding → EPSP → AChE removes transmitter. Transmission is one-way.
- The iris reflex uses antagonistic circular (constrict, parasympathetic) and radial (dilate, sympathetic) muscles; accommodation uses ciliary muscles and suspensory ligaments to change lens shape.
- Myopia (image in front of retina) is corrected by a concave lens; hypermetropia (image behind retina) by a convex lens.
- Hormones act via membrane receptors and second messengers (proteins) or intracellular receptors and gene activation (steroids); negative feedback keeps controlled variables near a set point.
- Tropisms are directional growth responses; auxin is redistributed and causes differential elongation (promotes in shoots, inhibits in roots).
- Nervous coordination is fast, brief and localised; hormonal coordination is slower, longer-lasting and widespread; the hypothalamus and adrenal medulla link the two systems.
- The ear detects sound (cochlea, organ of Corti) and balance (utricle, saccule, semi-circular canals) using hair cells.
- Plant hormones (auxins, gibberellins, cytokinins, ABA, ethylene) control growth and have major agricultural applications in Cameroon.

- Photoperiodism is controlled by night length via the phytochrome system ($Pr \leftrightarrow Pfr$); the last light quality received determines the flowering response.
- Rods (rhodopsin, dim light, peripheral retina, no colour) and cones (iodopsins, bright light, fovea, colour, high acuity); light hyperpolarises photoreceptors.

15 EXERCISES**15.1 I check my knowledge****EXERCISE 12 Multiple Choice Questions**

For each question, choose the correct answer (A, B, C or D).

1. Which of the following structures in the eye is responsible for detecting light intensity and initiating the pupillary reflex?
 - A Rods and cones in the retina
 - B Ganglion cells in the retina
 - C Bipolar cells in the retina
 - D Pigmented epithelium
2. During the depolarisation phase of an action potential in a myelinated axon, which ion movement occurs?
 - A Na^+ ions move out of the axon
 - B K^+ ions move into the axon
 - C Na^+ ions move into the axon
 - D Ca^{2+} ions move out of the axon
3. The hormone that stimulates the conversion of glycogen to glucose in the liver is:
 - A Insulin
 - B Glucagon
 - C Adrenaline
 - D Thyroxine
4. In a cholinergic synapse, the enzyme that hydrolyses acetylcholine in the synaptic cleft is:
 - A Acetylcholine synthetase
 - B Acetylcholinesterase
 - C Choline acetyltransferase
 - D Monoamine oxidase

EXERCISE 13 True / False with Justification

State whether each statement is true or false. Justify your answer with a brief biological explanation.

1. The fovea contains only rods and is responsible for vision in dim light.
2. The refractory period of a neurone ensures that action potentials travel in one direction only.
3. Auxins promote root elongation at high concentrations.
4. The semicircular canals detect linear acceleration and gravity.

EXERCISE 14 Definitions

Define the following terms precisely, as required in the GCE Advanced Level examination.

1. **Stimulus**
2. **Receptor**
3. **Action potential**
4. **Synaptic cleft**
5. **Phototropism**
6. **Second messenger** (in hormone action)

EXERCISE 15 Direct Calculations

1. The resting membrane potential of a neurone is -70 mV. During an action potential, the membrane potential rises to $+30$ mV. Calculate the magnitude of the change in membrane potential during depolarisation.
2. A student measures the conduction velocity of an action potential in a myelinated axon of length 1.2 m. The time taken for the impulse to travel this distance is 6 ms. Calculate the conduction velocity in m s^{-1} .
3. In an experiment on phototropism, a coleoptile bends 15° towards a unilateral light source after 2 hours. If the rate of bending is constant, what angle will it have bent after 5 hours?

15.2 I practise

EXERCISE 16 Eye Structure, Image Formation and Light Adaptation

The diagram below shows a longitudinal section of the human eye. (In the examination, a diagram would be provided; here you should draw and label your own simplified diagram.)

1. Draw a large, labelled diagram of the human eye showing the following structures: cornea, aqueous humour, pupil, lens, vitreous humour, retina, fovea, optic nerve, iris, ciliary muscle, suspensory ligaments, sclera, choroid.
2. Explain how the eye focuses on a near object when a person moves from a dimly lit room into bright sunlight. Your answer should include:
 - Changes in the lens shape and the role of the ciliary muscles and suspensory ligaments (accommodation).
 - Changes in pupil diameter and the role of the iris muscles (pupillary reflex).
 - The sequence of events from light hitting the retina to the brain interpreting the image.

EXERCISE 17 Analysis of an Action Potential

The graph below shows the change in membrane potential of a motor neurone during an action potential. (Sketch a typical action potential graph: resting potential at -70 mV, rapid rise to $+30$ mV, fall below resting potential, return to resting potential.)

1. On the graph, label and define: resting potential, depolarisation, repolarisation, hyperpolarisation (after-potential), refractory period.
2. Explain the ion movements responsible for each phase (depolarisation, repolarisation, hyperpolarisation). Mention the specific voltage-gated channels involved.
3. Why does the refractory period prevent the action potential from travelling backwards along the axon?

EXERCISE 18 Comparison of Rods and Cones

1. Complete the table below to compare rods and cones in the human retina.

Feature	Rods	Cones
Visual pigment		
Distribution in retina		
Sensitivity to light intensity		
Visual acuity (detail)		
Colour vision		
Convergence (neuronal pathways)		

2. Explain why the fovea provides the sharpest vision, referring to the distribution of photoreceptors and their connections to bipolar and ganglion cells.
3. Why is it difficult to see a faint star at night when looking directly at it, but easier when looking slightly to the side?

EXERCISE 19 Cholinergic Synapse and Drug Action

1. Describe the sequence of events at a cholinergic synapse from the arrival of an action potential at the presynaptic knob to the generation of a postsynaptic potential. Include the roles of voltage-gated Ca^{2+} channels, vesicles, acetylcholine (ACh), receptors, and acetylcholinesterase.
2. Explain why synaptic transmission is unidirectional.
3. A drug inhibits acetylcholinesterase. Predict and explain the effect of this drug on:
 - The concentration of ACh in the synaptic cleft.
 - The frequency of action potentials in the postsynaptic neurone.
 - The duration of the postsynaptic response.
4. Give one medical use and one toxic effect of an acetylcholinesterase inhibitor.

15.3 I prepare for the GCE

EXERCISE 20 Essay: Nervous vs Hormonal Coordination] [15 marks

Compare and contrast nervous coordination and hormonal (endocrine) coordination in mammals. Your answer should include:

- Nature of the signal (electrical/chemical) and speed of transmission.

- Duration of effect and reversibility.
- Pathway of the message (neurones vs bloodstream).
- Specificity of target cells.
- Two named examples of each type of coordination in humans, describing the stimulus, receptor, coordinator, effector and response.

EXERCISE 21 Data Analysis: Blood Glucose Regulation] [12 marks

The graph shows blood glucose concentration over time for two individuals, A (normal) and B (diabetic), after each consumed a glucose-rich meal at time $t = 0$. (Sketch typical curves: A rises to a peak at 30 min, returns to normal by 120 min; B rises higher, stays elevated longer.)

1. Describe the changes in blood glucose concentration for individual A between 0 and 120 minutes. [3]
2. Explain the hormonal mechanisms responsible for the return of blood glucose to normal in individual A. Name the hormones, their source, and their target tissues/actions. [5]
3. Suggest why individual B fails to regulate blood glucose effectively. Distinguish between Type 1 and Type 2 diabetes in your explanation. [4]

EXERCISE 22 Experimental Design: Phototropism in Coleoptiles] [13 marks

Design a controlled experiment to demonstrate that unilateral light causes phototropic curvature in oat coleoptiles (or maize seedlings) and that the tip is the site of perception/production of the hormone involved.

Your answer must include:

- A clear hypothesis.
- Description of at least four experimental groups (including controls) with diagrams/labels.
- The independent variable, dependent variable, and at least three controlled variables.
- The expected results for each group.
- A conclusion linking the results to the role of auxin (IAA) distribution.

16 SOLUTIONS TO THE EXERCISES**SOLUTION Exercise 1 — Multiple Choice Questions**

1. Answer: B — Ganglion cells in the retina.

A sub-population of retinal ganglion cells contains the photopigment **melanopsin** and is directly sensitive to light intensity. These cells send signals via the optic nerve to the pretectal nucleus of the midbrain, which initiates the **pupillary reflex** (constriction of the pupil in bright light). Rods and cones (A) detect light for image formation, not primarily for the pupillary reflex; bipolar cells (C) are intermediate relay neurones; the pigmented epithelium (D) absorbs stray light and regenerates visual pigments.

2. Answer: C — Na^+ ions move into the axon.

Depolarisation is caused by the opening of voltage-gated Na^+ channels. Because the electrochemical gradient favours Na^+ entry (higher external concentration and negative interior), Na^+ rushes into the axon, making the inside positive (up to about +30 mV). K^+ moving out (not in) causes repolarisation.

3. Answer: B — Glucagon.

Glucagon, secreted by the α -cells of the islets of Langerhans in the pancreas when blood glucose falls, binds to liver (hepatocyte) receptors and stimulates **glycogenolysis** (glycogen $\rightarrow$ glucose) and gluconeogenesis. Insulin (A) does the opposite (glucose $\rightarrow$ glycogen). Adrenaline (C) can also promote glycogen breakdown, but it is primarily a stress hormone; the specific blood-glucose-regulating hormone asked for here is glucagon. Thyroxine (D) controls metabolic rate.

4. Answer: B — Acetylcholinesterase.

Acetylcholinesterase (AChE), located in the synaptic cleft, hydrolyses acetylcholine into choline and ethanoic (acetic) acid, terminating the signal. Choline is reabsorbed into the presynaptic knob for resynthesis of ACh by choline acetyltransferase (C). Monoamine oxidase (D) acts on monoamine transmitters such as noradrenaline, not ACh.

SOLUTION Exercise 2 — True / False with Justification

1. False. The fovea (fovea centralis) contains only cones, not rods — rods are completely absent from it. Cones give high-acuity colour vision in bright light. Vision in dim light is the function of rods, which are concentrated in the peripheral retina and contain the pigment rhodopsin, which is very sensitive to low light intensities.

2. True. During the **absolute refractory period**, the voltage-gated Na^+ channels are inactivated and cannot reopen, so no new action potential can be generated in the region just behind the travelling impulse. The impulse can therefore only propagate forwards into the region where channels are still in their resting (closed but activatable) state. This ensures unidirectional transmission along the axon and keeps successive impulses separate.
3. False. Auxins (e.g. IAA) inhibit root elongation at high concentrations; roots are far more sensitive to auxin than shoots. Root elongation is promoted only by very low auxin concentrations. In shoots, by contrast, higher auxin concentrations promote cell elongation — this differential sensitivity explains the opposite curvatures in gravitropism of roots and shoots.
4. False. The semicircular canals detect rotational (angular) acceleration of the head: movement of endolymph deflects the cupula in each ampulla. Linear acceleration and gravity are detected by the **utricle and saccule** (the otolith organs), where otoliths embedded in a gelatinous membrane press on hair cells.

SOLUTION Exercise 3 — Definitions

1. Stimulus: A change in the internal or external environment of an organism which is detected by a receptor and which brings about a response.
2. Receptor: A specialised cell, or structure, that detects a specific stimulus and converts (transduces) the energy of that stimulus into an electrical signal — a generator potential / nerve impulse.
3. Action potential: A rapid, transient, self-propagating reversal of the potential difference across the membrane of a neurone (from about -70 mV to about $+30\text{ mV}$ and back), caused by the sequential opening of voltage-gated Na^+ and K^+ channels; it is the electrical signal by which information travels along a neurone.
4. Synaptic cleft: The narrow gap (about 20 nm) separating the presynaptic membrane of one neurone from the postsynaptic membrane of the next cell (neurone or effector), across which the signal is carried by diffusion of a chemical neurotransmitter.
5. Phototropism: A directional growth movement (curvature) of a plant organ in response to the direction of unilateral light; shoots are positively phototropic (grow towards light) because auxin accumulates on the shaded side and stimulates greater elongation there.
6. Second messenger: An intracellular molecule (e.g. cyclic AMP, cAMP) produced inside a target cell when a water-soluble hormone (the first messenger, e.g. adrenaline or glucagon) binds to a receptor on the cell surface membrane; the second messenger

ger relays and amplifies the signal by activating enzymes inside the cell, producing the cellular response.

SOLUTION Exercise 4 — Direct Calculations

1. The change in membrane potential is:

$$\Delta V = V_{\text{peak}} - V_{\text{rest}} = (+30) - (-70) = +100 \text{ mV}.$$

The magnitude of the change is 100 mV (an increase of 100 mV during depolarisation).

2. Conduction velocity $v = \frac{d}{t}$ with $d = 1.2 \text{ m}$ and $t = 6 \text{ ms} = 6 \times 10^{-3} \text{ s}$:

$$v = \frac{1.2}{6 \times 10^{-3}} = 200 \text{ m s}^{-1}.$$

Conduction velocity = 200 m s^{-1} . (This high value is typical of saltatory conduction in a myelinated axon.)

3. Rate of bending = $\frac{15^\circ}{2 \text{ h}} = 7.5^\circ \text{ h}^{-1}$. After 5 hours:


$$\theta = 7.5 \times 5 = 37.5^\circ.$$

The coleoptile will have bent 37.5° after 5 hours (assuming the rate remains constant).

SOLUTION Exercise 5 — Eye Structure, Image Formation and Light Adaptation

1. Labelled diagram of the human eye.

The diagram should show the three layers (sclera, choroid, retina), the optical pathway (cornea → aqueous humour → pupil → lens → vitreous humour → retina) and the accessory structures.



Longitudinal section through the human eye (schematic).

2. Response to moving into bright sunlight and focusing on a near object.

(a) Accommodation (focusing on the near object):

- The **ciliary muscles contract**, releasing tension on the suspensory ligaments.
- The **suspensory ligaments slacken**, so the elastic **lens** becomes more convex (shorter, fatter), increasing its refractive power.
- Light rays from the near object (which are diverging strongly) are now refracted enough to be focused sharply on the retina.

(b) Pupillary reflex (response to bright light):


- High light intensity is detected by retinal ganglion cells; impulses pass via the optic nerve to the midbrain and back along parasympathetic motor neurones.
- The **circular muscles of the iris contract** and the radial muscles relax.
- The **pupil constricts**, reducing the amount of light entering the eye and protecting the retina (especially the rods, whose rhodopsin is bleached).

(c) From retina to perception:

1. Light passes through the cornea, aqueous humour, pupil, lens and vitreous humour and is focused on the retina, forming a real, inverted, diminished image.
2. Photoreceptors (rods and cones) absorb light; visual pigments (rhodopsin/iodopsin) are bleached, generating a generator potential.
3. Impulses pass from photoreceptors to bipolar neurones, then to ganglion cells.
4. Ganglion cell axons form the optic nerve, which carries impulses to the visual cortex of the cerebrum (occipital lobe).
5. The brain interprets the impulses, re-inverting the image so that the object is perceived upright and in detail.

SOLUTION Exercise 6 — Analysis of an Action Potential

1. Labelled graph and definitions.



Membrane potential during an action potential.

- Resting potential: the stable potential difference across the membrane of an unstimulated neurone (about -70 mV, inside negative), maintained by the Na^+/K^+ pump and the greater permeability to K^+ .
- Depolarisation: the rapid reversal of the membrane potential from -70 mV to about $+30$ mV.
- Repolarisation: the rapid return of the membrane potential towards the resting value.
- Hyperpolarisation (after-potential): a brief period when the potential falls below -70 mV (about -90 mV) before returning to rest.
- Refractory period: the interval during and just after an action potential when a new action potential cannot (absolute) or can only with difficulty (relative) be generated.

2. Ionic basis of each phase.

- Depolarisation: the stimulus raises the potential above the threshold (about -55 mV); **voltage-gated Na^+ channels** open, Na^+ floods into the axon down its electrochemical gradient, making the inside positive ($+30$ mV) — positive feedback opens more Na^+ channels.
- Repolarisation: at the peak, Na^+ channels inactivate and **voltage-gated K^+ channels** open; K^+ diffuses out of the axon, making the inside negative again.
- Hyperpolarisation: the K^+ channels are slow to close, so excessive K^+ leaves the cell and the potential briefly falls below -70 mV. The channels then close and the Na^+/K^+ pump (3 Na^+ out : 2 K^+ in, using ATP) restores the resting distribution of ions.

3. Why the impulse cannot travel backwards.

Immediately behind the propagating action potential, the voltage-gated Na^+ channels are inactivated (absolute refractory period) and cannot reopen whatever the stimulus. The local currents from the active region can therefore only depolarise the membrane ahead of the impulse, where the channels are closed but activatable. Hence propagation is strictly unidirectional, and successive impulses remain discrete (they cannot merge).

SOLUTION Exercise 7 — Comparison of Rods and Cones

1. Completed table.

Feature	Rods	Cones
Visual pigment	Rhodopsin (one type only)	Iodopsin (three types, sensitive to red, green or blue light)
Distribution in retina	Peripheral retina; absent from the fovea	Concentrated in the fovea; few elsewhere
Sensitivity to light intensity	High — function in dim light (scotopic vision)	Low — require bright light (photopic vision)
Visual acuity (detail)	Low acuity, poor detail	High acuity, sharp detail
Colour vision	None — black/white/grey only	Responsible for colour vision (trichromatic)
Convergence (neuronal pathways)	High: many rods converge onto one bipolar/ganglion cell (summation)	Low: at the fovea each cone connects to its own bipolar and ganglion cell

2. Why the fovea gives the sharpest vision.

- The fovea contains only cones, packed very densely, and cones give high acuity and colour vision.
- Each foveal cone synapses with its own bipolar cell, which connects to its own ganglion cell — a 1:1:1 pathway with no convergence. The brain can therefore distinguish signals from two adjacent cones separately, resolving fine detail.
- Overlying neurones and blood vessels are displaced to the sides of the fovea, so light strikes the cones directly without scattering.

3. Why a faint star is seen better when looking slightly to the side.

Looking directly at the star focuses its image on the fovea, which contains only cones — and cones are insensitive to dim light, so the star is not detected. Looking slightly to the

side moves the image onto the peripheral retina, which is rich in rods. Rods are highly sensitive to low light intensities (and their convergence allows **summation** of weak signals), so the faint star becomes visible — although with poor detail and no colour.

SOLUTION Exercise 8 — Cholinergic Synapse and Drug Action

1. Sequence of events at a cholinergic synapse.

1. An action potential arrives at the presynaptic knob and depolarises the presynaptic membrane.
2. **Voltage-gated Ca^{2+} channels** open; Ca^{2+} ions diffuse into the knob down their electrochemical gradient.
3. The rise in Ca^{2+} causes synaptic **vesicles** containing acetylcholine (ACh) to move to and fuse with the presynaptic membrane, releasing ACh into the synaptic cleft by exocytosis.
4. ACh diffuses across the cleft (≈ 20 nm) and binds to specific **ACh receptors** (ligand-gated Na^+ channels) on the postsynaptic membrane.
5. The channels open; Na^+ enters the postsynaptic neurone, causing depolarisation — an excitatory postsynaptic potential (EPSP). If the EPSP reaches threshold, an action potential is generated in the postsynaptic neurone.
6. **Acetylcholinesterase** in the cleft rapidly hydrolyses ACh into choline and ethanoic acid, terminating the signal; choline is reabsorbed into the presynaptic knob and used to resynthesise ACh.

2. Why transmission is unidirectional.

Vesicles of neurotransmitter are present only in the presynaptic knob, and the specific receptors are located only on the postsynaptic membrane. Transmitter can therefore be released and detected in one direction only, so the impulse cannot travel backwards across the synapse.

3. Effect of an acetylcholinesterase inhibitor.

- Concentration of ACh in the cleft: it increases, because ACh is released normally but is no longer broken down, so it accumulates.
- Frequency of postsynaptic action potentials: it increases. Persisting ACh repeatedly opens the receptor channels, so the postsynaptic membrane is re-depolarised again and again, firing repeated action potentials.
- Duration of the postsynaptic response: it is prolonged, because each release of ACh continues to stimulate the receptors until the ACh finally diffuses away or is degraded by other routes.

4. Medical use and toxic effect.

- Medical use: treatment of myasthenia gravis (an autoimmune disease in which ACh receptors are destroyed); AChE inhibitors such as neostigmine raise ACh levels in the cleft so that the remaining receptors are adequately stimulated, restoring muscle contraction.
- Toxic effect: organophosphate insecticides and nerve gases irreversibly inhibit AChE; ACh accumulates, causing continuous stimulation of muscles — twitching, paralysis of respiratory muscles, excessive glandular secretion — which can be fatal.

SOLUTION Exercise 9 — Essay: Nervous vs Hormonal Coordination (15 marks)

Indicative mark scheme and model content.

Comparison table (up to 7 marks — one mark per correct contrasted pair).

Feature	Nervous coordination	Hormonal coordination
Nature of signal	Electrical impulses (plus chemical transmitter at synapses)	Chemical only (hormones)
Speed	Very fast (up to $\sim 120 \text{ m s}^{-1}$)	Slow (speed of blood circulation)
Duration of effect	Short-lived, brief	Longer-lasting, sustained
Reversibility	Rapidly reversed when stimulus stops	Effects persist while hormone circulates; slower to reverse
Pathway	Along specific neurones to a precise effector	Through the bloodstream to the whole body
Specificity of target	Highly localised (one muscle/gland)	Widespread; only cells with the correct receptors respond

Similarities (2 marks): both involve chemical messengers (neurotransmitters / hormones); both link a stimulus to a response via receptors and effectors; both maintain homeostasis.

Two named nervous examples (3 marks):

- Withdrawal reflex: stimulus — a pin prick on the finger; receptor — pain receptors in the skin; coordinator — spinal cord (relay neurone); effector — biceps muscle of the arm; response — rapid withdrawal of the hand.
- Pupillary reflex: stimulus — bright light; receptor — retinal ganglion cells; coordinator — midbrain; effector — circular muscles of the iris; response — pupil constricts.

Two named hormonal examples (3 marks):

- Blood glucose regulation: stimulus — rise in blood glucose after a meal; receptor — β -cells of the islets of Langerhans; coordinator — pancreas; effector — liver and mus-

cle cells; response — insulin secreted, glucose converted to glycogen, blood glucose falls.

- “Fight or flight”: stimulus — sudden danger; receptor/coordinator — hypothalamus stimulates the adrenal medulla; effector — heart, liver, bronchioles; response — adrenaline released, heart rate and blood glucose rise, preparing the body for action.

Examiner expectations: a clear table or paired statements (not two separate unlinked accounts); accurate terminology (neurone, effector, target cells, receptors); correct stimulus–receptor–coordinator–effector–response chains for each example; at least one similarity stated. Award full marks only if all four bullet points of the question are addressed.

SOLUTION Exercise 10 — Data Analysis: Blood Glucose Regulation (12 marks)

1. Description for individual A [3 marks].

- From $t = 0$ blood glucose rises steeply as glucose is absorbed from the gut, reaching a peak at about 30 minutes. [1]
- It then falls progressively as glucose is removed from the blood. [1]
- By about 120 minutes it has returned to (and is stabilised at) the normal resting level (about 90 mg per 100 cm³). [1]

2. Hormonal mechanism of the return to normal [5 marks].

- The rise in blood glucose is detected by the **β -cells of the islets of Langerhans** in the pancreas. [1]
- They secrete **insulin** into the blood. [1]
- Insulin binds to receptors on liver cells (hepatocytes) and muscle cells, stimulating the conversion of glucose to **glycogen** (glycogenesis) for storage. [1]
- It also increases the uptake of glucose by cells and its oxidation in respiration, and inhibits glycogen breakdown. [1]
- As blood glucose falls, insulin secretion falls (negative feedback); if glucose drops below normal, **glucagon** from the α -cells stimulates the liver to convert glycogen back to glucose, restoring the level. [1]

3. Why individual B fails to regulate [4 marks].

- B's glucose rises higher and remains elevated because the insulin mechanism is defective. [1]
- Type 1 diabetes: the β -cells are destroyed (autoimmune reaction), so no insulin is produced; blood glucose cannot be lowered despite the stimulus. Managed by insulin injections. [1.5]


- Type 2 diabetes: insulin is produced, but the target cells' receptors are insensitive/resistant to insulin (often associated with obesity and age), so glucose uptake and glycogenesis are inadequate. Managed by diet, exercise and drugs. [1.5]

Examiner expectations: precise description quoting times and trend; both hormones named with sources and targets; clear distinction between insulin deficiency (Type 1) and insulin resistance (Type 2).

SOLUTION Exercise 11 — Experimental Design: Phototropism in Coleoptiles (13 marks)

Hypothesis [1]: Unilateral light causes a coleoptile to bend towards the light because the tip perceives the light and produces a growth hormone (auxin/IAA) which is redistributed to the shaded side, where it stimulates greater cell elongation.

Experimental groups [4 — one per group with correct treatment]. Use uniform oat/maize coleoptiles of the same age and length, all exposed to unilateral light (except Group E):



Five groups of coleoptiles under unilateral light (E in darkness).

- Group A (control): intact coleoptile, unilateral light.
- Group B: tip removed, unilateral light.
- Group C: tip covered with an opaque cap, unilateral light.
- Group D: tip exposed but the base covered with an opaque sleeve, unilateral light.
- Group E (control): intact coleoptile kept in darkness (or uniform light).

Variables [3]:

- Independent variable: the treatment of the tip (present / removed / covered) and the direction of light. [1]
- Dependent variable: the angle/direction of curvature (measured with a protractor) and the growth in length. [1]
- Controlled variables (any three): same species and age of seedlings; same initial coleoptile length; same light intensity, duration and wavelength; same temperature; same humidity/water supply; same number of seedlings per group (replicates). [1]

Expected results [3]:


- A: grows and bends towards the light (positive phototropism).
- B: little or no growth and no bending — the tip is needed both for perception and as the source of auxin.
- C: grows straight up, no bending — the tip is present but cannot perceive the light direction, so auxin is distributed evenly.
- D: bends towards the light like A — covering the base does not matter, since perception occurs at the tip and the response (elongation) still occurs below.
- E: grows straight up (etiolated, tall and pale) with no curvature.

Conclusion [2]: Only coleoptiles with an illuminated tip bend towards unilateral light. The tip is therefore the site of light perception and of auxin (IAA) production. Unilateral light causes auxin to be transported to the shaded side, where the higher auxin concentration stimulates greater cell elongation; this differential growth curves the coleoptile towards the light. [2]

Examiner expectations: a testable hypothesis; at least four clearly labelled groups including a positive and a negative control; correct identification of all three variable types; results consistent with the auxin-redistribution theory; a conclusion explicitly linking tip, light and unequal auxin distribution to differential elongation.

17 SUMMARY SHEET

17.1 Summary Sheet



Mind map of Chapter BIO02: Response & Coordination in Organisms

FICHE BILAN L'essentiel à connaître pour l'examen

Key Definitions

- **Stimulus:** A detectable change in the internal or external environment.
- **Response:** The activity of an organism or part of an organism resulting from a stimulus.
- **Receptor:** A specialised cell or organ that converts a stimulus into a nerve impulse (transduction).
- **Effector:** A muscle or gland that brings about a response.
- **Neurone:** The structural and functional unit of the nervous system; transmits electrical impulses.
- **Synapse:** The junction between two neurones or a neurone and an effector; transmits-

sion is chemical (neurotransmitters).

- **Hormone:** A chemical messenger secreted by an endocrine gland into the blood, acting on specific target cells.
- **Tropism:** A directional growth response of a plant to a unilateral stimulus (e.g. phototropism, geotropism).
- **Nastic response:** A non-directional response to a stimulus (e.g. Mimosa leaf folding).

Core Principles & Formulas

- **Reflex Arc:** Receptor → Sensory neurone → Interneurone (spinal cord) → Motor neurone → Effector. Fast, involuntary, protective.
- **All-or-None Law:** An action potential is generated only if the stimulus reaches threshold; amplitude is constant.
- **Refractory Periods:** Absolute (no new AP possible) and Relative (stronger stimulus needed); ensures unidirectional propagation.
- **Accommodation (Eye):** Ciliary muscles contract → suspensory ligaments slacken → lens becomes more convex → increased refractive power for near objects.
- **Pupillary Light Reflex:** Circular muscles (parasympathetic) constrict pupil in bright light; radial muscles (sympathetic) dilate in dim light.
- **Phototransduction:** Light → Rhodopsin bleaching → Transducin activation → PDE activation → cGMP hydrolysis → Na⁺ channels close → Hyperpolarisation.
- **Hormone Mechanisms:**
 - **Peptide hormones:** Bind surface receptor → Second messenger (cAMP/IP₃) → Protein kinase cascade → Rapid response.
 - **Steroid hormones:** Diffuse through membrane → Bind intracellular receptor → Hormone-receptor complex enters nucleus → Gene transcription → Slow, long-lasting.
- **Plant Hormones:** Auxins (IAA) promote cell elongation, apical dominance, root initiation; Gibberellins break dormancy, stem elongation; Cytokinins promote cell division, delay senescence; Abscissic acid (ABA) closes stomata, induces dormancy; Ethene promotes fruit ripening, leaf abscission.

Key Methods (Exam Skills)

1. **Diagram Annotation:** Draw and label the eye (retina, fovea, blind spot, iris, lens, ciliary body), ear (cochlea, semi-circular canals, ossicles), neurone (myelin sheath, nodes of Ranvier, synaptic knob), reflex arc.
2. **Action Potential Graph:** Label axes (mV vs ms), phases: Resting potential (−70 mV), Depolarisation (Na⁺ influx), Overshoot (+30 mV), Repolarisation (K⁺ efflux), Hyperpolarisation, Return to rest.

3. Synaptic Transmission Steps: 1. AP arrives at knob → 2. Voltage-gated Ca^{2+} channels open → 3. Vesicles fuse → 4. Acetylcholine released → 5. Binds Na^+ channel receptors → 6. Depolarisation (EPSP) → 7. Acetylcholinesterase hydrolyses ACh.
4. Comparing Nervous vs Endocrine: Speed (ms vs mins/hrs), Duration (short vs long), Transmission (neurones vs blood), Specificity (precise vs widespread), Adaptation (rapid vs slow).
5. Plant Tropism Experiments: Know the classic Darwin/Coleoptile tip experiments: Tip perceives light → Auxin migrates to shaded side → Asymmetric elongation → Bending towards light.

Common Mistakes (Traps)

- Confusing **rods** (dim light, low acuity, no colour, rhodopsin) and **cones** (bright light, high acuity, colour vision, iodopsin).
- Stating the lens changes shape by muscle contraction directly; it is the release of tension on suspensory ligaments.
- Thinking the **cochlea** detects balance; it detects sound. Balance is via **semi-circular canals** (dynamic) and **utricle/saccule** (static).
- Writing “neurones secrete hormones” at synapses; they secrete **neurotransmitters**.
- Forgetting the **myelin sheath** is formed by Schwann cells (PNS) / Oligodendrocytes (CNS) and enables **saltatory conduction**.
- Mixing up **sympathetic** (fight/flight: dilates pupil, inhibits digestion, increases heart rate) and **parasympathetic** (rest/digest: constricts pupil, stimulates digestion, decreases heart rate).
- Claiming auxin inhibits root growth at high concentration but promotes shoot growth; the concentration optimum differs.
- Omitting **acetylcholinesterase** in synaptic transmission, leading to continuous stimulation.

GCE Advanced Level Tips

- Diagrams: Large (> 10 cm), clear lines, labels horizontal, label lines touching structure (no arrows), title + magnification if relevant.
- Definitions: Quote precisely: “A hormone is a chemical substance produced by an endocrine gland and transported by the blood to a target organ where it exerts its effect.”
- Comparison Tables: Use a ruler, clear columns (Feature / Nervous / Endocrine). Fill every cell.
- Calculations: Magnification = $\frac{\text{Image size}}{\text{Actual size}}$ (same units). Scale bars: $\frac{\text{Measured length}}{\text{Scale factor}} = \text{Actual size}$.

- Experimental Design: State **independent**, **dependent**, **controlled** variables. Mention **replicates** and **statistical test** (e.g. t-test, chi-squared) if asked.
- Contextual Knowledge: Link to Cameroon: Sickle cell trait (malaria resistance), Onchocerciasis (river blindness – eye), Goitre (iodine deficiency – thyroid), Agricultural use of auxins (rooting powders, weedkillers like 2,4-D).



Signed • OnBuch+

ONBUCH+ TEACHING TEAM
FOLLOWS THE OFFICIAL MINESEC SYLLABUS

OnBuch+