

GEOLOGY • UPPER SIXTH

UPPER SIXTH — GEOLOGY

CHAPTER 5

ECONOMIC GEOLOGY: ECONOMIC MINERALS

SUGGESTED TIME: 12 PERIODS

IN THIS CHAPTER

This chapter introduces the concept of economic minerals, the distinction between resources and reserves, and the classification, formation and occurrence of mineral deposits. Using Cameroonian examples such as the bauxite of Minim-Martap, the iron of Mbalam and the gold of the East Region, students learn how geologists evaluate and locate the mineral wealth of a country.

- By the end of this chapter I can define an economic mineral and distinguish it from a gangue mineral and an ordinary rock.
- By the end of this chapter I can explain the terms ore, grade, cut-off grade and tonnage with simple calculations.
- By the end of this chapter I can distinguish mineral resources from mineral reserves and explain the factors that convert one into the other.
- By the end of this chapter I can classify mineral deposits according to their morphology, their host rock and their mode of origin.
- By the end of this chapter I can describe the magmatic, hydrothermal, sedimentary and weathering processes that form ore deposits.
- By the end of this chapter I can relate the occurrence of major deposits to the geological context of Cameroon and the world.



DOWNLOAD ON- BUCH

Free • Google Play
AI tutor, past papers,
results



GO ONBUCH+

All signed courses, unlimited AI
tutor, PDF sheets — OnBuch+
tab in the app

THIS SIGNED COURSE CONTAINS

COURSE

complete, illustrated,
step by step

METHODS

and worked examples

EXERCISES

exam style + solutions

SUMMARY SHEET

the essentials to
remember



Signed • OnBuch+

ONBUCH+ TEACHING TEAM
FOLLOWS THE OFFICIAL MINESEC SYLLABUS

OnBuch+

CONTENTS

- | | |
|--|---|
| 1. The Notion of Economic Mineral | 6. Occurrence of Mineral Deposits: From the World to Cameroon |
| 2. Resources and Reserves | 7. Integration activity |
| 3. Classification of Mineral Deposits | 8. Methods and worked examples |
| 4. Formation of Ore Deposits I: Magmatic and Hydrothermal Processes | 9. Exercises |
| 5. Formation of Ore Deposits II: Sedimentary, Weathering and Metamorphic Processes | 10. Solutions to the exercises |
| | 11. Summary sheet |

CHAPTER OBJECTIVES

- By the end of this chapter I can define an economic mineral and distinguish it from a gangue mineral and an ordinary rock.
- By the end of this chapter I can explain the terms ore, grade, cut-off grade and tonnage with simple calculations.
- By the end of this chapter I can distinguish mineral resources from mineral reserves and explain the factors that convert one into the other.
- By the end of this chapter I can classify mineral deposits according to their morphology, their host rock and their mode of origin.
- By the end of this chapter I can describe the magmatic, hydrothermal, sedimentary and weathering processes that form ore deposits.
- By the end of this chapter I can relate the occurrence of major deposits to the geological context of Cameroon and the world.
- By the end of this chapter I can interpret simple maps, sections and grade-tonnage data to assess the economic potential of a deposit.
- By the end of this chapter I can discuss the economic and environmental stakes of mineral exploitation in Cameroon.

PREREQUISITES

- Minerals and rocks: definition of a mineral, common rock-forming minerals (Lower Sixth).
- The rock cycle and the three great families of rocks (igneous, sedimentary, metamorphic).
- Basic notions of plate tectonics and magmatic processes.
- Weathering and sedimentation processes (Lower Sixth, external geodynamics).
- Simple percentage and unit-conversion calculations from Mathematics.

1 THE NOTION OF ECONOMIC MINERAL

In 2023, a young entrepreneur from Batouri in the East Region of Cameroon sold a few kilograms of gold-bearing gravel for less than the price of a bag of cement, while a foreign company exporting the same gold earned millions of CFA francs. Why does the same mineral have such different values depending on who holds it and how much of it exists underground? How do geologists decide whether a hill of rock is worthless or a fortune? And why does Cameroon, rich in bauxite at Minim-Martap and iron at Mbalam, still import most of its metals? This chapter gives you the geological keys to answer these questions.

1.1 From Mineral to Ore: the Basic Vocabulary

In 2023, a young entrepreneur from Batouri in the East Region of Cameroon sold a few kilograms of gold-bearing gravel for less than the price of a bag of cement, while a foreign company exporting the same gold earned millions of CFA francs. Why does the same mineral have such different values depending on who holds it and how much of it exists underground? How do geologists decide whether a hill of rock is worthless or a fortune? And why does Cameroon, rich in bauxite at Minim-Martap and iron at Mbalam, still import most of its metals? This chapter gives you the geological keys to answer these questions.

1.2 From Mineral to Ore: the Basic Vocabulary

[illegible]

2 RESOURCES AND RESERVES

In the previous section we established that an **economic mineral** is a mineral substance whose concentration in the Earth's crust is high enough, and whose extraction is profitable enough, to be exploited. But between "there is iron in this hill" and "this iron mine is profitable", there is a long road of measurement, calculation and decision. Mining companies, banks and governments do not speak vaguely of "minerals in the ground": they use two precise, internationally standardised words — **resource** and **reserve**. Mastering the difference between them is one of the most frequently tested points of this chapter at the GCE Advanced Level.

2.1 The Two Fundamental Notions

Intuition first.

Imagine a farmer who knows that mangoes grow somewhere on his large farm. He has seen mangoes from a distance in some areas (low confidence), counted the trees in other areas (medium confidence), and weighed and priced the fruit on the trees nearest his house (high confidence). Only the fruit he can actually pick, transport and sell at a profit this season constitutes his real “harvestable stock”. The rest exists, certainly, but it is not yet money in the pocket. In the same

way, geologists distinguish what is probably there (a resource) from what is demonstrably mineable at a profit (a reserve).

DEFINITION Mineral resource

A **mineral resource** is a concentration or occurrence of solid mineral material in or on the Earth's crust, in such form, grade (quality) and quantity that there are reasonable prospects of eventual economic extraction. Its location, tonnage and characteristics are known, estimated or interpreted from geological evidence and sampling.

DEFINITION Mineral reserve (ore reserve)

A **mineral reserve** is the economically mineable part of a measured or indicated mineral resource. It is demonstrated by a feasibility study which takes into account mining, metallurgical, economic, marketing, legal, environmental and governmental factors. A reserve is what a company can legally and profitably extract today.

WARNING Resource ≠ reserve

In GCE answers, never use the two words as synonyms. A reserve is always a resource, but a resource is not always a reserve: it becomes one only when economic extraction is demonstrated. Writing "Cameroon has huge iron reserves at Mbalam" when the deposit is still at the resource stage is a factual error that examiners penalise.


2.2 Classification by Degree of Geological Confidence

As exploration progresses (outcrop mapping → geophysics → trenching → borehole drilling → bulk sampling), the geologist's confidence in the tonnage and grade increases. Three categories of resource are distinguished:

DEFINITION Inferred, indicated and measured resources

- **Inferred resource**: estimated from geological evidence and limited sampling; continuity of the orebody is assumed but not verified. Low confidence.
- **Indicated resource**: sampling (boreholes, trenches) is dense enough to assume continuity between observation points; tonnage and grade are estimated with reasonable confidence.
- **Measured resource**: sampling is so dense (close-spaced drilling, underground workings) that tonnage, grade, shape and mineral content are well established. Highest

confidence.



The resource–reserve pyramid: the largest tonnages are known with the least confidence (inferred), while the economically mineable, best-known part (proved and probable reserves) is the smallest.


2.3 Classification by Economic Viability

Geological confidence alone is not enough: the ore must be mineable at a profit. This gives the second axis of classification:

DEFINITION Proved and probable reserves

- **Proved (proven) reserve:** the economically mineable part of a measured resource; highest confidence in both geology and economics.
- **Probable reserve:** the economically mineable part of an indicated (and sometimes measured) resource; confidence is lower than for a proved reserve but extraction is still judged profitable.

Resources that are almost economic are called **marginal** (borderline profitable); those that cannot be mined profitably under present conditions are **sub-economic** resources.



The “McKelvey box”: resources and reserves classified simultaneously by geological certainty (vertical axis) and economic viability (horizontal axis). Only the upper-right zone constitutes true reserves.

KEY POINTS The McKelvey box

The **McKelvey diagram** combines the two classifications: as one moves upwards, geological certainty increases (inferred → indicated/measured); as one moves rightwards, economic viability increases (sub-economic → marginal → economic). Reserves occupy the upper-right corner. A deposit can migrate inside the box without any new drilling, simply because prices or technology change.

2.4 From Resource to Reserve: the Conversion Factors

What decides whether a resource crosses the boundary into the reserve zone? Five groups of factors:

1. **Metal and mineral prices.** If the world price of iron or gold rises, lower-grade ore that was previously worthless becomes profitable: the cut-off grade falls and the reserve grows.
2. **Mining and processing technology.** Cheaper open-pit machinery, better ore-dressing (e.g. magnetic separation of magnetite) or new hydrometallurgical methods lower costs and enlarge reserves.
3. **Infrastructure.** Railways, roads, deep-sea ports, water and electricity supply often decide everything for bulky, low-value commodities such as iron ore.
4. **Energy costs.** Smelting and crushing are energy-hungry; cheap hydroelectricity (a Cameroonian asset) improves viability.

5. **Legislation and environmental constraints.** Mining codes, taxes, land rights and environmental protection can either encourage or block conversion of a resource into a reserve.

WARNING A price rise can “create” a reserve

A rise in the world price of a metal can transform a sub-economic resource into a reserve without any new discovery and without moving a single rock. Reserves are therefore an economic stock, not a fixed physical quantity. Conversely, a price collapse can wipe out reserves overnight.

2.5 The Dynamic Nature of Reserves and Reserve Life

PROPERTY Reserves are a dynamic stock

Reserves shrink through extraction (depletion by mining) and grow through new discoveries, rises in commodity prices and technological progress. The “end of a mineral” is therefore rarely a physical exhaustion; it is usually an economic limit that moves with time. (Statement examinable; no mathematical proof required.)

A practical consequence is the notion of **reserve life**: the number of years a deposit (or a country, or the world) can continue producing at the current rate.

METHOD Calculating reserve life

1. Identify the reserve tonnage R (in tonnes, t) from the data.
2. Identify the annual production P (in t/yr), making sure the units match.
3. Compute: Reserve life = $\frac{R}{P}$ (in years).
4. Comment critically: the result assumes constant production, constant prices and no new discovery — a simplification.

WORKED EXAMPLE 1 Reserve life of an iron deposit

Statement: An iron deposit in the South Region of Cameroon has a proved reserve of 2.0×10^8 t of saleable ore. The planned mine will produce 1.0×10^7 t per year. (a) Calculate the reserve life. (b) Explain two reasons why the true mine life may differ from this figure.

SOLUTION

Solution. (a) Applying the method:

$$\text{Reserve life} = \frac{R}{P} = \frac{2.0 \times 10^8 \text{ t}}{1.0 \times 10^7 \text{ t yr}^{-1}} = 20 \text{ years.}$$

(b) The true life may be longer because further drilling may convert inferred resources into reserves, or a rise in iron prices may make lower-grade ore economic; it may be shorter if production is increased above the planned rate or if part of the reserve proves impossible to recover (sterile zones, environmental restrictions). Reserve life is therefore an estimate valid only under stated conditions.

2.6 Case Study: the Mbalam–Nabeba Iron Project

EXPERIMENT**Guided case study — Mbalam–Nabeba (East Region, Cameroon)**

The Mbalam–Nabeba iron deposit, straddling the Cameroon–Congo border, contains large tonnages of high-grade hematite and itabirite mineralisation hosted in Precambrian rocks of the Congo craton margin. Geological exploration established a resource of several hundred million tonnes of iron ore. Yet for years this resource could not be declared a full reserve, for one decisive reason: infrastructure. Iron ore is a bulky, low-value-per-tonne commodity; it cannot be exported profitably by lorry over hundreds of kilometres of forest roads. The project requires:

- a heavy-haul railway of roughly 500 km from the mine to the coast;
- a deep-sea port (mineral terminal near Kribi) capable of loading large bulk carriers;
- power and water supplies for the processing plant.

Only when financing and agreements for the railway and port are secured does the feasibility study demonstrate economic extraction, allowing the resource to be reclassified as a reserve. Conclusion: in Cameroon, as elsewhere, the passage from resource to reserve depends as much on engineers, bankers and ministers as on geologists.

DID YOU KNOW?**Did you know?**

The Mbalam–Nabeba project illustrates a general African paradox: the continent is extraordinarily rich in mineral resources (iron, bauxite, cobalt, gold) but converts them into reserves — and into revenue — slowly, because railways, ports and energy are lacking. Cameroon's bauxite at Minim–Martap and Ngaoundal, among the largest untapped resources in the world, faces exactly the same constraint.

EXERCISE 1 Resources, reserves and the McKelvey box

1. Define mineral resource and mineral reserve, and state the essential difference between them.
2. A gold prospect is known from only three boreholes 400 m apart. To which confidence category does its resource belong? Justify.
3. The world price of cobalt doubles. Using the McKelvey box, explain what happens to a marginal cobalt resource.
4. A mine holds a probable reserve of 5.0×10^7 t and produces 2.5×10^6 t yr⁻¹. Calculate its reserve life and state one assumption behind your answer.

SOLUTION Exercise 1

1. A mineral resource is a concentration of mineral material with reasonable prospects of eventual economic extraction; a mineral reserve is the economically mineable part of a measured or indicated resource, demonstrated by a feasibility study. The difference: a reserve is proven profitable now, a resource is only potentially economic.
2. Inferred resource: sampling is too sparse to confirm continuity of the orebody between boreholes; continuity is assumed, not verified.
3. The resource moves rightwards in the McKelvey box: higher price lowers the effective cut-off grade and raises revenue per tonne, so a marginal resource can become economic and be reclassified as a probable/proved reserve — with no new discovery.
4. Reserve life = $\frac{5.0 \times 10^7}{2.5 \times 10^6} = 20$ years. Assumption: production rate, prices and the reserve itself remain constant (no new discoveries, no reclassification).

KEY POINTS Key points of the section

- Resource = potentially economic concentration; reserve = demonstrated economically mineable part of a measured/indicated resource.
- Confidence increases: inferred → indicated → measured; viability increases: sub-economic → marginal → proved/probable reserve.
- Prices, technology, infrastructure, energy and legislation convert resources into reserves — or the reverse.
- Reserves are dynamic: depleted by mining, renewed by discovery, price rises and

technology.

- Reserve life = reserve tonnage ÷ annual production; the Mbalam–Nabeba case shows how railway and port condition the resource–to–reserve transition in Cameroon.

3

CLASSIFICATION OF MINERAL DEPOSITS

In the previous section we learned how to quantify what we find: **resources** tell us what exists in the ground, while **reserves** tell us what we can mine profitably today. But before we can even estimate tonnage and grade, the exploration geologist must answer a more fundamental question: what kind of deposit are we looking at? The answer dictates the exploration strategy, the mining method, the processing route and ultimately the economic viability. This section builds the classification toolkit that every geologist — and every GCE candidate — must master. It corresponds to **Lessons 91 and 92** of the official syllabus.

3.1 Why Classify Mineral Deposits?

Imagine you are handed a geological map showing a mineralised zone. Without a classification framework you would be facing a chaotic list of observations: “quartz veins in granite”, “banded iron layers in schist”, “gold grains in river sand”. Classification imposes order. It groups deposits that share **geometry**, **genetic origin**, **relationship to the host rock** or **commodity type**. Each criterion answers a different practical question:

- **Morphology (shape)** → determines the mining method (open pit vs underground, selective vs bulk mining).
- **Host–rock relation (syngenetic vs epigenetic)** → guides exploration targeting (look along stratigraphic horizons vs cross-cutting structures).
- **Genetic process** → predicts associated minerals, alteration halos, geochemical signatures and down-dip continuity.
- **Commodity group** → links to market dynamics, processing technology, strategic importance.

KEY POINTS

The Golden Rule of Classification

A single deposit can (and must) be described using all four criteria simultaneously. For example, the Mbalam iron deposit (Cameroon) is: morphologically stratiform; genetically sedimentary (BIF); host–relation syngenetic; commodity ferrous metallic. Never reduce a deposit to just one label.

3.2 Classification by Morphology (Shape and Geometry)

Morphology is the most immediately observable characteristic in the field or on a cross-section. It controls the **ore-body continuity** and the **stripping ratio**.

DEFINITION Stratiform Deposit

A tabular ore body that parallels the bedding or layering of the host rock. Thickness is small compared to lateral extent. Typical of sedimentary exhalative (SEDEX), volcanogenic massive sulphide (VMS) and banded iron formations (BIF).

DEFINITION Stratabound Deposit

An ore body confined to a specific stratigraphic unit but not necessarily parallel to bedding; it may be discordant within that unit. Example: Mississippi Valley Type (MVT) lead-zinc in a specific carbonate horizon.

DEFINITION Vein and Lode

A **vein** is a sheet-like body filling a fracture or fault; a **lode** is a mineralised fault zone (often multiple veins + altered wall rock). Both are **discordant** — they cut across host-rock layering.

DEFINITION Stockwork and Dissemination

A **stockwork** is a network of closely spaced, small veins; a **dissemination** consists of fine-grained ore minerals scattered through the host rock. Both imply bulk-mining methods (porphyry Cu, epithermal Au).


DEFINITION Pipe, Pod and Lens

Pipe: cylindrical, near-vertical body (kimberlite, breccia pipe). **Pod**: small, irregular, blob-like. **Lens**: lenticular, pinching out in all directions.

DEFINITION Placer Deposit

Concentration of heavy, resistant minerals (gold, cassiterite, diamond, ilmenite) in un-

consolidated sediments — river beds (alluvial), beaches, palaeo-channels. Morphology follows paleo-drainage.



Main morphological types of ore bodies in cross-section. (a) Stratiform layer parallel to bedding. (b) Discordant vein/lode cutting strata. (c) Stockwork of veinlets. (d) Placer concentrate in river channel. (e) Vertical pipe.

WARNING Morphology $\neq$ Genesis

A vein is a shape; a hydrothermal deposit is an origin. A hydrothermal deposit can be a vein, a stockwork, a replacement body or a stratiform exhalative layer. In exam questions, if asked "Classify the deposit by morphology", answer vein, stratiform, stockwork, etc. If asked "Classify by genesis", answer magmatic, hydrothermal, sedimentary, etc. Mixing the two criteria loses marks.

3.3 Classification by Relation to Host Rock: Syngenetic vs Epigenetic

This criterion asks: Did the ore form at the same time as the host rock, or later? It is the single most powerful predictor of where to drill next.

DEFINITION Syngenetic Deposit

Ore minerals crystallised simultaneously with the host rock. The ore is an integral part of the stratigraphy. Examples: Banded Iron Formations (BIF), sedimentary manganese, VMS lenses, magmatic chromite layers in layered intrusions.

DEFINITION Epigenetic Deposit

Ore minerals were introduced after the host rock had formed and usually after it had lithified. The ore cuts across, replaces or fills voids in the pre-existing rock. Examples: Hydrothermal veins, porphyry copper, skarns, Mississippi Valley Type (MVT), placer deposits (the host sediment is younger than the source rock).

METHOD Recognising Syngenetic vs Epigenetic on a Cross-Section

1. Step 1: Identify the primary layering/bedding of the host rock (sedimentary beds, volcanic flow bands, igneous layering).
2. Step 2: Trace the ore-body boundaries.
3. Step 3: Concordant (parallel to layering) → suggests syngenetic (but check for replacement!). Discordant (cross-cutting) → epigenetic.
4. Step 4: Look for alteration halos (epigenetic) vs primary sedimentary structures preserved in ore (syngenetic).
5. Step 5: Check age relationships: cross-cutting veins, contact metamorphism, unconformities.

EXAMPLE Syngenetic vs Epigenetic in the Field

Case A: In the Ntem Complex (South Cameroon), a layer of massive chromitite follows the magmatic layering of a dunite-peridotite sequence. No cross-cutting veins, no alteration halo. → Syngenetic magmatic.

Case B: At Batouri (East Cameroon), gold-bearing quartz veins cut across metasedimentary schists and are surrounded by sericite-pyrite alteration halos. → Epigenetic hydrothermal.

3.4 Classification by Genetic Process (Overview)

Genesis explains how the ore formed. The GCE syllabus recognises five main families. Detailed mechanisms are covered in Sections 4 and 5; here we give the recognition criteria for each family.

Genetic Family	Key Processes	Typical Examples (Cameroon / World)
Magmatic	Crystallisation from silicate melt: early crystallisation (chromite, PGE), immiscible sulphide liquids (Ni-Cu-PGE), residual liquids (pegmatites).	Nkamouna (Co-Ni-Mn laterite over ultramafic), Bushveld (PGE), Sudbury (Ni-Cu).
Hydrothermal	Precipitation from hot aqueous fluids: veins, stockworks, replacement, VMS, porphyry, epithermal, skarn.	Batouri (Au), Mayo Darle (W-Sn), Ok Tedi (Cu-Au porphyry), Kidd Creek (VMS).
Sedimentary	Chemical/biochemical precipitation from water: BIF, manganese, phosphorite, evaporites, placer (mechanical).	Mbalam/Nabeba (BIF Fe), Bafoussam (bauxite – residual), Witwatersrand (Au-U paleo-placer).
Residual / Weathering	In-situ chemical weathering concentrates insoluble residues: lateritic Ni-Co, bauxite, supergene enrichment.	Nkamouna (Co-Ni laterite), Minim-Martap (bauxite), New Caledonia (Ni laterite).
Metamorphic	Recrystallisation / remobilisation during metamorphism: graphite, kyanite, some Au deposits.	Mfou (graphite), Lom (kyanite).

DID YOU KNOW? Cameroon's Genetic Diversity

Cameroon hosts all five genetic families along the **Cameroon Volcanic Line (CVL)** and the **Congo Craton** margin: magmatic (chromite in ultramafics), hydrothermal (Au-W-Sn in Pan-African granites), sedimentary (BIF iron in Mbalam), residual (Co-Ni laterites at Nkamouna, bauxite on the Adamawa Plateau), metamorphic (graphite in Yaoundé Group). This makes Cameroon a natural laboratory for economic geology.

3.5 Classification by Commodity (Economic Grouping)

This classification speaks the language of economists and metallurgists.

DEFINITION Metallic Deposits

Ores from which metals are extracted. Sub-groups:

- Ferrous: Fe, Mn, Cr (steel industry).
- Base metals: Cu, Pb, Zn, Ni, Co (industrial bulk).
- Precious metals: Au, Ag, PGE (high value, low tonnage).

- Strategic / Critical: W, Sn, Ta, Nb, Li, REE, Co (high-tech, supply risk).

DEFINITION Non-Metallic (Industrial) Deposits

Valued for physical/chemical properties, not metal content: limestone, phosphate, potash, salt, gypsum, barite, fluorite, graphite, kaolin, dimension stone, aggregates.

DEFINITION Energy Minerals

Uranium, thorium, coal, oil shale, geothermal — sometimes treated separately.

WARNING Commodity $\neq$ Genesis

A base-metal deposit can be hydrothermal (VMS), sedimentary (SEDEX) or magmatic (Ni-Cu sulphide). Always state the genetic family and the commodity group when describing a deposit fully.

3.6 Concordant vs Discordant Ore Bodies: Mining Consequences

This geometric distinction, rooted in the syngenetic/epigenetic divide, has direct engineering impact.

PROPERTY Concordant Ore Bodies

Follow host-rock layering (stratiform, some stratabound, some replacement bodies).

- Mining: Amenable to open-pit (if near surface) or room-and-pillar / longwall underground methods. High continuity → predictable reserves.
- Dilution: Low if contacts are sharp; high if gradational.
- Example: Mbalam BIF iron ore — open pit, low stripping ratio.

PROPERTY Discordant Ore Bodies

Cut across layering (veins, lodes, pipes, stockworks, some breccias).

- Mining: Often selective underground (cut-and-fill, shrinkage stoping, sublevel caving for stockworks). Irregular geometry → higher exploration density needed.

- Dilution: High — wall rock must be broken with ore.
- Example: Batouri gold veins — narrow, high-grade, underground.

KEY POINTS Exam Checklist: Describe a Deposit Fully

When a question says “Classify the following deposit”, give four labels:


1. Morphology (vein, stratiform, stockwork, placer, pipe...)
2. Host-rock relation (syngenetic / epigenetic)
3. Genetic family (magmatic, hydrothermal, sedimentary, residual, metamorphic)
4. Commodity group (ferrous, base, precious, strategic, industrial, energy)

3.7 Summary Table: One Example per Morphological & Genetic Type

Morphology	Genetic Family	Deposit Example
Stratiform	Sedimentary (BIF)	Mbalam / Nabeba (Cameroon–Congo border)
Stratiform	Magmatic (layered intrusion)	Bushveld Complex (South Africa)
Vein / Lode	Hydrothermal (orogenic Au)	Batouri (East Cameroon)
Stockwork	Hydrothermal (porphyry Cu)	Ok Tedi (Papua New Guinea) / Mayo Darle (Cameroon)

3.8 Classification Tree: A Visual Synthesis

The following diagram organises the four classification criteria into a single decision tree. Use it as a mental checklist during field mapping or exam revision.



3.9 Worked Example: Full Classification of a Deposit

WORKED EXAMPLE 2 Classifying the Nkamouna Cobalt–Nickel Deposit (East Cameroon)

Statement: The Nkamouna deposit consists of a 10–30 m thick blanket of cobalt–nickel–manganese oxides (asbolane, lithiophorite) overlying weathered ultramafic rocks (serpentinised peridotite) of the Ntem Complex. The ore zone follows the paleo–topography and shows a clear vertical zonation from Fe–rich duricrust at surface to Co–Ni–Mn enrichment at depth. Classify this deposit using all four criteria.

Detailed solution:

1. Morphology: **Residual blanket / lateritic profile**. The ore forms a horizontal to gently undulating layer parallel to the paleo–surface, not to bedrock structure. It is a weathering mantle, not a primary igneous layer.
2. Host–rock relation: **Epigenetic** with respect to the bedrock (the ultramafic host is Archaean, the ore formed during Cenozoic weathering). However, it is **syngenetic** with respect to the lateritic profile itself (the Co–Ni enrichment formed concurrently with the laterite horizons). Always specify the reference frame!
3. Genetic family: **Residual / Weathering (supergene lateritic)**. Intense tropical weathering leached Mg, Si, Fe (partly) and concentrated immobile Co, Ni, Mn as oxide/hydroxide minerals in the saprolite and mottled zone.
4. Commodity group: **Strategic / Critical metals** (Co for batteries, Ni for stainless steel, Mn for alloys). Also base metals by traditional classification.

Final classification statement: “Nkamouna is a residual blanket (morphology), epigenetic–to–bedrock / syngenetic–to–profile (host relation), lateritic supergene (genesis), strategic Co–Ni–Mn (commodity) deposit.”

3.10 Exercises

EXERCISE 2 Classification Practice

For each of the following descriptions, give the four classification labels (Morphology, Host-relation, Genesis, Commodity).

1. A 2 m thick layer of magnetite–quartzite intercalated with amphibolites in the Yaoundé Group (Pan–African belt). The layer is folded with the host rocks.
2. A swarm of 5–20 cm wide quartz–tourmaline–cassiterite veins cutting a two–mica granite near Mayo Darle (Adamawa). Veins are surrounded by greisen alteration.
3. Rounded grains of gold and diamond concentrated in the gravels of the Sangha River palaeo–channels (East Cameroon).
4. A lenticular body of massive chromitite (5 m × 200 m) within a dunite layer of the Ntem Complex ultramafic suite. Contacts are sharp and parallel to magmatic layering.
5. A 500 m × 200 m × 50 m zone of disseminated chalcopyrite and molybdenite in a granodiorite stock, with a surrounding halo of propylitic alteration (chlorite–epidote).

SOLUTION Exercise 1

1. Morphology: Stratiform. Host-relation: Syngenetic (sedimentary/exhalative). Genesis: Sedimentary (BIF–type) + metamorphic overprint. Commodity: Ferrous (Fe).
2. Morphology: Vein / Lode (swarm). Host-relation: Epigenetic. Genesis: Hydrothermal (granite–related, Sn–W). Commodity: Strategic (Sn, W).
3. Morphology: Placer (alluvial / palaeo–channel). Host-relation: Epigenetic (host sediment younger than source). Genesis: Sedimentary (mechanical concentration). Commodity: Precious (Au) + Gem (diamond).
4. Morphology: Lens / Pod (within layer). Host-relation: Syngenetic (magmatic). Genesis: Magmatic (early crystallisation / immiscible liquid). Commodity: Strategic (Cr, PGE).
5. Morphology: Disseminated / Stockwork. Host-relation: Epigenetic. Genesis: Hydrothermal (porphyry Cu–Mo). Commodity: Base metals (Cu, Mo).

WARNING Common Exam Traps

- Trap 1: Calling a BIF “metamorphic” because it is in gneiss. Correction: The protolith is

sedimentary; metamorphism recrystallised it. Classify by primary genesis: Sedimentary (BIF).

- Trap 2: Calling a placer “syngenetic” because the gold grains are in the sediment. Correction: The sediment is the host; the gold was derived from an older source. → Epigenetic.
- Trap 3: Confusing stockwork (many small veins) with dissemination (individual grains). In a stockwork you see veinlets; in a dissemination you need a hand lens.
- Trap 4: Forgetting to state the reference frame for syngenetic/epigenetic (bedrock vs weathering profile vs sediment).

DID YOU KNOW? From Classification to Exploration: The Cameroon Example

The Minim-Martap bauxite deposit (Adamawa Plateau) was discovered by morphological reasoning: lateritic plateaux (residual blanket morphology) overlying Cretaceous basalts (host rock) in a tropical climate → target for residual/weathering genesis → bauxite (non-metallic/industrial commodity). The same logic guides exploration for Nkamouna-style Co-Ni laterites: look for ultramafic bedrock + deep weathering profile + plateau topography. Classification is not academic — it is the explorationist's compass.

3.11 Key Takeaways

KEY POINTS What You Must Remember

- Classification uses four independent criteria: Morphology, Host-rock relation (Syngenetic/Epigenetic), Genesis, Commodity. Always give all four.
- Morphology controls mining method: stratiform → open pit / longwall; veins/pipes → selective underground; stockworks → bulk underground.
- Syngenetic = ore formed with host rock (concordant, stratigraphic control). Epigenetic = ore introduced later (discordant, structural control).
- Five genetic families: Magmatic, Hydrothermal, Sedimentary, Residual/Weathering, Metamorphic. Cameroon has examples of all five.
- Commodity groups: Ferrous, Base, Precious, Strategic (metallic); Industrial (non-metallic); Energy.
- Concordant ore bodies are stratigraphically controlled; discordant bodies are structurally controlled.

- Exam tip: When asked to “classify”, explicitly list the four labels. Never mix morphology terms (vein) with genesis terms (hydrothermal) as if they were the same level of classification.

The next two sections will dive deep into the **formation processes** of each genetic family — first magmatic and hydrothermal (Section 4), then sedimentary, weathering and metamorphic (Section 5). Master this classification framework now; it is the scaffold on which all those genetic models will hang.

4 FORMATION OF ORE DEPOSITS I: MAGMATIC AND HYDROTHERMAL PROCESSES

4.1 From Classification to Genesis


In the previous section we learned how mineral deposits are classified according to their morphology, host rock and genetic affiliation. Classification tells us what a deposit looks like; genesis explains how it formed. Understanding the processes that concentrate metals from trace levels in the Earth’s crust into mineable ore bodies is the core of economic geology. Two great families of processes dominate the formation of the world’s major metal deposits: **magmatic processes** that operate while a melt is still largely liquid, and **hydrothermal processes** that involve hot, aqueous fluids moving through the crust. Both families are intimately linked — magmas supply the heat, the metals and often the fluids themselves. In this section we follow the journey of metals from a cooling magma chamber to the veins and stockworks that make up many of Cameroon’s gold and base-metal prospects.

4.2 Magmatic Processes: Segregation, Immiscibility and the Pegmatitic Stage

Magmatic segregation is the physical separation of crystals from a melt by gravity. As a mafic magma cools, dense oxide and sulphide minerals crystallise early and, because they are heavier than the surrounding silicate liquid, they sink toward the floor of the intrusion. This produces layered intrusions with rhythmic bands of chromite, magnetite, ilmenite and platinum-group minerals — classic examples are the Bushveld Complex (South Africa) and the Ntem Complex in southern Cameroon.

DEFINITION Magmatic Segregation

The gravitational settling of early-formed, dense crystals (oxides, sulphides, platinum-group minerals) from a convecting magma, producing layered concentrations at the base of an intrusion.



Cross-section of a cooling mafic intrusion showing crystal settling (chromite, magnetite, ilmenite, platinum layers) and an immiscible sulphide liquid accumulating at the base.

A second magmatic mechanism is **magmatic immiscibility**. When a mafic magma becomes saturated in sulphur, a separate sulphide liquid phase splits off from the silicate melt — just as oil separates from water. This sulphide liquid is an extremely efficient scavenger of chalcophile metals (Ni, Cu, Co, PGE). Because it is denser than the silicate melt, it also sinks and pools at the base of the intrusion, forming massive sulphide ore bodies (e.g. the Sudbury nickel-copper deposit in Canada). In Cameroon, the Ntem Complex shows evidence of such immiscible sulphide segregation in its gabbroic units.

DEFINITION Immiscible Sulphide Liquid

A dense, sulphur-rich liquid phase that separates from a silicate magma upon sulphur saturation, strongly partitioning chalcophile metals (Ni, Cu, Co, PGE) and sinking to form magmatic sulphide deposits.

The final magmatic stage is the **pegmatitic stage**. As crystallisation proceeds, the residual melt becomes enriched in water, fluxing components (B, F, P) and incompatible elements (Li, Be, Sn, Ta, Nb, rare earths). This highly fluid, low-viscosity melt intrudes fractures and crystallises as coarse-grained pegmatites. Pegmatites are the primary source of tin (cassiterite), tantalum (columbite-tantalite), lithium (spodumene, lepidolite) and gemstones (beryl, tourmaline). The Central African tin belt, which extends through the Dja River region of East Cameroon, owes its mineralisation to late-stage pegmatitic fluids derived from the Pan-African granites.

DEFINITION Pegmatite

An exceptionally coarse-grained igneous rock formed from the final, volatile-rich residual melt of a granitic magma, typically enriched in rare elements (Li, Be, Sn, Ta, Nb, gemstones).

DID YOU KNOW? Cameroon's Tin and Rare-Metal Potential

The Mbalam and Bipindi areas in the East Region host alluvial cassiterite derived from weathered pegmatites. Exploration targeting the parent granites and their pegmatitic apophyses could reveal primary Sn-Ta-Nb deposits similar to those of the Manono-Kitotolo belt in the DRC.

4.3 Hydrothermal Processes: Fluids, Transport and Deposition

When magmas crystallise, they release large volumes of hot, metal-bearing aqueous fluids — **hydrothermal solutions**. These fluids may also originate from meteoric water (rainwater that circulates deep into the crust) or connate water (trapped sedimentary brine). Regardless of origin, hydrothermal fluids are the principal agents that transport metals over kilometres and deposit them in fractures, pore spaces and reactive host rocks.

DEFINITION Hydrothermal Solution

A hot (typically 100 °C–600 °C), pressurised aqueous fluid, often saline and acidic, capable of dissolving, transporting and precipitating metallic ions in the Earth's crust.

PROPERTY Temperature–Pressure Control on Metal Solubility

The solubility of most ore metals (Cu, Pb, Zn, Au, Ag) in hydrothermal fluids decreases as temperature and pressure fall. Consequently, cooling, decompression, boiling, fluid mixing or reaction with wall rocks triggers metal precipitation.

Hydrothermal deposition occurs in three main geometric styles:


- **Vein filling** — fluids precipitate minerals in open fractures (e.g. quartz–gold veins).
- **Replacement** — fluids dissolve the host rock and simultaneously precipitate ore minerals, preserving the original rock texture (e.g. Mississippi Valley-type Pb–Zn in limestone).
- **Dissemination** — fine-grained ore minerals precipitate throughout a permeable rock volume (e.g. porphyry copper stockworks).

DEFINITION Replacement (Metasomatism)

A hydrothermal process in which a mineral assemblage is chemically replaced by a new assemblage, the fluid simultaneously removing the original components and supplying the new ones, often preserving the original rock fabric.

4.4 Metal Zoning Around Intrusions: Porphyry and Epithermal Systems

A classic illustration of hydrothermal zoning is the **porphyry copper–gold system**. A small, porphyritic granodiorite to diorite intrusion (the porphyry stock) emplaces at shallow crustal levels (1–3 km). It releases a magmatic–hydrothermal fluid that creates a dense network of quartz–veinlets — a stockwork — in the intrusion and the surrounding country rock. Metals precipitate in concentric shells reflecting decreasing temperature and changing fluid chemistry:



Schematic cross-section of a porphyry Cu–Au system showing the central stock, stockwork veinlets, and concentric metal zoning (Cu–Mo core → Zn–Pb shell → Ag–Au halo). Epithermal veins form above the porphyry environment.

EXAMPLE Porphyry–Epithermal Link in East Cameroon

The Batouri and Bertoua greenstone belts host numerous artisanal gold workings. Geological mapping reveals that many quartz–gold veins are spatially associated with small, porphyritic granodiorite intrusions — the roots of ancient porphyry systems. The high-temperature Cu–Mo core is often eroded away, leaving the lower-temperature Au–Ag epithermal veins exposed at surface. Recognising this genetic link guides exploration: soil geochemistry for Cu–Mo pathfinders can vector toward buried porphyry centres beneath the gold veins.

4.5 The Critical Role of Structures

Faults, fractures, shear zones and lithological contacts act as high-permeability pathways that channel hydrothermal fluids from their deep source to the depositional site. They also serve as traps: a change in rock competency, a fault bend or a lithological contact can cause a pressure

drop, fluid boiling or chemical reaction — all triggering precipitation. In the Adamawa Plateau, the Sanaga Fault and its splays control the localisation of both magmatic (Ni-Cu) and hydrothermal (Au) mineralisation.

METHOD Reconstructing Mineralisation Sequence from Cross-Cutting Veins

1. Map all vein sets, noting orientation, mineralogy and alteration halos.
2. Identify cross-cutting relationships: a vein that cuts another is younger.
3. Record the mineral assemblage of each vein generation (e.g. early quartz-pyrite-chalcopyrite → late quartz-gold-sericite).
4. Correlate vein generations with known regional deformation events (e.g. D1, D2).
5. Build a paragenetic sequence table linking structural event, fluid temperature (from fluid inclusions) and metal assemblage.

WARNING Epigenetic Nature of Hydrothermal Deposits

Even when hydrothermal veins occur inside the intrusion that supplied the fluid, they are epigenetic — they formed after the host rock solidified. Do not classify them as magmatic (syngenetic) simply because of their spatial association.

WARNING Gold: Primary vs. Placer

Most gold in the crust is deposited by hydrothermal fluids (primary gold). Placer gold is secondary, derived from the erosion of primary hydrothermal veins. In Cameroon, the alluvial gold of the Sangha and Boumba rivers originates from the primary quartz-gold veins of the East Region greenstone belts.

4.6 Worked Example: Interpreting a Porphyry Cross-Section

WORKED EXAMPLE 3 Porphyry Zoning Interpretation

Statement: A geological cross-section through a porphyry system shows (from centre outward): a stockwork of quartz-chalcopyrite-molybdenite veinlets in a granodiorite; a surrounding zone of quartz-sphalerite-galena veins in the contact aureole; and an outer halo of quartz-arsenopyrite-gold veins in the country rock. Fluid inclusion data give homogenisation temperatures of 420 °C (core), 300 °C (intermediate) and 200 °C (outer).

Explain the metal zoning and suggest an exploration vector for the buried Cu–Mo core.
Detailed solution:

1. Identify the three zones: The mineral assemblages and temperatures match the classic porphyry zoning: Cu–Mo (chalcopyrite + molybdenite) at highest T, Zn–Pb (sphalerite + galena) at intermediate T, Au–Ag–As (arsenopyrite + gold) at lowest T.
2. Explain the zoning: As the magmatic–hydrothermal fluid migrates outward, it cools and depressurises. Metal solubility drops sequentially: Cu and Mo precipitate first (high T, high fS_2), then Zn and Pb, finally Au and Ag (low T, lower fS_2 , possible boiling). Wall-rock reaction (e.g. with carbonate) can also shift pH and cause deposition.
3. Exploration vector: The Cu–Mo core is centred on the porphyry stock. Soil/stream sediment surveys for Cu, Mo and pathfinder elements (Re, Se, Te) should be focused up-gradient (toward the centre of the stockwork) from the known Zn–Pb and Au anomalies. Geophysical IP/resistivity surveys can delineate the stockwork chargeability high that marks the core.
4. Key takeaway: Metal zoning is a direct consequence of the temperature–solubility relationship (Property above). Mapping the zoning allows the geologist to vector toward the highest-value part of the system.

4.7 Key Points to Retain

KEY POINTS Formation of Ore Deposits I — Essentials

- Magmatic segregation (crystal settling) forms layered oxide/sulphide deposits in mafic intrusions (chromite, Pt).
- Magmatic immiscibility produces dense sulphide liquids that scavenge Ni, Cu, PGE and settle as massive sulphides.
- Pegmatites concentrate Li, Be, Sn, Ta, Nb and gemstones from the final volatile-rich melt.
- Hydrothermal fluids (magmatic, meteoric, connate) transport metals in solution; solubility falls with T and P, causing precipitation.
- Deposition styles: vein filling, replacement, dissemination.
- Porphyry Cu–Au systems show concentric metal zoning (Cu–Mo core → Zn–Pb shell → Ag–Au halo) linked to cooling fluid.
- Epithermal veins (low-sulphidation Au–Ag) form above porphyry environments; they are the exposed expression of deeper systems.

- Structures (faults, fractures, contacts) are fluid pathways and traps — essential for localising ore.
- Hydrothermal deposits are epigenetic; primary gold is hydrothermal, placers are secondary.

5 FORMATION OF ORE DEPOSITS II: SEDIMENTARY, WEATHERING AND METAMORPHIC PROCESSES

In the previous section we followed ore-forming fluids upwards from magmas and hot hydrothermal systems deep inside the crust. But not all ores are born in fire. Walk along the Wouri estuary, or across the red plateaux of the Adamaoua, and you are standing on ore deposits that formed at or near the Earth's surface, at ordinary temperatures, through the patient work of water, wind, organisms and time. Rivers sort and concentrate heavy minerals; tropical rain leaches away silica and leaves aluminium behind; seawater evaporates and abandons its salts; and when rocks are later buried and heated, metamorphism recrystallises old materials into new economic products. This section studies these **exogenous** (surface) and **metamorphic** processes, which together account for most of the world's iron, aluminium, salt, phosphate, placer gold and diamonds.

5.1 Sedimentary (Syngenetic) Deposits

Observation. When a shallow coastal lagoon or an enclosed salt lake evaporates under a hot sun, white crusts of salt appear at its margins. When certain ancient seas stagnated, iron precipitated layer upon layer on their floors. In both cases, the ore was deposited at the same time as the enclosing sediment: such deposits are called **syngenetic**.

DEFINITION Syngenetic sedimentary deposit

A **syngenetic sedimentary deposit** is an ore body formed by the direct precipitation or accumulation of useful material from surface waters (sea, lake, lagoon) at the same time as the sediments that enclose it. It typically forms extensive, regular beds or seams interstratified with normal sedimentary rocks.

Chemical precipitates. Three great families are distinguished.

1. **Banded iron formations (BIF).** In the early history of the Earth (roughly before 1.8 billion years ago), the oceans contained abundant dissolved iron and the atmosphere little oxygen. As photosynthetic organisms released oxygen, the iron oxidised and precipitated, producing alternations of iron-rich (haematite, magnetite) and silica-rich (chert) bands. These **banded iron formations** supply the greater part of the world's iron ore (e.g. the enormous deposits of Australia, Brazil, and the Kribi-Mbalam iron province of southern Cameroon).

2. **Evaporites.** When an arm of the sea or a salt lake is cut off and evaporation exceeds inflow, dissolved salts crystallise in a predictable order of increasing solubility: first carbonates, then **gypsum** ($\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$), then rock salt (**halite**, NaCl), and finally the highly soluble **potash** salts (sylvite, KCl). Evaporite beds are mined for salt, fertiliser and chemical industries.
3. **Biochemical accumulations: phosphates.** In warm, shallow seas with abundant life, phosphorus from organic remains (bones, shells, guano) accumulates on the sea floor and is converted to **phosphate** rock (apatite), the essential raw material of fertilisers.

WARNING Syngenetic versus epigenetic

Do not confuse syngenetic (ore formed with the host sediment, as a bed) with epigenetic (ore introduced later into pre-existing rocks, as veins). A salt bed is syngenetic; a gold-bearing quartz vein cutting across schists is epigenetic. GCE examiners regularly test this distinction.

5.2 Placer Deposits: Concentration by Moving Water


Observation. A gold panner on the River Lom or the Vina swirls gravel in a pan: the light sand washes away while dense, shiny grains settle at the bottom. The river has already done most of this sorting naturally. Minerals that are both dense and chemically resistant survive weathering and transport, and accumulate where the current slackens.

DEFINITION Placer deposit

A **placer** is a deposit formed by the mechanical concentration of dense, durable minerals (gold, diamond, cassiterite SnO_2 , ilmenite FeTiO_3 , platinum, monazite) by moving water, after the weathering and erosion of a primary source rock. Placers are classified by their setting:

- **eluvial** placers: concentrated by gravity and creep on slopes, just downhill from the source;
- **alluvial** placers: concentrated in river gravels, at the base of the gravel bed, in pot-holes and behind obstacles;
- **beach** placers: concentrated by wave action along shorelines (e.g. ilmenite and rutile sands).

The concentrating agent is hydraulic sorting: a current that can carry quartz (density 2.65) may be unable to move gold (density 19.3) or cassiterite (density 6.9). The heavy grains therefore sink through the moving gravel and collect at its base, on or near the bedrock, often in cracks, potholes and the lee of boulders or bars.



Placer concentration in a river: dense gold grains sink through the gravels and accumulate at the base, in bedrock hollows and behind obstacles.

METHOD Recognising a placer versus a primary vein

1. Grain shape: placer grains are rounded and worn by transport; vein minerals are angular, interlocking crystals.
2. Association: placers contain a mixture of dense, resistant minerals (gold + cassiterite + ilmenite + magnetite); veins contain quartz, sulphides and gangue minerals.
3. Geometry: placers follow sedimentary layers (gravel beds, terraces, beaches); veins cut across the rock structure.
4. Context: a placer lies in an unconsolidated sedimentary deposit near a drainage system; a vein lies within hard, often altered bedrock.

DID YOU KNOW? Placers of Cameroon

Alluvial gold is washed by local miners in the rivers of the East Region (Bétaré-Oya, Batouri) and the Adamaoua, and diamonds have long been recovered from placers in the Mobilong area near Yokadouma, close to the Central African border. These placers point to primary sources in the surrounding Precambrian basement rocks.

5.3 Residual (Weathering) Deposits: Laterites and Bauxite

Observation. Laterite road cuttings around Yaoundé show a striking vertical zonation: dark fresh rock at depth passes upward into soft, blotched, clay-rich material, then into a red, iron-rich horizon, sometimes capped by a hard crust. This is a **weathering profile**, and it is an ore factory.

Under a **humid tropical climate** — high temperature, abundant rainfall, dense vegetation — rain-water charged with CO_2 and organic acids attacks the bedrock. The most soluble elements (Ca, Na, K, and even SiO_2) are dissolved and carried away, while the least mobile elements (Al, Fe, Ti) remain and are relatively enriched. This is residual concentration: nothing is added, but


the removal of the soluble fraction concentrates what is left.

DEFINITION Laterite and bauxite

A **laterite** is a red, residual soil and weathering mantle rich in iron and aluminium oxides and hydroxides, formed by intense tropical weathering. **Bauxite** is the aluminium-rich end-product of this process: a rock composed essentially of aluminium hydroxide minerals (gibbsite $\text{Al}(\text{OH})_3$, boehmite, diaspore) with iron oxides, clay and residual quartz. A **residual deposit** is any ore formed by such in-situ enrichment during weathering.

PROPERTY Condition for residual concentration

Residual deposits form where soluble elements (Si, Ca, Na, K) are leached away faster than immobile elements (Al, Fe) are removed. They therefore require: (i) a warm, humid climate with strong chemical weathering; (ii) good drainage; (iii) a stable land surface (low erosion) preserved for a long time; and (iv) a suitable parent rock (aluminosilicate rocks for bauxite, ultrabasic rocks for nickel laterite).



Ideal lateritic weathering profile. Downward-percolating water removes soluble elements; immobile Al and Fe accumulate residually in the upper horizons.

Main residual deposits.

- **Bauxite (aluminium ore)**: forms by lateritic weathering of aluminosilicate rocks (granites, gneisses, syenites) under humid tropical climates. The great Cameroonian example is the **Minim-Martap / Ngaoundal bauxite** on the Adamaoua Plateau: a thick lateritic mantle developed over syenitic and granitic basement, preserved on a high, stable, well-drained peneplain.
- **Lateritic nickel**: weathering of ultrabasic rocks (peridotites) leaves a mantle enriched in nickel

(e.g. New Caledonia; nickel–cobalt laterites occur in the Lomíé region of East Cameroon).

- **Residual iron and manganese:** iron and manganese can be concentrated in lateritic caps and weathered zones above lower-grade protores.

WARNING Bauxite is a rock, not a mineral


A very frequent GCE error: bauxite is a rock — a mixture of aluminium hydroxide minerals (gibbsite, boehmite, diaspore) plus impurities — not a single mineral with a fixed formula. Similarly, laterite is a weathering product (a soil/rock type), not a mineral species.

5.4 Supergene Enrichment

Observation. Many copper sulphide deposits are too poor to mine in their primary state, yet their upper parts have been naturally upgraded by weathering. How? Descending rainwater dissolves copper from the oxidised top of the deposit and re-deposits it lower down, just below the water table, where it is safe from further oxidation.

DEFINITION Supergene enrichment

Supergene enrichment is the process by which a metal (especially copper) is leached from the oxidised upper part of a sulphide deposit, transported downward in solution, and re-precipitated as rich secondary sulphides (e.g. chalcocite Cu_2S , covellite CuS) just below the water table, thereby upgrading a low-grade primary ore into an economic one.



Supergene enrichment of a copper sulphide deposit: copper leached from the oxidised zone is re-precipitated as rich secondary sulphides just below the water table.

The process works because copper is soluble in oxidising, acidic surface waters but precipitates in the reducing environment below the water table. Without this natural upgrading, many of the world's great copper camps (e.g. in Zambia and the Congolese Copperbelt) would not have been economic.

5.5 Metamorphic and Metamorphosed Deposits

Finally, some deposits owe their economic value to **metamorphism** — the transformation of rocks by heat and pressure.

- **Metamorphosed deposits** are pre-existing ores modified by metamorphism: recrystallisation may improve quality (e.g. banded iron formations recrystallised into coarse, easily concentrated haematite ores).
- **Metamorphic deposits** are new economic materials created by metamorphism itself: **graphite** from carbon-rich sediments, **asbestos** and **talc** from ultrabasic rocks, **marble** from limestone, garnet and kyanite from aluminous sediments.
- **Skarn (contact-metasomatic) deposits** form where a hot intrusion invades carbonate rocks (limestone, dolomite): fluids from the magma react with the carbonate wall-rock, producing calcium-silicate minerals (garnet, pyroxene) together with ores of iron, copper, tungsten, lead and zinc.

DEFINITION Skarn

A **skarn** is a deposit formed at the contact between an igneous intrusion and a carbonate host rock, where hot magmatic fluids chemically replace the limestone or dolomite with calcium-magnesium silicate minerals and associated ores (magnetite, chalcopyrite, scheelite, sphalerite, galena).

5.6 Synthesis and Worked Exercise

Genetic process	Typical products	Deposit morphology	Example
Chemical precipitation (sedimentary)	Iron (BIF), salt, gypsum, potash	Extensive regular beds	BIF of the Kribi–Mbalam iron province
Biochemical accumulation	Phosphates	Beds and nodules in marine sediments	Phosphate basins of Morocco
Mechanical concentration (placer)	Gold, diamond, cassiterite, ilmenite	Gravel beds, terraces, beach sands	Gold placers of Bétaré-Oya (East Cameroon)
Residual weathering (lateritic)	Bauxite, lateritic Ni, Fe, Mn	Blanket-like weathering mantle on plateaux	Minim–Martap / Ngaoundal bauxite (Adamaoua)
Supergene enrichment	Copper (secondary sulphides)	Horizontal enriched horizon below water table	Zambian Copperbelt
Metamorphic / skarn	Graphite, talc, marble; Fe, Cu, W ores	Irregular bodies at intrusion contacts	Skarns of the Urals (Russia)

WORKED EXAMPLE 4 Explaining the Adamaoua bauxite

Statement. Using your knowledge of residual deposits, explain why economic bauxite occurs on the Adamaoua Plateau (Minim–Martap–Ngaoundal), and state two other products of tropical residual concentration.

SOLUTION

Solution.

1. Parent rock: the plateau is underlain by aluminosilicate rocks (granites, syenites, gneisses) rich in aluminium-bearing feldspars.
2. Climate: the region experiences a humid tropical climate with high rainfall and temperature, causing intense chemical weathering by CO₂-charged rainwater.
3. Process: soluble elements (SiO₂, Ca, Na, K) are leached downward and removed, while immobile aluminium accumulates as hydroxides (gibbsite, boehmite), producing a thick bauxitic laterite mantle.
4. Preservation: the plateau is a high, flat, well-drained and long-stable surface, so the weathering mantle was neither waterlogged nor eroded away, allowing a thick, high-

grade bauxite blanket to accumulate.

5. Other residual products: lateritic nickel (from ultrabasic rocks) and residual iron or manganese.

EXERCISE 3 Placer or vein?

A prospector finds gold as (a) angular grains locked inside a quartz vein cutting across schists, and (b) rounded flakes in the basal gravel of a river terrace. For each occurrence, state the type of deposit, the process that formed it, and one feature that justifies your answer.

SOLUTION Exercise 1

(a) Primary (hypogene) vein deposit: the gold crystallised from hydrothermal fluids within a fracture; justification — angular grains enclosed in vein quartz that cuts across the country rock (epigenetic geometry). (b) Alluvial placer deposit: the gold was liberated by weathering of a primary source and mechanically concentrated by river action; justification — rounded, water-worn flakes in a sedimentary gravel bed, at the base of the terrace where dense minerals settle.

KEY POINTS Key points

- Syngenetic sedimentary deposits (BIF, evaporites, phosphates) form as beds at the same time as their host sediments.
- Placers concentrate dense, resistant minerals (Au, diamond, cassiterite) by hydraulic sorting in rivers and on beaches.
- Lateritic weathering in humid tropical climates leaches Si, Ca, Na, K and residually enriches Al and Fe: this yields bauxite (Minim–Martap) and lateritic nickel.
- Supergene enrichment upgrades copper ores by leaching above and re-deposition below the water table.
- Metamorphism creates graphite, talc, marble and skarn ores at intrusion–limestone contacts.
- Exam tip: explicitly linking Cameroon’s humid tropical climate to lateritic bauxite (Adamaoua) and placer gold (East Region) earns marks in questions on national mineral wealth.

6 OCCURRENCE OF MINERAL DEPOSITS: FROM THE WORLD TO CAMEROON

In the previous two sections we learned how ore deposits form: by magmatic segregation, by hydrothermal circulation, by sedimentary concentration, by tropical weathering and by metamorphic remobilisation. A natural question now arises: where do these processes actually operate on the Earth's surface, and can we predict where to look for a given metal? A prospector in the East Region of Cameroon does not dig at random: he pans for gold in streams draining ancient greenstone belts, because experience and theory tell him that this is where gold occurs. This final section answers the question of **occurrence**: the geographical and geological distribution of mineral deposits, from the scale of the planet down to the mineral map of Cameroon.

6.1 Geological Controls on the Occurrence of Deposits

Observation. If you compare a world map of copper mines with a world map of gold mines, the two patterns do not coincide. Copper is concentrated along the western margins of the Americas and in central Africa; much of the world's gold comes from very old rocks in South Africa, Canada and Australia. This is not chance: each metal is tied to a particular **geological setting**.

DEFINITION Mineral occurrence and prospect

A **mineral occurrence** is any place where a useful mineral substance is present in anomalous concentration, whether or not it can be mined profitably. A **prospect** is an occurrence that has shown enough encouraging signs (good grades in samples, favourable geology) to justify detailed exploration. Only after full economic evaluation does a prospect become an **ore deposit** (a mineable reserve).

The occurrence of deposits is controlled by three great geological factors:

- **The type of crust and tectonic setting.** **Cratons** (old, stable continental nuclei such as the Congo craton) host Archaean gold, banded iron formations and diamonds. **Greenstone belts** within cratons concentrate gold and base metals. **Orogenic belts** (mountain chains) host hydrothermal veins and metamorphic deposits. **Rifts** and **sedimentary basins** host evaporites, phosphates, petroleum and stratabound copper–lead–zinc.
- **The age of the rocks.** Some deposits are linked to particular periods of Earth history: the great banded iron formations are essentially older than about 1.8 billion years, because their deposition required an ocean–atmosphere system still poor in free oxygen; Witwatersrand-type gold is late Archaean; most porphyry copper deposits are geologically young (Mesozoic to Cenozoic).
- **The tectonic and magmatic history.** A region that has experienced subduction, intrusion of granites, or rifting has been “prepared” by those events to receive the corresponding deposits: the heat, the fluids and the plumbing (faults, shear zones) are already in place.

PROPERTY Fundamental principle of metallogeny

The distribution of mineral deposits is controlled by **geology** (rock type, age, tectonic history), not by political boundaries. The copper belt crosses the Zambia–DRC frontier; the gold-bearing greenstones continue from Cameroon into Gabon and the Central African Republic. (Not examinable as a proof, but must be understood and illustrated with examples.)

6.2 Metallogenic Provinces and Metallogenic Epochs

Observation. Why does a single region, the Zambian–Congolese copper belt, supply a large share of the world’s copper and cobalt, while the Witwatersrand basin of South Africa has supplied a huge share of its gold? Because certain regions, endowed with the right rocks and the right history, concentrate certain metals.

DEFINITION Metallogenic province and epoch

A **metallogenic province** is a region within which mineral deposits of a particular type, or family of types, are unusually abundant because of a shared geological evolution. A **metallogenic epoch** is a period of geological time during which a particular family of deposits formed preferentially worldwide.

EXAMPLE Classic metallogenic provinces

- Copper belt of Zambia and the DRC: stratabound copper–cobalt in Neoproterozoic sedimentary rocks of the Katanga basin.
- Witwatersrand (South Africa): gold in late Archaean conglomerates (fossil placers), the richest gold province ever known.
- Banded iron formations: vast iron provinces in the Hamersley basin (Australia), the Transvaal (South Africa), Minas Gerais (Brazil) and Krivoy Rog (Ukraine), all of comparable great (Palaeoproterozoic) age — a fine example of a metallogenic epoch.




6.3 Plate Tectonics and Ore Deposits

Plate tectonics provides the unifying explanation: each tectonic setting creates the conditions for a characteristic family of deposits.

- **Subduction zones:** rising magmas release metal-rich fluids, forming **porphyry copper** (with Mo and Au) and epithermal gold veins, as along the Andes.
- **Mid-ocean ridges and ophiolites:** slices of oceanic lithosphere obducted onto continents

carry **chromite** (podiform deposits) and Cyprus-type massive sulphide copper.

- **Rift basins:** thick evaporite sequences (salt, potash), stratabound base metals, and the organic-rich sediments that generate **petroleum**.
- **Cratons and greenstone belts:** Archaean gold, banded iron, komatiite-hosted nickel, diamonds in kimberlite pipes.




Craton
Au, Fe (BIF), Ni

Plate-tectonic settings and their characteristic deposit types.

6.4 The Mineral Potential of Cameroon

Cameroon lies at the junction of three great geological domains: the **Congo craton** in the south (ancient, stable crust with greenstone belts), the **Pan-African mobile belts** in the centre and north (deformed and intruded about 600 million years ago), and the **Cameroon Volcanic Line** in the west (a much younger volcanic axis running from the Atlantic Ocean towards the Adamaoua plateau). Each domain carries its own mineral family, exactly as the principle of metallogeny predicts.



Schematic map of Cameroon: main mineral occurrences plotted on simplified geological domains.

Substance	Locality	Genetic type
Gold (Au)	East Region (Bétaré-Oya, Batouri)	Hydrothermal veins in greenstone belts; placers
Iron (Fe)	Mbalam, Nkout (South)	Banded iron formation (itabirite), enriched
Bauxite (Al)	Adamaoua plateau (Ngaoundal, Minim-Martap)	Tropical lateritic weathering
Nickel-cobalt	Nkamouna (East)	Lateritic weathering of ultramafic rocks
Rutile (TiO ₂)	Akonolinga (Centre)	Residual and alluvial concentration
Limestone	Figuil (North)	Sedimentary (marine carbonate)
Pozzolana	Volcanic line (West)	Volcanic scoria and ash (industrial mineral)
Petroleum	Rio del Rey, Logone basins	Sedimentary basin fill (fuel mineral)

DID YOU KNOW? Fuel minerals of Cameroon

Petroleum, exploited offshore in the **Rio del Rey** basin near Limbe and onshore in the **Logone** basin of the Far North, belongs to the family of **fuel minerals** (fossil energy resources), studied separately from metallic ores. It has long been the backbone of Cameroon's export earnings.

6.5 Reading a Mineral Occurrence Map

A mineral occurrence map plots symbols (each representing a substance) on a geological background. The skill tested at the GCE is to relate the symbols to the geology.

WARNING Do not read symbols in isolation

Never interpret a mineral symbol without relating it to the underlying geology. A gold symbol means little alone; a gold symbol inside a greenstone belt cut by granitic intrusions tells a coherent genetic story and points to prospective ground.

METHOD Interpreting a mineral occurrence map

1. Read the legend: identify each symbol and the substance it represents.

2. Study the geological background: rock types, ages, structures (faults, fold belts).
3. Note clusters of occurrences and ask which geological unit they follow.
4. Link each cluster to a genetic process (e.g. bauxite on a tropical plateau = lateritisation).
5. Identify **prospective zones**: areas with the same favourable geology but few recorded occurrences.

6.6 From Occurrence to Exploitation

METHOD Stages of mineral exploration

1. Regional survey: study of maps, satellite images and stream-sediment sampling over a large area.
2. Prospecting: fieldwork, trenching and pitting on anomalies; the occurrence becomes a prospect.
3. Geophysics and geochemistry: magnetic, electrical and soil-geochemical surveys to locate hidden ore.
4. Drilling: boreholes give samples at depth; grades and thicknesses are measured.
5. Reserve estimation and feasibility: tonnage and grade are computed; only then is the prospect declared an exploitable **deposit**.

WARNING An occurrence is not a deposit

A mineral showing, however spectacular, is not yet a mine. Economic evaluation — **grade**, **tonnage**, accessibility, market price, feasibility — must come first. Confusing “occurrence” with “reserve” is a classic GCE error.

Mining itself uses two great families of methods: **open-pit** mining for shallow, large or low-grade bodies (lateritic bauxite and nickel, iron of Mbalam) and **underground** mining for deep, high-grade bodies (vein gold). Finally, exploitation raises environmental and socio-economic stakes: uncontrolled **artisanal mining** (as in the gold fields of the East Region) degrades rivers and soils, while responsible projects plan **mine rehabilitation** (re-vegetation, backfilling) from the start.

WORKED EXAMPLE 5 Interpreting a prospecting report

A prospecting team reports: (i) gold values up to 4 g/t in quartz veins cutting a greenstone belt near Batouri; (ii) a bauxite cap averaging 45 % Al_2O_3 over a wide plateau near Ngaoundal. Classify each result (occurrence, prospect or reserve) and justify; state the genetic type of each.

SOLUTION

Solution.

- Batouri gold: encouraging grades in veins within a favourable greenstone setting, but no drilling or tonnage estimate is mentioned: it is a **prospect** (an occurrence under active exploration). Genetic type: hydrothermal (mesothermal) gold veins, typical of greenstone belts.
- Ngaoundal bauxite: high grade (45 % Al_2O_3) over a wide plateau suggests large tonnage, but without a full feasibility study it remains a well-advanced **prospect**, not yet a proven reserve. Genetic type: residual deposit from tropical lateritic weathering of aluminous rocks.
- Key point: neither result is yet a reserve, because reserves require demonstrated tonnage, grade and economic feasibility.

EXERCISE 4 Cameroon's mineral wealth

Name three Cameroonian mineral deposits, giving for each the substance, the locality and the genetic type. Then explain, in five lines, why the East Region is prospective for gold.

SOLUTION Exercise 1

Examples: gold at Bétaré-Oya (East) — hydrothermal veins in greenstone belts; iron at Mbalam (South) — enriched banded iron formation; bauxite at Minim-Martap (Adamaoua) — lateritic weathering; nickel-cobalt at Nkamouna — laterite on ultramafics; limestone at Figuil — marine sedimentary. The East Region is prospective for gold because it contains Archaean-to-Proterozoic greenstone belts cut by granitic intrusions and shear zones: the intrusions supplied heat and fluids, the shears channelled them, and gold precipitated in quartz veins, exactly the setting of the world's great gold provinces.

KEY POINTS Exam tip

Memorise at least three Cameroonians deposits with their substance, location and genetic type (e.g. Au-Bétaré-Oya-hydrothermal; Fe-Mbalam-BIF; Al-Adamaoua-lateritic; Ni-Co-Nkamouna-lateritic; limestone-Figuil-sedimentary). This is a classic GCE question. Remember also: deposits follow geology, not borders; an occurrence becomes a reserve only after economic evaluation; and always read a mineral symbol against the geological background.

7 INTEGRATION ACTIVITY

7.1 Aims of the activity

This integration activity closes the chapter by asking you to act as a **junior exploration geologist**. Working in teams, you will combine every notion studied in this chapter to answer one practical question: is a real iron deposit in southern Cameroon worth mining today? By the end of the activity you should be able to:

- recognise the **genetic type** and **morphology** of a deposit from map, cross-section and grade data;
- compute the **contained metal** in a deposit and the **reserve life** for a given production rate;
- distinguish **inferred** and **indicated resources** from a **proved reserve**, and state what additional work is needed to move from one category to the next;
- apply the notion of **economic mineral** by weighing grade, world market price, mining cost, technology and infrastructure;
- defend a reasoned recommendation orally, including one **environmental safeguard**, before a mock “Ministry of Mines” panel.

WARNING Skills being assessed

This activity mobilises all six sections of the chapter: the notion of economic mineral, resources and reserves, classification of deposits, magmatic/hydrothermal and sedimentary/weathering genesis, and the occurrence of deposits in Cameroon. Marks are awarded for correct reasoning, not for a single “right” verdict: two teams may reach opposite conclusions and both score highly if their arguments are rigorous. Every numerical answer must carry its unit, and every conclusion must be tied to a named notion from the course.

7.2 Situation

Dossier GEO05–NK: The Nkout Iron Project (South Region, Cameroon)

You are employed by a small exploration company, Mvila Minerals SARL. The company holds an exploration permit near Nkout, in the South Region, about 290 km inland from the Atlantic coast (Kribi). The Ministry of Mines, Industry and Technological Development has asked your company for an independent technical note before deciding whether to encourage a full feasibility study. Your team must write that note.

The dossier in your hands contains the following simplified data.

1. Geological setting. The permit lies within the **Congo craton**, on Archaean to Palaeoproterozoic basement (gneisses and greenstone belts, older than about 2.5 Ga). The geological map shows a ridge of **banded iron formation** (BIF) — alternating millimetric bands of quartz and iron oxides — traceable for about 8 km along strike. Where the BIF crops out at the surface, deep

tropical weathering has removed the silica and oxidised the iron minerals, producing a soft, reddish, enriched cap of **hematite** and **goethite** (a supergene enrichment developed on the BIF protore). Fresh BIF at depth contains only about 30 % Fe; the enriched cap contains 55–60 % Fe.

2. Morphology and dimensions (from 40 drill holes). The enriched orebody forms a **blanket** (a sub-horizontal stratabound layer) capping the ridge:


- average thickness: 40 m;
- lateral extent: 8000 m (strike) × 500 m (width);
- average dry density of the ore: $\rho = 3.2 \text{ t m}^{-3}$;
- average grade of the enriched cap: 45 % Fe (drill-hole grades range from 30 % Fe at the edges to 60 % Fe at the hilltop).

3. Drilling pattern. On the central 4 km of the ridge, holes are spaced 200 m apart (close enough to correlate the ore layer confidently from hole to hole). On the outer 4 km, only widely spaced holes ($\geq 800 \text{ m}$) exist, so continuity is assumed but not demonstrated. Roughly half of the total tonnage lies in each zone.

4. Economic data.

- world price of iron ore (62 % Fe benchmark): about 100 US dollars per tonne of ore;
- estimated mining and processing cost (open pit, crushing and upgrading to a saleable concentrate): 45 US dollars per tonne of ore;
- there is **no railway**; the ore would have to be trucked 290 km to Kribi, at an estimated transport cost of 35 US dollars per tonne;
- planned production rate if mined: 5 million tonnes of ore per year;
- the deep-water port of Kribi is operational and can handle bulk carriers.

5. Environmental and social notes. The ridge lies in the Congo Basin rainforest, near the Dja Faunal Reserve (a UNESCO World Heritage site). A river crossing the permit supplies water to three villages downstream.



Schematic cross-section of the Nkout ridge: a supergene-enriched hematite-goethite blanket (orange) caps the banded iron formation (blue), which rests in Archaean basement. Solid drill holes: close spacing in the central indicated zone; dashed holes: wide spacing in the inferred zones.

7.3 Tasks

Answer the following questions in order. Show all calculations and justify every conclusion with a notion from the chapter.

- Genesis and classification.** Using the dossier, identify the genetic type of the Nkout deposit. Name (a) the original sedimentary process that formed the BIF, and (b) the later process that raised the grade from 30 % to 45–60 % Fe. Classify the deposit by its morphology and state whether it is syngenetic or epigenetic with respect to the host BIF.
- Tonnage and contained metal.** Calculate the total tonnage of enriched ore, then the total tonnage of contained iron metal.
- Resource classification.** Using the drilling pattern, divide the tonnage between indicated resource and inferred resource. Explain, in two or three sentences, what additional work would be required before any of this material could be declared a proved reserve.
- Reserve life.** Assuming the whole tonnage were mined at 5 Mt of ore per year, calculate the life of the mine.
- Economic mineral?** Compute the total cost per tonne of ore delivered to the ship (mining + processing + transport) and compare it with the world price. Is the ore an economic mineral under present conditions? Discuss how the answer could change with (a) the world iron price, (b) construction of a railway, and (c) cut-off grade policy.
- Recommendation.** Write a one-page note to the Ministry panel: your verdict (mine now / keep exploring / wait), the two strongest arguments for it, and **one concrete environmental safeguard** your company would impose.
- Oral defence.** Present your note in five minutes and answer the panel's questions. Be ready to defend your resource classification and your economic threshold.

METHOD How to attack a tonnage–grade problem

1. Compute the volume of the orebody from its geometry: $V = \text{length} \times \text{width} \times \text{thickness}$.
2. Convert volume to tonnage using the density: $M_{\text{ore}} = V \times \rho$. Never skip this step.
3. Compute the contained metal: $M_{\text{metal}} = M_{\text{ore}} \times \text{grade}$ (grade as a decimal fraction).
4. Classify the tonnage by confidence (drilling density) before calling anything a reserve.
5. Only then discuss economics: compare total delivered cost with the sale price.

7.4 Model answer

1. Genesis and classification. The deposit is an **enriched banded iron formation** (BIF-type iron deposit). (a) The original BIF is a chemical sedimentary deposit: iron and silica were pre-precipitated from seawater in alternating bands on an Archaean–Palaeoproterozoic sea floor, so the primary iron concentration is of **sedimentary (chemical precipitation)** origin. (b) The high-grade cap results from **supergene enrichment by tropical weathering**: lateritic weathering dissolved and leached the quartz bands (residual concentration) and oxidised the iron minerals to hematite and goethite, roughly doubling the grade. Morphologically the orebody is a stratabound blanket (a horizontal layer) capping the ridge. The primary BIF is **syngenetic** (formed at the same time as the enclosing sediments); the enriched cap is **epigenetic** with respect to the BIF, since the enrichment post-dates the rock.

2. Tonnage and contained metal.

$$\begin{aligned}
 V &= 8000 \times 500 \times 40 = 1.6 \times 10^8 \text{ m}^3 \\
 M_{\text{ore}} &= V \times \rho = 1.6 \times 10^8 \times 3.2 = 5.12 \times 10^8 \text{ t} \approx 512 \text{ Mt} \\
 M_{\text{Fe}} &= 0.45 \times 512 \approx 230 \text{ Mt of contained iron}
 \end{aligned}$$

3. Resource classification. The central 4 km, drilled at 200 m spacing where continuity of the layer between holes is reasonably demonstrated, constitutes an **indicated resource**: about half the tonnage, i.e. $\approx 256 \text{ Mt}$. The outer 4 km, where continuity is only assumed from widely spaced holes, is an **inferred resource**: $\approx 256 \text{ Mt}$. To declare a **proved reserve**, the company would need: (i) closer-spaced drilling (e.g. 50–100 m) to demonstrate continuity in three dimensions; (ii) metallurgical tests proving the ore can be processed at the assumed cost; (iii) a full feasibility study including mining plan, infrastructure, market and environmental studies — because a reserve is the part of a resource that is economically mineable, not merely present.

Zone	Drilling	Category	Tonnage
Central 4 km	200 m spacing, continuity demonstrated	Indicated resource	$\approx 256 \text{ Mt}$
Outer 4 km	$\geq 800 \text{ m}$ spacing, continuity assumed	Inferred resource	$\approx 256 \text{ Mt}$
Whole ridge	after feasibility study only	(Potential) proved reserve	to be confirmed

4. Reserve life.

$$\text{Life} = \frac{512 \text{ Mt}}{5 \text{ Mt yr}^{-1}} \approx 102 \text{ years.}$$

Even using only the indicated tonnage (256 Mt), the life exceeds 50 years — comfortably long for a mining project.

5. Economic mineral? Total delivered cost per tonne of ore:

$$45 + 35 = 80 \text{ USD/t} \quad \text{against a sale price of 100 USD/t.}$$

The gross margin is only about 20 USD/t, and this ignores capital costs (building the mine, the plant and the road), taxes and the fact that a 45 % Fe ore must first be upgraded to near 62 % Fe to fetch the benchmark price. The deposit is therefore marginally economic at best under present conditions: it is a **resource**, but not yet clearly an **economic mineral**. (a) If the world price rose above roughly 120–130 USD/t, the margin would become comfortable. (b) A railway (as long discussed for the Mbalam–Nabeba–Kribi corridor) could cut transport from 35 to perhaps 10 USD/t, transforming the economics. (c) Raising the **cut-off grade** to mine only the 55–60 % Fe hilltop material would reduce tonnage but increase margin per tonne.

6. Recommendation (model). Verdict: keep exploring — do not mine yet. Arguments: (1) the tonnage and grade are real and large, but half of it is only an inferred resource and the economic margin (≈ 20 USD/t before capital costs) is too thin to justify investment without a railway or a higher iron price; (2) the indicated 256 Mt alone justifies further drilling and metallurgical testing to convert resources into proved reserves. Environmental safeguard: impose a protected buffer zone of at least 100 m of untouched forest along the river, with continuous water-quality monitoring downstream, so that the three villages' water supply is never contaminated by runoff from the open pit and waste dumps.

WARNING Marking pointers

Frequent errors: forgetting the density when converting volume to tonnage; calling the whole 512 Mt a “reserve” (it is a resource — no feasibility study exists); confusing the syngenetic BIF with the epigenetic enrichment; and concluding “economic” from grade alone while ignoring transport and infrastructure. A verdict of “wait” or “mine” is acceptable — only the quality of the argument is marked.

DID YOU KNOW? Iron in Cameroon: a real national stake

The Nkout deposit is not a fiction: it belongs to a belt of BIF-hosted iron occurrences (Mbalam, Nkout, Djadom and others) along the northern edge of the Congo craton in the South and East Regions of Cameroon. Their development has been discussed for decades, and the missing link has always been the same one you identified in this activity: a railway to the deep-water port of Kribi. This is a perfect illustration of the chapter's central message — a mineral deposit becomes an economic mineral only when geology, technology, infrastructure and market price all align.

8 METHODS AND WORKED EXAMPLES

9 ECONOMIC GEOLOGY: ECONOMIC MINERALS

9.1 Skill 1: Distinguishing ore, gangue, grade and economic mineral

DEFINITION Economic mineral, ore mineral, gangue, grade

An **economic mineral** is a mineral from which a valuable metal or substance can be extracted profitably under current economic, technological and legal conditions. The **ore mineral** is the specific mineral species that carries the metal of value (e.g. cassiterite SnO_2 for tin, chalcopyrite CuFeS_2 for copper). The **gangue** is the assemblage of valueless minerals (quartz, calcite, feldspars, micas...) that are extracted together with the ore mineral and must be separated during processing. The **grade** (or tenor) expresses the concentration of the valuable substance in the rock, usually in percent (%), grams per tonne (g/t) or parts per million (ppm); $1 \text{ g/t} = 1 \text{ ppm}$.

METHOD Identifying the components of an ore and computing a grade

When faced with a description of a mined rock, proceed in order:

1. Identify the useful mineral(s): the mineral species that contains the metal or substance of value (e.g. cassiterite SnO_2 for tin).
2. Identify the gangue: the valueless minerals extracted with the ore (quartz, calcite, feldspar...) which must be removed during processing.
3. Decide whether the mineral is “economic”: ask the three questions — is it concentrated enough (grade), is it present in sufficient quantity (tonnage), and can it be extracted profitably with existing technology and market prices?
4. Compute the grade if data are given:

$$\text{grade (\%)} = \frac{\text{mass of valuable substance}}{\text{total mass of rock}} \times 100$$

5. Convert between mineral grade and metal grade using the chemical formula and molar masses when required.

Reflexes: always state the unit of grade (% , g/t, ppm); remember that $1 \text{ g/t} = 1 \text{ ppm}$; an “economic mineral” is defined by economics, not only by chemistry — the same mineral may be ore in one place and waste in another.

WARNING Common mistakes in grade calculations

- Forgetting to convert % to a decimal fraction before multiplying by tonnage (e.g. using 5 instead of 0.05).
- Confusing mineral grade (e.g. % SnO_2) with metal grade (e.g. % Sn) — always specify which you are quoting.
- Using the wrong molar mass or forgetting that the metal fraction in the mineral is $M_{\text{metal}}/M_{\text{mineral}}$.
- Omitting units in the final answer (grade must carry %, g/t or ppm).

WORKED EXAMPLE 6 Cassiterite ore from the Mayo Darlé tin field

A prospector in the Mayo Darlé area (Adamawa Region, Cameroon) collects a 500 kg sample of mineralised granite. After concentration, the sample yields 8 kg of cassiterite (SnO_2).

- Define the terms ore mineral and gangue, and identify them in this sample.
- Calculate the grade of the rock in cassiterite.
- Given $M(\text{Sn}) = 118.7 \text{ g/mol}$ and $M(\text{O}) = 16.0 \text{ g/mol}$, calculate the grade of the rock in metallic tin.

SOLUTION

a) An **ore mineral** is a mineral from which a metal or useful substance can be extracted profitably. The **gangue** is the assemblage of valueless minerals (here quartz, feldspars and micas of the granite) extracted together with the ore mineral and discarded during processing. In this sample the ore mineral is cassiterite SnO_2 ; the gangue is the granitic material (quartz, feldspar, mica).

b) Grade in cassiterite:

$$\text{grade}_{\text{SnO}_2} = \frac{8 \text{ kg}}{500 \text{ kg}} \times 100 = 1.6 \%$$

The rock contains 1.6 % cassiterite, i.e. 16 kg of SnO_2 per tonne of rock.

c) Molar mass of cassiterite:

$$M(\text{SnO}_2) = 118.7 + 2 \times 16.0 = 150.7 \text{ g/mol}$$

Mass fraction of tin in cassiterite:

$$w_{\text{Sn}} = \frac{118.7}{150.7} = 0.7876 \approx 78.8 \%$$

Mass of tin in the sample:

$$m_{\text{Sn}} = 8 \text{ kg} \times 0.7876 = 6.30 \text{ kg}$$

Grade in metallic tin:

$$\text{grade}_{\text{Sn}} = \frac{6.30 \text{ kg}}{500 \text{ kg}} \times 100 = 1.26 \%$$

Conclusion: the rock grades 1.6 % SnO₂, equivalent to about 1.26 % Sn. Since typical economic tin ores grade around 0.5–1 % Sn, this sample is of economic interest.

KEY POINTS Key points: ore, gangue, grade

- Economic mineral = profitable extraction (geology + economics + technology).
- Ore mineral = carrier of the metal; gangue = waste minerals.
- Grade = concentration; units: %, g/t, ppm (1 g/t = 1 ppm).
- Metal grade = mineral grade × (metal molar mass / mineral molar mass).

9.2 Skill 2: Classifying a deposit as resource or reserve

DEFINITION Mineral resource and mineral reserve (CRIRSCO/JORC framework)

A **mineral resource** is a concentration of solid material of economic interest in or on the Earth's crust in such form, grade and quantity that there are reasonable prospects for eventual economic extraction. It is subdivided by geological confidence into Inferred, Indicated and Measured resources (increasing confidence). A **mineral reserve** is the economically mineable part of a Measured and/or Indicated Mineral Resource. It includes diluting materials and allowances for losses, and is subdivided into Probable and Proved reserves. All reserves are resources, but not all resources are reserves. The boundary is drawn by economics and technology, not by geology alone.

METHOD Applying the resource/reserve terminology

1. Read the level of geological knowledge: is the deposit only inferred (few data, wide spacing of samples), indicated (reasonably well sampled) or measured (densely sampled, well known)?
2. Ask the economic question: can the material be extracted legally and profitably at present with existing technology?
3. Place the material:

- **Resource:** concentration of economic interest, known or inferred, regardless of present profitability. Subdivided into inferred, indicated, measured resources by increasing confidence.
 - **Reserve:** the part of a measured or indicated resource that is economically mineable now. Subdivided into probable and proved reserves.
4. **Watch for changes of status:** a resource becomes a reserve if the metal price rises, technology improves, or infrastructure arrives; a reserve can revert to a resource if prices collapse.

Key idea to remember: All reserves are resources, but not all resources are reserves. The boundary is drawn by economics and technology, not by geology alone.

WARNING Resource vs Reserve — examiner's traps

- Never call a deposit a “reserve” just because it is large or well known. Profitability now is mandatory.
- “Measured resource” ≠ “Proved reserve”. A measured resource becomes a proved reserve only after a feasibility study demonstrates economic viability.
- Inferred resources can never be converted directly to reserves; they must first become indicated/measured.

WORKED EXAMPLE 7 The bauxite of Minim–Martap and Ngaoundal

The bauxite plateaux of Minim–Martap and Ngaoundal (Adamawa, Cameroon) are estimated to contain very large tonnages of aluminium-rich laterite. However, the deposits lie far from the coast, the alumina grade is moderate, and no railway currently serves the plateaux, so no mining has started.

- Define mineral resource and mineral reserve.
- Should the Minim–Martap bauxite be reported as a reserve? Justify.
- State two changes that could convert this material into a reserve.

SOLUTION

a) A **mineral resource** is a natural concentration of solid material of economic interest in or on the Earth's crust, whose grade and tonnage give reasonable prospects of eventual economic extraction; it is classified as inferred, indicated or measured according to geological confidence. A **mineral reserve** is the portion of a measured or indicated resource

that is economically, legally and technically mineable at the time of reporting; it is classified as probable or proved.

b) No. Although the tonnage is enormous and the geology is reasonably well known (so it is at least an indicated resource), the deposit cannot presently be mined profitably: transport costs to a port would exceed the value of the ore, and the moderate grade increases processing costs. Therefore it fails the economic test and must be reported as a resource, not a reserve.

c) Any two of:

- construction of a railway or improved road linking the plateaux to the deep-water port (e.g. Kribi), cutting transport costs;
- a sustained rise in the world price of aluminium/alumina;
- improved, cheaper beneficiation technology raising the recoverable grade;
- construction of a local alumina refinery or cheap energy source (e.g. hydroelectricity from the Lom Pangar dam area) reducing processing costs.

Examiner's remark: this example shows that "reserve" is a dynamic, economic category: the geology of Minim-Martap does not change, but its status can.

KEY POINTS Resource / Reserve hierarchy

Category	Geological confidence	Economic viability
Inferred Resource	Low	Not required
Indicated Resource	Moderate	Not required
Measured Resource	High	Not required
Probable Reserve	High (Indicated)	Demonstrated
Proved Reserve	Very high (Measured)	Demonstrated

9.3 Skill 3: Computing tonnage, contained metal and cut-off grade

METHOD Deposit evaluation calculations

1. Compute the volume of the ore body from its geometry (thickness $\times$ surface area for a tabular body; $\pi r^2 h$ for a cylinder; etc.).
2. Compute the tonnage: $T = V \times \rho$ where ρ is the bulk density of the ore (in t/m³).
3. Compute contained metal: $M = T \times g$ where g is the grade expressed as a decimal frac-

tion (for g/t grades, M in grams comes out directly: $M(g) = T(t) \times g(g/t)$).

4. Apply a cut-off grade: the **cut-off grade** is the minimum grade at which a block of rock pays for its extraction and processing. Material below cut-off is waste. Compare each block's grade with the cut-off before summing reserves.
5. Check profitability with a simple balance: value of recovered metal = contained metal $\times$ recovery $\times$ price; compare with extraction + processing costs.

Reflexes: convert % to decimals before multiplying; keep units consistent ($m^3 \times t/m^3 = t$); never confuse contained metal with recoverable metal (multiply by the recovery factor).

WARNING Cut-off grade and recovery

- The cut-off grade is not a geological constant — it depends on costs, metal price, recovery and dilution.
- “Contained metal” is the total metal in the ground; “Recoverable metal” = contained $\times$ metallurgical recovery (always $< 100\%$).
- In reserve estimation, only blocks above cut-off are summed; blocks below cut-off are waste even if they contain some metal.

WORKED EXAMPLE 8 Evaluating a small gold vein

A quartz vein near Bétaré Oya (East Region, Cameroon) is 200 m long, 2 m wide on average and is mined to a depth of 50 m. The bulk density of the ore is 2.7 t/m^3 and the average grade is 5 g/t Au. The cut-off grade is 2 g/t.

- a) Calculate the tonnage of ore in the vein.
- b) Calculate the mass of contained gold.
- c) Is the vein above cut-off grade? Comment.
- d) With a recovery of 90 % and a gold price of 60 000 FCFA/g, calculate the gross value of the recoverable gold.

SOLUTION

- a) Volume of the vein (tabular approximation):

$$V = \text{length} \times \text{width} \times \text{depth} = 200 \times 2 \times 50 = 20\,000 \text{ m}^3$$

Tonnage:

$$T = V \times \rho = 20\,000 \text{ m}^3 \times 2.7 \text{ t/m}^3 = 54\,000 \text{ t}$$

b) Contained gold (grade in g/t × tonnage in t gives grams):

$$M_{\text{Au}} = 54\,000 \text{ t} \times 5 \text{ g/t} = 270\,000 \text{ g} = 270 \text{ kg of gold}$$

c) The average grade (5 g/t) is above the cut-off grade (2 g/t): $5 > 2$. The whole vein therefore qualifies as ore and can be counted in the reserve, provided mining costs are covered.

d) Recoverable gold (metallurgical recovery = 90%):

$$M_{\text{rec}} = 270\,000 \text{ g} \times 0.90 = 243\,000 \text{ g}$$

Gross value:

$$V_{\text{gross}} = 243\,000 \text{ g} \times 60\,000 \text{ FCFA/g} = 1.458 \times 10^{10} \text{ FCFA} \approx 14.6 \text{ billion FCFA}$$

Conclusion: the vein contains 270 kg Au, of which 243 kg is recoverable, with a gross value of about 14.6 billion FCFA. Whether it is a reserve still depends on mining and processing costs being lower than this value.

EXERCISE 5 Tonnage and grade practice

A stratiform zinc–lead deposit in the Benue Trough has a horizontal extent of $1.2 \text{ km} \times 0.8 \text{ km}$ and an average thickness of 4 m. The bulk density is 3.2 t/m^3 . The average grade is 6% Zn and 2% Pb. The cut-off grade is 3% Zn equivalent.

1. Calculate the tonnage of the deposit.
2. Calculate the contained zinc and lead (in tonnes).
3. If the metallurgical recovery is 85% for Zn and 80% for Pb, and prices are 2500 FCFA/kg Zn and 2000 FCFA/kg Pb, estimate the gross in-situ value of the recoverable metals.

9.4 Skill 4: Classifying a mineral deposit from its description

DEFINITION Morphological and genetic classification

Mineral deposits are classified by morphology (form, geometry, spatial relationship to host rocks) and by genesis (the geological process that concentrated the metal). Examiners expect candidates to state both classifications and justify each with observed features.

METHOD Using the classification schemes

1. Extract the diagnostic facts from the statement: host rock, geometry of the body, mineral assemblage, associated igneous/sedimentary/metamorphic features.
2. Apply the morphological (form) classification: is the body stratiform/stratabound (layer-parallel), a vein/lode (cross-cutting fissure filling), a stockwork (network of veinlets), a dissemination, a pipe, or a placer (surficial accumulation)?
3. Apply the genetic classification: which process concentrated the metal?
 - Magmatic: crystallisation, segregation, immiscible liquids (e.g. Cr, Ni, Pt in ultramafics; Sn, W, rare metals in pegmatites).
 - Hydrothermal: hot aqueous fluids depositing veins, replacements, stockworks (e.g. Au, Cu, Zn, Pb, Sn, W).
 - Sedimentary: placers (mechanical), chemical precipitates (BIF, evaporites), residual (laterites).
 - Weathering/Residual: lateritic bauxite, nickel laterite, supergene enrichment.
 - Metamorphic: skarns, recrystallised deposits, metamorphosed massive sulphides.
4. Cross-check with the metal association: e.g. Cr–Ni–Pt point to ultramafic magmatic segregation; Sn–W–Mo point to granitic hydrothermal systems; Fe–Al laterites point to tropical weathering.
5. State both classifications in your answer: examiners reward candidates who give form and genesis.

KEY POINTS Common morphological terms

- Stratiform: layer-like, parallel to bedding/banding.
- Stratabound: confined to a specific stratigraphic unit but not necessarily layered.
- Vein / Lode: tabular, discordant, fracture-filling.
- Stockwork: network of small veinlets.
- Disseminated: fine-grained ore minerals scattered through the rock.
- Pipe / Chimney: vertical, cylindrical body.
- Placer: unconsolidated, surficial, hydraulically sorted.

WORKED EXAMPLE 9 Classifying four deposits

Classify each of the following deposits morphologically and genetically, justifying each answer:

- a) Layers of chromite interbedded with peridotite at the base of a large ultramafic intrusion.
- b) Gold-bearing quartz veins cutting across granitic rocks at Bétaré Oya.
- c) The bauxite capping the lateritic plateaux of Minim-Martap.
- d) Cassiterite and gold grains concentrated in the gravels of a river bed in the Kadey valley.

SOLUTION

a) Morphology: stratiform (the chromite forms layers parallel to the igneous layering). Genesis: magmatic segregation — early-crystallising dense chromite crystals settled and accumulated at the base of the crystallising ultramafic magma (gravitational settling during fractional crystallisation). Typical association: Cr with ultramafic rocks.

b) Morphology: veins/lodes (cross-cutting fissure fillings). Genesis: hydrothermal — hot aqueous fluids released during the late stages of granitic magmatism circulated through fractures; cooling and pressure drop caused quartz and gold to precipitate in the fissures. This is a mesothermal/epithermal-type Au-quartz vein system.

c) Morphology: blanket/stratiform residual cap (lateritic profile). Genesis: residual weathering (lateritisation) — intense tropical weathering of aluminous parent rocks leached silica and alkalis, concentrating insoluble aluminium hydroxides (gibbsite, boehmite) near the surface. This is a lateritic bauxite deposit.

d) Morphology: placer (surficial, unconsolidated accumulation in river gravels). Genesis: sedimentary/mechanical concentration (alluvial placer) — weathering liberated dense, chemically resistant minerals (cassiterite, gold) from primary sources; running water sorted and concentrated them by density in traps of the river bed.

Synthesis: the four deposits illustrate the four great genetic families — magmatic, hydrothermal, residual, and sedimentary (placer) — and show that the same metal (e.g. cassiterite) can occur both as a primary hydrothermal deposit and as a secondary placer.

9.5 Skill 5: Explaining ore-forming processes (magmatic and hydrothermal)**METHOD** Building a genetic explanation for magmatic/hydrothermal deposits

1. State the source of the metal (magma, hydrothermal fluid, country rock leached by fluids).

2. State the transport medium (silicate melt, immiscible sulphide liquid, hot aqueous fluid).
3. State the concentration mechanism:
 - Magmatic: crystal settling (Cr, Pt), immiscible sulphide liquid segregation (Ni–Cu), late-stage residual melt enrichment (rare metals in pegmatites, greisens).
 - Hydrothermal: dissolution at depth, transport as complexes in hot aqueous fluids, then precipitation caused by cooling, pressure drop, boiling, fluid mixing or reaction with wall rock.
4. State the trap/host: layered intrusion, fracture/vein, porous or reactive (e.g. carbonate) wall rock, contact zone (skarn).
5. Link each observation to the process in a cause–effect chain; examiners penalise bare lists of facts.

Key ideas: hydrothermal fluids are mostly magmatic or heated meteoric water; deposits are zoned (high-temperature minerals near the source, low-temperature ones farther away); porphyry and skarn deposits form around intrusive contacts.

EXPERIMENT**Magmatic sulphide segregation — a physical model**

Fill a transparent container with water (silicate melt analogue) and add a dense, immiscible liquid (e.g. glycerol with food colouring, or simply oil) to represent sulphide liquid. Shake to form droplets; observe them coalesce and sink to the bottom. This demonstrates gravitational accumulation of an immiscible dense liquid — the key step in magmatic Ni–Cu–PGE deposit formation.

WORKED EXAMPLE 10**From magma to a magmatic Ni–Cu sulphide deposit**

Describe and explain the sequence of processes by which a nickel–copper sulphide deposit can form from a mafic magma. Illustrate your answer with a labelled cross-section.

SOLUTION


Step 1 — Source and melting. A mantle-derived mafic/ultramafic magma rises into the crust. It carries Fe, Ni, Cu and sulphur dissolved in the silicate melt.

Step 2 — Sulphur saturation and immiscibility. As the magma cools (or assimilates sulphur from crustal rocks), it becomes saturated in sulphur. A dense immiscible sulphide liquid separates from the silicate melt, like oil separating from water. Because Ni, Cu and the platinum-group elements have a strong chemical affinity for sulphide (they are chalcophile), they partition preferentially into this sulphide liquid, scavenging the metals from

a large volume of magma (partition coefficients $D_{\text{Ni}}^{\text{sul/sil}} \sim 100\text{--}1000$).

Step 3 — Gravitational accumulation. The dense sulphide droplets coalesce and sink, collecting at the base of the magma chamber or in structural traps (embayments, feeder conduits, troughs).

Step 4 — Crystallisation. On final cooling the sulphide liquid crystallises as pyrrhotite (FeS), pentlandite ($(\text{Fe}, \text{Ni})_9\text{S}_8$) and chalcopyrite (CuFeS_2), forming a massive or disseminated sulphide ore body at the base of the intrusion.



Magmatic segregation: dense immiscible sulphide droplets sink and accumulate at the base of the magma chamber.

Conclusion: the deposit results from (i) sulphur saturation, (ii) immiscible sulphide liquid segregation which scavenges chalcophile metals, and (iii) gravitational settling — a classic magmatic segregation process.

DID YOU KNOW? Cameroon context: magmatic deposits

The ultramafic complexes of the Pan-African belt in southern Cameroon (e.g. Nkamouna, Lomié) host Ni-Co-Mn laterites developed on serpentinised peridotites. The primary magmatic sulphide mineralisation (if any) has been largely overprinted by tropical weathering, but the ultramafic protolith is the essential source. The Cameroon Volcanic Line itself is not a major magmatic sulphide province, but its alkaline rocks host rare-metal mineralisation (e.g. Nb, REE at Lomié, Akonolinga).

9.6 Skill 6: Explaining sedimentary, residual and supergene enrichment processes

METHOD Explaining surface and near-surface ore formation

1. Identify the climate and landscape context: tropical humid climate $\Rightarrow$ intense chem-

ical weathering, laterites, bauxite, nickel laterite; arid basins $\Rightarrow$ evaporites; rivers and coasts $\Rightarrow$ placers.

2. For residual deposits: describe leaching of soluble components (SiO_2 , Ca, Na, K, Mg) and residual concentration of insoluble ones (Al, Fe, Ti, Ni, Cr) in the weathering profile. Mention the lateritic profile: ferricrete / laterite duricrust $\rightarrow$ mottled zone $\rightarrow$ pallid zone $\rightarrow$ saprolite $\rightarrow$ bedrock.
3. For placers: apply the three conditions — (i) a primary source, (ii) liberation by weathering, (iii) hydraulic sorting concentrating dense and resistant minerals in traps (bedrock irregularities, inside bends, base of gravels, riffles).
4. For supergene enrichment: describe the vertical zonation — leached cap (gossan) at the top, supergene enriched zone below the water table where descending copper re-precipitates as chalcocite (Cu_2S), covellite (CuS), native copper; primary (hypogene) ore at depth (chalcopyrite, bornite). Explain that enrichment can upgrade a sub-economic protore into an ore.
5. For chemical sedimentary ores: evaporation (evaporites: halite, gypsum, potash) or chemical/biochemical precipitation (banded iron formations — BIF).

WARNING Placer vs residual — don't confuse them

- Residual (lateritic): forms in situ by chemical weathering; the ore stays where the parent rock was. No transport.
- Placer: forms by mechanical transport and hydraulic sorting; the ore moves from its source. Requires density contrast and durability.
- Supergene enrichment occurs below the water table in the saturated zone; the leached cap is above it.

WORKED EXAMPLE 11 Why is a placer rich only in certain minerals?

Alluvial workings in the East Region of Cameroon recover gold and cassiterite, but never feldspar or calcite, even though these are abundant in the source rocks.


- a) State the two physical-chemical properties a mineral must possess to form a placer deposit, and name the property that controls hydraulic sorting.
- b) Explain why feldspar and calcite are absent from the placers.
- c) Draw a labelled sketch of a river section showing where placer gold accumulates.

SOLUTION

a) A placer mineral must be (i) dense (high specific gravity) so that running water can separate it from lighter grains, and (ii) chemically resistant and mechanically durable so that it survives weathering and transport. Hydraulic sorting is controlled essentially by density (together with grain size and shape): when water velocity drops, the densest grains settle first.

b) Feldspar and calcite fail both tests: they have low densities (about $2.6\text{--}2.7\text{ g/cm}^3$, close to quartz), so they are not separated from ordinary sand; and they are chemically unstable — feldspar alters to clay minerals (kaolinite) during tropical weathering, and calcite simply dissolves in slightly acidic surface waters. They are therefore destroyed or dispersed long before they could be concentrated. Gold (19.3 g/cm^3 , inert) and cassiterite ($\approx 7.0\text{ g/cm}^3$, resistant) survive and are concentrated.

c)



Conclusion: placers are natural density-concentration plants operated by running water; only dense, resistant minerals such as gold, cassiterite, diamond or platinum can form them.

EXPERIMENT**Supergene enrichment profile — sketch and label**

Draw a vertical cross-section through a porphyry copper deposit showing: (1) leached cap (gossan) with iron oxides, (2) supergene enrichment zone (chalcocite blanket) at/near water table, (3) primary hypogene zone (chalcopyrite + pyrite) at depth. Label the water table and indicate the direction of metal migration (downward for Cu, upward for Fe). Explain why the chalcocite zone is richer than the primary ore.

9.7 Skill 7: Interpreting a geological cross-section to locate ore bodies

METHOD Reading occurrence geometry on a map or section

1. Identify the host structures: fractures and faults (veins), intrusive contacts (skarns, greisens), stratigraphic layers (stratiform ores), unconformities, palaeo-river channels (placers).
2. Note the spatial relationship of ore to geology: concordant (parallel to bedding $\Rightarrow$ syngenetic, formed with the rock) or discordant (cross-cutting $\Rightarrow$ epigenetic, formed later).
3. Use metal zoning: high-temperature metals (Sn, W, Mo, Bi) near the intrusion; intermediate (Cu, Zn, Pb, Fe) farther out; low-temperature (Ag, Hg, Sb, Au) at the periphery. Predict where each metal should be sought.
4. Reason on depth and continuity: a vein pinches and swells; a stratiform body is laterally continuous; a placer follows the channel.
5. Conclude with an exploration recommendation: where to drill or dig next, with justification.

DEFINITION Syngenetic vs Epigenetic

A **syngenetic** deposit forms at the same time as its host rock (e.g. a sedimentary iron bed, a magmatic chromite layer); it is usually concordant. An **epigenetic** deposit forms later than its host rock, introduced into pre-existing rocks (veins, replacements, skarns, placers); it is usually discordant.

WORKED EXAMPLE 12 Where should the exploration company drill?

The cross-section below shows a granite intruding schists and limestones. A tin-bearing greisen occurs at the granite roof (point A); skarn with copper occurs at the limestone contact (point B); quartz veins with silver occur in fractures far from the contact (point C).

- a) Define epigenetic and syngenetic deposits and state which applies here.
- b) Explain the observed metal zoning $\text{Sn} \rightarrow \text{Cu} \rightarrow \text{Ag}$ away from the granite.
- c) The company wants more tin. Where should it explore, and why?

SOLUTION

a) A **syngenetic** deposit forms at the same time as its host rock (e.g. a sedimentary iron bed); it is usually concordant. An **epigenetic** deposit forms later than its host rock, in-


roduced into pre-existing rocks (veins, replacements); it is usually discordant. Here all three occurrences (greisen, skarn, veins) were emplaced by fluids related to the granite after the schists and limestones existed: they are epigenetic.

b) The hydrothermal fluids were hottest at or within the granite and cooled as they migrated outward. Minerals precipitate in order of decreasing temperature of deposition:

- Sn (cassiterite) with tourmaline/topaz in the greisen: highest temperature (400–500°C), deposited at the granite roof (A);
- Cu (and Fe) in the skarn: high-to-intermediate temperature (300–400°C), deposited where the fluids reacted with the reactive limestone at the contact (B);
- Ag in distal quartz veins: lowest temperature (< 250°C), deposited far from the heat source (C).

This is the classic temperature-controlled hydrothermal zoning around an intrusion.

c) Tin is a high-temperature metal, so the company should explore close to and within the granite, especially along the roof and cupolas of the intrusion and along contacts with the schists (greisen-bordered veins), i.e. around point A and at depth beneath it — not in the distal fractures, which are the low-temperature silver zone. Drilling should target the granite margins and any unexposed apophyses (satellite intrusions) where greisenisation may have occurred.



Hydrothermal zoning around a granitic intrusion: Sn (greisen) at roof → Cu (skarn) at carbonate contact → Ag (veins) distally.

9.8 Skill 8: Writing a synoptic essay on Cameroon's mineral occurrences

METHOD Structuring a GCE essay on mineral occurrence

1. Define the key terms in one or two sentences (economic mineral, deposit, occurrence, resource, reserve).

2. Organise by genetic family or by commodity, never as a random list. A strong plan: magmatic/hydrothermal metals → residual deposits → sedimentary/placer deposits → industrial minerals and hydrocarbons.
3. Attach each commodity to a place and a process: e.g. bauxite – Minim-Martap/Ngaoundal – lateritisation; iron – Mbalam – BIF; gold – Bétaré Oya – hydrothermal veins and placers; tin – Mayo Darlé – granitic hydrothermal; rutile – Akonolinga; limestone – Figuil (cement); petroleum – Logone Birni and Rio del Rey basins.
4. Link occurrence to the geological framework: Congo craton (Fe, Au), Pan-African belts (Au, Sn, rutile), Cameroon Volcanic Line (volcanic rocks, industrial minerals, some REE), sedimentary basins (oil, gas, salt, limestone).
5. Conclude with the economic significance and the resource/reserve caution (many Cameroonian occurrences are resources awaiting infrastructure, investment or favourable prices).

KEY POINTS**Cameroon's geological domains and their mineral signatures**

Domain	Key commodities	Genetic process / Setting
Congo Craton (South)	Iron (Mbalan), Nk-out), Gold (primary & placer)	BIF (chemical sedimentary), shear-zone hydrothermal Au
Pan-African Belts (Centre, East, Adamawa)	Gold (Bé-taré, Oya, Colomine, Ba-touri), Tin (Mayo Darlé), Rutile (Akono-linga), Ni-Co laterites (Lomié, Nkamouna)	Hydrothermal veins (Au, Sn), granitic greisens (Sn), lateritic weathering on ultramafics (Ni-Co), placer marking
Cameroon Volcanic Line	Pozzolan, basalt, scoria, some REE/Nb	Alkaline volcanism, weathering
Sedimentary Basins (Logone Birni, Rio del Rey, Douala/Kribi-Campo)	Petroleum	Sedimentary

WORKED EXAMPLE 13 Guided essay: the mineral endowment of Cameroon

“Cameroon’s geology gives it a varied but unevenly exploited mineral endowment.” Discuss, with reference to the mode of occurrence and genesis of at least four mineral commodities.

SOLUTION

Introduction. An economic mineral is one that can be extracted at a profit; its occurrence is controlled by geology, its status by economics. Cameroon’s territory spans three great geological domains — the Congo craton in the south, the Pan-African mobile belts in the centre and north, and the Cameroon Volcanic Line crossing both — plus coastal and interior sedimentary basins. Each domain hosts a characteristic family of deposits.

1. Residual deposits of the cratonic south and Adamawa. Intense tropical weathering of aluminous rocks produced the great lateritic bauxite plateaux of Minim-Martap and Ngaoundal (Adamawa), among the largest known bauxite resources in Africa (several billion tonnes), and nickel-cobalt laterites near Lomié and Nkamoula. Genesis: leaching of silica and alkalis, residual enrichment of Al, Fe, Ni in a lateritic profile (duricrust → mottled → pallid → saprolite). Status: still essentially resources, for lack of a railway to the coast and moderate alumina grades.

2. Banded iron formations of the craton. The iron deposits of Mbalam and Nkout (South Region) are itabirite-type banded iron formations (BIF), chemical sediments of Palaeoproterozoic age, locally upgraded by metamorphism and supergene enrichment. Genesis: chemical/biochemical precipitation of iron in ancient oceans, later metamorphic recrystallisation and tropical weathering leaching silica. Status: Mbalam is a reserve (feasibility completed, rail-port infrastructure planned); Nkout remains a resource.

3. Hydrothermal and placer metals of the Pan-African belts. Gold occurs in quartz veins at Bétaré Oya, Colomine and Batouri (East Region), deposited from hydrothermal fluids in Pan-African shear zones, and in derived alluvial placers worked artisanally. Tin (cassiterite) at Mayo Darlé is associated with granitic magmatism (greisens and veins) and its placers. Rutile at Akonolinga occurs in hydrothermal veins and pegmatites related to Pan-African granites. Genesis: magmatic-hydrothermal concentration in shear zones and granitic cupolas, then mechanical re-concentration by rivers.

4. Sedimentary and industrial commodities. The Logone Birni (Chad Basin) and Rio del Rey / Douala / Kribi-Campo basins contain petroleum and natural gas trapped in structural and stratigraphic traps; limestone at Figuil (North) and Mfou (Centre) supports the cement industry; pozzolana and basalt from the Cameroon Volcanic Line are quarried for construction aggregates and cement additives.

Conclusion. Cameroon’s endowment is indeed varied — magmatic-hydrothermal metals (Au, Sn, W, rutile), residual bauxite and nickel laterites, sedimentary iron (BIF), hydrocarbons and industrial minerals — but unevenly exploited: geology has provided the occurrences, while economics (grade, infrastructure, prices, policy) decides which are reserves. The bauxite of Adamawa, a giant resource awaiting a railway, is the perfect illustration that an “economic mineral” is defined as much by economics as by geology.

EXERCISE 6 Essay practice

“The classification of a mineral occurrence as a resource or a reserve depends on factors beyond geology.” Discuss this statement with reference to at least two Cameroonian mineral occurrences.

SOLUTION Essay practice

Key points to include:

- Definitions: resource (geological confidence) vs reserve (economic viability now).
- Example 1: Minim-Martap bauxite — huge indicated/measured resource, but no reserve because transport costs > value; needs railway, price rise, or local refinery.
- Example 2: Mbalam iron — moved from resource to reserve after feasibility study, rail-port agreement (Kribi deep-water port), and offtake contracts.
- Example 3: Bétaré Oya gold — artisanal exploitation proves technical feasibility; reserve status depends on mechanised mining costs vs gold price.
- Conclusion: geology sets the stage (grade, tonnage, mineralogy), but economics, technology, infrastructure, law and politics draw the resource/reserve line.

10 EXERCISES

10.1 I check my knowledge

EXERCISE 7 MCQ: economic minerals and deposits

For each question, choose the single correct answer and justify your choice in one sentence.

1. An **ore** is:
 - (a) any rock containing a metallic element;
 - (b) a rock from which a useful substance can be extracted at a profit;
 - (c) the waste material surrounding a valuable mineral;
 - (d) a mineral deposit that has not yet been discovered.
2. The **grade** of a deposit is:
 - (a) the total tonnage of rock in the deposit;
 - (b) the concentration of the valuable substance in the ore;
 - (c) the depth of the deposit below the surface;
 - (d) the market price of the metal.
3. Which of the following is a syngenetic deposit?
 - (a) a gold-bearing quartz vein cutting granite;
 - (b) a bedded ironstone formed at the same time as the enclosing sediments;
 - (c) a skarn developed at a granite–limestone contact;
 - (d) a hydrothermal copper vein.
4. The Minim–Martap deposit of the Adamawa Plateau is mainly a deposit of:
 - (a) gold;
 - (b) iron;
 - (c) bauxite;
 - (d) limestone.

EXERCISE 8 True or false?

State whether each assertion is true or false, and justify your answer in one or two sentences.

1. A mineral resource automatically becomes a reserve as soon as it is discovered.
2. A rise in the world price of a metal can convert a sub-economic resource into a reserve without any new discovery.
3. Hydrothermal deposits are always older than the rocks that enclose them.
4. Residual (lateritic) concentration of aluminium is favoured by a hot, humid tropical climate such as that of the Adamawa Plateau.

EXERCISE 9 Key definitions

Define each of the following terms in one or two precise sentences, giving one example where appropriate:

1. **economic mineral**;
2. **gangue**;
3. **cut-off grade**;
4. **epigenetic deposit**;
5. **placer deposit**.

EXERCISE 10 Direct calculation: contained metal

A copper deposit contains 5 million tonnes of ore grading 0.8 % Cu.

1. Calculate the mass of copper metal contained in the deposit.
2. The mine produces 12 000 tonnes of copper metal per year. Calculate the life of the reserve, assuming total recovery.
3. State one reason why the real mine life may be shorter than your calculated value.

10.2 I practise

EXERCISE 11 Resource or reserve?


The table below summarises exploration results for three deposits A, B and C.

Deposit	Geological knowledge	Economic study	Estimated tonnage
A	Few boreholes; continuity assumed between widely spaced data points	No feasibility study	50 Mt
B	Dense drilling grid; continuity well established	Positive feasibility study at current prices	120 Mt
C	Moderate drilling; continuity reasonably assumed	Preliminary study only	80 Mt

1. Classify each deposit as an **inferred resource**, an **indicated resource** or a **proved reserve**, justifying each classification.
2. Explain what additional work would be needed to upgrade deposit C to reserve status.

EXERCISE 12 Interpreting a weathering profile

The diagram below is a schematic lateritic weathering profile developed on syenite at Minim-Martap (Adamawa, Cameroon).



Schematic lateritic profile, Minim-Martap.

1. Name the horizons labelled (a) to (d), from the surface downwards.
2. In which horizon is bauxite concentrated? Name the main aluminium minerals it con-

tains.

3. Explain, in three or four sentences, how intense chemical weathering removes silica and alkalis while concentrating aluminium.
4. State two features of Cameroon's climate that favour this process.

EXERCISE 13 Cross-cutting relations and genetic classification

A cross-section shows a granite intruding schists. Quartz vein V1, carrying chalcopyrite, cuts the schists but is itself cut and offset by quartz vein V2, which carries gold. Both veins stop against a younger dolerite dyke.

1. Draw a simple labelled sketch cross-section of these relationships.
2. Arrange the granite, schists, veins V1 and V2, and the dyke in order from oldest to youngest, justifying each step with the principle of cross-cutting relationships.
3. Classify the copper and gold mineralisations as **syngenetic** or **epigenetic**, and as **magmatic**, **hydrothermal**, **sedimentary** or **residual**, giving reasons.

EXERCISE 14 From resource to reserve

An iron deposit near Mbalam (East Region) is currently classified as a sub-economic resource.

1. Distinguish clearly between a **resource** and a **reserve**.
2. Explain two distinct ways in which this resource could become a reserve without any new discovery (think of prices, technology, infrastructure).
3. Suggest one reason why a deposit may remain unexploited even when it is technically a reserve.

10.3 I prepare for the GCE

EXERCISE 15 Structured question: ore, grade and reserves (20 marks)

1. Define the following terms, giving one example of each: ore, gangue, grade, cut-off grade. (8 marks)
2. A copper deposit contains 2 million tonnes of ore grading 1.2 % Cu. Calculate the mass

of copper metal contained in the deposit. (3 marks)

3. The mine produces 20 000 tonnes of copper metal per year. Calculate the life of the reserve, stating any assumption you make. (3 marks)
4. Explain three factors that determine whether a given mineral concentration can be worked as an ore. (6 marks)

EXERCISE 16 Map exercise: mineral occurrences of Cameroon (20 marks)

You are provided with an outline map of Cameroon showing the regions and the main towns.

1. On the map, place and label five major mineral occurrences: gold (East Region), bauxite (Minim–Martap), iron (Mbalam), nickel–cobalt (Lomie area), limestone (Figuil). (5 marks)
2. For each occurrence, state the genetic type of the deposit (magmatic, hydrothermal, sedimentary, residual/weathering or metamorphic) and justify your answer in one sentence. (10 marks)
3. Choose one region that you consider prospective for gold exploration. Justify your choice using two geological criteria (nature of the basement rocks, known occurrences, structural features). (5 marks)

EXERCISE 17 Essay: geology controls mineral distribution (20 marks)

“The distribution of mineral deposits is controlled by geology, not by politics.”

Discuss this statement with reference to:

1. at least three Cameroonian deposits (for example the bauxite of Minim–Martap, the iron of Mbalam, the gold of the East Region, the nickel–cobalt of Lomie, the limestone of Figuil), explaining in each case the geological process that localised the mineralisation;
2. at least one world metallogenic province (for example the copper belt of Central Africa or the gold provinces of cratonic shields);
3. the extent to which political boundaries, infrastructure and economic decisions nevertheless influence whether a deposit is actually mined.

Your essay should include an introduction, a developed argument with labelled diagrams where useful, and a conclusion. (20 marks)

11 SOLUTIONS TO THE EXERCISES

SOLUTION Exercise 1 — MCQ: economic minerals and deposits

1. Answer: (b). An ore is defined by economic extractability: a rock is only an ore if the valuable substance it contains can be extracted and sold at a profit under current technical and economic conditions. A rock merely containing a metal (a) is not necessarily an ore; (c) defines gangue; (d) describes an undiscovered resource.
2. Answer: (b). The grade is the concentration of the valuable substance in the ore, expressed as a percentage (e.g. % Cu) or in g/t for precious metals. Tonnage (a), depth (c) and price (d) are separate parameters of a deposit.
3. Answer: (b). A syngenetic deposit forms at the same time as the enclosing rock: a bedded ironstone precipitated during sedimentation is contemporaneous with its host sediments. Veins (a, d) and skarns (c) are introduced after the host rock and are therefore epigenetic.
4. Answer: (c). Minim–Martap on the Adamawa Plateau is Cameroon’s major bauxite (aluminium ore) deposit, formed by intense lateritic weathering of nepheline syenite under a tropical climate.

SOLUTION Exercise 2 — True or false?

1. False. A resource is a concentration whose existence and approximate size are known or inferred; it becomes a reserve only after detailed exploration (dense drilling) and a feasibility study proving that extraction is technically possible and economically profitable. Discovery alone is not sufficient.
2. True. Reserves are defined partly by economics: if the metal price rises, the cut-off grade falls and previously sub-economic material becomes profitable to mine, so tonnages can be reclassified from resource to reserve with no new discovery.
3. False. Hydrothermal fluids circulate through pre-existing rocks and deposit minerals in fractures and pores, so hydrothermal deposits are epigenetic — younger than their host rocks (principle of cross-cutting relationships).
4. True. Hot temperatures accelerate chemical reactions and abundant rainfall provides the percolating water that leaches silica and alkalis; this intense lateritic weathering concentrates residual aluminium hydroxides (bauxite), as at Minim–Martap on the Adamawa Plateau.

SOLUTION Exercise 3 — Key definitions

1. Economic mineral: any naturally occurring mineral substance (metallic or non-metallic) whose concentration in the Earth's crust is sufficient for it to be extracted and used at a profit; e.g. bauxite at Minim-Martap, gold in the East Region.
2. Gangue: the valueless minerals or rock material intimately mixed with the ore minerals in a deposit, which must be separated and discarded during processing; e.g. quartz accompanying gold in a vein.
3. Cut-off grade: the minimum grade above which material is economically worth mining and processing; material below the cut-off grade is treated as waste. It depends on metal price, mining cost and recovery.
4. Epigenetic deposit: a deposit formed after the enclosing rock, introduced into it by fluids or melts; e.g. a gold-bearing quartz vein cutting schists.
5. Placer deposit: a deposit formed by the mechanical concentration of dense, chemically resistant minerals (gold, diamond, cassiterite) in river gravels or beach sands after weathering, transport and sorting; e.g. alluvial gold in the rivers of the East Region of Cameroon.

SOLUTION Exercise 4 — Direct calculation: contained metal

1. Mass of copper = tonnage of ore × grade:

$$m_{\text{Cu}} = 5 \times 10^6 \text{ t} \times \frac{0.8}{100} = 5 \times 10^6 \times 0.008 = 40\,000 \text{ t Cu.}$$

The deposit contains 40 000 tonnes of copper metal.

2. Life of the reserve:

$$L = \frac{\text{contained metal}}{\text{annual production}} = \frac{40\,000}{12\,000} \approx 3.3 \text{ years.}$$

3. The real mine life may be shorter because recovery is never total: during mining and processing a fraction of the metal is lost (dilution by waste, metallurgical losses), so the recoverable metal is less than the contained metal. (Also acceptable: part of the ore may prove inaccessible or below cut-off grade.)

SOLUTION Exercise 5 — Resource or reserve?


1. Deposit A — inferred resource. Geological knowledge is poor (few boreholes, continuity only assumed between widely spaced points) and no economic study exists; tonnage and grade can only be estimated with low confidence.
2. Deposit B — proved reserve. The dense drilling grid establishes continuity with high confidence (measured category) and a positive feasibility study demonstrates that extraction is technically feasible and economically viable at current prices.
3. Deposit C — indicated resource. Moderate drilling gives reasonable confidence in continuity (better than inferred, less than measured), but only a preliminary economic study exists, so reserve status cannot yet be claimed.
4. To upgrade C to a reserve one must: (i) carry out additional drilling (infill drilling) to raise geological confidence to the measured category; (ii) perform metallurgical tests to establish recovery; (iii) complete a full feasibility study covering mining method, costs, infrastructure, market and environmental impact, demonstrating profitability.

SOLUTION Exercise 6 — Interpreting a weathering profile

1. From the surface downwards: (a) ferruginous crust / iron cap (cuirasse), rich in iron oxides; (b) bauxitic horizon, rich in aluminium hydroxides; (c) saprolite (altered, clay-rich, friable rock retaining the parent structure); (d) fresh parent rock (syenite).
2. Bauxite is concentrated in horizon (b), the bauxitic horizon. Its main aluminium minerals are the hydroxides gibbsite $\text{Al}(\text{OH})_3$ (dominant), with subordinate boehmite and diaspore ($\text{AlO}(\text{OH})$).
3. Under a hot, humid climate, rainwater charged with dissolved CO_2 percolates through the rock and hydrolyses the silicate minerals (feldspars) of the syenite. Alkalis and alkaline earths (Na^+ , K^+ , Ca_2^+) are removed in solution, and silica is leached away as soluble species. Aluminium and iron, being almost insoluble at near-neutral pH, remain behind and accumulate as residual hydroxides, so the residue is progressively enriched in Al — this is bauxitisation.
4. Favourable features of Cameroon's climate: (i) high temperatures all year round, which accelerate chemical weathering reactions; (ii) abundant, well-distributed rainfall (humid tropical climate of the Adamawa Plateau), which ensures continuous leaching and removal of soluble products.

SOLUTION Exercise 7 — Cross-cutting relations and genetic classification

1. Sketch cross-section:



Cross-section: granite in schists, veins V1 and V2, dolerite dyke.

2. Order from oldest to youngest, using the principle of cross-cutting relationships (a geological feature is younger than any feature it cuts):
 - (a) Schists — oldest, they host everything else;
 - (b) Granite — intrudes the schists, so it is younger;
 - (c) Vein V1 (chalcopyrite) — cuts the schists, so post-dates them;
 - (d) Vein V2 (gold) — cuts and offsets V1, so it is younger than V1;
 - (e) Dolerite dyke — cuts and terminates both veins, so it is the youngest.
3. Both mineralisations are epigenetic: they were introduced into pre-existing rocks after those rocks formed (the veins cut the schists). Genetically they are hydrothermal deposits: hot aqueous fluids circulated through fractures and precipitated quartz, chalcopyrite (Cu) and gold as the fluids cooled. They are neither syngenetic (not contemporaneous with the host), nor sedimentary/residual (no surface concentration process involved).

SOLUTION Exercise 8 — From resource to reserve

1. A resource is a concentration of a useful substance in the crust whose location, approximate tonnage and grade are known or reasonably inferred, but whose profitable extraction has not been demonstrated. A reserve is the portion of a resource that has been proved by detailed exploration and shown by a feasibility study to be extractable economically and technically under present conditions. Every reserve is a resource, but not every resource is a reserve.
2. Two ways the Mbalam resource could become a reserve without new discovery:
 - (a) Economic change: a sustained rise in the world price of iron ore lowers the effective cut-off grade and makes the deposit profitable; equally, a fall in costs (energy, reagents, labour) has the same effect.

- (b) Technical/infrastructural change: construction of a railway and a deep-water port (e.g. towards Kribi) drastically reduces transport costs, and improved mining/processing technology raises recovery — both can turn a sub-economic resource into a viable reserve.
3. A deposit may remain unexploited even as a reserve because of non-geological obstacles: lack of investment capital, political or administrative instability, land-use or environmental conflicts, or absence of the infrastructure needed to bring the product to market.

SOLUTION Exercise 9 — Structured question: ore, grade and reserves (20 marks)

(a) Definitions (8 marks — 2 per term: 1 for the definition, 1 for a valid example).

- Ore: a rock or aggregate of minerals from which a useful substance can be extracted at a profit; e.g. bauxite as the ore of aluminium.
- Gangue: the valueless minerals associated with the ore minerals, separated and discarded during processing; e.g. quartz in a gold vein.
- Grade: the concentration of the valuable substance in the ore, in % or g/t; e.g. a copper ore grading 1.2 % Cu.
- Cut-off grade: the minimum grade at which material can be mined and processed profitably; e.g. material below 0.4 % Cu sent to waste in a given mine.

(b) Contained metal (3 marks).

$$m_{\text{Cu}} = 2 \times 10^6 \text{ t} \times \frac{1.2}{100} = 24\,000 \text{ t Cu.}$$

Marking: 1 mark formula, 1 mark correct substitution, 1 mark answer with unit. (c) Life of the reserve (3 marks).

$$L = \frac{24\,000}{20\,000} = 1.2 \text{ years.}$$

Assumption stated: total recovery (all contained metal is mined and processed without loss). Marking: 1 mark method, 1 mark result, 1 mark the assumption. (d) Three factors (6 marks — 2 per factor, any three).

- Grade and tonnage: the concentration must be high enough and the volume large enough to repay investment.
- Market price and demand: the value of the metal must exceed the total cost of extraction and processing.
- Technology and cost of extraction: mining method (open pit vs underground), depth, metallurgical recoverability of the mineral.
- Infrastructure and location: roads, railway, port, energy and water supply (e.g. Mbalam iron awaiting a railway).

- Environmental, legal and political conditions: permits, stability, environmental costs.

Examiner expectation: precise vocabulary (ore, grade, cut-off grade), correct units in calculations, and factors clearly explained rather than listed.

SOLUTION Exercise 10 — Map exercise: mineral occurrences of Cameroon (20 marks)

(a) Placement (5 marks — 1 per correctly located and labelled occurrence).

- Gold — East Region: in the Batouri–Bétaré Oya–Kette area, eastern Cameroon, on the Archean–Proterozoic basement.
- Bauxite — Minim–Martap: on the Adamawa Plateau, north of Ngaoundéré.
- Iron — Mbalam: East Region, near the Congo/Gabon border, south-east of Batouri.
- Nickel–cobalt — Lomie: East Region, east of Yokadouma area, in the forest zone.
- Limestone — Figuil: North Region, in the Bénoué trough, south of Garoua.

(b) Genetic types (10 marks — 2 per occurrence: 1 for the type, 1 for the justification).

Occurrence	Genetic type	Justification
Gold (East Region)	Hydrothermal (or placer)	Gold occurs in quartz veins cutting basement rocks (epigenetic hydrothermal), reworked into alluvial placers in rivers.
Bauxite (Minim–Martap)	Residual / weathering	Intense lateritic weathering of syenite leached silica and alkalis, concentrating residual Al hydroxides.
Iron (Mbalam)	Sedimentary (BIF), enriched by weathering	Banded iron formations deposited with ancient sediments, then enriched by leaching of silica.
Ni–Co (Lomie)	Residual / weathering (lateritic)	Deep weathering of ultrabasic rocks concentrates Ni and Co in the lateritic profile.
Limestone (Figuil)	Sedimentary	Marine chemical/biochemical precipitation of CaCO_3 in the Cretaceous Bénoué trough.

(c) Prospective region (5 marks). The East Region is the most prospective for gold: (i) it is underlain by Archean to Proterozoic greenstone and schist belts of the Congo craton margin, a rock association known worldwide to host gold; (ii) known occurrences (Batouri, Bétaré Oya) and active artisanal workings prove the presence of a mineralised system; (iii) regional faults and shear zones provided the channels for gold-bearing hydrothermal fluids. Marking: 1 mark a coherent choice, 2 marks per justified geological criterion.

SOLUTION Exercise 11 — Essay: geology controls mineral distribution (20 marks)

Indicative mark scheme: introduction and definition of terms (3); three Cameroonian deposits with correct genetic process ($3 \times 3 = 9$); one world metallogenic province correctly explained (3); balanced discussion of political/economic factors (3); organisation, clarity, labelled diagram (2).

Model answer (outline).

Introduction. Mineral deposits are not evenly distributed: they occur where specific geological processes have concentrated useful elements. The statement is therefore largely true at the scale of where deposits exist, but must be qualified at the scale of whether they are mined.

Development — Cameroonian examples.

- Bauxite of Minim–Martap (Adamawa): residual deposit; intense lateritic weathering of syenite under a hot humid climate leached silica and alkalis and concentrated Al hydroxides on a stable, well-drained plateau surface.
- Iron of Mbalam (East Region): sedimentary banded iron formations of the Precambrian craton, enriched by tropical weathering which removed silica and raised the iron grade.
- Gold of the East Region: hydrothermal quartz veins localised by shear zones in Archean–Proterozoic greenstone/schist belts, reworked into placers.
- Nickel–cobalt of Lomie: lateritic concentration over ultrabasic bedrock.
- Limestone of Figuil: marine sedimentation in the Cretaceous Bénoué trough.

In every case the localising factor is geological: the parent rock, the climate-driven weathering, the sedimentary basin or the structural channelway — none of which respects administrative boundaries.

World province. The Central African Copper Belt (DRC–Zambia) illustrates this: stratiform copper mineralisation is confined to specific sedimentary horizons of the Katanga Supergroup, so the ore runs straight across the political border, following the geology. Similarly, gold provinces of the Kaapvaal, Canadian and West African cratons reflect the distribution of ancient greenstone belts.

Nuance. Geology fixes the location of deposits, but politics and economics decide their exploitation: a reserve requires investment, stable legislation, and infrastructure — the

Mbalam iron, though geologically proved, long awaited a railway and port; conversely the Figuil limestone is worked because a cement plant and market exist. Prices, technology and political decisions convert resources into reserves and mines.

Conclusion. The distribution of mineral deposits is indeed written in the rocks and controlled by geology; but the map of producing mines is drawn jointly by geology, economics and political choice.

Examiner expectations: precise genetic vocabulary (syngenetic/epigenetic, residual, hydrothermal, BIF), correctly located examples, a clear two-sided argument, and a structured essay with introduction and conclusion.

12 SUMMARY SHEET



Mind map of the chapter "Economic Minerals": six branches linking definitions, resource classification, deposit types, genetic processes and Cameroonian examples.

KEY POINTS Summary Sheet 1 — Key Definitions

- **Mineral**: a naturally occurring, inorganic solid with a definite chemical composition and an ordered internal (crystalline) structure.
- **Economic mineral**: any mineral (or rock) that can be extracted and sold at a profit under current technical and economic conditions. Profitability, not mere presence, is the deciding criterion.
- **Ore**: a rock containing one or more valuable minerals in sufficient concentration to be mined profitably. The **ore mineral** is the valuable mineral; the **gangue** is the worthless associated material (e.g. quartz, calcite) that must be separated during processing.
- **Grade (tenor)**: the concentration of the useful substance in the ore, expressed in % for common metals (e.g. 55 % Fe) or in g/t (equivalent to ppm) for precious metals (e.g. 2 g/t Au).
- **Cut-off grade**: the minimum grade below which exploitation is not profitable. It is not fixed: it varies with the world price of the metal, the available technology, energy and transport costs, and the location of the deposit.
- Scale of knowledge: **mineral occurrence** (anomalous concentration noted) → **prospect** (showing investigated, encouraging results) → **deposit** (body of ore of demonstrated economic interest).

KEY POINTS Summary Sheet 2 — Resources and Reserves

- **Resource**: the total quantity of a commodity known or suspected to exist in the ground, whether or not it is currently economic to extract.
- **Reserve**: the part of the resource that is economically extractable now with existing technology and at current prices.
- Increasing geological confidence (how well the body is known): inferred → indicated → measured resource.
- Increasing economic confidence (how sure the extraction is): probable → proven reserve.
- Reserves are dynamic: they grow when metal prices rise or technology improves (former resources become reserves), and shrink when prices fall or costs increase.
- **Reserve life** = reserves ÷ annual production; it estimates how many years a mine (or a country) can continue producing at the present rate.

KEY POINTS Summary Sheet 3 — Classification of Mineral Deposits

- **Syngenetic** deposits: formed at the same time as the host rock (chromite layers in basic intrusions, placers, banded iron formations, evaporites). They are typically concordant (parallel to bedding or layering).
- **Epigenetic** deposits: formed after the host rock, by introduction of material into it (hydrothermal veins, skarns, replacement bodies). They are typically discordant (cross-cutting).
- Genetic classes to master: magmatic, hydrothermal, sedimentary, residual/weathering, metamorphic.

KEY POINTS Summary Sheet 4 — Formation Processes to Know

- Magmatic: early crystallisation and settling of dense crystals → layered chromite and magnetite; liquid immiscibility → Ni–Cu sulphide ores; concentration of rare elements in late residual fluids → **pegmatites** (Sn, W, Li, gemstones).
- Hydrothermal: hot aqueous fluids circulating through rocks deposit dissolved minerals on cooling, especially in fractures → **veins** (Au, Ag, Pb, Zn); reaction of fluids with limestone → **skarn** (W, Fe, Cu); large low-grade disseminated bodies → **porphyry copper**.
- Sedimentary: mechanical sorting by water or wind → **placers** (gold, diamond, rutile, cassiterite); chemical precipitation → **banded iron formations** (BIF) and **evaporites** (halite, gypsum, potash); biochemical accumulation → phosphates, coal, petroleum.
- Weathering/residual: intense tropical leaching removes silica and bases and concentrates aluminium → **bauxite** (lateritisation); downward-percolating waters can upgrade sulphide ores → **supergene enrichment** of copper.
- Metamorphic: recrystallisation under heat and pressure → marble, graphite, garnet, kyanite; contact metasomatism around intrusions → skarn ores.

KEY POINTS Summary Sheet 5 — Occurrence: Cameroon to Remember

- **Bauxite**: Minim–Martap and Ngaoundal (Adamawa Plateau; residual, lateritic deposits on the plateau surface).
- **Iron**: Mbalam–Nabeba (South–East; itabirite/BIF–type ore at the margin of the Congo craton).

- **Nickel–cobalt:** Lomié (East Region; lateritic deposit developed by deep tropical weathering of ultrabasic rocks).
- **Gold:** Batouri–Bétaré–Oya (East Region; alluvial placers and primary lode occurrences).
- **Rutile:** Akonolinga (Centre); **limestone:** Figuil (North; cement manufacture); **oil and gas:** Rio del Rey and Douala sedimentary basins; **cassiterite:** Mayo–Darlé (North–West).

METHOD Summary Sheet 6 — Working Methods

To judge whether a body is an economic deposit:

1. Identify the commodity and its use.
2. Estimate the tonnage and multiply by the grade to obtain the contained metal.
3. Compare the grade with the cut-off grade under current prices and technology.
4. Conclude: uneconomic concentration (occurrence/resource) or exploitable ore (reserve).

To classify a deposit:

1. Ask: “Is the ore the same age as the host rock?” Yes → syngenetic; no → epigenetic.
2. Read the textures and geological setting (layering, veins, alteration, sedimentary structures).
3. Assign the genetic process (magmatic, hydrothermal, sedimentary, residual, metamorphic) and justify it with one observation.

WARNING Summary Sheet 7 — Common Mistakes

- Confusing mineral, ore mineral and ore: an ore is a rock (a mixture of ore minerals and gangue), not a single mineral.
- Stating that reserves are fixed quantities: they depend on price, technology and costs, and therefore change with time.
- Calling every deposit syngenetic: hydrothermal veins and skarns are epigenetic because they post-date their host rocks.
- Expressing the grade of gold in %: precious metals are quoted in g/t (ppm).
- Writing “bauxite mineral”: bauxite is a rock (a mixture of aluminium hydroxides such as

gibbsite), used as the ore of aluminium.

KEY POINTS Summary Sheet 8 — GCE Examination Tips

- Define each term precisely, then give one Cameroonian example per process — examiners reward local illustration (e.g. lateritisation → Minim–Martap bauxite).
- Be ready to draw quick labelled sketches: a hydrothermal vein cross-cutting sedimentary strata; a laterite profile (topsoil – bauxite/ferricrete – saprolite – fresh bedrock); a layered intrusion with a chromite seam.
- In resource/reserve questions, always structure the answer along the two axes: geological confidence (inferred–indicated–measured) and economic viability (resource–probable–proven reserve).
- Show the reasoning in calculations: contained metal = tonnage × grade; reserve life = reserves ÷ annual production; always carry the units.

**Signed • OnBuch+**ONBUCH+ TEACHING TEAM
FOLLOWS THE OFFICIAL MINESEC SYLLABUS**OnBuch+**