

Explaining Spatial Profiles of Line Emission in the Horsehead Nebula Using Cloud Surface Curvature

Student

Ducheng Lu

Supervisors

Franck Le Petit (LERMA)
Emeric Bron (LERMA)

Jan 2025

Photodissociation Regions (PDRs)

Interstellar medium (ISM): gas and dust between stars in galaxies

Credit: JWST

- $\sim 10\%$ of the total baryonic mass
- main site of star formation


PDR: regions of **neutral gas** in the ISM where **far-ultraviolet radiation dominates the chemical and heating processes**

- diagnostic of the ISM
- stellar feedback

Heating: photoelectric effect,
cosmic rays


Cooling: line emission

infrared and line emission
 \Rightarrow physical conditions in PDRs



Credit: Tielens 2005

The Horsehead Nebula




Credit: Hernández-Vera et al. 2023

- observed **edge-on** \Rightarrow observational access to the chemical stratification


part of a collaboration between JPL, Paris Observatory, IRAM, and CSIC Madrid to study the presence of water in the Horsehead Nebula

Data



observations along a cut through the Horsehead Nebula



- C⁺, C, CO and its isotopologues, and H₂O



Motivation

1D PDR models

- infinite and uniform in two dimensions, depth-dependent only
 ⇒ allows for a detailed study of the physical and chemical processes
- cannot be compared directly to edge-on observations





Motivation

1D PDR models

- infinite and uniform in two dimensions, depth-dependent only
⇒ allows for a detailed study of the physical and chemical processes
- cannot be compared directly to edge-on observations

Proposed Solution:

- approximate edge-on regions with **curvature radius**.
- a spherical geometry wrapper for the MeudonPDR code




Motivation

1D PDR models

- infinite and uniform in two dimensions, depth-dependent only
⇒ allows for a detailed study of the physical and chemical processes
- cannot be compared directly to edge-on observations

Proposed Solution:


- approximate edge-on regions with **curvature radius**.
- a spherical geometry wrapper for the MeudonPDR code
 - spatial profiles of **column densities**
 - solve **radiative transfer** for line intensities
 - **convolution** with the instrument resolution



The MeudonPDR Code

stationary 1D PDR code

- radiative transfer • chemical balance
- level populations • thermal balance




Cloud size ($A_{V,\text{max}}$)	40
Proton density (n_H)	$3 \times 10^4 - 3 \times 10^6 \text{ cm}^{-3}$
Pressure (P)	$1 \times 10^6 - 1 \times 10^7 \text{ K cm}^{-3}$
ISRF	shape: Mathis, geometry: beam_isot
ISRF scaling factor	$G_0^{\text{obs}} = 100, G_0^{\text{back}} = 0.04$
UV radiative transfer method	FGK approximation, or exact H ₂ self- and mutual shielding
Turbulent velocity dispersion	2 km s^{-1}
Extinction Curve	HD38087
$R_V = A_V/E(B-V)$	5.50
$C_D = N_H/E(B-V)$	1.57×10^{22}

⁰Parameter values are based on Maillard, 2023

The MeudonPDR Code

stationary 1D PDR code

- radiative transfer • chemical balance
- level populations • thermal balance



Cloud size ($A_{V,\text{max}}$)	40
Proton density (n_H)	$5 \times 10^6 \text{ K cm}^{-3}$
Pressure (P)	shape: Mathis, geometry: beam_isot
ISRF	$G_0^{\text{obs}} = 100$, $G_0^{\text{back}} = 0.04$
ISRF scaling factor	exact H ₂ self- and mutual shielding
UV radiative transfer method	
Turbulent velocity dispersion	2 km s^{-1}
Extinction Curve	HD38087
$R_V = A_V/E(B - V)$	5.50
$C_D = N_H/E(B - V)$	1.57×10^{22}

+ surface chemistry

⁰Parameter values are based on Maillard, 2023

From Slab to Spherical Geometry

Input:


- level number density $n_X(d)$
- cloud radius R (free parameter)
- LoS impact parameter b

Algorithm:

- interpolation of $n_X(d) = f(d)$ to allow computation of n_X at any depth
- $d = R - \sqrt{s^2 + b^2} \Rightarrow n_X(s)$
- integrate along the LoS

$$N_X(b) = 2 \int_0^{s_{\max}} n_X(s') ds'$$

For optically thin lines, $I_\nu \propto N_X$



Convolution with the Instrument Resolution

Algorithm

- interpolate the model grids to uniform ones `x_uniform`, `y_uniform`

Convolution with the Instrument Resolution

Algorithm

- interpolate the model grids to uniform ones `x_uniform`, `y_uniform`
- Gaussian kernel with the given resolution, full width half maximum (FWHM)

$$\sigma = \text{FWHM} / (2\sqrt{2 \ln 2}), \quad g(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{x^2}{2\sigma^2}\right)$$

Convolution with the Instrument Resolution

Algorithm

- interpolate the model grids to uniform ones `x_uniform`, `y_uniform`
- Gaussian kernel with the given resolution, full width half maximum (FWHM)

$$\sigma = \text{FWHM} / (2\sqrt{2 \ln 2}), \quad g(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{x^2}{2\sigma^2}\right)$$


Convolution with the Instrument Resolution

Algorithm

- interpolate the model grids to uniform ones `x_uniform`, `y_uniform`
- Gaussian kernel with the given resolution, full width half maximum (FWHM)


$$\sigma = \text{FWHM} / (2\sqrt{2 \ln 2}), \quad g(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{x^2}{2\sigma^2}\right)$$

- convolution with truncation at 3σ , padding `y_uniform` with 0




Convolved Column Densities from Spherical Models

step 1: slab vs spherical geometry




Convolved Column Densities from Spherical Models

spherical geometry: the peaks shift to deeper locations within the cloud




Convolved Column Densities from Spherical Models

spherical geometry: the peaks shift to deeper locations within the cloud




step 2: convolved vs unconvolved column densities




Convolved Column Densities from Spherical Models

spherical geometry: the peaks shift to deeper locations within the cloud




convolution: line spatial profiles are smoothed and further extended




Compare Column Densities with Observations

convolved column densities from **spherical models** match observations better



Compare Column Densities with Observations


convolved column densities from **spherical models** match observations better



profile width ✓, shape on the front side ✓, shape on the back side ✗






Compare Column Densities with Observations

convolved column densities from **spherical models** match observations better



profile width ✓, shape on the front side ✓, shape on the back side ✗






Can cloud radius make a difference?

$N(C+ \text{El}=2P, J=3/2)$  $N(C \text{ El}=3P, J=1)$  $N(\text{CO } v=0, J=2)$  $N(\text{CO } v=0, J=4)$  $N(\text{H}_2\text{O } J=1, ka=1, kc=0)$ 

$d_{\text{PDR}} = 50.98''$

- PDR layer
- $R_{\text{cloud}} = 0.5 \text{ pc}$
- $R_{\text{cloud}} = 1.0 \text{ pc}$
- $R_{\text{cloud}} = 3.0 \text{ pc}$
- $R_{\text{cloud}} = 5.0 \text{ pc}$
- observation








$N(C+ \text{El}=2P, J=3/2)$  $N(C \text{ El}=3P, J=1)$  $N(\text{CO } v=0, J=2)$  $N(\text{CO } v=0, J=4)$  $N(\text{H}_2\text{O } J=1, ka=1, kc=0)$ 

- $d_{\text{PDR}} = 50.98"$
- PDR layer
- $R_{\text{cloud}} = 0.5 \text{ pc}$
- $R_{\text{cloud}} = 1.0 \text{ pc}$
- $R_{\text{cloud}} = 3.0 \text{ pc}$
- $R_{\text{cloud}} = 5.0 \text{ pc}$
- observation








The differences in column density profiles are trivial

$N(C+ \text{EI}=2P, J=3/2)$  $N(C \text{ EI}=3P, J=1)$  $N(\text{CO } v=0, J=2)$  $N(\text{CO } v=0, J=4)$  $N(\text{H}_2\text{O } J=1, ka=1, kc=0)$ 

- $d_{\text{PDR}} = 50.98''$
- PDR layer
 - $R_{\text{cloud}}=0.5 \text{ pc}$
 - $R_{\text{cloud}}=1.0 \text{ pc}$
 - $R_{\text{cloud}}=3.0 \text{ pc}$
 - $R_{\text{cloud}}=5.0 \text{ pc}$
 - observation

The differences in column density profiles are trivial

Tail shape $\times, I_\nu \not\propto N_X?$

$N(C+ \text{El}=2P, J=3/2)$  $N(C \text{ El}=3P, J=1)$  $N(\text{CO } v=0, J=2)$  $N(\text{CO } v=0, J=4)$  $N(\text{H}_2\text{O } J=1, ka=1, kc=0)$ 

$d_{\text{PDR}} = 50.98"$
 PDR layer
 $R_{\text{cloud}} = 0.5 \text{ pc}$
 $R_{\text{cloud}} = 1.0 \text{ pc}$
 $R_{\text{cloud}} = 3.0 \text{ pc}$
 $R_{\text{cloud}} = 5.0 \text{ pc}$
 observation

The differences in column density profiles are trivial

⇒ need to solve the radiative transfer equation

Solving the Radiative Transfer Equation along LoS

The radiative transfer equation (neglecting dusts, scattering)

$$\frac{dI_\nu}{ds} = A_{ul} n_u \frac{h\nu}{4\pi} \phi(\nu) + B_{ul} n_u \frac{h\nu}{4\pi} I_\nu \phi(\nu) - B_{lu} n_l \frac{h\nu}{4\pi} I_\nu \phi(\nu),$$

with a thermal and turbulent broadening line profile $\phi(\nu)$

Solving the Radiative Transfer Equation along LoS


The radiative transfer equation (neglecting dusts, scattering)

$$\frac{dI_\nu}{ds} = A_{ul} n_u \frac{h\nu}{4\pi} \phi(\nu) + B_{ul} n_u \frac{h\nu}{4\pi} I_\nu \phi(\nu) - B_{lu} n_l \frac{h\nu}{4\pi} I_\nu \phi(\nu),$$

with a thermal and turbulent broadening line profile $\phi(\nu)$

For a toy problem with constant lower and upper level populations

$$\frac{dI_\nu}{ds} = c_1 + c_2 I_\nu, \text{ with } c_1, c_2 \text{ constants}$$



Conclusions

MeudonPDR wrapper

- **column densities in spherical geometry**
- **convolution with the instrument resolution**
- **comparison with observations**
- radiative transfer for line intensities

Conclusions

MeudonPDR wrapper

- **column densities in spherical geometry**
⇒ extended profiles with peaks shifted to greater depths
- **convolution with the instrument resolution**
- **comparison with observations**
- radiative transfer for line intensities

Conclusions

MeudonPDR wrapper

- **column densities in spherical geometry**
⇒ extended profiles with peaks shifted to greater depths
 - **convolution with the instrument resolution**
⇒ further smooths and extends the line spatial profiles
 - **comparison with observations**
-
- radiative transfer for line intensities

Conclusions

MeudonPDR wrapper

- **column densities in spherical geometry**
⇒ extended profiles with peaks shifted to greater depths
- **convolution with the instrument resolution**
⇒ further smooths and extends the line spatial profiles
- **comparison with observations**
⇒ convolved column density profiles match the observation better
cloud radius has a trivial effect
- radiative transfer for line intensities

Conclusions

MeudonPDR wrapper

- **column densities in spherical geometry**
⇒ extended profiles with peaks shifted to greater depths
- **convolution with the instrument resolution**
⇒ further smooths and extends the line spatial profiles
- **comparison with observations**
⇒ convolved column density profiles match the observation better
cloud radius has a trivial effect
radiative transfer equation needs to be solved
- radiative transfer for line intensities

Conclusions

MeudonPDR wrapper


- **column densities in spherical geometry**
⇒ extended profiles with peaks shifted to greater depths
- **convolution with the instrument resolution**
⇒ further smooths and extends the line spatial profiles
- **comparison with observations**
⇒ convolved column density profiles match the observation better
cloud radius has a trivial effect
radiative transfer equation needs to be solved
- radiative transfer for line intensities
 - solver for the radiative transfer equation
 - preliminary results at the line centers

Conclusions

MeudonPDR wrapper


- **column densities in spherical geometry**
⇒ extended profiles with peaks shifted to greater depths
- **convolution with the instrument resolution**
⇒ further smooths and extends the line spatial profiles
- **comparison with observations**
⇒ convolved column density profiles match the observation better
cloud radius has a trivial effect
radiative transfer equation needs to be solved
- radiative transfer for line intensities
 - solver for the radiative transfer equation
 - preliminary results at the line centers
 - full solution with line broadening

Thank you!

- 
- A background image of the Horsehead Nebula, a dark cloud of interstellar dust and gas in the constellation Orion. The nebula's distinct shape resembles a horse's head and neck, set against a backdrop of bright stars.
- ① Introduction
 - Photodissociation Regions (PDRs)
 - The Horsehead Nebula
 - ② Data
 - ③ Motivation
 - ④ Methods
 - The MeudonPDR Code
 - Column Densities in a Spherical PDR
 - Convolution with the Instrument Resolution
 - ⑤ Results
 - Convolved Column Densities from Spherical Models
 - Effect of Cloud Radius on Column Densities
 - ⑥ Solving the Radiative Transfer Equation along LoS


Questions?

Mutiple Peaks in the Observed Profiles




Credit: Maillard, 2023

Exact H₂ Self- and Mutual Shielding



Exact H₂ Self- and Mutual Shielding 2



Preliminary results of solving RTE at the line centers

