

# Explaining Spatial Profiles of Line Emission in the Horsehead Nebula Using Cloud Surface Curvature

Student

Ducheng Lu

Supervisors

Franck Le Petit (LERMA)

Emeric Bron (LERMA)

Jan 2025

# Photodissociation Regions (PDRs)

Interstellar medium (ISM): gas and dust between stars in galaxies

Credit: JWST

- $\sim 10\%$  of the total baryonic mass
- main site of star formation

PDR: regions of **neutral gas** in the ISM where **far-ultraviolet radiation dominates the chemical and heating processes**



- diagnostic of the ISM
- stellar feedback

Heating: photoelectric effect,  
cosmic rays

Cooling: line emission


infrared and line emission

$\Rightarrow$  physical conditions in PDRs



Credit: Tielens 2005

# The Horsehead Nebula




Credit: Hernández-Vera et al. 2023

- observed **edge-on**  $\Rightarrow$  observational access to the chemical stratification


part of a collaboration between JPL, Paris Observatory, IRAM, and CSIC Madrid to study the presence of water in the Horsehead Nebula

# Data



observations along a cut through the Horsehead Nebula


- $C^+$ , C, CO and its isotopologues, and  $H_2O$



# Motivation

## 1D PDR models

- infinite and uniform in two dimensions, depth-dependent only  
 $\Rightarrow$  allows for a detailed study of the physical and chemical processes
- cannot be compared directly to edge-on observations




# Motivation

## 1D PDR models

- infinite and uniform in two dimensions, depth-dependent only  
 $\Rightarrow$  allows for a detailed study of the physical and chemical processes
- cannot be compared directly to edge-on observations

## Proposed Solution:

- approximate edge-on regions with **curvature radius**.
- a spherical geometry wrapper for the MeudonPDR code




# Motivation

## 1D PDR models

- infinite and uniform in two dimensions, depth-dependent only  
 $\Rightarrow$  allows for a detailed study of the physical and chemical processes
- cannot be compared directly to edge-on observations

## Proposed Solution:


- approximate edge-on regions with **curvature radius**.
- a spherical geometry wrapper for the MeudonPDR code
- spatial profiles of **column densities**
- solve **radiative transfer** for line intensities
- **convolution** with the instrument resolution



# The MeudonPDR Code

## stationary 1D PDR code

- radiative transfer
- chemical balance
- level populations
- thermal balance



Cloud size ( $A_{V,\max}$ )	40
Proton density ( $n_H$ )	$3 \times 10^4 - 3 \times 10^6 \text{ cm}^{-3}$
Pressure ( $P$ )	$1 \times 10^6 - 1 \times 10^7 \text{ K cm}^{-3}$
ISRF	shape: Mathis, geometry: beam_isot
ISRF scaling factor	$G_0^{\text{obs}} = 100$ , $G_0^{\text{back}} = 0.04$
UV radiative transfer method	FGK approximation, or exact $\text{H}_2$ self- and mutual shielding
Turbulent velocity dispersion	$2 \text{ km s}^{-1}$
Extinction Curve	HD38087
$R_V = A_V/E(B - V)$	5.50
$C_D = N_H/E(B - V)$	$1.57 \times 10^{22}$


<sup>0</sup>Parameter values are based on Maillard, 2023



# The MeudonPDR Code

## stationary 1D PDR code

- radiative transfer
- chemical balance
- level populations
- thermal balance



Cloud size ( $A_{V,max}$ )	40
Proton density ( $n_H$ )	
Pressure ( $P$ )	$5 \times 10^6 \text{ K cm}^{-3}$
ISRF	shape: Mathis, geometry: beam_isot
ISRF scaling factor	$G_0^{obs} = 100$ , $G_0^{back} = 0.04$
UV radiative transfer method	exact H <sub>2</sub> self- and mutual shielding
Turbulent velocity dispersion	$2 \text{ km s}^{-1}$
Extinction Curve	HD38087
$R_V = A_V/E(B - V)$	5.50
$C_D = N_H/E(B - V)$	$1.57 \times 10^{22}$

+ surface chemistry

<sup>0</sup>Parameter values are based on Maillard, 2023

# From Slab to Spherical Geometry

Input:


- level number density  $n_X(d)$
- cloud radius  $R$  (free parameter)
- LoS impact parameter  $b$

Algorithm:

- interpolation of  $n_X(d) = f(d)$  to allow computation of  $n_X$  at any depth
- $d = R - \sqrt{s^2 + b^2} \Rightarrow n_X(s)$
- integrate along the LoS

$$N_X(b) = 2 \int_0^{s_{\max}} n_X(s') ds'$$

For optically thin lines,  $I_\nu \propto N_X$



# Convolution with the Instrument Resolution

## Algorithm

- interpolate the model grids to uniform ones `x_uniform`, `y_uniform`

# Convolution with the Instrument Resolution

## Algorithm

- interpolate the model grids to uniform ones `x_uniform`, `y_uniform`
- Gaussian kernel with the given resolution, full width half maximum (FWHM)

$$\sigma = \text{FWHM} / (2\sqrt{2\ln 2}), \quad g(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{x^2}{2\sigma^2}\right)$$

# Convolution with the Instrument Resolution

## Algorithm

- interpolate the model grids to uniform ones `x_uniform`, `y_uniform`
- Gaussian kernel with the given resolution, full width half maximum (FWHM)

$$\sigma = \text{FWHM} / (2\sqrt{2\ln 2}), \quad g(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{x^2}{2\sigma^2}\right)$$


# Convolution with the Instrument Resolution

## Algorithm

- interpolate the model grids to uniform ones `x_uniform`, `y_uniform`
- Gaussian kernel with the given resolution, full width half maximum (FWHM)


$$\sigma = \text{FWHM} / (2\sqrt{2\ln 2}), \quad g(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{x^2}{2\sigma^2}\right)$$

- convolution with truncation at  $3\sigma$ , padding `y_uniform` with 0




# Convolved Column Densities from Spherical Models

## step 1: slab vs spherical geometry



# Convolved Column Densities from Spherical Models


spherical geometry: the peaks shift to deeper locations within the cloud






# Convolved Column Densities from Spherical Models

spherical geometry: the peaks shift to deeper locations within the cloud




step 2: convolved vs unconvolved column densities




# Convolved Column Densities from Spherical Models

spherical geometry: the peaks shift to deeper locations within the cloud




convolution: line spatial profiles are smoothed and further extended




# Compare Column Densities with Observations

**convolved column densities** from **spherical models** match observations better



# Compare Column Densities with Observations


**convolved column densities** from **spherical models** match observations better



profile width ✓, shape on the front side ✓, shape on the back side ✗







# Compare Column Densities with Observations

**convolved column densities** from **spherical models** match observations better








profile width ✓, shape on the front side ✓, shape on the back side ✗

Can cloud radius make a difference?


$N(\text{C}^+ \text{ EI}=2\text{P}, J=3/2)$  $N(\text{C EI}=3\text{P}, J=1)$  $N(\text{CO } v=0, J=2)$  $N(\text{CO } v=0, J=4)$  $N(\text{H}_2\text{O } J=1, k_a=1, k_c=0)$  $d_{\text{PDR}} = 50.98''$ 

- PDR layer
- $R_{\text{cloud}}=0.5 \text{ pc}$
- $R_{\text{cloud}}=1.0 \text{ pc}$
- $R_{\text{cloud}}=3.0 \text{ pc}$
- $R_{\text{cloud}}=5.0 \text{ pc}$
- observation

$N(\text{C} + \text{EI}=2\text{P}, J=3/2)$  $N(\text{C EI}=3\text{P}, J=1)$  $N(\text{CO } v=0, J=2)$  $N(\text{CO } v=0, J=4)$  $N(\text{H}_2\text{O } J=1, k_a=1, k_c=0)$  $d_{\text{PDR}} = 50.98''$ 

- PDR layer
- $R_{\text{cloud}}=0.5 \text{ pc}$
- $R_{\text{cloud}}=1.0 \text{ pc}$
- $R_{\text{cloud}}=3.0 \text{ pc}$
- $R_{\text{cloud}}=5.0 \text{ pc}$
- ⋯ observation







The differences in column density profiles are trivial



The differences in column density profiles are trivial

Tail shape  $\propto I_\nu \propto N_X?$



$N(\text{C} + \text{EI}=2\text{P}, J=3/2)$  $N(\text{C EI}=3\text{P}, J=1)$  $N(\text{CO } v=0, J=2)$  $N(\text{CO } v=0, J=4)$  $N(\text{H}_2\text{O } J=1, k_a=1, k_c=0)$  $d_{\text{PDR}} = 50.98''$ 

PDR layer

 $R_{\text{cloud}}=0.5$  pc $R_{\text{cloud}}=1.0$  pc $R_{\text{cloud}}=3.0$  pc $R_{\text{cloud}}=5.0$  pc

..... observation

The differences in column density profiles are trivial

⇒ need to solve the radiative transfer equation

# Solving the Radiative Transfer Equation along LoS

The radiative transfer equation (neglecting dusts, scattering)

$$\frac{dI_\nu}{ds} = A_{ul}n_u \frac{h\nu}{4\pi} \phi(\nu) + B_{ul}n_u \frac{h\nu}{4\pi} I_\nu \phi(\nu) - B_{lu}n_l \frac{h\nu}{4\pi} I_\nu \phi(\nu),$$

with a thermal and turbulent broadening line profile  $\phi(\nu)$

# Solving the Radiative Transfer Equation along LoS


The radiative transfer equation (neglecting dusts, scattering)

$$\frac{dI_\nu}{ds} = A_{ul}n_u \frac{h\nu}{4\pi} \phi(\nu) + B_{ul}n_u \frac{h\nu}{4\pi} I_\nu \phi(\nu) - B_{lu}n_l \frac{h\nu}{4\pi} I_\nu \phi(\nu),$$

with a thermal and turbulent broadening line profile  $\phi(\nu)$

For a toy problem with constant lower and upper level populations

$$\frac{dI_\nu}{ds} = c_1 + c_2 I_\nu, \text{ with } c_1, c_2 \text{ constants}$$



# Conclusions

MeudonPDR wrapper

- **column densities in spherical geometry**
- **convolution with the instrument resolution**
- **comparison with observations**
- radiative transfer for line intensities

# Conclusions

## MeudonPDR wrapper

- **column densities in spherical geometry**  
⇒ extended profiles with peaks shifted to greater depths
- **convolution with the instrument resolution**
- **comparison with observations**
- radiative transfer for line intensities

# Conclusions

## MeudonPDR wrapper

- **column densities in spherical geometry**  
⇒ extended profiles with peaks shifted to greater depths
- **convolution with the instrument resolution**  
⇒ further smooths and extends the line spatial profiles
- **comparison with observations**
  
- radiative transfer for line intensities

# Conclusions

## MeudonPDR wrapper

- **column densities in spherical geometry**
  - ⇒ extended profiles with peaks shifted to greater depths
- **convolution with the instrument resolution**
  - ⇒ further smooths and extends the line spatial profiles
- **comparison with observations**
  - ⇒ convolved column density profiles match the observation better
  - cloud radius has a trivial effect
- radiative transfer for line intensities

# Conclusions

## MeudonPDR wrapper

- **column densities in spherical geometry**
  - ⇒ extended profiles with peaks shifted to greater depths
- **convolution with the instrument resolution**
  - ⇒ further smooths and extends the line spatial profiles
- **comparison with observations**
  - ⇒ convolved column density profiles match the observation better
  - cloud radius has a trivial effect
  - radiative transfer equation needs to be solved
- radiative transfer for line intensities



# Conclusions

## MeudonPDR wrapper

- **column densities in spherical geometry**
  - ⇒ extended profiles with peaks shifted to greater depths
- **convolution with the instrument resolution**
  - ⇒ further smooths and extends the line spatial profiles
- **comparison with observations**
  - ⇒ convolved column density profiles match the observation better
  - cloud radius has a trivial effect
  - radiative transfer equation needs to be solved
- radiative transfer for line intensities
  - solver for the radiative transfer equation
  - preliminary results at the line centers

# Conclusions

## MeudonPDR wrapper

- **column densities in spherical geometry**
  - ⇒ extended profiles with peaks shifted to greater depths
- **convolution with the instrument resolution**
  - ⇒ further smooths and extends the line spatial profiles
- **comparison with observations**
  - ⇒ convolved column density profiles match the observation better
  - cloud radius has a trivial effect
  - radiative transfer equation needs to be solved
- radiative transfer for line intensities
  - solver for the radiative transfer equation
  - preliminary results at the line centers
  - full solution with line broadening

# Thank you!

## 1 Introduction

- Photodissociation Regions (PDRs)
- The Horsehead Nebula

## 2 Data

## 3 Motivation

## 4 Methods

- The MeudonPDR Code
- Column Densities in a Spherical PDR
- Convolution with the Instrument Resolution


## 5 Results

- Convolved Column Densities from Spherical Models
- Effect of Cloud Radius on Column Densities

## 6 Solving the Radiative Transfer Equation along LoS


Questions?

# Mutiple Peaks in the Observed Profiles



Credit: Maillard, 2023

# Exact H2 Self- and Mutual Shielding



# Preliminary results of solving RTE at the line centers

