

Finite Model Theory

Heinz-Dieter Ebbinghaus
Jörg Flum

June 8, 2021

Contents

1	Preliminaries	1
1.1	Structures	1
1.1.1	Graph	2
1.1.2	Orderings	3
1.1.3	Operations on Structures	3
1.2	Syntax and Semantics of First-Order Logic	3
1.3	Some Classical Results of First-Order Logic	5
1.4	Model Classes and Global Relations	6
1.5	Relational Databases and Query Languages	7
2	Ehrenfeucht–Fraïssé Method	7
2.1	Elementary Classes	7
2.2	Ehrenfeucht’s Theorem	8
2.3	Examples and Fraïssé’s Theorem	13
3	Problem List	17

1 Preliminaries

1.1 Structures

Vocabularies are finite sets that consist of **relation symbols** and **constant symbols**. We denote vocabularies by τ, σ, \dots . A **vocabulary** is relational if it does not contain constants.

1.1.1 Graph

Let $\tau = \{E\}$ with a binary relation symbol E . A **graph** (or **undirected graph**) is a τ -structure $\mathcal{G} = (G, E^G)$ satisfying

1. for all $a \in G$: not $E^G aa$
2. for all $a, b \in G$: if $E^G ab$ then $E^G ba$

By GRAPH we denote the class of **finite** graphs. If only (1) is required, we speak of a **digraph**

A subset X of the universe of a graph \mathcal{G} is a **clique**, if $E^G ab$ for all $a, b \in X$, $a \neq b$

Let \mathcal{G} be a digraph. If $n \geq 1$ and

$$E^G a_0 a_1, E^G a_1 a_2, \dots, E^G a_{n-1} a_n$$

then a_0, \dots, a_n is a **path** from a_0 to a_n of **length** n . If $a_0 = a_n$, then a_0, \dots, a_n is a **cycle**. A path a_0, \dots, a_n is **Hamiltonian** if $G = \{a_0, \dots, a_n\}$ and $a_i \neq a_j$ for $i \neq j$. If, in addition, $E^G a_n a_0$ we speak of a **Hamiltonian circuit**

Let \mathcal{G} be a graph. Write $a \sim b$ if $a = b$ or if there is a path from a to b . The equivalence class of a is called the **(connected) component** of a . Let CONN be the class of finite connected graphs

Denote by $d(a, b)$ the length of a shortest path from a to b ; more precisely, define the **distance function** $d : G \times G \rightarrow \mathbb{N} \cup \{\infty\}$ by

$$d(a, b) = \infty \text{ iff } a \not\sim b, \quad d(a, b) = 0 \text{ iff } a = b$$

and otherwise

$$d(a, b) = \min\{n \geq 1 \mid \text{there is a path from } a \text{ to } b \text{ of length } n\}$$

We give the following definitions only for **finite** digraphs. A vertex b is a successor of a vertex a if $E^G ab$. The **in-degree** of a vertex is the number of its predecessors, the **out-degree** the number of its successors.

A **root** of a digraph is a vertex with in-degree 0 and a **leaf** a vertex with out-degree 0.

A **forest** is an acyclic digraph where each vertex has in-degree at most 1. A **tree** is a forest with connected underlying graph. Let TREE be the class of finite trees.

1.1.2 Orderings

Let $\tau = \{<\}$ with a binary relation symbol. A τ -structure $\mathcal{A} = (A, <^A)$ is called an **ordering** if for all $a, b, c \in A$:

1. not $a <^A a$
2. $a <^A b$ or $b <^A a$ or $a = b$
3. if $a <^A b$ and $b <^A c$ then $a <^A c$

1.1.3 Operations on Structures

Two τ -structures \mathcal{A} and \mathcal{B} are **isomorphic**, written $\mathcal{C} \cong \mathcal{B}$, if there is an isomorphism from \mathcal{A} to \mathcal{B} , i.e., a bijection $\pi : A \rightarrow B$ preserving relations and constants, that is

- for n -ary $R \in \tau$ and $a_1, \dots, a_n \in A$

$$R^A a_1 \dots a_n \quad \text{iff} \quad R^B \pi(a_1) \dots \pi(a_n)$$

- for $c \in \tau$, $\pi(c^A) = \pi(c^B)$

For relational τ we introduce the **union** (or, **disjoint union**) of structures. Assume that \mathcal{A} and \mathcal{B} are τ -structures with $A \cap B = \emptyset$. Then $\mathcal{A} \dot{\cup} \mathcal{B}$, the **union** of \mathcal{A} and \mathcal{B} , is the τ -structure with domain $A \cup B$ and

$$R^{\mathcal{A} \dot{\cup} \mathcal{B}} := R^A \cup R^B$$

for any R in τ

Note that the union of ordered structures is not an ordered structure. The situation is different for the so-called **ordered sum**: let τ with $< \in \tau$ be relational and let \mathcal{A} and \mathcal{B} be ordered τ -structures. Assume that $A \cap B = \emptyset$. Define $\mathcal{A} \triangleleft \mathcal{B}$, the **ordered sum** of \mathcal{A} and \mathcal{B} as $\mathcal{A} \dot{\cup} \mathcal{B}$ but setting

$$<^{\mathcal{A} \triangleleft \mathcal{B}} := <^A \cup <^B \cup \{(a, b) \mid a \in A, b \in B\}$$

1.2 Syntax and Semantics of First-Order Logic

Fix a vocabulary τ . Each formula of first-order logic will be a string of symbols taken from the alphabet consisting of

- variables

- connectives
- existential quantifier
- equality symbol
-)(
- the symbols in τ

Denote $\text{FO}[\tau]$ the set of formulas of first-order logic of vocabulary τ .

The axioms for graphs stated above have the following formalizations in $\text{FO}[\{E\}]$

$$\begin{aligned} & \forall x \neg Exx \\ & \forall x \forall y (Exy \rightarrow Eyx) \end{aligned}$$

When only taking into consideration finite structures, we use the notation $\Phi \models_{\text{fin}} \psi$

The **quantifier rank** $\text{qr}(\varphi)$ of a formula φ is the maximum number of nested quantifiers occurring in it

It can be shown that every first-order formula is logically equivalent to a formula in prenex normal form, that is, to a formula of the form $Q_1 x_1, \dots, Q_s x_s \psi$ where $Q_1, \dots, Q_s \in \{\forall, \exists\}$, and where ψ is quantifier-free. Such a formula is called Σ_n if the string of n consecutive blocks, where in each block all quantifiers all of the same type, adjacent blocks contain quantifiers of different type, and the first block is existential. Π_n formulas are defined in the same way but now we require that the first block consists of universal quantifiers. A Δ_n -formula is a formula logically equivalent to both a Σ_n -formula and a Π_n -formula

Given a formula $\varphi(x, \bar{z})$ and $n \geq 1$,

$$\exists^{\geq n} x \varphi(x, \bar{z})$$

is an abbreviation for the formula

$$\exists x_1, \dots, \exists x_n \left(\bigwedge_{1 \leq i \leq n} \varphi(x_i, \bar{z}) \wedge \bigwedge_{1 \leq i < j \leq n} \neg x_i = x_j \right)$$

We set

$$\varphi_{\geq n} := \exists^{\geq n} x x = x$$

Clearly

$$\mathcal{A} \models \varphi_{\geq n} \quad \text{iff} \quad \|\mathcal{A}\| \geq n$$

1.3 Some Classical Results of First-Order Logic

Theorem 1.1. *The set of logically valid sentences of first-order logic is r.e.*

Theorem 1.2 (Compactness Theorem). *Φ is satisfiable iff every finite subset of Φ is satisfiable*

Neither Theorem 1.1 nor 1.2 remain valid if one only considers finite structures. A counterexample for the Compactness Theorem is given by the set $\Phi_\infty := \{\varphi_{\geq n} \mid n \geq 1\}$: Each finite subset of Φ_∞ has a finite model, but Φ_∞ has no finite model

The failure of Theorem 1.1 is documented by

Theorem 1.3 (Trahtenbrot's Theorem). *The set of sentences of first-order logic valid in all finite structures is not r.e.*

Lemma 1.4. *Let $\varphi \in FO[\tau]$ and for $i \in I$, let $\Phi^i \subseteq FO[\tau]$. Assume that*

$$\models \varphi \leftrightarrow \bigvee_{i \in I} \bigwedge \Phi^i$$

Then there is a finite $I_0 \subseteq I$ and for every $i \in I_0$, a finite $\Phi_0^i \subseteq \Phi^i$ s.t.

$$\models \varphi \leftrightarrow \bigvee_{i \in I_0} \bigwedge \Phi_0^i$$

Proof. For simplicity we assume that φ is a sentence and that every Φ^i is a set of sentences. By hypothesis, for some $i \in I$, we have $\Phi^i \models \varphi$; hence, by the Compactness Theorem, $\Phi_0^i \models \varphi$ for some finite $\Phi_0^i \subseteq \Phi^i$.

If there is not such I_0 with $\models \varphi \rightarrow \bigvee_{i \in I_0} \bigwedge \Phi_0^i$, then each finite subset of $\{\varphi\} \cup \{\neg \bigwedge \Phi_0^i \mid i \in I\}$ has a model. Hence by the Compactness Theorem, there is a contradiction \square

Corollary 1.5. *Let Φ be a set of first-order sentences. Assume that any two structures that satisfy the same sentences of Φ are elementarily equivalent. Then any first-order sentence is equivalent to a boolean combination of sentences of Φ*

Proof. For any structure A set

$$\Phi(A) := \{\psi \mid \psi \in \Phi, A \models \psi\} \cup \{\neg\psi \mid \psi \in \Phi, A \models \neg\psi\}$$

Let φ be any first-order sentence. By the preceding lemma it suffices to show that

$$\models \varphi \leftrightarrow \bigvee_{A \models \varphi} \bigwedge \Phi(A)$$

If $\mathcal{B} \models \varphi$ then $\mathcal{B} \models \bigvee_{\mathcal{A} \models \varphi} \bigwedge \Phi(\mathcal{A})$. Suppose $\mathcal{A} \models \bigvee_{\mathcal{A} \models \varphi} \bigwedge \Phi(\mathcal{A})$. Then for some model \mathcal{A} of φ , $\mathcal{B} \models \Phi(\mathcal{A})$. By the definition of $\Phi(\mathcal{A})$, \mathcal{A} and \mathcal{B} satisfy the same sentences of Φ \square

1.4 Model Classes and Global Relations

Fix a vocabulary τ . For a sentence φ of $\text{FO}[\tau]$ we denote by $\text{Mod}(\varphi)$ the class of **finite** models of φ .

$\text{Mod}(\varphi)$ is closed under isomorphisms

For $\varphi(x_1, \dots, x_n) \in \text{FO}[\tau]$ and a structure \mathcal{A} let

$$\varphi^{\mathcal{A}}(-) := \{(a_1, \dots, a_n) \mid \mathcal{A} \models \varphi[a_1, \dots, a_n]\}$$

be the set of n -tuples **defined by φ in \mathcal{A}** . For $n = 0$ this be read as

$$\varphi^{\mathcal{A}} := \begin{cases} \text{TRUE} & \text{if } \mathcal{A} \models \varphi \\ \text{FALSE} & \text{if } \mathcal{A} \not\models \varphi \end{cases}$$

Use this notation we have

$$\text{if } \pi : \mathcal{A} \cong \mathcal{B} \text{ then } \pi(\varphi^{\mathcal{A}}(-)) = \varphi^{\mathcal{B}}(-)$$

where for $X \subseteq A^n$ we set $\pi(X) := \{\pi(a_1), \dots, \pi(a_n) \mid (a_1, \dots, a_n) \in X\}$

Throughout the book all classes K of structures considered will tacitly be assumed to be closed under isomorphisms, i.e.

$$\mathcal{A} \in K \text{ and } \mathcal{A} \cong \mathcal{B} \text{ implies } \mathcal{B} \in K$$

Definition 1.6. Let K be a class of τ -structures. An n -ary **global relation** Γ **on** K is a mapping assigning to each $\mathcal{A} \in K$ an n -ary relation $\Gamma(\mathcal{A})$ on \mathcal{A} satisfying

$$\Gamma(\mathcal{A})a_1 \dots a_n \quad \text{iff} \quad \Gamma(\mathcal{B})\pi(a_1) \dots \pi(a_n)$$

for every isomorphism $\pi : \mathcal{A} \cong \mathcal{B}$ and every $a_1, \dots, a_n \in \mathcal{A}$. If K is the class of all finite τ -structures, then we just speak of an n -ary **global relation**

Example 1.1. 1. Any formula $\varphi(x_1, \dots, x_n) \in \text{FO}[\tau]$ defines the global relation $\mathcal{A} \mapsto \varphi^{\mathcal{A}}(-)$

2. The "transitive closure relation" TC is the binary global relation on GRAPH with

$$\text{TC}(\mathcal{G}) := \{(a, b) \mid a, b \in G, \text{ there is a path from } a \text{ to } b\}$$

3. For $m \geq 0$, Γ_m is a unary global relation on GRAPH, where

$$\Gamma_m(\mathcal{G}) := \{a \mid \|\{b \in G \mid E^{\mathcal{G}}ab\}\| = m\}$$

is the set of elements of \mathcal{G} of degree m

An important issue in model theory is the study of properties of classes of structures that are axiomatizable in a given logic \mathcal{L} and in particular to determine what classes of structures are axiomatizable are what global relations are definable in \mathcal{L} .

1.5 Relational Databases and Query Languages

2 Ehrenfeucht–Fraïssé Method

2.1 Elementary Classes

Proposition 2.1. *Every finite structure can be characterized in first-order logic up to isomorphism, i.e., for every finite structure A there is a sentence φ_A of first-order logic s.t. for all structures B we have*

$$B \models \varphi_A \quad \text{iff} \quad A \cong B$$

Proof. Suppose $A = \{a_1, \dots, a_n\}$. Set $\bar{a} = a_1 \dots a_n$. Let

$$\Theta_n := \{\psi \mid \psi \text{ has the form } Rx_1 \dots x_k, x = y \text{ or } c = x, \\ \text{and variables among } v_1, \dots, v_n\}$$

and

$$\varphi_A := \exists v_1 \dots \exists v_n \left(\bigwedge \{\psi \mid \psi \in \Theta_n, A \models \psi[\bar{a}]\} \wedge \right. \\ \left. \bigwedge \{\neg\psi \mid \psi \in \Theta_n, A \models \neg\psi[\bar{a}]\} \wedge \forall v_{n+1} (v_{n+1} = v_n \vee \dots \vee v_{n+1} = v_n) \right)$$

□

Corollary 2.2. *Let K be a class of finite structures. Then there is a set Φ of first-order sentences s.t.*

$$K = \text{Mod}(\Phi)$$

that is, K is the class of finite models of Φ

Proof. For each n there is only a finite number of pairwise nonisomorphic structures of cardinality n . Let A_1, \dots, A_k be a maximal subset of K of pairwise nonisomorphic structures of cardinality n . Set

$$\psi_n := (\varphi_{=n} \rightarrow (\varphi_{A_1} \vee \dots \vee \varphi_{A_k}))$$

Then $K = \text{Mod}(\{\psi_n \mid n \geq 1\})$ □

Definition 2.3. Let K be a class of finite structures. K is called **axiomatizable in first-order logic** or **elementary** if there is a sentence φ of first-order logic s.t. $K = \text{Mod}(\varphi)$

For structures A and B and $m \in \mathbb{N}$ we write $A \equiv_m B$ and say that A and B are **m -equivalent** if A and B satisfy the same first-order sentences of quantifier rank $\leq m$

Theorem 2.4. Let K be a class of finite structures. Suppose that for every m there are finite structures A and B s.t.

$$A \in K, B \notin K, \text{ and } A \equiv_m B$$

Then K is not axiomatizable in first-order logic

Proof. Let φ be any first-order sentence. Set $m := \text{qr}(\varphi)$. By our assumption there are A and B s.t. $A \in K, B \notin K$, and $A \equiv_m B$. Hence $K \neq \text{Mod}(\varphi)$ □

2.2 Ehrenfeucht's Theorem

Definition 2.5. Assume A and B are structures. Let p be a map with $\text{dom}(p) \subseteq A$. Then p is said to be a **partial isomorphism** from A to B if

1. p is injective
2. for every $c \in \tau: c^A \in \text{dom}(p)$ and $p(c^A) = c^B$
3. for every n -ary $R \in \tau$ and all $a_1, \dots, a_n \in \text{dom}(p)$

$$R^A a_1 \dots a_n \quad \text{iff} \quad R^B p(a_1) \dots p(a_n)$$

We write $\text{Part}(A, B)$ for the set of partial isomorphisms from A to B

In the following we identify a map p with its graph $\{(a, p(a)) \mid a \in \text{dom}(p)\}$. Then $p \subseteq q$ means that q is an extension of p

- Remark.* 1. The empty map, $p = \emptyset$, is a partial isomorphism from A to B just in case the vocabulary contains no constants
2. If $p \neq \emptyset$ is a map with $\text{dom}(p) \subseteq A$ and $\text{ran}(p) \subseteq B$, then p is a partial isomorphism from A to B iff $\text{dom}(p)$ contains c^A for all constants $c \in \tau$ and $p : \text{dom}(p)^A \cong \text{ran}(p)^B$
3. For $\bar{a} = a_1 \dots a_s \in A$ and $\bar{b} = b_1 \dots b_s \in B$ the following statements are equivalent

(a) the clauses

$$p(a_i) = b_i \text{ for } i = 1, \dots, s$$

and

$$p(c^A) = c^B \text{ for } c \text{ in } \tau$$

define a map, which is a partial isomorphism from A to B (henceforth denoted by $\bar{a} \mapsto \bar{b}$)

(b) for all quantifier-free $\varphi(v_1, \dots, v_s)$: $A \models \varphi[\bar{a}]$ iff $B \models \varphi[\bar{b}]$

(c) for all atomic $\varphi(v_1, \dots, v_s)$: $A \models \varphi[\bar{a}]$ iff $B \models \varphi[\bar{b}]$

In general, a **partial isomorphism does not preserve the validity of formulas with quantifiers**: Let $\tau = \{<\}$, $A = (\{0, 1, 2\}, <)$, $B = (\{0, 1, 2, 3\}, <)$ where in both cases $<$ denotes the natural ordering. Then $p_0 := 02 \mapsto 01$ is a partial isomorphism from A to B s.t.

$$A \models \exists v_3 (v_1 < v_3 \wedge v_3 < v_2) [0, 2]$$

but

$$B \not\models \exists v_3 (v_1 < v_3 \wedge v_3 < v_2) [p_0(0), p_0(2)]$$

Let A and B be τ -structures, $\bar{a} \in A^s$, $\bar{b} \in B^s$, and $m \in \mathbb{N}$. The **Ehrenfeucht game** $G_m(A, \bar{a}, B, \bar{b})$ is played by two players called the **spoiler** and the **duplicator**. Each player has to make m moves in the course of a play. In his i -th move the spoiler first selects a structure, A or B , and an element in this structure. If the spoiler chooses e_i in A then the duplicator in his i -th move must choose an element f_i in B . If the spoiler chooses f_i in B then the duplicator must choose an element e_i in A .

	A, \bar{a}	B, \bar{b}
first move	e_1	f_1
second move	e_2	f_2
\vdots	\vdots	\vdots
m -th move	e_m	f_m

The duplicator **wins** iff $\bar{a}\bar{e} \mapsto \bar{b}\bar{f} \in \text{Part}(A, B)$.

Equivalently, the spoiler wins if after some $i \leq m$, $\bar{a}e_1 \dots e_i \mapsto \bar{b}f_1 \dots f_i$ is not a partial isomorphism. We say that a player, the spoiler or the duplicator, has a **winning strategy** in $G_m(A, \bar{a}, B, b)$, or shortly, that he **wins** $G_m(A, \bar{a}, B, \bar{b})$, if it is possible for him to win each play whatever choices are made by the opponent.

If $s = 0$, we denote the game by $G_m(A, B)$

Lemma 2.6. 1. If $A \cong B$ then the duplicator wins $G_m(A, B)$

2. If the duplicator wins $G_{m+1}(A, B)$ and $\|A\| \leq m$ then $A \cong B$

Lemma 2.7. Let A and B be structures, $\bar{a} \in A^s$, $\bar{b} \in B^s$, and $m \geq 0$

1. The duplicator wins $G_0(A, \bar{a}, B, \bar{b})$ iff $\bar{a} \mapsto \bar{b}$ is a partial isomorphism

2. For $m > 0$ the following are equivalent

- (a) The duplicator wins $G_m(A, \bar{a}, B, \bar{b})$
- (b) For all $a \in A$ there is $b \in B$ s.t. the duplicator wins the game $G_{m-1}(A, \bar{a}a, B, \bar{b}b)$ and for all $b \in B$ there is $a \in A$ s.t. the duplicator wins $G_{m-1}(A, \bar{a}a, B, \bar{b}b)$
- (c) If the duplicator wins $G_m(A, \bar{a}, B, \bar{B})$ and if $m' < m$ the duplicator wins $G_{m'}(A, \bar{a}, B, \bar{b})$

Let A be given. For $\bar{a} = a_1 \dots a_s \in A$ and $m \geq 0$ we introduce a formula $\varphi_{\bar{a}}^m(v_1, \dots, v_s)$ that describes the game-theoretic properties of \bar{a} in any game $G_m(A, \bar{a}, \dots)$ s.t. for any B and $\bar{b} = b_1 \dots b_s \in B$

$$B \models \varphi_{\bar{a}}^m[\bar{b}] \quad \text{iff} \quad \text{the duplicator wins } G_m(A, \bar{a}, B, \bar{b})$$

Definition 2.8. Let \bar{v} be v_1, \dots, v_s

$$\varphi_{\bar{a}}^0 := \bigwedge \{ \varphi(\bar{v}) \mid \varphi \text{ atomic or negated atomic}, A \models \varphi[\bar{a}] \}$$

(atomic diagram of A) and for $m > 0$

$$\varphi_{\bar{a}}^m(\bar{v}) := \bigwedge_{a \in A} \exists v_{s+1} \varphi_{\bar{a}a}^{m-1}(\bar{v}, v_{s+1}) \wedge \forall v_{s+1} \bigvee_{a \in A} \varphi_{\bar{a}a}^{m-1}(\bar{v}, v_{s+1})$$

$\varphi_{\bar{a}^0}$ describes the isomorphism type of the substructure generated by \bar{a} in A ; and for $m > 0$ the formula $\varphi_{\bar{a}}^m$ tells us to which isomorphism types the tuple \bar{a} can be extended in m steps adding one element in each step. $\varphi_{\bar{a}}^m$ is called the **m -isomorphism type** (or **m -Hintikka formula**) of \bar{a} in A

Since $\varphi(v_1, \dots, v_s) \mid \varphi$ atomic or negated atomic is finite, a simple induction on m shows

Lemma 2.9. *For $s, m \geq 0$, the set $\{\varphi_{A, \bar{a}}^m \mid A \text{ a structure and } \bar{a} \in A^s\}$ is finite*

Lemma 2.10. 1. $\text{qr}(\varphi_{\bar{a}}^m) = m$

2. $A \models \varphi_{\bar{a}}^m[\bar{a}]$

3. For any B and \bar{b} in B

$$B \models \varphi_{\bar{a}}^0[\bar{b}] \quad \text{iff} \quad \bar{a} \mapsto \bar{b} \in \text{Part}(A, B)$$

Theorem 2.11 (Ehrenfeucht's Theorem). *Given A and B , $\bar{a} \in A^s$ and $\bar{b} \in B^s$, and $m \geq 0$, the following are equivalent*

1. The duplicator wins $G_m(A, \bar{a}, B, \bar{b})$
2. $B \models \varphi_{\bar{a}}^m[\bar{b}]$
3. \bar{a} and \bar{b} satisfy the same formulas of quantifier rank $\leq m$, that is, if $\varphi(x_1, \dots, x_s)$ is of quantifier rank $\leq m$, then

$$A \models \varphi[\bar{a}] \quad \text{iff} \quad B \models \varphi[\bar{b}] \tag{1}$$

Proof. 1 \leftrightarrow 2. Induction on m . For $m = 0$

$$\begin{aligned} \text{the duplicator wins } G_0(A, \bar{a}, B, \bar{b}) & \quad \text{iff} \quad \bar{a} \mapsto \bar{b} \in \text{Part}(A, B) \\ & \quad \text{iff} \quad B \models \varphi_{\bar{a}}^0[\bar{b}] \end{aligned}$$

For $m > 0$

$$\begin{aligned} & \text{the duplicator wins } G_m(A, \bar{a}, B, \bar{b}) \\ \text{iff} & \text{ for all } a \in A, \text{ there is } b \in B \text{ s.t. the duplicator wins} \\ & G_{m-1}(A, \bar{a}a, B, \bar{b}b), \text{ and for all } b \in B \text{ there is } a \in A \text{ s.t.} \\ & \text{the duplicator wins } G_{m-1}(A, \bar{a}a, B, \bar{b}b) \\ \text{iff} & \text{ for all } a \in A, \text{ there is } b \in B \text{ with } B \models \varphi_{\bar{a}a}^{m-1}[\bar{b}b], \text{ and} \\ & \text{for all } b \in B, \text{ there is } a \in A \text{ with } B \models \varphi_{\bar{a}a}^{m-1}[\bar{b}b] \\ \text{iff} & B \models \bigwedge_{a \in A} \exists v_{s+1} \varphi_{\bar{a}a}^{m-1}(\bar{v}, v_{s+1}) \wedge \forall v_{s+1} \bigvee_{a \in A} \varphi_{\bar{a}a}^{m-1}(\bar{v}, v_{s+1})[\bar{b}] \\ \text{iff} & B \models \varphi_{\bar{a}}^m[\bar{b}] \end{aligned}$$

3 \rightarrow 1. $\text{qr}(\varphi_{\bar{a}}^m) = m$ and $A \models \varphi_{\bar{a}}^m[\bar{a}]$

1 \rightarrow 3. Induction on m . The case $m = 0$ is handled as above. Let $m > 0$ and suppose that the duplicator wins $G_m(A, \bar{a}, B, \bar{b})$. Clearly the set of formulas $\varphi(x_1, \dots, x_s)$ satisfying 1 contains the atomic formulas and is closed under \neg and \vee (Since duplicator wins the game, there are partial isomorphisms). Suppose that $\varphi(\bar{a}) = \exists y \psi$ and $\text{qr}(\varphi) \leq m$. Since $y \notin \text{free}(\varphi)$, we can assume that y is distinct from the variables in \bar{x} . Hence $\psi = \psi(\bar{x}, y)$. Assume, for instance, $A \models \varphi(\bar{a})$. Then there is $a \in A$ s.t. $A \models \psi[\bar{a}, a]$. As by 1, the duplicator wins $G_m(A, \bar{a}, B, \bar{b})$, there is $b \in B$ s.t. the duplicator wins $G_{m-1}(A, \bar{a}a, B, \bar{b}b)$. Since $\text{qr}(\psi) \leq m - 1$, the induction hypothesis yields $B \models \psi[\bar{b}, b]$, hence $B \models \varphi[\bar{b}]$ \square

Corollary 2.12. *For structures A, B and $m \geq 0$ the following are equivalent*

1. *The duplicator wins $G_m(A, B)$*
2. $B \models \varphi_A^m$
3. $A \equiv_m B$

Corollary 2.13. *Let A be a structure with $\|A\| \leq m$. Then for all B*

$$B \models \varphi_A^{m+1} \quad \text{iff} \quad A \cong B$$

The next result shows that the formulas $\varphi_{\bar{a}}^m$ give a clear picture of the expressive power of first-order logic

Theorem 2.14. *Let $\varphi(v_1, \dots, v_s)$ be a formula of quantifier rank $\leq m$. Then*

$$\models \varphi \leftrightarrow \bigvee \{ \varphi_{A, \bar{a}}^m \mid A \text{ a structure, } \bar{a} \in A, \text{ and } A \models \varphi[\bar{a}] \}$$

Proof. Suppose first that $B \models \varphi[\bar{b}]$. Then the formula $\varphi_{B, \bar{b}}^m$ is a member of the disjunction on the right side of the equivalence, which therefore is satisfied by \bar{b} .

Conversely, suppose $B \models \bigvee \{ \varphi_{A, \bar{a}}^m[\bar{a}] \}[\bar{b}]$. Then for some A and \bar{a} s.t. $A \models \varphi[\bar{a}]$ we have $B \models \varphi_{A, \bar{a}}^m[\bar{b}]$. By Theorem 2.11 \bar{a} and \bar{b} satisfy the same formulas of quantifier rank $\leq m$ and therefore $B \models \varphi[\bar{b}]$. \square

Theorem 2.15. *For a class K of finite structures the following are equivalent*

1. *K is not axiomatizable in first-order logic*
2. *For each m there are finite structures A and B s.t.*

$$A \in K, B \notin K \text{ and } A \equiv_m B$$

Proof. $2 \rightarrow 1$ is proved in theorem 2.4. For the converse, suppose that 2 doesn't hold, i.e., that for some m and all finite A and B

$$A \in K \text{ and } A \equiv_m B \text{ imply } B \in K$$

Then $K = \text{Mod}(\bigvee \{\varphi_A^m \mid A \in K\})$, and thus K is axiomatizable \square

2.3 Examples and Fraïssé's Theorem

Given structures A, B and $m \in \mathbb{N}$, let $W_m(A, B) :=$

$$\{\bar{a} \mapsto \bar{b} \mid s \geq 0, \bar{a} \in A^s, \bar{b} \in B^s, \text{ the duplicator wins } G_m(A, \bar{a}, B, \bar{b})\}$$

be the set of winning positions for the duplicator. The sequence of the $W_M(A, B)$ has the back and forth properties as introduced in the following definition

Definition 2.16. Structures A and B are said to be *m -isomorphic*, written $A \cong_m B$, if there is a sequence $(I_j)_{j \leq m}$ with the following properties

1. Every I_j is a nonempty set of partial isomorphisms from A to B
2. (**Forth property**) For every $j < m$, $p \in I_{j+1}$ and $a \in A$ there is $q \in I_j$ s.t. $q \supseteq p$ and $a \in \text{dom}(q)$
3. (**Back property**) For every $j < m$, $p \in I_{j+1}$, and $b \in B$ there is $q \in I_j$ s.t. $q \supseteq p$ and $b \in \text{ran}(q)$

If $(I_j)_{j \leq m}$ has the properties 1,2 and 3, we write $(I_j)_{j \leq m} : A \cong_m B$ and say that A and B are *m -isomorphic via $(I_j)_{j \leq m}$*

Exercise 2.3.1. Suppose $(I_j)_{j \leq m} : A \cong_m B$. Then $(\tilde{I}_j)_{j \leq m} : A \cong_m B$ with $\tilde{I}_j := \{q \in \text{Part}(A, B) \mid q \subseteq p \text{ for some } p \in I_j\}$. In particular, $\emptyset \mapsto \emptyset \in I_j$ for all $j \leq m$. Moreover

$$\widetilde{W_j(A, B)} = W_j(A, B)$$

Proof. Forth property: Suppose $j < m$, $p \in \tilde{I}_{j+1}$ and $a \in A$. Then $p \subseteq p' \in I_{j+1}$. Then we have $q' \in I_j$ with $a \in \text{dom}(q')$ and $p' \subseteq q'$. We construct $q = p \cup \{(a, q'(a))\}$. $q \in \text{Part}(A, B)$ since $q' \in \text{Part}(A, B)$. \square

Theorem 2.17. For structures A and B , $\bar{a} \in A^s$, $\bar{b} \in B^s$ and $m \geq 0$ the following are equivalent

1. The duplicator wins $G_m(A, \bar{a}, B, \bar{b})$

2. $\bar{a} \mapsto \bar{b} \in W_m(\mathcal{A}, \mathcal{B})$ and $(W_j(\mathcal{A}, \mathcal{B}))_{j \leq m} : \mathcal{A} \cong_m \mathcal{B}$
3. There is $(I_j)_{j \leq m}$ with $\bar{a} \mapsto \bar{b} \in I_m$ s.t. $(I_j)_{j \leq m} : \mathcal{A} \cong_m \mathcal{B}$
4. $\mathcal{B} \models \varphi_a^m[\bar{b}]$
5. \bar{a} satisfies in \mathcal{A} the same formulas of quantifier rank $\leq m$ as \bar{b} in \mathcal{B}

Proof. $1 \rightarrow 2$. For each $\bar{a} \mapsto \bar{b} \in W_m(\mathcal{A}, \mathcal{B})$, by Lemma 2.7 the duplicator wins $G_0(\mathcal{A}, \bar{a}, \mathcal{B}, \bar{b})$ as the duplicator wins $G_m(\mathcal{A}, \bar{a}, \mathcal{B}, \bar{b})$. For $j < m$, $\bar{a} \mapsto \bar{b} \in W_{j+1}(\mathcal{A}, \mathcal{B})$ and $a \in A$. We have $\bar{a}a \mapsto \bar{b}b \in W_j(\mathcal{A}, \mathcal{B})$ for some $b \in B$.

$2 \rightarrow 3$. Obvious

$3 \rightarrow 1$. Suppose that $(I_j)_{j \leq m} : \mathcal{A} \cong_m \mathcal{B}$ and $\bar{a} \mapsto \bar{b} \in I_m$. We describe a winning strategy in $G_m(\mathcal{A}, \bar{a}, \mathcal{B}, \bar{b})$ for the duplicator: in his i -th move he should choose the element e_i s.t. for $p_i : \bar{a}e_1 \dots e_i \mapsto \bar{b}f_1 \dots f_i$ it is true that $p_i \subseteq q$ for some $q \in I_{m-i}$; this is always possible because of the back and forth properties of $(I_j)_{j \leq m}$ \square

Corollary 2.18. For structures \mathcal{A}, \mathcal{B} and $m \geq 0$ the following are equivalent

1. the duplicator wins $G_m(\mathcal{A}, \mathcal{B})$
2. $(W_j(\mathcal{A}, \mathcal{B}))_{j \leq m} : \mathcal{A} \cong_m \mathcal{B}$
3. $\mathcal{A} \cong_m \mathcal{B}$
4. $\mathcal{B} \models \varphi_A^m$
5. $\mathcal{A} \equiv_m \mathcal{B}$

the equivalence of 3 and 5 is known as Fraïssé's Theorem.

Example 2.1. Let τ be the empty vocabulary and \mathcal{A} and \mathcal{B} be τ -structures. Suppose $\|\mathcal{A}\| \geq m$ and $\|\mathcal{B}\| \geq m$. Then $\mathcal{A} \cong_m \mathcal{B}$. In fact, $(I_j)_{j \leq m} : \mathcal{A} \cong_m \mathcal{B}$ with $I_j := \{p \in \text{Part}(\mathcal{A}, \mathcal{B}) \mid \|\text{dom}(p)\| \leq m - j\}$

As a consequence the class $\text{EVEN}[\tau]$ of finite τ -structures of even cardinality is not axiomatizable in first-order logic. In fact, for each $m > 0$, let \mathcal{A}_m be a structure of cardinality m . Then $\mathcal{A}_m \in \text{EVEN}[\tau]$ iff $\mathcal{A}_{m+1} \notin \text{EVEN}[\tau]$, but $\mathcal{A}_m \cong_m \mathcal{A}_{m+1}$. Now apply Theorem 2.15.

Prove that for arbitrary τ that $\text{EVEN}[\tau]$ is not axiomatizable.

Example 2.2. Let $\tau = \{<, \min, \max\}$ be a vocabulary for finite orderings. Suppose that \mathcal{A} and \mathcal{B} are finite orderings, $\|\mathcal{A}\| > 2^m$ and $\|\mathcal{B}\| > 2^m$ and

$m \geq 1$. Then $A \cong_m B$. Hence the class of finite orderings of even cardinality is not axiomatizable in first-order logic. If we consider orderings as $\{<, S, \min, \max\}$ -structures, the last statements remains true

Given any ordering C , we define its distance function d by

$$d(a, a') := \|\{b \in C \mid (a < b \leq a') \text{ or } (a' < b \leq a)\}\|$$

And for $j \geq 0$, we introduce the "truncated" j -distance function d_j on $C \times C$ by

$$d_j(a, a') := \begin{cases} d(a, a') & \text{if } d(a, a') < 2^j \\ \infty & \text{otherwise} \end{cases}$$

Now suppose that A and B are finite orderings with $\|A\|, \|B\| > 2^m$. For $j \leq m$ set

$$I_j := \{p \in \text{Part}(A, B) \mid d_j(a, a') = d_j(p(a), p(a')) \text{ for } a, a' \in \text{dom}(p)\}$$

Then $(I_j)_{j \leq m} : A \cong_m B$: by assumption on the cardinalities of A and B we have $\{(\min^A, \min^B), (\max^A, \max^B)\} \in I_j$ for every $j \leq m$. (As $d_j(\min, \max) = \infty$) (Ensure its nonempty and then we can continue by forth property?) To give a proof of the forth property of $(I_j)_{j \leq m}$, suppose $j < m, p \in I_{j+1}$ and $a \in A$. We distinguish two cases, depending on whether or the following condition



$$\text{there is an } a' \in \text{dom}(p) \text{ s.t. } d_j(a, a') < 2^j$$

is satisfied. If the condition holds then there is **exactly** one $b \in B$ for which $p \cup \{(a, b)\}$ is a partial isomorphism preserving d_j -distances. (Note the shape of an ordering) Now assume that the condition doesn't hold. let $\text{dom}(p) = \{a_1, \dots, a_r\}$ with $a_1 < \dots < a_r$. We restrict ourselves to the case $a_i < a < a_{i+1}$ for some i . Then $d_j(a_1, a) = \infty$ and $d_j(a, a_{i+1}) = \infty$; hence $d_{j+1}(a_i, a_{i+1}) = \infty$ and therefore $d_{j+1}(p(a_i), p(a_{i+1})) = \infty$. Thus there is a b s.t. $p(a_i) < b < p(a_{i+1})$, $d_j(p(a_i), b) = \infty$ and $d_j(b, p(a_{i+1})) = \infty$. We can verify that $q := p \cup \{(a, b)\}$ is a partial isomorphism

Example 2.3. Let $\tau = \{<, \min, \max\}$ be as in the preceding example and $\sigma = \tau \cup \{E\}$ with a binary relation symbol E . For $n \geq 3$ let A_n be the ordered τ -structure with $A_n = \{0, \dots, n\}$, $\min^{A_n} = 0$, $\max^{A_n} = n$, where $<^{A_n}$ is the natural ordering on A_n and

$$E^{A_n} = \{(i, j) \mid |i - j| = 2\} \cup \{(0, n), (n, 0), (1, n-1), (n-1, 1)\}$$

(A_n, E^{A_n}) is a graph that is connected iff n is odd. Now let $m \geq 2$ and $l, k \geq 2^m$



Let I_j be the set of partial isomorphism from $A_l|_{\tau}$ to $A_k|_{\tau}$ as introduced in the preceding example. We have $(I_j)_{j \leq m} : A_l|_{\tau} \cong_m A_k|_{\tau}$. For $j \geq 2$ any $p \in I_j$ preserves E too, that is, $I_j \subseteq \text{Part}(A_l, A_k)$. If $E^{A_l}ab$ for some $a, b \in A_l$. Then $d_j(a, b) = 4$ or ∞ (since $l \geq 2^m$). We need to ensure $4 < 2^j$ and this is why $j \geq 2$ (guess we loose the condition of the definition-.-). If $d_j(a, b) = 4$ then clearly $E^{A_k}p(a)p(b)$. If $d_j(a, b) = \infty$, a big gap: how to prove $E^{A_k}p(a)p(b)$

Hence $(I_{j+2})_{j \leq m-2} : A_l \cong_{m-2} A_k$ and we have

the class of finite connected ordered graphs is not first-order axiomatizable

Example 2.4. For $l \geq 1$, let G_l be the graph given by a cycle of length $l + 1$. To be precise, set

$$G_l := \{0, \dots, l\}, \dots E^{G_l} := \{(i, i+1) | i < l\} \cup \{(i+1, i) | i < l\} \cup \{(0, l), (l, 0)\}$$

Thus for $l, k \in \mathbb{N}$, the disjoint union $G_l \dot{\cup} G_k$ consists of a cycle of length $l + 1$ and of a cycle of length $k + 1$. We show

$$\text{if } l, k \geq 2^m \text{ then } G_l \cong_m G_k \text{ and } G_l \cong_m G_l \dot{\cup} G_l$$

In fact, for $j \in \mathbb{N}$, define the distance d_j on a graph G by

$$d_j(a, a') = \begin{cases} d(a, b) & \text{if } d(a, b) < 2^{j+1} \\ \infty & \end{cases}$$

We verifies $(I_j)_{j \leq m} : G_l \cong_m G_l \dot{\cup} G_l$ where I_j is the set of $p \in \text{Part}(G_l, G_l \dot{\cup} G_l)$ with

$$\|\text{dom}(p)\| \leq m - j \quad \text{and} \quad d_j(a, b) = d_j(p(a), p(b)) \text{ for } a, b \in \text{dom}(p)$$

Condition $d(a, b) < 2^{j+1}$ is only to ensure $\{(\min^A, \min^B), (\max^A, \max^B)\} \in I_j$ for every $j \leq m$ as $l, k \geq 2^m$.

For every $j < m$, $p \in I_{j+1}$ and $a \in G_l$, if there is a $a' \in \text{dom}(p)$ s.t. $d_j(a, a') < 2^{j+1}$ then there is a unique b for which $q = p \cup \{a, b\}$ is a partial isomorphism. Else, let $\text{dom}(p) = \{a_1, \dots, a_r\}$. Assume $a_i < a < a_{i+1}$ (abuse of notation of course)

We note two consequents.

- The class **CONN** of connected finite graphs is not axiomatizable in first-order logic

$$G_{2^m} \in \text{CONN}, G_{2^m} \dot{\cup} G_{2^m} \notin \text{CONN}, G_{2^m} \equiv_m G_{2^m} \dot{\cup} G_{2^m}$$

- The global relation **TC**, the relation of transitive closure on the class **GRAPH** of finite graphs, is not first-order definable

Suppose $\psi(x, y)$ is a first-order formula defining **TC** on **GRAPH**. Then **CONN** would be the class of finite models of $\forall x \forall y (\neg x = y \rightarrow \psi(x, y))$

Exercise 2.3.2. Set $\tau = \{E\}$. For $l \geq 1$ let \mathcal{B}_l and \mathcal{D}_l be the τ -structures given by

$$\begin{aligned} B_l &:= \{0, \dots, l\}, & E^{B_l} &:= \{(i, i+1) | i < l\} \\ D_l &:= \{0, \dots, l\}, & E^{D_l} &:= \{(i, i+1) | i < l\} \cup \{(l, 0)\} \end{aligned}$$

Given $m \geq 0$, show that $\mathcal{B}_l \cong_m \mathcal{B}_l \dot{\cup} \mathcal{D}_l$ for sufficiently large l . Conclude that the class of finite acyclic digraphs is not axiomatizable in first-order logic

Proof.

□

Proposition 2.19. *The product, the disjoint union and the ordered sum preserve \equiv_m*

Proof. Suppose $A_1 \equiv B_1$ and $A_2 \equiv_m B_2$. By Ehrenfeucht's Theorem there are winning strategies for the duplicator in the games $G_m(A_1, B_1)$ and $G_m(A_2, B_2)$. We refer to these strategies as S_1 and S_2

1. $G_m(A_1 \times A_2, B_1 \times B_2)$: we simultaneously play games in $G_m(A_1, B_1)$ and $G_m(A_2, B_2)$.

□

Corollary 2.20. 1. *If $(A_1, \bar{a}_1) \equiv_m (B_1, \bar{b}_1)$ and $(A_2, \bar{a}_2) \equiv_m (B_2, \bar{b}_2)$ then $(A_1 \dot{\cup} A_2, \bar{a}_1, \bar{a}_2) \equiv_m (B_1 \dot{\cup} B_2, \bar{b}_1, \bar{b}_2)$*

2. *If $(A_1, \bar{a}_1) \equiv_m (B_1, \bar{b}_1)$ and $(A_2, \bar{a}_2) \equiv_m (B_2, \bar{b}_2)$ then $(A_1 \triangleleft A_2, \bar{a}_1, \bar{a}_2) \equiv_m (B_1 \triangleleft B_2, \bar{b}_1, \bar{b}_2)$*

2.4 Hanf's Theorem

All vocabularies in this and the next section will be relational unless stated otherwise. For a nonempty subset M of a structure \mathcal{A} we denote by \mathcal{M} the substructure of \mathcal{A} with universe M

Given a structure \mathcal{A} , we define the binary relation $E^{\mathcal{A}}$ on A by

$$E^{\mathcal{A}} := \{(a, b) \mid a, \text{ and there are } R \text{ in } \tau \text{ and } \bar{c} \text{ s.t. } R^{\mathcal{A}}\bar{c} \\ \text{and } a \text{ and } b \text{ are components of the tuple } \bar{c}\}$$

3 Problem List

2.1

2.3