Introduction

False-Positiv

Discovery Rate

Lasso

Randon Forests

Hierarchica Clustering

Principal Componen Analysis

Comparison o

# Bladder Cancer - Survival Analysis

Arthur Lui Christine Ma

Department of Statistics Brigham Young University

April 22, 2014

False-Positi Discovery

Lasso

Randon Forests

Hierarchica Clustering

Principal Component Analysis

Comparison of

Entura

## Bladder Cancer

- USA 2014: 74690 new cases, 15580 deaths
- Interested in relationship between gene expression and bladder cancer
- Want to compare different statistical methods

False-Positiv

Lasso

Randon Forests

Hierarchica Clustering

Principal Componen Analysis

Comparison o Methods

Future

# Description of Data

- Biomarkers (43149)
- Survival Times
- Censoring Indicators
- Censoring Rate = 58%
- Number of Observations (Patients) = 165
- No dichotomization was done
- Removed the column of NA's in data set

#### Data

False-Position
Discovery
Rate

Lasso

Randon Forests

Hierarchica Clustering

Principal Componen Analysis

Comparison of Methods

e .

# Summary Statistics

## Summary Table of Survival Times

	Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
Censored	5.30	32.33	58.25	63.00	87.17	137.00
Died	1.03	10.40	16.67	28.05	35.70	135.00
Overall	1.03	17.13	36.57	48.38	74.17	137.00

Bladder Cancer -Survival Analysis

#### Arthur Lui Christine Ma

Introduction

False-Positiv

Rate

Lasso

Data

Random Forests

Hierarchica Clustering


Principal Componen

Comparison of


E .

# Histogram of Survival Times


#### Histogram of Time



#### Histogram of Censored Time



#### Histogram of Death Time



Note: We have more data on lower survival times. And more deaths occurred at lower survival times.

Introduction

## Data

alse-Positiv

Lacer


Random

Hierarchica

Principal Componen

Comparison of

Future



Median = 87.07 (33.97, 140.16)

False-Positive Discovery Rate

La330

Random Forests

Hierarchica Clustering

Principal Componer Analysis

Comparison o

Eutura

+ Appropriate for large size of independent and dependent coefficients

 Average fraction of false rejections has to be made or obtained using cross validation

Interaction terms were not included

False-Positive

Discovery Rate

# Cox Model Using Variables with FDR < .025

	coef	exp(coef)	se(coef)	Z	Pr(> z )
genelLMN_1666893	0.28	1.32	0.26	1.04	0.30
genelLMN_1689037	0.82	2.26	0.22	3.70	0.00
geneILMN_1690017	0.32	1.37	0.28	1.15	0.25
genelLMN_1702933	0.45	1.57	0.32	1.42	0.15
geneILMN_1714118	0.48	1.61	0.51	0.94	0.35
genelLMN_1714592	-0.15	0.86	0.37	-0.40	0.69
geneILMN_1718866	0.18	1.20	0.40	0.45	0.65
genelLMN_1745238	0.23	1.26	0.33	0.69	0.49
geneILMN_1757351	0.00	1.00	0.16	0.03	0.98
genelLMN_1767685	-0.26	0.77	0.48	-0.55	0.59
genelLMN_1807525	-0.01	0.99	0.56	-0.03	0.98
genelLMN_1809336	0.40	1.49	0.28	1.42	0.15
genelLMN_1889811	0.69	2.00	0.42	1.64	0.10

Rate

Lasso

Randon Forests

Hierarchica Clustering


Principal Componen Analysis

Comparison of Methods

Future

## FDR KM Plots

#### FDR Model: High-Risk vs. Low-Risk



Low Median = 18.2. High Median = 46.2

Likelihood ratio test=63 on 13 df, p=1.49e-08 n= 130.

False-Positive Discovery

Rate

Randon


Hierarchica Clustering

Principal Componen Analysis

Comparison o

E .

## Residuals Plot



False-Positive Discovery Rate

Lacco

Randon


Hierarchica

Principal Componen Analysis

Comparison o

## FDR AUC

## FDR Time-Dependent ROC



Introductio

.

False-Position Discovery

Rate

Lasso

Randon

Hierarchic

Clustering

Principal Componen Analysis

Comparison o Methods

Future

- + Performs model selection
  - Tuning parameter needs to be estimated

Interaction terms were not included

Introduction

IIILIOUUCLIOI

False-Positiv

Discovery

Lasso

Randon


Hierarchica

Principal
Componen

Comparison of

# Selecting Tuning Parameter $\lambda$





Introduction

IIILIOUUCLIO

False-Positive

Lasso

Random


Hierarchica

Principal Componen

Comparison o

Г. .....

## Selected Variables



	ILMN_1689037	ILMN_1702933
1	0.17	0.03

Data

Discovery

Lasso

Randon

Hierarchica Clustering


Principal Componen Analysis

Comparison o

.........

## Residuals Plot

#### Residuals Plot



Rate

Lasso

Forests

Clustering


Principal Componen Analysis

Comparison of Methods

**Future** 

## Lasso KM Plots

## Penalized Model: High-Risk vs. Low-Risk



Low Median = 36.3 High Median = 18.2

Likelihood ratio test=41.9 on 2 df, p=7.83e-10

Introductio

False-Positiv Discovery


Lasso

Random

Hierarchica Clustering

Principal Componen

Comparison o



#### Random Forests

## Random Forests Model

- 1 A regression tree is a model that predicts the response of an input based on a sequence of decisions
- 2 A Random Forest is created from many trees
- 3 The predicted response of the random forest is the mean of the predictions of the individual trees

False-Positiv

Rate

.

Random Forests

Hierarchica Clustering

Principal Componen Analysis

Comparison o

Future


 Good for modelling non-linear data (data assumed to be nonlinear)

- Lower prediction accuracy

Interaction terms were not included

## Random Forests

# Variable Importance



Introduction

False-Position Discovery

Lasso

Random Forests

Hierarchica Clustering

Principal Componer Analysis

Comparison of Methods

Enture

# Cox Model Using Important Variables from Random Forest

	coef	exp(coef)	se(coef)	Z	р
genesILMN_1689037	0.68	1.98	0.18	3.70	0.00
genesILMN_1702933	1.03	2.80	0.25	4.08	0.00
genesILMN_1704154	0.32	1.37	0.19	1.69	0.09
genesILMN_1749989	-2.32	0.10	1.33	-1.74	0.08

Likelihood ratio test=49.5 on 4 df, p=4.58e-10 n= 130, number of events= 49

Bladder Cancer -Survival Analysis

### Arthur Lui Christine Ma

Introduction

....

False-Positiv

Rate

Random

Forests

Clustering

Principal Componen Analysis


Comparison of Methods

Future

## Random Forest KM Plots

Important Markers: ILMN\_1689037, ILMN\_1702933, ILMN\_1704154, ILMN\_1749989

### Random Forest: High-Risk vs. Low-Risk



#### Median

	Estimate	CI.Lower	CI.Upper
Low-Risk Group	36.30	-39.43	112.03
High-Risk Group	25.83	-9.78	61.45

False-Positiv

\_\_\_\_\_


Random Forests

Hierarchica Clustering

Principal Componen Analysis

Comparison o

## Residuals Plot



Introduction

Introduction

False-Positiv

Discovery

Lasso

Random Forests


Hierarchica

Principal Componen

Comparison o

## Random Forest AUC

## Time-Dependent ROC



False-Positi

Lacco

Randon Forests

Hierarchical Clustering

Principal Componen Analysis

Comparison o

F----

# Hierarchical Clustering Model

 Identify hyperplane that provides maximum separation between clusters

Hierarchical Clustering

# Hierarchical Clustering

- Good result visualization
- Will obtain a hierarchy of clusters
- + Fast computation
- Helpful for identifying gene expression data patterns in time and space
  - Doesn't identify best clusters
  - Sensitive to noise and outliers
  - Might break for large clusters

False-Positiv

Rate

Lasso

Randon Forests

Hierarchical Clustering


Principal Componen Analysis

Comparison of Methods

Euturo

## H-Clust model

## Cluster Dendrogram



dist hclust (\*, "ward")

Hierarchical Clustering

# Cox Model Using Important Variables from H-Clust

	coef	exp(coef)	se(coef)	Z	р
genesILMN_1651236	-0.19	0.83	0.41	-0.45	0.65
genesILMN_1651260	0.44	1.55	0.38	1.16	0.25
genesILMN_1651429	0.26	1.29	0.14	1.78	0.08
genesILMN_1651433	-0.02	0.98	0.35	-0.05	0.96
genesILMN_1651438	0.47	1.60	0.29	1.65	0.10
genesILMN_1651557	-0.23	0.80	0.25	-0.92	0.36
genesILMN_1651574	-0.24	0.78	0.11	-2.21	0.03
genesILMN_1651611	0.17	1.19	0.16	1.04	0.30
genesILMN_1651652	-0.42	0.66	0.32	-1.31	0.19
genesILMN_1651694	0.32	1.38	0.25	1.27	0.20
genesILMN_1651799	0.24	1.27	0.20	1.18	0.24

 $(p-value < 10^{-5})$ 

False-Positiv

Rate

Lasso


Randon

Hierarchical Clustering

Principal
Componen

Comparison o

## Plot of Deviance Residuals



Rate

Forests

Hierarchical Clustering


Principal Componen Analysis

Comparison of Methods

Future

## H-Clust KM Plots

#### Unsupervised Hierarchical Clustering: High-Risk vs. Low-Risk



Low Risk Median = 36.3 (23.1, 49.5)

High Risk Median = 25.8

Likelihood ratio test= 22.25 on 11 df. p-value=0.0225

Introduction

False-Positiv

Discovery

Randon

Hierarchical


Clustering

Principal Componen Analysis

Comparison o

## H-Clust AUC

## Time-Dependent ROC



False-Positiv Discovery

LdSSU

Randon

Hierarchica Clustering

Principal Component Analysis

Comparison of Methods

Euturo

# Principal Component Analysis (PCA)

- 1 Orthogonal Transformation
- 2 Convert a set of observations of possibly correlated variables into a set of values of linearly uncorrelated variables

False-Positiv

Discovery Rate

Randon Forests

Hierarchica Clustering

Principal Component Analysis

Comparison Methods

Future

- + Lack of redundancy of data
- + Reduced complexity
- + Smaller database representation
- + Reduced noise b/c the maximum variation basis is chosen (small variations are ignored)
  - The covariance matrix is hard to evaluate
  - Ability to capture variance depends on the training data

Interaction terms were not included

False-Positi

Discovery

Lasso


Randon

Hierarchica

Principal Component Analysis

Comparison o

## PCA LRT Threshold



Lasso

Forests

Clustering

Principal Component Analysis

Comparison of Methods

Entire

# Cox Model Using Principal Components

	coef	exp(coef)	se(coef)	Z	Pr(> z )
geneILMN_1651574	-0.17	0.84	0.13	-1.37	0.17
geneILMN_1651429	0.23	1.25	0.21	1.08	0.28
genelLMN_1651237	-0.27	0.77	0.20	-1.32	0.19
genelLMN_1651611	-0.07	0.93	0.16	-0.46	0.65
genelLMN_1651832	-0.19	0.82	0.35	-0.55	0.58
genelLMN_1651428	0.16	1.18	0.27	0.60	0.55
genelLMN_1651496	0.02	1.02	0.23	0.10	0.92
genelLMN_1651776	0.23	1.26	0.32	0.70	0.48
genelLMN_1651745	-0.51	0.60	0.23	-2.24	0.02
geneILMN_1651364	0.39	1.48	0.38	1.03	0.30
genelLMN_1651789	0.46	1.58	0.22	2.07	0.04
genelLMN_1651538	0.02	1.02	0.36	0.05	0.96
genelLMN_1651872	0.74	2.09	0.39	1.90	0.06
genelLMN_1651254	-1.15	0.32	0.44	-2.61	0.01
genelLMN_1651336	-0.43	0.65	0.52	-0.83	0.41
genelLMN_1651544	-0.97	0.38	0.40	-2.42	0.02
genelLMN_1651375	0.30	1.35	0.27	1.09	0.28
geneILMN_1651517	-1.59	0.20	0.88	-1.82	0.07

Likelihood ratio test=3.76 on 3 df, p=0.288 n= 35

Lacer

Randon Forests

Hierarchica Clustering


Principal Component Analysis

Comparison of Methods

Future

## **PCA KM Plots**

#### PCA KM: High-Risk vs. Low-Risk



 $\label{eq:Low-Risk-Group-Median} \mbox{Low Risk Group Median} = 35.7. \mbox{ High Risk Group Median} = 25.8.$ 

Likelihood ratio test=59.8 on 18 df, p=2.21e-06

Lacer


Randon

Hierarchica Clustering

Principal Component Analysis

Comparison o

## Plot of Deviance Residuals



Data

Palse-Positiv

Lubbe

Random Forests


Hierarchica Clustering

Principal Component Analysis

Comparison of Methods

## **PCA AUC**


## Time-Dependent ROC




Comparison of Methods

# Comparison of Methods

## Time-Dependent ROC



#### Histogram of Time



Comparison of Methods

# Covariates that appeared most frequently

	FDR	Lasso	RF	H-Clust	PCA
ILMN_1702933	*	*	*		
ILMN_1689037	*	*	*		
ILMN_1651611				*	*
ILMN_1651574				*	*
ILMN_1651429				*	*

Introduction

False-Positiv

Discovery

racc

Lassu

Randon Forests

Hierarchica Clustering

Principal Componen Analysis

Comparison o

**Future** 

## Include other covariates