

Essays on the theory of numbers

Richard Dedekind

1901

Table of contents

Preface	1
Transcriber's Notes	3
I. Continuity and irrational numbers	5
1. Properties of rational numbers	9
2. Comparison of the rational numbers with the points of a straight line	13
3. Continuity of the straight line	15
4. Creation of irrational numbers	19
5. Continuity of the domain of real numbers	25
6. Operations with real numbers	27
7. Infinitesimal analysis	31
II. The nature and meaning of numbers	35
8. Systems of elements	39
9. Transformation of a system	45

Table of contents

10. Similarity of a transformation. Similar systems	49
11. Transformation of a system in itself	53
12. The finite and infinite	61
13. Simply infinite systems. Series of natural numbers	65
14. Greater and less numbers	69
15. Finite and infinite parts of the number-series	79
16. Definition of a transformation of the number-series by induction	83
17. The class of simply infinite systems	91
18. Addition of numbers	95
19. Multiplication of numbers	101
20. Involution of numbers	103
21. Number of the elements of a finite system	105
22. License	107
Bibliography	117

Preface

This eBook is for the use of anyone anywhere at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg License included with this eBook (see Section 22) or online at www.gutenberg.net. This digital edition presents Richard Dedekind's *Essays on the Theory of Numbers*, translated from the original German by Wooster Woodruff Beman.

Originally released on April 8, 2007 as Project Gutenberg [EBook #21016], this volume was prepared from source material encoded in TeX. The original source files, including the TeX sources (21016-t.tex, 21016-t.zip) and compiled documents (21016-pdf.pdf) along with all associated files across various formats, are available at <http://www.gutenberg.org/2/1/0/1/21016/>.

The preparation and digitisation of this text were carried out by Jonathan Ingram, Keith Eddins, and the Online Distributed Proofreading Team at <http://www.pgdp.net>.

This Quarto edition was prepared by Luis Francisco Gómez. The complete source code and project files are hosted on GitHub at <https://github.com/luifrancgom/essays-on-the-theory-of-numbers>.

Transcriber's Notes

In this edition, the symbol $\mathfrak{Z}$ is used as a typographical approximation for the author's original *part* of relational symbol, defined in [Section 8, par. 3](#), which should not be confused with the numerical digit 3. Correspondingly, the symbol $\mathfrak{E}$ serves as its inverse (*whole* of, defined in [Section 8, par. 3](#)), representing the converse relationship such that $S \mathfrak{E} A$ is logically equivalent to $A \mathfrak{Z} S$.

Additionally, the following typographical errors have been corrected:

- [Section 11, par. 42](#): “ $A' \mathfrak{Z} B, B' \mathfrak{Z} B, C' \mathfrak{Z} C$ ” in the original, amended to “ $A' \mathfrak{Z} A$ ”, etc.
- [Section 16, par. 125](#): “the transformation ψ , is therefore completely defined” in the original, amended to “ ψ_1 ”.

Part I.

Continuity and irrational numbers

My attention was first directed toward the considerations which form the subject of this pamphlet in the autumn of 1858. As professor in the Polytechnic School in Zürich I found myself for the first time obliged to lecture upon the elements of the differential calculus and felt more keenly than ever before the lack of a really scientific foundation for arithmetic. In discussing the notion of the approach of a variable magnitude to a fixed limiting value, and especially in proving the theorem that every magnitude which grows continually, but not beyond all limits, must certainly approach a limiting value, I had recourse to geometric evidences. Even now such resort to geometric intuition in a first presentation of the differential calculus, I regard as exceedingly useful, from the didactic standpoint, and indeed indispensable, if one does not wish to lose too much time. But that this form of introduction into the differential calculus can make no claim to being scientific, no one will deny. For myself this feeling of dissatisfaction was so overpowering that I made the fixed resolve to keep meditating on the question till I should find a purely arithmetic and perfectly rigorous foundation for the principles of infinitesimal analysis. The statement is so frequently made that the differential calculus deals with continuous magnitude, and yet an explanation of this continuity is nowhere given; even the most rigorous expositions of the differential calculus do not base their proofs upon continuity but, with more or less consciousness of the fact, they either appeal to geometric notions or those suggested by geometry, or depend upon theorems which are never established in a purely arithmetic manner. Among these, for example, belongs the above-mentioned theorem, and a more careful investigation convinced me that this theorem, or any one equivalent to it, can be regarded in some way as a sufficient basis for infinitesimal analysis. It then only remained to discover its true origin in the elements of arithmetic and thus at the same time to secure a real definition of the essence of continuity. I succeeded Nov. 24, 1858, and a few days afterward I communicated the results of my meditations to my dear friend Durège with whom I had a long and lively discussion. Later I explained these views of a scientific basis of arithmetic to a few of my pupils, and here in Braunschweig read a paper upon the subject before the scientific club of professors, but I could not make up my mind to its publication, because, in the first place, the presentation did not seem altogether simple, and further, the theory itself had little promise.

Nevertheless I had already half determined to select this theme as subject for this occasion, when a few days ago, March 14, by the kindness of the author, the paper *Die Elemente der Funktionenlehre* by E. Heine (*Crelle's Journal*, Vol. 74) [1] came into my hands and confirmed me in my decision. In the main I fully agree with the substance of this memoir, and indeed I could hardly do otherwise, but I will frankly acknowledge that my own presentation seems to me to be simpler in form and to bring out the vital point more clearly. While writing this preface (March 20, 1872), I am just in receipt of the interesting paper *Ueber die Ausdehnung eines Satzes aus der Theorie der trigonometrischen Reihen*, by G. Cantor (*Math. Annalen*, Vol. 5) [2], for which I owe the ingenious author my hearty thanks. As I find on a hasty perusal, the axiom given in Section II. of that paper, aside from the form of presentation, agrees with what I designate in Section 3 as the essence of continuity. But what advantage will be gained by even a purely abstract definition of real numbers of a higher type, I am as yet unable to see, conceiving as I do of the domain of real numbers as complete in itself.

1. Properties of rational numbers

The development of the arithmetic of rational numbers is here presupposed, but still I think it worth while to call attention to certain important matters without discussion, so as to show at the outset the standpoint assumed in what follows. I regard the whole of arithmetic as a necessary, or at least natural, consequence of the simplest arithmetic act, that of counting, and counting itself as nothing else than the successive creation of the infinite series of positive integers in which each individual is defined by the one immediately preceding; the simplest act is the passing from an already-formed individual to the consecutive new one to be formed. The chain of these numbers forms in itself an exceedingly useful instrument for the human mind; it presents an inexhaustible wealth of remarkable laws obtained by the introduction of the four fundamental operations of arithmetic. Addition is the combination of any arbitrary repetitions of the above-mentioned simplest act into a single act; from it in a similar way arises multiplication. While the performance of these two operations is always possible, that of the inverse operations, subtraction and division, proves to be limited. Whatever the immediate occasion may have been, whatever comparisons or analogies with experience, or intuition, may have led thereto; it is certainly true that just this limitation in performing the indirect operations has in each case been the real motive for a new creative act; thus negative and fractional numbers have been created by the human mind; and in the system of all rational numbers there has been gained an instrument of infinitely greater perfection. This system, which I shall denote by R , possesses first of all a completeness and self-containedness which I have designated in another place¹ as characteristic of a *body of numbers* [Zahlkörper] and which consists

¹*Vorlesungen über Zahlentheorie*, by P. G. Lejeune Dirichlet. 2d ed. §159 [3].

1. Properties of rational numbers

in this that the four fundamental operations are always performable with any two individuals in R , i.e., the result is always an individual of R , the single case of division by the number zero being excepted.

For our immediate purpose, however, another property of the system R is still more important; it may be expressed by saying that the system R forms a well-arranged domain of one dimension extending to infinity on two opposite sides. What is meant by this is sufficiently indicated by my use of expressions borrowed from geometric ideas; but just for this reason it will be necessary to bring out clearly the corresponding purely arithmetic properties in order to avoid even the appearance as if arithmetic were in need of ideas foreign to it.

To express that the symbols a and b represent one and the same rational number we put $a = b$ as well as $b = a$. The fact that two rational numbers a , b are different appears in this that the difference $a - b$ has either a positive or negative value. In the former case a is said to be *greater* than b , b *less* than a ; this is also indicated by the symbols $a > b$, $b < a$.² As in the latter case $b - a$ has a positive value it follows that $b > a$, $a < b$. In regard to these two ways in which two numbers may differ the following laws will hold:

I. If $a > b$, and $b > c$, then $a > c$. Whenever a , c are two different (or unequal) numbers, and b is greater than the one and less than the other, we shall, without hesitation because of the suggestion of geometric ideas, express this briefly by saying: b lies between the two numbers a , c .

II. If a , c are two different numbers, there are infinitely many different numbers lying between a , c .

III. If a is any definite number, then all numbers of the system R fall into two classes, A_1 and A_2 , each of which contains infinitely many individuals; the first class A_1 comprises all numbers a_1 that are $< a$, the second class A_2 comprises all numbers a_2 that are $> a$; the number a itself may be assigned at pleasure to the first or second class, being respectively the greatest number of the first class or the least of the second. In every case the separation

²Hence in what follows the so-called “algebraic” greater and less are understood unless the word “absolute” is added.

of the system R into the two classes A_1, A_2 is such that every number of the first class A_1 is less than every number of the second class A_2 .

2. Comparison of the rational numbers with the points of a straight line

The above-mentioned properties of rational numbers recall the corresponding relations of position of the points of a straight line L . If the two opposite directions existing upon it are distinguished by “right” and “left,” and p , q are two different points, then either p lies to the right of q , and at the same time q to the left of p , or conversely q lies to the right of p and at the same time p to the left of q . A third case is impossible, if p , q are actually different points. In regard to this difference in position the following laws hold:

I. If p lies to the right of q , and q to the right of r , then p lies to the right of r ; and we say that q lies between the points p and r .

II. If p , r are two different points, then there always exist infinitely many points that lie between p and r .

III. If p is a definite point in L , then all points in L fall into two classes, P_1 , P_2 , each of which contains infinitely many individuals; the first class P_1 contains all the points p_1 , that lie to the left of p , and the second class P_2 contains all the points p_2 that lie to the right of p ; the point p itself may be assigned at pleasure to the first or second class. In every case the separation of the straight line L into the two classes or portions P_1 , P_2 , is of such a character that every point of the first class P_1 lies to the left of every point of the second class P_2 .

2. *Comparison of the rational numbers with the points of a straight line*

This analogy between rational numbers and the points of a straight line, as is well known, becomes a real correspondence when we select upon the straight line a definite origin or zero-point o and a definite unit of length for the measurement of segments. With the aid of the latter to every rational number a a corresponding length can be constructed and if we lay this off upon the straight line to the right or left of o according as a is positive or negative, we obtain a definite end-point p , which may be regarded as the point corresponding to the number a ; to the rational number zero corresponds the point o . In this way to every rational number a , i.e., to every individual in R , corresponds one and only one point p , i.e., an individual in L . To the two numbers a , b respectively correspond the two points, p , q , and if $a > b$, then p lies to the right of q . To the laws I., II., III. of the previous Section correspond completely the laws I., II., III. of the present.

3. Continuity of the straight line

Of the greatest importance, however, is the fact that in the straight line L there are infinitely many points which correspond to no rational number. If the point p corresponds to the rational number a , then, as is well known, the length op is commensurable with the invariable unit of measure used in the construction, i.e., there exists a third length, a so-called common measure, of which these two lengths are integral multiples. But the ancient Greeks already knew and had demonstrated that there are lengths incommensurable with a given unit of length, e.g., the diagonal of the square whose side is the unit of length. If we lay off such a length from the point o upon the line we obtain an end-point which corresponds to no rational number. Since further it can be easily shown that there are infinitely many lengths which are incommensurable with the unit of length, we may affirm: The straight line L is infinitely richer in point-individuals than the domain R of rational numbers in number-individuals.

If now, as is our desire, we try to follow up arithmetically all phenomena in the straight line, the domain of rational numbers is insufficient and it becomes absolutely necessary that the instrument R constructed by the creation of the rational numbers be essentially improved by the creation of new numbers such that the domain of numbers shall gain the same completeness, or as we may say at once, the same *continuity*, as the straight line.

The previous considerations are so familiar and well known to all that many will regard their repetition quite superfluous. Still I regarded this recapitulation as necessary to prepare properly for the main question. For, the way in which the irrational numbers are usually introduced is based directly upon the conception of extensive magnitudes—which itself is nowhere carefully

3. Continuity of the straight line

defined—and explains number as the result of measuring such a magnitude by another of the same kind.¹ Instead of this I demand that arithmetic shall be developed out of itself.

That such comparisons with non-arithmetic notions have furnished the immediate occasion for the extension of the number-concept may, in a general way, be granted (though this was certainly not the case in the introduction of complex numbers); but this surely is no sufficient ground for introducing these foreign notions into arithmetic, the science of numbers. Just as negative and fractional rational numbers are formed by a new creation, and as the laws of operating with these numbers must and can be reduced to the laws of operating with positive integers, so we must endeavor completely to define irrational numbers by means of the rational numbers alone. The question only remains how to do this.

The above comparison of the domain R of rational numbers with a straight line has led to the recognition of the existence of gaps, of a certain incompleteness or discontinuity of the former, while we ascribe to the straight line completeness, absence of gaps, or continuity. In what then does this continuity consist? Everything must depend on the answer to this question, and only through it shall we obtain a scientific basis for the investigation of *all* continuous domains. By vague remarks upon the unbroken connection in the smallest parts obviously nothing is gained; the problem is to indicate a precise characteristic of continuity that can serve as the basis for valid deductions. For a long time I pondered over this in vain, but finally I found what I was seeking. This discovery will, perhaps, be differently estimated by different people; the majority may find its substance very commonplace. It consists of the following. In the preceding section attention was called to the fact that every point p of the straight line produces a separation of the same into two portions such that every point of one portion lies to the left of every point of the other. I find the essence of continuity in the converse, i.e., in the following principle:

¹The apparent advantage of the generality of this definition of number disappears as soon as we consider complex numbers. According to my view, on the other hand, the notion of the ratio between two numbers of the same kind can be clearly developed only after the introduction of irrational numbers.

“If all points of the straight line fall into two classes such that every point of the first class lies to the left of every point of the second class, then there exists one and only one point which produces this division of all points into two classes, this severing of the straight line into two portions.”

As already said I think I shall not err in assuming that every one will at once grant the truth of this statement; the majority of my readers will be very much disappointed in learning that by this commonplace remark the secret of continuity is to be revealed. To this I may say that I am glad if every one finds the above principle so obvious and so in harmony with his own ideas of a line; for I am utterly unable to adduce any proof of its correctness, nor has any one the power. The assumption of this property of the line is nothing else than an axiom by which we attribute to the line its continuity, by which we find continuity in the line. If space has at all a real existence it is *not* necessary for it to be continuous; many of its properties would remain the same even were it discontinuous. And if we knew for certain that space was discontinuous there would be nothing to prevent us, in case we so desired, from filling up its gaps, in thought, and thus making it continuous; this filling up would consist in a creation of new point-individuals and would have to be effected in accordance with the above principle.

4. Creation of irrational numbers

From the last remarks it is sufficiently obvious how the discontinuous domain R of rational numbers may be rendered complete so as to form a continuous domain. In Section 1 it was pointed out that every rational number a effects a separation of the system R into two classes such that every number a_1 of the first class A_1 is less than every number a_2 of the second class A_2 ; the number a is either the greatest number of the class A_1 or the least number of the class A_2 . If now any separation of the system R into two classes A_1, A_2 is given which possesses only *this* characteristic property that every number a_1 in A_1 is less than every number a_2 in A_2 , then for brevity we shall call such a separation a *cut* [Schnitt] and designate it by (A_1, A_2) . We can then say that every rational number a produces one cut or, strictly speaking, two cuts, which, however, we shall not look upon as essentially different; this cut possesses, *besides*, the property that either among the numbers of the first class there exists a greatest or among the numbers of the second class a least number. And conversely, if a cut possesses this property, then it is produced by this greatest or least rational number.

But it is easy to show that there exist infinitely many cuts not produced by rational numbers. The following example suggests itself most readily.

Let D be a positive integer but not the square of an integer, then there exists a positive integer λ such that

$$\lambda^2 < D < (\lambda + 1)^2.$$

4. Creation of irrational numbers

If we assign to the second class A_2 , every positive rational number a_2 whose square is $> D$, to the first class A_1 all other rational numbers a_1 , this separation forms a cut (A_1, A_2) , i.e., every number a_1 is less than every number a_2 . For if $a_1 = 0$, or is negative, then on that ground a_1 is less than any number a_2 , because, by definition, this last is positive; if a_1 is positive, then is its square $\leq D$, and hence a_1 is less than any positive number a_2 whose square is $> D$.

But this cut is produced by no rational number. To demonstrate this it must be shown first of all that there exists no rational number whose square $= D$. Although this is known from the first elements of the theory of numbers, still the following indirect proof may find place here. If there exist a rational number whose square $= D$, then there exist two positive integers t, u , that satisfy the equation

$$t^2 - Du^2 = 0,$$

and we may assume that u is the *least* positive integer possessing the property that its square, by multiplication by D , may be converted into the square of an integer t . Since evidently

$$\lambda u < t < (\lambda + 1)u,$$

the number $u' = t - \lambda u$ is a positive integer certainly *less* than u . If further we put

$$t' = Du - \lambda t,$$

t' is likewise a positive integer, and we have

$$t'^2 - Du'^2 = (\lambda^2 - D)(t^2 - Du^2) = 0,$$

which is contrary to the assumption respecting u .

Hence the square of every rational number x is either $< D$ or $> D$. From this it easily follows that there is neither in the class A_1 a greatest, nor in the class A_2 a least number. For if we put

$$y = \frac{x(x^2 + 3D)}{3x^2 + D},$$

we have

$$y - x = \frac{2x(D - x^2)}{3x^2 + D}$$

and

$$y^2 - D = \frac{(x^2 - D)^3}{(3x^2 + D)^2}.$$

If in this we assume x to be a positive number from the class A_1 , then $x^2 < D$, and hence $y > x$ and $y^2 < D$. Therefore y likewise belongs to the class A_1 . But if we assume x to be a number from the class A_2 , then $x^2 > D$, and hence $y < x$, $y > 0$, and $y^2 > D$. Therefore y likewise belongs to the class A_2 . This cut is therefore produced by no rational number.

In this property that not all cuts are produced by rational numbers consists the incompleteness or discontinuity of the domain R of all rational numbers.

Whenever, then, we have to do with a cut (A_1, A_2) produced by no rational number, we create a new, an *irrational* number α , which we regard as completely defined by this cut (A_1, A_2) ; we shall say that the number α corresponds to this cut, or that it produces this cut. From now on, therefore, to every definite cut there corresponds a definite rational or irrational number, and we regard two numbers as *different* or *unequal* always and only when they correspond to essentially different cuts.

In order to obtain a basis for the orderly arrangement of all *real*, i.e., of all rational and irrational numbers we must investigate the relation between any two cuts (A_1, A_2) and (B_1, B_2) produced by any two numbers α and β . Obviously a cut (A_1, A_2) is given completely when one of the two classes, e.g., the first A_1 is known, because the second A_2 consists of all rational numbers not contained in A_1 , and the characteristic property of such a first class lies in this that if the number a_1 is contained in it, it also contains all

4. Creation of irrational numbers

numbers less than a_1 . If now we compare two such first classes A_1, B_1 with each other, it may happen

1. That they are perfectly identical, i.e., that every number contained in A_1 is also contained in B_1 , and that every number contained in B_1 is also contained in A_1 . In this case A_2 is necessarily identical with B_2 , and the two cuts are perfectly identical, which we denote in symbols by $\alpha = \beta$ or $\beta = \alpha$.

But if the two classes A_1, B_1 are not identical, then there exists in the one, e.g., in A_1 , a number $a'_1 = b'_2$ not contained in the other B_1 and consequently found in B_2 ; hence all numbers b_1 contained in B_1 are certainly less than this number $a'_1 = b'_2$ and therefore all numbers b_1 are contained in A_1 .

2. If now this number a'_1 is the only one in A_1 that is not contained in B_1 , then is every other number a_1 contained in A_1 also contained in B_1 and is consequently $< a'_1$, i.e., a'_1 is the greatest among all the numbers a_1 , hence the cut (A_1, A_2) is produced by the rational number $a = a'_1 = b'_2$. Concerning the other cut (B_1, B_2) we know already that all numbers b_1 in B_1 are also contained in A_1 and are less than the number $a'_1 = b'_2$ which is contained in B_2 ; every other number b_2 contained in B_2 must, however, be greater than b'_2 , for otherwise it would be less than a'_1 , therefore contained in A_1 and hence in B_1 ; hence b'_2 is the least among all numbers contained in B_2 , and consequently the cut (B_1, B_2) is produced by the same rational number $\beta = b'_2 = a'_1 = \alpha$. The two cuts are then only unessentially different.

3. If, however, there exist in A_1 at least two different numbers $a'_1 = b'_2$ and $a''_1 = b''_2$, which are not contained in B_1 , then there exist infinitely many of them, because all the infinitely many numbers lying between a'_1 and a''_1 are obviously contained in A_1 [Section 1, II.](#) but not in B_1 . In this case we say that the numbers α and β corresponding to these two essentially different cuts (A_1, A_2) and (B_1, B_2) are *different*, and further that α is *greater* than β , that β is *less* than α , which we express in symbols by $\alpha > \beta$ as well as $\beta < \alpha$. It is to be noticed that this definition coincides completely with the one given earlier, when α, β are rational.

The remaining possible cases are these:

4. If there exists in B_1 one and only one number $b'_1 = a'_2$, that is not contained in A_1 then the two cuts (A_1, A_2) and (B_1, B_2) are only unessentially different and they are produced by one and the same rational number $\alpha = a'_2 = b'_1 = \beta$.

5. But if there are in B_1 at least two numbers which are not contained in A_1 , then $\beta > \alpha$, $\alpha < \beta$.

As this exhausts the possible cases, it follows that of two different numbers one is necessarily the greater, the other the less, which gives two possibilities. A third case is impossible. This was indeed involved in the use of the *comparative* (greater, less) to designate the relation between α , β ; but this use has only now been justified. In just such investigations one needs to exercise the greatest care so that even with the best intention to be honest he shall not, through a hasty choice of expressions borrowed from other notions already developed, allow himself to be led into the use of inadmissible transfers from one domain to the other.

If now we consider again somewhat carefully the case $\alpha > \beta$ it is obvious that the less number β , if rational, certainly belongs to the class A_1 ; for since there is in A_1 a number $a'_1 = b'_2$ which belongs to the class B_2 , it follows that the number β , whether the greatest number in B_1 or the least in B_2 is certainly $\leq a'_1$ and hence contained in A_1 . Likewise it is obvious from $\alpha > \beta$ that the greater number α , if rational, certainly belongs to the class B_2 , because $\alpha \geq a'_1$. Combining these two considerations we get the following result: If a cut is produced by the number α then any rational number belongs to the class A_1 or to the class A_2 according as it is less or greater than α ; if the number α is itself rational it may belong to either class.

From this we obtain finally the following: If $\alpha > \beta$, i.e., if there are infinitely many numbers in A_1 not contained in B_1 then there are infinitely many such numbers that at the same time are different from α and from β ; every such rational number c is $< \alpha$, because it is contained in A_1 and at the same time it is $> \beta$ because contained in B_2 .

5. Continuity of the domain of real numbers

In consequence of the distinctions just established the system $\mathfrak{R}$ of all real numbers forms a well-arranged domain of one dimension; this is to mean merely that the following laws prevail:

I. If $\alpha > \beta$, and $\beta > \gamma$, then is also $\alpha > \gamma$. We shall say that the number β lies between α and γ .

II. If α, γ are any two different numbers, then there exist infinitely many different numbers β lying between α, γ .

III. If α is any definite number then all numbers of the system $\mathfrak{R}$ fall into two classes $\mathfrak{A}_1$ and $\mathfrak{A}_2$ each of which contains infinitely many individuals; the first class $\mathfrak{A}_1$ comprises all the numbers α_1 that are less than α , the second $\mathfrak{A}_2$ comprises all the numbers α_2 that are greater than α ; the number α itself may be assigned at pleasure to the first class or to the second, and it is respectively the greatest of the first or the least of the second class. In each case the separation of the system $\mathfrak{R}$ into the two classes $\mathfrak{A}_1, \mathfrak{A}_2$ is such that every number of first class $\mathfrak{A}_1$ is smaller than every number of the second class $\mathfrak{A}_2$ and we say that this separation is produced by the number α .

For brevity and in order not to weary the reader I suppress the proofs of these theorems which follow immediately from the definitions of the previous section.

Beside these properties, however, the domain $\mathfrak{R}$ possesses also *continuity*; i.e., the following theorem is true:

5. Continuity of the domain of real numbers

IV. If the system $\mathfrak{R}$ of all real numbers breaks up into two classes $\mathfrak{A}_1, \mathfrak{A}_2$ such that every number α_1 of the class $\mathfrak{A}_1$ is less than every number α_2 of the class $\mathfrak{A}_2$ then there exists one and only one number α by which this separation is produced.

Proof. By the separation or the cut of $\mathfrak{R}$ into $\mathfrak{A}_1$ and $\mathfrak{A}_2$ we obtain at the same time a cut (A_1, A_2) of the system R of all rational numbers which is defined by this that A_1 contains all rational numbers of the class $\mathfrak{A}_1$ and A_2 all other rational numbers, i.e., all rational numbers of the class $\mathfrak{A}_2$. Let α be the perfectly definite number which produces this cut (A_1, A_2) . If β is any number different from α , there are always infinitely many rational numbers c lying between α and β . If $\beta < \alpha$, then $c < \alpha$; hence c belongs to the class A_1 and consequently also to the class $\mathfrak{A}_1$, and since at the same time $\beta < c$ then β also belongs to the same class $\mathfrak{A}_1$, because every number in $\mathfrak{A}_2$ is greater than every number c in $\mathfrak{A}_1$. But if $\beta > \alpha$, then is $c > \alpha$; hence c belongs to the class A_2 and consequently also to the class $\mathfrak{A}_2$, and since at the same time $\beta > c$, then β also belongs to the same class $\mathfrak{A}_2$, because every number in $\mathfrak{A}_1$ is less than every number c in $\mathfrak{A}_2$. Hence every number β different from α belongs to the class $\mathfrak{A}_1$ or to the class $\mathfrak{A}_2$ according as $\beta < \alpha$ or $\beta > \alpha$; consequently α itself is either the greatest number in $\mathfrak{A}_1$ or the least number in $\mathfrak{A}_2$, i.e., α is one and obviously the only number by which the separation of R into the classes $\mathfrak{A}_1, \mathfrak{A}_2$ is produced. Which was to be proved. $\square$

6. Operations with real numbers

To reduce any operation with two real numbers α, β to operations with rational numbers, it is only necessary from the cuts $(A_1, A_2), (B_1, B_2)$ produced by the numbers α and β in the system R to define the cut (C_1, C_2) which is to correspond to the result of the operation, γ . I confine myself here to the discussion of the simplest case, that of addition.

If c is any rational number, we put it into the class C_1 , provided there are two numbers one a_1 in A_1 and one b_1 in B_1 such that their sum $a_1 + b_1 \geq c$; all other rational numbers shall be put into the class C_2 . This separation of all rational numbers into the two classes C_1, C_2 evidently forms a cut, since every number c_1 in C_1 is less than every number c_2 in C_2 . If both α and β are rational, then every number c_1 contained in C_1 is $\leq \alpha + \beta$, because $a_1 \leq \alpha, b_1 \leq \beta$, and therefore $a_1 + b_1 \leq \alpha + \beta$; further, if there were contained in C_2 a number $c_2 < \alpha + \beta$, hence $\alpha + \beta = c_2 + p$, where p is a positive rational number, then we should have

$$c_2 = (\alpha - \frac{1}{2}p) + (\beta - \frac{1}{2}p),$$

which contradicts the definition of the number c_2 , because $\alpha - \frac{1}{2}p$ is a number in A_1 , and $\beta - \frac{1}{2}p$ a number in B_1 ; consequently every number c_2 contained in C_2 is $\geq \alpha + \beta$. Therefore in this case the cut (C_1, C_2) is produced by the sum $\alpha + \beta$. Thus we shall not violate the definition which holds in the arithmetic of rational numbers if in all cases we understand by the sum $\alpha + \beta$ of any two real numbers α, β that number γ by which the cut (C_1, C_2) is produced. Further, if only one of the two numbers α, β is rational, e.g., α , it is easy to see that it makes no difference with the sum

6. Operations with real numbers

$\gamma = \alpha + \beta$ whether the number α is put into the class A_1 or into the class A_2 .

Just as addition is defined, so can the other operations of the so-called elementary arithmetic be defined, viz., the formation of differences, products, quotients, powers, roots, logarithms, and in this way we arrive at real proofs of theorems (as, e.g., $\sqrt{2} \cdot \sqrt{3} = \sqrt{6}$), which to the best of my knowledge have never been established before. The excessive length that is to be feared in the definitions of the more complicated operations is partly inherent in the nature of the subject but can for the most part be avoided. Very useful in this connection is the notion of an *interval*, i.e., a system A of rational numbers possessing the following characteristic property: if a and a' are numbers of the system A , then are all rational numbers lying between a and a' contained in A . The system R of all rational numbers, and also the two classes of any cut are intervals. If there exist a rational number a_1 which is less and a rational number a_2 which is greater than every number of the interval A , then A is called a finite interval; there then exist infinitely many numbers in the same condition as a_1 and infinitely many in the same condition as a_2 ; the whole domain R breaks up into three parts A_1 , A , A_2 and there enter two perfectly definite rational or irrational numbers α_1 , α_2 which may be called respectively the lower and upper (or the less and greater) *limits* of the interval; the lower limit α_1 is determined by the cut for which the system A_1 forms the first class and the upper α_2 by the cut for which the system A_2 forms the second class. Of every rational or irrational number α lying between α_1 and α_2 it may be said that it lies *within* the interval A . If all numbers of an interval A are also numbers of an interval B , then A is called a portion of B .

Still lengthier considerations seem to loom up when we attempt to adapt the numerous theorems of the arithmetic of rational numbers (as, e.g., the theorem $(a + b)c = ac + bc$) to any real numbers. This, however, is not the case. It is easy to see that it all reduces to showing that the arithmetic operations possess a certain continuity. What I mean by this statement may be expressed in the form of a general theorem:

“If the number λ is the result of an operation performed on

the numbers $\alpha, \beta, \gamma, \dots$ and λ lies within the interval L , then intervals $A, B, C, \dots$ can be taken within which lie the numbers $\alpha, \beta, \gamma, \dots$ such that the result of the same operation in which the numbers $\alpha, \beta, \gamma, \dots$ are replaced by arbitrary numbers of the intervals $A, B, C, \dots$ is always a number lying within the interval L ."

The forbidding clumsiness, however, which marks the statement of such a theorem convinces us that something must be brought in as an aid to expression; this is, in fact, attained in the most satisfactory way by introducing the ideas of *variable magnitudes*, *functions*, *limiting values*, and it would be best to base the definitions of even the simplest arithmetic operations upon these ideas, a matter which, however, cannot be carried further here.

7. Infinitesimal analysis

Here at the close we ought to explain the connection between the preceding investigations and certain fundamental theorems of infinitesimal analysis.

We say that a variable magnitude x which passes through successive definite numerical values approaches a fixed limiting value α when in the course of the process x lies finally between two numbers between which α itself lies, or, what amounts to the same, when the difference $x - \alpha$ taken absolutely becomes finally less than any given value different from zero.

One of the most important theorems may be stated in the following manner:

“If a magnitude x grows continually but not beyond all limits it approaches a limiting value.”

I prove it in the following way. By hypothesis there exists one and hence there exist infinitely many numbers α_2 such that x remains continually $< \alpha_2$; I designate by $\mathfrak{A}_2$ the system of all these numbers α_2 , by $\mathfrak{A}_1$ the system of all other numbers α_1 ; each of the latter possesses the property that in the course of the process x becomes finally $\geq \alpha_1$, hence every number α_1 is less than every number α_2 and consequently there exists a number α which is either the greatest in $\mathfrak{A}_1$ or the least in $\mathfrak{A}_2$ ([Section 5, IV.](#)). The former cannot be the case since x never ceases to grow, hence α is the least number in $\mathfrak{A}_2$. Whatever number α_1 be taken we shall have finally $\alpha_1 < x < \alpha$, i.e., x approaches the limiting value α .

This theorem is equivalent to the principle of continuity, i.e., it loses its validity as soon as we assume a single real number not to be contained in

7. Infinitesimal analysis

the domain $\mathfrak{R}$; or otherwise expressed: if this theorem is correct, then is also theorem [Section 5, IV.](#) correct.

Another theorem of infinitesimal analysis, likewise equivalent to this, which is still oftener employed, may be stated as follows:

“If in the variation of a magnitude x we can for every given positive magnitude δ assign a corresponding position from and after which x changes by less than δ then x approaches a limiting value.”

This converse of the easily demonstrated theorem that every variable magnitude which approaches a limiting value finally changes by less than any given positive magnitude can be derived as well from the preceding theorem as directly from the principle of continuity. I take the latter course. Let δ be any positive magnitude (i.e., $\delta > 0$), then by hypothesis a time will come after which x will change by less than δ , i.e., if at this time x has the value a , then afterwards we shall continually have $x > a - \delta$ and $x < a + \delta$. I now for a moment lay aside the original hypothesis and make use only of the theorem just demonstrated that all later values of the variable x lie between two assignable finite values. Upon this I base a double separation of all real numbers. To the system $\mathfrak{A}_2$ I assign a number α_2 (e.g., $a + \delta$) when in the course of the process x becomes finally $\leq \alpha_2$; to the system $\mathfrak{A}_1$ I assign every number not contained in $\mathfrak{A}_2$; if α_1 is such a number, then, however far the process may have advanced, it will still happen infinitely many times that $x > \alpha_2$. Since every number α_1 is less than every number α_2 there exists a perfectly definite number α which produces this cut $(\mathfrak{A}_1, \mathfrak{A}_2)$ of the system $\mathfrak{R}$ and which I will call the upper limit of the variable x which always remains finite. Likewise as a result of the behavior of the variable x a second cut $(\mathfrak{B}_1, \mathfrak{B}_2)$ of the system $\mathfrak{R}$ is produced; a number β_2 (e.g., $a - \delta$) is assigned to $\mathfrak{B}_2$ when in the course of the process x becomes finally $\geq \beta_2$; every other number β_1 , to be assigned to $\mathfrak{B}_1$, has the property that x is never finally $\geq \beta_1$; therefore infinitely many times x becomes $< \beta_1$; the number β by which this cut is produced I call the lower limiting value of the variable x . The two numbers α, β are obviously characterised by the following property: if ϵ is an arbitrarily small positive magnitude

then we have always finally $x < \alpha + \epsilon$ and $x > \beta - \epsilon$, but never finally $x < \alpha - \epsilon$ and never finally $x > \beta + \epsilon$. Now two cases are possible. If α and β are different from each other, then necessarily $\alpha > \beta$, since continually $\alpha_2 \geq \beta_2$; the variable x oscillates, and, however far the process advances, always undergoes changes whose amount surpasses the value $(\alpha - \beta) - 2\epsilon$ where ϵ is an arbitrarily small positive magnitude. The original hypothesis to which I now return contradicts this consequence; there remains only the second case $\alpha = \beta$ since it has already been shown that, however small be the positive magnitude ϵ , we always have finally $x < \alpha + \epsilon$ and $x > \beta - \epsilon$, x approaches the limiting value α , which was to be proved.

These examples may suffice to bring out the connection between the principle of continuity and infinitesimal analysis.

Part II.

The nature and meaning of numbers

Preface to the first edition

Preface to the first edition

ἀεί ὁ ἄνθρωπος ἀριθμητίζει.

R. Dedekind.

HARZBURG, October 5, 1887.

Preface to the second edition

R. Dedekind.

HARZBURG, August 24, 1893.

8. Systems of elements

1. In what follows I understand by *thing* every object of our thought. In order to be able easily to speak of things, we designate them by symbols, e.g., by letters, and we venture to speak briefly of the thing a or of a simply, when we mean the thing denoted by a and not at all the letter a itself. A thing is completely determined by all that can be affirmed or thought concerning it. A thing a is the same as b (identical with b), and b the same as a , when all that can be thought concerning a can also be thought concerning b , and when all that is true of b can also be thought of a . That a and b are only symbols or names for one and the same thing is indicated by the notation $a = b$, and also by $b = a$. If further $b = c$, that is, if c as well as a is a symbol for the thing denoted by b , then is also $a = c$. If the above coincidence of the thing denoted by a with the thing denoted by b does not exist, then are the things a , b said to be different, a is another thing than b , b another thing than a ; there is some property belonging to the one that does not belong to the other.

2. It very frequently happens that different things, a , b , c , ... for some reason can be considered from a common point of view, can be associated in the mind, and we say that they form a *system* S ; we call the things a , b , c , ... *elements* of the system S , they are *contained* in S ; conversely, S *consists* of these elements. Such a system S (an aggregate, a manifold, a totality) as an object of our thought is likewise a thing (1); it is completely determined when with respect to every thing it is determined whether it is an element of S or not.¹ The system S is hence the same as the system T , in symbols

¹In what manner this determination is brought about, and whether we know a way of deciding upon it, is a matter of indifference for all that follows; the general laws to be developed in no way depend upon it; they hold under all circumstances. I mention

8. Systems of elements

$S = T$, when every element of S is also element of T , and every element of T is also element of S . For uniformity of expression it is advantageous to include also the special case where a system S consists of a *single* (one and only one) element a , i.e., the thing a is element of S , but every thing different from a is not an element of S . On the other hand, we intend here for certain reasons wholly to exclude the empty system which contains no element at all, although for other investigations it may be appropriate to imagine such a system.

3. Definition. A system A is said to be *part* of a system S when every element of A is also element of S . Since this relation between a system A and a system S will occur continually in what follows, we shall express it briefly by the symbol $A \mathfrak{Z} S$. The inverse symbol $S \mathfrak{E} A$, by which the same fact might be expressed, for simplicity and clearness I shall wholly avoid, but for a lack of a better word I shall sometimes say that S is *whole* of A , by which I mean to express that among the elements of S are found all the elements of A . Since further every element s of a system S by (2) can be itself regarded as a system, we can hereafter employ the notation $s \mathfrak{Z} S$.

4. Theorem. $A \mathfrak{Z} A$, by reason of (3).

5. Theorem. If $A \mathfrak{Z} B$ and $B \mathfrak{Z} A$, then $A = B$.

The proof follows from (3), (2).

6. Definition. A system A is said to be a *proper* [*echter*] part of S , when A is part of S , but different from S . According to (5) then S is not a part of A , i.e., there is in S an element which is not an element of A .

7. Theorem. If $A \mathfrak{Z} B$ and $B \mathfrak{Z} C$, which may be denoted briefly by $A \mathfrak{Z} B \mathfrak{Z} C$, then is $A \mathfrak{Z} C$, and A is certainly a proper part of C , if A is a proper part of B or if B is a proper part of C .

this expressly because Kronecker not long ago (*Crelle's Journal*, Vol. 99, pp. 334-336) [4] has endeavored to impose certain limitations upon the free formation of concepts in mathematics which I do not believe to be justified; but there seems to be no call to enter upon this matter with more detail until the distinguished mathematician shall have published his reasons for the necessity or merely the expediency of these limitations.

The proof follows from (3), (6).

8. Definition. By the system *compounded* out of any systems $A, B, C, \dots$ to be denoted by $\mathfrak{M}(A, B, C, \dots)$ we mean that system whose elements are determined by the following prescription: a thing is considered as element of $\mathfrak{M}(A, B, C, \dots)$ when and only when it is element of some one of the systems $A, B, C, \dots$, i.e., when it is element of A , or B , or C , We include also the case where only a single system A exists; then obviously $\mathfrak{M}(A) = A$. We observe further that the system $\mathfrak{M}(A, B, C, \dots)$ compounded out of $A, B, C, \dots$ is carefully to be distinguished from the system whose elements are the systems $A, B, C, \dots$ themselves.

9. Theorem. The systems $A, B, C, \dots$ are parts of $\mathfrak{M}(A, B, C, \dots)$.

The proof follows from (8), (3).

10. Theorem. If $A, B, C, \dots$ are parts of a system S , then is $\mathfrak{M}(A, B, C, \dots) \mathfrak{J} S$.

The proof follows from (8), (3).

11. Theorem. If P is part of one of the systems $A, B, C, \dots$ then is $P \mathfrak{J} \mathfrak{M}(A, B, C, \dots)$.

The proof follows from (9), (7).

12. Theorem. If each of the systems $P, Q, \dots$ is part of one of the systems $A, B, C, \dots$ then is $\mathfrak{M}(P, Q, \dots) \mathfrak{J} \mathfrak{M}(A, B, C, \dots)$.

The proof follows from (11), (10).

13. Theorem. If A is compounded out of any of the systems $P, Q, \dots$ then is $A \mathfrak{J} \mathfrak{M}(P, Q, \dots)$.

Proof. For every element of A is by (8) element of one of the systems $P, Q, \dots$, consequently by (8) also element of $\mathfrak{M}(P, Q, \dots)$, whence the theorem follows by (3). $\square$

8. Systems of elements

14. Theorem. If each of the systems $A, B, C, \dots$ is compounded out of any of the systems $P, Q, \dots$ then is

$$\mathfrak{M}(A, B, C, \dots) \supset \mathfrak{M}(P, Q, \dots)$$

The proof follows from (13), (10).

15. Theorem. If each of the systems $P, Q, \dots$ is part of one of the systems $A, B, C, \dots$, and if each of the latter is compounded out of any of the former, then is

$$\mathfrak{M}(P, Q, \dots) = \mathfrak{M}(A, B, C, \dots).$$

The proof follows from (12), (14), (5).

16. Theorem. If

$$\begin{aligned} A &= \mathfrak{M}(P, Q) \text{ and } B = \mathfrak{M}(Q, R) \\ \text{then is } \mathfrak{M}(A, R) &= \mathfrak{M}(P, B). \end{aligned}$$

Proof. For by the preceding theorem (15)

$$\mathfrak{M}(A, R) \text{ as well as } \mathfrak{M}(P, B) = \mathfrak{M}(P, Q, R).$$

□

17. Definition. A thing g is said to be *common* element of the systems $A, B, C, \dots$, if it is contained in each of these systems (that is in A and in B and in $C \dots$). Likewise a system T is said to be a *common part* of $A, B, C, \dots$ when T is part of each of these systems; and by the *community* [*Gemeinheit*] of the systems $A, B, C, \dots$ we understand the perfectly determinate system $\mathfrak{G}(A, B, C, \dots)$ which consists of all the common elements g of $A, B, C, \dots$ and hence is likewise a common part of those systems. We again include the case where only a single system A occurs; then $\mathfrak{G}(A)$ (is to be put) $= A$. But the case may also occur that the systems $A, B, C, \dots$ possess no

common element at all, therefore no common part, no community; they are then called systems *without* common part, and the symbol $\mathfrak{G}(A, B, C, \dots)$ is meaningless (compare the end of (2)). We shall however almost always in theorems concerning communities leave it to the reader to add in thought the condition of their existence and to discover the proper interpretation of these theorems for the case of non-existence.

18. Theorem. Every common part of $A, B, C, \dots$ is part of $\mathfrak{G}(A, B, C, \dots)$.

The proof follows from (17).

19. Theorem. Every part of $\mathfrak{G}(A, B, C, \dots)$ is common part of $A, B, C, \dots$

The proof follows from (17), (7).

20. Theorem. If each of the systems $A, B, C, \dots$ is whole (3) of one of the systems $P, Q, \dots$ then is

$$\mathfrak{G}(P, Q, \dots) \supset \mathfrak{G}(A, B, C, \dots)$$

Proof. For every element of $\mathfrak{G}(P, Q, \dots)$ is common element of $P, Q, \dots$, therefore also common element of $A, B, C, \dots$, which was to be proved. $\square$

9. Transformation of a system

21. Definition.¹ By a *transformation* [*Abbildung*] ϕ of a system S we understand a law according to which to every determinate element s of S there *belongs* a determinate thing which is called the *transform* of s and denoted by $\phi(s)$; we say also that $\phi(s)$ *corresponds* to the element s , that $\phi(s)$ *results* or is *produced* from s by the transformation ϕ , that s is *transformed* into $\phi(s)$ by the transformation ϕ . If now T is any part of S , then in the transformation ϕ of S is likewise contained a determinate transformation of T , which for the sake of simplicity may be denoted by the same symbol ϕ and consists in this that to every element t of the system T there corresponds the same transform $\phi(t)$, which t possesses as element of S ; at the same time the system consisting of all transforms $\phi(t)$ shall be called the transform of T and be denoted by $\phi(T)$, by which also the significance of $\phi(S)$ is defined. As an example of a transformation of a system we may regard the mere assignment of determinate symbols or names to its elements. The simplest transformation of a system is that by which each of its elements is transformed into itself; it will be called the *identical* transformation of the system. For convenience, in the following theorems (22), (23), (24), which deal with an arbitrary transformation ϕ of an arbitrary system S , we shall denote the transforms of elements s and parts T respectively by s' and T' ; in addition we agree that small and capital italics without accent shall always signify elements and parts of this system S .

22. Theorem.² If $A \mathfrak{Z} B$ then $A' \mathfrak{Z} B'$.

¹See Dirichlet's *Vorlesungen über Zahlentheorie*, 3d edition, 1879, § 163 [3].

²See theorem 27.

9. Transformation of a system

Proof. For every element of A' is the transform of an element contained in A , and therefore also in B , and is therefore element of B' , which was to be proved. $\square$

23. Theorem. The transform of $\mathfrak{M}(A, B, C, \dots)$ is $\mathfrak{M}(A', B', C', \dots)$.

Proof. If we denote the system $\mathfrak{M}(A, B, C, \dots)$ which by (10) is likewise part of S by M , then is every element of its transform M' the transform m' of an element m of M ; since therefore by (8) m is also element of one of the systems $A, B, C, \dots$ and consequently m' element of one of the systems $A', B', C', \dots$, and hence by (8) also element of $\mathfrak{M}(A', B', C', \dots)$, we have by (3)

$$M' \supset \mathfrak{M}(A', B', C', \dots).$$

On the other hand, since $A, B, C, \dots$ are by (9) parts of M , and hence $A', B', C', \dots$ by (22) parts of M' , we have by (10)

$$\mathfrak{M}(A', B', C', \dots) \supset M'.$$

By combination with the above we have by (5) the theorem to be proved

$$M' = \mathfrak{M}(A', B', C', \dots).$$

$\square$

24. Theorem.³ The transform of every common part of $A, B, C, \dots$, and therefore that of the community $\mathfrak{G}(A, B, C, \dots)$ is part of $\mathfrak{G}(A', B', C', \dots)$.

Proof. For by (22) it is common part of $A', B', C', \dots$, whence the theorem follows by (18). $\square$

³See theorem 29.

25. Definition and theorem. If ϕ is a transformation of a system S , and ψ a transformation of the transform $S' = \phi(S)$, there always results a transformation θ of S , compounded⁴ out of ϕ and ψ , which consists of this that to every element s of S there corresponds the transform

$$\theta(s) = \psi(s') = \psi(\phi(s)),$$

where again we have put $\phi(s) = s'$. This transformation θ can be denoted briefly by the symbol $\psi \cdot \phi$ or $\psi\phi$, the transform $\theta(s)$ by $\psi\phi(s)$ where the order of the symbols ϕ, ψ is to be considered, since in general the symbol $\phi\psi$ has no interpretation and actually has meaning only when $\psi(s') \exists s$. If now χ signifies a transformation of the system $\psi(s') = \psi\phi(s)$ and η the transformation $\chi\psi$ of the system S' compounded out of ψ and χ , then is $\chi\theta(s) = \chi\psi(s') = \eta(s') = \eta\phi(s)$; therefore the compound transformations $\chi\theta$ and $\eta\phi$ coincide for every element s of S , i.e., $\chi\theta = \eta\phi$. In accordance with the meaning of θ and η this theorem can finally be expressed in the form

$$\chi \cdot \psi\phi = \chi\psi \cdot \phi,$$

and this transformation compounded out of ϕ, ψ, χ can be denoted briefly by $\chi\psi\phi$.

⁴A confusion of this compounding of transformations with that of systems of elements is hardly to be feared.

10. Similarity of a transformation. Similar systems

26. Definition. A transformation ϕ of a system S is said to be *similar* [*ähnlich*] or *distinct*, when to different elements a, b of the system S there always correspond different transforms $a' = \phi(a), b' = \phi(b)$. Since in this case conversely from $s' = t'$ we always have $s = t$, then is every element of the system $S' = \phi(S)$ the transform s' of a single, perfectly determinate element s of the system S , and we can therefore set over against the transformation ϕ of S an *inverse* transformation of the system S' , to be denoted by $\bar{\phi}$, which consists in this that to every element s' of S' there corresponds the transform $\bar{\phi}(s') = s$, and obviously this transformation is also similar. It is clear that $\bar{\phi}(S') = S$, that further $\bar{\phi}$ is the inverse transformation belonging to ϕ and that the transformation $\bar{\phi}\phi$ compounded out of ϕ and $\bar{\phi}$ by (25) is the identical transformation of S (21). At once we have the following additions to Section 9, retaining the notation there given.

27. Theorem.¹ If $A' \mathfrak{Z} B'$, then $A \mathfrak{Z} B$.

Proof. For if a is an element of A then is a' an element of A' , therefore also of B' , hence $a' = b'$, where b is an element of B ; but since from $a' = b'$ we always have $a = b$, then is every element of A also element of B , which was to be proved. $\square$

¹See theorem 22.

10. *Similarity of a transformation. Similar systems*

28. Theorem. If $A' = B'$, then $A = B$.

The proof follows from (27), (4), (5).

29. Theorem.² If $G = \mathfrak{G}(A, B, C, \dots)$, then

$$G' = \mathfrak{G}(A', B', C', \dots).$$

Proof. Every element of $\mathfrak{G}(A', B', C', \dots)$ is certainly contained in S' , and is therefore the transform g' of an element g contained in S ; but since g' is common element of $A', B', C', \dots$ then by (27) must g be common element of $A, B, C, \dots$ therefore also element of G ; hence every element of $\mathfrak{G}(A', B', C', \dots)$ is transform of an element g of G , therefore element of G' , i.e., $\mathfrak{G}(A', B', C', \dots) \subseteq G'$, and accordingly our theorem follows from (24), (5). $\square$

30. Theorem. The identical transformation of a system is always a similar transformation.

31. Theorem. If ϕ is a similar transformation of S and ψ a similar transformation of $\phi(S)$, then is the transformation $\psi\phi$ of S , compounded of ϕ and ψ , a similar transformation, and the associated inverse transformation $\overline{\psi\phi} = \overline{\phi\psi}$.

Proof. For to different elements a, b of S correspond different transforms $a' = \phi(a), b' = \phi(b)$, and to these again different transforms $\psi(a') = \psi\phi(a), \psi(b') = \psi\phi(b)$ and therefore $\psi\phi$ is a similar transformation. Besides every element $\psi\phi(s) = \psi(s')$ of the system $\psi\phi(S)$ is transformed by $\overline{\psi}$ into $s' = \phi(s)$ and this by $\overline{\phi}$ into s , therefore $\psi\phi(s)$ is transformed by $\overline{\phi\psi}$ into s , which was to be proved. $\square$

32. Definition. The systems R, S are said to be *similar* when there exists such a similar transformation ϕ of S that $\phi(S) = R$, and therefore $\overline{\phi}(R) = S$. Obviously by (30) every system is similar to itself.

²See theorem 24.

33. Theorem. If R, S are similar systems, then every system Q similar to R is also similar to S .

Proof. For if ϕ, ψ are similar transformations of S, R such that $\phi(S) = R$, $\psi(R) = Q$, then by (31) $\psi\phi$ is a similar transformation of S such that $\psi\phi(S) = Q$, which was to be proved. $\square$

34. Definition. We can therefore separate all systems into *classes* by putting into a determinate class all systems $Q, R, S, \dots$, and only those, that are similar to a determinate system R , the *representative* of the class; according to (33) the class is not changed by taking as representative any other system belonging to it.

35. Theorem. If R, S are similar systems, then is every part of S also similar to a part of R , every proper part of S also similar to a proper part of R .

Proof. For if ϕ is a similar transformation of S , $\phi(S) = R$, and $T \subsetneq S$, then by (22) is the system similar to T $\phi(T) \subsetneq R$; if further T is proper part of S , and s an element of S not contained in T , then by (27) the element $\phi(s)$ contained in R cannot be contained in $\phi(T)$; hence $\phi(T)$ is proper part of R , which was to be proved. $\square$

11. Transformation of a system in itself

36. Definition. If ϕ is a similar or dissimilar transformation of a system S , and $\phi(S)$ part of a system Z , then ϕ is said to be a transformation of S in Z , and we say S is transformed by ϕ in Z . Hence we call ϕ a transformation of the system S in itself, when $\phi(S) \mathfrak{Z} S$, and we propose in this paragraph to investigate the general laws of such a transformation ϕ . In doing this we shall use the same notations as in II. and again put $\phi(s) = s'$, $\phi(T) = T'$. These transforms s' , T' are by (22), (7) themselves again elements or parts of S , like all things designated by italic letters.

37. Definition. K is called a *chain* [*Kette*], when $K' \mathfrak{Z} K$. We remark expressly that this name does not in itself belong to the part K of the system S , but is given only with respect to the particular transformation ϕ ; with reference to another transformation of the system S in itself K can very well not be a chain.

38. Theorem. S is a chain.

39. Theorem. The transform K' of a chain K is a chain.

Proof. For from $K' \mathfrak{Z} K$ it follows by (22) that $(K')' \mathfrak{Z} K'$, which was to be proved. $\square$

40. Theorem. If A is part of a chain K , then is also $A' \mathfrak{Z} K$.

Proof. For from $A \mathfrak{Z} K$ it follows by (22) that $A' \mathfrak{Z} K'$, and since by (37) $K' \mathfrak{Z} K$, therefore by (7) $A' \mathfrak{Z} K$, which was to be proved. $\square$

11. Transformation of a system in itself

41. Theorem. If the transform A' is part of a chain L , then is there a chain K , which satisfies the conditions $A \mathfrak{Z} K$, $K' \mathfrak{Z} L$; and $\mathfrak{M}(A, L)$ is just such a chain K .

Proof. If we actually put $K = \mathfrak{M}(A, L)$, then by (9) the one condition $A \mathfrak{Z} K$ is fulfilled. Since further by (23) $K' = \mathfrak{M}(A', L')$ and by hypothesis $A' \mathfrak{Z} L$, $L' \mathfrak{Z} L$, then by (10) is the other condition $K' \mathfrak{Z} L$ also fulfilled and hence it follows because by (9) $L \mathfrak{Z} K$, that also $K' \mathfrak{Z} K$, i.e., K is a chain, which was to be proved. $\square$

42. Theorem. A system M compounded simply out of chains $A, B, C, \dots$ is a chain.

Proof. Since by (23) $M' = \mathfrak{M}(A', B', C', \dots)$ and by hypothesis $A' \mathfrak{Z} A$, $B' \mathfrak{Z} B$, $C' \mathfrak{Z} C$, ... therefore by (12) $M' \mathfrak{Z} M$, which was to be proved. $\square$

43. Theorem. The community G of chains $A, B, C, \dots$ is a chain.

Proof. Since by (17) G is common part of $A, B, C, \dots$, therefore by (22) G' common part of $A', B', C', \dots$, and by hypothesis $A' \mathfrak{Z} A$, $B' \mathfrak{Z} B$, $C' \mathfrak{Z} C$, ..., then by (7) G' is also common part of $A, B, C, \dots$ and hence by (18) also part of G , which was to be proved. $\square$

44. Definition. If A is any part of S , we will denote by A_0 the community of all those chains (e.g., S) of which A is part; this community A_0 exists (17) because A is itself common part of all these chains. Since further by (43) A_0 is a chain, we will call A_0 the *chain of the system* A , or briefly the chain of A . This definition too is strictly related to the fundamental determinate transformation ϕ of the system S in itself, and if later, for the sake of clearness, it is necessary we shall at pleasure use the symbol $\phi_0(A)$ instead of A_0 , and likewise designate the chain of A corresponding to another transformation ω by $\omega_0(A)$. For this very important notion the following theorems hold true.

45. Theorem. $A \mathfrak{Z} A_0$.

Proof. For A is common part of all those chains whose community is A_0 , whence the theorem follows by (18). $\square$

46. Theorem. $(A_0)' \supseteq A_0$.

Proof. For by (44) A_0 is a chain (37). $\square$

47. Theorem. If A is part of a chain K , then is also $A_0 \supseteq K$.

Proof. For A_0 is the community and hence also a common part of all the chains K , of which A is part. $\square$

48. Remark. One can easily convince himself that the notion of the chain A_0 defined in (44) is completely characterised by the preceding theorems, (45), (46), (47).

49. Theorem. $A' \supseteq (A_0)'$.

The proof follows from (45), (22).

50. Theorem. $A' \supseteq A_0$.

The proof follows from (49), (46), (7).

51. Theorem. If A is a chain, then $A_0 = A$.

Proof. Since A is part of the chain A , then by (47) $A_0 \supseteq A$, whence the theorem follows by (45), (5). $\square$

52. Theorem. If $B \supseteq A$, then $B \supseteq A_0$.

The proof follows from (45), (7).

53. Theorem. If $B \supseteq A_0$, then $B_0 \supseteq A_0$, and conversely.

11. Transformation of a system in itself

Proof. Because A_0 is a chain, then by (47) from $B \mathfrak{Z} A_0$, we also get $B_0 \mathfrak{Z} A_0$; conversely, if $B_0 \mathfrak{Z} A_0$, then by (7) we also get $B \mathfrak{Z} A_0$, because by (45) $B \mathfrak{Z} B_0$. $\square$

54. Theorem. If $B \mathfrak{Z} A$, then is $B_0 \mathfrak{Z} A_0$.

The proof follows from (52), (53).

55. Theorem. If $B \mathfrak{Z} A_0$, then is also $B' \mathfrak{Z} A_0$.

Proof. For by (53) $B_0 \mathfrak{Z} A_0$, and since by (50) $B' \mathfrak{Z} B_0$, the theorem to be proved follows by (7). The same result, as is easily seen, can be obtained from (22), (46), (7), or also from (40). $\square$

56. Theorem. If $B \mathfrak{Z} A_0$, then is $(B_0)' \mathfrak{Z} (A_0)'$.

The proof follows from (53), (22).

57. Theorem and definition. $(A_0)' = (A')_0$, i.e., the transform of the chain of A is at the same time the chain of the transform of A . Hence we can designate this system in short by A'_0 and at pleasure call it the *chain-transform* or *transform-chain* of A . With the clearer notation given in (44) the theorem might be expressed by $\phi(\phi_0(A)) = \phi_0(\phi(A))$.

Proof. If for brevity we put $(A')_0 = L$, L is a chain (44) and by (45) $A' \mathfrak{Z} L$; hence by (41) there exists a chain K satisfying the conditions $A \mathfrak{Z} K$, $K' \mathfrak{Z} L$; hence from (47) we have $A_0 \mathfrak{Z} K$, therefore $(A_0)' \mathfrak{Z} K'$, and hence by (7) also $(A_0)' \mathfrak{Z} L$, i.e.,

$$(A_0)' \mathfrak{Z} (A')_0.$$

Since further by (49) $A' \mathfrak{Z} (A_0)'$, and by (44), (39) $(A_0)'$ is a chain, then by (47) also

$$(A')_0 \mathfrak{Z} (A_0)',$$

whence the theorem follows by combining with the preceding result (5). $\square$

58. Theorem. $A_0 = \mathfrak{M}(A, A'_0)$, i.e., the chain of A is compounded out of A and the transform-chain of A .

Proof. If for brevity we again put

$$L = A'_0 = (A_0)' = (A')_0 \text{ and } K = \mathfrak{M}(A, L),$$

then by (45) $A' \mathfrak{Z} L$, and since L is a chain, by (41) the same thing is true of K ; since further $A \mathfrak{Z} K$ (9), therefore by (47)

$$A_0 \mathfrak{Z} K.$$

On the other hand, since by (45) $A \mathfrak{Z} A_0$, and by (46) also $L \mathfrak{Z} A_0$, then by (10) also

$$K \mathfrak{Z} A_0,$$

whence the theorem to be proved $A_0 = K$ follows by combining with the preceding result (5). $\square$

59. Theorem of complete induction. In order to show that the chain A_0 is part of any system Σ —be this latter part of S or not—it is sufficient to show,

ρ . that $A \mathfrak{Z} \Sigma$, and

σ . that the transform of every common element of A_0 and Σ is likewise element of Σ .

11. Transformation of a system in itself

Proof. For if ρ is true, then by (45) the community $G = \mathfrak{G}(A_0, \Sigma)$ certainly exists, and by (18) $A \mathfrak{Z} G$; since besides by (17)

$$G \mathfrak{Z} A_0,$$

then is G also part of our system S , which by ϕ is transformed in itself and at once by (55) we have also $G' \mathfrak{Z} A_0$. If then σ is likewise true, i.e. if $G' \mathfrak{Z} \Sigma$, then must G' as common part of the systems A_0, Σ by (18) be part of their community G , i.e. G is a chain (37), and since, as above noted, $A \mathfrak{Z} G$, then by (47) is also

$$A_0 \mathfrak{Z} G,$$

and therefore by combination with the preceding result $G = A_0$, hence by (17) also $A_0 \mathfrak{Z} \Sigma$, which was to be proved. $\square$

60. The preceding theorem, as will be shown later, forms the scientific basis for the form of demonstration known by the name of complete induction (the inference from n to $n+1$); it can also be stated in the following manner: In order to show that all elements of the chain A_0 possess a certain property $\mathfrak{E}$ (or that a theorem $\mathfrak{S}$ dealing with an undetermined thing n actually holds good for all elements n of the chain A_0) it is sufficient to show

ρ . that all elements a of the system A possess the property $\mathfrak{E}$ (or that $\mathfrak{S}$ holds for all a 's) and

σ . that to the transform n' of every such element n of A_0 possessing the property $\mathfrak{E}$, belongs the same property $\mathfrak{E}$ (or that the theorem $\mathfrak{S}$, as soon as it holds for an element n of A_0 , certainly must also hold for its transform n').

Indeed, if we denote by Σ the system of all things possessing the property $\mathfrak{E}$ (or for which the theorem $\mathfrak{S}$ holds) the complete agreement of the present manner of stating the theorem with that employed in (59) is immediately obvious.

61. Theorem. The chain of $\mathfrak{M}(A, B, C, \dots)$ is $\mathfrak{M}(A_0, B_0, C_0, \dots)$.

Proof. If we designate by M the former, by K the latter system, then by (42) K is a chain. Since then by (45) each of the systems $A, B, C, \dots$ is part of one of the systems $A_0, B_0, C_0, \dots$, and therefore by (12) $M \supset K$, then by (47) we also have

$$M_0 \supset K.$$

On the other hand, since by (9) each of the systems $A, B, C, \dots$ is part of M , and hence by (45), (7) also part of the chain M_0 , then by (47) must also each of the systems $A_0, B_0, C_0, \dots$ be part of M_0 , therefore by (10)

$$K \supset M_0$$

whence by combination with the preceding result follows the theorem to be proved $M_0 = K$ (5). $\square$

62. Theorem. The chain of $\mathfrak{G}(A, B, C, \dots)$ is part of $\mathfrak{G}(A_0, B_0, C_0, \dots)$.

Proof. If we designate by G the former, by K the latter system, then by (43) K is a chain. Since then each of the systems $A_0, B_0, C_0, \dots$ by (45) is whole of one of the systems $A, B, C, \dots$, and hence by (20) $G \supset K$, therefore by (47) we obtain the theorem to be proved $G_0 \supset K$. $\square$

63. Theorem. If $K' \supset L \supset K$, and therefore K is a chain, L is also a chain. If the same is proper part of K , and U the system of all those elements of K which are not contained in L , and if further the chain U_0 is proper part of K , and V the system of all those elements of K which are not contained in U_0 , then is $K = \mathfrak{M}(U_0, V)$ and $L = \mathfrak{M}(U'_0, V)$. If finally $L = K'$ then $V \supset V'$.

The proof of this theorem of which (as of the two preceding) we shall make no use may be left for the reader.

12. The finite and infinite

64. Definition.¹ A system S is said to be *infinite* when it is similar to a proper part of itself (32); in the contrary case S is said to be a *finite* system.

65. Theorem. Every system consisting of a single element is finite.

Proof. For such a system possesses no proper part (2), (6). □

66. Theorem. There exist infinite systems.

Proof. ² My own realm of thoughts, i.e., the totality S of all things, which can be objects of my thought, is infinite. For if s signifies an element of S , then is the thought s' , that s can be object of my thought, itself an element of S . If we regard this as transform $\phi(s)$ of the element s then has the transformation ϕ of S , thus determined, the property that the transform S' is part of S ; and S' is certainly proper part of S , because there are elements in S (e.g., my own ego) which are different from such thought s' and therefore are not contained in S' . Finally it is clear that if a, b

¹If one does not care to employ the notion of similar systems (32) he must say: S is said to be infinite, when there is a proper part of S (6) in which S can be distinctly (similarly) transformed (26), (36). In this form I submitted the definition of the infinite which forms the core of my whole investigation in September, 1882, to G. Cantor and several years earlier to Schwarz and Weber. All other attempts that have come to my knowledge to distinguish the infinite from the finite seem to me to have met with so little success that I think I may be permitted to forego any criticism of them.

²A similar consideration is found in § 13 of the *Paradoxien des Unendlichen* by Bolzano (Leipzig, 1851) [5].

12. The finite and infinite

are different elements of S , their transforms a' , b' are also different, that therefore the transformation ϕ is a distinct (similar) transformation (26). Hence S is infinite, which was to be proved. $\square$

67. Theorem. If R , S are similar systems, then is R finite or infinite according as S is finite or infinite.

Proof. If S is infinite, therefore similar to a proper part S' of itself, then if R and S are similar, S' by (33) must be similar to R and by (35) likewise similar to a proper part of R , which therefore by (33) is itself similar to R ; therefore R is infinite, which was to be proved. $\square$

68. Theorem. Every system S , which possesses an infinite part is likewise infinite; or, in other words, every part of a finite system is finite.

Proof. If T is infinite and there is hence such a similar transformation ψ of T , that $\psi(T)$ is a proper part of T , then, if T is part of S , we can extend this transformation ψ to a transformation ϕ of S in which, if s denotes any element of S , we put $\phi(s) = \psi(s)$ or $\phi(s) = s$ according as s is element of T or not. This transformation ϕ is a similar one; for, if a , b denote different elements of S , then if both are contained in T , the transform $\phi(a) = \psi(a)$ is different from the transform $\phi(b) = \psi(b)$, because ψ is a similar transformation; if further a is contained in T , but b not, then is $\phi(a) = \psi(a)$ different from $\phi(b) = b$, because $\psi(a)$ is contained in T ; if finally neither a nor b is contained in T then also is $\phi(a) = a$ different from $\phi(b) = b$, which was to be shown. Since further $\psi(T)$ is part of T , because by (7) also part of S , it is clear that also $\phi(S) \not\supset S$. Since finally $\psi(T)$ is proper part of T there exists in T and therefore also in S , an element t , not contained in $\psi(T) = \phi(T)$; since then the transform $\phi(s)$ of every element s not contained in T is equal to s , and hence is different from t , t cannot be contained in $\phi(S)$; hence $\phi(S)$ is proper part of S and consequently S is infinite, which was to be proved. $\square$

69. Theorem. Every system which is similar to a part of a finite system, is itself finite.

Proof. The proof follows from (67), (68). □

70. Theorem. If a is an element of S , and if the aggregate T of all the elements of S different from a is finite, then is also S finite.

Proof. We have by (64) to show that if ϕ denotes any similar transformation of S in itself, the transform $\phi(S)$ or S' is never a proper part of S but always $= S$. Obviously $S = \mathfrak{M}(a, T)$ and hence by (23), if the transforms are again denoted by accents, $S' = \mathfrak{M}(a', T')$, and, on account of the similarity of the transformation ϕ , a' is not contained in T' (26). Since further by hypothesis $S' \supset S$, then must a' and likewise every element of T' either $= a$, or be element of T . If then—a case which we will treat first— a is not contained in T' , then must $T' \supset T$ and hence $T' = T$, because ϕ is a similar transformation and because T is a finite system; and since a' , as remarked, is not contained in T' , i.e., not in T , then must $a' = a$, and hence in this case we actually have $S' = S$ as was stated. In the opposite case when a is contained in T' and hence is the transform b' of an element b contained in T , we will denote by U the aggregate of all those elements u of T , which are different from b ; then $T = \mathfrak{M}(b, U)$ and by (15) $S = \mathfrak{M}(a, b, U)$, hence $S' = \mathfrak{M}(a', a, U')$. We now determine a new transformation ψ of T in which we put $\psi(b) = a'$, and generally $\psi(u) = u'$, whence by (23) $\psi(T) = \mathfrak{M}(a', U')$. Obviously ψ is a similar transformation, because ϕ was such, and because a is not contained in U and therefore also a' not in U' . Since further a and every element u is different from b then (on account of the similarity of ϕ) must also a' and every element u' be different from a and consequently contained in T ; hence $\psi(T) \supset T$ and since T is finite, therefore must $\psi(T) = T$, and $\mathfrak{M}(a', U') = T$. From this by (15) we obtain

$$\mathfrak{M}(a', a, U') = \mathfrak{M}(a, T)$$

12. *The finite and infinite*

i.e., according to the preceding $S' = S$. Therefore in this case also the proof demanded has been secured. $\square$

13. Simply infinite systems. Series of natural numbers

71. Definition. A system N is said to be *simply infinite* when there exists a similar transformation ϕ of N in itself such that N appears as chain (44) of an element not contained in $\phi(N)$. We call this element, which we shall denote in what follows by the symbol 1, the *base-element* of N and say the simply infinite system N is *set in order* [*geordnet*] by this transformation ϕ . If we retain the earlier convenient symbols for transforms and chains (Section 11) then the essence of a simply infinite system N consists in the existence of a transformation ϕ of N and an element 1 which satisfy the following conditions α , β , γ , δ :

α . $N' \not\supset N$.

β . $N = 1_0$.

γ . The element 1 is not contained in N' .

δ . The transformation ϕ is similar.

Obviously it follows from α , γ , δ that every simply infinite system N is actually an infinite system (64) because it is similar to a proper part N' of itself.

72. Theorem. In every infinite system S a simply infinite system N is contained as a part.

Proof. By (64) there exists a similar transformation ϕ of S such that $\phi(S)$ or S' is a proper part of S ; hence there exists an element 1 in S which

13. Simply infinite systems. Series of natural numbers

is not contained in S' . The chain $N = 1_0$, which corresponds to this transformation ϕ of the system S in itself (44), is a simply infinite system set in order by ϕ ; for the characteristic conditions $\alpha, \beta, \gamma, \delta$ in (71) are obviously all fulfilled. $\square$

73. Definition. If in the consideration of a simply infinite system N set in order by a transformation ϕ we entirely neglect the special character of the elements; simply retaining their distinguishability and taking into account only the relations to one another in which they are placed by the order-setting transformation ϕ , then are these elements called *natural numbers* or *ordinal numbers* or simply *numbers*, and the base-element 1 is called the *base-number* of the *number-series* N . With reference to this freeing the elements from every other content (abstraction) we are justified in calling numbers a free creation of the human mind. The relations or laws which are derived entirely from the conditions $\alpha, \beta, \gamma, \delta$ in (71) and therefore are always the same in all ordered simply infinite systems, whatever names may happen to be given to the individual elements (compare 134), form the first object of the *science of numbers* or *arithmetic*. From the general notions and theorems of Section 11 about the transformation of a system in itself we obtain immediately the following fundamental laws where $a, b, \dots m, n, \dots$ always denote elements of N , therefore numbers, $A, B, C, \dots$ parts of N , $a', b', \dots m', n', \dots A', B', C', \dots$ the corresponding transforms, which are produced by the order-setting transformation ϕ and are always elements or parts of N ; the transform n' of a number n is also called the number *following* n .

74. Theorem. Every number n by (45) is contained in its chain n_0 and by (53) the condition $n \mathfrak{Z} m_0$ is equivalent to $n_0 \mathfrak{Z} m_0$.

75. Theorem. By (57) $n'_0 = (n_0)' = (n')_0$.

76. Theorem. By (46) $n'_0 \mathfrak{Z} n_0$.

77. Theorem. By (58) $n_0 = \mathfrak{M}(n, n'_0)$.

78. Theorem. $N = \mathfrak{M}(1, N')$, hence every number different from the base-number 1 is element of N' , i.e., transform of a number.

The proof follows from (77) and (71).

79. Theorem. N is the only number-chain containing the base-number 1.

Proof. For if 1 is element of a number-chain K , then by (47) the associated chain $N \mathfrak{Z} K$, hence $N = K$, because it is self-evident that $K \mathfrak{Z} N$. $\square$

80. Theorem of complete induction (inference from n to n'). In order to show that a theorem holds for all numbers n of a chain m_0 , it is sufficient to show,

ρ . that it holds for $n = m$, and

σ . that from the validity of the theorem for a number n of the chain m_0 its validity for the following number n' always follows.

This results immediately from the more general theorem (59) or (60). The most frequently occurring case is where $m = 1$ and therefore m_0 is the complete number-series N .

14. Greater and less numbers

81. Theorem. Every number n is different from the following number n' .

Proof by complete induction (80):

ρ . The theorem is true for the number $n = 1$, because it is not contained in N' (71), while the following number $1'$ as transform of the number 1 contained in N is element of N' .

σ . If the theorem is true for a number n and we put the following number $n' = p$, then is n different from p , whence by (26) on account of the similarity (71) of the order-setting transformation ϕ it follows that n' , and therefore p , is different from p' . Hence the theorem holds also for the number p following n , which was to be proved.

82. Theorem. In the transform-chain n'_0 of a number n by (74), (75) is contained its transform n' , but not the number n itself.

Proof by complete induction (80):

ρ . The theorem is true for $n = 1$, because $1'_0 = N'$, and because by (71) the base-number 1 is not contained in N' .

σ . If the theorem is true for a number n , and we again put $n' = p$, then is n not contained in p_0 , therefore is it different from every number q contained in p_0 , whence by reason of the similarity of ϕ it follows that n' , and therefore p , is different from every number q' contained in p'_0 , and is hence not contained in p'_0 . Therefore the theorem holds also for the number p following n , which was to be proved.

83. Theorem. The transform-chain n'_0 is proper part of the chain n_0 .

14. Greater and less numbers

The proof follows from (76), (74), (82).

84. Theorem. From $m_0 = n_0$ it follows that $m = n$.

Proof. Since by (74) m is contained in m_0 , and

$$m_0 = n_0 = \mathfrak{M}(n, n'_0)$$

by (77), then if the theorem were false and hence m different from n , m would be contained in the chain n'_0 , hence by (74) also $m_0 \mathfrak{Z} n'_0$, i.e., $n_0 \mathfrak{Z} n'_0$; but this contradicts theorem (83). Hence our theorem is established. $\square$

85. Theorem. If the number n is not contained in the number-chain K , then is $K \mathfrak{Z} n'_0$.

Proof by complete induction (80):

ρ . By (78) the theorem is true for $n = 1$.

σ . If the theorem is true for a number n , then is it also true for the following number, $p = n'$; for if p is not contained in the number-chain K , then by (40) n also cannot be contained in K and hence by our hypothesis $K \mathfrak{Z} n'_0$; now since by (77) $n'_0 = p_0 = \mathfrak{M}(p, p'_0)$, hence $K \mathfrak{Z} \mathfrak{M}(p, p'_0)$ and p is not contained in K , then must $K \mathfrak{Z} p'_0$, which was to be proved.

86. Theorem. If the number n is not contained in the number-chain K , but its transform n' is, then $K = n'_0$.

Proof. Since n is not contained in K , then by (85) $K \mathfrak{Z} n'_0$, and since $n' \mathfrak{Z} K$, then by (47) is also $n'_0 \mathfrak{Z} K$, and hence $K = n'_0$, which was to be proved. $\square$

87. Theorem. In every number-chain K there exists one, and by (84) only one, number k , whose chain $k_0 = K$.

Proof. If the base-number 1 is contained in K , then by (79) $K = N = 1_0$. In the opposite case let Z be the system of all numbers not contained in K ; since the base-number 1 is contained in Z , but Z is only a proper part of the number-series N , then by (79) Z cannot be a chain, i.e., Z' cannot be part of Z ; hence there exists in Z a number n , whose transform n' is not contained in Z , and is therefore certainly contained in K ; since further n is contained in Z , and therefore not in K , then by (86) $K = n'_0$, and hence $k = n'$, which was to be proved. $\square$

88. Theorem. If m, n are different numbers then by (83), (84) one and only one of the chains m_0, n_0 is proper part of the other and either $n_0 \supset m'_0$ or $m_0 \supset n'_0$.

Proof. If n is contained in m_0 , and hence by (74) also $n_0 \supset m_0$, then m can not be contained in the chain n_0 (because otherwise by (74) we should have $m_0 \supset n_0$, therefore $m_0 = n_0$, and hence by (84) also $m = n$) and thence it follows by (85) that $n_0 \supset m'_0$. In the contrary case, when n is not contained in the chain m_0 , we must have by (85) $m_0 \supset n'_0$, which was to be proved. $\square$

89. Definition. The number m is said to be *less* than the number n and at the same time n *greater* than m , in symbols

$$m < n, \quad n > m,$$

when the condition

$$n_0 \supset m'_0$$

is fulfilled, which by (74) may also be expressed

$$n \supset m'_0.$$

14. Greater and less numbers

90. Theorem. If m, n are any numbers, then always one and only one of the following cases λ, μ, ν occurs:

$$\begin{aligned}\lambda. \quad & m = n, \quad n = m, \text{ i.e., } \quad m_0 = n_0 \\ \mu. \quad & m < n, \quad n > m, \text{ i.e., } \quad n_0 \mathfrak{Z} m'_0 \\ \nu. \quad & m > n, \quad n < m, \text{ i.e., } \quad m_0 \mathfrak{Z} n'_0.\end{aligned}$$

Proof. For if λ occurs (84) then can neither μ nor ν occur because by (83) we never have $n_0 \mathfrak{Z} n'_0$. But if λ does not occur then by (88) one and only one of the cases μ, ν occurs, which was to be proved. $\square$

91. Theorem. $n < n'$.

Proof. For the condition for the case ν in (90) is fulfilled by $m = n'$. $\square$

92. Definition. To express that m is either $= n$ or $< n$, hence not $> n$ (90) we use the symbols

$$m \leq n \text{ or also } n \geq m$$

and we say m is *at most equal* to n , and n is *at least equal* to m .

93. Theorem. Each of the conditions

$$m \leq n, \quad m < n', \quad n_0 \mathfrak{Z} m_0$$

is equivalent to each of the others.

Proof. For if $m \leq n$, then from λ, μ in (90) we always have $n_0 \mathfrak{Z} m_0$, because by (76) $m'_0 \mathfrak{Z} m_0$. Conversely, if $n_0 \mathfrak{Z} m_0$, and therefore by (74) also $n \mathfrak{Z} m_0$, it follows from $m_0 = \mathfrak{M}(m, m'_0)$ that either $n = m$, or $n \mathfrak{Z} m'_0$, i.e., $n > m$. Hence the condition $m \leq n$ is equivalent to $n_0 \mathfrak{Z} m_0$. Besides it follows from (22), (27), (75) that this condition $n_0 \mathfrak{Z} m_0$ is again equivalent to $n'_0 \mathfrak{Z} m'_0$, i.e., by μ in (90) to $m < n'$, which was to be proved. $\square$

94. Theorem. Each of the conditions

$$m' \leq n, \quad m' < n', \quad m < n$$

is equivalent to each of the others.

The proof follows immediately from (93), if we replace in it m by m' , and from μ in (90).

95. Theorem. If $l < m$ and $m \leq n$ or if $l \leq m$, and $m < n$, then is $l < n$. But if $l \leq m$ and $m \leq n$, then is $l \leq n$.

Proof. For from the corresponding conditions (89), (93) $m_0 \mathfrak{Z} l'_0$ and $n_0 \mathfrak{Z} m_0$, we have by (7) $n_0 \mathfrak{Z} l'_0$ and the same thing comes also from the conditions $m_0 \mathfrak{Z} l_0$ and $n_0 \mathfrak{Z} m'_0$, because in consequence of the former we have also $m'_0 \mathfrak{Z} l'_0$. Finally from $m_0 \mathfrak{Z} l_0$ and $n_0 \mathfrak{Z} m_0$ we have also $n_0 \mathfrak{Z} l_0$ which was to be proved. $\square$

96. Theorem. In every part T of N there exists one and only one *least* number k , i.e., a number k which is less than every other number contained in T . If T consists of a single number, then is it also the least number in T .

Proof. Since T_0 is a chain (44), then by (87) there exists one number k whose chain $k_0 = T_0$. Since from this it follows by (45), (77) that $T \mathfrak{Z} \mathfrak{M}(k, k'_0)$, then first must k itself be contained in T (because otherwise $T \mathfrak{Z} k'_0$, hence by (47) also $T_0 \mathfrak{Z} k'_0$, i.e., $k_0 \mathfrak{Z} k'_0$, which by (83) is impossible), and besides every number of the system T , different from k , must be contained in k'_0 , i.e., be $> k$ (89), whence at once from (90) it follows that there exists in T one and only one least number, which was to be proved. $\square$

97. Theorem. The least number of the chain n_0 is n , and the base-number 1 is the least of all numbers.

14. Greater and less numbers

Proof. For by (74), (93) the condition $m \mathfrak{Z} n_0$ is equivalent to $m \geq n$. Or our theorem also follows immediately from the proof of the preceding theorem, because if in that we assume $T = n_0$, evidently $k = n$ (51). $\square$

98. Definition. If n is any number, then will we denote by Z_n the system of all numbers that are *not greater* than n , and hence *not* contained in n'_0 . The condition

$$m \mathfrak{Z} Z_n$$

by (92), (93) is obviously equivalent to each of the following conditions:

$$m \leq n, \quad m < n', \quad n_0 \mathfrak{Z} m_0.$$

99. Theorem. $1 \mathfrak{Z} Z_n$ and $n \mathfrak{Z} Z_n$.

The proof follows from (98) or from (71) and (82).

100. Theorem. Each of the conditions equivalent by (98)

$$m \mathfrak{Z} Z_n, \quad m \leq n, \quad m < n', \quad n_0 \mathfrak{Z} m_0$$

is also equivalent to the condition

$$Z_m \mathfrak{Z} Z_n.$$

Proof. For if $m \mathfrak{Z} Z_n$, and hence $m \leq n$, and if $l \mathfrak{Z} Z_m$, and hence $l \leq m$, then by (95) also $l \leq n$, i.e., $l \mathfrak{Z} Z_n$; if therefore $m \mathfrak{Z} Z_n$, then is every element l of the system Z_m also element of Z_n , i.e., $Z_m \mathfrak{Z} Z_n$. Conversely, if $Z_m \mathfrak{Z} Z_n$, then by (7) must also $m \mathfrak{Z} Z_n$, because by (99) $m \mathfrak{Z} Z_m$, which was to be proved. $\square$

101. Theorem. The conditions for the cases λ , μ , ν in (90) may also be put in the following form:

$$\begin{array}{lll} \lambda. & m = n, & n = m, \quad Z_m = Z_n \\ \mu. & m < n, & n > m, \quad Z_{m'} \mathfrak{Z} Z_n \\ \nu. & m > n, & n < m, \quad Z_{n'} \mathfrak{Z} Z_m \end{array}$$

The proof follows immediately from (90) if we observe that by (100) the conditions $n_0 \mathfrak{Z} m_0$ and $Z_m \mathfrak{Z} Z_n$ are equivalent.

102. Theorem. $Z_1 = 1$.

Proof. For by (99) the base-number 1 is contained in Z_1 , while by (78) every number different from 1 is contained in $1'_0$, hence by (98) not in Z_1 , which was to be proved. $\square$

103. Theorem. By (98) $N = \mathfrak{M}(Z_n, n'_0)$.

104. Theorem. $n = \mathfrak{G}(Z_n, n_0)$, i.e., n is the only common element of the system Z_n and n_0 .

Proof. From (99) and (74) it follows that n is contained in Z_n and n_0 ; but every element of the chain n_0 , different from n by (77) is contained in n'_0 , and hence by (98) not in Z_n , which was to be proved. $\square$

105. Theorem. By (91), (98) the number n' is not contained in Z_n .

106. Theorem. If $m < n$, then is Z_m proper part of Z_n and conversely.

Proof. If $m < n$, then by (100) $Z_m \mathfrak{Z} Z_n$, and since the number n , by (99) contained in Z_n , can by (98) not be contained in Z_m because $n > m$, therefore Z_m is proper part of Z_n . Conversely if Z_m is proper part of Z_n then by (100) $m \leq n$, and since m cannot be $= n$, because otherwise $Z_m = Z_n$, we must have $m < n$, which was to be proved. $\square$

14. Greater and less numbers

107. Theorem. Z_n is proper part of $Z_{n'}$.

The proof follows from (106), because by (91) $n < n'$.

108. Theorem. $Z_{n'} = \mathfrak{M}(Z_n, n')$.

Proof. For every number contained in $Z_{n'}$, by (98) is $\leq n'$, hence either $= n'$ or $< n'$, and therefore by (98) element of Z_n . Therefore certainly $Z_{n'} \supset \mathfrak{M}(Z_n, n')$. Since conversely by (107) $Z_n \supset Z_{n'}$ and by (99) $n' \supset Z_{n'}$, then by (10) we have

$$\mathfrak{M}(Z_n, n') \supset Z_{n'},$$

whence our theorem follows by (5). □

109. Theorem. The transform Z'_n of the system Z_n is proper part of the system $Z_{n'}$.

Proof. For every number contained in Z'_n is the transform m' of a number m contained in Z_n , and since $m \leq n$, and hence by (94) $m' \leq n'$, we have by (98) $Z'_n \supset Z_{n'}$. Since further the number 1 by (99) is contained in $Z_{n'}$, but by (71) is not contained in the transform Z'_n , then is Z'_n proper part of $Z_{n'}$, which was to be proved. □

110. Theorem. $Z_{n'} = \mathfrak{M}(1, Z'_n)$.

Proof. Every number of the system $Z_{n'}$ different from 1 by (78) is the transform m' of a number m and this must be $\leq n$, and hence by (98) contained in Z_n (because otherwise $m > n$, hence by (94) also $m' > n'$ and consequently by (98) m' would not be contained in $Z_{n'}$); but from $m \supset Z_n$ we have $m' \supset Z'_n$, and hence certainly

$$Z_{n'} \supset \mathfrak{M}(1, Z'_n).$$

Since conversely by (99) $1 \mathfrak{Z} Z_{n'}$, and by (109) $Z'_n \mathfrak{Z} Z_{n'}$, then by (10) we have $\mathfrak{M}(1, Z'_n) \mathfrak{Z} Z_{n'}$ and hence our theorem follows by (5). $\square$

111. Definition. If in a system E of numbers there exists an element g , which is greater than every other number contained in E , then g is said to be the *greatest* number of the system E , and by (90) there can evidently be only one such greatest number in E . If a system consists of a single number, then is this number itself the greatest number of the system.

112. Theorem. By (98) n is the greatest number of the system Z_n .

113. Theorem. If there exists in E a greatest number g , then is $E \mathfrak{Z} Z_g$.

Proof. For every number contained in E is $\leq g$, and hence by (98) contained in Z_g , which was to be proved. $\square$

114. Theorem. If E is part of a system Z_n , or what amounts to the same thing, there exists a number n such that all numbers contained in E are $\leq n$, then E possesses a greatest number g .

Proof. The system of all numbers p satisfying the condition $E \mathfrak{Z} Z_p$ —and by our hypothesis such numbers exist—is a chain (37), because by (107), (7) it follows also that $E \mathfrak{Z} Z_{p'}$, and hence by (87) $= g_0$, where g signifies the least of these numbers (96), (97). Hence also $E \mathfrak{Z} Z_g$, therefore by (98) every number contained in E is $\leq g$, and we have only to show that the number g is itself contained in E . This is immediately obvious if $g = 1$, for then by (102) Z_g , and consequently also E consists of the single number 1. But if g is different from 1 and consequently by (78) the transform f' of a number f , then by (108) is $E \mathfrak{Z} \mathfrak{M}(Z_f, g)$; if therefore g were not contained in E , then would $E \mathfrak{Z} Z_f$, and there would consequently be among the numbers p a number f by (91) $< g$, which is contrary to what precedes; hence g is contained in E , which was to be proved. $\square$

14. Greater and less numbers

115. Definition. If $l < m$ and $m < n$ we say the number m *lies between* l and n (also between n and l).

116. Theorem. There exists no number lying between n and n' .

Proof. For as soon as $m < n'$, and hence by (93) $m \leq n$, then by (90) we cannot have $n < m$, which was to be proved. $\square$

117. Theorem. If t is a number in T , but not the least (96), then there exists in T one and only one *next less* number s , i.e., a number s such that $s < t$, and that there exists in T no number lying between s and t . Similarly, if t is not the greatest number in T (111) there always exists in T one and only one *next greater* number u , i.e., a number u such that $t < u$, and that there exists in T no number lying between t and u . At the same time in T t is next greater than s and next less than u .

Proof. If t is not the least number in T , then let E be the system of all those numbers of T that are $< t$; then by (98) $E \mathfrak{Z} Z_t$, and hence by (114) there exists in E a greatest number s obviously possessing the properties stated in the theorem, and also it is the only such number. If further t is not the greatest number in T , then by (96) there certainly exists among all the numbers of T , that are $> t$, a least number u , which and which alone possesses the properties stated in the theorem. In like manner the correctness of the last part of the theorem is obvious. $\square$

118. Theorem. In N the number n' is next greater than n , and n next less than n' .

The proof follows from (116), (117).

15. Finite and infinite parts of the number-series

119. Theorem. Every system Z_n in (98) is finite.

Proof by complete induction (80).

ρ . By (65), (102) the theorem is true for $n = 1$.

σ . If Z_n is finite, then from (108) and (70) it follows that $Z_{n'}$ is also finite, which was to be proved.

120. Theorem. If m, n are different numbers, then are Z_m, Z_n dissimilar systems.

Proof. By reason of the symmetry we may by (90) assume that $m < n$; then by (106) Z_m is proper part of Z_n , and since by (119) Z_n is finite, then by (64) Z_m and Z_n cannot be similar, which was to be proved. $\square$

121. Theorem. Every part E of the number-series N , which possesses a greatest number (111), is finite.

The proof follows from (113), (119), (68).

122. Theorem. Every part U of the number-series N , which possesses no greatest number, is simply infinite (71).

15. *Finite and infinite parts of the number-series*

Proof. If u is any number in U , there exists in U by (117) one and only one next greater number than u , which we will denote by $\psi(u)$ and regard as transform of u . The thus perfectly determined transformation ψ of the system U has obviously the property

$$\alpha. \quad \psi(U) \supset U,$$

i.e., U transformed in itself by ψ . If further u, v are different numbers in U , then by symmetry we may by (90) assume that $u < v$; thus by (117) it follows from the definition of ψ that $\psi(u) \leq v$ and $v < \psi(v)$, and hence by (95) $\psi(u) < \psi(v)$; therefore by (90) the transforms $\psi(u), \psi(v)$ are different, i.e.,

$\delta.$ the transformation ψ is similar.

Further, if u_1 denotes the least number (96) of the system U , then every number u contained in U is $\geq u_1$, and since generally $u < \psi(u)$, then by (95) $u_1 < \psi(u)$, and therefore by (90) u_1 is different from $\psi(u)$, i.e.,

$\gamma.$ the element u_1 of U is not contained in $\psi(U)$.

Therefore $\psi(U)$ is proper part of U and hence by (64) U is an infinite system. If then in agreement with (44) we denote by $\psi_0(V)$, when V is any part of U , the chain of V corresponding to the transformation ψ , we wish to show finally that

$$\beta. \quad U = \psi_0(u_1).$$

In fact, since every such chain $\psi_0(V)$ by reason of its definition (44) is a part of the system U transformed in itself by ψ , then evidently is $\psi_0(u_1) \supset U$; conversely it is first of all obvious from (45) that the element u_1 contained in U is certainly contained in $\psi_0(u_1)$; but if we assume that there exist elements of U , that are not contained in $\psi_0(u_1)$, then must there be among them by (96) a least number w , and since by what precedes this is different from the least number u_1 of the system U , then by (117) must there exist in U also a number v which is next less than w , whence it follows at once that $w = \psi(v)$; since therefore $v < w$, then must v by reason of the definition of w certainly be contained in $\psi_0(u_1)$; but from this by (55) it follows that also $\psi(v)$, and hence w must be contained in $\psi_0(u_1)$, and since this is contrary

to the definition of w , our foregoing hypothesis is inadmissible; therefore $U \not\leq \psi_0(u_1)$ and hence also $U = \psi_0(u_1)$ as stated. From $\alpha, \beta, \gamma, \delta$ it then follows by (71) that U is a simply infinite system set in order by ψ , which was to be proved. $\square$

123. Theorem. In consequence of (121), (122) any part T of the number-series N is finite or simply infinite, according as a greatest number exists or does not exist in T .

16. Definition of a transformation of the number-series by induction

124. In what follows we denote numbers by small Italics and retain throughout all symbols of the previous Section 13 to Section 15, while Ω designates an arbitrary system whose elements are not necessarily contained in N .

125. Theorem. If there is given an arbitrary (similar or dissimilar) transformation θ of a system Ω in itself, and besides a determinate element ω in Ω , then to every number n corresponds one transformation ψ_n and one only of the associated number-system Z_n explained in (98), which satisfies the conditions:¹

I. $\psi_n Z_n \ni \Omega$

II. $\psi_n(1) = \omega$

III. $\psi_n(t') = \theta\psi_n(t)$, if $t < n$, where the symbol $\theta\psi_n$ has the meaning given in (25).

Proof by complete induction (80).

ρ . The theorem is true for $n = 1$. In this case indeed by (102) the system Z_n consists of the single number 1, and the transformation ψ_1 is therefore completely defined by II alone so that I is fulfilled while III drops out entirely.

¹For clearness here and in the following theorem (126) I have especially mentioned condition I., although properly it is a consequence of II. and III.

16. Definition of a transformation of the number-series by induction

σ . If the theorem is true for a number n then we show that it is also true for the following number $p = n'$, and we begin by proving that there can be only a single corresponding transformation ψ_p of the system Z_p . In fact, if a transformation ψ_p satisfies the conditions

$$\text{I'. } \psi_p(Z_p) \ni \Omega$$

$$\text{II'. } \psi_p(1) = \omega$$

III'. $\psi_p(m') = \theta\psi_p(m)$, when $m < p$, then there is also contained in it by (21), because $Z_n \ni Z_p$ (107) a transformation of Z_n which obviously satisfies the same conditions I, II, III as ψ_n , and therefore coincides throughout with ψ_n ; for all numbers contained in Z_n , and hence (98) for all numbers m which are $< p$, i.e., $\leq n$, must therefore

$$\psi_p(m) = \psi_n(m) \tag{m}$$

whence there follows, as a special case,

$$\psi_p(n) = \psi_n(n); \tag{n}$$

since further by (105), (108) p is the only number of the system Z_p not contained in Z_n , and since by III' and (n) we must also have

$$\psi_p(p) = \theta\psi_n(n) \tag{p}$$

there follows the correctness of our foregoing statement that there can be only one transformation ψ_p of the system Z_p satisfying the conditions I, II', III', because by the conditions (m) and (p) just derived ψ_p is completely reduced to ψ_n . We have next to show conversely that this transformation ψ_p of the system Z_p completely determined by (m) and (p) actually satisfies the conditions I', II', III'. Obviously I' follows from (m) and (p) with reference to I, and because $\theta(\Omega) \ni \Omega$. Similarly II' follows from (m) and II, since by (99) the number 1 is contained in Z_n . The correctness of III' follows first for those numbers m which are $< n$ from (m) and III, and for the

single number $m = n$ yet remaining it results from (p) and (n) . Thus it is completely established that from the validity of our theorem for the number n always follows its validity for the following number p , which was to be proved.

126. Theorem of the definition by induction. If there is given an arbitrary (similar or dissimilar) transformation θ of a system Ω in itself, and besides a determinate element ω in Ω , then there exists one and only one transformation ψ of the number-series N , which satisfies the conditions

I. $\psi(N) \ni \Omega$

II. $\psi(1) = \omega$

III. $\psi(n') = \theta\psi(n)$, where n represents every number.

Proof. Since, if there actually exists such a transformation ψ , there is contained in it by (21) a transformation ψ_n , of the system Z_n , which satisfies the conditions I, II, III stated in (125), then because there exists one and only one such transformation ψ_n must necessarily

$$\psi(n) = \psi_n(n). \quad (n)$$

Since thus ψ is completely determined it follows also that there can exist only one such transformation ψ (see the closing remark in (130)). That conversely the transformation ψ determined by (n) also satisfies our conditions I, II, III, follows easily from (n) with reference to the properties I, II and (p) shown in (125), which was to be proved. $\square$

127. Theorem. Under the hypotheses made in the foregoing theorem,

$$\psi(T') = \theta\psi(T),$$

where T denotes any part of the number-series N .

16. Definition of a transformation of the number-series by induction

Proof. For if t denotes every number of the system T , then $\psi(T')$ consists of all elements $\psi(t')$, and $\theta\psi(T)$ of all elements $\theta\psi(t)$; hence our theorem follows because by III in (126) $\psi(t') = \theta\psi(t)$. $\square$

128. Theorem. If we maintain the same hypotheses and denote by θ_0 the chains (44) which correspond to the transformation θ of the system Ω in itself, then is

$$\psi(N) = \theta_0(\omega).$$

Proof. We show first by complete induction (80) that

$$\psi(N) \supset \theta_0(\omega),$$

i.e., that every transform $\psi(n)$ is also element of $\theta_0(\omega)$. In fact,

ρ . this theorem is true for $n = 1$, because by (126, II) $\psi(1) = \omega$, and because by (45) $\omega \supset \theta_0(\omega)$.

σ . If the theorem is true for a number n , and hence $\psi(n) \supset \theta_0(\omega)$, then by (55) also $\theta(\psi(n)) \supset \theta_0(\omega)$, i.e., by (126, III) $\psi(n') \supset \theta_0(\omega)$, hence the theorem is true for the following number n' , which was to be proved.

In order further to show that every element ν of the chain $\theta_0(\omega)$ is contained in $\psi(N)$, therefore that

$$\theta_0(\omega) \supset \psi(N)$$

we likewise apply complete induction, i.e., theorem (59) transferred to Ω and the transformation θ . In fact,

ρ . the element $\omega = \psi(1)$, and hence is contained in $\psi(N)$.

σ . If ν is a common element of the chain $\theta_0(\omega)$ and the system $\psi(N)$, then $\nu = \psi(n)$, where n denotes a number, and by (126, III) we get $\theta(\nu) =$

$\theta\psi(n) = \psi(n')$, and therefore $\theta(\nu)$ is contained in $\psi(N)$, which was to be proved.

From the theorems just established, $\psi(N) \supset \theta_0(\omega)$ and $\theta_0(\omega) \supset \psi(N)$, we get by (5) $\psi(N) = \theta_0(\omega)$, which was to be proved. $\square$

129. Theorem. Under the same hypotheses we have generally:

$$\psi(n_0) = \theta_0(\psi(n)).$$

By complete induction (80). For

ρ . By (128) the theorem holds for $n = 1$, since $1_0 = N$ and $\psi(1) = \omega$.

σ . If the theorem is true for a number n , then

$$\theta(\psi(n_0)) = \theta(\theta_0(\psi(n)));$$

since by (127), (75)

$$\theta(\psi(n_0)) = \psi(n'_0),$$

and by (57), (126, III)

$$\theta(\theta_0(\psi(n))) = \theta_0(\theta(\psi(n))) = \theta_0(\psi(n')),$$

we get

$$\psi(n'_0) = \theta_0(\psi(n')),$$

i.e., the theorem is true for the number n' following n , which was to be proved.

16. Definition of a transformation of the number-series by induction

130. Remark. Before we pass to the most important applications of the theorem of definition by induction proved in (126), (Section 17 — Section 21), it is worth while to call attention to a circumstance by which it is essentially distinguished from the theorem of demonstration by induction proved in (80) or rather in (59), (60), however close may seem the relation between the former and the latter. For while the theorem (59) is true quite generally for every chain A_0 where A is any part of a system S transformed in itself by any transformation ϕ (Section 11), the case is quite different with the theorem (126), which declares only the existence of a consistent (or one-to-one) transformation ψ of the simply infinite system 1_0 . If in the latter theorem (still maintaining the hypotheses regarding Ω and θ) we replace the number-series 1_0 by an arbitrary chain A_0 out of such a system S , and define a transformation ψ of A_0 in Ω in a manner analogous to that in (126, II, III) by assuming that

ρ . to every element a of A there is to correspond a determinate element $\psi(a)$ selected from Ω , and

σ . for every element n contained in A_0 and its transform $n' = \phi(n)$, the condition $\psi(n') = \theta\psi(n)$ is to hold, then would the case very frequently occur that such a transformation ψ does not exist, since these conditions ρ , σ may prove incompatible, even though the freedom of choice contained in ρ be restricted at the outset to conform to the condition σ . An example will be sufficient to convince one of this. If the system S consisting of the different elements a and b is so transformed in itself by ϕ that $a' = b$, $b' = a$, then obviously $a_0 = b_0 = S$; suppose further the system Ω consisting of the different elements α , β , and γ be so transformed in itself by θ that $\theta(\alpha) = \beta$, $\theta(\beta) = \gamma$, $\theta(\gamma) = \alpha$; if we now demand a transformation ψ of a_0 in Ω such that $\psi(a) = \alpha$, and that besides for every element n contained in a_0 always $\psi(n') = \theta\psi(n)$, we meet a contradiction; since for $n = a$, we get $\psi(b) = \theta(\alpha) = \beta$, and hence for $n = b$, we must have $\psi(a) = \theta(\beta) = \gamma$, while we had assumed $\psi(a) = \alpha$.

But if there exists a transformation ψ of A_0 in Ω , which satisfies the foregoing conditions ρ , σ without contradiction, then from (60) it follows easily that it is completely determined; for if the transformation χ satisfies the

same conditions, then we have, generally, $\chi(n) = \psi(n)$, since by ρ this theorem is true for all elements $n = a$ contained in A , and since if it is true for an element n of A_0 it must by σ be true also for its transform n' .

131. In order to bring out clearly the import of our theorem (126), we will here insert a consideration which is useful for other investigations also, e.g., for the so-called group-theory.

We consider a system Ω , whose elements allow a certain combination such that from an element ν by the effect of an element ω , there always results again a determinate element of the same system Ω , which may be denoted by $\omega \cdot \nu$ or $\omega\nu$, and in general is to be distinguished from $\nu\omega$. We can also consider this in such a way that to every determinate element ω , there corresponds a determinate transformation of the system Ω in itself (to be denoted by $\dot{\omega}$), in so far as every element ν furnishes the determinate transform $\dot{\omega}(\nu) = \omega\nu$. If to this system Ω and its element ω we apply theorem (126), designating by $\dot{\omega}$ the transformation there denoted by θ , then there corresponds to every number n a determinate element $\psi(n)$ contained in Ω , which may now be denoted by the symbol ω^n and sometimes called the n th power of ω ; this notion is completely defined by the conditions imposed upon it

$$\text{II. } \omega^1 = \omega$$

$$\text{III. } \omega^{n'} = \omega \omega^n,$$

and its existence is established by the proof of theorem (126).

If the foregoing combination of the elements is further so qualified that for arbitrary elements μ, ν, ω , we always have $\omega(\nu\mu) = \omega\nu(\mu)$, then are true also the theorems

$$\omega^{n'} = \omega^n \omega, \quad \omega^m \omega^n = \omega^n \omega^m,$$

16. Definition of a transformation of the number-series by induction

whose proofs can easily be effected by complete induction and may be left to the reader.

The foregoing general consideration may be immediately applied to the following example. If S is a system of arbitrary elements, and Ω the associated system whose elements are all the transformations ν of S in itself (36), then by (25) can these elements be continually compounded, since $\nu(S) \supset S$, and the transformation $\omega\nu$ compounded out of such transformations ν and ω is itself again an element of Ω . Then are also all elements ω^n transformations of S in itself, and we say they arise by repetition of the transformation ω . We will now call attention to a simple connection existing between this notion and the notion of the chain $\omega_0(A)$ defined in (44), where A again denotes any part of S . If for brevity we denote by A_n the transform $\omega^n(A)$ produced by the transformation ω^n , then from III and (25) it follows that $\omega(A_n) = A_{n'}$. Hence it is easily shown by complete induction (80) that all these systems A_n are parts of the chain $\omega_0(A)$; for

ρ . by (50) this statement is true for $n = 1$, and

σ . if it is true for a number n , then from (55) and from $A_{n'} = \omega(A_n)$ it follows that it is also true for the following number n' , which was to be proved. Since further by (45) $A \supset \omega_0(A)$, then from (10) it results that the system K compounded out of A and all transforms A_n is part of $\omega_0(A)$. Conversely, since by (23) $\omega(K)$ is compounded out of $\omega(A) = A_1$ and all systems $\omega(A_n) = A_{n'}$, therefore by (78) out of all systems A_n , which by (9) are parts of K , then by (10) is $\omega(K) \supset K$, i.e., K is a chain (37), and since by (9) $A \supset K$, then by (47) it follows also that that $\omega_0(A) \supset K$. Therefore $\omega_0(A) = K$, i.e., the following theorem holds: If ω is a transformation of a system S in itself, and A any part of S , then is the chain of A corresponding to the transformation ω compounded out of A and all the transforms $\omega^n(A)$ resulting from repetitions of ω . We advise the reader with this conception of a chain to return to the earlier theorems (57), (58).

17. The class of simply infinite systems

132. Theorem. All simply infinite systems are similar to the number-series N and consequently by (33) also to one another.

Proof. Let the simply infinite system Ω be set in order (71) by the transformation θ , and let ω be the base-element of Ω thus resulting; if we again denote by θ_0 the chains corresponding to the transformation θ (44), then by (71) is the following true:

$\alpha.$ $\theta(\Omega) \supset \Omega$.

$\beta.$ $\Omega = \theta_0(\omega)$.

$\gamma.$ ω is not contained in $\theta(\Omega)$.

$\delta.$ The transformation θ is similar.

If then ψ denotes the transformation of the number-series N defined in (126), then from β and (128) we get first

$$\psi(N) = \Omega,$$

and hence we have only yet to show that ψ is a similar transformation, i.e., (26) that to different numbers m, n correspond different transforms $\psi(m), \psi(n)$. On account of the symmetry we may by ((90)) assume that $m > n$, hence $m \supset n'_0$, and the theorem to prove comes to this that $\psi(n)$ is not

17. The class of simply infinite systems

contained in $\psi(n'_0)$, and hence by (127) is not contained in $\theta\psi(n_0)$. This we establish for every number n by complete induction (80). In fact,

ρ . this theorem is true by γ for $n = 1$, since $\psi(1) = \omega$ and $\psi(1_0) = \psi(N) = \Omega$.

σ . If the theorem is true for a number n , then is it also true for the following number n' ; for if $\psi(n')$, i.e., $\theta\psi(n)$, were contained in $\theta\psi(n'_0)$, then by δ and (27), $\psi(n)$ would also be contained in $\psi(n'_0)$ while our hypothesis states just the opposite; which was to be proved. $\square$

133. Theorem. Every system which is similar to a simply infinite system and therefore by (132), (33) to the number-series N is simply infinite.

Proof. If Ω is a system similar to the number-series N , then by (32) there exists a similar transformation ψ of N such that

$$\text{I. } \psi(N) = \Omega;$$

then we put

$$\text{II. } \psi(1) = \omega.$$

If we denote, as in (26), by $\bar{\psi}$ the inverse, likewise similar transformation of Ω , then to every element ν of Ω there corresponds a determinate number $\bar{\psi}(\nu) = n$, viz., that number whose transform $\psi(n) = \nu$. Since to this number n there corresponds a determinate following number $\phi(n) = n'$, and to this again a determinate element $\psi(n')$ in Ω there belongs to every element ν of the system Ω a determinate element $\psi(n')$ of that system which as transform of ν we shall designate by $\theta(\nu)$. Thus a transformation θ of Ω in itself is completely determined,¹ and in order to prove our theorem we will show that by $\theta\Omega$ is set in order (71) as a simply infinite system, i.e., that the conditions α , β , γ , δ stated in the proof of (132) are all fulfilled.

¹Evidently θ is the transformation $\psi\phi\bar{\psi}$ compounded by (25) out of $\bar{\psi}$, ϕ , ψ .

First α is immediately obvious from the definition of θ . Since further to every number n corresponds an element $\nu = \phi(n)$, for which $\theta(\nu) = \psi(n')$, we have generally,

$$\text{III. } \psi(n') = \theta\psi(n),$$

and thence in connection with I, II, α it results that the transformations θ , ψ fulfill all the conditions of theorem (126); therefore β follows from (128) and I. Further by (127) and I

$$\psi(N') = \theta\psi(N) = \theta(\Omega),$$

and thence in combination with II and the similarity of the transformation ψ follows γ , because otherwise $\psi(1)$ must be contained in $\psi(N')$, hence by (27) the number 1 in N' , which by (71, γ) is not the case. If finally μ, ν denote elements of Ω and m, n the corresponding numbers whose transforms are $\psi(m) = \mu$, $\psi(n) = \nu$, then from the hypothesis $\theta(\mu) = \theta(\nu)$ it follows by the foregoing that $\psi(m') = \psi(n')$, thence on account of the similarity of ψ, ϕ that $m' = n'$, $m = n$, therefore also $\mu = \nu$; hence also δ is true, which was to be proved. $\square$

134. Remark. By the two preceding theorems (132), (133) all simply infinite systems form a class in the sense of (34). At the same time, with reference to (71), (73) it is clear that every theorem regarding numbers, i.e., regarding the elements n of the simply infinite system N set in order by the transformation ϕ , and indeed every theorem in which we leave entirely out of consideration the special character of the elements n and discuss only such notions as arise from the arrangement ϕ , possesses perfectly general validity for every other simply infinite system Ω set in order by a transformation θ and its elements ν , and that the passage from N to Ω (e.g., also the translation of an arithmetic theorem from one language into another) is effected by the transformation ψ considered in (132), (133), which changes every element n of N into an element ν of Ω , i.e., into $\psi(n)$. This element ν can be called the n th element of Ω and accordingly the number n is itself

17. The class of simply infinite systems

the n th number of the number-series N . The same significance which the transformation ϕ possesses for the laws in the domain N , in so far as every element n is followed by a determinate element $\phi(n) = n'$, is found, after the change effected by ψ , to belong to the transformation θ for the same laws in the domain Ω , in so far as the element $\nu = \psi(n)$ arising from the change of n is followed by the element $\theta(\nu) = \psi(n')$ arising from the change of n' ; we are therefore justified in saying that by $\psi\phi$ is changed into θ , which is symbolically expressed by $\theta = \psi\phi\bar{\psi}$, $\phi = \bar{\psi}\theta\psi$. By these remarks, as I believe, the definition of the notion of numbers given in (73) is fully justified. We now proceed to further applications of theorem (126).

18. Addition of numbers

135. Definition. It is natural to apply the definition set forth in theorem (126) of a transformation ψ of the number-series N , or of the *function* $\psi(n)$ determined by it to the case, where the system there denoted by Ω in which the transform $\psi(N)$ is to be contained, is the number-series N itself, because for this system Ω a transformation θ of Ω in itself already exists, viz., that transformation ϕ by which N is set in order as a simply infinite system (71), (73). Then is also $\Omega = N$, $\theta(n) = \phi(n) = n'$, hence

$$\text{I. } \psi(N) \text{ } \mathfrak{Z} \text{ } N,$$

and it remains in order to determine ψ completely only to select the element ω from Ω , i.e., from N , at pleasure. If we take $\omega = 1$, then evidently ψ becomes the identical transformation (21) of N , because the conditions

$$\psi(1) = 1, \quad \psi(n') = (\psi(n))'$$

are generally satisfied by $\psi(n) = n$. If then we are to produce another transformation ψ of N , then for ω we must select a number m' different from 1, by (78) contained in N , where m itself denotes any number; since the transformation ψ is obviously dependent upon the choice of this number m , we denote the corresponding transform $\psi(n)$ of an arbitrary number n by the symbol $m + n$, and call this number the *sum* which arises from the number m by the *addition* of the number n , or in short the sum of the

18. Addition of numbers

numbers m, n . Therefore by (126) this sum is completely determined by the conditions¹

$$\text{II. } m + 1 = m',$$

$$\text{III. } m + n' = (m + n)'.$$

136. Theorem. $m' + n = m + n'$.

Proof by complete induction (80). For

ρ . the theorem is true for $n = 1$, since by ((135), II)

$$m' + 1 = (m')' = (m + 1)',$$

and by (135, III) $(m + 1)' = m + 1'$.

σ . If the theorem is true for a number n , and we put the following number $n' = p$ then is $m' + n = m + p$, hence also $(m' + n)' = (m + p)'$, whence by (135, III) $m' + p = m + p'$; therefore the theorem is true also for the following number p , which was to be proved.

137. Theorem. $m' + n = (m + n)'$.

The proof follows from (136) and (135, III).

138. Theorem. $1 + n = n'$.

Proof by complete induction (80). For

¹The above definition of addition based immediately upon theorem (126) seems to me to be the simplest. By the aid of the notion developed in (131) we can, however, define the sum $m + n$ by $\phi^n(m)$ or also by $\phi^m(n)$, where ϕ has again the foregoing meaning. In order to show the complete agreement of these definitions with the foregoing, we need by (126) only to show that if $\phi^n(m)$ or $\phi^m(n)$ is denoted by $\psi(n)$, the conditions $\psi(1) = m'$, $\psi(n') = \phi\psi(n)$ are fulfilled which is easily done with the aid of complete induction (80) by the help of (131).

ρ . by (135, II) the theorem is true for $n = 1$.

σ . If the theorem is true for a number n and we put $n' = p$, then $1 + n = p$, therefore also $(1 + n)' = p'$, whence by (135, III) $1 + p = p'$, i.e., the theorem is true also for the following number p , which was to be proved.

139. Theorem. $1 + n = n + 1$.

The proof follows from (138) and (135, II).

140. Theorem. $m + n = n + m$.

Proof by complete induction (80). For

ρ . by (139) the theorem is true for $n = 1$.

σ . If the theorem is true for a number n , then there follows also $(m + n)' = (n + m)'$, i.e., by (135, III) $m + n' = n + m'$, hence by (136) $m + n' = n' + m$; therefore the theorem is also true for the following number n' , which was to be proved.

141. Theorem. $(l + m) + n = l + (m + n)$.

Proof by complete induction (80). For

ρ . the theorem is true for $n = 1$, because by (135, II, III, II) $(l + m) + 1 = (l + m)' = l + m' = l + (m + 1)$.

σ . If the theorem is true for a number n , then there follows also $((l + m) + n)' = (l + (m + n))'$, i.e., by (135, III)

$$(l + m) + n' = l + (m + n)' = l + (m + n'),$$

therefore the theorem is also true for the following number n' , which was to be proved.

142. Theorem. $m + n > m$.

Proof by complete induction (80). For

ρ . by (135, II) and (91) the theorem is true for $n = 1$.

18. Addition of numbers

σ . If the theorem is true for a number n , then by (95) it is also true for the following number n' , because by (135, III) and (91)

$$m + n' = (m + n)' > m + n,$$

which was to be proved.

143. Theorem. The conditions $m > a$ and $m + n > a + n$ are equivalent.

Proof by complete induction (80). For

ρ . by (135, II) and (94) the theorem is true for $n = 1$.

σ . If the theorem is true for a number n , then is it also true for the following number n' , since by (94) the condition $m + n > a + n$ is equivalent to $(m + n)' > (a + n)'$, hence by (135, III) also equivalent to

$$m + n' > a + n',$$

which was to be proved.

144. Theorem. If $m > a$ and $n > b$, then is also

$$m + n > a + b.$$

Proof. For from our hypotheses we have by (143) $m + n > a + n$ and $n + a > b + a$ or, what by (140) is the same, $a + n > a + b$, whence the theorem follows by (95). $\square$

145. Theorem. If $m + n = a + n$ then $m = a$.

Proof. For if m does not $= a$, hence by (90) either $m > a$ or $m < a$, then by (143) respectively $m + n > a + n$ or $m + n < a + n$, therefore by (90) we surely cannot have $m + n = a + n$ which was to be proved. $\square$

146. Theorem. If $l > n$, then there exists one and by (157) only one number m which satisfies the condition $m + n = l$.

Proof by complete induction (80). For

ρ . the theorem is true for $n = 1$. In fact, if $l > 1$, i.e., (89) if l is contained in N' , and hence is the transform m' of a number m , then by (135, II) it follows that $l = m + 1$, which was to be proved.

σ . If the theorem is true for a number n , then we show that it is also true for the following number n' . In fact, if $l > n'$, then by (91), (95) also $l > n$, and hence there exists a number k which satisfies the condition $l = k + n$; since by (138) this is different from 1 (otherwise l would be $= n'$) then by (78) is it the transform m' of a number m , consequently $l = m' + n$, therefore also by (136) $l = m + n'$, which was to be proved.

19. Multiplication of numbers

20. Involution of numbers

21. Number of the elements of a finite system

22. License

THE FULL PROJECT GUTENBERG LICENSE

PLEASE READ THIS BEFORE YOU DISTRIBUTE OR USE THIS WORK

To protect the Project Gutenberg-tm mission of promoting the free distribution of electronic works, by using or distributing this work (or any other work associated in any way with the phrase “Project Gutenberg”), you agree to comply with all the terms of the Full Project Gutenberg-tm License (available with this file or online at www.gutenberg.org/license).

22.1. General Terms of Use and Redistributing Project Gutenberg-tm electronic works

1.A. By reading or using any part of this Project Gutenberg-tm electronic work, you indicate that you have read, understand, agree to and accept all the terms of this license and intellectual property (trademark/copyright) agreement. If you do not agree to abide by all the terms of this agreement, you must cease using and return or destroy all copies of Project Gutenberg-tm electronic works in your possession. If you paid a fee for obtaining a copy of or access to a Project Gutenberg-tm electronic work and you do not agree to be bound by the terms of this agreement, you may obtain a refund from the person or entity to whom you paid the fee as set forth in paragraph 1.E.8.

22. License

1.B. “Project Gutenberg” is a registered trademark. It may only be used on or associated in any way with an electronic work by people who agree to be bound by the terms of this agreement. There are a few things that you can do with most Project Gutenberg-tm electronic works even without complying with the full terms of this agreement. See paragraph 1.C below. There are a lot of things you can do with Project Gutenberg-tm electronic works if you follow the terms of this agreement and help preserve free future access to Project Gutenberg-tm electronic works. See paragraph 1.E below.

1.C. The Project Gutenberg Literary Archive Foundation (“the Foundation” or PGLAF), owns a compilation copyright in the collection of Project Gutenberg-tm electronic works. Nearly all the individual works in the collection are in the public domain in the United States. If an individual work is in the public domain in the United States and you are located in the United States, we do not claim a right to prevent you from copying, distributing, performing, displaying or creating derivative works based on the work as long as all references to Project Gutenberg are removed. Of course, we hope that you will support the Project Gutenberg-tm mission of promoting free access to electronic works by freely sharing Project Gutenberg-tm works in compliance with the terms of this agreement for keeping the Project Gutenberg-tm name associated with the work. You can easily comply with the terms of this agreement by keeping this work in the same format with its attached full Project Gutenberg-tm License when you share it without charge with others.

¹This particular work is one of the few individual works restricted by copyright law in the United States and most of the remainder of the world, included in the Project Gutenberg collection with the permission of the copyright holder. Information on the copyright owner for this particular work and the terms of use imposed by the copyright holder on this work are set forth at the beginning of this work.

¹This paragraph, after 1.C., is included only for copyrighted works. For those, you must contact the copyright holder before any non-free use or removal of the Project Gutenberg header.

22.1. General Terms of Use and Redistributing Project Gutenberg-tm electronic works

1.D. The copyright laws of the place where you are located also govern what you can do with this work. Copyright laws in most countries are in a constant state of change. If you are outside the United States, check the laws of your country in addition to the terms of this agreement before downloading, copying, displaying, performing, distributing or creating derivative works based on this work or any other Project Gutenberg-tm work. The Foundation makes no representations concerning the copyright status of any work in any country outside the United States.

1.E. Unless you have removed all references to Project Gutenberg:

1.E.1. The following sentence, with active links to, or other immediate access to, the full Project Gutenberg-tm License must appear prominently whenever any copy of a Project Gutenberg-tm work (any work on which the phrase “Project Gutenberg” appears, or with which the phrase “Project Gutenberg” is associated) is accessed, displayed, performed, viewed, copied or distributed:

This eBook is for the use of anyone anywhere at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg License included with this eBook or online at www.gutenberg.org. If you are not located in the United States, you'll have to check the laws of the country where you are located before using this ebook.

1.E.2. If an individual Project Gutenberg-tm electronic work is derived from the public domain (does not contain a notice indicating that it is posted with permission of the copyright holder), the work can be copied and distributed to anyone in the United States without paying any fees or charges. If you are redistributing or providing access to a work with the phrase “Project Gutenberg” associated with or appearing on the work, you must comply either with the requirements of paragraphs 1.E.1 through 1.E.7 or obtain permission for the use of the work and the Project Gutenberg-tm trademark as set forth in paragraphs 1.E.8 or 1.E.9.

22. License

1.E.3. If an individual Project Gutenberg-tm electronic work is posted with the permission of the copyright holder, your use and distribution must comply with both paragraphs 1.E.1 through 1.E.7 and any additional terms imposed by the copyright holder. Additional terms will be linked to the Project Gutenberg-tm License for all works posted with the permission of the copyright holder found at the beginning of this work.

1.E.4. Do not unlink or detach or remove the full Project Gutenberg-tm License terms from this work, or any files containing a part of this work or any other work associated with Project Gutenberg-tm.

1.E.5. Do not copy, display, perform, distribute or redistribute this electronic work, or any part of this electronic work, without prominently displaying the sentence set forth in paragraph 1.E.1 with active links or immediate access to the full terms of the Project Gutenberg-tm License.

1.E.6. You may convert to and distribute this work in any binary, compressed, marked up, nonproprietary or proprietary form, including any word processing or hypertext form. However, if you provide access to or distribute copies of a Project Gutenberg-tm work in a format other than “Plain Vanilla ASCII” or other format used in the official version posted on the official Project Gutenberg-tm web site (www.gutenberg.net), you must, at no additional cost, fee or expense to the user, provide a copy, a means of exporting a copy, or a means of obtaining a copy upon request, of the work in its original “Plain Vanilla ASCII” or other form. Any alternate format must include the full Project Gutenberg-tm License as specified in paragraph 1.E.1.

1.E.7. Do not charge a fee for access to, viewing, displaying, performing, copying or distributing any Project Gutenberg-tm works unless you comply with paragraph 1.E.8 or 1.E.9.

1.E.8. You may charge a reasonable fee for copies of or providing access to or distributing Project Gutenberg-tm electronic works provided that:

- You pay a royalty fee of 20% of the gross profits you derive from the use of Project Gutenberg-tm works calculated using the method you already use to calculate your applicable taxes. The fee is owed to the

22.1. *General Terms of Use and Redistributing Project Gutenberg-tm electronic works*

owner of the Project Gutenberg-tm trademark, but he has agreed to donate royalties under this paragraph to the Project Gutenberg Literary Archive Foundation. Royalty payments must be paid within 60 days following each date on which you prepare (or are legally required to prepare) your periodic tax returns. Royalty payments should be clearly marked as such and sent to the Project Gutenberg Literary Archive Foundation at the address specified in Section 22.4, “Information about donations to the Project Gutenberg Literary Archive Foundation.”

- You provide a full refund of any money paid by a user who notifies you in writing (or by e-mail) within 30 days of receipt that s/he does not agree to the terms of the full Project Gutenberg-tm License. You must require such a user to return or destroy all copies of the works possessed in a physical medium and discontinue all use of and all access to other copies of Project Gutenberg-tm works.
- You provide, in accordance with paragraph 1.F.3, a full refund of any money paid for a work or a replacement copy, if a defect in the electronic work is discovered and reported to you within 90 days of receipt of the work.
- You comply with all other terms of this agreement for free distribution of Project Gutenberg-tm works.

1.E.9. If you wish to charge a fee or distribute a Project Gutenberg-tm electronic work or group of works on different terms than are set forth in this agreement, you must obtain permission in writing from both the Project Gutenberg Literary Archive Foundation and Michael Hart, the owner of the Project Gutenberg-tm trademark. Contact the Foundation as set forth in Section 22.3 below.

1.F.

1.F.1. Project Gutenberg volunteers and employees expend considerable effort to identify, do copyright research on, transcribe and proofread public domain works in creating the Project Gutenberg-tm collection. Despite

22. *License*

these efforts, Project Gutenberg-tm electronic works, and the medium on which they may be stored, may contain “Defects,” such as, but not limited to, incomplete, inaccurate or corrupt data, transcription errors, a copyright or other intellectual property infringement, a defective or damaged disk or other medium, a computer virus, or computer codes that damage or cannot be read by your equipment.

1.F.2. LIMITED WARRANTY, DISCLAIMER OF DAMAGES

- Except for the “Right of Replacement or Refund” described in paragraph 1.F.3, the Project Gutenberg Literary Archive Foundation, the owner of the Project Gutenberg-tm trademark, and any other party distributing a Project Gutenberg-tm electronic work under this agreement, disclaim all liability to you for damages, costs and expenses, including legal fees. YOU AGREE THAT YOU HAVE NO REMEDIES FOR NEGLIGENCE, STRICT LIABILITY, BREACH OF WARRANTY OR BREACH OF CONTRACT EXCEPT THOSE PROVIDED IN PARAGRAPH F3. YOU AGREE THAT THE FOUNDATION, THE TRADEMARK OWNER, AND ANY DISTRIBUTOR UNDER THIS AGREEMENT WILL NOT BE LIABLE TO YOU FOR ACTUAL, DIRECT, INDIRECT, CONSEQUENTIAL, PUNITIVE OR INCIDENTAL DAMAGES EVEN IF YOU GIVE NOTICE OF THE POSSIBILITY OF SUCH DAMAGE.

1.F.3. LIMITED RIGHT OF REPLACEMENT OR REFUND -

If you discover a defect in this electronic work within 90 days of receiving it, you can receive a refund of the money (if any) you paid for it by sending a written explanation to the person you received the work from. If you received the work on a physical medium, you must return the medium with your written explanation. The person or entity that provided you with the defective work may elect to provide a replacement copy in lieu of a refund. If you received the work electronically, the person or entity providing it to you may choose to give you a second opportunity to receive the work electronically in lieu of a refund. If the second copy is also defective, you may demand a refund in writing without further opportunities to fix the problem.

1.F.4. Except for the limited right of replacement or refund set forth in

22.2. Information about the Mission of Project Gutenberg-tm

paragraph 1.F.3, this work is provided to you 'AS-IS', WITH NO OTHER WARRANTIES OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR ANY PURPOSE.

1.F.5. Some states do not allow disclaimers of certain implied warranties or the exclusion or limitation of certain types of damages. If any disclaimer or limitation set forth in this agreement violates the law of the state applicable to this agreement, the agreement shall be interpreted to make the maximum disclaimer or limitation permitted by the applicable state law. The invalidity or unenforceability of any provision of this agreement shall not void the remaining provisions.

1.F.6. INDEMNITY - You agree to indemnify and hold the Foundation, the trademark owner, any agent or employee of the Foundation, anyone providing copies of Project Gutenberg-tm electronic works in accordance with this agreement, and any volunteers associated with the production, promotion and distribution of Project Gutenberg-tm electronic works, harmless from all liability, costs and expenses, including legal fees, that arise directly or indirectly from any of the following which you do or cause to occur: (a) distribution of this or any Project Gutenberg-tm work, (b) alteration, modification, or additions or deletions to any Project Gutenberg-tm work, and (c) any Defect you cause.

22.2. Information about the Mission of Project Gutenberg-tm

Project Gutenberg-tm is synonymous with the free distribution of electronic works in formats readable by the widest variety of computers including obsolete, old, middle-aged and new computers. It exists because of the efforts of hundreds of volunteers and donations from people in all walks of life.

22. License

Volunteers and financial support to provide volunteers with the assistance they need, is critical to reaching Project Gutenberg-tm's goals and ensuring that the Project Gutenberg-tm collection will remain freely available for generations to come. In 2001, the Project Gutenberg Literary Archive Foundation was created to provide a secure and permanent future for Project Gutenberg-tm and future generations. To learn more about the Project Gutenberg Literary Archive Foundation and how your efforts and donations can help, see Section 22.3 and Section 22.4 and the Foundation information page at www.gutenberg.org.

22.3. Information about the Project Gutenberg Literary Archive Foundation

The Project Gutenberg Literary Archive Foundation is a non profit 501(c)(3) educational corporation organized under the laws of the state of Mississippi and granted tax exempt status by the Internal Revenue Service. The Foundation's EIN or federal tax identification number is 64-6221541. Contributions to the Project Gutenberg Literary Archive Foundation are tax deductible to the full extent permitted by U.S. federal laws and your state's laws.

The Foundation's principal office is located in Montclair, New Jersey, but its volunteers and employees are scattered throughout numerous locations. Email contact links and up to date contact information can be found at the Foundation's web site and official page at www.gutenberg.org/contact.

For additional contact information:

Eric Hellman
Executive Director
[eric \(at\) pglaf.org](mailto:eric(at)pglaf.org)

22.4. Information about Donations to the Project Gutenberg Literary Archive Foundation

Project Gutenberg-tm depends upon and cannot survive without wide spread public support and donations to carry out its mission of increasing the number of public domain and licensed works that can be freely distributed in machine readable form accessible by the widest array of equipment including outdated equipment. Many small donations (\$5 to \$5,000) are particularly important to maintaining tax exempt status with the IRS.

The Foundation is committed to complying with the laws regulating charities and charitable donations in all 50 states of the United States. Compliance requirements are not uniform and it takes a considerable effort, much paperwork and many fees to meet and keep up with these requirements. We do not solicit donations in locations where we have not received written confirmation of compliance. To SEND DONATIONS or determine the status of compliance for any particular state visit www.gutenberg.org/donate.

While we cannot and do not solicit contributions from states where we have not met the solicitation requirements, we know of no prohibition against accepting unsolicited donations from donors in such states who approach us with offers to donate.

International donations are gratefully accepted, but we cannot make any statements concerning tax treatment of donations received from outside the United States. U.S. laws alone swamp our small staff.

Please check the Project Gutenberg Web pages for current donation methods and addresses. Donations are accepted in a number of other ways including including checks, online payments and credit card donations. To donate, please visit: www.gutenberg.org/donate.

22.5. General Information About Project Gutenberg-tm electronic works

Professor Michael S. Hart is the originator of the Project Gutenberg-tm concept of a library of electronic works that could be freely shared with anyone. For thirty years, he produced and distributed Project Gutenberg-tm eBooks with only a loose network of volunteer support.

Project Gutenberg-tm eBooks are often created from several printed editions, all of which are confirmed as Public Domain in the U.S. unless a copyright notice is included. Thus, we do not necessarily keep eBooks in compliance with any particular paper edition.

Most people start at our Web site which has the main PG search facility:
www.gutenberg.org

This Web site includes information about Project Gutenberg-tm, including how to make donations to the Project Gutenberg Literary Archive Foundation, how to help produce our new eBooks, and how to subscribe to our email newsletter to hear about new eBooks.

Bibliography

- [1] E. Heine, “Die Elemente der Functionenlehre,” *Journal für die reine und angewandte Mathematik (Crelles Journal)*, vol. 1872, no. 74, pp. 172–188, 1872, doi: [10.1515/crll.1872.74.172](https://doi.org/10.1515/crll.1872.74.172).
- [2] G. Cantor, “Ueber die Ausdehnung eines Satzes aus der Theorie der trigonometrischen Reihen,” *Mathematische Annalen*, vol. 5, no. 1, pp. 123–132, 1872, doi: [10.1007/BF01446327](https://doi.org/10.1007/BF01446327).
- [3] P. Lejeune Dirichlet, *Vorlesungen über Zahlentheorie / von P. G. Lejeune Dirichlet. Hrsg. Und mit Zusätzen versehen von R. Dedekind*. Vieweg, 1879. doi: [10.24355/DBBS.084-200909111325-0](https://doi.org/10.24355/DBBS.084-200909111325-0).
- [4] L. Kronecker, “Ueber einige Anwendungen der Modulsysteme auf elementare algebraische Fragen,” *Journal für die reine und angewandte Mathematik (Crelles Journal)*, vol. 1886, no. 99, pp. 329–371, Jan. 1886, doi: [10.1515/crll.1886.99.329](https://doi.org/10.1515/crll.1886.99.329).
- [5] B. Bolzano and P. Frantisek, *Paradoxien des Unendlichen*, 1st ed. Leipzig: C.H. Reclam sen., 1851. Available: <https://archive.org/details/details/paradoxiendesune00bolz/>

