Analysis report

Labelled mediation · ML · listwise · N = 400 · lavaan 0.7.3

Table 1: Parameter estimates

Group	Left	Relation	Right	Label / value	Estimate	SE	z	p	95% CI lower	95% CI upper	std.all
1	M	~	X	a	0.596	0.038	15.683	< .001	0.518	0.685	0.620
1	Y	~	M	b	0.473	0.038	12.342	< .001	0.379	0.541	0.494
1	Y	~	X	cp	0.183	0.037	4.929	< .001	0.098	0.243	0.199
1	M	~~	M		0.584	0.027	21.976	< .001	0.543	0.647	0.615
1	Y	~~	Y		0.517	0.021	24.765	< .001	0.473	0.558	0.594
1	X	~~	X		1.027	0.056	18.439	< .001	0.917	1.137	1.000
1	ind	:=	a*b	ind	0.282	0.029	9.585	< .001	0.229	0.350	0.307
1	total	:=	cp+ind	total	0.465	0.039	11.861	< .001	0.383	0.550	0.505

Note. Estimates and intervals are unstandardised; Std. all is fully standardised. Values are rounded only for display. A dash indicates unavailable output. 95% CI. Percentile confidence intervals are unstandardised. SEs are bootstrap SEs; z and p remain normal approximations. lavaan excludes failed draws from the intervals and reports inadmissible draws separately.

Table 2: Fit measures

Measure	Value
χ^2	0.000
df	0.000
p (χ^2)	—
CFI	1.000
TLI	1.000
RMSEA	0.000
RMSEA 90% CI lower	0.000
RMSEA 90% CI upper	0.000
SRMR	0.000
Scaled χ^2	—
Scaled df	—
p (scaled χ^2)	—
Robust CFI	—
Robust TLI	—
Robust RMSEA	—
Robust RMSEA 90% CI lower	—
Robust RMSEA 90% CI upper	—
Scaled CFI	—
Scaled TLI	—
Scaled RMSEA	—
Scaled RMSEA 90% CI lower	—
Scaled RMSEA 90% CI upper	—

Note. Scaled and robust statistics are labelled explicitly. Compare the appropriate version for your estimator.

Table 3: Explained variance (R^2)

Group	Left	R^2
1	M	0.385
1	Y	0.406

Table 4: Bootstrap summary

Requested	Returned draws	Failed draws	Inadmissible draws	Random seed	Confidence level
100	100	0	0	12345	95%

Note. Percentile confidence intervals are unstandardised. SEs are bootstrap SEs; z and p remain normal approximations. lavaan excludes failed draws from the intervals and reports inadmissible draws separately.