# Analysis report

Labelled mediation · ML · listwise · N = 400 · lavaan 0.7.3

Table 1: Parameter estimates

| Group | Left | Relation | Right | Label / value | Estimate | SE | z | p | 95% CI lower | 95% CI upper | std.all |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | M | ~ | X | a | 0.596 | 0.038 | 15.683 | < .001 | 0.518 | 0.685 | 0.620 |
| 1 | Y | ~ | M | b | 0.473 | 0.038 | 12.342 | < .001 | 0.379 | 0.541 | 0.494 |
| 1 | Y | ~ | X | cp | 0.183 | 0.037 | 4.929 | < .001 | 0.098 | 0.243 | 0.199 |
| 1 | M | ~~ | M | | 0.584 | 0.027 | 21.976 | < .001 | 0.543 | 0.647 | 0.615 |
| 1 | Y | ~~ | Y | | 0.517 | 0.021 | 24.765 | < .001 | 0.473 | 0.558 | 0.594 |
| 1 | X | ~~ | X | | 1.027 | 0.056 | 18.439 | < .001 | 0.917 | 1.137 | 1.000 |
| 1 | ind | := | a*b | ind | 0.282 | 0.029 | 9.585 | < .001 | 0.229 | 0.350 | 0.307 |
| 1 | total | := | cp+ind | total | 0.465 | 0.039 | 11.861 | < .001 | 0.383 | 0.550 | 0.505 |

Note. Estimates and intervals are unstandardised; Std. all is fully standardised. Values are rounded only for display. A dash indicates unavailable output. 95% CI. Percentile confidence intervals are unstandardised. SEs are bootstrap SEs; z and p remain normal approximations. lavaan excludes failed draws from the intervals and reports inadmissible draws separately.

Table 2: Fit measures

| Measure | Value |
|---|---|
| $\chi^2$ | 0.000 |
| df | 0.000 |
| p ($\chi^2$) | — |
| CFI | 1.000 |
| TLI | 1.000 |
| RMSEA | 0.000 |
| RMSEA 90% CI lower | 0.000 |
| RMSEA 90% CI upper | 0.000 |
| SRMR | 0.000 |
| Scaled $\chi^2$ | — |
| Scaled df | — |
| p (scaled $\chi^2$) | — |
| Robust CFI | — |
| Robust TLI | — |
| Robust RMSEA | — |
| Robust RMSEA 90% CI lower | — |
| Robust RMSEA 90% CI upper | — |
| Scaled CFI | — |
| Scaled TLI | — |
| Scaled RMSEA | — |
| Scaled RMSEA 90% CI lower | — |
| Scaled RMSEA 90% CI upper | — |

Note. Scaled and robust statistics are labelled explicitly. Compare the appropriate version for your estimator.

Table 3: Explained variance ($R^2$)

| Group | Left | $R^2$ |
|---|---|---|
| 1 | M | 0.385 |
| 1 | Y | 0.406 |

Table 4: Bootstrap summary

| Requested | Returned draws | Failed draws | Inadmissible draws | Random seed | Confidence level |
|---|---|---|---|---|---|
| 100 | 100 | 0 | 0 | 12345 | 95% |

Note. Percentile confidence intervals are unstandardised. SEs are bootstrap SEs; z and p remain normal approximations. lavaan excludes failed draws from the intervals and reports inadmissible draws separately.