



Original article

Pristine and UV-aged polyethylene microplastics' impact on gut microbiome and reproduction of earthworm *Eisenia andrei*Elise Quigley^{a,*}, Ana L. Patrício Silva^b, Sónia Chelinho^c, Luís Cunha^d, Maria JI. Briones^a, José P. Sousa^d^a Departamento de Ecología y Biología Animal, Universidad de Vigo, 36310, Vigo, Spain^b Centre for Environmental and Marine Studies (CESAM) & Department of Biology, University of Aveiro, Portugal^c Polytechnic Institute of Coimbra, Coimbra, Portugal^d Centre for Functional Ecology, Associate Laboratory TERRA, Department of Life Sciences, University of Coimbra, Coimbra, Portugal

ARTICLE INFO

Handling editor: Guénola Pérès

Keywords:

Photodegradation

UV irradiation

LDPE

Dysbiosis

Microbiota

ABSTRACT

The impact of microplastics (MPs) on soil organisms is still a growing field, yet very little is known about the exposure of ultraviolet (UV) aging of MPs to soil organisms. In this study, we explored the response of the epigeic earthworm *Eisenia andrei* to pristine and UV-aged polyethylene (PE-MP) exposure at a wide range of environmentally relevant concentrations (0, 0.2, 2, 20, 200 and 2000 mg kg⁻¹) in an organic farm soil for 56d and assessed changes in reproduction, ingestion, egestion and gut microbiome. Results showed that exposure to 20 mg kg⁻¹ pristine PE-MP significantly increased earthworm reproduction by 39 % but the same concentration decreased reproduction by 29 % when they were exposed to UV-aged PE-MP. Ingestion of PE was verified by staining the whole worm body and their casts after 48h of starvation. The amounts of PE-MP found in the body and the casts were positively correlated with PE-MP concentrations in the soil, however only significantly so with pristine PE-MP. A decline in *E. andrei* gut microbiome alpha diversity and a significantly different community composition were observed in UV-aged PE-MP exposures compared to pristine PE-MP. Relative to the control treatments, Proteobacteria increased up to 135 %, Actinobacteria increased up to 35 %, and Firmicutes decreased up to 38 % under UV-aged PE-MP exposure, whereas Cyanobacteria increased up to 19 times in the pristine PE-MP treatments. These results confirm the negative effect of UV-aged PE-MP on earthworms even at low concentrations and could have important implications in the well-functioning of agricultural soils.

1. Introduction

In the current plastic age, the production and management of plastic and its residues have become unsustainable [1]. In 2021, plastic production reached a global amount of 390.7 Mt [2] and is still rising. The OECD estimates, as of 2019, that globally 22 % of plastics are mismanaged and end up as litter [2]. These staggering numbers result in plastic debris in every environmental compartment on earth [3], characterized by its own biogeochemical cycle [4]. Terrestrial environments are a prominent and growing sink for plastic debris, with increasing concern towards smaller size fractions [5]. Concentrations of microplastics (plastic particles between 1 µm and 5 mm in size [6]) in the soil have been reported globally ranging from $4 \times 10^{-6}\%$ –6.7 % (0.004–67000 mg kg⁻¹) of soil weight [7]. The accumulation and persistence of microplastics in soils pose a potential threat to the

pedosphere, the organisms that inhabit it, and the functions they play a role in.

A considerable part of plastic debris present in soils is represented by secondary microplastics, which are formed from macroplastics subjected to weathering and degradation in the environment, predominantly by ultraviolet (UV) irradiation [8–10]. The importance of studying the effects of aged plastic debris/microplastics has been urged [11], because the process of aging is inevitable in the environment and can affect the particle's physicochemistry [12–14]. In addition, weathering processes alter the polymer's interactions with its surroundings biologically, geologically, and molecularly [15–18], altering the severity and mechanisms of toxicity posed to organisms. For example, UV-aged microplastics have been shown to exacerbate toxicity in a wide range of mostly aquatic organisms, from intestinal injury to cellular dysfunction, among other effects [19–22]. In soil organisms, however,

* Corresponding author.

E-mail address: elisemarie.quigley@uvigo.gal (E. Quigley).<https://doi.org/10.1016/j.ejsobi.2024.103640>

Received 31 July 2023; Received in revised form 31 March 2024; Accepted 9 June 2024

1164-5563/© 2024 Published by Elsevier Masson SAS.

knowledge of aged microplastic toxicity is more limited. Li et al. [23] reported that the aging process (H_2O_2 or UV) did not change the toxicity of polyethylene microplastics (PE-MPs) in the earthworm *Eisenia fetida* when exposed to 0.1 %, 1 %, and 10 % dw soil, indicated by negligible differences in oxidative stress levels. Whereas oxidative stress was exacerbated in *E. fetida* exposed to field aged LDPE film (0.25 %) [24], and in *E. andrei* exposed to mixtures of plastics (predominantly PE) at 0.1 mg kg⁻¹ [25], both in agriculturally sourced soils. Although the effects of MPs at environmentally relevant concentrations on apical endpoints are still variable among test organisms, especially regarding earthworms [26], there is a great consensus when evaluating physiological and biochemical parameters (e.g., oxidative stress biomarkers and gut microbiome composition [15,26]) as they could be considered early indications of microplastic's toxicity.

Comprising the most abundant animal biomass in most terrestrial systems, the ecosystem services earthworms provide, such as litter decomposition, nutrient cycling, and pollutant transformation [27], are mediated by their gut microbiome [28]. Plus, being closely linked to the host's health, the gut microbiota is essential to vital processes in the earthworm body [29,30] and thus can serve as a powerful biological indicator of health [31]. Previous studies have shown that *E. fetida* exposed to pristine polystyrene (PS) nanoplastics at 200 mg kg⁻¹ experienced a reduction in diversity and species richness of gut microbiome along with community composition shifts [32]. Similar changes were found in the gut microbiota of *E. fetida* exposed to polystyrene (PS) MPs (0.1–10 µm) at 10 and 100 mg kg⁻¹ [33]. These authors also found more negative interactions among gut communities when exposed to larger-sized particles [33]. With regards to aged MPs, H_2O_2 degraded PE-MPs have been shown to intensify gut microbial disruptions where 500 mg kg⁻¹ increased pathogenic bacteria and inhibited probiotic bacteria that promote gut homeostasis [34]. In addition, aged farmland residual PE mulch film MPs (0.25 %) showed a notable effect on *Metaphire guillelmi* gut community structure compared to pristine PE-MP exposure resulting in more simplified microbial networks with greater antagonistic interactions [35]. This disruption in mutualistic processes could ripple throughout the soil ecosystem where they reside [15], making the potential risks of altered gut microbial function a pressing matter to investigate.

Polyethylene (PE) is one of the most common and prevalent polymers in agricultural soils, due to its intensive application in mulching and greenhouse films, and irrigation systems, among others [36]. These uses increase exposure to UV photodegradation, where further fragmentation and accumulation in agricultural soils is inevitable [13,37]. This study assessed *E. andrei* survival, growth, reproduction, and gut microbiome composition after being exposed to pristine and UV-aged polyethylene microplastics (PE-MP) at a wide range of environmentally relevant concentrations (0, 0.2, 2, 20, 200 and 2000 mg kg⁻¹) for 56 days. *E. andrei* is a model organism in soil ecotoxicology [38], due to its sensitivity to a wide range of toxic substances, including microplastics (e.g., Ref. [33]), high reproduction rate, short life cycle, and easy maintenance in the laboratory environment. It was hypothesized that increasing concentrations of PE-MPs added to the soil would negatively impact earthworms, with UV-aged PE-MP inducing more pronounced effects than pristine PE-MP exposures. To the best of our knowledge, this is the first study addressing the ecotoxicity of pristine vs. UV-aged PE-MPs on earthworm reproduction and gut microbiome and serves as a steppingstone for further research in this area, considering the prevalence of photodegraded plastics in the terrestrial environment.

2. Materials and methods

2.1. Microplastics and contamination contingency measures

Polyethylene (PE) particles with an average size of 125 µm (ultra-high molecular weight powder, CAS No. 9002-88-4, density 0.94 g mL⁻¹

(at 25 °C)), were obtained from Sigma-Aldrich UK without any pre-treatment, and hence considered as “pristine” in the experimental design. UV-aged PE-MPs were prepared by subjecting the pristine particles to an ultraviolet light schedule of 24 h per day for 16 days in a dark box. Briefly, particles were placed in aluminum trays roughly 15 cm below the UV light and gently stirred twice a day for more consistent exposure. UV-C light (VL-6.LC Vilber, Germany), with a wavelength of 254 nm, was used for irradiation and was monitored twice daily, each with 3 readings for intensity consistency using a lux meter wand (Delta Ohm HD 9221).

The particle size was chosen based on average microplastics size ranges in previous studies that used *E. andrei* as the model organism [39–41]. Particle size distribution of the PE-MP particles was determined by Silva et al. [11]. Briefly, 100 g of PE-MP particles were sieved in a vibratory sieve shaker (mesh pore-sizes: 500, 250, 125, 63 and 32 µm); the majority of the particles fell into the size ranges ³125 - < 250 µm (50.15 ± 2.35 %) and ³63 - <125 (39.75 ± 1.75).

All equipment was either acid washed or thoroughly washed with soap and water then dried and additionally wiped down with ethanol (96 %) before and after each use. All solutions used were previously filtrated, and fiberglass filters were previously burnt for 3h at 500 °C. Procedural blanks (n = 4) were considered for contamination assessment. The filtering cup from the glassware filtration apparatus was rinsed with 10 % nitric acid, followed by ultra-pure water to avoid cross-contamination.

2.2. Test soil

Agricultural soil was collected from an organic farm (Herdade do Freixo do Meio, 440ha, 38°42'13.2" N, 8°19'31.4" W) in southern Portugal (Montemor-o-Novo, in the region of Alentejo). The soil was sieved with a 5 mm mesh in the field and defaunated by freezing, followed by thawing (3 cycles). Detailed soil physicochemical information (texture, phosphorus, effective cation exchange capacity, pH, humic acids, nitrate, ammonium, C and N) are described by Reis et al. [42]. In brief, the soil presented a sandy loam texture, low pH (average pH of 5.4 ± 0.04 (mean ± SE)), and high organic matter content (4.0 ± 0.7 %). Water holding capacity (WHC; measured according to ISO 11267 [43]) was 35.56 ± 0.55 % (mean ± SE).

2.3. Test organisms

The earthworms *Eisenia andrei* (Oligochaeta: Lumbricidae) Bouché, 1972 were used as model organisms and were obtained from cultures maintained in the laboratory at a temperature of 20 ± 2 °C, 40–50 % humidity and 16h:8h light:dark cycle. The earthworms were acclimated in the experimental soil two weeks prior to the beginning of the incubation period and fed weekly with defaunated cow manure. The individuals selected for the tests were adults (clitellated) with an average fresh weight of 308.2 ± 2.3 mg and had an individual fresh weight between 187 and 507 mg.

2.4. MP exposure

Experimental units consisted of mason glass test jars containing 500 g (dw) of soil and PE-MP particles added with distilled water at a volume corresponding to 40 % of soil water holding capacity (WHC), and thoroughly mixed for approximately 5 min for an even dispersion. Nominal concentrations of 0, 0.2, 2, 20, 200 and 2000 mg PE kg⁻¹ of soil dw were prepared for both types of PE-MP treatments (pristine and UV-aged). These concentrations corresponded to 0.00002, 0.0002, 0.002, 0.02 and 0.2 % of dw soil and fall within realistic concentrations found in soil samples [44,45,].

Four replicates were prepared for each PE-MP concentration and control treatments (48 jars in total). Water content and pH were measured at the beginning and end of the incubation period. Controls

were prepared in the same manner with no PE-MP (pristine nor UV-aged) addition; pristine and UV-aged exposures began on different dates resulting in two sets of controls. Exposure followed ISO 11268–2 [38] guidelines. The earthworms (ten clitellate adults per treatment) were washed, dried with paper towels, and weighed before being randomly added to the experimental jars. Each week, water was added by mass loss to every replicate to maintain a constant moisture level; additionally, earthworms were fed weekly with initially 10g fresh weight of cow manure (moistened and homogenized), then increased to 20 g in later weeks by demand.

After 28 days of exposure, surviving adults were removed, scored, washed, dried, and weighed. Three adults from each replicate were placed in separate Petri dishes lined with moist filter paper for gut depuration. After 48 h in the dark, adult worms were collected then frozen (-4°C), while the filter paper with casts were dried in an incubator for further MP quantification described below. For further gut microbiome analysis, three additional adults from each replicate were fixated in 96 % ethanol. Adults were dissected under sterile conditions from the clitellum to the tail and intact gastrointestinal tracts were removed, pooled and separated into 4 eppendorfs (1.5 mL) and frozen (-4°C).

The experimental soil was incubated once more, without adult earthworms, for 28 additional days to allow the cocoons to hatch. On day 56, the experimental jars were placed in a water bath at 50°C to coax the juveniles to the soil surface for collection.

2.5. MP quantification and chemical changes during UV-aging

PE-MP extraction and quantification were performed on both the casts and the earthworm bodies adapting the methodology from Silva et al. [11] for the digestion procedure and Prata et al. [46] for PE-MP's quantification. Briefly, dried casts were scraped from the filter paper, where earthworms depurated their guts, into glass beakers where earthworm bodies and casts were digested separately in covered beakers in the oven at 50°C in a 5 mL solution of nitric acid (63 %) for 3 h, followed by digestion in 5 mL H_2O_2 (30 %) for another 24 h at room temperature under a fume hood to eliminate any biological matter from MP surface. The reaction was terminated with deionized water, and the solution was filtered on 1.2 μm glass fiber filters, followed by incubation for 10 min in 10 mL of acetone, stained for 5 min with 3 drops of Nile Red (0.01 mg mL^{-1} ethanol), then washed again with deionized water. The filters containing suspected microplastics were allowed to dry completely before fluorescent observation using the 470 nm FOCUS LED (SPEX Forensic, USA) with an orange camera lens filter (Standard Pro-Master® Orange Filter) attached to the microscope. Imaging was performed using the Nikon SMZ-U Zoom 1:10 microscope paired with the Nikon Digital Sight DS-L1 camera attachment.

Fourier transform infrared spectroscopy (FTIR Spectrometer Nicolet 6700 from Thermo Nicolet Corp., Madison, WI, USA) was used to analyze the change in the surface functional groups of the PE-MPs ($n = 3$). FTIR spectra analysis is a valuable tool, and commonly used in microplastic's ecotoxicology, for MP aging assessment [47]. During UV irradiation/aging, FTIR can detect the formation of new functional groups or changes in the intensity of existing ones, indicating i) oxidation with the formation of carbonyl groups ($\text{C}=\text{O}$), which is a classic sign of oxidation due to UV exposure; and ii) chain scission, as breaking of polymer chains can change the relative intensities of functional group peaks [48]. The carbonyl index is used to monitor the absorption band of the carbonyl species formed on the surface of a material. In the context of microplastics, it indicates the presence of oxygen-containing functional groups, such as carbonyl ($\text{C}=\text{O}$) groups. Carbonyl index was calculated by the area of $1650\text{--}1800\text{ cm}^{-1}$ over the area of $1420\text{--}1500\text{ cm}^{-1}$ (modified calculation from Almond et al. [49]).

2.6. 16S rRNA gene amplification, sequencing, and bioinformatics analysis

To compare the effects of pristine and UV-aged PE-MP on the gut microbiome of *E. andrei*, 16S rRNA V4–V5 variable regions of the gut bacteria were sequenced by high-throughput sequencing performed on an Illumina Novaseq platform (Novogene, Cambridge, UK). DNA of *E. andrei* gut content was extracted using DNeasy Blood and Tissue Kit (Qiagen, Valencia, CA) according to the manufacturer's instructions with an added initial tissue homogenization step using metal beads in a TissueLyser LT (Qiagen, Valencia, CA). From the 3 adult worms per treatment, a total of 48 composite samples were obtained from combined gut contents (4 per treatment). The quality and concentration of DNA were determined using a Nanodrop 2000 spectrophotometer (Thermo Fisher, USA). Extracted DNA was stored at -20°C for downstream analysis. Further quality control, PCR, purification, library preparation, and sequencing were performed at Novogene (Cambridge, UK). Forward primer 515F (GTGCCAGCMGCCGCGG) and reverse primer 907R (CCGTCAATCMTTTRAGTTT), labeled with unique barcodes that target the hypervariable V4–V5 region of the bacterial 16S rRNA gene, were selected. PCR amplification was done with Phusion® High-Fidelity PCR Master Mix (New England Biolabs). Successful amplification was detected with electrophoresis on 2 % agarose gel; samples with bright bands between 400 and 450bp chosen for further analysis. Then, PCR products were purified with a Qiagen Gel Extraction Kit (Qiagen, Germany).

The same amount of PCR products from each experimental treatment sample were pooled, end-repaired, A-tailed and further ligated with Illumina adapters, then sequenced on an Illumina Novaseq platform (Novogene, Cambridge, UK). DADA2 pipeline [50] was used to infer the amplicon sequence variants (ASVs) present in each sample; a higher accuracy feature sequence analogous to operational taxonomic units (OTUs) [51]. Briefly, paired-end reads were matched to samples and truncated, then merged using FLASH (v1.2.11, <http://ccb.jhu.edu/software/FLASH/>) [52]. High-quality clean tags were obtained using fastp software [53]. Then, chimeras were identified and removed using Vsearch software. The resulting tags were denoised using DADA2 software with less than 5 abundance sequences filtered out to obtain the final ASVs. Finally, species annotation of each ASV was carried out by the QIIME2's classify-sklearn module [54,55].

2.7. Statistical analyses

Effects on survival, growth, reproduction, gut microbiota phyla and family abundances, and PE-MP particles found in worm body and casts were analyzed using a two-way ANOVA with PE-MP type (pristine and UV-aged) and concentration level as fixed factors, followed by a post-hoc LSD test. Change in carbonyl index between pristine and UV-aged PE-MP was analyzed using an unpaired *t*-test. Normality was tested using the Shapiro-Wilk test, and homogeneity of variances was tested using Levene's test. Tests were performed using GraphPad Prism 9. Correlation analyses were performed on the PE-MP count found in the body and casts of the earthworms using the Pearson correlation coefficient or Spearman rank correlation coefficient, depending on the normality of the dataset. For microbiome data, QIIME2 software was used to calculate alpha diversity indices and UniFrac distance for beta diversity between treatments. Weighted UniFrac and Unweighted UniFrac distances were used to perform a Principal Coordinate Analysis (PCoA) to explore the differences between microbial communities (beta diversity) [56,57], and visualized using the R “ggplot2” package [58]. Differences in microbiome composition between pristine and UV-aged PE-MP and between exposure concentrations were analyzed using Bray-Curtis distance with a PERMANOVA implemented in the “adonis2” function from R “vegan” package. All values are expressed as mean \pm standard deviation (SD).

3. Results

3.1. Characterization of pristine and UV-aged PE

FTIR spectra from pristine and UV aged particles presented similar peak distribution, yet there were significantly different carbonyl indices (Fig. S1, Supplementary Material). Carbonyl index significantly increased from pristine to UV-aged PE-MP with values of 0.0078–0.038, respectively ($t = 8.693$; $df = 4$; $p < 0.001$). This formation of new carbonyl species, demonstrated by increasing CI, indicates oxidation and degradation. Distinction in FTIR spectra between pristine and UV-aged PE-MP can be appreciated in the magnified regions used in the calculation of CI (Fig. S2, Supplementary Material).

3.2. Effects of pristine and UV-aged polyethylene on the survival, growth and reproduction of *E. andrei*

After 28d, there was 100 % survival in all treatments. Likewise, neither pristine nor UV-aged PE-MP had any significant effect on the earthworm biomass compared to the control (Fig. S3).

After 56d of exposure the number of juveniles recorded was significantly different between pristine and UV-aged PE-MP (Fig. 1) ($F(1, 36) = 5.629$; $p < 0.05$), yet increasing concentrations did not impact reproduction. Earthworm reproduction increased under pristine PE-MP exposures of 2 mg kg^{-1} and above, but it decreased under UV-aged PE-MP exposures, apart from a slight increase of $0.77 \pm 13.4 \%$ in reproduction at 2 mg kg^{-1} UV-aged PE-MP observed when compared to the control. The difference in reproduction between pristine and UV-aged PE-MP was particularly evident at the concentration 20 mg kg^{-1} , where the pristine PE-MPs treatments had $39.3 \pm 7.3 \%$ more juveniles ($p < 0.05$) and those with UV-aged PE-MPs registered $21.9 \pm 3.8 \%$ less juveniles ($p < 0.05$) compared to the respective controls.

Validity of *E. andrei* reproduction tests based on ISO 11268-2 [38] standard was confirmed and further described in Supplementary Material.

3.3. PE-MP concentration in *E. andrei* body and casts

Both pristine and UV-aged PE-MP particles were found in earthworm bodies and casts at all exposure concentrations (Fig. 2), confirming both ingestion and egestion. However, the two PE-MP types did not differ in the number of particles recovered. Nevertheless, the number of particles in the bodies and casts increased with increasing PE-MP concentrations in the soil (Fig. 2), yet only pristine PE-MP particle concentrations




Fig. 2. Concentration of PE-MP particles present in the body and casts of *E. andrei* exposed to pristine or UV-aged polyethylene microplastics after 48h of depuration. Data are expressed as mean number of particles (PE-MP #) \pm standard deviation (SD) ($n = 4$).

showed a significant positive correlation with particles found in the bodies and the casts (Spearman $r = 1$, $p < 0.05$ in both cases). This positive trend was not significant in the case of UV-aged PE-MP (Pearson $r = 0.6954$, and Spearman $r = 0.8208$, respectively, $p > 0.05$).

3.4. Effect of PE-MP on *E. andrei* gut microbiota

After assembling and filtering, a total of 2,822,147 clean reads were acquired from 48 samples, ranging from 43,787 to 79,339 per sample and with an average of 58,795 reads. After denoising, 2,6034 ASVs were obtained in each sample. The total amount of ASVs detected across all samples was not significantly different between treatments. Firmicutes (42.09 %), Actinobacteriota (28.69 %), Proteobacteria (12.54 %), Bacteroidota (7.78 %) and Planctomycetota (4.72 %) were the predominant bacterial phyla, comprising 95.82 % of the total microbiome community abundance among the control treatments.

A significant decline in alpha diversity of the gut microbiome occurred in UV-aged PE-MP exposure compared to pristine treatments, according to Chao1 index ($F(1, 36) = 6.679$; $p < 0.05$), however there were no significant differences between the different exposure concentrations ($F(5, 36) = 1.672$; $p > 0.05$).

Fig. 3a shows the ten most abundant phyla and their trends in relative abundance when exposed to increasing concentrations of pristine or UV-aged PE-MP. It can be seen that exposure to UV-aged PE-MP resulted in a larger increase of Actinobacteria compared to the pristine exposed worms ($F(1, 36) = 19.58$; $p < 0.001$), despite not being correlated with the concentration of PE-MPs in the soil. Actinobacteria increased absolute abundance by 0.81–23.74 % under the majority of pristine PE-MP exposures compared to the control, except for slight decreases of $1.7 \pm 20.16 \%$ and $7.12 \pm 22.78 \%$ at 2 and 2000 mg kg^{-1} , respectively. In contrast, exposure to UV-aged PE-MP increased Actinobacteria absolute abundance by 5.17–35.28 % compared to the control.

Firmicute absolute abundance remained relatively unchanged in the pristine PE-MP exposures, but decreased by 16.76–37.89 % in response to UV-aged PE-MP by showing a relatively steady decline at higher concentrations. This led to Firmicute abundance being the bacteria phyla that showed the most drastic changes between pristine and UV-aged exposures compared to any other phylum ($F(1, 36) = 65.05$; $p < 0.001$).

Proteobacteria exhibited a modest increasing trend when exposed to pristine PE-MP, but a more marked increase in absolute abundance when exposed to UV-aged MPs, with the highest increase of $135.47 \pm 130.3 \%$ at 2000 mg kg^{-1} ($p < 0.001$), resulting in PE-MP concentration being a significant factor ($F(5, 36) = 3.984$; $p < 0.05$).

Cyanobacteria showed the largest increase of any phyla with 19




Fig. 1. *E. andrei* reproduction as indicated by the number of juveniles produced after 56 days of exposure to pristine or UV-aged polyethylene microplastics in natural soil. Data are expressed as mean (juvenile #) \pm standard deviation (SD) ($n = 4$). Significant differences to control (two-way ANOVA): * ($p < 0.05$); ** ($p < 0.01$).




Fig. 3. Relative abundance of *E. andrei* gut microbiota at phylum (A) and family (B) levels present in the earthworm gut among different concentration (mg kg^{-1}) and treatments (prefix “P” signifies pristine and “UV” signifies UV-aged PE-MPs). Stacked bar plot of the top 10 microbial phyla (A) and top 21 families (B) with the largest mean relative abundance (%) in earthworm gut; categories outside of this were grouped into “Other”.

times the absolute abundance compared to the control at 2000 mg kg^{-1} exposure to pristine PE-MP ($p < 0.05$).

At family and genus level, it was observed that absolute abundance of the family Bacillaceae decreased when exposed to UV-aged PE-MP, but increased in pristine PE-MP exposures ($F(1, 36) = 68.25$; $p < 0.001$). The absolute abundance of *Aeromonas* was significantly different between pristine and UV-aged PE-MPs ($F(1, 36) = 6.89$; $p < 0.05$) and among concentrations ($F(5, 36) = 8.5$; $p < 0.001$), as a result of an abrupt increase in numbers at the highest concentration of UV-aged PE-MP whereas abundance was virtually nonexistent in other concentrations and pristine exposure. More bacterial family results can be found in the Supplementary Material.

The results of the Principal Coordinate Analysis (Fig. 4) showed a clear distinction between samples exposed to UV-aged PE-MP and those exposed to pristine PE-MP. The first two principal components (Fig. 4a) explained 49.11 % and 14.87 % of the variation between the treatments (weighted UniFrac distance), respectively. The unweighted UniFrac distance calculated for the first two components of the PCoA explained 21.2 % and 5.6 % of the variation (Fig. 4b), and indicated the existence

of important differences in gut microbial communities when the worms were exposed to the highest concentration of UV-aged PE-MP (2000 mg kg^{-1}) compared to those in the pristine PE-MP and control treatments. The PERMANOVA, using Bray-Curtis distance, confirmed these differences in community structure of *E. andrei* gut microbiome between pristine and UV-aged PE-MP exposures ($df = 1$; $F = 5.6896$; $R^2 = 0.10260$; $p < 0.05$).

4. Discussion

4.1. Contrasting effects of pristine PE-MP and UV-aged PE-MP on *E. andrei* reproduction

Pristine PE-MP exposure at concentrations of 2 mg kg^{-1} and above stimulated reproduction in earthworms, whereas UV-aged PE-MP at 20 mg kg^{-1} and above inhibited it. This opposite response was most pronounced at environmentally relevant concentrations of 20 mg kg^{-1} for both PE-MP types. Various studies have found that MP exposures can hinder reproduction in earthworms [59–63], especially at unrealistically




Fig. 4. Principal Coordinate Analysis (PCoA) based on Weighted UniFrac distance (a) and Unweighted UniFrac distance (b) showing the distribution pattern of *E. andrei* gut microbiome among different polyethylene microplastic treatments (0 mg kg^{-1} (C0), 0.2 mg kg^{-1} (C1), 2 mg kg^{-1} (C2), 20 mg kg^{-1} (C3), 200 mg kg^{-1} (C4), 2000 mg kg^{-1} (C5)). Data are based on 16S rRNA amplification and high-throughput sequencing. The microbiome of each sample is shown as a point. The percentages of the axis represent the percentage of variation explained.

high concentrations. However, the majority of research indicates that earthworm reproduction is not significantly impacted [40,62,64–66]. Some of these studies even describe a slight increase in earthworm reproduction exposed to pristine MPs [40,62,64], as observed in this study. Interestingly, a study using agricultural soil reported very minor increases in *E. fetida* reproduction at low concentrations of PE-MP exposure ($0.05\text{--}2.5\text{ mg kg}^{-1}$) after 20d, and subsequent measurements showed reproduction to decrease over time and with increasing concentrations, especially at 20 mg kg^{-1} after 60d [67]. In the present study, this concentration showed the greatest increases in reproduction with pristine PE-MP after 56d. Stimulation and hormetic responses to low levels of contamination of MPs have been extensively reported in aquatic organisms [68,69,70] and from this, the reproduction results of *E. andrei* observed here can be interpreted as stimulatory responses to low-risk contamination (i.e., pristine PE-MP exposure) with inversely inhibitory responses when faced with high-risk contamination (i.e., UV-aged PE-MPs). Sun et al. [69] found an overall 31 % increase in reproduction in aquatic organisms exposed to environmentally relevant concentrations of MPs ($\leq 1\text{ mg L}^{-1}$) regardless of habitat, polymer type, shape, size, exposure period or concentration. They concluded that the stimulation effect acts as a conditioning agent to strengthen biological performance and resistance [71,70]. Furthermore, these MP exposures had inhibitory effects on oxidative stress biomarkers, therefore alleviating oxidative stress, reflecting an adaptive response to maintain redox homeostasis [72]. In agreement with this, a later review [73] of dose responses among a wider range of organisms (22 diverse taxa from aquatic and terrestrial environments) exposed to various types of nanoplastics/microplastics (with differing polymers, sizes, shapes, and concentrations) identified 65 studies that presented hormetic responses in a wide range of endpoints from molecular to individual level. Our results are also supported by previous studies showing an increase in earthworm reproduction (albeit non-significant) after exposure to MP concentrations of low to midrange [40,62,64], which is comparable with increased reproduction seen here in *E. andrei* that was most prominent when exposed to pristine PE-MP up to a mid-level concentration (20 mg kg^{-1}) and subsequently decreased, yet not below the rates observed in the control treatment.

The effects of UV-aged MPs on organisms' reproduction have only been investigated in aquatic media [74–76], and shown to deepen the severity of toxicity, ranging from molecular to individual-level responses [19–22]. For example, nematodes exhibited reproductive toxicity after exposure to pristine and UV-aged PS-MP in an aqueous medium at environmental concentrations ($0.1\text{--}100\text{ }\mu\text{g L}^{-1}$); however, significantly more so in UV-aged PS-MPs, which was attributed to DNA damage-induced cell apoptosis [75]. The aged particles used in that study had increased carbonyl indices indicating UV photodegradation causing surface oxidation, and they were substantially dimpled and wrinkled compared to the smooth surface of pristine PS-MPs, creating a larger surface area to potentially interact with cellular membranes [77, 78]. Similarly, *Daphnia magna*, when exposed to UV-aged PS-MP showed chronic toxicity in the form of reproductive inhibition, which was linked to smaller-sized particle generation of PS-MPs through the UV-aging process resulting in 3 times the initial pristine concentration of particles [76]. This would reinforce the idea that smaller-sized particles increase specific surface area and biological interaction. In the present study, the carbonyl index increased substantially in UV-aged particles, indicative of oxidation [49], and progressive deterioration [79]. Along with the changing chemical structure, the process of oxidation causes significant changes in the physical properties of the material as well [49].

4.2. UV-aged PE-MP altered *E. andrei* gut bacterial communities

The UV-aged PE-MP exposure led to decreased alpha diversity (Chao1 index) and to a distinct community when compared to the pristine PE-MP exposure, in agreement with Kim et al. [80]. The stability

of the earthworm gut microbiome has been linked to health, pathogen defense, immunity, and metabolism; therefore, any derivation in the community composition can be seen as an indicator of contamination [81,82]. Compared to other microplastic exposures of terrestrial oligochaetes, earthworm gut microbiota richness and evenness was also shown to decrease when exposed to various sizes of PS-MPs ($0.1\text{--}100\text{ }\mu\text{m}$, $10\text{--}100\text{ mg kg}^{-1}$ soil) [33]. A similar response was observed in the gut microbiome of enchytraeids exposed to 10 % PS nanoplastics ($0.05\text{--}0.1\text{ }\mu\text{m}$) in oatmeal [83]. However, in an incubation mesocosm study of natural soil spiked with the equivalent of 340 mg kg^{-1} PE-MP placed on the surface saw no change in bacterial richness, diversity, or community composition in *Lumbricus terrestris* gut microbiota [84].

Aged microplastics have been shown to have more severe impacts on the gut microbiome of other organisms [85–88]. For earthworms, exposure to 500 mg kg^{-1} pristine and H_2O_2 -aged PE-MPs caused intestinal microbial disturbances, yet the effects were more severe with aged PE-MP exposure by promoting pathogenic bacteria and inhibiting probiotic bacteria [34]. Similarly, aged farmland residual PE mulch film MPs (0.25 %) impacted *Metaphire guillelmi* gut community structure compared to pristine PE-MP, and showed an enrichment in pathogenic bacteria [35].

Structural and functional damages in the gut have been linked to an imbalance in redox status (i.e. oxidative stress), where free radicals develop on the surface of MPs through photoaging in turn triggering cytotoxicity and greater reactive oxygen species (ROS) produced by the organism [89,90]. In relation to this, Zhu et al. [89] found a strong correlation between the oxidative potential of UV-aged MPs and increased carbonyl index or CI (namely conjugated carbonyl content). Accordingly, Jiang et al. [34] observed intestinal damage, ROS production, and shifted gut microbial communities in *E. fetida* after exposure to H_2O_2 aged PE-MP with increased CI. Similarly, UV-aged PE-MPs in this study, which also had increased CI, could have exerted higher oxidative potential in the gut lumen of *E. andrei* with negative effects on its gut microbiome. Increased oxidative process in the earthworm gut lumen has been associated with shifting dominance towards facultative anaerobes, aerobes and pathogenetic species and away from obligate anaerobes essential to earthworm gut health [91–95].

The most significant changes in *E. andrei* intestinal phyla absolute abundance, when exposed to UV-aged PE-MP, was the increasing abundances of Actinobacteria and Proteobacteria and decreasing ones of Firmicutes. Intestinal disorders have been characterized by a decline in bacterial diversity coupled with an increase in Actinobacteria and Proteobacteria [96]. As stated above, the chemical and physical alterations of UV-aged PE-MP might have promoted certain microbial colonization. Actinobacteria are known for their ability to degrade complex organic compounds, specifically with an efficiency in polyethylene-derived MP biodegradation [97], while Proteobacteria encompass diverse bacterial groups with versatile metabolic capabilities [98]. Both have been found to have enriched abundances on the surfaces of PE-MPs in soil [99]. Additionally, the presence of PE in soil has been shown to enrich Proteobacteria and Actinobacteria communities in the soil, and PE weathering favors the abundance of Proteobacteria [100]. Moreover, previous research has shown that field sourced PE-MPs with roughened and indented surfaces served as a suitable microhabitat for Actinobacteria and Proteobacteria, among other phyla [101]. Therefore, earthworms passively ingesting soil with aged-PE MPs would have gradually altered their gut microbial communities reflecting the change in the microbial composition occurring in the soil.

In the current study, there were increases in Proteobacteria absolute abundance with both pristine and UV-aged PE-MP as exposure concentrations rose but much more prominently with the aged material where a 135.5 % increase at the highest concentration was observed, in agreement with previous studies [88]. The increase in Proteobacteria with increasing PE-MP concentrations could also support the idea that gut microbiome analysis is a more sensitive toxicological indicator, as Proteobacterial increases serve as an indicator for gut dysbiosis [102].

Gut microbiome changes in *E. fetida* exposed to PS-MPs were most prominent with Proteobacteria growing more abundant [33,95]. Moreover, network analysis of gut microbial interaction in *E. fetida* revealed that most negative interactions (competition, amensalism, or predation) induced by PS-MP exposure were involved with increasing Proteobacteria, further hastening dysbiosis [103]. A specific genus of Proteobacteria (*Aeromonas*), a facultative anaerobe, was almost non-existent in *E. andrei* gut microbiome in the present study until it was exposed to the highest concentration of UV-aged PE-MP where it significantly increased absolute abundance with over a 500-fold increase. The genus *Aeromonas* was considered a main contributor to negative interactions in the gut microbiome network of *E. fetida*, mentioned above, after exposure to 100 mg kg⁻¹ PS-MPs (10 µm) [103] and caused the most severe intestinal damage, highest oxidative toxicity, immune system and energy metabolism disturbance and increased abundance of antibiotic-resistant genes [95]. This is in line with other studies indicating that increase in facultative anaerobes like *Aeromonas* [94,95], and simultaneously decreasing obligate anaerobes [93] is linked to gut complications.

Firmicutes, on the other hand, are obligate anaerobes and are rich in the oxygen depleted gut of Oligochaeta [104]. Firmicute absolute abundance declined with increasing UV-aged PE-MP exposure concentrations compared to control, but not with pristine PE-MP exposure. A study found that ozone-aged PS-MPs reduced the diversity of intestinal microbes, especially Firmicutes, while interfering with hormones related to energy metabolism in mice, compared to pristine PS-MPs [87]. In the present study, the concurrent increase in Proteobacteria as Firmicutes decreased was most notable during UV-aged PE-MP exposures, suggesting gut dysbiosis [93,105]. In addition, an important genus in the phylum Firmicutes, *Bacillus*, key in organic matter degradation and nutrient cycling [106], was significantly reduced when earthworms were exposed to UV-aged PE-MP compared to pristine. *Bacillus* suppression has been linked to C and N metabolic disorders [82], and when reintroduced to earthworm coelomic cells it promoted immunity regulation and antimicrobial defense [107]. This indicates that its inhibition by UV-aged PE-MP exposure in the present study possibly hindered *E. andrei* immune defenses and metabolism.

5. Conclusions

Pristine polyethylene microplastics at concentrations of 2 mg kg⁻¹ and above induced stimulation of reproduction in *E. andrei* while UV irradiated PE-MPs at 20 mg kg⁻¹ and above inhibited reproduction, this dichotomous response was significant at 20 mg kg⁻¹ for both PE-MP types. Furthermore, UV irradiated PE-MP exposure to *E. andrei* in soil not only decreased gut microbiome species richness and diversity, but also altered its balance in abundance when compared to the gut microbiome of earthworms exposed to pristine PE-MP, where demonstrably beneficial groups were decreased and potentially opportunistic and/or pathogenetic species were promoted at higher PE-MP concentrations. A positive correlation between gut microbiome function and reproduction has been seen in springtails exposed to PE-MPs [108], and enchytraeids exposed to nano-PS [83] and tire particle MPs [109]. In line with this, Ding et al. [109] confirmed a link between changes in gut bacterial communities and inhibited reproduction, which supports the idea that the gut microbiome is fundamentally tied to reproduction [110]. Other studies exploring the effects of various contaminants have also observed this trend [82,111], with one of them arguing that effects on the gut microbiome precede reproductive inhibition at lower contaminant concentrations [112], possibly being a precursor. This indicates that there could be an interplay between gut microbiome composition and reproduction of earthworms. Gut microbial imbalance can lead to health deterioration, such as increased susceptibility to pathogens and metabolic disorders [113]. Thus, because earthworm gut microbiome is inextricable from the roles they play in the surrounding soil environment, further investigation is necessary assess how the

toxicity of UV-aged microplastics could impact ecosystem functioning.

CRedit authorship contribution statement

Elise Quigley: Writing – review & editing, Writing – original draft, Visualization, Investigation, Formal analysis, Data curation. **Ana L. Patrício Silva:** Writing – review & editing, Validation, Resources, Methodology. **Sónia Chelinho:** Supervision, Methodology, Conceptualization. **Luís Cunha:** Writing – review & editing, Methodology. **Maria JI. Briones:** Writing – review & editing, Supervision, Project administration, Funding acquisition, Conceptualization. **José P. Sousa:** Writing – review & editing, Supervision, Resources, Project administration, Conceptualization.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Data will be made available on request.

Acknowledgments

This work was carried out within the SOPLAS project financed by the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No. 955334.

Special thanks to Patrícia Ferreira and Camila Campello for their help in implementing the experiment. Thank you to Carla Maria Nobre Maleita for allowing the use of her laboratory and equipment. Thank you to Daniel García Souto for providing statistical assistance.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ejsobi.2024.103640>.

References

- [1] T.R. Walker, L. Fequet, Current trends of unsustainable plastic production and micro(nano)plastic pollution, *TrAC, Trends Anal. Chem.* 160 (2023) 116984, <https://doi.org/10.1016/j.trac.2023.116984>.
- [2] P. Europe, Plastics – the facts 2022, plastics europe. <https://plasticseurope.org/knowledge-hub/plastics-the-facts-2022/>, 2022.
- [3] M. MacLeod, H.P.H. Arp, M.B. Tekman, A. Jahnke, The global threat from plastic pollution, *Science* 373 (2021) 61–65, <https://doi.org/10.1126/science.abg5433>.
- [4] J. Brahney, N. Mahowald, M. Prank, G. Cornwell, Z. Klimont, H. Matsui, K. A. Prather, Constraining the atmospheric limb of the plastic cycle, in: *Proceedings of the National Academy of Sciences*, vol. 118, 2021, <https://doi.org/10.1073/pnas.2020719118>.
- [5] A.A. de Souza Machado, W. Kloas, C. Zarfl, S. Hempel, M.C. Rillig, Microplastics as an emerging threat to terrestrial ecosystems, *Global Change Biol.* 24 (2018) 1405–1416, <https://doi.org/10.1111/gcb.14020>.
- [6] J. Gigault, A. Halle, M. Baudrimont, P.-Y. Pascal, F. Gauffre, T.-L. Phi, H.El Hadri, B. Grassl, S. Reynaud, Current opinion: what is a nanoplastic? *Environ. Pollut.* 235 (2018) 1030–1034, <https://doi.org/10.1016/j.envpol.2018.01.024>.
- [7] S. Fuller, A. Gautam, A procedure for measuring microplastics using pressurized fluid extraction, *Environ. Sci. Technol.* 50 (2016) 5774–5780, <https://doi.org/10.1021/acs.est.6b00816>.
- [8] F. Cheng, T. Zhang, Y. Liu, Y. Zhang, J. Qu, Non-negligible effects of UV irradiation on transformation and environmental risks of microplastics in the water environment, *Journal of Xenobiotics* 12 (2022) 1–12, <https://doi.org/10.3390/jox12010001>.
- [9] A.L. Andraday, The plastic in microplastics: a review, *Mar. Pollut. Bull.* 119 (2017) 12–22, <https://doi.org/10.1016/j.marpolbul.2017.01.082>.
- [10] K. Zhang, A.H. Hamidian, A. Tubić, Y. Zhang, J.K.H. Fang, C. Wu, P.K.S. Lam, Understanding plastic degradation and microplastic formation in the environment: a review, *Environ. Pollut.* 274 (2021) 116554, <https://doi.org/10.1016/j.envpol.2021.116554>.
- [11] C.A. Silva, A. Maria, J.L.T. Pestana, Ingestion of small-sized and irregularly shaped polyethylene microplastics affect *Chironomus riparius* life-history traits,

- Sci. Total Environ. 672 (2019) 862–868, <https://doi.org/10.1016/j.scitotenv.2019.04.017>.
- [12] J. Liu, T. Zhang, L. Tian, X. Liu, Z. Qi, Y. Ma, R. Ji, W. Chen, Aging significantly affects mobility and contaminant-mobilizing ability of nanoplastics in saturated loamy sand, Environ. Sci. Technol. 53 (2019) 5805–5815, <https://doi.org/10.1021/acs.est.9b00787>.
- [13] Z. Steinmetz, C. Wollmann, M. Schaefer, C. Buchmann, J. David, J. Tröger, K. Muñoz, O. Frör, G.E. Schaumann, Plastic mulching in agriculture. Trading short-term agronomic benefits for long-term soil degradation? Sci. Total Environ. 550 (2016) 690–705, <https://doi.org/10.1016/j.scitotenv.2016.01.153>.
- [14] A. Ammala, S. Bateman, K. Dean, E. Petinakis, P. Sangwan, S. Wong, Q. Yuan, L. Yu, C. Patrick, K.H. Leong, An overview of degradable and biodegradable polyolefins, Prog. Polym. Sci. 36 (2011) 1015–1049, <https://doi.org/10.1016/j.proppolymsci.2010.12.002>.
- [15] F. Vaccari, B. Forestieri, G. Papa, F. Bandini, E. Huerta-Lwanga, I. Boughattas, O. Missawi, M. Banni, I. Negri, P.S. Cocconcelli, E. Puglisi, Effects of micro and nanoplastics on soil fauna gut microbiome: an emerging ecological risk for soil health, Current Opin Environ. Sci. Health 30 (2022) 100402, <https://doi.org/10.1016/j.coesh.2022.100402>.
- [16] J. Li, Y. Song, Y. Cai, Focus topics on microplastics in soil: analytical methods, occurrence, transport, and ecological risks, Environ. Pollut. 257 (2020) 113570, <https://doi.org/10.1016/j.envpol.2019.113570>.
- [17] Y. Cao, M. Zhao, X. Ma, Y. Song, S. Zuo, H. Li, W. Deng, A critical review on the interactions of microplastics with heavy metals: mechanism and their combined effect on organisms and humans, Sci. Total Environ. 788 (2021) 147620, <https://doi.org/10.1016/j.scitotenv.2021.147620>.
- [18] T. Hüffer, A.-K. Weniger, T. Hofmann, Data on sorption of organic compounds by aged polystyrene microplastic particles, Data Brief 18 (2018) 474–479, <https://doi.org/10.1016/j.dib.2018.03.053>.
- [19] P. Liu, Y. Shi, X. Wu, H. Wang, H. Huang, X. Guo, S. Gao, Review of the artificially-accelerated aging technology and ecological risk of microplastics, Sci. Total Environ. 768 (2021) 144969, <https://doi.org/10.1016/j.scitotenv.2021.144969>.
- [20] P. Ding, C. Xiang, X. Li, H. Chen, X. Shi, X. Li, C. Huang, Y. Yu, J. Qi, A. Jing, G. Hu, Photoaged microplastics induce neurotoxicity via oxidative stress and abnormal neurotransmission in zebrafish larvae (*Danio rerio*), Sci. Total Environ. 881 (2023) 163480, <https://doi.org/10.1016/j.scitotenv.2023.163480>.
- [21] H. Yang, X. Li, M. Guo, X. Cao, X. Zheng, D. Bao, UV-induced microplastics (MPs) aging leads to comprehensive toxicity, Mar. Pollut. Bull. 189 (2023) 114745, <https://doi.org/10.1016/j.marpolbul.2023.114745>.
- [22] W. Song, C. Fu, Y. Fang, Z. Wang, J. Li, X. Zhang, K. Bhatt, L. Liu, N. Wang, F. Liu, S. Zhu, Single and combined toxicity assessment of primary or UV-aged microplastics and adsorbed organic pollutants on microalgae *Chlorella pyrenoidosa*, Environ. Pollut. 318 (2023) 120925, <https://doi.org/10.1016/j.envpol.2022.120925>.
- [23] M. Li, H. Jia, Q. Gao, S. Han, Y. Yu, L. Sun, Influence of aged and pristine polyethylene microplastics on bioavailability of three heavy metals in soil: toxic effects to earthworms (*Eisenia fetida*), Chemosphere 311 (2023) 136833, <https://doi.org/10.1016/j.chemosphere.2022.136833>.
- [24] Y. Cheng, L. Zhu, W. Song, C. Jiang, B. Li, Z. Du, J. Wang, J. Wang, D. Li, K. Zhang, Combined effects of mulch film-derived microplastics and atrazine on oxidative stress and gene expression in earthworm (*Eisenia fetida*), Sci. Total Environ. 746 (2020) 141280, <https://doi.org/10.1016/j.scitotenv.2020.141280>.
- [25] I. Boughattas, S. Hattab, N. Zitouni, M. Mkhinini, O. Missawi, N. Bousserhine, M. Banni, Assessing the presence of microplastic particles in Tunisian agriculture soils and their potential toxicity effects using *Eisenia andrei* as bioindicator, Sci. Total Environ. 796 (2021) 148959, <https://doi.org/10.1016/j.scitotenv.2021.148959>.
- [26] W. Cui, P. Gao, M. Zhang, L. Wang, H. Sun, C. Liu, Adverse effects of microplastics on earthworms: a critical review, Sci. Total Environ. 850 (2022) 158041, <https://doi.org/10.1016/j.scitotenv.2022.158041>.
- [27] J. Ding, D. Zhu, B. Hong, H.T. Wang, G. Li, Y.B. Ma, Y.T. Tang, Q.L. Chen, Long-term application of organic fertilization causes the accumulation of antibiotic resistance in earthworm gut microbiota, Environ. Int. 124 (2019) 145–152, <https://doi.org/10.1016/j.envint.2019.01.017>.
- [28] M. Sun, H. Zhao, X. Zheng, S. Deng, M. Ye, F. Hu, Ecological role of earthworm intestinal bacteria in terrestrial environments: a review, Sci. Total Environ. 740 (2020) 140008, <https://doi.org/10.1016/j.scitotenv.2020.140008>.
- [29] L.V. Hooper, D.R. Littman, A.J. Macpherson, Interactions between the microbiota and the immune system, Science 336 (2012) 1268–1273, <https://doi.org/10.1038/s41422-020-0332-7>.
- [30] K.A. Khan, M.J. Ansari, A. Al-Ghamdi, A. Nuru, S. Harakeh, J. Iqbal, Investigation of gut microbial communities associated with indigenous honey bee (*Apis mellifera jemenitica*) from two different eco-regions of Saudi Arabia, Saudi J. Biol. Sci. 24 (2017) 1061–1068, <https://doi.org/10.1016/j.sjbs.2017.01.055>.
- [31] J. Ding, C. Liu, Q. Chen, Z. Zhang, J. Han, A. Liang, D. Zhu, H. Wang, M. Lv, L. Chen, Extractable additives in microplastics: a hidden threat to soil fauna, Environ. Pollut. 294 (2022) 118647, <https://doi.org/10.1016/j.envpol.2021.118647>.
- [32] R. Tang, D. Zhu, Y. Luo, D. He, H. Zhang, A. El-Naggar, K.N. Palansooriya, K. Chen, Y. Yan, X. Lu, M. Ying, T. Sun, Y. Cao, Z. Diao, Y. Zhang, Y. Lian, S. X. Chang, Y. Cai, Nanoplastics induce molecular toxicity in earthworm: integrated multi-omics, morphological, and intestinal microorganism analyses, J. Hazard Mater. 442 (2023) 130034, <https://doi.org/10.1016/j.jhazmat.2022.130034>.
- [33] Y. Yang, G. Xu, Y. Yu, Microplastics impact the accumulation of metals in earthworms by changing the gut bacterial communities, Sci. Total Environ. 831 (2022) 154848, <https://doi.org/10.1016/j.scitotenv.2022.154848>.
- [34] X. Jiang, J. Cao, Z. Ye, G. Klobučar, M. Li, Microplastics - back to reality: impact of pristine and aged microplastics in soil on earthworm *Eisenia fetida* under environmentally relevant conditions, Environ. Sci. Technol. 57–44 (2023) 16788–16799, <https://doi.org/10.1021/acs.est.3c04097>.
- [35] H. Tian, C. Zheng, X. Huang, C. Qi, B. Li, Z. Du, L. Zhu, J. Wang, J. Wang, Effects of farmland residual mulch film-derived microplastics on the structure and function of soil and earthworm *Metaphire guillelmi* gut microbiota, Sci. Total Environ. 915 (2024) 170094, <https://doi.org/10.1016/j.scitotenv.2024.170094>.
- [36] L. Tian, C. Jinjin, R. Ji, Y. Ma, X. Yu, Microplastics in agricultural soils: sources, effects, and their fate, Current Opin Environ. Sci. Health 25 (2022) 100311, <https://doi.org/10.3389/fenvs.2022.855292>.
- [37] A.A. Horton, A. Walton, D.J. Spurgeon, E. Lahive, C. Svendsen, Microplastics in freshwater and terrestrial environments: evaluating the current understanding to identify the knowledge gaps and future research priorities, Sci. Total Environ. 586 (2017) 127–141, <https://doi.org/10.1016/j.scitotenv.2017.01.190>.
- [38] I.S. O, Soil Quality - Effects of Pollutants on Earthworms - Part 2: Determination of Effects on Reproduction to *Eisenia fetida*/*Eisenia andrei*, International Organization for Standardization, 2012.
- [39] Y. Chen, X. Liu, Y. Leng, J. Wang, Defense responses in earthworms (*Eisenia fetida*) exposed to low-density polyethylene microplastics in soils, Ecotoxicol. Environ. Saf. 187 (2020) 109788, <https://doi.org/10.1016/j.ecoenv.2019.109788>.
- [40] E. Huerta Lwanga, H. Gertsen, H. Gooren, P. Peters, T. Salánki, M. Ploeg, E. Besseling, A.A. Koelmans, V. Geissen, Microplastics in the terrestrial ecosystem: implications for *Lumbricus terrestris* (Oligochaeta, Lumbricidae), Environ. Sci. Technol. 50 (2016) 2685–2691, <https://doi.org/10.1021/acs.est.5b05478>.
- [41] Y. Zhou, X. Liu, J. Wang, Ecotoxicological effects of microplastics and cadmium on the earthworm *Eisenia foetida*, J. Hazard Mater. 392 (2020) 122273, <https://doi.org/10.1016/j.jhazmat.2020.122273>.
- [42] F. Reis, E. Nascimento, H. Castro, C. Canhoto, A.L. Gonçalves, S. Simões, P. García-Palacios, R. Milla, J.P. Sousa, P. Silva, Land management impacts on the feeding preferences of the woodlouse *Porcellio dilatatus* (Isopoda: oniscidea) via changes in plant litter quality, Appl. Soil Ecol.: Sect. Agricult.Ecosyst. Environ. 132 (2018) 45–52, <https://doi.org/10.1016/j.apsoil.2018.08.018>.
- [43] I.S. O, Soil Quality - Inhibition of Reproduction of *Collembola (Folsomia candida)* by Soil Pollutants, International Organization for Standardization, 1999.
- [44] Y. Zhou, J. Wang, M. Zou, Z. Jia, S. Zhou, Y. Li, Microplastics in soils: a review of methods, occurrence, fate, transport, ecological and environmental risks, Sci. Total Environ. 748 (2020) 141368, <https://doi.org/10.1016/j.scitotenv.2020.141368>.
- [45] F. Büks, M. Kaupenjohann, Global concentrations of microplastics in soils – a review, Soil 6 (2020) 649–662, <https://doi.org/10.5194/soil-6-649-2020>.
- [46] J.C. Prata, I.F. Sequeira, S.S. Monteiro, A.L.P. Silva, J.P. Costa, P. Dias-Pereira, A. J.S. Fernandes, F.M. Costa, A.C. Duarte, T. Rocha-Santos, Preparation of biological samples for microplastic identification by Nile Red, Sci. Total Environ. 783 (2021) 147065, <https://doi.org/10.1016/j.scitotenv.2021.147065>.
- [47] A.L.P. Silva, S.A.M. Silva, A. Duarte, D. Barceló, T. Rocha-Santos, Analytical methodologies used for screening micro(nano)plastics in (eco)toxicity tests, Green, Anal. Chem. 3 (2022) 100037, <https://doi.org/10.1016/j.greac.2022.100037>.
- [48] A.K. Rodriguez, B. Mansoor, G. Ayoub, X. Colin, A.A. Benzerga, Effect of UV-aging on the mechanical and fracture behavior of low density polyethylene, Polym. Degrad. Stabil. 180 (2020) 109185, <https://doi.org/10.1016/j.polymerdegradstab.2020.109185>.
- [49] J. Almond, P. Sugumaar, M.N. Wenzel, G. Hill, C. Wallis, Determination of the carbonyl index of polyethylene and polypropylene using specified area under band methodology with ATR-FTIR spectroscopy, E-Polymers 20 (2020) 369–381, <https://doi.org/10.1515/epoly-2020-0041>.
- [50] B.J. Callahan, P.J. McMurdie, M.J. Rosen, A.W. Han, A.J.A. Johnson, S.P. Holmes, DADA2: high-resolution sample inference from Illumina amplicon data, Nat. Methods 13 (2016) 581–583, <https://doi.org/10.1038/nmeth.3869>.
- [51] B.J. Callahan, P.J. McMurdie, S.P. Holmes, Exact sequence variants should replace operational taxonomic units in marker-gene data analysis, ISME J. 11 (2017) 2639–2643, <https://doi.org/10.1038/ismej.2017.119>.
- [52] T. Magoč, S.L. Salzberg, FLASH: fast length adjustment of short reads to improve genome assemblies, Bioinformatics 27 (2011) 2957–2963, <https://doi.org/10.1093/bioinformatics/btr507>.
- [53] S. Chen, Y. Zhou, Y. Chen, J. Gu, Fastp: an ultra-fast all-in-one fastq preprocessor, Bioinformatics 34 (2018) i884–i890, <https://doi.org/10.1093/bioinformatics/bty560>.
- [54] N.A. Bokulich, B.D. Kaehler, J.R. Rideout, M. Dillon, E. Bolyen, R. Knight, G. A. Huttley, G. Caporaso, Optimizing taxonomic classification of marker-gene amplicon sequences with QIIME 2's q2-feature-classifier plugin, Microbiome 6 (2018) 90, <https://doi.org/10.1186/s40168-018-0470-z>.
- [55] E. Bolyen, J.R. Rideout, M.R. Dillon, N.A. Bokulich, C.C. Abnet, G.A. Al-Ghalith, H. Alexander, E.J. Alm, M. Arumugam, F. Asnicar, Y. Bai, J.E. Bisanz, K. Bittinger, A. Brejnrod, C.J. Brislawn, C.T. Brown, B.J. Callahan, A.M. Caraballo-Rodríguez, J. Chase, J.G. Caporaso, Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2, Nat. Biotechnol. 37 (2019) 852–857, <https://doi.org/10.1038/s41587-019-0209-9>.

- [56] C. Lozupone, R. Knight, UniFrac: a new phylogenetic method for comparing microbial communities, *Appl. Environ. Microbiol.* 71 (2005) 8228–8235, <https://doi.org/10.1128/AEM.71.12.8228-8235.2005>.
- [57] A. Lozupone Catherine, H. Micah, K.S. T., K. Rob, Quantitative and qualitative β diversity measures lead to different insights into factors that structure microbial communities, *Appl. Environ. Microbiol.* 73 (2007) 1576–1585, <https://doi.org/10.1128/AEM.01996-06>.
- [58] H. Wickham, *ggplot2: Elegant Graphics for Data Analysis*, Springer-Verlag, New York, 2016, <https://ggplot2.tidyverse.org>.
- [59] M. Liu, J. Feng, Y. Shen, B. Zhu, Microplastics effects on soil biota are dependent on their properties: a meta-analysis, *Soil Biol. Biochem.* 178 (2023) 108940, <https://doi.org/10.1016/j.soilbio.2023.108940>.
- [60] T. Trakić, F. Popović, J. Sekulić, D.K. Hackenberger, Ecotoxicological effects of commercial microplastics on earthworm *Eisenia fetida*, Savigny, 1826) (Clitellata; Lumbricidae), *Agriculture* 14 (2024) 267, <https://doi.org/10.3390/agriculture14020267>.
- [61] D. Angmo, R. Dutta, J. Singh, A.B. Chowdhary, J. Quadar, B. Thakur, H. Kaur, M. Sharma, S. Singh, A.P. Vig, Biochemical responses, growth and reproduction of earthworm in low density polyethylene (LDPE), *Environ. Qual. Manag.* 33–1 (2023) 223–237, <https://doi.org/10.1002/tqem.22041>.
- [62] Z. Sobhani, C. Fang, R. Naidu, M. Megharaj, Microplastics as a vector of toxic chemicals in soil: enhanced uptake of perfluorooctane sulfonate and perfluorooctanoic acid by earthworms through sorption and reproductive toxicity, *Environ. Technol. Innovat.* 22 (2021) 101476, <https://doi.org/10.1016/j.eti.2021.101476>.
- [63] Z. Sobhani, L. Panneerselvam, C. Fang, R. Naidu, M. Megharaj, Chronic and transgenerational effects of polystyrene microplastics at environmentally relevant concentrations in earthworms (*Eisenia fetida*), *Environ. Toxicol. Chem./SETAC* 40 (2021) 2240–2246, <https://doi.org/10.1016/j.eti.2021.102226Get.rightsandcontent>.
- [64] A. Rodríguez-Seijo, J. Lourenço, T.A.P. Rocha-Santos, J. Costa, A.C. Duarte, H. Vala, R. Pereira, Histopathological and molecular effects of microplastics in *Eisenia andrei* Bouché, *Environ. Pollut.* 220 (2017) 495–503, <https://doi.org/10.1016/j.envpol.2016.09.092>.
- [65] T.B. Masset, F. Breider, W. Duféoi, K. Schirmer, A. Bergmann, E. Vermeirssen, B. J.D. Ferrari (Eds.), Effects of Tire Particles on the Reproduction of Earthworms (*Eisenia fetida*) and Bioaccumulation Potential of Tire Related Chemicals, 2022. <https://infoscience.epfl.ch/record/292168?ln=en>. (Accessed 30 November 2023).
- [66] T. Mondal, E.H. Jho, S.K. Hwang, Y. Hyeon, C. Park, Responses of earthworms exposed to low-density polyethylene microplastic fragments, *Chemosphere* 333 (2023) 138945, <https://doi.org/10.1016/j.chemosphere.2023.138945>.
- [67] X. Yang, X. Zhang, X. Shu, J. Gong, J. Yang, B. Li, J. Lin, Y. Chai, J. Liu, The effects of polyethylene microplastics on the growth, reproduction, metabolic enzymes, and metabolomics of earthworms *Eisenia fetida*, *Ecotoxicol. Environ. Saf.* 263 (2023) 115390, <https://doi.org/10.1016/j.ecoenv.2023.115390>.
- [68] R.R. Rix, R.N.C. Guedes, C. Cutler, Hormesis dose–response contaminant-induced hormesis in animals, *Current Opinion Toxicol.* 30 (2022) 100336, <https://doi.org/10.1016/j.cotox.2022.02.009>.
- [69] T. Sun, J. Zhan, F. Li, C. Ji, H. Wu, Effect of microplastics on aquatic biota: a hormetic perspective, *Environ. Pollut.* 285 (2021) 117206, <https://doi.org/10.1016/j.envpol.2021.117206>.
- [70] Q. Chen, Y. Li, B. Li, Is color a matter of concern during microplastic exposure to *Scenedesmus obliquus* and *Daphnia magna*? *J. Hazard Mater.* 383 (2020) 121224, <https://doi.org/10.1016/j.jhazmat.2019.121224>.
- [71] X. Sun, B. Chen, Q. Li, N. Liu, B. Xia, L. Zhu, K. Qu, Toxicities of polystyrene nano- and microplastics toward marine bacterium *Halomonas alkalicola*, *Sci. Total Environ.* 642 (2018) 1378–1385, <https://doi.org/10.1016/j.scitotenv.2018.06.141>.
- [72] L. Jia, X. He, W. Chen, Z. Liu, Y. Huang, S. Yu, Hormesis phenomena under Cd stress in a hyperaccumulator–*Lonicera japonica* Thunb, *Ecotoxicology* 22 (2013) 476–485, <https://doi.org/10.1007/s10646-013-1041-5>.
- [73] E. Agathokleous, I. Iavicoli, D. Barceló, E.J. Calabrese, Micro/nanoplastics effects on organisms: a review focusing on “dose”, *J. Hazard Mater.* 417 (2021) 126084, <https://doi.org/10.1016/j.jhazmat.2021.126084>.
- [74] H. Cui, W. Yang, Y. Cui, L. Qi, X. Jiang, M. Li, Adverse effects of pristine and aged polystyrene microplastics in mice and their Nrf2-mediated defense mechanisms with tissue specificity, *Environ. Sci. Pollut. Res. Int.* 30 (2023) 39894–39906, <https://doi.org/10.1007/s11356-022-24918-1>.
- [75] H. Chen, Y. Yang, C. Wang, X. Hua, H. Li, D. Xie, M. Xiang, Y. Yu, Reproductive toxicity of UV-photodegraded polystyrene microplastics induced by DNA damage-dependent cell apoptosis in *Caenorhabditis elegans*, *Sci. Total Environ.* 811 (2022) 152350, <https://doi.org/10.1016/j.scitotenv.2021.152350>.
- [76] Z. Liu, Y. Zhu, S. Lv, Y. Shi, S. Dong, D. Yan, X. Zhu, R. Peng, A.A. Keller, Y. Huang, Quantifying the dynamics of polystyrene microplastics UV-Aging process, *Environ. Sci. Technol. Lett.* 9 (2022) 50–56, <https://doi.org/10.1021/acs.estlett.1c00888>.
- [77] Q. Wang, X. Wangjin, Y. Zhang, N. Wang, Y. Wang, G. Meng, Y. Chen, The toxicity of virgin and UV-aged PVC microplastics on the growth of freshwater algae *Chlamydomonas reinhardtii*, *Sci. Total Environ.* 749 (2020) 141603, <https://doi.org/10.1016/j.scitotenv.2020.141603>.
- [78] H. Luo, C. Liu, D. He, J. Xu, J. Sun, J. Li, X. Pan, Environmental behaviors of microplastics in aquatic systems: a systematic review on degradation, adsorption, toxicity and biofilm under aging conditions, *J. Hazard Mater.* 423 (2022) 126915, <https://doi.org/10.1016/j.jhazmat.2021.126915>.
- [79] M. Celik, H. Nakano, K. Uchida, A. Isobe, H. Arakawa, Comparative evaluation of the carbonyl index of microplastics around the Japan coast, *Mar. Pollut. Bull.* 190 (2023) 114818, <https://doi.org/10.1016/j.marpolbul.2023.114818>.
- [80] B.-R. Kim, J. Shin, R. Guevarra, J.H. Lee, D.W. Kim, K.-H. Seol, J.-H. Lee, H. B. Kim, R. Isaacson, Deciphering diversity indices for a better understanding of microbial communities, *J. Microbiol. Biotechnol.* 27 (2017) 2089–2093, <https://doi.org/10.4014/jmb.1709.09027>.
- [81] D.A. Pass, A.J. Morgan, D.S. Read, D. Field, A.J. Weightman, P. Kille, The effect of anthropogenic arsenic contamination on the earthworm microbiome, *Environ. Microbiol.* 17 (2015) 1884–1896, <https://doi.org/10.1111/1462-2920.12712>.
- [82] L. Ma, Y. Xie, Z. Han, J.P. Giesy, X. Zhang, Responses of earthworms and microbial communities in their guts to Triclosan, *Chemosphere* 168 (2017) 1194–1202, <https://doi.org/10.1016/j.chemosphere.2016.10.079>.
- [83] B.-K. Zhu, Y.-M. Fang, D. Zhu, P. Christie, X. Ke, Y.-G. Zhu, Exposure to nanoplastics disturbs the gut microbiome in the soil oligochaete *Enchytraeus crypticus*, *Environ. Pollut.* 239 (2018) 408–415, <https://doi.org/10.1016/j.envpol.2018.04.017>.
- [84] K. Adhikari, A.F. Astner, J.M. DeBruyn, Y. Yu, D.G. Hayes, B.T. O’Callahan, M. Flury, Earthworms exposed to polyethylene and biodegradable microplastics in soil: microplastic characterization and microbial community analysis, *ACS Agricult. Sci. Technol.* 3–4 (2023) 340–349, <https://doi.org/10.1021/acscagritech.2c00333>.
- [85] P. Zhang, G. Lu, Y. Sun, J. Zhang, J. Liu, Z. Yan, Aged microplastics change the toxicological mechanism of roxithromycin on *Carassius auratus*: size-dependent interaction and potential long-term effects, *Environ. Int.* 169 (2022) 107540, <https://doi.org/10.1016/j.envint.2022.107540>.
- [86] P. Zhang, G. Lu, Y. Sun, Z. Yan, T. Dang, J. Liu, Metagenomic analysis explores the interaction of aged microplastics and roxithromycin on gut microbiota and antibiotic resistance genes of *Carassius auratus*, *J. Hazard Mater.* 425 (2022) 127773, <https://doi.org/10.1016/j.jhazmat.2021.127773>.
- [87] J. Jin, W. Biying, W. Di, A comparative study on the effects of short-term oral exposure to ozone-aged polystyrene microplastics and standard polystyrene microplastic beads on intestinal flora structure and energy metabolism-related hormones in mice, *Asian J. Ecotoxicol.* 17 (2022) 95–105, <https://doi.org/10.7524/AJE.1673-5897.20220411003>.
- [88] Y. Zhong, Q. Ding, Z. Huang, X. Xiao, X. Han, Y. Su, D. Wang, J. You, Influence of ultraviolet-aging and adsorbed pollutants on toxicological effects of polyvinyl chloride microplastics to zebrafish, *Environ. Pollut.* 316 (2023) 120617, <https://doi.org/10.1016/j.envpol.2022.120617>.
- [89] K. Zhu, H. Jia, Y. Sun, Y. Dai, C. Zhang, X. Guo, T. Wang, L. Zhu, Enhanced cytotoxicity of photoaged phenol-formaldehyde resins microplastics: combined effects of environmentally persistent free radicals, reactive oxygen species, and conjugated carbonyls, *Environ. Int.* 145 (2020) 106137, <https://doi.org/10.1016/j.envint.2020.106137>.
- [90] X. Zhang, M. Xia, J. Zhao, Z. Cao, W. Zou, Q. Zhou, Photoaging enhanced the adverse effects of polyamide microplastics on the growth, intestinal health, and lipid absorption in developing zebrafish, *Environ. Int.* 158 (2022) 106922, <https://doi.org/10.1016/j.envint.2021.106922>.
- [91] F. Rivera-Chávez, C.A. Lopez, A.J. Bäuml, Oxygen as a driver of gut dysbiosis, *Free Radical Biol. Med.* 105 (2017) 93–101, <https://doi.org/10.1016/j.freeradbiomed.2016.09.022>.
- [92] G. Rizzatti, L.R. Lopetuso, G. Gibiino, C. Binda, A. Gasbarrini, Proteobacteria: a common factor in human diseases, *BioMed Res. Int.* (2017) 9351507, <https://doi.org/10.1155/2017/9351507>.
- [93] L. Rigottier-Gois, Dysbiosis in inflammatory bowel diseases: the oxygen hypothesis, *ISME J.* 7 (2013) 1256–1261, <https://doi.org/10.1038/ismej.2013.80>.
- [94] Z. Xing, H. Li, M. Li, R. Gao, C. Guo, S. Mi, Disequilibrium in chicken gut microflora with avian colibacillosis is related to microenvironment damaged by antibiotics, *Sci. Total Environ.* 762 (2020) 143058, <https://doi.org/10.1016/j.scitotenv.2020.143058>.
- [95] G. Xu, Y. Yu, Polystyrene microplastics impact the occurrence of antibiotic resistance genes in earthworms by size-dependent toxic effects, *J. Hazard Mater.* 416 (2021) 125847, <https://doi.org/10.1016/j.jhazmat.2021.125847>.
- [96] P. Lepage, R. Häslér, M.E. Spehlmann, A. Rehman, A. Zvirbliene, A. Begun, S. Ott, L. Kupcinskas, J. Doré, A. Raedler, S. Schreiber, Twin study indicates loss of interaction between microbiota and mucosa of patients with ulcerative colitis, *Gastroenterology* 141 (2011) 227–236, <https://doi.org/10.1053/j.gastro.2011.04.011>.
- [97] A. Parsaeimehr, C.M. Miller, G. Ozbay, Microplastics and their interactions with microbiota, *Heliyon* 9 (2023) 15104, <https://doi.org/10.1016/j.heliyon.2023.e15104>.
- [98] A. Gupta, R. Gupta, R.L. Singh, Microbes and environment, in: R.L. Singh (Ed.), Principles and Applications of Environmental Biotechnology for a Sustainable Future, Springer Singapore, 2017, pp. 43–84, https://doi.org/10.1007/978-981-10-1866-4_3.
- [99] H. Ya, Y. Xing, T. Zhang, M. Lv, B. Jiang, LDPE microplastics affect soil microbial community and form a unique plastisphere on microplastics, *Appl. Soil Ecol.: Sect. Agricult. Ecosyst. Environ.* 180 (2022) 104623, <https://doi.org/10.1016/j.apsoil.2022.104623>.
- [100] D. Huang, Y. Xu, F. Lei, X. Yu, Z. Ouyang, Y. Chen, H. Jia, X. Guo, Degradation of polyethylene plastic in soil and effects on microbial community composition, *J. Hazard Mater.* 416 (2021) 126173, <https://doi.org/10.1016/j.jhazmat.2021.126173>.

- [101] M. Zhang, Y. Zhao, X. Qin, W. Jia, L. Chai, M. Huang, Y. Huang, Microplastics from mulching film is a distinct habitat for bacteria in farmland soil, *Sci. Total Environ.* 688 (2019) 470–478, <https://doi.org/10.1016/j.scitotenv.2019.06.108>.
- [102] I. Mukhopadhyay, R. Hansen, E.M. El-Omar, G.L. Hold, IBD-what role do Proteobacteria play? *Nat. Rev. Gastroenterol. Hepatol.* 9 (2012) 219–230, <https://doi.org/10.1038/nrgastro.2012.14>.
- [103] T. Li, M. Lu, B. Xu, H. Chen, J. Li, Z. Zhu, M. Yu, J. Zheng, P. Peng, S. Wu, Multiple perspectives reveal the gut toxicity of polystyrene microplastics on *Eisenia fetida*: insights into community signatures of gut bacteria and their translocation, *Sci. Total Environ.* 838 (2022) 156352, <https://doi.org/10.1016/j.scitotenv.2022.156352>.
- [104] R. Sapkota, S. Santos, P. Farias, P.H. Krogh, A. Winding, Insights into the earthworm gut multi-kingdom microbial communities, *Sci. Total Environ.* 727 (2020) 138301, <https://doi.org/10.1016/j.scitotenv.2020.138301>.
- [105] Q. He, L. Wang, F. Wang, C. Wang, C. Tang, Q. Li, J. Li, Q. Zhao, Microbial fingerprinting detects intestinal microbiota dysbiosis in Zebrafish models with chemically-induced enterocolitis, *BMC Microbiol.* 13 (2013) 289, <https://doi.org/10.1186/1471-2180-13-289>.
- [106] I. Mandic-Mulec, P. Stefanic, J.D. Elsas, Ecology of Bacillaceae, *Microbiol. Spectr.* 3 (2015) 17–2013, <https://doi.org/10.1128/microbiolspec.tbs-0017-2013>.
- [107] E.S. Tak, S.-J. Cho, S.C. Park, Gene expression profiling of coelomic cells and discovery of immune-related genes in the earthworm, *Eisenia andrei*, using expressed sequence tags, *Biosci., Biotechnol., Biochem.* 79 (2015) 367–373, <https://doi.org/10.1080/09168451.2014.988677>.
- [108] H. Ju, D. Zhu, M. Qiao, Effects of polyethylene microplastics on the gut microbial community, reproduction and avoidance behaviors of the soil springtail, *Folsomia candida*, *Environ. Pollut.* 247 (2019) 890–897, <https://doi.org/10.1016/j.envpol.2019.01.097>.
- [109] J. Ding, D. Zhu, H.-T. Wang, S.B. Lassen, Q.-L. Chen, G. Li, M. Lv, Y.-G. Zhu, Dysbiosis in the gut microbiota of soil fauna explains the toxicity of tire tread particles, *Environ. Sci. Technol.* 54 (2020) 7450–7460, <https://doi.org/10.1021/acs.est.0c00917>.
- [110] P. Lavelle, C. Lattaud, D. Trigo, I. Barois, Mutualism and biodiversity in soils, *Plant Soil* 170 (1995) 23–33, <https://doi.org/10.1007/BF02183052>.
- [111] D. Zhu, F. Zheng, Q.-L. Chen, X.-R. Yang, P. Christie, X. Ke, Y.-G. Zhu, Exposure of a soil collembolan to Ag nanoparticles and AgNO₃ disturbs its associated microbiota and lowers the incidence of antibiotic resistance genes in the gut, *Environ. Sci. Technol.* 52 (2018) 12748–12756, <https://doi.org/10.1021/acs.est.8b02825>.
- [112] E. Swart, T. Goodall, P. Kille, D.J. Spurgeon, C. Svendsen, The earthworm microbiome is resilient to exposure to biocidal metal nanoparticles, *Environ. Pollut.* 267 (2020) 115633, <https://doi.org/10.1016/j.envpol.2020.115633>.
- [113] A.K. DeGruttola, D. Low, A. Mizoguchi, E. Mizoguchi, Current understanding of dysbiosis in disease in human and animal models, *Inflamm. Bowel Dis.* 22 (2016) 1137–1150, <https://doi.org/10.1097/MIB.0000000000000750>.