


Resolutions of matroid ideals

Luis David Garcia-Puente


lgarcia@math.vt.edu

Virginia Tech

- Let $M = \langle a^2b, ac, b^2, bc^2 \rangle$ **monomial ideal** in $S = \mathbf{k}[a, b, c]$.
- Let X be the **finite regular CW-complex**:




- Let $M = \langle a^2b, ac, b^2, bc^2 \rangle$ **monomial ideal** in $S = \mathbf{k}[a, b, c]$.
- Let X be the **finite regular CW-complex**:



- Let \mathbb{F}_X be the **complex** of \mathbb{Z}^n -graded free S -modules


$$\begin{aligned}
 0 \rightarrow & S[-(2, 2, 1)] \oplus S[-(1, 2, 2)] \xrightarrow{\partial_2} S[-(2, 1, 1)] \oplus S[-(2, 2, 0)] \oplus \\
 & S[-(1, 2, 1)] \oplus S[-(1, 1, 2)] \oplus S[-(0, 2, 2)] \xrightarrow{\partial_1} S[-(2, 1, 0)] \oplus \\
 & S[-(1, 0, 1)] \oplus S[-(0, 1, 2)] \oplus S[-(0, 2, 0)] \xrightarrow{\partial_0} S
 \end{aligned}$$



● The differential ∂ acts on basis vectors

$$\partial(a^2b^2c) = -b \cdot a^2bc + c \cdot a^2b^2 - a \cdot ab^2c,$$

$$\partial(F) = \sum_{\text{facets } G \text{ of } F} \text{sign}(G, F) \frac{\mathbf{x}^{\mathbf{a}_F}}{\mathbf{x}^{\mathbf{a}_G}} G.$$



● The differential ∂ acts on basis vectors


$$\partial(a^2b^2c) = -b \cdot a^2bc + c \cdot a^2b^2 - a \cdot ab^2c,$$

$$\partial(F) = \sum_{\text{facets } G \text{ of } F} \text{sign}(G, F) \frac{\mathbf{x}^{\mathbf{a}_F}}{\mathbf{x}^{\mathbf{a}_G}} G.$$

Observe $\partial_0 = \begin{bmatrix} a^2b & ac & bc^2 & b^2 \end{bmatrix}$. Thus $\text{Coker}(\partial_0) = S/M$.


For $\mathbf{b} \in \mathbb{N}^n$, let $X_{\preceq \mathbf{b}}$ be the subcomplex of X consisting of all faces whose degrees are coordinatewise at most \mathbf{b} .

Example. Let $\mathbf{b} = (1, 1, 2)$, then $X_{\preceq \mathbf{b}}$ is the subcomplex



For $\mathbf{b} \in \mathbb{N}^n$, let $X_{\preceq \mathbf{b}}$ be the subcomplex of X consisting of all faces whose degrees are coordinatewise at most \mathbf{b} .

Example. Let $\mathbf{b} = (1, 1, 2)$, then $X_{\preceq \mathbf{b}}$ is the subcomplex



Theorem. \mathbb{F}_X is *exact* if and only if $X_{\preceq \mathbf{b}}$ is *acyclic* over k for all $\mathbf{b} \in \mathbb{N}^n$.

For $\mathbf{b} \in \mathbb{N}^n$, let $X_{\preceq \mathbf{b}}$ be the subcomplex of X consisting of all faces whose degrees are coordinatewise at most \mathbf{b} .

Example. Let $\mathbf{b} = (1, 1, 2)$, then $X_{\preceq \mathbf{b}}$ is the subcomplex $\bullet \xrightarrow{ac} \bullet \xleftarrow{abc^2} \bullet \xleftarrow{bc^2}$

Theorem. \mathbb{F}_X is *exact* if and only if $X_{\preceq \mathbf{b}}$ is *acyclic* over \mathbf{k} for all $\mathbf{b} \in \mathbb{N}^n$.

● The \mathbb{Z}^n -graded degree \mathbf{b} part of \mathbb{F}_X is the *chain complex*

$$0 \rightarrow \mathbf{k} \cdot 1 \rightarrow \mathbf{k} \cdot bc \oplus \mathbf{k} \cdot a \rightarrow \mathbf{k} \cdot abc^2$$

● ∂ is a degree 0 map. Then $(\mathbb{F}_X)_{\mathbf{b}}$ is the *reduced chain complex*

$$\tilde{C}(X_{\preceq \mathbf{b}}; \mathbf{k}) = 0 \rightarrow \mathbf{k} \rightarrow \mathbf{k}^2 \rightarrow \mathbf{k}$$

For $\mathbf{b} \in \mathbb{N}^n$, let $X_{\preceq \mathbf{b}}$ be the subcomplex of X consisting of all faces whose degrees are coordinatewise at most \mathbf{b} .

Example. Let $\mathbf{b} = (1, 1, 2)$, then $X_{\preceq \mathbf{b}}$ is the subcomplex $\bullet \xrightarrow{ac} \bullet \xleftarrow{abc^2} \bullet \xleftarrow{bc^2}$

Theorem. \mathbb{F}_X is *exact* if and only if $X_{\preceq \mathbf{b}}$ is *acyclic* over \mathbf{k} for all $\mathbf{b} \in \mathbb{N}^n$.

• The \mathbb{Z}^n -graded degree \mathbf{b} part of \mathbb{F}_X is the *chain complex*

$$0 \rightarrow \mathbf{k} \cdot 1 \rightarrow \mathbf{k} \cdot bc \oplus \mathbf{k} \cdot a \rightarrow \mathbf{k} \cdot abc^2$$

• ∂ is a degree 0 map. Then $(\mathbb{F}_X)_{\mathbf{b}}$ is the *reduced chain complex*

$$\tilde{C}(X_{\preceq \mathbf{b}}; \mathbf{k}) = 0 \rightarrow \mathbf{k} \rightarrow \mathbf{k}^2 \rightarrow \mathbf{k}$$

• $X_{\preceq \mathbf{b}}$ is *contractible*, so it has no reduced homology.

An ideal M is a **matroid ideal** if and only if the following conditions hold


- M is a square-free monomial ideal,
- for every pair of monomials $m_1, m_2 \in M$ and any $i \in \{1, \dots, n\}$ such that x_i divides both m_1 and m_2 , the monomial $\text{lcm}(m_1, m_2)/x_i$ is in M as well.

An ideal M is a **matroid ideal** if and only if the following conditions hold


- M is a square-free monomial ideal,
- for every pair of monomials $m_1, m_2 \in M$ and any $i \in \{1, \dots, n\}$ such that x_i divides both m_1 and m_2 , the monomial $\text{lcm}(m_1, m_2)/x_i$ is in M as well.

$$M = \langle x_1x_2, x_1x_3x_4, x_1x_3x_5, x_2x_3x_4, x_2x_3x_5, x_4x_5 \rangle.$$

Let \mathcal{A} be the hyperplane arrangement in \mathbb{R}^2




Let \mathcal{A} be the hyperplane arrangement in \mathbb{R}^2



- The hyperplanes partition \mathbb{R}^2 into **cells**.
- The set of **bounded cells** forms a cell complex, denoted $B_{\mathcal{A}}$, which resolves $S/M_{\mathcal{A}}$ minimally.

Let \mathcal{A} be the hyperplane arrangement in \mathbb{R}^2



- The hyperplanes partition \mathbb{R}^2 into **cells**.
- The set of **bounded cells** forms a cell complex, denoted $B_{\mathcal{A}}$, which resolves $S/M_{\mathcal{A}}$ minimally.
- Hence $0 \longrightarrow S^4 \longrightarrow S^9 \longrightarrow S^6 \longrightarrow S$ is a **minimal free resolution** for $S/M_{\mathcal{A}}$.

- We say that a monomial m' **strictly divides** m if $\text{supp}(\frac{m}{m'}) = \text{supp}(m)$.

- We say that a monomial m' **strictly divides** m if $\text{supp}(\frac{m}{m'}) = \text{supp}(m)$.
- A monomial ideal $M = \langle m_1, \dots, m_r \rangle$ is called **generic** if, whenever two distinct minimal generators m_i and m_j have the same positive degree in some variable x_s , there is a third generator m_l which strictly divides $\text{lcm}(m_i, m_j)$.

Example. $M = \langle x^2y^2, x^2z^2, yz \rangle$ is generic.


- We say that a monomial m' **strictly divides** m if $\text{supp}(\frac{m}{m'}) = \text{supp}(m)$.
- A monomial ideal $M = \langle m_1, \dots, m_r \rangle$ is called **generic** if, whenever two distinct minimal generators m_i and m_j have the same positive degree in some variable x_s , there is a third generator m_l which strictly divides $\text{lcm}(m_i, m_j)$.

Example. $M = \langle x^2y^2, x^2z^2, yz \rangle$ is generic.


- For $I \subseteq \{1, \dots, r\}$, let $m_I = \text{lcm}(m_i, i \in I)$.
- The **Scarf complex** of M consists of the following subsets:

$$\Delta_M := \{I \subseteq \{1, \dots, r\} \mid m_I = m_J \Rightarrow I = J\}.$$

Let $M = \langle a^2b, ac, b^2, bc^2 \rangle$. The Scarf complex of M is




Let $M = \langle a^2b, ac, b^2, bc^2 \rangle$. The Scarf complex of M is



Theorem. For M *generic*, the cellular complex \mathbb{F}_{Δ_M} is a *minimal free resolution* of S/M .

Let $M = \langle a^2b, ac, b^2, bc^2 \rangle$. The Scarf complex of M is



Theorem. For M *generic*, the cellular complex \mathbb{F}_{Δ_M} is a *minimal free resolution* of S/M .

● The minimal free resolution of S/M is

$$0 \rightarrow S^2 \rightarrow S^5 \rightarrow S^4 \rightarrow S$$

- From a monomial ideal M we can get a generic monomial ideal \widetilde{M} by deforming the exponents of the minimal generators of M .
- $\mathbb{F}_{\Delta_{\widetilde{M}}}$ is a resolution of M .

- From a monomial ideal M we can get a generic monomial ideal \widetilde{M} by deforming the exponents of the minimal generators of M .
- $\mathbb{F}_{\Delta_{\widetilde{M}}}$ is a resolution of M .
- Let $S = \mathbf{k}[x_1, \dots, x_5]$, and M be the **matroid ideal**

$$M = \langle x_1x_2, x_1x_3x_4, x_2x_3x_4, x_1x_3x_5, x_2x_3x_5, x_4x_5 \rangle$$

- From a monomial ideal M we can get a generic monomial ideal \widetilde{M} by deforming the exponents of the minimal generators of M .
- $\mathbb{F}_{\Delta_{\widetilde{M}}}$ is a resolution of M .
- Let $S = \mathbf{k}[x_1, \dots, x_5]$, and M be the **matroid ideal**

$$M = \langle x_1x_2, x_1x_3x_4, x_2x_3x_4, x_1x_3x_5, x_2x_3x_5, x_4x_5 \rangle$$

- Fix the order $\pi = 1, 2, 3, 4, 5$ in the indeterminates of S .

- From a monomial ideal M we can get a generic monomial ideal \widetilde{M} by deforming the exponents of the minimal generators of M .
- $\mathbb{F}_{\Delta_{\widetilde{M}}}$ is a resolution of M .
- Let $S = \mathbf{k}[x_1, \dots, x_5]$, and M be the **matroid ideal**

$$M = \langle x_1x_2, x_1x_3x_4, x_2x_3x_4, x_1x_3x_5, x_2x_3x_5, x_4x_5 \rangle$$

- Fix the order $\pi = 1, 2, 3, 4, 5$ in the indeterminates of S .
- $M_i = \langle m^i \mid m \in \text{Gen}(M), m \mid x_1x_2 \cdots x_i, m^i \notin \sum_{j=1}^{i-1} M_j \rangle$

- From a monomial ideal M we can get a generic monomial ideal \widetilde{M} by deforming the exponents of the minimal generators of M .
- $\mathbb{F}_{\Delta_{\widetilde{M}}}$ is a resolution of M .
- Let $S = \mathbf{k}[x_1, \dots, x_5]$, and M be the **matroid ideal**

$$M = \langle x_1x_2, x_1x_3x_4, x_2x_3x_4, x_1x_3x_5, x_2x_3x_5, x_4x_5 \rangle$$

- Fix the order $\pi = 1, 2, 3, 4, 5$ in the indeterminates of S .
- $M_i = \langle m^i \mid m \in \text{Gen}(M), m \mid x_1x_2 \cdots x_i, m^i \notin \sum_{j=1}^{i-1} M_j \rangle$
- $M^\pi = M_1 + M_2 + \cdots + M_n$
- M^π is a **Cohen-Macaulay generic deformation** of M

- From a monomial ideal M we can get a generic monomial ideal \widetilde{M} by deforming the exponents of the minimal generators of M .
- $\mathbb{F}_{\Delta_{\widetilde{M}}}$ is a resolution of M .
- Let $S = \mathbf{k}[x_1, \dots, x_5]$, and M be the **matroid ideal**

$$M = \langle x_1x_2, x_1x_3x_4, x_2x_3x_4, x_1x_3x_5, x_2x_3x_5, x_4x_5 \rangle$$

- Fix the order $\pi = 1, 2, 3, 4, 5$ in the indeterminates of S .
- $M_i = \langle m^i \mid m \in \text{Gen}(M), m \mid x_1x_2 \cdots x_i, m^i \notin \sum_{j=1}^{i-1} M_j \rangle$
- $M^\pi = M_1 + M_2 + \cdots + M_n$
- M^π is a **Cohen-Macaulay generic deformation** of M

$$M^\pi = \langle x_1^2x_2^2, x_1^4x_3^4x_4^4, x_2^4x_3^4x_4^4, x_1^5x_3^5x_5^5, x_2^5x_3^5x_5^5, x_4^5x_5^5 \rangle$$

- From a monomial ideal M we can get a generic monomial ideal \widetilde{M} by deforming the exponents of the minimal generators of M .
- $\mathbb{F}_{\Delta_{\widetilde{M}}}$ is a resolution of M .
- Let $S = \mathbf{k}[x_1, \dots, x_5]$, and M be the **matroid ideal**

$$M = \langle x_1x_2, x_1x_3x_4, x_2x_3x_4, x_1x_3x_5, x_2x_3x_5, x_4x_5 \rangle$$

- Fix the order $\pi = 1, 2, 3, 4, 5$ in the indeterminates of S .
- $M_i = \langle m^i \mid m \in \text{Gen}(M), m \mid x_1x_2 \cdots x_i, m^i \notin \sum_{j=1}^{i-1} M_j \rangle$
- $M^\pi = M_1 + M_2 + \cdots + M_n$
- M^π is a **Cohen-Macaulay generic deformation** of M


$$M^\pi = \langle x_1^2x_2^2, x_1^4x_3^4x_4^4, x_2^4x_3^4x_4^4, x_1^5x_3^5x_5^5, x_2^5x_3^5x_5^5, x_4^5x_5^5 \rangle$$

$$M^\pi = \langle x_1x_2, x_1^2x_3^2x_4^2, x_2^2x_3^2x_4^2, x_1^3x_3^3x_5^3, x_2^3x_3^3x_5^3, x_4^3x_5^3 \rangle$$

$$M^\pi = \langle x_1 x_2, x_1^2 x_3^2 x_4^2, x_2^2 x_3^2 x_4^2, x_1^3 x_3^3 x_5^3, x_2^3 x_3^3 x_5^3, x_4^3 x_5^3 \rangle$$

$$M^\pi = \langle x_1 x_2, x_1^2 x_3^2 x_4^2, x_2^2 x_3^2 x_4^2, x_1^3 x_3^3 x_5^3, x_2^3 x_3^3 x_5^3, x_4^3 x_5^3 \rangle$$

• Δ_{M^π} has facets $\{1, 2, 4\}$, $\{1, 3, 5\}$, $\{1, 2, 6\}$, $\{1, 3, 6\}$



• The cellular complex $\mathbb{F}_{\Delta_{M^\pi}}$ is equal to

$$0 \longrightarrow S^4 \longrightarrow S^9 \longrightarrow S^6 \longrightarrow S$$

- Fix the order $\sigma = 1, 3, 4, 2, 5$. Then

$$M^\sigma = \langle x_1 x_3 x_4, x_1^2 x_2^2, x_2^2 x_3^2 x_4^2, x_1^3 x_3^3 x_5^3, x_2^3 x_3^3 x_5^3, x_4^3 x_5^3 \rangle$$


is the **CM generic deformation** of M associated to the given order.

- Fix the order $\sigma = 1, 3, 4, 2, 5$. Then

$$M^\sigma = \langle x_1 x_3 x_4, x_1^2 x_2^2, x_2^2 x_3^2 x_4^2, x_1^3 x_3^3 x_5^3, x_2^3 x_3^3 x_5^3, x_4^3 x_5^3 \rangle$$

is the **CM generic deformation** of M associated to the given order.

- Δ_{M^σ} has facets $\{1, 2, 4\}$, $\{1, 2, 5\}$, $\{1, 3, 5\}$, $\{1, 2, 6\}$, $\{1, 3, 6\}$




- Fix the order $\sigma = 1, 3, 4, 2, 5$. Then

$$M^\sigma = \langle x_1 x_3 x_4, x_1^2 x_2^2, x_2^2 x_3^2 x_4^2, x_1^3 x_3^3 x_5^3, x_2^3 x_3^3 x_5^3, x_4^3 x_5^3 \rangle$$


is the **CM generic deformation** of M associated to the given order.

- Δ_{M^σ} has facets $\{1, 2, 4\}$, $\{1, 2, 5\}$, $\{1, 3, 5\}$, $\{1, 2, 6\}$, $\{1, 3, 6\}$



- $\mathbb{F}_{\Delta_{M^\sigma}}$ is equal to $0 \longrightarrow S^5 \longrightarrow S^{10} \longrightarrow S^6 \longrightarrow S$


Theorem. *Let $\dim(\Delta_{M^\pi}) = n$, and let $h(\pi)$ be the unique monomial generator of M_1 . Then $\mathbb{F}_{\Delta_{M^\pi}}$ is a **minimal** free resolution if and only if any facet F , with $h(\pi) \in F$, lies in a hyperplane.*




• $\langle x_1, x_2x_3, x_2x_4, x_2x_5, x_3x_4, x_3x_5, x_4x_5 \rangle$

• Minimal free resolution

$$0 \longrightarrow S^3 \longrightarrow S^{11} \longrightarrow S^{14} \longrightarrow S^7 \longrightarrow S$$




● Give $\pi = 1, 2, 3, 5, 4$, the Scarf complex Δ_{M^π} is



● Give $p = 1, 2, 3, 5, 4$, the Scarf complex Δ_{M^p} is


$$\{x_1, x_2x_3, x_2x_5, x_2x_4\},$$



• Give $p = 1, 2, 3, 5, 4$, the Scarf complex Δ_{M^p} is

$$\{x_1, x_2x_3, x_2x_5, x_2x_4\},$$

$$\{x_1, x_2x_3, x_3x_5, x_3x_4\},$$




● Give $p = 1, 2, 3, 5, 4$, the Scarf complex Δ_{M^p} is

$$\{x_1, x_2x_3, x_2x_5, x_2x_4\},$$

$$\{x_1, x_2x_3, x_3x_5, x_3x_4\},$$

$$\{x_1, x_2x_3, x_2x_5, x_4x_5\},$$




● Give $p = 1, 2, 3, 5, 4$, the Scarf complex Δ_{M^p} is

$$\{x_1, x_2x_3, x_2x_5, x_2x_4\},$$

$$\{x_1, x_2x_3, x_3x_5, x_3x_4\},$$

$$\{x_1, x_2x_3, x_2x_5, x_4x_5\},$$

$$\{x_1, x_2x_3, x_3x_5, x_4x_5\}$$



- There does not exist a minimal \mathcal{A} -deformation.
- Generic deformation which is not CM

$$\langle x_1, x_2x_4^3, x_3x_5^3, x_2^2x_3^2, x_4^2x_5^2, x_2^3x_5, x_3^3x_4 \rangle$$

- Scarf complex has dimension 4.