

Chapter 9

AI in Mental Health Services



Yuyi Yang and Dorian Traube

1 Introduction

1.1 *The Digital Revolution in Mental Health*

Rapid advances in digital technology have positioned artificial intelligence (AI) as a catalyst for new approaches to predicting, detecting, and treating mental disorders. Digital mental health is now an established field that continues to expand through rigorous research and a growing portfolio of AI-enabled interventions (D'Alfonso, 2020). Yet, uptake of such technologies in routine psychiatric practice has lagged behind other medical specialties, largely because mental health care depends heavily on interpersonal relationships and qualitative assessments (Graham et al., 2019).

Recent evidence indicates that AI can enhance accessibility, efficiency, and evidence-based decision-making across the mental-health spectrum (Olawade et al., 2024). Machine-learning models improve diagnostic precision, while natural language processing (NLP) tools help clinicians monitor treatment response in real time (Dehbozorgi et al., 2025; Graham et al., 2019). By extracting meaningful patterns from large, heterogeneous data sets, AI supports population-level surveillance, explains biological mechanisms, and informs personalized care plans. Remote delivery platforms and self-assessment applications further extend the reach of evidence-based interventions.

Despite these opportunities, implementation remains constrained by technological, clinical, and ethical challenges. Limitations include sparse real-world training

Y. Yang (✉)

Division of Computational and Data Sciences, Washington University in St. Louis,
St. Louis, MO, USA

e-mail: y.yuyi@wustl.edu

D. Traube

Brown School, Washington University in St. Louis, St. Louis, MO, USA

e-mail: traube@wustl.edu

data, gaps between algorithmic outputs and clinical workflows, and the need to safeguard patient privacy while ensuring transparency and fairness (Koutsouleris et al., 2022; Tavory, 2024). Addressing these issues is critical to realizing AI's potential in mental-health services.

1.2 Aims and Objectives

This chapter critically surveys how contemporary AI—ranging from multimodal deep-learning diagnostics and digital phenotyping to large-language-model chatbots and socially assistive robots (SAR)—is reshaping mental-health care across early detection, diagnosis, treatment personalization, monitoring, crisis response, and population-specific services. It synthesizes current evidence on accuracy, clinical benefit, user engagement, and cost-effectiveness, while interrogating persistent barriers such as data bias, privacy threats, opacity, workforce readiness, and epistemic blind spots. Building on these insights, it delineates research, implementation, and governance priorities needed to integrate next-generation systems, particularly retrieval-augmented large language models (LLMs), into routine practice. Throughout, the analysis underscores that early and sustained co-production with patients, caregivers, clinicians, technologists, and policymakers is indispensable for developing AI that is clinically effective, ethically sound, and socially just.

1.3 Defining AI in Mental Health

AI encompasses computational techniques, such as machine learning, deep learning, and reinforcement learning, that enable computer systems to perform tasks that conventionally require human cognition (Carr, 2020). While widely applied to diagnostic imaging, patient monitoring, and electronic health records (EHR), mental-health contexts demand sensitivity to emotion, culture, and lived experience (Gabbard & Crisp-Han, 2017). Accordingly, AI systems should augment rather than replace therapeutic relationships, providing transparent, interpretable, and ethically grounded decision support.

2 Scope and Core Concepts

2.1 Evolution of AI Applications in Mental Health

The integration of AI in mental healthcare has evolved significantly over recent decades, beginning with simple cognitive models such as the early chatbot ELIZA and progressing to sophisticated AI-driven interventions currently available (Bassett, 2019). Machine-learning-driven tools now underpin screening instruments, relapse-prevention systems, and adaptive digital therapeutics, signaling a shift from trial-and-error approaches to data-informed precision psychiatry (D’Alfonso, 2020).

2.2 Cognitive and Clinical Foundations

By modeling complex interactions among biological, psychological, and social determinants, machine-learning algorithms support more nuanced risk stratification and earlier detection of prodromal states (Koutsouleris et al., 2022). Continued refinement may eventually yield diagnostic taxonomies that supplement—or surpass—conventional nosology such as the DSM-5 (Alhuwaydi, 2024).

2.3 Digital Phenotyping and Ecological Momentary Assessment

Smartphones, wearables, and other sensor-laden devices now enable two complementary approaches to real-world data capture. Digital phenotyping converts passively collected streams, such as GPS traces, accelerometry, and heart-rate variability (HRV), into objective markers of mood, cognition, and social activity (Rogan et al., 2024). Ecological momentary assessment (EMA), by contrast, relies on brief active self-reports (e.g., “How anxious do you feel right now?”) delivered several times a day. When combined, these passive and active signals reveal patterns—such as irregular mobility, fragmented sleep, and fluctuating self-rated affect—that precede depressive or bipolar episodes and can trigger just-in-time (JIT) interventions (de Lacy et al., 2023; Patel et al., 2024).

2.4 NLP in Mental Health

NLP models analyze textual and spoken language to detect linguistic cues associated with psychopathology, including suicidal ideation and affective instability (Flanagan et al., 2021). Large-language-model-based conversational agents (e.g.,

GPT variants) can deliver psychoeducation and preliminary triage, but there are still unresolved concerns about privacy, bias, and clinical validity (Hua et al., 2024; Park & Han, 2018). Rigorous evaluation and stakeholder oversight are essential before widespread deployment.

2.5 Stakeholder-Centered Co-Production

Meaningful AI solutions are co-produced through sustained collaboration between technical developers and those who use or are impacted by the technology. Patients contribute lived experience and behavioral data; clinicians bridge AI insights with clinical practice; data scientists integrate domain knowledge with technical modeling; and policymakers safeguard equity through governance. As depicted in Fig. 9.1, this interdependence forms a dynamic closed-loop system. The Data Loop: Patient-generated data fuel AI model development. The Support Loop: AI-generated insights trigger clinician actions, which in turn yield new behavioral data to refine models. The Governance Loop: Policymakers iteratively regulate the system in response to researcher-identified risks and feedback. Early stakeholder involvement—via participatory design or advisory boards—surfaces critical priorities (e.g., workflow constraints, equity barriers) that shape the AI lifecycle. Empirical evidence confirms that such co-production enhances feasibility, sharpens risk identification, and mitigates algorithmic harm (Broadbent et al., 2024; Patel et al., 2024; Zidaru et al., 2021).




Fig. 9.1 Stakeholder loops governing AI-enabled mental-health services

3 Key Applications and Use Cases

Table 9.1 consolidates AI applications that operate not only across the core stages of the mental-health care continuum—ranging from early detection to crisis response—but also within population-specific domains such as adolescent care, geriatric care, PTSD, and addiction.

Table 9.1 AI applications and representative tools across the mental-health care spectrum

Subsection	Primary AI modalities	Illustrative use cases / tools
Early detection & prediction	Multimodal ML on speech, text, facial affect, EHR; micro-expression & paralinguistic analysis	Cogito voice analytics; Affectiva vision models; EHR suicide-risk DL models; multimodal genomic-wearable models
Clinical diagnosis & decision support	High-dimensional pattern recognition; decision-support dashboards; automated EHR summarization	FDA-cleared autism diagnostic aid; EHR mood-disorder detectors
Treatment & therapy enhancements	Pharmacogenomic classifiers; adaptive digital therapeutics; typing-dynamics wearables; LLM chatbots	Wysa CBT; Mindstrong typing-dynamics monitor; Woebot postpartum-depression chatbot
Monitoring & follow-up	Passive sensing (wearables, smartphones); EMA fusion; anomaly detection vs. personal baseline	Oura ring HRV early-warning; alcohol-use relapse EMA predictor
Chatbots & conversational agents	LLM-based context-aware dialogue platforms	Kai.AI (ACT); Woebot (CBT: Substance use and depression)
Crisis response & suicide prevention	Real-time NLP triage classifiers on crisis text/voice	Crisis text line
Adolescent mental health	Social-media sentiment; sleep & activity tracking; school digital screeners	Youth social-media mood map; school referral screener (co-designed)
Elderly mental health care	Socially assistive robots; smart-home sensors + NLP companions	Elliq robot adherence coach; internet-of-things framework
PTSD & trauma treatment	Physio-adaptive VR exposure therapy; biosensor feedback; NLP analysis of therapy transcripts	VR exposure auto-titration; biosensor anxiety classifier; NLP trauma-cue tracker
Addiction therapy	JIT adaptive interventions; geospatial pattern recognition in recovery apps	Addict free context-aware prompts

3.1 Early Detection and Prediction

The early identification of mental health disorders increasingly depends on AI-driven analysis of speech, text, facial affect, and longitudinal EHRs. For example, micro-expression recognition algorithms now detect prodromal symptoms that frequently escape clinician observation (Zhao et al., 2023). Commercial platforms such as Cogito employ paralinguistic analysis during tele-psychiatry visits to flag subtle vocal changes, whereas Affectiva's vision models screen for depressive affective states (Affectiva, 2025; Cogito, 2025). By integrating heterogeneous data streams, machine-learning pipelines generate personalized risk scores that outperform traditional single-modality screeners.

Population-scale studies likewise demonstrate the value of multimodal modeling. Hilbert et al. (2017) distinguished generalized anxiety disorder from major depressive disorder by fusing endocrine markers with structural MRI data. Deep-learning models trained on EHR predict suicide attempts up to 1 year in advance, facilitating proactive outreach (Walsh et al., 2021; Zheng et al., 2020). Most recently, de Lacy et al. (2023) achieved area-under-the-curve (AUC—a 0-to-1 measure of overall diagnostic accuracy) values above 0.90 by jointly modeling clinical, genomic, and wearable-sensor data for three common disorders. Although external validation remains limited, these findings underscore the feasibility of moving from reactive to preventive psychiatry (Ehtemam et al., 2024; Olawade et al., 2024).

3.2 Clinical Diagnosis and Decision Support

In diagnostic practice, pattern-recognition capabilities enable AI systems to process the high-dimensional data generated by routine care. Decision-support dashboards display differential diagnoses, estimate prognoses, and recommend evidence-based treatment pathways (Dehbozorgi et al., 2025). When embedded into shared decision-making workflows, such systems have shortened assessment time, reduced waiting lists, and improved recovery rates (Rollwage et al., 2023). Importantly, clinicians report that automated summarization of patient-reported outcomes frees them to focus on the therapeutic relationship rather than data entry (De Angel et al., 2023).

Regulatory agencies are beginning to recognize clinical AI. In 2021, the U.S. Food and Drug Administration (FDA) granted breakthrough designation to an autism diagnostic aid that combines caregiver questionnaires with clinician inputs and achieves 98% sensitivity (Broadbent et al., 2024). Comparable gains have been documented for automated detection of depressive and anxiety disorders in EHRs (Higgins & Wilson, 2025). To sustain clinician trust, however, model transparency and interpretability remain essential (Warrier et al., 2023).

3.3 Treatment and Therapy Enhancements

Precision therapeutics leverage machine-learning models to match interventions with individual biological and psychosocial profiles. Pharmacogenomic classifiers, for instance, forecast antidepressant response by analyzing single-nucleotide variants (Bobo et al., 2022). Digital companions such as Wysa deliver evidence-informed cognitive-behavioral content and mood-tracking functions (Wysa, 2025), while adaptive internet-delivered cognitive behavioral therapy (CBT) adjusts session sequencing in real time (Thieme et al., 2023). Wearable-enabled platforms like Mindstrong infer cognitive fluctuations from typing dynamics, enabling just-in-time clinical review (Gruber, 2025). The FDA's breakthrough designation for the Woebot postpartum depression chatbot illustrates regulators' growing openness to digital therapeutics (Patel et al., 2024).

3.4 Monitoring and Follow-Up

Continuous passive sensing—via smartphones, wearables, and ambient sensors—provides high-frequency behavioral and physiological data. AI models detect departures from personal baselines and trigger alerts well before clinical deterioration becomes obvious (Khoo et al., 2024; Rollwage et al., 2023). For example, Oura Ring analytics link reduced HRV and fragmented sleep to impending depressive episodes (Kristiansson et al., 2023). Ecological momentary-assessment apps that fuse mood diaries with sensor streams can predict relapse in alcohol-use disorder with high precision (Alowais et al., 2023). By continuously monitoring patient data and automatically prioritizing follow-up, AI frees clinicians to devote their time to complex therapeutic work.

3.5 Chatbots and Conversational Agents

Conversational agents have evolved from rule-based scripts to LLM-based systems capable of context-aware dialogue (Thapa & Adhikari, 2024). Randomized trials report moderate reductions in depressive and anxiety symptoms after 4–8 weeks of guided interaction (Loveys et al., 2023). Special-purpose chatbots expand access to mental health support for underserved groups. For example, Kai.ai, an acceptance and commitment therapy (ACT)-based texting companion, improved well-being in more than 10,000 adolescents awaiting psychiatric intake (Vertsberger et al., 2022). Moreover, a study of a CBT-based postpartum chatbot showed modest reductions in depressive symptoms among new mothers, suggesting such tools can widen access to perinatal mental-health support (Suharwardy et al., 2023). Preliminary evidence indicates that users perceive these sophisticated interactions as beneficial for

managing stress and loneliness (Yang et al., 2025). While promising, large-scale effectiveness studies and robust safeguarding protocols are still required.

3.6 Crisis Response and Suicide Prevention

Natural language classifiers embedded in crisis-text platforms can triage users within seconds, prioritizing those at imminent risk of self-harm. For example, Crisis Text Line's NLP-driven triage model correctly flags 86% of texters at severe imminent risk on their very first message (Filbin, 2018) and has cut the median wait for a counselor response from 8 min under human-only triage to just 3 min—almost a 50% reduction (Swaminathan et al., 2023). Reliance on language alone, however, poses ethical and technical dilemmas. Sarcasm, cultural idioms, and code-switching can confuse models and lead to both missed crises and false alarms; recent benchmark studies show markedly lower accuracy on African American English and other underrepresented dialects (Mire et al., 2025). False positives may trigger unwarranted police wellness checks that traumatize callers or exhaust scarce emergency resources, whereas false negatives leave texters waiting in silence.

Privacy risks further complicate deployment. In 2022, Crisis Text Line faced public backlash for sharing anonymized conversation data with its for-profit spin-off, prompting the hotline to terminate the arrangement and request deletion of all previously transferred data (Crisis Text Line, 2022). The episode underscores why crisis platforms must adopt transparent governance, opt-in data policies, human-in-the-loop overrides, and regular bias audits to complement the speed and scale of algorithms.

3.7 Adolescent Mental Health

Recent work shows that multimodal, developmentally tuned AI can flag emerging mood problems in teens days, sometimes weeks, before they meet clinical cut-offs. In a U.S. cohort of 9- to 17-year-olds, de Lacy et al. (2023) trained meta-learning ensembles on 160 psychosocial, physiological, and environmental features and achieved held-out AUCs ≥ 0.94 for depression, anxiety, attention-deficit/hyperactivity disorder (ADHD), disruptive behavior, and post-traumatic stress disorder (PTSD), with everyday behavioral signals outperforming neuro-imaging inputs. When these passive streams (social-media sentiment, sleep–wake regularity, and activity rhythms) are layered into real-time dashboards, clinicians can intervene earlier and more precisely.

In school settings, digital screeners have moved beyond static questionnaires: Olawade et al. (2024) report that an adaptive triage engine increased referral precision without adding false positives, ensuring counselor time goes to students who need it most. Sustained engagement, however, hinges on participatory governance.

A mixed-methods study of AI-enabled mental-health apps for young people found that adolescents will share sensitive data only when they help set privacy rules, see clear value, and can veto unwanted personalization (Nicol et al., 2022); similar themes emerged in a German co-design study that highlighted transparency and user control as non-negotiables (Götzl et al., 2022). Best practice now pairs youth-led design reviews with parent–teacher steering groups so that alert thresholds, escalation protocols, and tone remain both developmentally appropriate and culturally sensitive.

3.8 Elderly Mental Health Care

SARs and voice-enabled virtual agents are moving from pilot studies to real-world elder-care programs. A systematic review of eight trials by Boumans et al. (2022) found that socially assistive systems most often deliver timed medication reminders, brief cognitive games, and conversational check-ins. Home deployments achieved the highest usability scores, yet hearing-related speech-recognition errors remained a persistent barrier. Building on this evidence, Broadbent et al. (2024) describe the roll-out of the tabletop robot ElliQ across 15 U.S. and Canadian ageing-service agencies. Early field data show sustained daily engagement and self-reported reductions in loneliness, while weekly summary reports allow clinicians to integrate robot-captured insights into routine care.

Ambient sensing provides a complementary safety net. Feng et al. (2017) describe a cognitive Internet-of-Things framework that integrates heterogeneous smart-home sensors, such as motion, acoustic, light, temperature, and even furniture-pressure, into a unified data stream. An adaptive control layer then analyses these real-time traces, flags behavioral deviations, and sends plain-language alerts to caregivers. When paired with the dialogue capabilities highlighted by Boumans et al., such systems help older adults remain independent while keeping families informed. Crucially, all three studies stress that age-inclusive usability testing—covering interface legibility, error recovery, and user-tunable alert thresholds—is essential to prevent technology-driven exclusion as these AI tools scale.

3.9 PTSD and Trauma Treatment

Machine-learning tools now support trauma care across physiological and textual modalities. A systematic review of clinical studies by Ehtemam et al. (2024) found that suicide-risk prediction algorithms—many trained on PTSD-enriched cohorts—achieve moderate-to-high discrimination, laying the groundwork for automated safety-planning flags. Within virtual-reality exposure therapy, Mevlevioğlu et al. (2024) showed that biosensor streams captured inside the headset (e.g., electrodermal activity, HRV) classify moment-to-moment anxiety with 75% accuracy,

enabling physiologically adaptive stimulus dosing. Complementing these physiological signals, NLP pipelines mine free text—from adolescent earthquake narratives (Yuan et al., 2024) to electronic medical records (Miranda et al., 2024)—to surface trauma cues and track symptom trajectories, provided that stringent confidentiality safeguards remain in place throughout the data lifecycle.

3.10 *Addiction Therapy*

JIT adaptive interventions that fuse smartphone sensing with reinforcement-learning policies can issue coping prompts exactly when relapse risk peaks. Gönül et al. (2021) showed that a personalized scheduling algorithm consistently outperformed fixed-rule baselines in simulated craving scenarios, demonstrating the value of continuous risk forecasting. Geospatial context adds another layer: Yang et al. (2019) use GPS and temporal patterns in the Addict Free app to detect high-risk “smoking zones” and nudge users toward alternative activities, while a deep-learning early-warning engine for suicide attempts built on EHRs (Zheng et al., 2020) illustrates how similar predictive frameworks transfer to comorbid self-harm risks.

Automated chatbots employing motivational-interviewing scripts can maintain therapeutic continuity outside clinic hours, but experience so far indicates they work best when tightly integrated with peer and clinician support. Long-term recovery, therefore, hinges on combining machine-generated prompts with human accountability, regular professional check-ins, and access to community resources that reinforce behavior change.

4 Limitations, Challenges, and Ethical Considerations

4.1 *Technical and Implementation Barriers*

Developing robust AI tools for mental health is constrained foremost by data quality and scope. Small, homogeneous samples promote over-fitting and undermine external validity (Schnack & Kahn, 2016), while missing or inconsistent records can distort predictions and jeopardize clinical recommendations (Tsvetanova et al., 2021). Even well-performing models often falter once deployed beyond the datasets on which they were trained (Graham et al., 2019). Under-representation of racial and ethnic minorities, older adults, and socio-economically diverse populations compounds these issues by embedding structural bias in algorithmic outputs (Franklin et al., 2024).

Despite rapid methodological advances, precision psychiatry remains in its infancy. Many machine-learning studies are conceived without the early involvement of clinicians, patients, or health-system leaders, leading to tools that are

technically impressive yet clinically awkward (Wiens et al., 2019). Current prototypes rarely align with standard psychiatric assessment frameworks and cannot reproduce the empathetic exchanges fundamental to therapeutic practice (Ray et al., 2022). When model outputs are derived from questionable sensor data or self-reports, clinicians' skepticism is justified: over-reliance can delay intervention or overlook clinical deterioration (Malik et al., 2025; Rogan et al., 2024). Ensuring transparent reporting, continuous monitoring, and human oversight is therefore imperative.

4.2 Ethical and Regulatory Considerations

4.2.1 Algorithmic Bias and Data Protection

Training data mirror the inequities of the societies that generate them, and algorithms can magnify those inequities if safeguards are absent (Chen et al., 2019). Discriminatory outputs have been observed when diagnostic labels correlate spuriously with ethnicity or income. Regulatory regimes are beginning to respond: the EU Artificial Intelligence Act, for example, mandates bias audits and minimum performance thresholds for high-risk medical systems (Higgins & Wilson, 2025). Given the sensitivity of mental-health information, robust encryption, differential-privacy techniques, and explicit consent protocols are non-negotiable (Cruz-Gonzalez et al., 2025).

4.2.2 Explainability and Clinical Trust

Deep learning architectures often operate as opaque “black boxes (their internal reasoning is hidden from users),” limiting clinicians' willingness to rely on them in high-stakes decisions (Olawade et al., 2024). Explainable-AI (XAI) methods—such as attention maps, counterfactual explanations, and Shapley values—can expose the logic of a model's prediction, enabling users to detect bias or error (Albahri et al., 2023). Future research must prioritize solutions that balance predictive accuracy with interpretability to sustain accountability and promote adoption.

4.2.3 Global Governance Gaps

While the FDA's Software-as-a-Medical-Device framework and the EU AI Act provide a starting point, neither addresses the nuances of mental-health contexts—ranging from crisis-hotline triage to longitudinal mood monitoring (Broadbent et al., 2024; Warrier et al., 2023). The World Health Organization (WHO) has urged member states to establish sector-specific standards that encompass privacy, safety, and cultural appropriateness. Harmonized international guidelines would help

ensure that vulnerable populations benefit from AI without incurring disproportionate risks.

4.3 Clinical and Human-Centered Concerns

Therapeutic effectiveness in mental health depends on empathy, rapport, and subtle non-verbal communication—capacities that current AI systems only approximate (Gabbard & Crisp-Han, 2017). When assessment or treatment planning is delegated wholesale to automated decision aids, care can become depersonalized, and the clinician–patient alliance weakened. Misalignment between algorithmic recommendations and professional judgment or lived experience may, in turn, erode trust (Koutsouleris et al., 2022; Rashid & Sharma, 2025). A growing body of user-centered design research demonstrates that engaging service users, carers, and frontline clinicians throughout development mitigates these risks. Participatory workshops, iterative prototyping, and in-situ simulations surface context-specific safety concerns, improve perceived helpfulness, and reduce attrition in digital interventions (Bernaerts et al., 2024). Such co-production has proved especially important for tailoring chatbot tone, escalation thresholds, and interface accessibility for older adults, adolescents, and linguistically minoritized populations (Peng et al., 2024).

As conversational agents adopt naturalistic voices and avatars, users increasingly attribute human-like understanding and intentions to them—a variant of the “ELIZA effect.” Experimental comparisons of anthropomorphic “digital humans” with text-only chatbots show higher perceived empathy but also inflated expectations of clinical competence, which can lead to unsafe reliance during crises (Peter et al., 2025). Designers therefore face a trade-off: richer social cues may enhance engagement, yet excessive anthropomorphism can blur the boundary between tool and therapist. LLMs further complicate the picture. Recent evaluations reveal that state-of-the-art systems still generate inaccurate or unsafe advice in a minority of self-harm scenarios despite guardrails (Kolding et al., 2024). Embedding explicit usage boundaries, real-time risk-detection triggers, and human-in-the-loop escalation pathways is therefore essential. Transparent disclosure of an agent’s non-human status, combined with continuous bias audits and user-centered design principles, can help preserve relational quality while minimizing the dangers of both depersonalized care and misplaced anthropomorphic trust.

4.4 Practical and Workforce Implications

Successful implementation depends on clinician literacy in data science and on thoughtfully re-engineered workflows. Targeted training should equip mental-health professionals to interpret model outputs while maintaining patient-centered values

(Higgins & Wilson, 2025). Participatory design, iterative prototyping, and simulation exercises can surface bottlenecks before deployment (Scheinker & Brandeau, 2020). Where appropriately integrated, AI may automate administrative tasks, streamline triage, and expand capacity for high-need patients (Warrier et al., 2023).

Attention to health equity is essential. People with intellectual or developmental disabilities, for example, are frequently excluded from digital-health research, thereby widening existing disparities (Zidaru et al., 2021). Inclusive data strategies and culturally attuned interface design can help ensure that AI reinforces rather than undermines equitable access to mental-health services. Emerging work on intelligent resource matching—such as directing users to linguistically appropriate crisis lines—illustrates how algorithmic tools can advance, rather than hinder, justice in mental-health care (Ehtemam et al., 2024).

4.5 Epistemological Reflexivity

Despite the growing enthusiasm for data-driven precision psychiatry, scholars in critical psychiatry and disability studies traditions caution that an exclusive focus on quantifiable markers of distress risks marginalizing lived experiences that resist easy instrumentation. Qualitative narratives of trauma, cultural idioms of distress, and personal recovery stories can be flattened when reduced to sensor streams or symptom counts. Embracing epistemological pluralism—where algorithmic outputs are complemented by participatory, phenomenological, and narrative methods—can mitigate this blind spot and foster more inclusive mental health AI (Kirmayer & Gómez-Carrillo, 2019; Shakespeare, 2013).

5 Future Directions

5.1 Research Agendas and Mechanistic Understanding

Prospective investigations should elucidate the mechanisms through which AI-enabled interventions improve mental-health outcomes. While conventional predictive models may be adequate for discrete, point-of-care tasks, the multifactorial nature of mental disorders demands interpretable approaches that facilitate accountability and clinician trust (Koutsouleris et al., 2022). Accordingly, forthcoming work ought to privilege methods that achieve both analytic precision and explanatory clarity. Leveraging large-scale multimodal datasets in this manner will allow researchers to surface clinically actionable patterns, refine treatment pathways, and identify opportunities for earlier intervention.

Despite increasing experimentation, the science of AI-supported mental health-care remains emergent. Few studies track long-term implementation or cross-site

generalizability (Auf et al., 2025). Rigorous longitudinal and multi-center designs—supplemented by rich case studies—are required to generate evidence that informs sustainable deployment for diverse populations and settings.

5.2 Implementation Science and Stakeholder Involvement

Patient and public involvement are increasingly recognized as prerequisites for responsible AI innovation. Treating service users and caregivers as domain experts throughout the design and evaluation cycle promotes equity, fosters trust, and aligns tools with lived experience (Zidaru et al., 2021). Empirical work demonstrates that early stakeholder engagement—via prototype testing, advisory boards, or citizen-science methods—enhances feasibility and acceptability (Broadbent et al., 2024; Patel et al., 2024).

Standardized research methodologies and reporting guidelines remain urgently needed to facilitate evidence synthesis and cross-study comparability (Dehbozorgi et al., 2025). Interdisciplinary partnerships—linking computational scientists, clinicians, implementation researchers, and ethicists—can help ensure methodological rigor and smooth translation from algorithm to workflow.

5.3 Integration of Large Language Models

LLMs have rapidly expanded the frontier of conversational support tools in psychiatry. By modeling nuanced linguistic cues, contemporary systems can simulate empathic dialogue, deliver psychoeducation, and triage distress (Kolding et al., 2024). Promising lines of work, such as multi-agent reinforcement learning (Liu et al., 2024), further suggest avenues for collaborative, context-aware interventions.

These advances, however, are counter-balanced by substantive concerns around privacy, factual reliability, and clinical safety. LLM outputs may be linguistically persuasive yet substantively inaccurate. Retrieval-augmented generation (RAG), which first fetches supporting documents from a knowledge base and then conditions the language model on them, can mitigate hallucinations (Li et al., 2025), while calibration techniques seek to temper unwarranted confidence (Huang et al., 2025; Leng et al., 2024). Nevertheless, robust validation in real-world mental-health contexts is still limited.

Future evaluations must articulate transparent development pipelines, benchmark performance in routine clinical environments, and incorporate iterative feedback from service users and clinicians. Special attention is warranted in high-risk scenarios—including suicide-prevention chatbots—where escalation protocols and external oversight are indispensable (Heston, 2023; Kiseleva et al., 2022).

5.4 Beyond the Clinic—Education and Capacity Building

Effective translation of algorithms into routine care is hindered by gaps between algorithmic performance and clinical utility, limited access to representative datasets, and insufficient integration into existing workflows (Koutsouleris et al., 2022). Addressing these issues requires blended human–machine systems co-designed by clinicians and data scientists. Curriculum reform and continuous professional development should equip mental-health professionals to interpret AI outputs, while sensitizing technologists to ethical, cultural, and relational dimensions of care. Shared infrastructure, common standards, and open benchmarking will accelerate the shift from proof-of-concept to sustainable practice.

5.5 Governance Futures for Ethical AI

Looking ahead, regulatory landscapes are crystallizing yet remain uneven. In the United States, the FDA already treats most mental health software as a Software-as-a-Medical-Device. Across the Atlantic, the European Artificial Intelligence Act—Regulation (EU) 2024/1689, in force since 1 August 2024, formally classifies clinical-decision support, digital phenotyping, and conversational agents used in mental-health care as high-risk. By 2026, manufacturers seeking access to the European market must complete conformity assessments, implement post-market monitoring plans, and provide documented human-oversight safeguards (European Union, 2024). Future governance will depend on translating such statutory duties into day-to-day practice. The process-oriented ISO/IEC 42001:2023 management system standard provides a ready template for continuous risk assessment and quality improvement. Pilot programs such as NHS England’s AI & Digital Regulations Service—which maps ISO 42001 clauses onto national clinical-safety standards—demonstrate how health systems can operationalize global norms while retaining local accountability (NHS England, 2025).

At the multilateral level, the International Telecommunication Union (ITU) and WHO Focus Group on AI for Health have launched an open benchmarking pipeline that supplies reference datasets and performance thresholds; extending this model to mental-health use-cases would create a common evidentiary baseline for regulators, payers, and developers (ITU & WHO, 2023). Complementing these technical mechanisms, the WHO’s 2021 guidance on the ethics and governance of AI for health articulates cross-cutting principles—safety, inclusiveness, explainability, and sustainability—that can anchor future national strategies (WHO, 2021).

Taken together, these layered instruments—binding regulation, voluntary international standards, sector-specific benchmarking, and ethics frameworks—sketch a path toward convergent governance, in which mental-health AI systems are accompanied by interoperable “safety cases” that follow each product across jurisdictions, are revised through post-market surveillance, and are audited against globally

shared metrics. Realizing such a system will depend on continued investment in regulatory science, the expansion of shared test beds, and participatory oversight mechanisms capable of keeping pace with rapid technical change.

6 Conclusion

6.1 Key Takeaways

Across detection, diagnosis, treatment, personalization, and population surveillance, AI-enabled tools are reshaping mental-health services. Mobile applications, conversational agents, and immersive platforms demonstrate the capacity to analyze multimodal data streams and deliver timely, targeted interventions. Use cases spanning adolescent care, ageing populations, PTSD, and substance-use disorders illustrate both breadth and depth of application. Yet persistent challenges—algorithmic bias, limited generalizability, data inequities, and tension with relational models of care—underscore the need for cautious, evidence-driven adoption.

6.2 Final Remarks and Call to Action

Harnessing AI for mental-health benefit will demand sustained interdisciplinary collaboration, rigorous evaluation, and meaningful stakeholder partnership. Priorities include:

- *Interpretability and Safety*: Develop transparent models, calibrate uncertainty, and embed escalation pathways for crisis contexts.
- *Inclusive Governance*: Implement regulatory and professional frameworks that balance innovation with patient protection and data justice.
- *Capacity Building*: Provide clinicians, researchers, and technologists with training that bridges technical and human-centered competencies.
- *Longitudinal Evidence*: Conduct multi-site, real-world studies that track clinical effectiveness, implementation barriers, and equity impacts over time.

By foregrounding equity, trust, and therapeutic alliance, AI can augment—not supplant—human expertise, contributing to more responsive, accessible, and just mental-health systems.

Competing Interests The authors have no conflicts of interest to declare that are relevant to the content of this chapter.

References

- Affectiva. (2025). *Humanizing technology to bridge the gap between humans and machines*. Affectiva. <https://www.affectiva.com/>
- Albahri, A. S., Duhaime, A. M., Fadhel, M. A., Alnoor, A., Baqer, N. S., Alzubaidi, L., et al. (2023). A systematic review of trustworthy and explainable artificial intelligence in healthcare: Assessment of quality, bias risk, and data fusion. *Information Fusion*, 96, 156–191. <https://doi.org/10.1016/j.inffus.2023.03.008>
- Alhuwaydi, A. M. (2024). Exploring the role of artificial intelligence in mental healthcare: Current trends and future directions—A narrative review for a comprehensive insight. *Risk Management and Healthcare Policy*, 17, 1339–1348. <https://doi.org/10.2147/RMHP.S461562>
- Alowais, S. A., Alghamdi, S. S., Alsuhbany, N., Alqahtani, T., Alshaya, A. I., Almohareb, S. N., et al. (2023). Revolutionizing healthcare: The role of artificial intelligence in clinical practice. *BMC Medical Education*, 23(1), 689. <https://doi.org/10.1186/s12909-023-04698-z>
- Auf, H., Svedberg, P., Nygren, J., Nair, M., & Lundgren, L. E. (2025). The use of AI in mental health services to support decision-making: Scoping review. *Journal of Medical Internet Research*, 27, e63548. <https://doi.org/10.2196/63548>
- Bassett, C. (2019). The computational therapeutic: Exploring Weizenbaum's ELIZA as a history of the present. *AI & Society*, 34(4), 803–812. <https://doi.org/10.1007/s00146-018-0825-9>
- Bernaerts, S., Van Daele, T., Carlsen, C. K., Nielsen, S. L., Schaap, J., & Roke, Y. (2024). User involvement in digital mental health: Approaches, potential and the need for guidelines. *Frontiers in Digital Health*, 6, 1440660. <https://doi.org/10.3389/fgdth.2024.1440660>
- Bobo, W. V., Van Ommeren, B., & Athreya, A. P. (2022). Machine learning, pharmacogenomics, and clinical psychiatry: Predicting antidepressant response in patients with major depressive disorder. *Expert Review of Clinical Pharmacology*, 15(8), 927–944. <https://doi.org/10.1080/017512433.2022.2112949>
- Boumans, R., van de Sande, Y., Thill, S., & Bosse, T. (2022). Voice-enabled intelligent virtual agents for people with amnesia: Systematic review. *JMIR Aging*, 5(2), e32473. <https://doi.org/10.2196/32473>
- Broadbent, E., Loveys, K., Ilan, G., Chen, G., Chilukuri, M. M., Boardman, S. G., et al. (2024). ElliQ, an ai-driven social robot to alleviate loneliness: Progress and lessons learned. *The Journal of Aging Research & Lifestyle*, 13, 22–28. <https://doi.org/10.14283/jarlife.2024.2>
- Carr, S. (2020). 'AI gone mental': Engagement and ethics in data-driven technology for mental health. *Journal of Mental Health*, 29(2), 125–130. <https://doi.org/10.1080/09638237.2020.1714011>
- Chen, I. Y., Szolovits, P., & Ghassemi, M. (2019). Can AI help reduce disparities in general medical and mental health care? *AMA Journal of Ethics*, 21(2), 167–179. <https://doi.org/10.1001/amajethics.2019.167>
- Cogito. (2025). *Training data solutions for AI and machine learning*. Cogitotech. <https://www.cogitotech.com/>
- Crisis Text Line. (2022, January 31). *An update on data privacy, our community and our service*. Crisis Text Line. <https://www.crisistextline.org/blog/2022/01/31/an-update-on-data-privacy-our-community-and-our-service/>
- Cruz-Gonzalez, P., He, A. W. J., Lam, E. P., Ng, I. M. C., Li, M. W., Hou, R., et al. (2025). Artificial intelligence in mental health care: A systematic review of diagnosis, monitoring, and intervention applications. *Psychological Medicine*, 55, e18. <https://doi.org/10.1017/S0033291724003295>
- D'Alfonso, S. (2020). AI in mental health. *Current Opinion in Psychology*, 36, 112–117. <https://doi.org/10.1016/j.copsyc.2020.04.005>
- De Angel, V., Adeleye, F., Zhang, Y., Cummins, N., Munir, S., Lewis, S., et al. (2023). The feasibility of implementing remote measurement technologies in psychological treatment for depression: Mixed methods study on engagement. *JMIR Mental Health*, 10, e42866. <https://doi.org/10.2196/42866>

- de Lacy, N., Ramshaw, M. J., McCauley, E., Kerr, K. F., Kaufman, J., & Nathan Kutz, J. (2023). Predicting individual cases of major adolescent psychiatric conditions with artificial intelligence. *Translational Psychiatry*, 13(1), 314. <https://doi.org/10.1038/s41398-023-02599-9>
- Dehbozorgi, R., Zangeneh, S., Khooshab, E., Nia, D. H., Hanif, H. R., Samian, P., et al. (2025). The application of artificial intelligence in the field of mental health: A systematic review. *BMC Psychiatry*, 25, 132. <https://doi.org/10.1186/s12888-025-06483-2>
- Ehtemam, H., Sadeghi Esfahlani, S., Sanaei, A., Ghaemi, M. M., Hajesmaeel-Gohari, S., Rahimisadegh, R., et al. (2024). Role of machine learning algorithms in suicide risk prediction: A systematic review-meta analysis of clinical studies. *BMC Medical Informatics and Decision Making*, 24(1), 138. <https://doi.org/10.1186/s12911-024-02524-0>
- European Union. (2024). Regulation (EU) 2024/1689 of the European Parliament and of the Council of 13 June 2024 laying down harmonised rules on artificial intelligence (Artificial Intelligence Act). *Official Journal of the European Union*, L 2024/1689, 12 July 2024. <http://data.europa.eu/eli/reg/2024/1689/oj>
- Feng, S., Setoodeh, P., & Haykin, S. (2017). Smart home: Cognitive interactive people-centric internet of things. *IEEE Communications Magazine*, 55(2), 34–39. <https://doi.org/10.1109/MCOM.2017.1600682CM>
- Filbin, B. (2018, March 28). Detecting Crisis: An AI Solution. *Crisis Text Line Blog*. <https://www.crisistextline.org/blog/2018/03/28/detecting-crisis-an-ai-solution/>
- Flanagan, O., Chan, A., Roop, P., & Sundram, F. (2021). Using acoustic speech patterns from smartphones to investigate mood disorders: Scoping review. *JMIR mHealth and uHealth*, 9(9), e24352. <https://doi.org/10.2196/24352>
- Franklin, G., Stephens, R., Piracha, M., Tiosano, S., Lehouillier, F., Koppel, R., & Elkin, P. L. (2024). The sociodemographic biases in machine learning algorithms: A biomedical informatics perspective. *Life*, 14(6), 652. <https://doi.org/10.3390/life14060652>
- Gabbard, G. O., & Crisp-Han, H. (2017). The early career psychiatrist and the psychotherapeutic identity. *Academic Psychiatry*, 41, 30–34. <https://doi.org/10.1007/s40596-016-0627-7>
- Gönül, S., Namlı, T., Coşar, A., & Toroslu, İ. H. (2021). A reinforcement learning based algorithm for personalization of digital, just-in-time, adaptive interventions. *Artificial Intelligence in Medicine*, 115, 102062. <https://doi.org/10.1016/j.artmed.2021.102062>
- Götzl, C., Hiller, S., Rauschenberg, C., Schick, A., Fechtelpeter, J., Fischer Abaigar, U., et al. (2022). Artificial intelligence-informed mobile mental health apps for young people: A mixed-methods approach on users' and stakeholders' perspectives. *Child and Adolescent Psychiatry and Mental Health*, 16(1), 86. <https://doi.org/10.1186/s13034-022-00522-6>
- Graham, S., Depp, C., Lee, E. E., Nebeker, C., Tu, X., Kim, H. C., & Jeste, D. V. (2019). Artificial intelligence for mental health and mental illnesses: An overview. *Current Psychiatry Reports*, 21, 1–18. <https://doi.org/10.1007/s11920-019-1094-0>
- Gruber, T. (2025). Patient, know thyself: AI-powered digital brain biomarkers. <https://tomgruber.org/mindstrong-story/>
- Heston, T. F. (2023). Safety of large language models in addressing depression. *Cureus*, 15(12). <https://doi.org/10.7759/cureus.50729>
- Higgins, O., & Wilson, R. L. (2025). Integrating artificial intelligence (AI) with workforce solutions for sustainable care: A follow up to artificial intelligence and machine learning (ML) based decision support Systems in Mental Health. *International Journal of Mental Health Nursing*, 34(2), e70019. <https://doi.org/10.1111/inm.70019>
- Hilbert, K., Lueken, U., Muehlhan, M., & Beesdo-Baum, K. (2017). Separating generalized anxiety disorder from major depression using clinical, hormonal, and structural MRI data: A multimodal machine learning study. *Brain and Behavior*, 7(3), e00633. <https://doi.org/10.1002/brb3.633>
- Hua, Y., Na, H., Li, Z., Liu, F., Fang, X., Clifton, D., & Torous, J. (2024). Applying and evaluating large language models in mental health care: A scoping review of human-assessed generative tasks. *arXiv*. <https://doi.org/10.48550/arXiv.2408.11288>
- Huang, C., Huang, L., Leng, J., Liu, J., & Huang, J. (2025). Efficient test-time scaling via self-calibration. *arXiv*. <https://doi.org/10.48550/arXiv.2503.00031>

- International Telecommunication Union & World Health Organization. (2023). *Focus Group on Artificial Intelligence for Health (FG-AI4H)*. <https://www.itu.int/go/fgai4h>
- Khoo, L. S., Lim, M. K., Chong, C. Y., & McNaney, R. (2024). Machine learning for multimodal mental health detection: A systematic review of passive sensing approaches. *Sensors*, 24(2), 348. <https://doi.org/10.3390/s24020348>
- Kirmayer, L. J., & Gómez-Carrillo, A. (2019). Agency, embodiment and enactment in psychosomatic theory and practice. *Medical Humanities*, 45(2) <https://mh.bmj.com/content/45/2/169>
- Kiseleva, A., Kotzinos, D., & De Hert, P. (2022). Transparency of AI in healthcare as a multilayered system of accountabilities: Between legal requirements and technical limitations. *Frontiers in Artificial Intelligence*, 5, 879603. <https://doi.org/10.3389/frai.2022.879603>
- Kolding, S., Lundin, R. M., Hansen, L., & Østergaard, S. D. (2024). Use of generative artificial intelligence (AI) in psychiatry and mental health care: A systematic review. *Acta Neuropsychiatrica*, 1–14. <https://doi.org/10.1017/neu.2024.50>
- Koutsouleris, N., Hauser, T. U., Skvortsova, V., & De Choudhury, M. (2022). From promise to practice: Towards the realisation of AI-informed mental health care. *The Lancet Digital Health*, 4(11), e829–e840. [https://doi.org/10.1016/S2589-7500\(22\)00153-4](https://doi.org/10.1016/S2589-7500(22)00153-4)
- Kristiansson, E., Fridolfsson, J., Arvidsson, D., Holmäng, A., Börjesson, M., & Andersson-Hall, U. (2023). Validation of Oura ring energy expenditure and steps in laboratory and free-living. *BMC Medical Research Methodology*, 23(1), 50. <https://doi.org/10.1186/s12874-023-01868-x>
- Leng, J., Huang, C., Zhu, B., & Huang, J. (2024). Taming overconfidence in LLMs: Reward calibration in RLHF. *arXiv*. <https://doi.org/10.48550/arXiv.2410.09724>
- Li, H., Huang, J., Ji, M., Yang, Y., & An, R. (2025). Use of retrieval-augmented large language model for COVID-19 fact-checking: Development and usability study. *Journal of Medical Internet Research*, 27, e66098. <https://doi.org/10.2196/66098>
- Liu, Q., Ye, J., Ma, X., Yang, J., Liang, B., & Zhang, C. (2024). Efficient multi-agent reinforcement learning by planning. In *Proceedings of the twelfth international conference on learning representations (ICLR)*. <https://openreview.net/forum?id=CpnKq3UJwp>
- Loveys, K., Antoni, M., Donkin, L., Sagar, M., & Broadbent, E. (2023). Comparing the feasibility and acceptability of a virtual human, teletherapy, and an e-manual in delivering a stress management intervention to distressed adult women: Pilot study. *JMIR Formative Research*, 7, e42390. <https://doi.org/10.2196/42390>
- Malik, P. K., Bhatt, H., & Sharma, M. (2025). AI integration in healthcare systems—A review of the problems and potential associated with integrating AI in healthcare for disease detection and diagnosis. *AI in Disease Detection: Advancements and Applications*, 191–213. <https://doi.org/10.1002/9781394278695.ch9>
- Mevlevioğlu, D., Tabircan, S., & Murphy, D. (2024). Real-time classification of anxiety in virtual reality therapy using biosensors and a convolutional neural network. *Biosensors*, 14(3), 131. <https://doi.org/10.3390/bios14030131>
- Miranda, O., Kiehl, S. M., Qi, X., Brannock, M. D., Kosten, T., Ryan, N. D., et al. (2024). Enhancing post-traumatic stress disorder patient assessment: Leveraging natural language processing for research of domain criteria identification using electronic medical records. *BMC Medical Informatics and Decision Making*, 24(1), 154. <https://doi.org/10.1186/s12911-024-02554-8>
- Mire, J., Aysola, Z. T., Chechelnitsky, D., Deas, N., Zerva, C., & Sap, M. (2025). Rejected dialects: Biases against African American language in reward models. *arXiv*. <https://arxiv.org/pdf/2502.12858>
- NHS England. (2025, May 2). *AI knowledge repository*. NHS Digital. <https://digital.nhs.uk/services/ai-knowledge-repository>
- Nicol, G., Wang, R., Graham, S., Dodd, S., & Garbutt, J. (2022). Chatbot-delivered cognitive behavioral therapy in adolescents with depression and anxiety during the COVID-19 pandemic: Feasibility and acceptability study. *JMIR Formative Research*, 6(11), e40242. <https://doi.org/10.2196/40242>
- Olawade, D. B., Wada, O. Z., Odetayo, A., David-Olawade, A. C., Asaolu, F., & Eberhardt, J. (2024). Enhancing mental health with artificial intelligence: Current trends and future prospects. *Journal of Medicine, Surgery, and Public Health*, 100099. <https://doi.org/10.1016/j.glmed.2024.100099>

- Park, S. H., & Han, K. (2018). Methodologic guide for evaluating clinical performance and effect of artificial intelligence technology for medical diagnosis and prediction. *Radiology*, 286(3), 800–809. <https://doi.org/10.1148/radiol.2017171920>
- Patel, J., Hung, C., & Katapally, T. R. (2024). Evaluating predictive artificial intelligence approaches used in mobile health platforms to forecast mental health symptoms among youth: A systematic review. *Psychiatry Research*, 116277. <https://doi.org/10.1016/j.psychres.2024.116277>
- Peng, W., Lee, H. R., & Lim, S. (2024). Leveraging chatbots to combat health misinformation for older adults: Participatory design study. *JMIR Formative Research*, 8(1), e60712. <https://doi.org/10.2196/60712>
- Peter, S., Riemer, K., & West, J. D. (2025). The benefits and dangers of anthropomorphic conversational agents. *Proceedings of the National Academy of Sciences*, 122(22), e2415898122. <https://doi.org/10.1073/pnas.2415898122>
- Rashid, M., & Sharma, M. (2025). AI-assisted diagnosis and treatment planning—A discussion of how AI can assist healthcare professionals in making more accurate diagnoses and treatment plans for diseases. *AI in Disease Detection: Advancements and Applications*, 313–336. <https://doi.org/10.1002/9781394278695.ch14>
- Ray, A., Bhardwaj, A., Malik, Y. K., Singh, S., & Gupta, R. (2022). Artificial intelligence and psychiatry: An overview. *Asian Journal of Psychiatry*, 70, 103021. <https://doi.org/10.1016/j.ajp.2022.103021>
- Rogan, J., Bucci, S., & Firth, J. (2024). Health care professionals' views on the use of passive sensing, AI, and machine learning in mental health care: Systematic review with meta-synthesis. *JMIR Mental Health*, 11, e49577. <https://doi.org/10.2196/49577>
- Rollwage, M., Habicht, J., Juechems, K., Carrington, B., Viswanathan, S., Stylianou, M., et al. (2023). Using conversational AI to facilitate mental health assessments and improve clinical efficiency within psychotherapy services: Real-world observational study. *JMIR AI*, 2(1), e44358. <https://doi.org/10.2196/44358>
- Scheinker, D., & Brandeau, M. L. (2020). Implementing analytics projects in a hospital: Successes, failures, and opportunities. *INFORMS Journal on Applied Analytics*, 50(3), 176–189. <https://doi.org/10.1287/inte.2020.1036>
- Schnack, H. G., & Kahn, R. S. (2016). Detecting neuroimaging biomarkers for psychiatric disorders: Sample size matters. *Frontiers in Psychiatry*, 7, 50. <https://doi.org/10.3389/fpsy.2016.00050>
- Shakespeare, T. (2013). *Disability rights and wrongs revisited*. Routledge. <https://doi.org/10.4324/9781315887456>
- Suharwardy, S., Ramachandran, M., Leonard, S. A., Gunaseelan, A., Lyell, D. J., Darcy, A., et al. (2023). Feasibility and impact of a mental health chatbot on postpartum mental health: A randomized controlled trial. *AJOG Global Reports*, 3(3), 100165. <https://doi.org/10.1016/j.xagr.2023.100165>
- Swaminathan, A., López, I., Mar, R. A. G., Heist, T., McClintock, T., Caoili, K., et al. (2023). Natural language processing system for rapid detection and intervention of mental health crisis chat messages. *npj Digital Medicine*, 6(1), 213. <https://doi.org/10.1038/s41746-023-00951-3>
- Tavory, T. (2024). Regulating AI in mental health: Ethics of care perspective. *JMIR Mental Health*, 11(1), e58493. <https://doi.org/10.2196/58493>
- Thapa, S., & Adhikari, S. (2024). GPT-4o and multimodal large language models as companions for mental wellbeing. *Asian Journal of Psychiatry*, 99, 104157. <https://doi.org/10.1016/j.ajp.2024.104157>
- Thieme, A., Hanratty, M., Lyons, M., Palacios, J., Marques, R. F., Morrison, C., & Doherty, G. (2023). Designing human-centered AI for mental health: Developing clinically relevant applications for online CBT treatment. *ACM Transactions on Computer-Human Interaction*, 30(2), 1–50. <https://doi.org/10.1145/3564752>
- Tsvetanova, A., Sperrin, M., Peek, N., Buchan, I., Hyland, S., & Martin, G. P. (2021). Missing data was handled inconsistently in UK prediction models: A review of method used. *Journal of Clinical Epidemiology*, 140, 149–158. <https://doi.org/10.1016/j.jclinepi.2021.09.008>
- Vertsberger, D., Naor, N., & Winsberg, M. (2022). Adolescents' Well-being while using a mobile artificial intelligence-powered acceptance commitment therapy tool: Evidence from a longitudinal study. *JMIR AI*, 1(1), e38171. <https://doi.org/10.2196/38171>

- Walsh, C. G., Johnson, K. B., Ripperger, M., Sperry, S., Harris, J., Clark, N., et al. (2021). Prospective validation of an electronic health record–based, real-time suicide risk model. *JAMA Network Open*, 4(3), e211428–e211428. <https://doi.org/10.1001/jamanetworkopen.2021.1428>
- Warrier, U., Warrier, A., & Khandelwal, K. (2023). Ethical considerations in the use of artificial intelligence in mental health. *The Egyptian Journal of Neurology, Psychiatry and Neurosurgery*, 59(1), 139. <https://doi.org/10.1186/s41983-023-00735-2>
- Wiens, J., Saria, S., Sendak, M., Ghassemi, M., Liu, V. X., Doshi-Velez, F., et al. (2019). Do no harm: A roadmap for responsible machine learning for health care. *Nature Medicine*, 25(9), 1337–1340. <https://doi.org/10.1038/s41591-019-0548-6>
- World Health Organization. (2021). *Ethics and governance of artificial intelligence for health*. <https://www.who.int/publications/i/item/9789240029200>
- Wysa. (2025). *Wysa: AI for mental health support*. <https://www.wysa.com/>
- Yang, Y., Wang, C., Xiang, X., & An, R. (2025, February). AI applications to reduce loneliness among older adults: A systematic review of effectiveness and technologies. *Healthcare*, 13(5), 446. <https://doi.org/10.3390/healthcare13050446>
- Yang, Z., Reddy, V. J., Kesidi, R., & Jin, F. (2019, August). Addict free—a smart and connected relapse intervention mobile app. In *Proceedings of the 16th international symposium on spatial and temporal databases* (pp. 202–205). doi:10.1145/3340964.3340986.
- Yuan, Y., Liu, Z., Miao, W., & Tian, X. (2024). Automatic screening for posttraumatic stress disorder in early adolescents following the ya'an earthquake using text mining techniques. *Frontiers in Psychiatry*, 15, 1439720. <https://doi.org/10.3389/fpsyt.2024.1439720>
- Zhao, G., Li, X., Li, Y., & Pietikäinen, M. (2023). Facial micro-expressions: An overview. *Proceedings of the IEEE*, 111(10), 1215–1235. <https://doi.org/10.1109/JPROC.2023.3275192>
- Zheng, L., Wang, O., Hao, S., Ye, C., Liu, M., Xia, M., et al. (2020). Development of an early-warning system for high-risk patients for suicide attempt using deep learning and electronic health records. *Translational Psychiatry*, 10(1), 72. <https://doi.org/10.1038/s41398-020-0684-2>
- Zidar, T., Morrow, E. M., & Stockley, R. (2021). Ensuring patient and public involvement in the transition to AI-assisted mental health care: A systematic scoping review and agenda for design justice. *Health Expectations*, 24(4), 1072–1124. <https://doi.org/10.1111/hex.13299>

Open Access This chapter is licensed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

The images or other third party material in this chapter are included in the chapter's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the chapter's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.

