

Sampling People, Records, & Networks


Jim Lepkowski, PhD

Professor & Research Professor *Emeritus*

Institute for Social Research, University of Michigan


Research Professor,

Joint Program in Survey Methodology, University of Maryland




Unit 4


- **1 Forming groups**
 - **2 Sampling variance**
 - **3 More on grouping**
 - **4 Allocate sample**
 - **5 Other allocations**
 - **6 Weights**
- Unit 1: Sampling as a research tool
 - Unit 2: Mere randomization
 - Unit 3: Saving money
 - Unit 4: Being more efficient
 - Forming groups
 - Sampling variance
 - More on grouping
 - Allocate sample
 - **Other allocations**
 - Weights to combine across strata
 - Unit 5: Simplifying sampling
 - Unit 6: Some extensions & applications



- Other allocations
 - Proportionate
 - Equal sample size
 - Domain estimation
 - Effect on precision
- Unit 1: Sampling as a research tool
 - Unit 2: Mere randomization
 - Unit 3: Saving money
 - Unit 4: Being more efficient
 - Forming groups
 - Sampling variance
 - More on grouping
 - Allocate sample
 - **Other allocations**
 - Weights to combine across strata
 - Unit 5: Simplifying sampling
 - Unit 6: Some extensions & applications




- Equal sample size
 - Domain estimation
 - Effect on precision
- **Another allocation that may make sense in other situations is to take the same or about the same number in each stratum:**




- Equal sample size
- Domain estimation
- Effect on precision

h	Stratum	N_h	W_h	n_h
1	Female, Assistant	40	0.1000	13
2	Female, Associate	25	0.0625	14
3	Female, Full	20	0.0500	13
4	Male, Assistant	75	0.1875	14
5	Male, Associate	50	0.1250	13
6	Male, Full	190	0.4750	13
Total		400	1.0000	80



- Equal sample size
 - Domain estimation
 - Effect on precision
- Here, we end up with 27 assistant, 27 associate, and 26 full professors
 - Compared to the proportionate allocation of 23, 15, and 38
 - Or we have 40 females and 40 males



- Equal sample size
 - Domain estimation
 - Effect on precision
- Here, we end up with 27 assistant, 27 associate, and 26 full professors
 - Compared to the proportionate allocation of 23, 15, and 38
 - Or we have 40 females and 40 males
 - These are of course quite different than the proportionate allocation (17 females and 63 males)



- Equal sample size
 - Domain estimation
 - Effect on precision
- Here, we end up with 27 assistant, 27 associate, and 26 full professors
 - Compared to the proportionate allocation of 23, 15, and 38
 - Or we have 40 females and 40 males
 - These are of course quite different than the proportionate allocation (17 females and 63 males)
 - Purposes of the 'equal' allocation
 - **Comparisons** of different sized subgroups
 - **Better estimates** for small sized groups



- Equal sample size
 - Domain estimation
 - Effect on precision
- Here, we end up with 27 assistant, 27 associate, and 26 full professors
 - Compared to the proportionate allocation of 23, 15, and 38
 - Or we have 40 females and 40 males
 - These are of course quite different than the proportionate allocation (17 females and 63 males)
 - Purposes of the 'equal' allocation
 - **Comparisons** of different sized subgroups
 - **Better estimates** for small sized groups
 - Weighting the sample is necessary if we are going to combine across subgroups to get back to conclusions about the total population




- Equal sample size
- Domain estimation
- Effect on precision



- Equal sample size
- **Domain estimation**
- Effect on precision

2011 Population by Province



Source: Statistics Canada, Created by: Mack D. Male

License: <http://creativecommons.org/licenses/by-sa/2.5/ca/>



- Equal sample size
- Domain estimation
- Effect on precision
- On occasion, use an allocation that gives us the smallest sampling variance of all allocations – **optimum allocation**



- Equal sample size
- Domain estimation
- Effect on precision
- On occasion, use an allocation that gives us the smallest sampling variance of all allocations – **optimum allocation**
- These purposes require us to look at the size of strata, and the variability within strata, and sometimes even the cost within strata



- Equal sample size
- Domain estimation
- Effect on precision
- On occasion, use an allocation that gives us the smallest sampling variance of all allocations – **optimum allocation**
- These purposes require us to look at the size of strata, and the variability within strata, and sometimes even the cost within strata
- This kind of minimum variance estimation is beyond the scope of this course



- Equal sample size
 - Domain estimation
 - Effect on precision
- On occasion, use an allocation that gives us the smallest sampling variance of all allocations – **optimum allocation**
 - These purposes require us to look at the size of strata, and the variability within strata, and sometimes even the cost within strata
 - This kind of minimum variance estimation is beyond the scope of this course
 - It arises in studies of a single variable, and when there is a lot of variance in the data – things like income, or expenditures, or wealth, and so on



- Equal sample size
- Domain estimation
- Effect on precision

- On occasion, use an allocation that gives us the smallest sampling variance of all allocations – **optimum allocation**
- These purposes require us to look at the size of strata, and the variability within strata, and sometimes even the cost within strata
- This kind of minimum variance estimation is beyond the scope of this course
- It arises in studies of a single variable, and when there is a lot of variance in the data – things like income, or expenditures, or wealth, and so on
- Minimum variance allocation does not arise often in much of the social, public health, medical, or other sciences



- Equal sample size
 - Domain estimation
 - Effect on precision
- **Let's consider one other simpler example to see how these allocation can affect the sampling variance:**

- Equal sample size
- Domain estimation
- Effect on precision

Population	Stratum 1 Qatari	Stratum 2 White & Blue Collar Expatriate (Other)
Size N 1,000,000	N_1 200,000	N_2 800,000
Variance S^2 1,800,000	S_1^2 4,000,000	S_2^2 1,000,000
Mean \bar{Y} 1,400	\bar{Y}_1 3,000	\bar{Y}_2 1,000

- Equal sample size
- Domain estimation
- Effect on precision

Suppose $n_1 = 240, n_2 = 960$

What will be $Var(\bar{y})$?

$$\begin{aligned} Var(\bar{y}) &= \sum_{h=1}^2 W_h^2 \frac{(1-f_h)}{n_h} S_h^2 \approx \frac{W_1^2 S_1^2}{n_1} + \frac{W_2^2 S_2^2}{n_2} \\ &= \frac{(0.2)^2 (4000000)}{240} + \frac{(0.8)^2 (1000000)}{960} \\ &= 666.7 + 666.7 \\ &= 1333 \end{aligned}$$

- Equal sample size
- Domain estimation
- **Effect on precision**

For $n = 1200$ what will be $Var_{SRS}(\bar{y})$?

$$\begin{aligned}
 Var_{SRS}(\bar{y}) &= \frac{(1-f)}{n} S^2 \\
 &= \frac{\left(1 - \frac{1,200}{1,000,000}\right)}{1,200} (1800000) \\
 &\approx \frac{1800000}{1200} = 1500
 \end{aligned}$$

- Equal sample size
- Domain estimation
- Effect on precision

- **As for cluster sampling,**

$$deff(\bar{y}) = \frac{Var(\bar{y}) \text{ for a given design}}{Var_{SRS}(\bar{y}) \text{ of same size}}$$

- **For this example,**

$$deff(\bar{y}) = \frac{Var(\bar{y})}{Var_{SRS}(\bar{y})} = \frac{1333}{1500} = 0.89$$

- Equal sample size
- Domain estimation
- **Effect on precision**

What about for each of the following combinations of sample sizes across the two strata?

(1) $n_1 = 100$ $n_2 = 1100$

(2) $n_1 = 240$ $n_2 = 960$

(3) $n_1 = 400$ $n_2 = 800$

(4) $n_1 = 600$ $n_2 = 600$

(5) $n_1 = 960$ $n_2 = 240$

- Equal sample size
- Domain estimation
- Effect on precision

What about for each of the following combinations of sample sizes across the two strata?

(1) $n_1 = 100$ $n_2 = 1100$: $Var(\bar{y}) = 2133$ & $deff(\bar{y}) = 1.45$

(2) $n_1 = 240$ $n_2 = 960$

(3) $n_1 = 400$ $n_2 = 800$

(4) $n_1 = 600$ $n_2 = 600$

(5) $n_1 = 960$ $n_2 = 240$

- Equal sample size
- Domain estimation
- Effect on precision

What about for each of the following combinations of sample sizes across the two strata?

- (1) $n_1 = 100$ $n_2 = 1100$: $Var(\bar{y}) = 2133$ & $deff(\bar{y}) = 1.45$
- (2) $n_1 = 240$ $n_2 = 960$: $Var(\bar{y}) = 1333$ & $deff(\bar{y}) = \mathbf{0.89}$
- (3) $n_1 = 400$ $n_2 = 800$
- (4) $n_1 = 600$ $n_2 = 600$
- (5) $n_1 = 960$ $n_2 = 240$

- Equal sample size
- Domain estimation
- Effect on precision

What about for each of the following combinations of sample sizes across the two strata?

- (1) $n_1 = 100$ $n_2 = 1100$: $Var(\bar{y}) = 2133$ & $deff(\bar{y}) = 1.45$
(2) $n_1 = 240$ $n_2 = 960$: $Var(\bar{y}) = 1333$ & $deff(\bar{y}) = \mathbf{0.89}$
(3) $n_1 = 400$ $n_2 = 800$: $Var(\bar{y}) = 1200$ & $deff(\bar{y}) = 0.80$
(4) $n_1 = 600$ $n_2 = 600$: $Var(\bar{y}) = 1333$ & $deff(\bar{y}) = 0.89$
(5) $n_1 = 960$ $n_2 = 240$

- Equal sample size
- Domain estimation
- Effect on precision

What about for each of the following combinations of sample sizes across the two strata?

- (1) $n_1 = 100$ $n_2 = 1100$: $Var(\bar{y}) = 2133$ & $deff(\bar{y}) = 1.45$
- (2) $n_1 = 240$ $n_2 = 960$: $Var(\bar{y}) = 1333$ & $deff(\bar{y}) = \mathbf{0.89}$
- (3) $n_1 = 400$ $n_2 = 800$: $Var(\bar{y}) = 1200$ & $deff(\bar{y}) = 0.80$
- (4) $n_1 = 600$ $n_2 = 600$: $Var(\bar{y}) = 1333$ & $deff(\bar{y}) = 0.89$
- (5) $n_1 = 960$ $n_2 = 240$: $Var(\bar{y}) = 2833$ & $deff(\bar{y}) = 2.36$

Unit 4

- **1 Forming groups**
 - **2 Sampling variance**
 - **3 More on grouping**
 - **4 Allocate sample**
 - **5 Other allocations**
 - **6 Weights**
- Unit 1: Sampling as a research tool
 - Unit 2: Mere randomization
 - Unit 3: Saving money
 - Unit 4: Being more efficient
 - Forming groups
 - Sampling variance
 - More on grouping
 - Allocate sample
 - Other allocations
 - **Weights to combine across strata**
 - Unit 5: Simplifying sampling
 - Unit 6: Some extensions & applications

