Sampling People, Records, & Networks

Jim Lepkowski, PhD **Professor & Research Professor Emeritus** Institute for Social Research, University of Michigan Research Professor. Joint Program in Survey Methodology, University of Maryland






Unit 4

- I Forming groups
- 2 Sampling variance
- 3 More on grouping
- 4 Allocate sample
- 5 Other allocations
- 6 Weights


- Unit 1: Sampling as a research tool
- Unit 2: Mere randomization
- Unit 3: Saving money
- Unit 4: Being more efficient
 - Forming groups
 - Sampling variance
 - More on grouping
 - Allocate sample
 - Other allocations
 - Weights to combine across strata
- Unit 5: Simplifying sampling
- Unit 6: Some extensions & applications




- Principles
- Example
- Confidence interval
- Design effect
- Effective sample size

- Unit I: Sampling as a research tool
- Unit 2: Mere randomization
- Unit 3: Saving money
- Unit 4: Being more efficient
 - Forming groups
 - Sampling variance
 - More on grouping
 - Allocate sample
 - Other allocations
 - Weights to combine across strata
- Unit 5: Simplifying sampling
- Unit 6: Some extensions & applications







- Principles
- Example
- Confidence interval
- Design effect
- Effective sample size

• In theory,

$$Var(\overline{y}) = \sum_{h=1}^{H} W_h^2 Var(\overline{y}_h)$$



- Principles
- Example
- Confidence interval
- Design effect
- Effective sample size

In theory,

$$Var(\overline{y}) = \sum_{h=1}^{H} W_h^2 Var(\overline{y}_h)$$

• Estimate this variance by

$$var(\overline{y}) = \sum_{h=1}^{H} W_h^2 var(\overline{y}_h)$$



- Principles
- Example
- Confidence interval
- Design effect
- Effective sample size

• In theory,

$$Var(\overline{y}) = \sum_{h=1}^{H} W_h^2 Var(\overline{y}_h)$$

• Estimate this variance by

$$var(\overline{y}) = \sum_{h=1}^{H} W_h^2 var(\overline{y}_h)$$

- And what is $var(\bar{y}_h)$?
- For SRS within strata, $var(\bar{y}_h) = \frac{(1-f_h)}{n_h} s_h^2$



- Principles
- Example
- Confidence interval
- Design effect
- Effective sample size

• In theory,

$$Var(\overline{y}) = \sum_{h=1}^{H} W_h^2 Var(\overline{y}_h)$$

• Estimate this variance by

$$var(\overline{y}) = \sum_{h=1}^{H} W_h^2 var(\overline{y}_h)$$

- And what is $var(\bar{y}_h)$?
- For SRS within strata, $var(\bar{y}_h) = \frac{(1-f_h)}{n_h} s_h^2$
- We thus need the within stratum variances:



Survey Data Collection & Analytic Specialization

Sampling People, Records, & Networks

- Principles
- Example
- Confidence interval
- Design effect
- Effective sample size

h	Stratum	N_h	$W_{_h}$	n_h	f_h	$\overline{\mathcal{Y}}_h$	s_h^2
1	Assistant	115	0.2875	23	0.2	50	125
2	Associate	75	0.1875	15	0.2	70	250
3	Full	210	0.5250	42	0.2	90	500
Total		400	1.0000	80	0.2	\$74.75	



- Principles
- Example
- Confidence interval
- Design effect
- Effective sample size

- This simplifies a little, since here $I-f_h = 0.8$
- The calculation is really beyond the scope of this course, but since we've come this far ...



- Principles
- Example
- Confidence interval
- Design effect
- Effective sample size

- This simplifies a little, since here $I-f_h = 0.8$
- The calculation is really beyond the scope of this course, but since we've come this far ...

•
$$var(\bar{y}) = \frac{(0.2875)^2(0.8)(125)}{23} + \frac{(0.1875)^2(0.8)250}{15} + \frac{(0.5250)^2(0.8)500}{42} = 3.453$$



- Principles
- Example
- Confidence interval
- Design effect
- Effective sample size

- This simplifies a little, since here $I-f_h = 0.8$
- The calculation is really beyond the scope of this course, but since we've come this far ...

•
$$var(\bar{y}) = \frac{(0.2875)^2(0.8)(125)}{23} + \frac{(0.1875)^2(0.8)250}{15} + \frac{(0.5250)^2(0.8)500}{42} = 3.453$$

• And $se(\bar{y}) = 1.858$




- Principles
- Example
- Confidence interval
- Design effect
- Effective sample size

- This simplifies a little, since here $I-f_h = 0.8$
- The calculation is really beyond the scope of this course, but since we've come this far ...

•
$$var(\bar{y}) = \frac{(0.2875)^2(0.8)(125)}{23} + \frac{(0.1875)^2(0.8)250}{15} + \frac{(0.5250)^2(0.8)500}{42} = 3.453$$

- And $se(\bar{y}) = 1.858$
- Thus, we have completed step 6a & 6b within stratum sampling variances and combining across strata





6 Standard error

$$se(\overline{y}) = \sqrt{\sum_{h=1}^{2} W_h^2 \frac{\left(1 - f_h\right)}{n_h} s_h^2}$$

7 Confidence interval

4 Estimate
$$\overline{y}_{1} = \frac{1}{n_{1}} \sum_{i=1}^{n_{1}} y_{1i}$$
 $\overline{y}_{2} = \frac{1}{n_{2}} \sum_{i=1}^{n_{2}} y_{2i}$ $\overline{y}_{w} = \sum_{h=1}^{2} W_{h} \overline{y}_{h}$

Survey Data Collection & Analytic Specialization

- Principles
- Example
- Confidence interval
- Design effect
- Effective sample size

- The last step, step 7, is the confidence interval
- Here, let's use the t-distribution
- We have n_h 1 degrees of freedom for each stratum and n H for combining across strata
- For a 95% confidence interval, then, $t_{(0.975,80-3)} = 1.991$



Survey Data Collection & Analytic Specialization

- Principles
- Example
- Confidence interval
- Design effect
- Effective sample size

- The last step, step 7, is the confidence interval
- Here, let's use the t-distribution
- We have n_h I degrees of freedom for each stratum and n H for combining across strata
- For a 95% confidence interval, then, $t_{(0.975,80-3)} = 1.991$
- The 95% confidence interval is then

•
$$(\bar{y} - t_{(0.975,77)}se(\bar{y}), \bar{y} + t_{(0.975,77)}se(\bar{y}))$$

- $(74.75 1.991 \times 1.858, 74.75 + 1.991 \times 1.858)$
- (71.05, 78.45)



- Principles
- Example
- Confidence interval
- Design effect
- Effective sample size

- Here, as in cluster sampling, there is another issue to be addressed – how does the sampling variance from stratified sampling compare to simple random sampling?
- That is, what is the design effect, deff?



- Principles
- Example
- Confidence interval
- Design effect
- Effective sample size

- Here, as in cluster sampling, there is another issue to be addressed – how does the sampling variance from stratified sampling compare to simple random sampling?
- That is, what is the design effect, deff?
- For SRS, we need to calculate $var(\bar{y}) = (1-f)\frac{s^2}{n}$
- From a separate calculation, $s^2 = 647.8$



- Principles
- Example
- Confidence interval
- Design effect
- Effective sample size

• For our sample of size n=80, from a population of size N=400, or a sampling fraction f=0.2, $var(\bar{y})=(1-0.2)\frac{647.8}{80}=6.478$

- Principles
- Example
- Confidence interval
- Design effect
- Effective sample size

• For our sample of size n = 80, from a population of size N = 400, or a sampling fraction f = 0.2,

$$var(\bar{y}) = (1 - 0.2) \frac{647.8}{80} = 6.478$$

• Compared to the stratified sample, $deff(\bar{y}) =$

$$\frac{var(\bar{y})}{var_{SRS}(\bar{y})} = \frac{3.453}{6.478} = 0.5331$$



- Principles
- Example
- Confidence interval
- Design effect
- Effective sample size

• For our sample of size n = 80, from a population of size N = 400, or a sampling fraction f = 0.2,

$$var(\bar{y}) = (1 - 0.2) \frac{647.8}{80} = 6.478$$

- Compared to the stratified sample, $deff(\bar{y}) = \frac{var(\bar{y})}{var_{SRS}(\bar{y})} = \frac{3.453}{6.478} = 0.5331$
- A 47% reduction in sampling variance



- Principles
- Example
- Confidence interval
- Design effect
- Effective sample size

 As we did for cluster samples, this design effect can be thought of in several ways



- Principles
- Example
- Confidence interval
- Design effect
- Effective sample size

- As we did for cluster samples, this design effect can be thought of in several ways
- For example, the actual sample size of 400 in the stratified proportionately allocated sample is the equivalent of having a simple random sample of

$$n_{eff} = \frac{400}{0.5331} = 750$$



- Principles
- Example
- Confidence interval
- Design effect
- Effective sample size

- As we did for cluster samples, this design effect can be thought of in several ways
- For example, the actual sample size of 400 in the stratified proportionately allocated sample is the equivalent of having a simple random sample of

$$n_{eff} = \frac{400}{0.5331} = 750$$

 That is, the gains in precision are the equivalent of adding 350 cases to the sample



- Principles
- Example
- Confidence interval
- Design effect
- Effective sample size

- As we did for cluster samples, this design effect can be thought of in several ways
- For example, the actual sample size of 400 in the stratified proportionately allocated sample is the equivalent of having a simple random sample of


$$n_{eff} = \frac{400}{0.5331} = 750$$

- That is, the gains in precision are the equivalent of adding 350 cases to the sample
- Alternatively, the standard errors are smaller and the confidence intervals are narrower by a factor of

$$1 - \sqrt{0.5331} = 1 - 0.7301 = 0.2699$$
, 27%




- Principles
- Example
- Confidence interval
- Design effect
- Effective sample size






- Principles
- Example
- Confidence interval
- Design effect
- Effective sample size





- Principles
- Example
- Confidence interval
- Design effect
- Effective sample size





Unit 4

- I Forming groups
- 2 Sampling variance
- 3 More on grouping
- 4 Allocate sample
- 5 Other allocations
- 6 Weights

- Unit I: Sampling as a research tool
- Unit 2: Mere randomization
- Unit 3: Saving money
- Unit 4: Being more efficient
 - Forming groups
 - Sampling variance
 - More on grouping
 - Allocate sample
 - Other allocations
 - Weights to combine across strata
- Unit 5: Simplifying sampling
- Unit 6: Some extensions & applications

