Sampling People, Records, & Networks


Jim Lepkowski, PhD
Professor & Research Professor Emeritus
Institute for Social Research, University of Michigan
Research Professor,
Joint Program in Survey Methodology, University of Maryland



Unit 2

- I Simple random sampling
- 2 History
- 3 Sampling distribution
- 4 Sample size
- 5 Margin of error
- 6 Sample & population size

- Unit I: Sampling as a research tool
- Unit 2: Mere randomization
 - Lecture 1: Simple Random Sampling (SRS)
 - Lecture 2: A short history
 - Lecture 3: The SRS sampling distribution
 - Lecture 4: Sample size
 - Lecture 5: Margin of error
 - Lecture 6: Sample size & population size
- Unit 3: Saving money
- Unit 4: Being more efficient
- Unit 5: Simplifying sampling
- Unit 6: Some extensions & applications



Survey Data Collection & Analytic Specialization

Unit 2


- I Simple random sampling
- 2 History
- 3 Sampling distribution
- 4 Sample size
- 5 Margin of error
- 6 Sample & population size

- Unit I: Sampling as a research tool
- Unit 2: Mere randomization
 - Lecture 1: Simple Random Sampling (SRS)
 - Lecture 2: A short history
 - Lecture 3: The SRS sampling distribution
 - Lecture 4: Sample size
 - Lecture 5: Margin of error
 - Lecture 6: Sample size & population size
- Unit 3: Saving money
- Unit 4: Being more efficient
- Unit 5: Simplifying sampling
- Unit 6: Some extensions & applications



- Basic framework
- Properties of the sampling distribution
- Confidence intervals

- Unit I: Sampling as a research tool
- Unit 2: Mere randomization
 - Lecture I: Simple Random Sampling (SRS)
 - Lecture 2: A short history
 - Lecture 3: The SRS sampling distribution
 - Lecture 4: Sample size
 - Lecture 5: Margin of error
 - Lecture 6: Sample size & population size
- Unit 3: Saving money
- Unit 4: Being more efficient
- Unit 5: Simplifying sampling
- Unit 6: Some extensions & applications



Survey Data Collection & Analytic Specialization

- Basic framework
- Properties of the sampling distribution
- Confidence intervals



- A sample design for which the unit of selection is the population element
- Basic framework: Neyman, 1934
 - Must be applicable to all populations
 - Must not depend on assumptions about the population structure
 - Appropriate for large populations of elements

Survey Data Collection & Analytic Specialization

- Basic framework
- Properties of the sampling distribution
- Confidence intervals

- Repeated sampling
 - Objective (mechanical) selection of elements
 - Consider possible outcomes of the sampling process
 - Evaluation of the whole set of possible outcomes



- Basic framework
- Properties of the sampling distribution
- Confidence intervals



- The set of all possible values of the estimator that can be obtained with a given sample design
 - For a given sample we obtain a particular value, the estimate (such as \overline{y})
- We want to know ...
 - ... how likely is the estimate to be close to the population value?

- Basic framework
- Properties of the sampling distribution
- Confidence intervals

- In fact, we select just one sample
- The estimate may be correct, or incorrect
- Want to maximize the probability of a satisfactory estimate

- Basic framework
- Properties of the sampling distribution
- Confidence intervals

- Unbiasedness
 - Expected value (average value): $E(\overline{y})$

- Basic framework
- Properties of the sampling distribution
- Confidence intervals

- Unbiasedness
 - Expected value (average value): $E(\overline{y})$
 - Meaning of expected value:

$$E(\overline{y}) = \frac{1}{\binom{N}{n}} \sum_{s=1}^{\binom{N}{n}} \overline{y}_{s}$$

- Basic framework
- Properties of the sampling distribution
- Confidence intervals

- Unbiasedness
 - Expected value (average value): $E(\overline{y})$
 - Meaning of expected value:

$$E(\overline{y}) = \frac{1}{\binom{N}{n}} \sum_{s=1}^{\binom{N}{n}} \overline{y}_{s}$$

• Meaning of unbiasedness:

$$E(\overline{y}) = \frac{1}{\binom{N}{n}} \sum_{s=1}^{\binom{N}{n}} \overline{y}_s = \overline{Y}$$

Standard errors

For our SRS of n = 20,

$$var(\overline{y}) = \frac{(1-f)}{n}s^{2}$$

$$= \frac{\left(1 - \frac{20}{370}\right)}{20}766.62$$

$$= 36.26$$

$$se(\overline{y}) = \sqrt{var(\overline{y})} = 6.02$$

- Basic framework
- Properties of the sampling distribution
- Confidence intervals

- Variability from one sample to another
 - Variance of the estimator: $Var(\overline{y})$

- Basic framework
- Properties of the sampling distribution
- Confidence intervals

- Variability from one sample to another
 - Variance of the estimator: $Var(\overline{y})$
 - Meaning of the variance:

$$Var(\overline{y}) = \frac{1}{\binom{N}{n}} \sum_{s=1}^{\binom{N}{n}} (\overline{y}_s - E(\overline{y}))^2$$

- Basic framework
- Properties of the sampling distribution
- Confidence intervals

- Variability from one sample to another
 - Variance of the estimator: $Var(\overline{y})$
 - Meaning of the variance:

$$Var(\overline{y}) = \frac{1}{\binom{N}{n}} \sum_{s=1}^{\binom{N}{n}} (\overline{y}_s - E(\overline{y}))^2$$

• Algebraically equivalent formula:

$$Var(\overline{y}) = \frac{1}{\binom{N}{n}} \sum_{s=1}^{\binom{N}{n}} \left(\overline{y}_s - E(\overline{y})\right)^2 = \left(1 - \frac{n}{N}\right) \frac{S^2}{n}$$

- Basic framework
- Properties of the sampling distribution
- Confidence intervals

- Variability from one sample to another
 - Components:

$$S^{2}$$

$$\left(1 - \frac{n}{N}\right) = \left(1 - f\right)$$


$$\frac{1}{n}$$

- Basic framework
- Properties of the sampling distribution
- Confidence intervals

- Variability from one sample to another
 - Components:

$$S^{2} \left(1 - \frac{n}{N}\right) = \left(1 - f\right)$$

 $\frac{1}{n}$



- Basic framework
- Properties of the sampling distribution
- Confidence intervals

- Variability from one sample to another
 - Scale conversion:

$$SE(\overline{y}) = \sqrt{Var(\overline{y})} = \sqrt{\left(1 - \frac{n}{N}\right)\frac{S^2}{n}} = \frac{S}{\sqrt{n}}\sqrt{\left(1 - \frac{n}{N}\right)}$$

- Basic framework
- Properties of the sampling distribution
- Confidence intervals

- Estimating variability from one sample to another
 - Element variance: S^2
 - Estimated element variance:

$$s^{2} = \frac{1}{n-1} \sum_{i=1}^{n} (y_{i} - \overline{y})^{2} \qquad \left(= \frac{n}{n-1} p(1-p) \right)$$

- Basic framework
- Properties of the sampling distribution
- Confidence intervals

- Estimating variability from one sample to another
 - Element variance: S^2
 - Estimated element variance:

$$s^{2} = \frac{1}{n-1} \sum_{i=1}^{n} (y_{i} - \overline{y})^{2} \qquad \left(= \frac{n}{n-1} p(1-p) \right)$$

• Estimated sampling variance & standard error

$$var(\overline{y}) = \left(1 - \frac{n}{N}\right) \frac{s^2}{n} \qquad \left(= \left(1 - \frac{n}{N}\right) \frac{p(1-p)}{n-1}\right)$$

- Basic framework
- Properties of the sampling distribution
- Confidence intervals

- Estimating variability from one sample to another
 - Element variance: S^2
 - Estimated element variance:

$$s^{2} = \frac{1}{n-1} \sum_{i=1}^{n} (y_{i} - \overline{y})^{2} \qquad \left(= \frac{n}{n-1} p(1-p) \right)$$

Estimated sampling variance & standard error:

$$var(\overline{y}) = \left(1 - \frac{n}{N}\right) \frac{s^2}{n} \qquad \left(= \left(1 - \frac{n}{N}\right) \frac{p(1-p)}{n-1}\right)$$

$$se(\overline{y}) = \frac{s}{\sqrt{n}} \sqrt{\left(1 - \frac{n}{N}\right)}$$



- Basic framework
- Properties of the sampling distribution
- Confidence intervals

- For large samples, the sampling distribution of \overline{y} is Normal
 - Law of large numbers or Central limit theorem

- Basic framework
- Properties of the sampling distribution
- Confidence intervals

- For large samples, the sampling distribution of \overline{y} is Normal
 - Law of large numbers or Central limit theorem
- Form an interval around \overline{y} :

$$\overline{y} \pm z_{(1-\alpha/2)} \times se(\overline{y})$$

- Basic framework
- Properties of the sampling distribution
- Confidence intervals

- For large samples, the sampling distribution of \overline{y} is Normal
 - Law of large numbers or Central limit theorem
- Form an interval around \overline{y} :

$$\overline{y} \pm 1.96 \times se(\overline{y})$$

- $(1-\alpha)\%$ or 95% confidence interval
- A statement of uncertainty about our estimated mean

95% Confidence interval

For our SRS of
$$n = 20$$
,
$$\overline{y} \pm t_{(1-\alpha/2,n-1)} se(\overline{y})$$

$$78.6 \pm t_{(0.975,19)} \times 6.02$$

$$78.6 \pm 2.09 \times 6.02$$

$$(66.0,98.2)$$

Unit 2

- I Simple random sampling
- 2 History
- 3 Sampling distribution
- 4 Sample size
- 5 Margin of error
- 6 Sample & population size

- Unit I: Sampling as a research tool
- Unit 2: Mere randomization
 - Lecture 1: Simple Random Sampling (SRS)
 - Lecture 2: A short history
 - Lecture 3: The SRS sampling distribution
 - Lecture 4: Sample size
 - Lecture 5: Margin of error
 - Lecture 6: Sample size & population size
- Unit 3: Saving money
- Unit 4: Being more efficient
- Unit 5: Simplifying sampling
- Unit 6: Some extensions & applications

