

1 Notation

- n : number of teams;
- N : number of matches;
- T_k^q : last minute of q -th half of match;
- m_k^q : total goals in q -th half of match k ;
- $t_{k,l}^q$: minute of the l -th goal in the q -th half of match k ;
- $t_{k,0}^1 = 0$, $t_{k,m_k^1+1}^1 = T_k^1$, $t_{k,0}^2 = 45$ and $t_{k,m_k^2+1}^2 = T_k^2$;
- $H_{k,l}^q = \begin{cases} 1, & \text{if the home team scored a goal in interval } [t_{k,l}, t_{k,l+1}); \\ 0, & \text{otherwise;} \end{cases}$
- $A_{k,l}^q = \begin{cases} 1, & \text{if the away team scored a goal in interval } [t_{k,l}, t_{k,l+1}); \\ 0, & \text{otherwise;} \end{cases}$
- $\lambda_k(t)$: rate of goals for the home team in match k ;
- $\mu_k(t)$: rate of goals for the away team in match k ;
- $\rho = [t_k^1, t_k^2, H_k^1, H_k^2, A_k^1, A_k^2]$;
- $\Lambda(t_1, t_2) = \int_{t_1}^{t_2} \lambda_k(t) dt$;
- $\mathbf{Y}(t_1, t_2) = \int_{t_1}^{t_2} \mu_k(t) dt$.

2 Likelihood for match k

We define $T_k^1 = 45 + U_k^1$ and $T_k^2 = 90 + U_k^2$ and model U_k^1 and U_k^2 as

$$U_k^1 \sim \text{Poisson}(\pi_1)$$

$$U_k^2 \sim \text{Poisson}(\pi_2)$$

therefore,

$$L(\rho \mid T_k^1, T_k^2) = \prod_{l=0}^{m_k^1} \left[\exp \left\{ -\Lambda(t_{k,l}^1, t_{k,l+1}^1) \right\} \exp \left\{ -\mathbf{Y}(t_{k,l}^1, t_{k,l+1}^1) \right\} \left(\Lambda(t_{k,l}^1, t_{k,l+1}^1) \right)^{H_{k,l}^1} \left(\mathbf{Y}(t_{k,l}^1, t_{k,l+1}^1) \right)^{A_{k,l}^1} \right] \\ \prod_{l=0}^{m_k^2} \left[\exp \left\{ -\Lambda(t_{k,l}^2, t_{k,l+1}^2) \right\} \exp \left\{ -\mathbf{Y}(t_{k,l}^2, t_{k,l+1}^2) \right\} \left(\Lambda(t_{k,l}^2, t_{k,l+1}^2) \right)^{H_{k,l}^2} \left(\mathbf{Y}(t_{k,l}^2, t_{k,l+1}^2) \right)^{A_{k,l}^2} \right]$$

and

$$L(\rho, T_k^1, T_k^2) = \prod_{l=0}^{m_k^1} \left[\exp \left\{ -\Lambda(t_{k,l}^1, t_{k,l+1}^1) \right\} \exp \left\{ -\mathbf{Y}(t_{k,l}^1, t_{k,l+1}^1) \right\} \left(\Lambda(t_{k,l}^1, t_{k,l+1}^1) \right)^{H_{k,l}^1} \left(\mathbf{Y}(t_{k,l}^1, t_{k,l+1}^1) \right)^{A_{k,l}^1} \right] \\ \prod_{l=0}^{m_k^2} \left[\exp \left\{ -\Lambda(t_{k,l}^2, t_{k,l+1}^2) \right\} \exp \left\{ -\mathbf{Y}(t_{k,l}^2, t_{k,l+1}^2) \right\} \left(\Lambda(t_{k,l}^2, t_{k,l+1}^2) \right)^{H_{k,l}^2} \left(\mathbf{Y}(t_{k,l}^2, t_{k,l+1}^2) \right)^{A_{k,l}^2} \right] \\ \frac{\pi_1^{U_k^1} \exp\{-\pi_1\}}{U_k^1!} \frac{\pi_2^{U_k^2} \exp\{-\pi_2\}}{U_k^2!}.$$

$$\begin{aligned}
\lambda_k(t) &= \alpha_i \beta_j \gamma_h \lambda_{xy} e^{f(t)\xi_1} \\
&= \exp\{\alpha_i^* + \beta_j^* + \gamma_h^* + \lambda_{xy}^* + f(t)\xi_1\}
\end{aligned}$$

where $\alpha_i^* = \log(\alpha_i)$, $\beta_j^* = \log(\beta_j)$, $\gamma_h^* = \log(\gamma_h)$ and $\lambda_{xy}^* = \log(\lambda_{xy})$.

If $f(t) = \log(t+1)$ then

$$\begin{aligned}
\Lambda(t_1, t_2) &= \int_{t_1}^{t_2} \lambda_k(t) dt = \int_{t_1}^{t_2} \exp\{\alpha_i^* + \beta_j^* + \gamma_h^* + \lambda_{xy}^* + \log(t+1)\xi_1\} dt \\
&= \exp\{\alpha_i^* + \beta_j^* + \gamma_h^* + \lambda_{xy}^*\} \int_{t_1}^{t_2} \exp\{\log(t+1)\xi_1\} dt \\
&= \exp\{\alpha_i^* + \beta_j^* + \gamma_h^* + \lambda_{xy}^*\} \int_{t_1}^{t_2} (t+1)^{\xi_1} dt \\
&= \exp\{\alpha_i^* + \beta_j^* + \gamma_h^* + \lambda_{xy}^*\} \left(\frac{(t_2+1)^{\xi_1+1} - (t_1+1)^{\xi_1+1}}{\xi_1+1} \right)
\end{aligned}$$

If $f(t) = t$ then

$$\begin{aligned}
\Lambda(t_1, t_2) &= \int_{t_1}^{t_2} \lambda_k(t) dt = \int_{t_1}^{t_2} \exp\{\alpha_i^* + \beta_j^* + \gamma_h^* + \lambda_{xy}^* + t\xi_1\} dt \\
&= \exp\{\alpha_i^* + \beta_j^* + \gamma_h^* + \lambda_{xy}^*\} \int_{t_1}^{t_2} \exp\{t\xi_1\} dt \\
&= \exp\{\alpha_i^* + \beta_j^* + \gamma_h^* + \lambda_{xy}^*\} \left(\frac{\exp\{t_2\xi_1\} - \exp\{t_1\xi_1\}}{\xi_1} \right) \\
&= \exp\{\alpha_i^* + \beta_j^* + \gamma_h^* + \lambda_{xy}^*\} \left(\frac{\exp\{t_2\xi_1\}(1 - \exp\{\xi_1(t_1 - t_2)\})}{\xi_1} \right)
\end{aligned}$$