

Master's Thesis Presentation

On the Parameterized Complexity of SEMITOTAL DOMINATING SET On Graph Classes

Lukas Retschmeier

Theoretical Foundations of Artificial Intelligence
School of Computation
Technical University of Munich

February 28th, 2023

Creative Introduction

Motivation

Theory

Intractability
 ω_2 hardness

Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References

Our Plan for Today

1 Motivation

2 Theory

Intractability

ω_2 hardness

3 Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

4 Conclusions

Motivation

Theory

Intractability

ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

Motivation

Motivation

Theory

Intractability
 ω_2 hardness

Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References

DOMINATING SET

Input

Graph $G = (V, E)$, $k \in \mathbb{N}$

Question

Exists $D \subseteq V$ with $|D| \leq k$ such that $N[D] = V$?

- The domination number is the minimum cardinality of a ds of G , denotes as $\gamma(G)$
- **Observation:** In connected G every $v \in D$ has another $z \in D$ with $d(v, z) \leq 3$.

Motivation

Motivation

Theory

Intractability
 ω_2 hardness

Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References

DOMINATING SET

Input

Graph $G = (V, E)$, $k \in \mathbb{N}$

Question

Exists $D \subseteq V$ with $|D| \leq k$ such that $N[D] = V$?

- The domination number is the minimum cardinality of a ds of G , denotes as $\gamma(G)$
- **Observation:** In connected G every $v \in D$ has another $z \in D$ with $d(v, z) \leq 3$.

Motivation

Motivation

Theory

Intractability
 ω_2 hardness

Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References

DOMINATING SET

Input

Graph $G = (V, E)$, $k \in \mathbb{N}$

Question

Exists $D \subseteq V$ with $|D| \leq k$ such that $N[D] = V$?

- The domination number is the minimum cardinality of a ds of G , denotes as $\gamma(G)$
- **Observation:** In connected G every $v \in D$ has another $z \in D$ with $d(v, z) \leq 3$.

Motivation

Motivation

Theory

Intractability
 ω_2 hardness

Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References

TOTAL DOMINATING SET

Input

Graph $G = (V, E)$, $k \in \mathbb{N}$

Question

Exists $D \subseteq V$ with $|D| \leq k$ such that

$\forall d_1 \in X : \exists d_2 \in D \setminus \{d_1\}$ with $d(d_1, d_2) \leq 1$?

- The total domination number is the minimum cardinality of a tds of G , denoted as $\gamma_t(G)$.
- We say d_1 witnesses d_2 (and vice versa)

Motivation

Motivation

Theory

Intractability
 ω_2 hardness

Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References

TOTAL DOMINATING SET

InputGraph $G = (V, E)$, $k \in \mathbb{N}$ **Question**Exists $D \subseteq V$ with $|D| \leq k$ such that $\forall d_1 \in X : \exists d_2 \in D \setminus \{d_1\} \text{ with } d(d_1, d_2) \leq 1?$

- The total domination number is the minimum cardinality of a tds of G , denoted as $\gamma_t(G)$.
- We say d_1 witnesses d_2 (and vice versa)

Motivation

Motivation

Theory

Intractability
 ω_2 hardness

Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References

TOTAL DOMINATING SET

InputGraph $G = (V, E)$, $k \in \mathbb{N}$ **Question**Exists $D \subseteq V$ with $|D| \leq k$ such that $\forall d_1 \in X : \exists d_2 \in D \setminus \{d_1\} \text{ with } d(d_1, d_2) \leq 1?$

- The total domination number is the minimum cardinality of a tds of G , denoted as $\gamma_t(G)$.
- We say d_1 witnesses d_2 (and vice versa)

Motivation

Motivation

Theory

Intractability

 ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

SEMITOTAL DOMINATING SET

InputGraph $G = (V, E)$, $k \in \mathbb{N}$ **Question**Exists $D \subseteq V$ with $|D| \leq k$ such that $\forall d_1 \in X : \exists d_2 \in D \setminus \{d_1\}$ with $d(d_1, d_2) \leq 2$?

- The semitotal domination number is the minimum cardinality of an sds of G , denoted as $\gamma_{2t}(G)$.
- **Observation:** $\gamma(G) \leq \gamma_{2t}(G) \leq \gamma_t(G)$
- We say d_1 witnesses d_2 (and vice versa)

Motivation

Motivation

Theory

Intractability

 ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

SEMITOTAL DOMINATING SET

InputGraph $G = (V, E)$, $k \in \mathbb{N}$ **Question**Exists $D \subseteq V$ with $|D| \leq k$ such that $\forall d_1 \in X : \exists d_2 \in D \setminus \{d_1\}$ with $d(d_1, d_2) \leq 2$?

- The semitotal domination number is the minimum cardinality of an sds of G , denoted as $\gamma_{2t}(G)$.
- **Observation:** $\gamma(G) \leq \gamma_{2t}(G) \leq \gamma_t(G)$
- We say d_1 witnesses d_2 (and vice versa)

Motivation

Motivation

Theory

Intractability

 ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

SEMITOTAL DOMINATING SET

InputGraph $G = (V, E)$, $k \in \mathbb{N}$ **Question**Exists $D \subseteq V$ with $|D| \leq k$ such that $\forall d_1 \in X : \exists d_2 \in D \setminus \{d_1\}$ with $d(d_1, d_2) \leq 2$?

- The semitotal domination number is the minimum cardinality of an sds of G , denoted as $\gamma_{2t}(G)$.
- **Observation:** $\gamma(G) \leq \gamma_{2t}(G) \leq \gamma_t(G)$
- We say d_1 witnesses d_2 (and vice versa)

Motivation

Motivation

Theory

Intractability

 ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

SEMITOTAL DOMINATING SET

InputGraph $G = (V, E)$, $k \in \mathbb{N}$ **Question**Exists $D \subseteq V$ with $|D| \leq k$ such that $\forall d_1 \in X : \exists d_2 \in D \setminus \{d_1\}$ with $d(d_1, d_2) \leq 2$?

- The semitotal domination number is the minimum cardinality of an sds of G , denoted as $\gamma_{2t}(G)$.
- **Observation:** $\gamma(G) \leq \gamma_{2t}(G) \leq \gamma t(G)$
- We say d_1 witnesses d_2 (and vice versa)

Example: $\gamma(G) < \gamma_{2t}(G) < \gamma_t(G)$

Motivation

Theory

Intractability
 ω_2 hardness


Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size


Conclusions

References


DOMINATING SET



SEMITOTAL DOMINATING SET



TOTAL DOMINATING SET



Parameterized Complexity

Motivation

Theory

Intractability
 ω_2 hardness

Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References

- NP-hard? We expect problem to be **at least** exponential
- **Idea:** Limit combinatorial explosion to some aspect of the problem
- **Goal:** Find an algorithm running in time $\mathcal{O}(f(k) \cdot n^c)$ for **some** parameter k
- In this work: by solution size
- **Techniques:** Kernelization, Bounded Search Trees, ...

If possible, the problem is **fixed-parameter tractable**.

Parameterized Complexity

Motivation

Theory

Intractability

ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

- NP-hard? We expect problem to be **at least** exponential
 - **Idea:** Limit combinatorial explosion to some aspect of the problem
 - **Goal:** Find an algorithm running in time $\mathcal{O}(f(k) \cdot n^c)$ for **some** parameter k
 - In this work: by solution size
 - **Techniques:** Kernelization, Bounded Search Trees, ...
- If possible, the problem is **fixed-parameter tractable**.

Parameterized Complexity

Motivation

Theory

Intractability
 ω_2 hardness

Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References

- NP-hard? We expect problem to be **at least** exponential
 - **Idea:** Limit combinatorial explosion to some aspect of the problem
 - **Goal:** Find an algorithm running in time $\mathcal{O}(f(k) \cdot n^c)$ for **some** parameter k
 - In this work: by solution size
 - **Techniques:** Kernelization, Bounded Search Trees, ...
- If possible, the problem is **fixed-parameter tractable**.

Parameterized Complexity

Motivation

Theory

Intractability
 ω_2 hardness

Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References

- NP-hard? We expect problem to be **at least** exponential
 - **Idea:** Limit combinatorial explosion to some aspect of the problem
 - **Goal:** Find an algorithm running in time $\mathcal{O}(f(k) \cdot n^c)$ for **some** parameter k
 - In this work: by solution size
 - **Techniques:** Kernelization, Bounded Search Trees, ...
- If possible, the problem is **fixed-parameter tractable**.

Parameterized Complexity

Motivation

Theory

Intractability

ω_1 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

- NP-hard? We expect problem to be **at least** exponential
- **Idea:** Limit combinatorial explosion to some aspect of the problem
- **Goal:** Find an algorithm running in time $\mathcal{O}(f(k) \cdot n^c)$ for **some** parameter k
- In this work: by solution size
- **Techniques:** Kernelization, Bounded Search Trees, ...

If possible, the problem is **fixed-parameter tractable**.

Parameterized Complexity

Motivation

Theory

Intractability
 ω_2 hardness

Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References

- NP-hard? We expect problem to be **at least** exponential
- **Idea:** Limit combinatorial explosion to some aspect of the problem
- **Goal:** Find an algorithm running in time $\mathcal{O}(f(k) \cdot n^c)$ for **some** parameter k
- In this work: by solution size
- **Techniques:** Kernelization, Bounded Search Trees, ...

If possible, the problem is **fixed-parameter tractable**.

Fixed-Parameter Intractability

Motivation

Theory

Intractability

ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

- Class NP corresponds to whole hierarchy $W[i]$ in parameterized setting.
- Problems at least $W[1]$ -hard considered **fixed-parameter intractable**
- DOMINATING SET is $W[2]$ -complete
- **Tool for Proving Hardness:** FPT Reductions, preserving the parameter

Fixed-Parameter Intractability

Motivation

Theory

Intractability

ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

- Class NP corresponds to whole hierarchy $W[i]$ in parameterized setting.
- Problems at least $W[1]$ -hard considered **fixed-parameter intractable**
- DOMINATING SET is $W[2]$ -complete
- **Tool for Proving Hardness:** FPT Reductions, preserving the parameter

Fixed-Parameter Intractability

Motivation

Theory

Intractability

ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

- Class NP corresponds to whole hierarchy $W[i]$ in parameterized setting.
- Problems at least $W[1]$ -hard considered **fixed-parameter intractable**
- DOMINATING SET is $W[2]$ -complete
- **Tool for Proving Hardness:** FPT Reductions, preserving the parameter

Fixed-Parameter Intractability

- Class NP corresponds to whole hierarchy $W[i]$ in parameterized setting.
- Problems at least $W[1]$ -hard considered **fixed-parameter intractable**
- DOMINATING SET is $W[2]$ -complete
- **Tool for Proving Hardness:** FPT Reductions, preserving the parameter

Fixed-Parameter Intractability

Motivation

Theory

Intractability

ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

- Class NP corresponds to whole hierarchy $W[i]$ in parameterized setting.
- Problems at least $W[1]$ -hard considered **fixed-parameter intractable**
- DOMINATING SET is $W[2]$ -complete
- **Tool for Proving Hardness:** FPT Reductions, preserving the parameter

Complexity Comparison

Motivation

Theory

Intractability
 ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

Graph Class	DOMINATING SET		SEMITOTAL DOMINATING SET		TOTAL DOMINATING SET	
	classical	Parameterized	classical	Parameterized	classical	Parameterized
bipartite	NPc [4]	$W[2]$ [39]	NPc [25]	$W[2]$ (this)	NPc [32]	$W[2]$ (cite!)
line graph of bipartite	NPc [28]	?	NPc [19]	? (?)	NPc [35]	?
circle	NPc [26]	$W[1]$ [7]	NPc [27]	? (?)	NPc [35]	$W[1]$ [7]
chordal	NPc [6]	$W[2]$ [39]	NPc [25]	$W[2]$ (this)	NPc [37]	$W[1]$ [11] by <i>split</i>
s -chordal , $s > 3$	NPc [33]	$W[2]$ [33]	? (?)	? (?)	NPc [33]	$W[1]$ [33]
split	NPc [4]	$W[2]$ [39]	NPc [25]	$W[2]$ this	NPc [37]	$W[1]$ [11]
3-claw-free	NPc [14]	FPT [14]	Prob. Unk	Prob. Unk	NPc [35]	Unknown
t -claw-free, $t > 3$	NPc [14]	$W[2]$ [14]	Prob. Unknown	Unknown	NPc [35]	Prob. Unknown
chordal bipartite	NPc [36]	? (?)	NPc [25]	?		P [15]
planar	NPc (Sources!)	FPT [2]	NPc	FPT (this)	NPc	FPT [20]
undirected path	NPc [6]	FPT [18]	NPc [24]	?	NPc [31]	?
dually chordal		P [8]		? (attempted [19])		P [30]
strongly chordal		P [17]		P [40]	NPc [17]	
AT-free		P [29]		P [27]		P [29]
tolerance		P [22]		?		?
block		P [17]		P [24]		P [10]
interval		P [12]		P [38]		P [5]
<hr/>						
bounded clique-width		P [13]		P [13]		P [13]
bounded mim-width		P [3, 9]		P [19]		P [3, 9]

Status SEMITOTAL DOMINATING SET

Motivation

Theory

Intractability

ω_2 hardness

Kernel

Definitions

Rule 1


Rule 2

Rule 3

Kernel Size

Conclusions

References




Warmup: Intractability Results

ω_2 hard on split, chordal and bipartite graphs

- **Split Graph:** $G = \text{Clique} + \text{IndependentSet}$

Split Graphs

SEMITOTAL DOMINATING SET on *split* and *chordal* graphs is ω_2 -hard




Proof by fpt-reduction from PLANAR DOMINATING SET on split graphs:

- 1 Construct G^* by adding v with pendant z to clique. G^* split
- 2 If ds D in G , $D' = D \cup \{v\}$ is sds D' .
- 3 If sds D' in G' , $D \setminus \{v\}$ is D in G
- 4 Parameter k only changed by constant

Split Graphs

SEMITOTAL DOMINATING SET on *split* and *chordal* graphs is ω_2 -hard




Proof by fpt-reduction from PLANAR DOMINATING SET on split graphs:

- 1 **Construct** G^* by adding v with pendant z to clique. G^* split
- 2 If ds D in G , $D' = D \cup \{v\}$ is sds D' .
- 3 If sds D' in G' , $D \setminus \{v\}$ is D in G
- 4 Parameter k only changed by constant

Split Graphs

SEMITOTAL DOMINATING SET on *split* and *chordal* graphs is ω_2 -hard




Proof by fpt-reduction from PLANAR DOMINATING SET on split graphs:

- 1 **Construct** G^* by adding v with pendant z to clique. G^* split
- 2 If ds D in G , $D' = D \cup \{v\}$ is sds D' .
- 3 If sds D' in G' , $D \setminus \{v\}$ is D in G
- 4 Parameter k only changed by constant

Split Graphs

SEMITOTAL DOMINATING SET on *split* and *chordal* graphs is ω_2 -hard




Proof by fpt-reduction from PLANAR DOMINATING SET on split graphs:

- 1 **Construct** G^* by adding v with pendant z to clique. G^* split
- 2 If ds D in G , $D' = D \cup \{v\}$ is sds D' .
- 3 If sds D' in G' , $D \setminus \{v\}$ is D in G
- 4 Parameter k only changed by constant

Split Graphs

SEMITOTAL DOMINATING SET on *split* and *chordal* graphs is ω_2 -hard




Proof by fpt-reduction from PLANAR DOMINATING SET on split graphs:

- 1 **Construct** G^* by adding v with pendant z to clique. G^* split
- 2 If ds D in G , $D' = D \cup \{v\}$ is sds D' .
- 3 If sds D' in G' , $D \setminus \{v\}$ is D in G
- 4 Parameter k only changed by constant

Bipartite Graphs

SEMITOTAL DOMINATING SET on *bipartite* graphs is ω_2 -hard




Proof by fpt-reduction from PLANAR DOMINATING SET on bipart. graphs:

- 1 **Construct** Add new neighbor to each vertex and add d_1, d_2, u_1, u_2
- 2 If ds D in G , then $D' = D \cup \{d_1, d_2\}$ is sds in G'
- 3 Assume sds D' in G' . If $a_i \in D'$ (b_i), flip. $D = D' \setminus \{d_1, d_2\}$ is ds in G

Bipartite Graphs

SEMITOTAL DOMINATING SET on *bipartite* graphs is ω_2 -hard




Proof by fpt-reduction from PLANAR DOMINATING SET on bipart. graphs:

- 1 **Construct** Add new neighbor to each vertex and add d_1, d_2, u_1, u_2
- 2 If ds D in G , then $D' = D \cup \{d_1, d_2\}$ is sds in G'
- 3 Assume sds D' in G' . If $a_i \in D'$ (b_i), flip. $D = D' \setminus \{d_1, d_2\}$ is ds in G

Bipartite Graphs

SEMITOTAL DOMINATING SET on *bipartite* graphs is ω_2 -hard




Proof by fpt-reduction from PLANAR DOMINATING SET on bipart. graphs:

- 1 **Construct** Add new neighbor to each vertex and add d_1, d_2, u_1, u_2
- 2 If ds D in G , then $D' = D \cup \{d_1, d_2\}$ is sds in G'
- 3 Assume sds D' in G' . If $a_i \in D'$ (b_i), flip. $D = D' \setminus \{d_1, d_2\}$ is ds in G

Bipartite Graphs

SEMITOTAL DOMINATING SET on *bipartite* graphs is ω_2 -hard



Proof by fpt-reduction from PLANAR DOMINATING SET on bipart. graphs:

- 1 **Construct** Add new neighbor to each vertex and add d_1, d_2, u_1, u_2
- 2 If ds D in G , then $D' = D \cup \{d_1, d_2\}$ is sds in G'
- 3 Assume sds D' in G' . If $a_i \in D'$ (b_i), flip. $D = D' \setminus \{d_1, d_2\}$ is ds in G

A Linear Kernel for PLANAR SEMITOTAL DOMINATING SET

Another Explicit kernel for a Dominating Problem

Kernelization

Motivation

Theory

Intractability
 ω_2 hardness

Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References

- **Idea:** Preprocess an instance using *Reduction Rules* until hard *kernel* is found.



Kernelization

Motivation

Theory

Intractability
 ω_2 hardness


Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References

- **Idea:** Preprocess an instance using *Reduction Rules* until hard *kernel* is found.



Kernelization

Motivation

Theory

Intractability
 ω_2 hardness


Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References

- **Idea:** Preprocess an instance using *Reduction Rules* until hard *kernel* is found.



Related Works

Problem	Size	Source
PLANAR DOMINATING SET	$67k$	[16]
PLANAR TOTAL DOMINATING SET	$410k$	[20]
PLANAR SEMITOTAL DOMINATING SET	$359k$	This work
PLANAR EDGE DOMINATING SET	$14k$	[23]
PLANAR EFFICIENT DOMINATING SET	$84k$	[23]
PLANAR RED-BLUE DOMINATING SET	$43k$	[21]
PLANAR CONNECTED DOMINATING SET	$130k$	[34]
PLANAR DIRECTED DOMINATING SET	Linear	[1]

Main Theorem

Motivation

Theory

Intractability

ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

The Main Theorem

PLANAR SEMITOTAL DOMINATING SET parameterized by solution size admits a linear kernel. There exists a polynomial-time algorithm that, given a planar graph (G, k) , either correctly reports that (G, k) is a NO-instance or returns an equivalent instance (G', k) such that $|V(G')| \leq 359 \cdot k$.

The Big Picture

Motivation

Theory

Intractability

ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

Given a planar graph $G = (V, E)$, we will:

- 1 Split the neighborhoods of the graph;
- 2 Define reduction Rules
- 3 Use the region decomposition to analyze the size of each region

The Big Picture

Motivation

Theory

Intractability

ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

Given a planar graph $G = (V, E)$, we will:

- 1 Split the neighborhoods of the graph;
- 2 Define reduction Rules
- 3 Use the region decomposition to analyze the size of each region

The Big Picture

Motivation

Theory

Intractability

ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

Given a planar graph $G = (V, E)$, we will:

- 1 Split the neighborhoods of the graph;
- 2 Define reduction Rules
- 3 Use the region decomposition to analyze the size of each region

The Big Picture

Motivation

Theory

Intractability

ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

Given a planar graph $G = (V, E)$, we will:


- 1 Split the neighborhoods of the graph;
- 2 Define reduction Rules
- 3 Use the region decomposition to analyze the size of each region

The Basic Principle: Regions

Region (Simplified)

Given plane G and $v, w \in V$, a region is a closed subset, such that

- there are two non-crossing (but possibly overlapping) boundary paths
- Every vertex in R belongs to $N(v, w)$




The Basic Principle: Regions

Region (Simplified)

Given plane G and $v, w \in V$, a region is a closed subset, such that

- there are two non-crossing (but possibly overlapping) boundary paths
- Every vertex in R belongs to $N(v, w)$



Motivation

Theory

Intractability

ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions


References

The Basic Principle: Regions

Region (Simplified)

Given plane G and $v, w \in V$, a region is a closed subset, such that

- there are two non-crossing (but possibly overlapping) boundary paths
- Every vertex in R belongs to $N(v, w)$



D-region decomposition

Motivation

Theory

Intractability
 ω_2 hardness

Kernel

Definitions

Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References

D-region decomposition [2]

Given $G = (V, W)$ and $D \subseteq V$, a *D-region decomposition* is a set \mathfrak{R} with poles in D such that:

- for any vw -region $R \in \mathfrak{R}$: $D \cap V(R) = \{v, w\}$
- Regions are disjunct, but can share border vertices

A region is **maximal**, if no $R \in \mathfrak{R}$ such that $\mathfrak{R}' = \mathfrak{R} \cup \{R\}$ is a *D-region decomposition* with $V(\mathfrak{R}) \subsetneq V(\mathfrak{R}')$.

D-region decomposition

Motivation

Theory

Intractability
 ω_2 hardness

Kernel

Definitions

Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References

D-region decomposition [2]

Given $G = (V, W)$ and $D \subseteq V$, a *D-region decomposition* is a set \mathfrak{R} with poles in D such that:

- for any vw -region $R \in \mathfrak{R}$: $D \cap V(R) = \{v, w\}$
- Regions are disjunct, but can share border vertices

A region is **maximal**, if no $R \in \mathfrak{R}$ such that $\mathfrak{R}' = \mathfrak{R} \cup \{R\}$ is a *D-region decomposition* with $V(\mathfrak{R}) \subsetneq V(\mathfrak{R}')$.

Maximal D -region decomposition

Motivation

Theory


Intractability
 ω_2 hardness

Kernel


Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References



Splitting Up $N(v)$

 $N(v)$ 

Motivation

Theory

Intractability

 ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

Splitting Up $N(v)$

Motivation

Theory

Intractability
 ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

We split $N(v)$ into three subsets:

$$N_1(v) = \{u \in N(v) : N(u) \setminus N[v] \neq \emptyset\} \quad (1)$$

$$N_2(v) = \{u \in N(v) \setminus N_1(v) : N(u) \cap N_1(v) \neq \emptyset\} \quad (2)$$

$$N_3(v) = N(v) \setminus (N_1(v) \cup N_2(v)) \quad (3)$$

For $i, j \in [1, 3]$, we denote $N_{i,j}(v) := N_i(v) \cup N_j(v)$.

Splitting Up $N(v)$

Motivation

Theory

Intractability
 ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

We split $N(v)$ into three subsets:

$$N_1(v) = \{u \in N(v) : N(u) \setminus N[v] \neq \emptyset\} \quad (1)$$

$$N_2(v) = \{u \in N(v) \setminus N_1(v) : N(u) \cap N_1(v) \neq \emptyset\} \quad (2)$$

$$N_3(v) = N(v) \setminus (N_1(v) \cup N_2(v)) \quad (3)$$

For $i, j \in [1, 3]$, we denote $N_{i,j}(v) := N_i(v) \cup N_j(v)$.

Splitting Up $N(v)$

Motivation

Theory

Intractability
 ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

We split $N(v)$ into three subsets:

$$N_1(v) = \{u \in N(v) : N(u) \setminus N[v] \neq \emptyset\} \quad (1)$$

$$N_2(v) = \{u \in N(v) \setminus N_1(v) : N(u) \cap N_1(v) \neq \emptyset\} \quad (2)$$

$$N_3(v) = N(v) \setminus (N_1(v) \cup N_2(v)) \quad (3)$$

For $i, j \in [1, 3]$, we denote $N_{i,j}(v) := N_i(v) \cup N_j(v)$.

Splitting Up $N(v)$

Motivation

Theory

Intractability
 ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

We split $N(v)$ into three subsets:

$$N_1(v) = \{u \in N(v) : N(u) \setminus N[v] \neq \emptyset\} \quad (1)$$

$$N_2(v) = \{u \in N(v) \setminus N_1(v) : N(u) \cap N_1(v) \neq \emptyset\} \quad (2)$$


$$N_3(v) = N(v) \setminus (N_1(v) \cup N_2(v)) \quad (3)$$

For $i, j \in [1, 3]$, we denote $N_{i,j}(v) := N_i(v) \cup N_j(v)$.

Rule 1: Shrinking $N_3(v)$

Let $G = (V, E)$ be a graph and let $v \in V$. If $|N_3(v)| \geq 1$:

- remove $N_{2,3}(v)$ from G ,
- add a vertex v' and an edge $\{v, v'\}$.



- **Idea:** v better choice than $N_{2,3}$

Splitting up $N(v, w)$

Motivation

Theory


Intractability
 ω_2 hardness

Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References



Splitting up $N(v, w)$

Motivation

Theory

Intractability
 ω_2 hardness

Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References

$$N_1(v, w) = \{u \in N(v, w) \mid N(u) \setminus (N(v, w) \cup \{v, w\}) \neq \emptyset\} \quad (4)$$

$$N_2(v, w) = \{u \in N(v, w) \setminus N_1(v, w) \mid N(u) \cap N_1(v, w) \neq \emptyset\} \quad (5)$$

$$N_3(v, w) = N(v, w) \setminus (N_1(v, w) \cup N_2(v, w)) \quad (6)$$

For $i, j \in [1, 3]$, we denote $N_{i,j}(v, w) = N_i(v, w) \cup N_j(v, w)$.

Splitting up $N(v, w)$

Motivation

Theory

Intractability
 ω_2 hardness

Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References

$$N_1(v, w) = \{u \in N(v, w) \mid N(u) \setminus (N(v, w) \cup \{v, w\}) \neq \emptyset\} \quad (4)$$

$$N_2(v, w) = \{u \in N(v, w) \setminus N_1(v, w) \mid N(u) \cap N_1(v, w) \neq \emptyset\} \quad (5)$$

$$N_3(v, w) = N(v, w) \setminus (N_1(v, w) \cup N_2(v, w)) \quad (6)$$

For $i, j \in [1, 3]$, we denote $N_{i,j}(v, w) = N_i(v, w) \cup N_j(v, w)$.

Splitting up $N(v, w)$

Motivation

Theory

Intractability
 ω_2 hardness

Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References

$$N_1(v, w) = \{u \in N(v, w) \mid N(u) \setminus (N(v, w) \cup \{v, w\}) \neq \emptyset\} \quad (4)$$

$$N_2(v, w) = \{u \in N(v, w) \setminus N_1(v, w) \mid N(u) \cap N_1(v, w) \neq \emptyset\} \quad (5)$$

$$N_3(v, w) = N(v, w) \setminus (N_1(v, w) \cup N_2(v, w)) \quad (6)$$

For $i, j \in [1, 3]$, we denote $N_{i,j}(v, w) = N_i(v, w) \cup N_j(v, w)$.

Splitting up $N(v, w)$

Motivation

Theory

Intractability
 ω_2 hardness

Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References

$$N_1(v, w) = \{u \in N(v, w) \mid N(u) \setminus (N(v, w) \cup \{v, w\}) \neq \emptyset\} \quad (4)$$

$$N_2(v, w) = \{u \in N(v, w) \setminus N_1(v, w) \mid N(u) \cap N_1(v, w) \neq \emptyset\} \quad (5)$$

$$N_3(v, w) = N(v, w) \setminus (N_1(v, w) \cup N_2(v, w)) \quad (6)$$

For $i, j \in [1, 3]$, we denote $N_{i,j}(v, w) = N_i(v, w) \cup N_j(v, w)$.

Rule 2: Setting Up Our Weapons

Motivation

Theory

Intractability
 ω_3 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

Key Idea: $N_{2,3}(v, w)$ can **always** be semitotally dominated with 4 vertices.

$$\mathcal{D} = \{\tilde{D} \subseteq N_{2,3}(v, w) \mid N_3(v, w) \subseteq \bigcup_{v \in \tilde{D}} N(v), |\tilde{D}| \leq 3\} \quad (7)$$

$$\mathcal{D}_v = \{\tilde{D} \subseteq N_{2,3}(v, w) \cup \{v\} \mid N_3(v, w) \subseteq \bigcup_{v \in \tilde{D}} N(v), |\tilde{D}| \leq 3, v \in \tilde{D}\} \quad (8)$$

$$\mathcal{D}_w = \{\tilde{D} \subseteq N_{2,3}(v, w) \cup \{w\} \mid N_3(v, w) \subseteq \bigcup_{v \in \tilde{D}} N(v), |\tilde{D}| \leq 3, w \in \tilde{D}\} \quad (9)$$

Rule 2: Setting Up Our Weapons

Motivation

Theory

Intractability
 ω_3 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

Key Idea: $N_{2,3}(v, w)$ can **always** be semitotally dominated with 4 vertices.

$$\mathcal{D} = \{\tilde{D} \subseteq N_{2,3}(v, w) \mid N_3(v, w) \subseteq \bigcup_{v \in \tilde{D}} N(v), |\tilde{D}| \leq 3\} \quad (7)$$

$$\mathcal{D}_v = \{\tilde{D} \subseteq N_{2,3}(v, w) \cup \{v\} \mid N_3(v, w) \subseteq \bigcup_{v \in \tilde{D}} N(v), |\tilde{D}| \leq 3, v \in \tilde{D}\} \quad (8)$$

$$\mathcal{D}_w = \{\tilde{D} \subseteq N_{2,3}(v, w) \cup \{w\} \mid N_3(v, w) \subseteq \bigcup_{v \in \tilde{D}} N(v), |\tilde{D}| \leq 3, w \in \tilde{D}\} \quad (9)$$

Rule 2

If $\mathcal{D} = \emptyset$ we apply the following:

Case 1: if $\mathcal{D}_v = \emptyset$ and $\mathcal{D}_w = \emptyset$

- Remove $N_{2,3}(v, w)$
- Add vertices v' and w' and two edges $\{v, v'\}$ and $\{w, w'\}$
- Preserve $d(v, w)$

Case 2: if $\mathcal{D}_v \neq \emptyset$ and $\mathcal{D}_w = \emptyset$

- Remove $N_{2,3}(v)$
- Add $\{v, v'\}$

Case 3: if $\mathcal{D}_v = \emptyset$ and $\mathcal{D}_w \neq \emptyset$

Symmetric

Rule 2

If $\mathcal{D} = \emptyset$ we apply the following:

Case 1: if $\mathcal{D}_v = \emptyset$ and $\mathcal{D}_w = \emptyset$

- Remove $N_{2,3}(v, w)$
- Add vertices v' and w' and two edges $\{v, v'\}$ and $\{w, w'\}$
- Preserve $d(v, w)$

Case 2: if $\mathcal{D}_v \neq \emptyset$ and $\mathcal{D}_w = \emptyset$

- Remove $N_{2,3}(v)$
- Add $\{v, v'\}$

Case 3: if $\mathcal{D}_v = \emptyset$ and $\mathcal{D}_w \neq \emptyset$

Symmetric

Rule 2

If $\mathcal{D} = \emptyset$ we apply the following:

Case 1: if $\mathcal{D}_v = \emptyset$ and $\mathcal{D}_w = \emptyset$

- Remove $N_{2,3}(v, w)$
- Add vertices v' and w' and two edges $\{v, v'\}$ and $\{w, w'\}$
- Preserve $d(v, w)$

Case 2: if $\mathcal{D}_v \neq \emptyset$ and $\mathcal{D}_w = \emptyset$

- Remove $N_{2,3}(v)$
- Add $\{v, v'\}$

Case 3: if $\mathcal{D}_v = \emptyset$ and $\mathcal{D}_w \neq \emptyset$

Symmetric

Rule 2: Case 1



Motivation

Theory

Intractability
 ω_2 hardness

Kernel

Definitions

Rule 1


Rule 2

Rule 3

Kernel Size

Conclusions

References



Rule 2: Case 2

Motivation

Theory


Intractability
 ω_2 hardness

Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References




Simple Regions

The Main Theorem

A simple vw -region is a vw -region such that:

- 1 its boundary paths have length at most 2, and
- 2 $V(R) \setminus \{v, w\} \subseteq N(v) \cap N(w)$.




Simple Regions

The Main Theorem

A simple vw -region is a vw -region such that:

- 1 its boundary paths have length at most 2, and
- 2 $V(R) \setminus \{v, w\} \subseteq N(v) \cap N(w)$.



Rule 3: Shrinking the Size of Simple Regions

Motivation

Theory

Intractability
 ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

Let $G = (V, E)$ be a plane graph, $v, w \in V$ and R be a simple region between v and w . If $|V(R) \setminus \{v, w\}| \geq 5$ apply the following:

Case 1: If $G[R \setminus \partial R] \cong P_3$, then:

- remove $V(R \setminus \partial R)$
- add vertex y with edges $\{v, y\}$ and $\{y, w\}$

Case 2: If $G[R \setminus \partial R] \not\cong P_3$, then

- remove $V(R \setminus \partial R)$
- add vertices y, y' and four edges $\{v, y\}, \{v, y'\}, \{y, w\}$ and $\{y', w\}$


Rule 3: Shrinking the Size of Simple Regions

Case 1: If $G[R \setminus \partial R] \cong P_3$, then:

- remove $V(R \setminus \partial R)$
- add vertex y with edges $\{v, y\}$ and $\{y, w\}$

Case 2: If $G[R \setminus \partial R] \not\cong P_3$, then

- remove $V(R \setminus \partial R)$
- add vertices y, y' and four edges $\{v, y\}, \{v, y'\}, \{y, w\}$ and $\{y', w\}$




Rule 3: Shrinking the Size of Simple Regions

Case 1: If $G[R \setminus \partial R] \cong P_3$, then:

- remove $V(R \setminus \partial R)$
- add vertex y with edges $\{v, y\}$ and $\{y, w\}$

Case 2: If $G[R \setminus \partial R] \not\cong P_3$, then

- **remove** $V(R \setminus \partial R)$
- **add vertices** y, y' **and four edges** $\{v, y\}, \{v, y'\}, \{y, w\}$ **and** $\{y', w\}$



Notes

Motivation

Theory

Intractability

ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

- We proved that all these rules are sound,
- change the solution size by only a constant factor
- and can be applied in poly-time.

Notes

Motivation

Theory

Intractability

ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

- We proved that all these rules are sound,
- change the solution size by only a constant factor
- and can be applied in poly-time.

Notes

Motivation

Theory

Intractability

ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

- We proved that all these rules are sound,
- change the solution size by only a constant factor
- and can be applied in poly-time.

Notes

Motivation

Theory

Intractability

ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

- We proved that all these rules are sound,
- change the solution size by only a constant factor
- and can be applied in poly-time.

Bounding the Kernel: Vertices Outside any Region

Motivation

Theory


Intractability
 ω_2 hardness

Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References



For each d in $\text{sds } D$:

- ① $|N_1(v) \setminus V(\mathfrak{R})| = 0$ [2], On Border
- ② $|N_2(v) \setminus V(\mathfrak{R})| = 96$ [2]: TODO Reasoning
- ③ $|N_3(v) \setminus V(\mathfrak{R})| = 1$, by Rule 1

Bounding the Kernel: Vertices Outside any Region

Motivation

Theory


Intractability
 ω_2 hardness

Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References



For each d in sds D :

- 1 $|N_1(v) \setminus V(\mathfrak{R})| = 0$ [2], On Border
- 2 $|N_2(v) \setminus V(\mathfrak{R})| = 96$ [2]: TODO Reasoning
- 3 $|N_3(v) \setminus V(\mathfrak{R})| = 1$, by Rule 1

Bounding the Kernel: Vertices Outside any Region

Motivation

Theory


Intractability
 ω_2 hardness

Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References



For each d in $\text{sds } D$:

- 1 $|N_1(v) \setminus V(\mathfrak{R})| = 0$ [2], On Border
- 2 $|N_2(v) \setminus V(\mathfrak{R})| = 96$ [2]: TODO Reasoning
- 3 $|N_3(v) \setminus V(\mathfrak{R})| = 1$, by Rule 1

Bounding the Kernel: Vertices Outside any Region

Motivation

Theory


Intractability
 ω_2 hardness

Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References



For each d in $\text{sds } D$:

- 1 $|N_1(v) \setminus V(\mathfrak{R})| = 0$ [2], On Border
- 2 $|N_2(v) \setminus V(\mathfrak{R})| = 96$ [2]: TODO Reasoning
- 3 $|N_3(v) \setminus V(\mathfrak{R})| = 1$, by Rule 1

Bounding the Kernel: Vertices Outside any Region

Motivation

Theory


Intractability
 ω_2 hardness

Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions


References



For each d in $\text{sds } D$:

- 1 $|N_1(v) \setminus V(\mathfrak{R})| = 0$ [2], On Border
- 2 $|N_2(v) \setminus V(\mathfrak{R})| = 96$ [2]: TODO Reasoning
- 3 $|N_3(v) \setminus V(\mathfrak{R})| = 1$, by Rule 1

Bounding the Kernel: Inside a region



For each vw -region, we have

- ① $|N_1(v, w)| \leq 4$ [2] (vertices on border)
 - ② $|N_2(v, w)| \leq 6 \cdot 4$ (simple regions to $N_1(v, w)$, Rule 3)
 - ③ $|N_3(v, w)| \leq \max(27, 44, 4, 57) \cdot 4$ (proof omitted depending on Rule 2)
- Total:** $|V(R)| = |\{v, w\} \cup (N_1(v, w) \cup N_2(v, w) \cup N_3(v, w))| \leq 87$

Bounding the Kernel: Inside a region

Motivation

Theory


Intractability
 ω_2 hardness

Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions


References



For each vw -region, we have

- 1 $|N_1(v, w)| \leq 4$ [2] (vertices on border)
 - 2 $|N_2(v, w)| \leq 6 \cdot 4$ (simple regions to $N_1(v, w)$, Rule 3)
 - 3 $|N_3(v, w)| \leq \max(27, 44, 4, 57) \cdot 4$ (proof omitted depending on Rule 2)
- Total:** $|V(R)| = |\{v, w\} \cup (N_1(v, w) \cup N_2(v, w) \cup N_3(v, w))| \leq 87$

Bounding the Kernel: Inside a region



For each vw -region, we have

- 1 $|N_1(v, w)| \leq 4$ [2] (vertices on border)
 - 2 $|N_2(v, w)| \leq 6 \cdot 4$ (simple regions to $N_1(v, w)$, Rule 3)
 - 3 $|N_3(v, w)| \leq \max(27, 44, 4, 57) \cdot 4$ (proof omitted depending on Rule 2)
- Total:** $|V(R)| = |\{v, w\} \cup (N_1(v, w) \cup N_2(v, w) \cup N_3(v, w))| \leq 87$

Bounding the Kernel: Inside a region

Motivation

Theory

Intractability
 ω_2 hardness

Kernel

Definitions

Rule 1


Rule 2

Rule 3

Kernel Size

Conclusions

References



For each vw -region, we have

- 1 $|N_1(v, w)| \leq 4$ [2] (vertices on border)
 - 2 $|N_2(v, w)| \leq 6 \cdot 4$ (simple regions to $N_1(v, w)$, Rule 3)
 - 3 $|N_3(v, w)| \leq \max(27, 44, 4, 57) \cdot 4$ (proof omitted depending on Rule 2)
- Total:** $|V(R)| = |\{v, w\} \cup (N_1(v, w) \cup N_2(v, w) \cup N_3(v, w))| \leq 87$

Bounding the Kernel: Number of Regions

Motivation

Theory


Intractability
 ω_2 hardness

Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References



Number of Regions [2]

Let G be a plane graph and let D be a SEMITOTAL DOMINATING SET with $|D| \geq 3$. There is a maximal D -region decomposition of G such that $|\mathfrak{R}| \leq 3 \cdot |D| - 6$.

Bounding the Kernel: Number of Regions

Motivation

Theory


Intractability
 ω_2 hardness

Kernel

Definitions
Rule 1
Rule 2
Rule 3
Kernel Size

Conclusions

References



Number of Regions [2]

Let G be a plane graph and let D be a SEMITOTAL DOMINATING SET with $|D| \geq 3$. There is a maximal D -region decomposition of G such that $|\mathfrak{R}| \leq 3 \cdot |D| - 6$.

Summary: Bounding Kernel Size

Motivation

Theory

Intractability

ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

Let D be sds of size k . There exists a maximal D -region decomposition \mathfrak{R} such that:

- 1 \mathfrak{R} has only at most $3k - 6$ regions ([2]);
- 2 There are at most $97 \cdot k$ vertices outside of any region;
- 3 Each region $R \in \mathfrak{R}$ contains at most 87 vertices.

Hence: $|V| = \bigcup_{v \in D} N(v) = 87 \cdot (3k - 6) + 97 \cdot k + k < 359 \cdot k$

Main Theorem

Motivation

Theory

Intractability

ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

All reduction rules can be applied in poly/time, hence:

The Main Theorem

The SEMITOTAL DOMINATING SET problem parameterized by solution size admits a linear kernel on planar graphs. There exists a polynomial-time algorithm that, given a planar graph (G, k) , either correctly reports that (G, k) is a NO-instance or returns an equivalent instance (G', k) such that $|V(G')| \leq 359 \cdot k$.

Proof: Add Proof here.

Conclusions

Results:

- Given an overview over the status
- SEMITOTAL DOMINATING SET is $W[1]$ for *chordal*, *split* and *bipartite* graphs
- exists linear kernel of size $359 \cdot k$ when parameterized by solution size

Future Work:

- Improve kernel size and do empirical evaluation
- Solve parameterized complexities for *Circle*, *chordal bipartite* and *undirected path graphs*

Conclusions

Results:

- Given an overview over the status
- SEMITOTAL DOMINATING SET is $W[1]$ for *chordal*, *split* and *bipartite* graphs
- exists linear kernel of size $359 \cdot k$ when parameterized by solution size

Future Work:

- Improve kernel size and do empirical evaluation
- Solve parameterized complexities for *Circle*, *chordal bipartite* and *undirected path graphs*

Conclusions

Results:

- Given an overview over the status
- SEMITOTAL DOMINATING SET is $W[1]$ for *chordal*, *split* and *bipartite* graphs
- exists linear kernel of size $359 \cdot k$ when parameterized by solution size

Future Work:

- Improve kernel size and do empirical evaluation
- Solve parameterized complexities for *Circle*, *chordal bipartite* and *undirected path graphs*

Conclusions

Results:

- Given an overview over the status
- SEMITOTAL DOMINATING SET is $W[1]$ for *chordal*, *split* and *bipartite* graphs
- exists linear kernel of size $359 \cdot k$ when parameterized by solution size

Future Work:

- Improve kernel size and do empirical evaluation
- Solve parameterized complexities for *Circle*, *chordal bipartite* and *undirected path graphs*

Conclusions

Results:

- Given an overview over the status
- SEMITOTAL DOMINATING SET is $W[1]$ for *chordal*, *split* and *bipartite* graphs
- exists linear kernel of size $359 \cdot k$ when parameterized by solution size

Future Work:

- Improve kernel size and do empirical evaluation
- Solve parameterized complexities for *Circle*, *chordal bipartite* and *undirected path graphs*

Conclusions

Results:

- Given an overview over the status
- SEMITOTAL DOMINATING SET is $W[1]$ for *chordal*, *split* and *bipartite* graphs
- exists linear kernel of size $359 \cdot k$ when parameterized by solution size

Future Work:

- Improve kernel size and do empirical evaluation
- Solve parameterized complexities for *Circle*, *chordal bipartite* and *undirected path graphs*

Motivation

Theory

Intractability

ω_2 hardness

Kernel

Definitions

Rule 1

Rule 2

Rule 3

Kernel Size

Conclusions

References

? Any Questions ?
... And Thank You For Your Attention ...

References I



Jochen Alber, Britta Dorn, and Rolf Niedermeier. “A General Data Reduction Scheme for Domination in Graphs”. In: *SOFSEM 2006: Theory and Practice of Computer Science, 32nd Conference on Current Trends in Theory and Practice of Computer Science, Merin, Czech Republic, January 21-27, 2006, Proceedings*. Ed. by Jiri Wiedermann et al. Vol. 3831. Lecture Notes in Computer Science. Springer, 2006, pp. 137–147.



Jochen Alber, Michael R. Fellows, and Rolf Niedermeier. “Polynomial-time data reduction for dominating set”. In: (May 2004), pp. 363–384.



Rémy Belmonte and Martin Vatshelle. “Graph Classes with Structured Neighborhoods and Algorithmic Applications”. In: *Proceedings of the 37th International Conference on Graph-Theoretic Concepts in Computer Science. WG’11*. Teplá Monastery, Czech Republic: Springer-Verlag, 2011, pp. 47–58.



Alan A. Bertossi. “Dominating sets for split and bipartite graphs”. English. In: *Information Processing Letters* 19 (1984), pp. 37–40.



Alan A. Bertossi. “Total domination in interval graphs”. In: *Information Processing Letters* 23.3 (1986), pp. 131–134.

References II



Kellogg S. Booth and J. Howard Johnson. "Dominating Sets in Chordal Graphs". In: *SIAM J. Comput.* 11.1 (Feb. 1982), pp. 191–199.



Nicolas Bousquet et al. "Parameterized Domination in Circle Graphs". In: *Proceedings of the 38th International Conference on Graph-Theoretic Concepts in Computer Science*. WG'12. Jerusalem, Israel: Springer-Verlag, 2012, pp. 308–319.



Andreas Brandstädt, Victor D. Chepoi, and Feodor F. Dragan. "The Algorithmic Use of Hypertree Structure and Maximum Neighbourhood Orderings". In: *Discrete Appl. Math.* 82.1–3 (Mar. 1998), pp. 43–77.



Binh-Minh Bui-Xuan, Jan Arne Telle, and Martin Vatshelle. "Fast Dynamic Programming for Locally Checkable Vertex Subset and Vertex Partitioning Problems". In: *Theor. Comput. Sci.* 511 (Nov. 2013), pp. 66–76.



Gerard J Chang. "Total domination in block graphs". In: *Operations Research Letters* 8.1 (1989), pp. 53–57.



Gerard J. Chang. "Algorithmic Aspects of Domination in Graphs". In: *Handbook of Combinatorial Optimization: Volume 1–3*. Ed. by Ding-Zhu Du and Panos M. Pardalos. Boston, MA: Springer US, 1998, pp. 1811–1877.

References III



Maw-Shang Chang. “Efficient Algorithms for the Domination Problems on Interval and Circular-Arc Graphs”. In: *SIAM Journal on Computing* 27.6 (1998), pp. 1671–1694. eprint: <https://doi.org/10.1137/S0097539792238431>.



Bruno Courcelle. “The Monadic Second-Order Logic of Graphs. I. Recognizable Sets of Finite Graphs”. In: *Inf. Comput.* 85.1 (Mar. 1990), pp. 12–75.



Marek Cygan et al. “Dominating set is fixed parameter tractable in claw-free graphs”. In: *Theoretical Computer Science* 412.50 (2011), pp. 6982–7000.



Peter Damaschke, Haiko Müller, and Dieter Kratsch. “Domination in Convex and Chordal Bipartite Graphs”. In: *Inf. Process. Lett.* 36.5 (Dec. 1990), pp. 231–236.



Volker Diekert and Bruno Durand, eds. *STACS 2005, 22nd Annual Symposium on Theoretical Aspects of Computer Science, Stuttgart, Germany, February 24-26, 2005, Proceedings*. Vol. 3404. *Lecture Notes in Computer Science*. Springer, 2005.



Martin Farber. “Domination, independent domination, and duality in strongly chordal graphs”. In: *Discrete Applied Mathematics* 7.2 (1984), pp. 115–130.

References IV



Celina M. H. de Figueiredo et al. “Parameterized Algorithms for Steiner Tree and Dominating Set: Bounding the Leafage by the Vertex Leafage”. In: *WALCOM: Algorithms and Computation: 16th International Conference and Workshops, WALCOM 2022, Jember, Indonesia, March 24–26, 2022, Proceedings*. Jember, Indonesia: Springer-Verlag, 2022, pp. 251–262.



Esther Galby, Andrea Munaro, and Bernard Ries. “Semitotal Domination: New Hardness Results and a Polynomial-Time Algorithm for Graphs of Bounded Mim-Width”. In: *Theor. Comput. Sci.* 814.C (Apr. 2020), pp. 28–48.



Valentin Garnero and Ignasi Sau. “A Linear Kernel for Planar Total Dominating Set”. In: *Discrete Mathematics & Theoretical Computer Science* Vol. 20 no. 1 (May 2018). Sometimes we explicitly refer to the arXiv preprint version: <https://doi.org/10.48550/arXiv.1211.0978>. eprint: 1211.0978.



Valentin Garnero, Ignasi Sau, and Dimitrios M. Thilikos. “A linear kernel for planar red-blue dominating set”. In: *Discret. Appl. Math.* 217 (2017), pp. 536–547.



Archontia C. Giannopoulou and George B. Mertzios. “New Geometric Representations and Domination Problems on Tolerance and Multitolerance Graphs”. In: *SIAM Journal on Discrete Mathematics* 30.3 (2016), pp. 1685–1725. eprint: <https://doi.org/10.1137/15M1039468>.

References V



Jiong Guo and Rolf Niedermeier. “Linear Problem Kernels for NP-Hard Problems on Planar Graphs”. In: *Automata, Languages and Programming*. Ed. by Lars Arge et al. Berlin, Heidelberg: Springer Berlin Heidelberg, 2007, pp. 375–386.



Michael A. Henning, Saikat Pal, and D. Pradhan. “The semitotal domination problem in block graphs”. English. In: *Discussiones Mathematicae. Graph Theory* 42.1 (2022), pp. 231–248.



Michael A. Henning and Arti Pandey. “Algorithmic aspects of semitotal domination in graphs”. In: *Theoretical Computer Science* 766 (2019), pp. 46–57.



J. Mark Keil. “The Complexity of Domination Problems in Circle Graphs”. In: *Discrete Appl. Math.* 42.1 (Feb. 1993), pp. 51–63.



Ton Kloks and Arti Pandey. “Semitotal Domination on AT-Free Graphs and Circle Graphs”. In: *Algorithms and Discrete Applied Mathematics: 7th International Conference, CALDAM 2021, Rupnagar, India, February 11–13, 2021, Proceedings*. Rupnagar, India: Springer-Verlag, 2021, pp. 55–65.



D. V. Korobitsin. “On the complexity of domination number determination in monogenic classes of graphs”. In: 2.2 (1992), pp. 191–200.

References VI



Dieter Kratsch. “Domination and Total Domination on Asteroidal Triple-Free Graphs”. In: *Proceedings of the 5th Twente Workshop on on Graphs and Combinatorial Optimization*. Enschede, The Netherlands: Elsevier Science Publishers B. V., 2000, pp. 111–123.



Dieter Kratsch and Lorna Stewart. “Total domination and transformation”. In: *Information Processing Letters* 63.3 (1997), pp. 167–170.



James K. Lan and Gerard Jennhwa Chang. “On the algorithmic complexity of k-tuple total domination”. In: *Discrete Applied Mathematics* 174 (2014), pp. 81–91.



J. Pfaff; R. Laskar and S.T. Hedetniemi. *NP-completeness of Total and Connected Domination, and Irredundance for bipartite graphs*. Technical Report 428. Department of Mathematical Sciences: Clemson University, 1983.



Chunmei Liu and Yinglei Song. “Parameterized Complexity and Inapproximability of Dominating Set Problem in Chordal and near Chordal Graphs”. In: *J. Comb. Optim.* 22.4 (Nov. 2011), pp. 684–698.



Weizhong Luo et al. “Improved linear problem kernel for planar connected dominating set”. In: *Theor. Comput. Sci.* 511 (2013), pp. 2–12.

References VII



Alice Anne McRae. “Generalizing NP-Completeness Proofs for Bipartite Graphs and Chordal Graphs”. UMI Order No. GAX95-18192. PhD thesis. USA, 1995.



Haiko Müller and Andreas Brandstädt. “The NP-Completeness of Steiner Tree and Dominating Set for Chordal Bipartite Graphs”. In: *Theor. Comput. Sci.* 53.2 (June 1987), pp. 257–265.



R. Laskar;J. Pfaff. *Domination and irredundance in split graphs*. Technical Report 428. Department of Mathematical Sciences: Clemson University, 1983.



D. Pradhan and Saikat Pal. “An $O(n+m)$ time algorithm for computing a minimum semitotal dominating set in an interval graph”. In: *Journal of Applied Mathematics and Computing* 66.1 (June 2021), pp. 733–747.



Venkatesh Raman and Saket Saurabh. “Short Cycles Make W-hard Problems Hard: FPT Algorithms for W-hard Problems in Graphs with no Short Cycles”. In: *Algorithmica* 52.2 (2008), pp. 203–225.



Vikash Tripathi, Arti Pandey, and Anil Maheshwari. *A linear-time algorithm for semitotal domination in strongly chordal graphs*. 2021.