



DEPARTMENT OF INFORMATICS

TECHNICAL UNIVERSITY MUNICH

Master Thesis

# **On the Parametrized Complexity of Semitotal Domination on Graph Classes**

Lukas Retschmeier







DEPARTMENT OF INFORMATICS

TECHNICAL UNIVERSITY MUNICH

Master Thesis

# **On the Parametrized Complexity of Semitotal Domination on Graph Classes**

Author:	Lukas Retschmeier
Supervisor:	Prof. Debarghya Ghoshdastidar (TUM)
Advisor:	Prof. Paloma T. Lima (ITU)
Submission Date:	September 17, 2022



I confirm that this master thesis is my own work and I have documented all sources and material used.

*København S*  
September 17, 2022

Lukas Retschmeier

## Acknowledgments



---

## CONTENTS

<b>Acknowledgments</b>	<b>v</b>
<b>Abstract</b>	<b>ix</b>
<b>1 Introduction</b>	<b>2</b>
1.1 Content of the thesis . . . . .	2
<b>2 Preliminaries</b>	<b>4</b>
2.1 Graph Theory . . . . .	4
2.2 Parametrized Complexity . . . . .	5
2.2.1 Fixed Parameter Tractability . . . . .	5
2.2.2 Kernelization . . . . .	5
<b>3 On Parametrized Semitotal Domination</b>	<b>6</b>
3.1 Semitotal Domination . . . . .	6
3.2 $w[i]$ -Intractibility . . . . .	6
3.2.1 Warm-Up: $W[2]$ -hard on General Graphs . . . . .	6
3.2.2 $W[2]$ -hard on Bipartite Graphs . . . . .	6
3.2.3 $W[2]$ -hard on Chordal Graphs . . . . .	8
3.2.4 $W[2]$ -hard on Split Graphs . . . . .	8
<b>4 A Linear Kernel for Planar Semitotal Domination</b>	<b>9</b>
4.1 The Main Idea . . . . .	10
4.2 Definitions . . . . .	11
4.2.1 Regions in Planar Graphs . . . . .	13
4.3 The Big Picture . . . . .	16
4.4 Reduction Rules . . . . .	16
4.4.1 Reduction Rule I: Getting Rid of unnecessary $N_3(v)$ vertices . .	16
4.4.2 Reduction Rule II: Shrinking the Size of a Region . . . . .	17
4.4.3 Reduction Rule III: Shrinking Simple Regions . . . . .	19
4.4.4 Computing Maximal Simple Regions between two vertices . . .	19
4.5 Bounding the Size of the Kernel . . . . .	20
4.5.1 Bounding the Size of a Region . . . . .	20
4.5.2 Number of Vertices outside the Decomposition . . . . .	22

*Contents*

4.5.3 Bounding the Number of Regions . . . . .	25
<b>5 Open Questions and Further Research</b>	<b>27</b>
<b>Bibliography</b>	<b>29</b>
<b>List of Figures</b>	<b>32</b>
<b>List of Tables</b>	<b>33</b>



---

## ABSTRACT

*Abstract all the way*





---

# CHAPTER 1

---

## INTRODUCTION

Parametrized Complexity emerging branch. Books about that

Semitotal domination introduced by

Idea: Lake with stones, and family of ducks of fixed size wants to occupy the lake so that no other clan tries to take it over. Rules: \* A duck can quack freeing up neighboring stones. \* Ducks don't like to be alone and want to quack together. So for every duck their must be another duck that is not further than two stones away. Q: Can our ducklings occupy the whole lake while not feeling lonely?

### 1.1 Content of the thesis

In this thesis we continue the systematic analysis of the SEMITOTAL DOMINATING SET problem by focusing on the parametrized complexity of the problem.

Although the problem already had a lot of attention regarding classical complexity (CITE), only a few results are currently known for the parametrized variant.

As far as we have seen, even the  $w$ -hardness of the general case has not been explicitly been proven in the literature.

In this thesis, we continue the journey towards a systematic analysis by stating some hardness results for specific graph classes for the problem.

**Our contributions** Our main contributions consist of first showing the  $w[2]$ -hardness of SEMITOTAL DOMINATING SET for XXXX graphs.

As the DOMINATING SET problem and the TOTAL DOMINATING SET problem both admit a linear kernel for planar graphs, it is interesting to analyse whether this result also holds for the SEMITOTAL DOMINATING SET problem which lies in between these two.

Having these kernels also for other variants like EDGE DOMINATING SET, EFFICIENT DOMINATING SET, CONNECTED DOMINATING SET, PLANAR RED-BLUE DOMINATING SET

lent us a great confidence that the result will also work for SEMITOTAL DOMINATING SET on planar graphs.

Following the approach from ... which already relies on the technique given in, we give some simple data reduction rules for SEMITOTAL DOMINATING SET on planar graphs leading to a linear kernel. More precisely, we are going to prove the following central theorem of this thesis:

With some modifications we were able to transfer the approach given by Garnero and Stau in [GS18] to the SEMITOTAL DOMINATING SET problem.

**Theorem 1.** *The SEMITOTAL DOMINATING SET problem parametrized by solution size admits a linear kernel on planar graphs. There exists a polynomial-time algorithm that given a planar graph  $(G, k)$ , either correctly reports that  $(G, k)$  is a NO-instance or returns an equivalent instance  $(G', k)$  such that  $|V(G')| \leq 561 \cdot k$ .*

DOMINATING SET problem and TOTAL DOMINATING SET problem, both already

---

## CHAPTER 2

---

### PRELIMINARIES

We start by recapping some basic notation in Graph Theory and Parametrized Complexity.

Continuing an intensive study of parametrized complexity of that problem.

#### 2.1 Graph Theory

We quickly state the following definitions given by [Die10, p. xxx].

**Definition 1** (Graph [Die10, p. 3]). A graph is a pair  $G = (V, E)$  of two sets where  $V$  denotes the vertices and  $E \subseteq V \times V$  the edges of the graph. A vertex  $v \in V$  is incident with an edge  $e \in E$  if  $v \in e$ . Two vertices  $x, y$  are adjacent, or neighbours, if  $\{x, y\} \in E$ .

**Definition 2** (Vertex Degrees). The degree  $d_G(v)$  (If  $G$  is clear, also  $d(v)$ ) of a vertex  $v$  is the number of neighbors of  $v$ . We call a vertex of degree 0 as isolated and one of degree 1 as a pendant.

**Definition 3** (Special Graph Notations [Die10, p. 27]). A simple Graph

A directed Graph is a graph

A Multi Graph

A Planar Graph

**Definition 4** (Closed and Open Neighborhoods [BR12]). Let  $G = (V, E)$  be a (non-empty) graph. The set of all neighbors of  $v$  is the open neighborhood of  $v$  and denoted by  $N(v)$ ; the set  $N[v] = N(v) \cup \{v\}$  is the closed neighborhood of  $v$  in  $G$ . When  $G$  needs to be made explicit, those open and closed neighborhoods are denoted by  $N_G(v)$  and  $N_G[v]$ .

**Definition 5** (Induced Subgraph). asd

**Definition 6** (Isomorphic Graph). asd

## Special Graph Classes

We call the class of graphs without any special restrictions “General Graphs”.

**Definition 7 (r-partite Graphs).** Let  $r \geq 2$  be an integer. A Graph  $G = (V, E)$  is called “r-partite” if  $V$  admits a partition into  $r$  classes such that every edge has its ends in different classes: Vertices in the same partition class must not be adjacent.

For the case  $r = 2$  we say that the  $G$  is “bipartite”

**Definition 8 (Chordal Graphs).**

**Definition 9 (Split Graphs).**

## 2.2 Parametrized Complexity

### 2.2.1 Fixed Parameter Tractability

Fixed Parameter Intractability: The W Hierarchy

### 2.2.2 Kernelization

---

## CHAPTER 3

---

### ON PARAMETRIZED SEMITOTAL DOMINATION

#### 3.1 Semitotal Domination

##### SEMITOTAL DOMINATING SET

For two dominating vertices  $d_1, d_2$  we say that they are witnesses for each other if  $d(d_1, d_2) \leq 2$

Definition, dominating number

##### Complexity Status of Semitotal Dominating Set

#### 3.2 $w[i]$ -Intractibility

Now some  $w[i]$  hard classes.

##### 3.2.1 Warm-Up: $W[2]$ -hard on General Graphs

As any bipartite graphs with bipartition can be split further into  $r$ -partite graphs this results also implies the  $w[1]$ -hardness of  $r$ -partite graphs

##### 3.2.2 $W[2]$ -hard on Bipartite Graphs

**Definition 10** (Bipartite Graph, [BM08, p.5]). A bipartite graph is a Graph  $G$  whose vertex set can be partitioned into two subsets  $X$  and  $Y$ , so that each edge has one end in  $X$  and one end in  $Y$ . Such a partition  $(X, Y)$  is called a bipartition of  $G$ .

**Theorem 2.** Semitotal Dominating Set is  $\omega[2]$  hard for bipartite Graphs

*Proof.* Given a bipartite Graph  $G = (\{X \cup Y\}, E)$ , we construct a bipartite Graph  $G'$  in the following way:






Figure 3.1: Constructing  $G'$  from a bipartite Graph  $G$  by duplicating the vertices and adding a dominating tail

1. For each vertex  $x_i \in X$  we add a new vertex  $x'_i$  and an edge  $(x_i, x'_i)$  in between.
2. For each vertex  $y_j \in Y$  we add a new vertex  $y'_j$  and an edge  $(y_j, y'_j)$  in between.
3. We add two  $P_1$ , namely  $(u_1, d_1)$  and  $(u_2, d_2)$ , and connect them with all  $(d_1, x'_i)$  and  $(d_2, y'_j)$  respectively.

**Observation:**  $G'$  is clearly bipartite as all  $y'_j$  and  $x'_i$  form again an Independent Set. Setting  $X' = X \cup \{u_2\} \cup \bigcup y'_i$  and  $Y' = Y \cup \{u_1\} \cup \bigcup x'_i$  form the partitions of bipartite  $G'$ .

**Corollary 1.**  $G$  has a Dominating Set of size  $k$  iff  $G$  has a Semitotal Dominating Set of size  $k' = k + 2$

$\Rightarrow$ : Assume there exists a Dominating Set  $D$  in  $G$  with size  $k$ .  $DS = D \cup \{d_1, d_2\}$  is a Semitotal Dominating Set in  $G'$  with size  $k' = k + 2$ , because  $d_1$  dominates  $u_1$  and all  $x'_i$ ;  $d_2$  dominates  $u_2$  and all  $y'_i$ . Hence, it is a Semitotal Dominating Set, because  $\forall v \in (D \cap X) : d(v, d_1) = 2$  and  $\forall v \in (D \cap Y) : d(v, d_2) = 2$

$\Leftarrow$ : On the contrary, assume any Semitotal Dominating Set  $SD$  in  $G'$  with size  $k'$ . WLOG we can assume that  $u_1, u_2 \notin DS$ .

Our construction forces  $d_1, d_2 \in DS$ . Because all  $x'_i$  are only important in dominating  $x_i$  ( $y'_i$  for  $y_i$  resp.) as  $d_1, d_2 \in DS$ . If  $x'_i \in DS$  we simply exchange it with  $x_i$  (for  $y'_i$  and  $y_i$  respectively) in our DS keeping the size of the dominating set.  $D = DS \setminus \{d_1, d_2\}$  give us a Dominating Set in  $G$  with size  $k = k' - 2$

As  $G'$  can be constructed in  $\mathcal{O}(n)$  and parameter  $k$  is only blown up by a constant, this reduction is a FPT reduction. As Dominating Set is  $w[2]$  hard for bipartite Graphs<sup>1</sup> so is Semitotal Dominating Set.  $\square$

<sup>1</sup>Citation needed!

### *3 On Parametrized Semitotal Domination*

#### **3.2.3 $W[2]$ -hard on Chordal Graphs**

#### **3.2.4 $W[2]$ -hard on Split Graphs**

---

## CHAPTER 4

---

### A LINEAR KERNEL FOR PLANAR SEMITOTAL DOMINATION

*The best way to explain it is to do it.*

Lewis Carroll, *Alice in Wonderland*

We are going to present a polynomial-time preprocessing procedure which gives a linear kernel for PLANAR SEMITOTAL DOMINATING SET parametrized by solution size. Based on the technique first introduced by Alber et al. ([AFN04]) in 2004, an abundance of similar results to other domination problems emerged which gave us the believe that we can also transfer these results to SEMITOTAL DOMINATING SET. Table 4.1 gives an overview about the status of kernels for the planar case on other domination problems. All of these results introduce reduction rules bounding the number of vertices inside so-called “regions” which can be obtained by a special decomposition of the graph.

In the following years, this approach bore fruits in other planar problems as well like CONNECTED VERTEX COVER ( $11/3k$  in [KPS13]), MAXIMUM TRIANGLE PACKING ( $624k$  in [Wan+11]) INDUCED MATCHING ( $40k$  in [Kan+11]) FULL-DEGREE SPANNING TREE (TODO in [GNW06]) FEEDBACK VERTEX SET ( $13k$  in [BK16]) and CYCLE PACKING ([Gar+19])

In the upcoming years, many results could generalize the approach to larger graph classes. Fomin and Thilikos started this journey by directly proofing in the same year that the initial reduction rules given by Alber et al. [AFN04] can also be used to obtain a linear kernel on graphs with bounded genus  $g$  ([FT04]). Alon and Gutner advanced in 2008 with showing that the problem has a linear kernel on  $K_{3,h}$ -topological-minor-free graph classes and a polynomial kernel for  $K_h$ -topological-minor-free graph classes ([Gut09]). In 2007 they extended this result to show that graphs of bounded degeneracy are FPT ([AG07]). Finally, in 2012 Philip et al. showed that even  $K_{i,j}$ -free graph classes admit a polynomial kernel for DOMINATING SET ([PRS12]). In an attempt to extend these ideas to other problems as well, Bodlaender et al. ([Bod+16]) proofed that all problems

Problem	Best Known Kernel	Source
DOMINATING SET	67k	[DD05] <sup>1</sup>
TOTAL DOMINATING SET	410k	[GS18] <sup>2</sup>
SEMITOTAL DOMINATING SET	TODO	This Work
EDGE DOMINATING SET	14k	[Arg+07, p. 375 - 386, Theorem 2]
EFFICIENT DOMINATING SET	84k	[Arg+07, p. 375 - 386, Theorem 4]
PLANAR RED-BLUE DOMINATING SET	43k	[GST17a]
CONNECTED DOMINATING SET	130k	[Luo+13]
DIRECTED DOMINATING SET	?	[ADN06]

---

There is also a masters thesis claiming a bound of 43k [Hal16], but a conference or journal version was not found.

Table 4.1: An overview about existing kernels for planar dominating set variants

expressible in counting monadic second-order logic who satisfy a coverability property admit a polynomial kernel on graphs of bounded genus  $g$ .

Although these results are very interesting from a theoretical point of view, the constants for the kernels obtained by these methods are so large that they are not of practical interest. The question is how such a kernel can explicitly and efficiently be constructed.

We are going to show that, with some slight modifications, the kernel described by Garnero and Stau ([GS14]) for PLANAR TOTAL DOMINATING SET can also be used for PLANAR SEMITOTAL DOMINATING SET giving us an explicitly constructed kernel with “reasonable” small constants.

## 4.1 The Main Idea

The main idea is to use the fact that a plane graph  $G = (V, E)$  and a set  $D$ ,  $G$  can be decomposed into at most  $(3 \cdot |D| - 6)$  so-called “regions” (Definition 17). If  $D$  is now a given SEMITOTAL DOMINATING SET of size  $k := |D|$ , we know that the number of regions will depend linearly on  $k$ . If we define *reduction rules* (Rules 1 to 3) that try to minimize the number of vertices in and around a region we can overall bound the size of  $G$  in  $k$ .

The reduction rules

Currently, our reduction rules do not rely on the decomposition itself, but rather consider the neighborhood of every pair of vertices.

[TODO SUM UP THE STRATEGY]

## 4.2 Definitions

We are now giving ....

First, we would like to define what we consider to be a *reduced* graph.

**Definition 11** (Reduced Graph [GS18, p. 13] and [GST17b]). *A Graph  $G$  is reduced under a set of rules if either none of these rules can be applied to  $G$  or the application of any of them creates a graph isomorphic to  $G$ .*

From an algorithmic point of view, all our given reduction rules are local and only concern the neighborhood of at most two vertices and replace them partially with gadgets of constant size. Now checking whether a graph after applying the rule has been changed can be trivially accomplished in constant time.

In our case, we say  $G$  is reduced if none of the Rules 1 to 3 are modifying  $G$  any more.

This differs from the definition usually given in literature where a graph  $G$  is *reduced* under a set of reduction rules, if none of them can be applied to  $G$  anymore (Compare e.g. [Fom+19]). The reason is that we are giving reduction rules (see Rule 1 or Rule 2) that could be applied *ad infinitum* sending us into an endless loop without ever changing  $G$  any more. Our definition guarantees termination in that case.

In order to achieve this, we will first split up the neighborhood of one vertice and a pair of vertices into three distinct subsets, give some nice properties about them, and then state the corresponding reduction rules.

In this section, we are giving some key definitions that are used in our reduction rules for obtaining the linear kernel. These are inspired by those given by Garnero and Stau (PLANAR TOTAL DOMINATING SET in [GS14] or PLANAR RED-BLUE DOMINATING SET in [GST17a]) and already relied on those given by Alber et al. in [AFN04] for PLANAR DOMINATING SET.

The idea is to split up the neighborhood of a single vertex and a pair of vertices into three (disjoint) subsets that make a statement about how strongly the neighborhood is connected with the rest of the graph.

**Definition 12.** *Let  $G = (V, E)$  be a graph and let  $v \in V$ . We denote by  $N(v) = \{u \in V : \{u, v\} \in E\}$  the neighborhood of  $v$ . We split  $N(v)$  into three subsets:*

$$N_1(v) = \{u \in N(v) : N(u) \setminus N[v] \neq \emptyset\} \quad (4.1)$$

$$N_2(v) = \{u \in N(v) \setminus N_1(v) : N(u) \cap N_1(v) \neq \emptyset\} \quad (4.2)$$

$$N_3(v) = N(v) \setminus (N_1(v) \cup N_2(v)) \quad (4.3)$$

*In order to enhance future readability, for  $i, j \in [1, 3]$ , we denote  $N_{i,j}(v) := N_i(v) \cup N_j(v)$ .*

Intuitively, these sets are classifying neighbors of  $v$  by how much they can interact with the rest of the graph and how much they are locally centered around  $v$ :




Figure 4.1: The neighborhood of a vertex split to  $N_1(v)$  (blue),  $N_2(v)$  (purple), and  $N_3(v)$  (green). Note that all purple vertices have at least one blue neighbor making setting them in-between the green and blue vertices.

$N_1(v)$  are all neighbors of  $v$  which have at least one adjacent vertex that is outside of  $N(v)$  and therefore connect  $v$  with the rest of the graph. They could possibly belong to a solution.

$N_2(v)$  are all neighbors of  $v$  that have at least one neighbor from  $N_1(v)$ . These vertices do not have any function as a dominating vertex and can be seen as a *buffer* bridging  $N_1(v)$ -vertices with those from  $N_3(v) \cup \{v\}$ . Furthermore, they are useless as witnesses, because either we can replace them by  $v$  (sharing the same neighborhood) or when being a witness for  $v$ , we replace it with one  $z \in N_1(v)$ .

Vertices from  $N_3(v)$  are unmitigated and sealed off from the rest of the graph. They are useless as dominating vertices: For all  $z \in N_3(v)$  it holds that  $N(z) \subseteq N(v)$  by definition and thus, we would always prefer  $v$  as a dominating vertex instead of  $z$ . Nevertheless, they can be important as a witness for  $v$  in the case that  $N_1(v) \cup N_2(v) = \emptyset$ . We are using this observation in Rule 1 where we shrink  $|N_3(v)| \leq 1$ .

In the following, we are going to further extend this definition to a pair of vertices. Using this, Rule 2 will later try to reduce the neighborhood of two vertices, and similar to 12, we can deduce some properties.

**Definition 13.** Let  $G = (V, E)$  be a graph and  $v, w \in V$ . We denote by  $N(v, w) = N(v) \cup N(w)$  the neighborhood of the pair  $v, w$ . We split  $N(v, w)$  into three subsets:

$$N_1(v, w) = \{u \in N(v, w) \mid N(u) \setminus (N(v, w) \cup \{v, w\}) \neq \emptyset\} \quad (4.4)$$

$$N_2(v, w) = \{u \in N(v, w) \setminus N_1(v, w) \mid N(u) \cap N_1(v, w) \neq \emptyset\} \quad (4.5)$$

$$N_3(v, w) = N(v, w) \setminus (N_1(v, w) \cup N_2(v, w)) \quad (4.6)$$

Again, for  $i, j \in [1, 3]$ , we denote  $N_{i,j}(v, w) = N_i(v, w) \cup N_j(v, w)$ .

Again,  $N_1(v, w)$  are those connected with the rest of the graph,  $N_2(v, w)$  are a *buffer* between  $N_3(v, w) \cup \{v, w\}$  and  $N_3(v, w)$  are those isolated from the rest of the graph, but can still be useful as a witness for  $v$  or  $w$ .




Figure 4.2: TODO

Note that for example a vertex  $z \in N_1(v) \not\Rightarrow z \in N_1(v, w)$ . Figure 4.3 gives an example, where  $z$  belongs to  $N_1(v)$ , but not to  $N_1(v, w)$ .




Figure 4.3: The vertex  $z$  is in  $N_1(v)$ , because there is an edge pointing outside of  $N(v)$  to  $w$ . Contrary, it is not in  $N_1(v, w)$ , but now belongs to  $N_3(v, w)$ , because we are considering the “shared” neighborhood

### 4.2.1 Regions in Planar Graphs

We will introduce a concept that leads towards a new perspective looking at planar graphs, regions

As it is possible to bound the number of total  $vw$ -regions in a planar graph, we can analyze the local impacts to these regions from our future reduction rules.

It might be interesting to note that the authors of [GST17b] have revised their original definitions to set the basis for a more formal analysis.

**Definition 14.** Two simple paths  $p_1, p_2$  in a plane graph  $G$  are confluent if:

1. they are vertex-disjoint

#### 4 A Linear Kernel for Planar Semitotal Domination

2. they are edge-disjoint and for every common vertex  $u$ , if  $v_i, w_i$  are the neighbors of  $u$  in  $p_i$ , for  $i \in [1, 2]$ , it holds that  $[v_1, w_1, v_2, w_2]$ , or
3. they are confluent after contracting common edges

**Definition 15.** Let  $G = (V, E)$  be a plane graph and let  $v, w \in V$  be two distinct vertices. A region  $R(v, w)$  (also denoted as  $vw$ -region) is a closed subset of the plane, such that:

1. the boundary of  $R$  is formed by two confluent simple  $vw$ -paths with length at most 3
2. every vertex in  $R$  belongs to  $N(v, w)$ , and
3. the complement of  $R$  in the plane is connected.

We denote by  $\partial R$  the boundary of  $R$  and by  $V(R)$  the set of vertices which lay (with the plane embedding) in  $R$ . Furthermore, we call  $|V(R)|$  the size of the region.

The poles of  $R$  are the vertices  $v$  and  $w$ . The boundary paths are the two  $vw$ -paths that form  $\partial R$

We now have all the definitions ready to formally define a decomposition technique for planar graphs:

**Definition 16.** Two regions  $R_1$  and  $R_2$  are non-crossing, if:

1.  $(R_1 \setminus \partial R_1) \cap R_2 = (R_2 \setminus \partial R_2) \cap R_1 = \emptyset$ , and
2. the boundary paths of  $R_1$  are pairwise confluent with the ones in  $R_2$

**Definition 17.** Given a plane graph  $G = (V, E)$  and  $D \subseteq V$ , a  $D$  – region Decomposition of  $G$  is a set  $\mathfrak{R}$  of regions with poles in  $D$  such that:

1. for any  $vw$ -region  $R \in \mathfrak{R}$ , it holds that  $D \cap V(R) = \{v, w\}$ , and
2. all regions are pairwise non-crossing.

We denote  $V(\mathfrak{R}) = \bigcup_{R \in \mathfrak{R}} V(R)$ .

A  $D$ -region decomposition is maximal if there is no region  $R \notin \mathfrak{R}$  such that  $\mathfrak{R}' = \mathfrak{R} \cup \{R\}$  is a  $D$ -region decomposition with  $V(\mathfrak{R}) \subsetneq V(\mathfrak{R}')$

Fig. 4.4 gives an example of how to decompose a graph into a maximal  $D$  – region decomposition with a given SEMITOTAL DOMINATING SET of size 3.

We are introducing a special subset of a region, a *simple region* where every vertex is a common neighbor of  $v$  and  $w$ . They will appear in many unexpected astonishing places and are an important tool to operate on small parts of a plane graph. The upcoming Rule 3 will bound the size of these *simple regions*.

**Definition 18.** A *simple  $vw$ -region* is a  $vw$ -region such that:






Figure 4.4: left: A maximal  $D$ -region decomposition  $\mathfrak{R}$ , where  $D = \{d_1, d_2, d_3\}$  form a SEMI-TOTAL DOMINATING SET. There are two regions between  $d_2$  and  $d_1$ , one region between  $d_1$  and  $d_3$  and one region between  $d_2$  and  $d_3$ . Observe that some neighbors of  $d_1$  are not part of any  $vw$ -region. For those, our reduction rules are going to take care of that and bound these number of vertices to obtain the kernel. Right: The corresponding underlying multigraph  $G_{\mathfrak{R}}$




Figure 4.5: A simple region with two vertices from  $N_1(v, w)$  setting the boundary, two vertices from  $N_2(v, w)$  and some vertices from  $N_3(v, w)$  in between

1. its boundary paths have length at most 2, and
2.  $V(R) \setminus \{v, w\} \subseteq N(v) \cap N(w)$ .

Fig. 4.5 shows an example of a simple region containing 9 vertices.

Later we will use properties of the underlying multigraph of a  $D$ -region decomposition. Refer to Fig. 4.4 for an example.

**Definition 19.** Let  $G = (V, E)$  be a plane graph, let  $D \subseteq V$  and let  $\mathfrak{R}$  be a  $D$ -region decomposition of  $G$ . The underlying multigraph  $G_{\mathfrak{R}} = (V_{\mathfrak{R}}, E_{\mathfrak{R}})$  of  $\mathfrak{R}$  is such that  $V_{\mathfrak{R}} = D$  and there is an edge  $\{v, w\} \in E_{\mathfrak{R}}$  for each  $vw$ -region  $R(v, w) \in \mathfrak{R}$

### 4.3 The Big Picture

TODO Text here




Figure 4.6: The plan for obtaining a linear kernel for PLANAR SEMITOTAL DOMINATING SET.  
 TODO: Check whether R1-3 apply for gray lemmas

### 4.4 Reduction Rules

Following the approach by [GS14], we are now stating reduction rules that after exhaustive application will expose a linear kernel.

#### 4.4.1 Reduction Rule I: Getting Rid of unnecessary $N_3(v)$ vertices

An exemplarly application of the rule is shown in figure 4.7

**Rule 1.** Let  $G = (V, E)$  be a graph and let  $v \in V$ . If  $|N_3(v)| \geq 1$ :




Figure 4.7: TODO

- remove  $N_3(v)$  from  $G$ ,
- add a vertex  $v'$  and an edge  $\{v, v'\}$

**Lemma 1.** Let  $G = (V, E)$  be a graph and let  $v \in V$ . If  $G'$  is the graph obtained by applying Rule 1 on  $V$ , then  $G$  has SDS of size  $k$  if and only if  $G'$  has one.

*Proof.* This will be the proof for this lemma X □



Note, that we need our definition of a reduced instance given in 4.2. If Rule 3 is being applied, it will still leave us with a vertex  $z \in N_3(v)$  allowing this rule to be applied again.

#### 4.4.2 Reduction Rule II: Shrinking the Size of a Region

Extending the approach for a linear kernel for DOMINATING SET proposed by Alber et al. in [AFN04], Garnero and Stau transferred these results in [GS18] to the TOTAL DOMINATING SET problem.

Their idea was to strengthen the reduction rules in such a way that the witness properties for total domination are being preserved.

Following their approach in one of the first versions of [GS14], we stating reduction rules that. Interestingly, the reduction rules given in the latest version of this paper was not directly be transferable to SEMITOTAL DOMINATING SET, but an older version giving slightly easier reduction rules could be adjusted to our problem.

which relies on the technique first introduced by Alber et al we try to reduce the neighborhood for two given vertices  $v$  and  $w$

Before we give the concrete reduction rule, we will define three sets

$$\mathcal{D} = \{\tilde{D} \subseteq N_{2,3}(v, w) \mid N_3(v, w) \subseteq \bigcup_{v \in \tilde{D}} N(v), |\tilde{D}| \leq 3\} \quad (4.7)$$

$$\mathcal{D}_v = \{\tilde{D} \subseteq N_{2,3}(v, w) \cup \{v\} \mid N_3(v, w) \subseteq \bigcup_{v \in \tilde{D}} N(v), |\tilde{D}| \leq 3, v \in \tilde{D}\} \quad (4.8)$$

$$\mathcal{D}_w = \{\tilde{D} \subseteq N_{2,3}(v, w) \cup \{w\} \mid N_3(v, w) \subseteq \bigcup_{v \in \tilde{D}} N(v), |\tilde{D}| \leq 3, w \in \tilde{D}\} \quad (4.9)$$

**Rule 2.** Let  $G = (V, E)$  be a graph and two distinct  $v, w \in V$ . If  $\mathcal{D} = \emptyset$  we apply the following:

**Case 1:** if  $\mathcal{D}_v = \emptyset$  and  $\mathcal{D}_w = \emptyset$

- Remove  $N_{2,3}(v, w)$
- Add vertices  $v'$  and  $w'$  and two edges  $\{v, v'\}$  and  $\{w, w'\}$
- If there was a common neighbor of  $v$  and  $w$  in  $N_{2,3}(v, w)$  add another vertex  $y$  and two connecting edges  $\{v, y\}$  and  $\{y, w\}$

**Case 2:** if  $\mathcal{D}_v \neq \emptyset$  and  $\mathcal{D}_w \neq \emptyset$

Do nothing<sup>3</sup>

**Case 3:** if  $\mathcal{D}_v \neq \emptyset$  and  $\mathcal{D}_w = \emptyset$

- Remove  $N_{2,3}(v) \cap N_3(v, w)$
- Add  $\{v, v'\}$

**Case 4:** if  $\mathcal{D}_v = \emptyset$  and  $\mathcal{D}_w \neq \emptyset$

This case is symmetrical to **Case 3**.

Before proofing Rule 2 we will deduce some *Facts* which are implied by the definitions above.

**Fact 1.** Let  $G = (V, E)$  be a graph, let  $v, w \in V$ , and let  $G'$  be the graph obtained by the application of Rule 2 on  $v, w$ . If  $\mathcal{D} = \emptyset$ , then  $G$  has a solution if and only if it has a solution containing at least one of the two vertices  $\{v, w\}$ .

*Proof.*

□

Now we are ready to proof the correctness of Rule 2

---

<sup>3</sup>Originally, reduce Simple Regions [STAU]

**Fact 2.** Let  $G = (V, E)$  be a graph, let  $v, w \in V$ , and let  $G'$  be the graph obtained by the application of Rule 2 on  $v, w$ . If  $\mathcal{D} = \emptyset$  and  $\mathcal{D}_v = \emptyset$  (resp.  $\mathcal{D}_w = \emptyset$ ) then  $G'$  has a solution if and only if it has a solution containing  $v$  (resp.  $w$ ).

*Proof.* □

**Lemma 2.** Let  $G = (V, E)$  be a plane graph,  $v, w \in V$  and  $G' = (V', E')$  be the graph obtained after application of Rule 2 on the pair  $\{v, w\}$ . Then  $G$  has SDS of size  $k$  if and only if  $G'$  has SDS of size  $k$ .

*Proof.* We will proof the claim by analysing the different cases separately. □

#### 4.4.3 Reduction Rule III: Shrinking Simple Regions

**Rule 3.** Let  $G = (V, E)$  be a plane graph,  $v, w \in V$  and  $R$  be a simple region between  $v$  and  $w$ . If  $|V(R) \setminus \{v, w\}| \geq 7$

- Remove  $N_3(v, w)$
- Add two vertices  $h_1$  and  $h_2$  and four edges  $\{v, h_1\}$ ,  $\{v, h_2\}$ ,  $\{w, h_1\}$  and  $\{w, h_2\}$

**Lemma 3 (Correctness of Rule 3).** Let  $G = (V, E)$  be a plane graph,  $v, w \in V$  and  $G' = (V', E')$  be the graph obtained after application of Rule 3 on the pair  $\{v, w\}$ . Then  $G$  has SDS of size  $k$  if and only if  $G'$  has SDS of size  $k$ .

The application of Rule 3 gives us a bound on the number of vertices inside a simple region.

**Corollary 2.** Let  $G = (V, E)$  be a graph,  $v, w \in V$  and  $R$  a simple region between  $v$  and  $w$ . If Rule 3 has been applied, this simple region has size at most 6.

*Proof.* Clearly, if  $|V(R) \setminus \{v, w\}| < 7$  then the rule would not have changed  $G$  and the size of the region would already be bounded by 6. Assuming  $|V(R) \setminus \{v, w\}| \geq 7$  we note that every simple region can have at most two distinct vertices from  $N_1(v, w)$  and two distinct ones from  $N_2(v, w)$  without breaking planarity. These vertices are not touched by the reduction. Adding the two vertices that are being added between  $v$  and  $w$  gives us the desired upper bound. □

#### 4.4.4 Computing Maximal Simple Regions between two vertices

For the sake of completeness, we state an algorithm how a maximal simple region between two vertices  $v, w \in V$  can be computed in time  $\mathcal{O}(d(v) + d(w))$ :




Figure 4.8: TO BE DONE

## 4.5 Bounding the Size of the Kernel

We are now putting all our pieces together in order to prove our main result: A linear bound on the kernel size. In order to do so, we distinguish between those vertices that are covered by a maximal *D-region decomposition* and those that are not. In both cases our reduction rules bound the number of vertices to a constant size which means the kernel size does only depend on the number of regions of these decomposition. Fig. 4.4 states that for any solution  $D$ , we only have a linear number of regions that cover the whole graph. In particular, we show that given a SEMITOTAL DOMINATING SET  $D$  of size  $k$ , there exist a maximal *D-region decomposition*  $\mathfrak{R}$  such that:

1.  $\mathfrak{R}$  has only at most  $3|D| - 6$  regions
2.  $V(\mathfrak{R})$  covers most vertices of  $V$ . There are at most  $144 \cdot |D|$  vertices outside of any region.
3. each region of  $\mathfrak{R}$  contains at most XX vertices

Combining these three parts will give us a linear kernel.

### 4.5.1 Bounding the Size of a Region

We start with a more fine-grained analysis of the impact of the different cases of Rule 2 on a  $vw$ -region. The main idea is to count the number of simple regions in the  $vw$ -region and then use the bound on the size of a simple region after Rule 3 was applied exhaustively and which was obtained in Corollary 2.

We start by giving

**Lemma 4.** *Given a plane Graph  $G = (V, E)$  and a  $vw$ -region  $R$   $|N_1(v, w) \cap V(R)| \leq 4$  and these vertices lay exactly on the boundary  $\partial R$  of  $R$ .*

*Proof.*

□

**Lemma 5.** *[GS18, See Fact 5] Given a reduced plane graph  $G = (V, E)$  and a region  $R(v, w)$ ,  $N_2(v, w) \cap V(R)$  can be covered by at most 6 simple regions.*

*Proof.* Let  $(v, u_1, u_2, w)$  and  $(v, u_3, u_4, w)$  be the two boundary paths of  $R(v, w)$ . (A shorter path would only lead to a smaller bound). By definition of  $N_2(v, w)$ , vertices from  $N_2(v, w) \cap V(R)$  are common neighbors of  $v$  and  $w$  and  $u_i, i \in [4]$ . By planarity, we can cover  $N_2(v, w) \cap V(R)$  with at most 6 simple regions. To see this, imagine the graph where edges denote all possible simple  $vw$ -regions (See fig. 4.9). There are at most 8 simple regions possible. but we have to remove at least two of them to maintain planarity.

Furthermore, assuming the graph to be reduced, any intermediate  $N_3(v, w)$  which could possibly separate multiple simple regions between  $v$  and  $u_i$  has been deleted by Rule 1 already.  $\square$




Figure 4.9: Bounding the maximum number of simple regions inside a region  $R(v, w)$ .  $N_2(v, w)$  is covered by 6 green (simple) regions. A dashed edge would also be an option, but would contradict planarity. Note that the gray vertex  $g$  was reduced by Rule 1 allowing the formation of exactly one simple region between  $v$  and  $u_1$

We continue by giving a constant bound on the number of simple regions that cover all  $N_3(v, w)$  vertices in a given region.

**Lemma 6.** Given a plane Graph  $G = (V, E)$  reduced under Rule 2 and a region  $R(v, w)$ , if  $\mathcal{D}_v \neq \emptyset$  (resp.  $\mathcal{D}_w \neq \emptyset$ ),  $N_3(v, w) \cap V(R)$  can be covered by:

1. 11 simple regions if  $\mathcal{D}_w \neq \emptyset$ ,
2. 14 simple regions if  $N_{2,3}(v) \cap N_3(v, w) = \emptyset$

Note, that the first case applies, when Case 2 & 3 of Rule 2 have been applied and the second one, when Case 4 of Rule 2 was applied.

*Proof.* We will just give some intuition, because the proof of Garnero and Stau in [GS14, Fact 6] does not use any special property exposed by the reduction rules. Figure (Add picture about figures) gives a visualization worst case scenarios to cover  $N_3(v, w) \cap V(R)$  with simple regions in the relevant cases.<sup>4</sup>

<sup>4</sup>Note: In a newer revision of their paper [GS18], Stau und Garnero removed this proof, because they changed Rule 2 and a more fine-grained analysis was made possible.

□

**Lemma 7 (#Vertices inside a Region after Rules 1 to 3).** *Let  $G = (V, E)$  be a plane graph reduced under Rules 1 to 3. Furthermore, let  $D$  be a SDS of  $G$  and let  $v, w \in D$ . Any  $vw$ -region  $R$  contains at most 139 vertices distinct from its poles.*

*Proof.* By Lemmas 4 and 5 and Corollary 2 to bound the number of vertices inside a simple region, we know that  $|N_1(v, w) \cap V(R)| \leq 4$  and  $|N_2(v, w) \cap V(R)| \leq 6 \cdot 7 = 42$ .

It is still remaining to bound for  $|N_3(v, w) \cap V(R)|$ , but gladly we have Rule 2, which took care about them! Fig. 4.10 shows worst case amount of simple regions the individual cases can have.

**Case 0:** If Rule 2 has **not** been applied

As  $\mathcal{D} \neq \emptyset$ , there exists a set  $\tilde{D} = \{d_1, d_2, d_3\} \in \mathcal{D}$  of at most three vertices dominating  $N_3(v, w)$ . We observe that vertices from  $|N_3(v, w) \cap V(R)|$  are common neighbors of either  $v$  or  $w$  (by the definition of a  $vw$ -region) and at least one vertex from  $\tilde{D}$ . Without violating planarity, we can span at most 6 simple regions. Using Corollary 2 and adding  $|\tilde{D}| = 3$ , we can conclude  $|N_3(v, w) \cap V(R)| \leq 6 \cdot 6 + 3 = 39$ .

**Case 1:** If Rule 2 Case 1 has been applied

In that case  $|N_2(v, w) \cap V(R)|$  was entirely removed and at  $|N_3(v, w) \cap V(R)|$  replaced by at most three vertices ( $v', w'$  and  $y$ ) added. Hence  $|N_3(v, w) \cap V(R)| \leq 3$ .

**Case 2:** If Rule 2 Case 2 has been applied

As we know that  $\mathcal{D}_v \neq \emptyset$  and  $\mathcal{D}_w \neq \emptyset$ , we can apply Lemma 6 and although Rule 2 has not changed the  $G$ , we can cover  $R$  with at most 11 simple regions giving as  $|N_3(v, w) \cap V(R)| \leq 11 \cdot 6 = 66$  vertices.

**Case 3:** If Rule 2 Case 3 (sym. 4) has been applied

We know that in this case  $N_{2,3}(v) \cap N_3(v, w)$  was entirely removed and replaced by a single possible witness. Using Lemma 6, we can cover  $(N_3(v, w) \setminus \{v'\}) \cap V(R)$  with (at most) 14 simple regions giving us  $||N_3(v, w) \cap V(R)|| \leq 14 \cdot 6 + 1 = 85$ .

All in all, as  $V(R) = \{v, w\} \cup (N_1(v, w) \cup N_2(v, w) \cup N_3(v, w)) \cap V(R)$  we get

$$V(R) \leq 2 + 4 + 42 + \max(39, 3, 66, 85) = 139$$

□

#### 4.5.2 Number of Vertices outside the Decomposition

We continue to bound the number of vertices that do not lay inside any region of a maximal  $D$ -region decomposition  $\mathfrak{R}$ , that is, we bound the size of  $V \setminus V(\mathfrak{R})$ . Rule 1 ensures that we only have a small amount of  $N_3(v)$ -pendants. We then try to cover the rest with as few simple regions as possible, because, by application of Rule 3, these simple regions are of constant size.







**Case 0:** Maximal 6 Simple Regions

**Case 1:** Exactly 3 vertices

**Case 2:** All Maximal 11 Simple Regions **STILL WRONG** **Case 3:** All Maximal 9 Simple Regions **STILL WRONG**


Figure 4.10: TODO

**Lemma 8.** [AFN04] (Deprecated) Every vertex in  $u \in V \setminus V(\mathfrak{R})$  is either in  $D$  or belongs to a set  $N_2(v) \cup N_3(v)$ .

The following lemma states that no vertices from a set  $N_1(v)$  will be outside of a maximal  $D$ -region decomposition.

**Lemma 9.** [AFN04, Lemma 6] Let  $G = (V, E)$  be a plane graph and  $\mathfrak{R}$  be a maximal  $D$ -region decomposition of a DS  $D$ . If  $u \in N_1(v)$  for some vertex  $v \in D$  then  $u \in V(\mathfrak{R})$

In the following, we define  $d_{G_{\mathfrak{R}}}(v) = |\{R(v, w) \in \mathfrak{R}, w \in D\}|$  to be the number of regions in  $\mathfrak{R}$  having  $v$  as a pole.

**Corollary 3.** Let  $G = (V, E)$  be a graph and  $D$  be a set. For any maximal  $D$ -region decomposition  $\mathfrak{R}$  on  $G$  it holds that  $\sum_{v \in D} d_{G_{\mathfrak{R}}}(v) = 2 \cdot |\mathfrak{R}|$ .

*Proof.* The proof follows directly from the handshake lemma applied to the underlying multigraph  $G_{\mathfrak{R}}$ .  $\square$

**Proposition 1 (#Vertices outside a Region).** *Let  $G = (V, E)$  be a plane graph reduced under Rules 1 and 2 and let  $D$  be a SDS of  $G$ . If  $G$  has a maximal  $D$ -region decomposition, then  $|V \setminus (V(\mathfrak{R}) \cup D)| \leq 144|D|$*

With slight modifications, the proof given in [GS14] will also work in our case. Note that although assuming the graph to be entirely reduced, the following proof only relies on Rules 1 and 3. The proof uses the observation that vertices from  $N_2(v)$  span simple regions between those from  $\{v\} \cup N_1(v)$ .

*Proof.* Again, we will follow the proof proposed by Alber et al. [AFN04, Proposition 2].

The proof does only rely on Rules 1 and 3 and we can use the number of vertices in a simple region we have proven in Corollary 2. In particular, we are going to proof that  $|V \setminus V(\mathfrak{R})| \leq 48 \cdot |\mathfrak{R}| + 2 \cdot |D|$ . Directly placing in Lemma 10 will give as the desired bound.

let  $\mathfrak{R}$  be a maximal  $D$ -region decomposition and let  $v \in D$ . Since  $D$  dominates all vertices from  $V$ , we can consider  $V$  as  $\bigcup_{v \in D} N(v)$  and thus, we only need to bound the sizes of  $N_1(v) \setminus V(\mathfrak{R})$ ,  $N_2(v) \setminus V(\mathfrak{R})$  and  $N_3(v) \setminus V(\mathfrak{R})$  separately. In the following, let  $v \in D$ :

**N<sub>3</sub>(v):** As we know that Rule 1 has been exhaustively applied, we trivially see that  $|N_3(v)| \leq 1$  and hence,

$$\left| \bigcup_{v \in D} N_3(v) \setminus V(\mathfrak{R}) \right| \leq |D|$$

**N<sub>2</sub>(v):** According to Garnero and Stau ([GS18, Proposition 2]), we know that  $N_2(v) \setminus V(\mathfrak{R})$  can be covered by at most  $4d_{G_{\mathfrak{R}}}(v)$  simple regions between  $v$  and some vertices from  $N_1(v)$  on the boundary of a region in  $\mathfrak{R}$ . Figure 4.11 gives some intuition.

Because  $G$  is reduced by assumption, we know by Corollary 2 that a simple region can only have at least 6 vertices distinct from its poles and hence,

$$\begin{aligned} \left| \bigcup_{v \in D} N_2(v) \setminus V(\mathfrak{R}) \right| &\leq 6 \sum_{v \in D} 4 \cdot d_{G_{\mathfrak{R}}}(v) \\ &= 24 \cdot \sum_{v \in D} d_{G_{\mathfrak{R}}}(v) \\ &\stackrel{\text{Cor. 4.5.2}}{\leq} 48|\mathfrak{R}| \end{aligned} \tag{4.10}$$

**N<sub>1</sub>(v):** By Lemma 9, we know that  $N_1(v) \subseteq V(\mathfrak{R})$  and hence,

$$\left| \bigcup_{v \in D} N_1(v) \setminus V(\mathfrak{R}) \right| = 0$$

Summing up these three upper bounds for each  $v \in D$  we obtain the result using the equation from Lemma 10:

$$\begin{aligned}
|V \setminus V(\mathfrak{R}) \cup D| &\leq 48 \cdot |\mathfrak{R}| + |D| && \text{(Lemma 10)} \\
&\leq 48 \cdot (3|D| - 6) + |D| \\
&\leq 144|D| + |D| \\
&= 145|D|
\end{aligned} \tag{4.11}$$

□




Figure 4.11: Bounding the number of  $N_2(v)$ -vertices around a dominating vertex  $v$  given a maximal  $D$ -region decomposition  $\mathfrak{R}$ .  $v$  is a pole of  $R_1, R_2, \dots, R_j$  and can span simple regions with the help of  $N_2(v)$ -vertices to at most two  $N_1(v)$ -vertices per  $R_i$ . Each region has at most four vertices in  $N_1(v, w) \subseteq N_1(v)$  on the boundary of  $R_j$ , but only at most two can be used for a simple region: Observe that trying to build a simple region between  $v$  and  $n_2$  in this example would contradict the maximality of  $\mathfrak{R}$ . Furthermore, the size of these simple regions is bounded after the application of Rule 3.

### 4.5.3 Bounding the Number of Regions

Alber et al. [AFN04, Proposition 1] gave a greedy algorithm to construct a maximal  $D$ -region decomposition for a DOMINATING SET. Building up on these results, Garnero and Stau gave decomposition procedures for both PLANAR RED-BLUE DOMINATING SET ([GST17a]) and TOTAL DOMINATING SET ([GS18]) relying on linear kernelization, because it states that given a DOMINATING SET  $D$ , we can decompose the graph into a linear number of regions.

The following lemma corresponds to [AFN04, Proposition 1 and Lemma 5]. Although the authors gave different reduction rules and require a *reduced* instance as an assumption for the following lemma, they do not use any specific properties exposed by these rules. As any SEMITOTAL DOMINATING SET is also a DOMINATING SET, we can safely apply it for our problem as well. For a more detailed and formal proof, one can also refer to [GS18, Proposition 1].

**Lemma 10.** *Let  $G$  be a reduced plane graph and let  $D$  be a SEMITOTAL DOMINATING SET with  $|D| \geq 3$ . There is a maximal  $D$ -region decomposition of  $G$  such that  $|R| \leq 3 \cdot |D| - 6$*

*Proof.* Follows directly from [AFN04, Proposition 1 and Lemma 5] □

**Lemma 11 (Running Time of Reduction Procedure).** *TODO Runsi in polynomial Time.*

*Proof.* □

By utilizing all the previous results, we are now finally ready to proof the Theorem 1:

**Theorem 1.** *The SEMITOTAL DOMINATING SET problem parametrized by solution size admits a linear kernel on planar graphs. There exists a polynomial-time algorithms that given a planar graph  $(G, k)$ , either correctly reports that  $(G, k)$  is a NO-instance or returns an equivalent instance  $(G', k)$  such that  $|V(G')| \leq 561 \cdot k$ .*

*Proof.* Let  $G = (V, E)$  be the plane input graph and  $G' = (V', E')$  be the graph obtained by the exhaustive application of the Rules 1 to 3. As none of our rules change the size of a possible solution  $D'$  in  $G'$ , we know by Lemma 1, Lemma 2 and Lemma 3 that  $G'$  has a SEMITOTAL DOMINATING SET of size  $k$  if and only if  $G$  has a SEMITOTAL DOMINATING SET of size  $k$ . In Lemma 11, we have proven that this preprocessing procedure runs in polynomial time.

Assume that  $G'$  admits a solution  $D'$ .

By taking the size of each region proven in Proposition 1, the total number of regions in a maximal  $D$ -region decomposition (Lemma 10) and the number of vertices that can lay outside of any region (Proposition 1), we obtain the following bound:

$$139 \cdot (3k - 6) + 145 \cdot k + k \leq 562 \cdot k \quad (4.12)$$

If  $|V(G')| > 562 \cdot k$   $G$  is a NO-instance and we replace  $G'$  by two single disconnected vertices (trivial NO-instance). Then the kernel is of the claimed size. □

---

## CHAPTER 5

---

### OPEN QUESTIONS AND FURTHER RESEARCH

\* Chordal Bipartite Graphs have a very interesting case. \* Improve the Kernel Bound

## 5 *Open Questions and Further Research*

---

## BIBLIOGRAPHY

- [ADN06] J. Alber, B. Dorn, and R. Niedermeier. “A General Data Reduction Scheme for Domination in Graphs.” In: *SOFSEM 2006: Theory and Practice of Computer Science, 32nd Conference on Current Trends in Theory and Practice of Computer Science, Merin, Czech Republic, January 21-27, 2006, Proceedings*. Ed. by J. Wiedermann, G. Tel, J. Pokorný, M. Bieliková, and J. Stuller. Vol. 3831. Lecture Notes in Computer Science. Springer, 2006, pp. 137–147. doi: [10.1007/11611257\\_11](https://doi.org/10.1007/11611257_11).
- [AFN04] J. Alber, M. R. Fellows, and R. Niedermeier. “Polynomial-time data reduction for dominating set.” In: (May 2004), pp. 363–384. doi: [10.1145/990308.990309](https://doi.org/10.1145/990308.990309).
- [AG07] N. Alon and S. Gutner. “Linear time algorithms for finding a dominating set of fixed size in degenerated graphs.” In: *Proceedings of the 13th annual international conference on Computing and Combinatorics. COCOON’07*. Berlin, Heidelberg: Springer-Verlag, July 2007, pp. 394–405. ISBN: 9783540735441.
- [Arg+07] L. Arge, C. Cachin, T. Jurdzinski, and A. Tarlecki, eds. *Automata, Languages and Programming, 34th International Colloquium, ICALP 2007, Wroclaw, Poland, July 9-13, 2007, Proceedings*. Vol. 4596. Lecture Notes in Computer Science. Springer, 2007. ISBN: 978-3-540-73419-2. doi: [10.1007/978-3-540-73420-8](https://doi.org/10.1007/978-3-540-73420-8).
- [BK16] M. Bonamy and Ł. Kowalik. “A 13k-kernel for planar feedback vertex set via region decomposition.” en. In: *Theoretical Computer Science* 645 (Sept. 2016), pp. 25–40. ISSN: 0304-3975. doi: [10.1016/j.tcs.2016.05.031](https://doi.org/10.1016/j.tcs.2016.05.031).
- [BM08] J. A. Bondy and U. S. R. Murty. *Graph Theory*. Graduate Texts in Mathematics. Springer, 2008. ISBN: 978-1-84628-970-5. doi: [10.1007/978-1-84628-970-5](https://doi.org/10.1007/978-1-84628-970-5).
- [Bod+16] H. L. Bodlaender, F. V. Fomin, D. Lokshantov, E. Penninkx, S. Saurabh, and D. M. Thilikos. “(Meta) Kernelization.” In: *J. ACM* 63.5 (2016), 44:1–44:69. doi: [10.1145/2973749](https://doi.org/10.1145/2973749).

## Bibliography

- [BR12] R. Balakrishnan and K. Ranganathan. *A textbook of graph theory*. English. 2nd ed. Universitext. New York, NY: Springer, 2012. ISBN: 978-1-4614-4528-9; 978-1-4614-4529-6. DOI: [10.1007/978-1-4614-4529-6](https://doi.org/10.1007/978-1-4614-4529-6).
- [DD05] V. Diekert and B. Durand, eds. *STACS 2005, 22nd Annual Symposium on Theoretical Aspects of Computer Science, Stuttgart, Germany, February 24-26, 2005, Proceedings*. Vol. 3404. Lecture Notes in Computer Science. Springer, 2005. ISBN: 3-540-24998-2. DOI: [10.1007/b106485](https://doi.org/10.1007/b106485).
- [Die10] R. Diestel. *Graph Theory*. Fourth. Vol. 173. Graduate Texts in Mathematics. Heidelberg; New York: Springer, 2010. ISBN: 9783642142789 3642142788 9783642142796 3642142796.
- [Fom+19] F. V. Fomin, D. Lokshtanov, S. Saurabh, and M. Zehavi. *Kernelization. Theory of parameterized preprocessing*. English. Cambridge: Cambridge University Press, 2019. ISBN: 978-1-107-05776-0; 978-1-107-41515-7. DOI: [10.1017/9781107415157](https://doi.org/10.1017/9781107415157).
- [FT04] F. V. Fomin and D. M. Thilikos. “Fast Parameterized Algorithms for Graphs on Surfaces: Linear Kernel and Exponential Speed-Up.” en. In: *Automata, Languages and Programming*. Ed. by J. Díaz, J. Karhumäki, A. Lepistö, and D. Sannella. Lecture Notes in Computer Science. Berlin, Heidelberg: Springer, 2004, pp. 581–592. ISBN: 9783540278368. DOI: [10.1007/978-3-540-27836-8\\_50](https://doi.org/10.1007/978-3-540-27836-8_50).
- [Gar+19] V. Garnero, C. Paul, I. Sau, and D. M. Thilikos. “Explicit Linear Kernels for Packing Problems.” en. In: *Algorithmica* 81.4 (Apr. 2019), pp. 1615–1656. ISSN: 1432-0541. DOI: [10.1007/s00453-018-0495-5](https://doi.org/10.1007/s00453-018-0495-5).
- [GNW06] J. Guo, R. Niedermeier, and S. Wernicke. “Fixed-Parameter Tractability Results for Full-Degree Spanning Tree and Its Dual.” en. In: *Parameterized and Exact Computation*. Ed. by H. L. Bodlaender and M. A. Langston. Lecture Notes in Computer Science. Berlin, Heidelberg: Springer, 2006, pp. 203–214. ISBN: 9783540391012. DOI: [10.1007/11847250\\_19](https://doi.org/10.1007/11847250_19).
- [GS14] V. Garnero and I. Sau. “A Linear Kernel for Planar Total Dominating Set.” In: *Discrete Mathematics & Theoretical Computer Science, Vol. 20 no. 1, Discrete Algorithms (May 16, 2018) dmtcs:4487* (Nov. 2014). First Revision of the Paper. DOI: [10.23638/DMTCS-20-1-14](https://doi.org/10.23638/DMTCS-20-1-14). arXiv: [1211.0978](https://arxiv.org/abs/1211.0978) [cs.DS].
- [GS18] V. Garnero and I. Sau. “A Linear Kernel for Planar Total Dominating Set.” In: *Discret. Math. Theor. Comput. Sci.* 20.1 (2018).
- [GST17a] V. Garnero, I. Sau, and D. M. Thilikos. “A linear kernel for planar red-blue dominating set.” In: *Discret. Appl. Math.* 217 (2017), pp. 536–547. DOI: [10.1016/j.dam.2016.09.045](https://doi.org/10.1016/j.dam.2016.09.045).



- [GST17b] V. Garnero, I. Sau, and D. M. Thilikos. “A linear kernel for planar red–blue dominating set.” In: 217 (2017), pp. 536–547. issn: 0166-218X. doi: [10.1016/j.dam.2016.09.045](https://doi.org/10.1016/j.dam.2016.09.045).
- [Gut09] S. Gutner. “Polynomial Kernels and Faster Algorithms for the Dominating Set Problem on Graphs with an Excluded Minor.” In: *Parameterized and Exact Computation: 4th International Workshop, IWPEC 2009, Copenhagen, Denmark, September 10-11, 2009, Revised Selected Papers*. Berlin, Heidelberg: Springer-Verlag, Dec. 2009, pp. 246–257. isbn: 9783642112683.
- [Hal16] J. T. Halseth. “A 43k Kernel for Planar Dominating Set using Computer-Aided Reduction Rule Discovery.” In: *University of Bergen, University Library* (Feb. 2016).
- [Kan+11] I. Kanj, M. J. Pelsmayer, M. Schaefer, and G. Xia. “On the induced matching problem.” en. In: *Journal of Computer and System Sciences* 77.6 (Nov. 2011), pp. 1058–1070. issn: 0022-0000. doi: [10.1016/j.jcss.2010.09.001](https://doi.org/10.1016/j.jcss.2010.09.001).
- [KPS13] Ł. Kowalik, M. Pilipczuk, and K. Suchan. “Towards optimal kernel for connected vertex cover in planar graphs.” en. In: *Discrete Applied Mathematics* 161.7 (May 2013), pp. 1154–1161. issn: 0166-218X. doi: [10.1016/j.dam.2012.12.001](https://doi.org/10.1016/j.dam.2012.12.001).
- [Luo+13] W. Luo, J. Wang, Q. Feng, J. Guo, and J. Chen. “Improved linear problem kernel for planar connected dominating set.” In: *Theor. Comput. Sci.* 511 (2013), pp. 2–12. doi: [10.1016/j.tcs.2013.06.011](https://doi.org/10.1016/j.tcs.2013.06.011).
- [PRS12] G. Philip, V. Raman, and S. Sikdar. “Polynomial kernels for dominating set in graphs of bounded degeneracy and beyond.” In: *ACM Transactions on Algorithms* 9.1 (Dec. 2012), 11:1–11:23. issn: 1549-6325. doi: [10.1145/2390176.2390187](https://doi.org/10.1145/2390176.2390187).
- [Wan+11] J. Wang, Y. Yang, J. Guo, and J. Chen. “Linear Problem Kernels for Planar Graph Problems with Small Distance Property.” en. In: *Mathematical Foundations of Computer Science 2011*. Ed. by F. Murlak and P. Sankowski. Lecture Notes in Computer Science. Berlin, Heidelberg: Springer, 2011, pp. 592–603. isbn: 9783642229930. doi: [10.1007/978-3-642-22993-0\\_53](https://doi.org/10.1007/978-3-642-22993-0_53).

---

## LIST OF FIGURES

3.1	Constructing $G'$ from a bipartite Graph $G$ by duplicating the vertices and adding a dominating tail . . . . .	7
4.1	The neighborhood of a Single Vertex $v$ . . . . .	12
4.2	$N_i(v, w)$ . . . . .	13
4.3	Counterexample . . . . .	13
4.4	Region Decomposition . . . . .	15
4.5	A Simple Region . . . . .	15
4.6	Structure of the Proof . . . . .	16
4.7	<i>TODO</i> . . . . .	17
4.8	Application of Rule 3 . . . . .	20
4.9	Bounding number of simple regions with $N_2(v, w)$ inside a $vw$ -region $R$ . . . . .	21
4.10	Bounding number of simple regions with inside a $vw$ -region $R$ . . . . .	23
4.11	Vertices from $N_2(v)$ laying outside . . . . .	25

---

## LIST OF TABLES

4.1	An overview about existing kernels for planar dominating set variants .	10
-----	---	----