

Master's Thesis Presentation

On the Parameterized Complexity of Semitotal Dominating Set On Graph Classes

Lukas Retschmeier

Informatik 7 - Theoretical Foundations of Artificial Intelligence
Faculty of Informatics
Technical University of Munich

February 28th, 2023

Creative Introduction



Motivation

Theory

Kernel

Definitions

Reduction Rules

References

Our Plan for Today



① Motivation

② Theory

③ Kernel

Definitions

Reduction Rules

Motivation

Motivation

Theory

Kernel

Definitions

Reduction Rules

References

DOMINATING SET

Input

Graph $G = (V, E)$, $k \in \mathbb{N}$

Question

Is there a set $D \subseteq V$ of size at most k such that $N[D] = V$?

- The domination number is the minimum cardinality of a ds of G , denotes as $\gamma(G)$
- **Observation:** In connected G every $v \in D$ has another $z \in D$ with $d(v, z) \leq 3$.

Motivation

Motivation

Theory

Kernel

Definitions

Reduction Rules

References

TOTAL DOMINATING SET

Input

Graph $G = (V, E)$, $k \in \mathbb{N}$

Question

Is there a set $D \subseteq V$ of size at most k such that for all $d_1 \in X$ exists $d_2 \in X \setminus \{d_1\}$ s.t. $d(d_1, d_2) \leq 1$?

- The total domination number is the minimum cardinality of a tds of G , denoted as $\gamma_t(G)$.

Motivation

Motivation

Theory

Kernel

Definitions

Reduction Rules

References

SEMITOTAL DOMINATING SET

Input

Graph $G = (V, E)$, $k \in \mathbb{N}$

Question

Is there a subset $D \subseteq V$ with $|D| \leq k$ such that $N[D] = V$ and for all $d_1 \in D$ there exists another $d_2 \in D$ such that $d(d_1, d_2) \leq 2$?

- The semitotal domination number is the minimum cardinality of a sds of G , denoted as $\gamma_{2t}(G)$.
- **Observation:** $\gamma(G) \leq \gamma_{2t}(G) \leq \gamma_t(G)$

Example: $\gamma(G) < \gamma_{2t}(G) < \gamma_t(G)$

Motivation


Theory

Kernel


- Definitions
- Reduction Rules

References


DOMINATING SET



SEMITOTAL DOMINATING SET



TOTAL DOMINATING SET



Parameterized Complexity

- Developed by Downey and Fellows
- **Idea:** Limit combinatorial explosion to some aspect of the problem
-

Fixed-Parameter Tractability

Motivation

Theory

Kernel

Definitions

Reduction Rules

References

Kernelization

Motivation

Theory

Kernel

Definitions

Reduction Rules


References

- **Idea:** Preprocess an instance using *Reduction Rules* until hard *kernel* is found.




Kernelization

- **Idea:** Preprocess an instance using *Reduction Rules* until hard *kernel* is found.



Complexity Status



A Linear Kernel for PLANAR SEMITOTAL DOMINATING SET

The main result of the thesis

Related Works



Problem	Size	Source
PLANAR DOMINATING SET	$67k$	Diekert and Durand 2005
PLANAR TOTAL DOMINATING SET	$410k$	Garnero and Sau 2018
PLANAR SEMITOTAL DOMINATING SET	$xxxxk$	This work
PLANAR EDGE DOMINATING SET	$14k$	Guo and Niedermeier 2007
PLANAR EFFICIENT DOMINATING SET	$84k$	Guo and Niedermeier 2007
PLANAR RED-BLUE DOMINATING SET	$43k$	Garnero, Sau, and Thilikos 2017
PLANAR CONNECTED DOMINATING SET	$130k$	Luo et al. 2013
PLANAR DIRECTED DOMINATING SET	Linear	Alber, Dorn, and Niedermeier 2006

Main Theorem



Motivation

Theory

Kernel

Definitions

Reduction Rules

References

Introducing Region Decompositions

Master's
Thesis
Presentation

Lukas
Retschmeier

Motivation

Theory

Kernel

Definitions

Reduction Rules

References

Splitting up $N(v)$

Master's
Thesis
Presentation

Lukas
Retschmeier

Motivation

Theory

Kernel

Definitions

Reduction Rules

References

Rule 1: Shrinking $N_3(v)$

Motivation

Theory

Kernel

Definitions

Reduction Rules

References

Rule 2



Motivation

Theory

Kernel

Definitions

Reduction Rules

References


Rule 3: Shrinking the size of simple regions



Bounding Kernel Size

Let D be sds of size k . There exists a maximal D -region decomposition \mathfrak{R} such that

- 1 \mathfrak{R} has only at most $3k - 6$ regions (Alber, Fellows, and Niedermeier 2004);
- 2 There are at most $97 \cdot k$ vertices outside of any region;
- 3 Each region $R \in \mathfrak{R}$ contains at most 87 vertices.



Hence: $87 \cdot (3k - 6) + 97 \cdot k + k < 359 \cdot k$

Future Work



Motivation

Theory

Kernel

Definitions

Reduction Rules

References

References I



Alber, Jochen, Britta Dorn, and Rolf Niedermeier (2006). “A General Data Reduction Scheme for Domination in Graphs”. In: *SOFSEM 2006: Theory and Practice of Computer Science, 32nd Conference on Current Trends in Theory and Practice of Computer Science, Merin, Czech Republic, January 21-27, 2006, Proceedings*. Ed. by Jiri Wiedermann et al. Vol. 3831. Lecture Notes in Computer Science. Springer, pp. 137–147. DOI: 10.1007/11611257_11. URL: https://doi.org/10.1007/11611257_11.



Alber, Jochen, Michael R. Fellows, and Rolf Niedermeier (May 2004). “Polynomial-time data reduction for dominating set”. In: pp. 363–384. DOI: 10.1145/990308.990309. URL: <https://doi.org/10.1145/990308.990309>.



Diekert, Volker and Bruno Durand, eds. (2005). *STACS 2005, 22nd Annual Symposium on Theoretical Aspects of Computer Science, Stuttgart, Germany*,

References II

February 24-26, 2005, *Proceedings*. Vol. 3404. Lecture Notes in Computer Science. Springer. ISBN: 3-540-24998-2. DOI: 10.1007/b106485.



Garnero, Valentin and Ignasi Sau (May 2018). “A Linear Kernel for Planar Total Dominating Set”. In: *Discrete Mathematics & Theoretical Computer Science* Vol. 20 no. 1. Sometimes we explicitly refer to the arXiv preprint version: <https://doi.org/10.48550/arXiv.1211.0978>. DOI: 10.23638/DMTCS-20-1-14. eprint: 1211.0978. URL: <https://dmtcs.episciences.org/4487>.



Garnero, Valentin, Ignasi Sau, and Dimitrios M. Thilikos (2017). “A linear kernel for planar red-blue dominating set”. In: *Discret. Appl. Math.* 217, pp. 536–547. DOI: 10.1016/j.dam.2016.09.045.



Guo, Jiong and Rolf Niedermeier (2007). “Linear Problem Kernels for NP-Hard Problems on Planar Graphs”. In: *Automata, Languages and Programming*.

References III

Ed. by Lars Arge et al. Berlin, Heidelberg: Springer Berlin Heidelberg, pp. 375–386. ISBN: 978-3-540-73420-8.



Luo, Weizhong et al. (2013). “Improved linear problem kernel for planar connected dominating set”. In: *Theor. Comput. Sci.* 511, pp. 2–12. DOI: 10.1016/j.tcs.2013.06.011.