



A model of macroalgal decomposition

Journal:	<i>Ecology</i>
Manuscript ID	Draft
Wiley - Manuscript type:	Statistical Innovation
Date Submitted by the Author:	n/a
Complete List of Authors:	Wright, Luka Seamus; UWA Oceans Institute, ; UWA School of Biological Sciences,
Abstract:	Macroalgae are currently the focus of blue carbon, biological carbon dioxide (CO₂) removal in the ocean. Carbon stored in seaweed forest biomass is inconsequential to CO₂ removal for climate change mitigation. However, carbon exported as seaweed detritus represents ~7.5% of anthropogenic CO₂ emissions annually. Macroalgal blue carbon hinges on the fate of this detritus, so a reliable model of its decomposition is key. Macroalgal decomposition has been neither consistently nor reliably modelled. Here I present the first model tailored to macroalgal decomposition. Informed by the theory of detrital photosynthesis, it treats detritus as an entity in limbo that can initially either grow or decompose. This is achieved by replacing the rate constant of the classic exponential decay model with a logistic function of time, mirroring detrital photosynthesis. The resulting model has three parameters that describe the initial relative growth rate, the half-life of detrital photosynthesis, and the final exponential decay rate which is comparable to the rate constant of the classic model. Using a Bayesian implementation in R with Stan, I show that the model performs well given empirical data from a variety of contexts. I anticipate that it will prove instrumental to resolving the debate around macroalgal CO₂ removal.

Ecology

Statistical Innovations

A model of macroalgal decomposition

Luka Seamus Wright^{1,2*}

¹Oceans Institute, University of Western Australia, Perth, Australia

²School of Biological Sciences, University of Western Australia, Perth, Australia

*luka.wright@research.uwa.edu.au, luka@wright.it, <https://orcid.org/0000-0002-1273-6256>

Open Research Statement

Data and code to reproduce this paper and apply my model to your own data are available at github.com/lukaseamus/limbodeco.

Keywords

Bayesian statistics, calculus, decay, decomposition, degradation, exponential model, macroalgae, Markov Chain Monte Carlo, mathematical ecology, ordinary differential equation, seaweed, Stan

Abstract

Macroalgae are currently the focus of blue carbon, biological carbon dioxide (CO₂) removal in the ocean. Carbon stored in seaweed forest biomass is inconsequential to CO₂ removal for climate change mitigation. However, carbon exported as seaweed detritus represents ~7.5% of anthropogenic CO₂ emissions annually. Macroalgal blue carbon hinges on the fate of this detritus, so a reliable model of its decomposition is key. Macroalgal decomposition has been neither consistently nor reliably modelled. Here I present the first model tailored to macroalgal decomposition. Informed by the theory of detrital photosynthesis, it treats detritus as an entity in limbo that can initially either grow or decompose. This is achieved by replacing the rate constant of the classic exponential decay model with a logistic function of time, mirroring detrital photosynthesis. The resulting model has three parameters that describe the initial relative growth rate, the half-life of detrital photosynthesis, and the final exponential decay rate which is comparable to the rate constant of the classic model. Using a Bayesian implementation in R with Stan, I show that the model performs well given empirical data from a variety of contexts. I anticipate that it will prove instrumental to resolving the debate around macroalgal CO₂ removal.

Introduction

In a seminal paper published in this journal 62 years ago, Olson (1963) popularised the ordinary differential equation of leaf litter decomposition

$$\frac{dm}{dt} = -km \tag{1}$$

which describes the rate of change in litter mass m proportional to itself, a relationship which is parameterised by the decay constant k . This reflects our ecological understanding that given a theoretically limitless pool of decomposers, decomposition is proportional to the detrital pool. The equation integrates to

$$\ln \frac{m(t)}{m_0} = -kt \tag{2}$$

and simplifies to the familiar exponential decay function

$$m(t) = m_0 e^{-kt} \tag{3}$$

where $m(t)$ is a function of the initial litter mass m_0 , the decay constant k and time t . This function has prevailed as the model of choice for decomposition.

Interest in decomposition has been rekindled due to the hope that plants can mitigate anthropogenic climate change by fixing carbon dioxide (CO₂), assimilating it into carbohydrates, and sequestering it in their biomass. Carbon sequestration timescales of decades to centuries are being considered the minimum and ten millennia or longer are preferred (Babiker et al., 2022). Plant biomass can be split into two distinct carbon sinks:

living plants and the detrital pool, of which the latter is considered the more promising. Often surprising to plant ecologists, plant detritus amounts to a biomass that is generally comparable and often far greater than that of living plants (Cebrián & Duarte, 1995, Wright & Kregting, 2023). But more importantly, living plants, even when considering stable or increasing populations, are limited by biomass-density constraints (Creed et al., 2019) and lifespans of typically less than a millennium (Thomas, 2013). Detritus on the other hand is believed to have the potential of cumulative sequestration over much longer timescales. Fossil fuels were created via the Carboniferous detrital pathway, mostly of scale trees (Hibbett et al., 2016), and the hope of biological CO₂ removal essentially boils down to re-enacting fossil fuel generation.

Biological CO₂ removal in the ocean, which amounts to somewhat less than on land (Friedlingstein et al., 2025), is referred to by the shorthand blue carbon (Babiker et al., 2022). Macroalgae have recently moved into the focus of blue carbon (Pessarrodona et al., 2023), primarily due to their substantial global area (Duarte et al., 2022), biomass packing (Creed et al., 2019), productivity (Pessarrodona et al., 2022) and detrital export (Pessarrodona et al., 2023, Filbee-Dexter et al., 2024). Notably, decomposition is disproportionately consequential for macroalgal blue carbon. Unlike their fellow blue carbon ecosystems (seagrass meadows, mangrove forests and salt marshes), seaweed forests cannot accumulate carbon in below-ground biomass and soil. There is thus a set limit for the living seaweed carbon sink which can be estimated as the product of 6.1 million km² global cover (Duarte et al., 2022), 1.2 ± 3 kg m⁻² (mean $\pm$ standard deviation) standing biomass (Creed et al., 2019) and 28 ± 4.5 % carbon content (Pessarrodona et al., 2022). This yields 0.34 ± 0.87 kg C m⁻² and 2.1 ± 5.3 Gt C globally, or only 19 ± 49 % of annual anthropogenic CO₂ emissions (Friedlingstein et al., 2025). However, seaweeds are thought to assimilate 0.32 ± 0.54 kg C m⁻² y⁻¹ (Pessarrodona

et al., 2022) and 1.3 Gt C y^{-1} globally (Duarte et al., 2022) of which $61 \pm 4.2 \%$ is exported as detritus (Pessarrodona et al., 2023). Hence $0.19 \pm 0.33 \text{ kg C m}^{-2} \text{ y}^{-1}$ and $0.81 \pm 0.055^* \text{ Gt C y}^{-1}$ globally, or $7.5 \pm 0.51^* \%$ of annual anthropogenic CO_2 emissions, may enter the detrital pool yearly. Viewed cumulatively this is clearly the more substantial carbon sink. Decomposition alone therefore determines whether macroalgal blue carbon is significant or not, as is illustrated by the central role it takes in the latest macroalgal blue carbon models (Filbee-Dexter et al., 2024). Despite this, macroalgal decomposition remains comparatively poorly understood.

Phycologists have never been able to properly model macroalgal decomposition. Decomposition trajectories rarely match Equation 3 reasonably well (e.g. de Bettignies et al., 2020, Pedersen et al., 2021). Most studies have therefore either refrained from fitting a model (e.g. Birch et al., 1983, Hamersley et al., 2015) or most commonly fit linear models (e.g. Bedford & Moore, 1984, Brouwer, 1996, Frontier et al., 2021, Filbee-Dexter et al., 2022, Frontier et al., 2022, Wright et al., 2022). Meta-analyses of macroalgal k (Equation 3) are numerous (Enríquez et al., 1993, Banta et al., 2004, Cebrián & Lartigue, 2004, Filbee-Dexter et al., 2024, Kennedy & Blain, 2025, White & Norkko, 2025) but have generally assumed Equation 3 by default and not assessed the validity of this model in the macroalgal context. It is evident that these meta-analyses are often hampered by the complexity and diversity of macroalgal decomposition trajectories (Kennedy & Blain, 2025, White & Norkko, 2025). The only serious attempts at modelling diverse trajectories have invariably invoked the logistic function alongside the exponential (Bourguès et al., 1996, Kennedy & Blain, 2025). This

* This standard deviation is an underestimate because no uncertainty was provided for the global primary production estimate (Duarte et al., 2022).

defeats the purpose of a model since comparing logistic and exponential rate constants is as effective as squinting at the data. Kennedy & Blain (2025) realised this when trying to compare half-lives ($\frac{\ln 2}{k}$ for Equation 3) across exponential and logistic models and ultimately resorted to forcing an exponential model to derive this parameter, which undermines fitting the logistic model in the first place.

We now know what causes the observed diversity in macroalgal decomposition trajectories. Early phycologists noticed the need to “pre-kill” macroalgal detritus (Birch et al., 1983, Smith & Foreman, 1984, Rieper-Kirchner, 1989, Brouwer, 1996) if it was to behave according to the expectations of the classic model (Equation 3). Recently it became evident that this is because excised macroalgal tissue can continue to photosynthesise for months (de Bettignies et al., 2020), which may allow macroalgal detritus *sensu lato*[†] to grow rather than decompose over the short term (Frontier et al., 2021, Pedersen et al., 2021, Filbee-Dexter et al., 2022, Frontier et al., 2022, Wright et al., 2022). There is now extensive investigation into what has since been termed detrital photosynthesis (Frontier et al., 2021, Wright & Foggo, 2021, Frontier et al., 2022, Wright et al., 2022, Wright & Kregting, 2023, Wright et al., 2024) and macroalgae have been found to be unique among plants in having detritus *sensu lato* that photosynthesises to an extent that is likely to influence its decomposition (Wright, in review). This knowledge enables the creation of a model that accurately describes macroalgal decomposition by taking into account the effect of detrital

[†] I have repeatedly faced semantic challenges when communicating this phenomenon since detritus has traditionally been used only to refer to dead matter. To avoid confusion, I have suggested detritus *sensu lato* for all detritus and detritus *sensu stricto* for dead detritus (Wright, in review). I use this terminology hereinafter.

photosynthesis. It is probable that, ignorant of detrital photosynthesis, previous models have substantially underestimated macroalgal decomposition with obvious consequences for blue carbon accounting. To remedy this I here present a model with only three parameters that is tailored to the unique case of macroalgal decomposition. First I describe its mathematical form, then I implement it as a Bayesian statistical model in R (R Core Team, 2022) with Stan (Carpenter et al., 2017), and finally I validate the model against a variety of empirical examples.

The mathematical model

...tossed and swept like dead leaves from one spot to another, never resting, never giving back their goodness to the earth, never fully dead but in some vast limbo between life and extinction, tossing and tumbling without end... — Philip Pullman (1994) *The Tin Princess*

Macroalgal detritus *sensu lato* is essentially in limbo. It may grow or decay, but not indefinitely. The goal is to model this temporary limbo. There are several possible approaches to describing this mathematically. As mentioned above, the logistic function has been the only attempt so far, but its rate constant is not comparable to that of Equation 3 and it cannot describe the full spectrum of possible decomposition trajectories (e.g. an increase followed by a decrease). Hence we must dismiss it. Another approach could be a composite function of a saturating function (e.g. rectangular hyperbola, hyperbolic tangent, exponential) and Equation 3. This may describe all possible trajectories but is somewhat of an artificial construct and more importantly would have more parameters than is practical. The best way forward is to stick with our trusty Equation 3 but rebuild it from its differential form (Equation 1). Instead of treating k as constant, we can let it be a function of time:

$$\frac{dm}{dt} = k(t)m \quad (4)$$

168

169 To my knowledge this approach has only been applied twice before, by Rovira & Rovira
 170 (2010) to describe the variety of terrestrial leaf litter decomposition trajectories and by
 171 Manojlovic (2021) to model seagrass decomposition (Trevathan-Tackett et al., 2020). Rovira
 172 & Rovira (2010) chose the form of $k(t)$ based on context while Manojlovic (2021) used a
 173 random walk, the simplest autoregression which defines k only in terms of k at the previous
 174 timepoint plus normal error:

175

$$k(t) \sim N(k(t - \Delta t), \sigma) \quad (5)$$

177

178 where σ is the standard deviation of the normal distribution around the mean $k(t - \Delta t)$ and
 179 determines the k step size of the random walk. The timestep size Δt is typically set to 1.
 180 While this approach is the most flexible, it requires sophisticated Bayesian state space
 181 modelling with an ordinary differential equation solver. Manojlovic (2021) used LibBi
 182 (Murray, 2015) and to my knowledge there is currently no way to implement this in the more
 183 popular Stan language (Manojlovic, personal communication, [discourse.mc-](https://discourse.mc-stan.org/t/random-walk-in-rstan/34308)
 184 [stan.org/t/random-walk-in-rstan/34308](https://discourse.mc-stan.org/t/random-walk-in-rstan/34308)). Furthermore, being the simplest type of order 1
 185 autoregressive process, a random walk's only parameter is σ which has no inherent
 186 directionality and precludes informative prediction beyond the range of the data. Finally, no
 187 comparison can be made between studies beyond overlaying plots of $k(t)$, and formal
 188 comparison with studies that use the traditional model (Equation 3) is out of question. A
 189 model with informative parameters which can be formally compared is the way forward.
 190 Hence I here opt for the path laid out by Rovira & Rovira (2010).

191

Now the most important task is to find an appropriate function to describe k . Since the underlying process affecting k is detrital photosynthesis, it makes sense to start there. The change in detrital photosynthesis with time post excision, or detrital age, has mostly been described using various linear models (de Bettignies et al., 2020, Frontier et al., 2021, Wright & Foggo, 2021, Frontier et al., 2022, Wright et al., 2022, Wright & Kregting, 2023, Wright et al., 2024). In truth there is no reasonable mechanism for photosynthesis to increase post excision[‡], but it is often maintained near maximal levels until a breakpoint is reached and it decays to a minimum beyond which there logically is no further decline. Voilà, logistic decay! In Wright (in review) I have described detrital photosynthesis (p) as

$$p(t) = \frac{\alpha + \tau}{1 + e^{r(t - \mu)}} - \tau \tag{6}$$

where α and τ may be read as the maximal CO₂ assimilation and production of the plant and its detritus respectively, r is the logistic rate of decay and μ is the time of photosynthetic death, more specifically the half-life of photosynthesis. The naming of parameters is inspired by the word מא (graecised $\tau\mu\alpha$), meaning truth. Composed of the first, middle and last letters of the Hebrew alphabet, it symbolises life ($\aleph$), transition (μ) and death (τ). In legend this word lends life to the golem, but removing $\aleph$ changes its meaning to death (מט) and the golem dies. As in the story of the golem, the logistic function is optimal for modelling a shift between two alternate states and is thus the logical choice here.

[‡] We sometimes see this due to some unrelated factor such as a change in light environment (Frontier et al., 2021, Wright & Kregting, 2023).

213 In practice, I found the full four-parameter form of Equation 6 difficult to work with because
 214 r and μ are not identifiable and the model affords little control over the intercept, for which
 215 we known $p(0) \approx \alpha$ to be true. This usually requires a workaround, such as parameterising the
 216 model in terms of the logistic intercept ($r\mu$) or constraining the model with a joint prior on $\ln r$
 217 $+ \ln \mu$ (Wright, in review). There are two ways to simplify Equation 6 to a three-parameter
 218 form: (1) set r constant and (2) describe r relative to μ , which is effectively the same as
 219 setting $r\mu$ constant. Reasonable values are $r = 1$ for option 1, which is roughly equivalent to
 220 photosynthetic death over one week, and $r = \frac{5}{\mu}$ for option 2 which implies that $r\mu = 5$, so $p(0)$
 221 $= 0.99(\alpha + \tau) - \tau \approx \alpha$. Theory would favour option 2 because we cannot assume a universal
 222 logistic rate of decay for detrital photosynthesis but we know $p(0) \approx \alpha$. Nonetheless, option 1
 223 allows rapid photosynthetic decay independent of μ while option 2 implies that if detrital
 224 photosynthesis persists it must be due to slow decay initiated at $t = 0$. Accordingly the
 225 interpretation of α changes from detrital photosynthesis until photosynthetic death in option 1
 226 to initial photosynthesis in option 2. While we know that the ultimate cause of photosynthetic
 227 decay is that initiating event (e.g. excision) and this agrees well with detrital photosynthesis
 228 trajectories (Wright, in review), data on macroalgal decomposition often seem to suggest that
 229 detrital photosynthesis is unperturbed until some more proximate mechanism causes it to fail
 230 (de Bettignies et al., 2020, Frontier et al., 2021, Frontier et al., 2022). I therefore tested both
 231 decomposition models on two empirical datasets (Brouwer, 1996, Frontier et al., 2022) given
 232 identical priors to compare performance. Option 2 has marginally better predictive
 233 performance according to leave-one-out cross-validation (Vehtari et al., 2017) but more
 234 importantly has smoother posteriors, while option 1 shows signs of bimodality, non-
 235 identifiability or jaggedness in posteriors (github.com/lukaseamus/limbodeco). Thus I
 236 proceed with option 2 to describe k over time as
 237

$$k(t) = \frac{\alpha + \tau}{1 + e^{\frac{5(t-\mu)}{\mu}}} - \tau \quad (7)$$

239

240 where α are the initial exponential growth rate (aka relative or specific growth rate) of
 241 detritus *sensu lato* at detachment or excision, τ is the final exponential decay rate of detritus
 242 *sensu stricto* once detrital photosynthesis has ceased, and μ is the time of transition between
 243 the two states when the influence of detrital photosynthesis on decomposition ceases. The
 244 influence of each parameter on the trajectory of k is visualised in Figure 1. Inserting Equation
 245 7 into Equation 4 yields

246

$$\frac{dm}{dt} = m \left(\frac{\alpha + \tau}{1 + e^{\frac{5(t-\mu)}{\mu}}} - \tau \right) \quad (8)$$

248

249 This equation could be used directly with statistical software that allows solving ordinary
 250 differential equations on the fly. But there is a simpler, generally applicable solution. Rovira
 251 & Rovira (2010) have shown that the logistic function can simply be integrated alongside the
 252 ordinary differential equation to yield a simpler model akin to the integrated form of Equation
 253 3. Since we know our function is bounded by $t = 0$, we use an accumulation function. The
 254 definite integral of Equation 7 is

255

$$\int_0^t k(t) dt = t\alpha + \frac{\mu(\alpha + \tau)}{5} \ln \frac{1 + e^{\frac{5(t-\mu)}{\mu}}}{1 + e^{-5}} \quad (9)$$

257

258 Thus when Equation 8 is integrated as a whole, we get the final form

259

$$m(t) = m_0 e^{t\alpha + \frac{\mu(\alpha + \tau)}{5} \ln \frac{1 + e^{\frac{5(t-\mu)}{\mu}}}{1 + e^{-5}}} \quad (10)$$

261

Since decomposition data are best analysed as proportions of initial mass, $m_0 = 1$ which simplifies the function to one with only three parameters, the now familiar α , μ and τ . The influence of each parameter on the shape of the curve m describes over time is illustrated in Figure 1. Note that these parameters may be formally compared within and between studies, including those that used the traditional model (Equation 3). τ is the exponential decay rate of detritus *sensu stricto* which is not under the influence of detrital photosynthesis. Since the traditional model assumes that physiology does not play a role in detrital dynamics, k in Equation 3 also represents the exponential decay rate of detritus *sensu stricto*. Thus τ may be directly compared to k . In the next section I turn this mathematical model into a fully functional statistical model.

The statistical model

Maximum likelihood solutions to nonlinear modelling invariably demand the tedious task of guessing starting values. Rovira & Rovira (2010) followed this approach using the Levenberg-Marquardt algorithm which is more robust to poor starting values than the usual Gauss-Newton algorithm. Despite this advantage, they still needed to manually approximate the starting values before passing them to the algorithm. Bayesian inference using Markov Chain Monte Carlo is more flexible in that there is no practical difference between fitting linear and nonlinear models (McElreath, 2019). Rather than starting values which can make or break the model, a prior probability distribution is provided, which at worst can lead to inefficient sampling and uncertain prediction. With the increase in computing power, every laptop can now run Markov Chain Monte Carlo. For these and other reasons Bayesian inference is quickly becoming the *status quo* in applied statistics across disciplines (McElreath, 2019). The most popular statistical language is R (R Core Team, 2022) and the most popular Bayesian language is Stan (Carpenter et al., 2017) which integrates nicely with

287 R through various packages. Hence I developed the statistical model within this joint
288 framework.

289

290 Several hierarchical functions are needed to describe a full statistical model. The one to rule
291 them all is the likelihood function. It describes the probability distribution of the data. Most
292 probability distributions are fully described by two parameters, usually the expected value
293 and spread. For the normal distribution this is the mean and variance. Each of these
294 likelihood parameters needs to be a function of the predictor. Finally the functions describing
295 the likelihood parameters in turn have parameters, which in the Bayesian framework are
296 expressed as probability distribution functions, the priors which once conditioned on the data
297 turn into posteriors.

298

299 The mathematical model (Equation 10) describes the expected value of the likelihood. But
300 what likelihood? We know that the data are proportions of initial mass that can exceed 1 due
301 to growth caused by detrital photosynthesis. Several distributions are restricted to positive
302 real numbers, such as the popular gamma distribution. But we can do better by thinking
303 generatively. The proportions are ratios of mass to initial mass. Mass itself can only take
304 positive real numbers and may be thought of as the sum of cells, growth increments etc. Thus
305 the gamma distribution describes mass well. The ratio of two gamma distributions is a beta
306 prime distribution[§]. Voilà, we have our likelihood!

[§] Technically this is only true if both gamma distributions share the same rate or scale, i.e. size of summand, and are independent. We can assume the former because both measures of mass can be broken down into the same components, e.g. cells. It is difficult to assume the latter because the fundamental assumption of the model is that the change in mass depends on

307

308 The beta prime distribution, typically parameterised by the two shape parameters α and β , is309 most intuitively parameterised in terms of mean $\mu = \frac{\alpha}{\beta - 1}$ and precision $\nu = \beta - 2$, where310 variance $\sigma^2 = \frac{\mu(1 + \mu)}{\nu}$ (Bourguignon et al., 2021). To reparametrise the likelihood, α and β 311 need to be expressed in terms of μ and ν as $\alpha = \mu(1 + \nu)$ and $\beta = 2 + \nu$. We can either assume312 ν to be a constant (homoskedasticity) or a function of the predictor (heteroskedasticity).313 Given that for proportional data $m(0) = 1$ is predetermined, there is no uncertainty at $t = 0$, so314 ν is theoretically infinite to begin with. But with time comes uncertainty and we expect

315 precision to decline. Sometimes remaining mass is calculated as a proportion of the mean

316 rather than sample-specific initial mass so there is variability at $t = 0$ (e.g. Brouwer, 1996),317 but even then the variability at $t = 0$ is naturally lowest. Since $\nu > 0$ an appropriate model is

318

319
$$\nu(t) = (\epsilon - \theta)e^{-\lambda t} + \theta \quad (11)$$

320

321 where ϵ is ν at $t = 0$, λ is the rate of exponential decay in ν , and θ is ν as $t \rightarrow \infty$, inspired by its

322 representation of Thánatos. Combining the beta prime likelihood and Equations 10 and 11

323 yields the full statistical model:

324

325
$$m \sim \text{Beta}'(\mu_m(1 + \nu), 2 + \nu) \quad (12)$$

326
$$\mu_m = e^{t\alpha + \frac{\mu(\alpha + \tau)}{5}} \ln \frac{1 + e^{\frac{5(t - \mu)}{\mu}}}{1 + e^{-5}}$$

327
$$\nu = (\epsilon - \theta)e^{-\lambda t} + \theta$$

mass (Equation 1, 3). Nonetheless, beta prime remains the best approximation of the true data distribution I can think of.

328 $\alpha \sim N(\dots)$, $\text{Exp}(\dots)$ or $\text{Gamma}(\dots)$

329 $\mu, \tau, \epsilon, \lambda, \theta \sim \text{Exp}(\dots)$ or $\text{Gamma}(\dots)$

330

331 where m is the proportion of remaining mass and μ_m is the mean of m , the remaining

332 parameters being the same as in Equations 10 and 11. α may have a normal prior since it can

333 take positive (initial growth) or negative (initial decomposition) values. However, in some

334 cases it may be desirable to enforce initial growth with an exponential or gamma prior. All

335 other parameters are restricted to positive values, so the exponential or gamma distribution is

336 most suitable. I found generally suitable priors to be $\epsilon \sim \text{Gamma}(4, 10^{-4})$, $\lambda \sim \text{Exp}(1)$ and θ

337 $\sim \text{Gamma}(4, 0.008)$, especially because these parameters are a bit abstract. I use these

338 priors in all examples in the next section. For α , μ and τ I cannot recommend such generally

339 applicable priors. As is usual with Bayesian workflow (McElreath, 2019), priors have to be

340 chosen based on prior simulation. Furthermore, α , μ and τ are biologically meaningful and

341 can thus be strongly informed by the literature. Some of the simplest examples that may be a

342 reasonable starting point are $\alpha \sim N(0, 0.01)$, $\mu \sim \text{Exp}(0.1)$ and $\tau \sim \text{Exp}(10)$. But in most

343 cases these will have to be further constrained to ensure smooth sampling. The best next step

344 would be to use a tighter normal prior for α , a gamma prior centred on half the range of t for

345 μ and something along the lines of $\tau \sim \text{Gamma}(4, 40)$ which has mean 0.1 and standard

346 deviation 0.05. But in most cases you will be able to do far better.

347

348 Other likelihoods may also yield reasonable predictions if Equation 12 cannot be

349 implemented in your statistical framework of choice. The gamma distribution may only get

350 extreme values wrong due to its differently shaped tail and the normal distribution may still

351 provide valid predictions of the mean even though it is not constrained to positive real

352 numbers. If necessary, the model can also be simplified by assuming homoskedasticity,

which may have minimal impact on mean predictions albeit inflating uncertainty near $t = 0$ and potentially underestimating uncertainty as $t \rightarrow \infty$.

There are some final considerations for the statistical model that are purely computational. Stan uses Hamiltonian Monte Carlo to explore probability space. Some of the expressions I have used in Equations 10 and 11 are easier to read but not friendly to the sampler. These can be improved for computational efficiency using the quotient and product rules of logarithms as

$$m(t) = e^{t\alpha + \frac{\mu(\alpha + \nu)}{5}(\ln(1 + e^{\frac{5(t-\mu)}{\mu}}) - \ln(1 + e^{-5}))} \quad (13)$$

$$\nu(t) = e^{\ln(\epsilon - \theta) - \lambda t + \theta} \quad (14)$$

The $\ln(1 + e^x)$ expressions in Equation 13 are further simplified in Stan using the wrapper function `log1p_exp` which is numerically stabler. An equivalent function doesn't exist in R but can be written as `log1p_exp <- function(x) ifelse( x > 0 , x + log1p( exp(-x) ) , log1p( exp(x) ) )`.

In terms of software, I use R v4.5.1 with the tidyverse package family v2.0.0 (Wickham et al., 2019) within the integrated development environment RStudio v2025.09.0+387 (RStudio Team, 2022) and run Stan with the R interface cmdstanr v0.9.0 (Gabry et al., 2024) via CmdStan v2.37.0 (Lee et al., 2017). All models have 8 Markov chains with 10^4 warmup and sampling iterations each. To assess quality, I use $\hat{R}$ scores and visually scrutinise trace rank and pair plots with bayesplot v1.14.0 (Gabry et al., 2019). All reported results are posterior

probabilities and derived central tendencies and intervals calculated with tidybayes v3.0.7 (Kay, 2024). See github.com/lukaseamus/limbodeco for further details.

In the next section I will apply Equation 12 to four ecological effects in six published studies (Bourguès et al., 1996, Brouwer, 1996, Hamersley et al., 2015, de Bettignies et al., 2020, Frontier et al., 2021, Frontier et al., 2022). I either received data from the lead author (Frontier et al., 2021, Frontier et al., 2022) or digitised figures (Bourguès et al., 1996, Brouwer, 1996, Hamersley et al., 2015, de Bettignies et al., 2020) using WebPlotDigitizer v4.6 (Rohatgi, 2022). When several samples were collected at each timepoint, figures only yielded mean and standard deviation of remaining mass. The typical approach in such cases is measurement error modelling to account for the uncertainty of each mean so that means with larger standard deviations carry less weight in influencing the prediction (McElreath, 2019). However, at each timepoint de Bettignies et al. (2020) collected 5 samples, summarised as mean and standard deviation, for one treatment and only 1 sample for the other, resulting in a mixture of sample sizes and data levels. A measurement error model is not applicable in such a case. Furthermore, I attempted building beta prime, gamma and lognormal measurement error models and they all result in poor sampling when paired with the beta prime likelihood. Only a zero-truncated normal measurement error distribution was feasible but this does not necessarily represent the data distribution. I therefore simulated observations from summary statistics as random draws from the beta prime distribution in R using the *rbetapr* function from extraDistr v1.10.0 (Wolodzko, 2023) as

```
rbetapr (
  n ,
  mean * ( 1 + mean * ( 1 + mean ) / sd^2 ) ,
  (15)
```

402 $2 + \text{mean} * (1 + \text{mean}) / \text{sd}^2$
 403)

404

405 where n , mean and sd are the reported sample size, sample mean and sample standard
 406 deviation. I derived this alternative parameterisation of the beta prime distribution by
 407 inserting $v = \frac{\mu(1+\mu)}{\sigma^2}$ into $\alpha = \mu(1+v)$ and $\beta = 2+v$ (Bourguignon et al., 2021).

408

409 **Examples**

410 **Dead or alive**

411 Brouwer (1996) provides a nice example of detrital photosynthesis influencing
 412 decomposition of the feathery brown seaweed *Desmarestia anceps* at Signy Island in
 413 Antarctica. She killed some of the detritus *sensu lato* by heating to 50°C for 10–15 min prior
 414 to *in situ* decomposition (Brouwer, 1996). The resulting effect is remarkable but Brouwer
 415 (1996) was unable to quantify it due to lack of a suitable model at the time, so only provided
 416 an analysis of variance F statistic for change over time in both treatments and described the
 417 divergence in the decomposition trajectories.

418

419 The statistical model outlined in Equation 12 given this dataset and priors $\alpha \sim N(-$
 420 $0.005, 0.003)$, $\mu \sim \text{Gamma}(\frac{200^2}{150^2}, \frac{200}{150^2})$ and $\tau \sim \text{Gamma}(\frac{0.1^2}{0.05^2}, \frac{0.1}{0.05^2})$ is visualised in Figure
 421 2. Predictions across t for new observations of m based on the full likelihood ($\tilde{m}$) are in the
 422 top panel, followed by the likelihood parameters v and μ_m , and k in the bottom panel. One
 423 could also overlay the probability distributions for the basal parameters α , μ and τ and ϵ and θ
 424 onto the plots of k and v respectively to present the full hierarchy of the model. To avoid
 425 clutter I have opted to only display μ , since this is the key parameter, marking the half-life of
 426 detrital photosynthesis, which, unlike α , τ , ϵ and θ , cannot be guessed by looking at

predictions of k and v . Summaries of posterior probability distributions for the basal parameters describing μ_m and k are given in Table 1. I decided to pool τ , ϵ and θ across treatments. The experimental duration is not long enough to inform τ for the control treatment and it is reasonable to assume that untreated detritus decomposes at the same rate as pre-killed detritus once physiology has ceased. There is also no reason to suspect ϵ and θ to vary across treatments, since ϵ is only governed by the variability in initial experimental mass, which should be unaffected by treatment, and θ goes hand in hand with τ .

The model suggests that pre-killed *D. anceps* detritus decomposes much faster than the control (Figure 2). Untreated detritus has an initial relative growth rate of $0.037 \pm 0.017 \%$ d^{-1} (mean $\pm$ standard deviation, $P_{\alpha > 0} = 0.99$) while pre-killed detritus has an initial decay rate of $0.6 \pm 0.29 \%$ d^{-1} ($P_{\alpha < 0} = 0.98$) (Table 1). Detrital photosynthesis of *D. anceps* only ceases to influence decomposition after $778 \pm 108 \text{ d}$ (Table 1). Pre-killed detritus starts decomposing at the ultimate rate $750 \pm 106 \text{ d}$ earlier ($P_{\Delta\mu > 0} = 1$) (Table 1). Without physiology, detritus of *D. anceps* therefore decomposes in about $3.6 \pm 0.56 \%$ of the expected time.

Senescence

Hamersley et al. (2015) and de Bettignies et al. (2020) both studied the effect of tissue senescence on kelp decomposition. Hamersley et al. (2015) collected attached fresh and senescent fronds of the giant kelp *Macrocystis pyrifera* at Catalina Island in California along with naturally detached wrack and raft material which they regarded as detritus *sensu stricto*. de Bettignies et al. (2020) collected tissue from above and below the May cast abscission border of the kelp *Laminaria hyperborea* at Roscoff in Brittany. Their senescent blade thus represents tissue that the plant is in the process of abscising. Hamersley et al. (2015) refrained

from fitting a model but de Bettignies et al. (2020) are among the few that fit Equation 3 to kelp decomposition data.

My models for *M. pyrifera* (Figure 3a) and *L. hyperborea* (Figure 3b) were both given priors $\alpha \sim N(0, 0.01)$ and $\mu \sim \text{Exp}(0.1)$. But since de Bettignies et al. (2020) reported k values (Equation 3) of 0.0366 d^{-1} for senescent and 0.0107 d^{-1} for fresh lamina, I was able to better constrain the τ prior for this dataset. I therefore used $\tau \sim \text{Gamma}(\frac{0.2^2}{0.1^2}, \frac{0.2}{0.1^2})$ for *M. pyrifera* and $\tau \sim \text{Gamma}(\frac{0.0366^2}{0.02^2}, \frac{0.0366}{0.02^2})$ for *L. hyperborea*. For clarity I here only show predictions for k and μ_m with posterior probability distributions for μ , but the model provides the full set of information as in Figure 2. While species is constant across treatments in each study, the detritus *sensu lato* has various natural origins rather than having been subjected to different experimental treatments as in the previous example. So I allowed all parameters except for ϵ to vary by treatment. I still pooled ϵ because, as is usual but unlike the previous example, there are no data to inform variability at $t = 0$ in these datasets.

Both models state that senescent fronds decompose faster than fresh ones, and detached ones decompose faster still (Figure 3). Initial decomposition rates for all treatments of *M. pyrifera* (Table 1) are not distinguishable from zero ($P_{\alpha < 0} = 0.53\text{--}0.74$) or from one another ($P_{\Delta\alpha > 0} = 0.53\text{--}0.58$). However, the drop in k takes place $1.3 \pm 0.81 \text{ d}$ ($P_{\Delta\mu > 0} = 0.97$) earlier for detached than senescent tissue for which in turn it happens $3.6 \pm 1.3 \text{ d}$ ($P_{\Delta\mu > 0} = 1$) earlier than for fresh tissue (Figure 3a). The final decomposition rate for fresh tissue is $2.8 \pm 1.2 \text{ \% d}^{-1}$ ($P_{\Delta\tau > 0} = 0.99$) and $0.47 \pm 1.2 \text{ \% d}^{-1}$ ($P_{\Delta\tau > 0} = 0.65$) greater than for senescent and detached tissue. Fresh *L. hyperborea* has an initial growth rate of $1 \pm 0.6 \text{ \% d}^{-1}$ ($P_{\alpha > 0} = 0.97$) while senescent tissue of the same plant has an initial decay rate of $0.27 \pm 1 \text{ \% d}^{-1}$ ($P_{\alpha < 0} = 0.6$) (Table 1). Detrital photosynthesis in fresh tissue apparently ceases relatively early—after

only 10 ± 3.1 d which is 6.7 ± 4.9 d ($P_{\Delta\mu > 0} = 0.93$) after the senescent tissue—but its final decomposition rate is less than half that of senescent tissue ($\Delta\tau = 2 \pm 0.38$ % d⁻¹, $P_{\Delta\tau > 0} = 1$) (Table 1). Due to this apparently relatively minor influence of detrital photosynthesis, decomposition rates are ultimately similar to those reported by (de Bettignies et al., 2020): 0.013 ± 0.0011 d⁻¹ vs. 0.0107 d⁻¹ for fresh and 0.032 ± 0.0036 vs. 0.0366 d⁻¹ for senescent *L. hyperborea* (Table 1).

Light

The effect of depth on decomposition rates has repeatedly been evidenced for *Laminaria* species (Frontier et al., 2021, Frontier et al., 2022). Detritus *sensu lato* of *L. hyperborea* and *L. ochroleuca* was either exposed to three simulated depths *ex situ* at Roscoff in Brittany (Frontier et al., 2021) or deployed at three depths *in situ* across the Channel at Plymouth in Devon (Frontier et al., 2022). Photophysiological measurements across detrital age have clarified that the amount of light available to detrital photosynthesis is the mechanism behind this effect (Frontier et al., 2021, Frontier et al., 2022). However, detrital photosynthesis and decomposition have not been quantitatively linked. Furthermore remaining mass was analysed using linear models which don't do a very good job given the nature of the data (Frontier et al., 2021, Frontier et al., 2022).

Given identical priors $\alpha \sim \text{Exp}(100)$, $\mu \sim \text{Gamma}(\frac{50^2}{30^2}, \frac{50}{30^2})$ and $\tau \sim \text{Gamma}(\frac{0.1^2}{0.05^2}, \frac{0.1}{0.05^2})$, the results for both datasets are shown in Figure 4. Note that I am here making use of the possibility of constraining α to positive real numbers. All experimental tissue was excised from fresh parts of the lamina (Frontier et al., 2021, Frontier et al., 2022) so we expect initial growth. Due to the complexity of this dataset with three predictor variables, I only show $\tilde{m}$, the posterior we would use to ultimately make predictions. In both cases I let α vary by

species but not treatment, since there is no reasonable way for treatment to affect initial growth, and μ vary by species and treatment. As with α , for the first dataset I let τ vary by species but not by treatment following the rationale outlined above that detritus of the same origin can be expected to ultimately decompose at the same rate. For the second dataset I pooled τ across species and treatment since the experimental duration is not long enough to inform τ for *L. hyperborea* (Figure 4b). A more elegant solution would be partial pooling using multilevel modelling but that is beyond the scope of this paper. ϵ again was completely pooled and θ followed the pooling of τ for reasons outlined before.

Both models suggest that decomposition is faster at greater depths (Figure 4) for reasons outlined above. Decomposition also tends to be faster for *L. ochroleuca* than *L. hyperborea* (Figure 4), which is well established (Wright et al., 2022), but this does not hold true at greater depths (Figure 4a). Initial relative growth rates (Table 1) are not different between species for the first model ($P_{\Delta\alpha > 0} = 0.51$) but are $0.34 \pm 0.13 \text{ \% d}^{-1}$ ($P_{\Delta\alpha > 0} = 1$) greater for *L. ochroleuca* than *L. hyperborea* in the second model. In the first model, detrital photosynthesis of *L. hyperborea* ceases $175 \pm 38 \text{ d}$ ($P_{\Delta\mu > 0} = 1$) later at the surface but $50 \pm 17 \text{ d}$ ($P_{\Delta\mu > 0} = 1$) and $12 \pm 3.9 \text{ d}$ ($P_{\Delta\mu > 0} = 1$) earlier at 15 and 30 m. Such a reversal in physiological persistence with depth is not evident in the second model, where *L. hyperborea* detritus *sensu lato* photosynthesises $89 \pm 35 \text{ d}$ ($P_{\Delta\mu > 0} = 1$), $81 \pm 32 \text{ d}$ ($P_{\Delta\mu > 0} = 1$) and $27 \pm 17 \text{ d}$ ($P_{\Delta\mu > 0} = 1$) longer at 0.5, 1.5 and 3 m, probably due to the lesser depth range. In both models the maximal negative effect of depth on the longevity of detrital photosynthesis is stronger for *L. hyperborea* ($\Delta\mu = 87 \pm 37 - 230 \pm 38 \text{ d}$, $P_{\Delta\mu > 0} = 0.99-1$) than *L. ochroleuca* ($\Delta\mu = 25 \pm 12 - 43 \pm 6.4 \text{ d}$, $P_{\Delta\mu > 0} = 1$). So the detrital physiology of *L. hyperborea* is less tolerant to low light, which agrees with observed kelp forest zonation (Wright et al., 2022). The first model, with which it was possible to quantify differences in τ (Table 1), suggests that *L. ochroleuca*

ultimately decomposes $0.6 \pm 0.89 \% \text{ d}^{-1}$ ($P_{\Delta\tau > 0} = 0.75$) faster than *L. hyperborea*, independent of physiology. However, contrasting detrital physiology is clearly the main driver of the interspecific difference in decomposition timescales, as has been previously hinted at (Wright et al., 2022).

Season

Bourguès et al. (1996) provide perhaps the only data of macroalgal decomposition across all seasons. They studied decomposition of the ephemeral green seaweed *Ulvaria obscura* in the intertidal Arcachon Bay in France. Bourguès et al. (1996) collected fresh *U. obscura* but buried it below 1 cm of sediment where anoxia apparently persisted and there clearly was no light for detrital photosynthesis. Given these experimental conditions, it would be surprising to see decomposition affected by physiology. Yet Bourguès et al. (1996) were faced with the challenge of having to model decomposition trajectories of sigmoidal appearance for which they chose the logistic function. Excised macroalgae can survive exceptional periods of darkness (Wright et al., 2024) so it may be that saprotrophs were initially inhibited by chemical defence despite the absence of photosynthesis. Even if this was not the case because anoxia immediately killed the seaweed, and the model therefore is conceptually wrong, it is still useful because it avoids having to fit fundamentally incomparable models as Bourguès et al. (1996) have done.

Given this dataset and priors $\alpha \sim \text{Exp}(100)$, $\mu \sim \text{Gamma}(\frac{25^2}{10^2}, \frac{25}{10^2})$ and $\tau \sim \text{Gamma}(\frac{0.1^2}{0.05^2}, \frac{0.1}{0.05^2})$, $\tilde{m}$ is visualised in Figure 5. Again I make use of constraining α to positive values because the seaweed was initially alive. Season, which for all intents and purposes is temperature, affects detrital physiology as well as other aspects that may drive decomposition post-physiology such as saprotroph metabolic rates (Filbee-Dexter et al., 2022, Wright et al.,

2024). So all parameters are expected to be affected by season. I therefore only pooled ϵ across treatments for reasons outlined above.

According to the model decomposition is generally faster in warmer seasons (Figure 5). This agrees well with the known effect of temperature on decomposition (Filbee-Dexter et al., 2022, Wright et al., 2024). Initial growth rate is not strikingly different between any two seasons ($P_{\Delta\alpha > 0} = 0.54\text{--}79$), perhaps unsurprisingly given data sparsity immediately after the start of the experiment (Figure 5). Differences are more pronounced in the decline of k over time. k shifts at a similar time in autumn and winter ($P_{\Delta\mu > 0} = 0.52$) but does so 9 ± 3.6 d ($P_{\Delta\mu > 0} = 1$) and 19 ± 3.3 d ($P_{\Delta\mu > 0} = 1$) earlier in spring and summer than in autumn (Table 1). This may be the detrimental effect of increased temperature on detrital physiology (Wright et al., 2024) since spring and summer are generally warmer than autumn and winter, unless we believe the experimental conditions immediately killed the seaweed. The final value of k is 6 ± 6.4 % d⁻¹ ($P_{\Delta\tau > 0} = 0.84$), 2.8 ± 5.5 % d⁻¹ ($P_{\Delta\tau > 0} = 0.73$) and 5 ± 6 % d⁻¹ ($P_{\Delta\tau > 0} = 0.81$) lower—i.e. decomposition is faster and τ is greater—in spring, summer and autumn than in winter (Table 1). This is the effect of seasonal temperature differences on factors other than detrital physiology. Again the physiological effect, albeit putative in this case, is more pronounced, as has been suggested in the context of temperature (Wright et al., 2024).

Conclusion

Macroalgal decomposition has neither been consistently nor reliably modelled. But a realistic model is the key to understanding macroalgal blue carbon. I have presented the first decomposition model tailored to macroalgae by taking into account the theory of detrital photosynthesis. This is made possible by treating the exponential decay rate as a function of time rather than a constant, a function which is informed by the latest understanding of

detrital photosynthesis. My model only has three parameters which describe the initial relative growth rate (α), the half-life of detrital photosynthesis (μ), and the final exponential decay rate (τ) which is directly comparable to that of the classic model (k). I have shown that these parameters are successfully estimated and meaningful for empirical data from a variety of contexts. Given a set of parameter estimates, the model can reliably predict new observations, even beyond the duration of the experiment. Although I have written the model in R with Stan, it can be implemented in any statistical framework that allows robust nonlinear modelling. Failing to account for detrital photosynthesis almost invariably leads to underestimation of macroalgal decomposition. I therefore expect that this model will increase estimates of macroalgal decomposition and thus have major implications for macroalgal blue carbon accounting. I will be using this model to clarify the role of kelp forests in CO₂ removal and hope others find it useful too.

Acknowledgements

Thanks go to Andy Foggo and Nadia Frontier for inspiring discussions over the years and Nadia for sharing her data with me. Thanks also to Pere Rovira for the email exchange and Bojana Manojlovic for the chat.

Funding source acknowledgements

This work was supported by the Forrest Research Foundation Scholarship and Australian Government Research Training Program Scholarship. Open access publishing was facilitated by The University of Western Australia, as part of the Wiley–The University of Western Australia agreement via the Council of Australian University Librarians.

Conflict of interest statement

I declare no conflicts of interest.

Data availability statement

Data and code to reproduce this paper and apply my model to your own data are available at github.com/lukaseamus/limbodeco. Please cite this paper as well as the model (Wright, 2025).

References

- Babiker, M., G. Berndes, K. Blok, B. Cohen, A. Cowie, O. Geden, V. Ginzburg, A. Leip, P. Smith, M. Sugiyama, and F. Yamba. 2022. "Cross-sectoral perspectives." in Shukla, P. R., J. Skea, R. Slade, A. Al Khourdajie, R. van Diemen, D. McCollum, M. Pathak, S. Some, P. Vyas, R. Fradera, M. Belkacemi, A. Hasija, G. Lisboa, S. Luz, and J. Malley, editors. "IPCC, 2022: Climate Change 2022: Mitigation of Climate Change. Contribution of Working Group III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change." Cambridge University Press, Cambridge and New York. <https://doi.org/10.1017/9781009157926.005>.
- Banta, G. T., M. F. Pedersen, and S. L. Nielsen. 2004. "Decomposition of marine primary producers: consequences for nutrient recycling and retention in coastal ecosystems." Pages 187-216 in Nielsen, S. L., G. T. Banta, and M. F. Pedersen, editors. "Estuarine nutrient cycling: the influence of primary producers." Springer, Dordrecht. https://doi.org/10.1007/978-1-4020-3021-5_7.
- Bedford, A. P., and P. G. Moore. 1984. "Macrofaunal involvement in the sublittoral decay of kelp debris: the detritivore community and species interactions." *Estuarine, Coastal and Shelf Science* 18(1): 97-111. [https://doi.org/10.1016/0272-7714\(84\)90009-X](https://doi.org/10.1016/0272-7714(84)90009-X).

- 625 Birch, P. B., J. O. Gabrielson, and K. S. Hamel. 1983. "Decomposition of *Cladophora*. I.
626 Field studies in the Peel-Harvey estuarine system, Western Australia." *Botanica Marina*
627 26(4): 165-172. <https://doi.org/10.1515/botm.1983.26.4.165>.
- 628 Bourguès, S., I. Auby, R. de Wit, and P.-J. Labourg. 1996. "Differential anaerobic
629 decomposition of seagrass (*Zostera noltii*) and macroalgal (*Monostroma obscurum*)
630 biomass from Arcachon Bay (France)." *Hydrobiologia* 329:121-131.
631 <https://doi.org/10.1007/bf00034552>.
- 632 Bourguignon, M., M. Santos-Neto, and M. D. Castro. 2021. "A new regression model for
633 positive random variables with skewed and long tail." *Metron* 79:33-55.
634 <https://doi.org/10.1007/s40300-021-00203-y>.
- 635 Brouwer, P. E. M. 1996. "Decomposition in situ of the sublittoral Antarctic macroalga
636 *Desmarestia anceps* Montagne." *Polar Biology* 16(2): 129-137.
637 <https://doi.org/10.1007/BF02390433>.
- 638 Carpenter, B., A. Gelman, M. D. Hoffman, D. Lee, B. Goodrich, M. Betancourt, M.
639 Brubaker, J. Guo, P. Li, and A. Riddell. 2017. "Stan: a probabilistic programming
640 language." *Journal of Statistical Software* 76(1): 1-32.
641 <https://doi.org/10.18637/jss.v076.i01>.
- 642 Cebrián, J., and C. M. Duarte. 1995. "Plant growth-rate dependence of detrital carbon storage
643 in ecosystems." *Science* 268(5217): 1606-1608.
644 <https://doi.org/10.1126/science.268.5217.1606>.
- 645 Cebrián, J., and J. Lartigue. 2004. "Patterns of herbivory and decomposition in aquatic and
646 terrestrial ecosystems." *Ecological Monographs* 74(2): 237-259.
647 <https://doi.org/10.1890/03-4019>.

- 648 Creed, J. C., V. M. N. C. S. Vieira, T. A. Norton, and D. Caetano. 2019. "A meta-analysis
649 shows that seaweeds surpass plants, setting life-on-Earth's limit for biomass packing."
650 *BMC Ecology* 19(1): 6. <https://doi.org/10.1186/s12898-019-0218-z>.
- 651 de Bettignies, F., P. Dauby, F. Thomas, A. Gobet, L. Delage, O. Bohnner, S. Loisel, and D.
652 Davoult. 2020. "Degradation dynamics and processes associated with the accumulation
653 of *Laminaria hyperborea* (Phaeophyceae) kelp fragments: an in situ experimental
654 approach." *Journal of Phycology* 56(6): 1481-1492. <https://doi.org/10.1111/jpy.13041>.
- 655 Duarte, C. M., J.-P. Gattuso, K. Hancke, H. Gundersen, K. Filbee-Dexter, M. F. Pedersen, J.
656 J. Middelburg, M. T. Burrows, K. A. Krumhansl, and T. Wernberg. 2022. "Global
657 estimates of the extent and production of macroalgal forests." *Global Ecology and*
658 *Biogeography* 01-18. <https://doi.org/10.1111/geb.13515>.
- 659 Enríquez, S., C. M. Duarte, and K. Sand-Jensen. 1993. "Patterns in decomposition rates
660 among photosynthetic organisms: the importance of detritus C:N:P content." *Oecologia*
661 94(4): 457-471. <https://doi.org/10.1007/BF00566960>.
- 662 Filbee-Dexter, K., C. J. Feehan, D. A. Smale, K. A. Krumhansl, S. Augustine, F. de
663 Bettignies, M. T. Burrows, J. E. K. Byrnes, J. Campbell, D. Davoult, K. H. Dunton, J.
664 N. Franco, I. Garrido, S. P. Grace, K. Hancke, L. E. Johnson, B. Konar, P. J. Moore, K.
665 M. Norderhaug, A. O'Dell, M. F. Pedersen, A. K. Salomon, I. Sousa-Pinto, S. Tiegs, D.
666 Yiu, and T. Wernberg. 2022. "Kelp carbon sink potential decreases with warming due
667 to accelerating decomposition." *PLoS Biology* 20(8): e3001702.
668 <https://doi.org/10.1371/journal.pbio.3001702>.
- 669 Filbee-Dexter, K., A. Pessarrodona, M. F. Pedersen, T. Wernberg, C. M. Duarte, J. Assis, T.
670 Bekkby, M. T. Burrows, D. F. Carlson, and J.-P. Gattuso. 2024. "Carbon export from
671 seaweed forests to deep ocean sinks." *Nature Geoscience* 17(6): 552-559.
672 <https://doi.org/10.1038/s41561-024-01449-7>.

- 673 Friedlingstein, P., M. O'sullivan, M. W. Jones, R. M. Andrew, J. Hauck, P. Landschützer, C.
674 Le Quéré, H. Li, I. T. Luijkx, and A. Olsen. 2025. "Global carbon budget 2024." *Earth*
675 *System Science Data Discussions* 17(3): 965-1039. [https://doi.org/10.5194/essd-17-965-](https://doi.org/10.5194/essd-17-965-2025)
676 2025.
- 677 Frontier, N., F. de Bettignies, A. Foggo, and D. Davoult. 2021. "Sustained productivity and
678 respiration of degrading kelp detritus in the shallow benthos: detached or broken, but
679 not dead." *Marine Environmental Research* 166105277.
680 <https://doi.org/10.1016/j.marenvres.2021.105277>.
- 681 Frontier, N., M. Mulas, A. Foggo, and D. A. Smale. 2022. "The influence of light and
682 temperature on detritus degradation rates for kelp species with contrasting thermal
683 affinities." *Marine Environmental Research* 173105529.
684 <https://doi.org/10.1016/j.marenvres.2021.105529>.
- 685 Gabry, J., R. Češnovar, A. Johnson, and S. Bronder. 2024. "cmdstanr: R Interface to
686 'CmdStan'." R package version 0.5.3. <https://mc-stan.org/cmdstanr>.
- 687 Gabry, J., D. Simpson, A. Vehtari, M. Betancourt, and A. Gelman. 2019. "Visualization in
688 Bayesian workflow." *Journal of the Royal Statistical Society Series A: Statistics in*
689 *Society* 182(2): 389-402. <https://doi.org/10.1111/rssa.12378>.
- 690 Hamersley, M. R., J. A. Sohm, J. A. Burns, and D. G. Capone. 2015. "Nitrogen fixation
691 associated with the decomposition of the giant kelp *Macrocystis pyrifera*." *Aquatic*
692 *Botany* 12557-63. <https://doi.org/10.1016/j.aquabot.2015.05.003>.
- 693 Hibbett, D., R. Blanchette, P. Kenrick, and B. Mills. 2016. "Climate, decay, and the death of
694 the coal forests." *Current Biology* 26(13): R563-R567.
695 <https://doi.org/10.1016/j.cub.2016.01.014>.
- 696 Kay, M. 2024. "tidybayes: tidy data and geoms for Bayesian models v3.0.7." *Zenodo*.
697 <https://doi.org/10.5281/zenodo.13770114>.

- 698 Kennedy, J. R., and C. O. Blain. 2025. "A systematic review of marine macroalgal
699 degradation: toward a better understanding of macroalgal carbon sequestration
700 potential." *Journal of Phycology* 61(3): 399-432. <https://doi.org/10.1111/jpy.70031>.
- 701 Lee, D., Stan buildbot, seantalts, B. Carpenter, M. Morris, A. Kucukelbir, M. Betancourt, D.
702 Tran, M. Brubaker, B. Bales, bgoodri, A. Vehtari, wds15, maverickg, evelynmitchell,
703 weberse2, and P. Li. 2017. "stan-dev/cmdstan: v2.17.1." *Zenodo*.
704 <https://doi.org/10.5281/zenodo.1117248>.
- 705 Manojlovic, B. 2021. "Data assimilation techniques for aquatic ecological processes."
706 <https://opus.lib.uts.edu.au/handle/10453/150726>.
- 707 McElreath, R. 2019. Statistical rethinking²: a Bayesian course with examples in R and Stan.
708 Second Edition. Chapman & Hall/CRC, Boca Raton.
- 709 Murray, L. M. 2015. "Bayesian state-space modelling on high-performance hardware using
710 LibBi." *Journal of Statistical Software* 67(10): 1-36.
711 <https://doi.org/10.18637/jss.v067.i10>.
- 712 Olson, J. S. 1963. "Energy storage and the balance of producers and decomposers in
713 ecological systems." *Ecology* 44(2): 322-331. <https://doi.org/10.2307/1932179>.
- 714 Pedersen, M. F., K. Filbee-Dexter, N. L. Frisk, Z. Sárossy, and T. Wernberg. 2021. "Carbon
715 sequestration potential increased by incomplete anaerobic decomposition of kelp
716 detritus." *Marine Ecology Progress Series* 66053-67.
717 <https://doi.org/10.3354/meps13613>.
- 718 Pessarrodona, A., J. Assis, K. Filbee-Dexter, M. T. Burrows, J.-P. Gattuso, C. M. Duarte, D.
719 Krause-Jensen, P. J. Moore, D. A. Smale, and T. Wernberg. 2022. "Global seaweed
720 productivity." *Science Advances* 8(37): eabn2465.
721 <https://doi.org/10.1126/sciadv.abn2465>.

- 722 Pessarrodona, A., R. M. Franco-Santos, L. S. Wright, M. A. Vanderkluft, J. Howard, E.
723 Pidgeon, T. Wernberg, and K. Filbee-Dexter. 2023. "Carbon sequestration and climate
724 change mitigation using macroalgae: a state of knowledge review." *Biological Reviews*
725 98(6): 1945-1971. <https://doi.org/10.1111/brv.12990>.
- 726 R Core Team. 2022. R: a language and environment for statistical computing. R Foundation
727 for Statistical Computing, Vienna.
- 728 Rieper-Kirchner, M. 1989. "Microbial degradation of North Sea macroalgae: field and
729 laboratory studies." *Botanica Marina* 32(3): 241-252.
730 <https://doi.org/10.1515/botm.1989.32.3.241>.
- 731 Rohatgi, A. 2022. "WebPlotDigitizer v4.6." <https://automeris.io>.
- 732 Rovira, P., and R. Rovira. 2010. "Fitting litter decomposition datasets to mathematical
733 curves: towards a generalised exponential approach." *Geoderma* 155(3-4): 329-343.
734 <https://doi.org/10.1016/j.geoderma.2009.11.033>.
- 735 RStudio Team. 2022. RStudio: integrated development environment for R. RStudio, Inc.,
736 Boston.
- 737 Smith, B. D., and R. E. Foreman. 1984. "An assessment of seaweed decomposition within a
738 southern Strait of Georgia seaweed community." *Marine Biology* 84(2): 197-205.
739 <https://doi.org/10.1007/BF00393005>.
- 740 Thomas, H. 2013. "Senescence, ageing and death of the whole plant." *New Phytologist*
741 197(3): 696-711. <https://doi.org/10.1111/nph.12047>.
- 742 Trevathan-Tackett, S. M., T. C. Jeffries, P. I. Macreadie, B. Manojlovic, and P. Ralph. 2020.
743 "Long-term decomposition captures key steps in microbial breakdown of seagrass
744 litter." *Science of The Total Environment* 705135806.
745 <https://doi.org/10.1016/j.scitotenv.2019.135806>.

- Vehtari, A., A. Gelman, and J. Gabry. 2017. "Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC." *Statistics and Computing* 27:1413-1432. <https://doi.org/10.1007/s11222-016-9696-4>.
- White, L. J., and A. Norkko. 2025. "A path towards appropriate degradation experiments for assessing carbon sequestration potential of macroalgae." *Journal of Ecology* 113(10): 2730-2743. <https://doi.org/10.1111/1365-2745.70107>.
- Wickham, H., M. Averick, J. Bryan, W. Chang, L. D. McGowan, R. François, G. Grolemund, A. Hayes, L. Henry, and J. Hester. 2019. "Welcome to the Tidyverse." *Journal of Open Source Software* 4(43): 1686. <https://doi.org/10.21105/joss.01686>.
- Wolodzko, T. 2023. "extraDistr: additional univariate and multivariate distributions." R package version 1.10.0. <https://doi.org/10.32614/CRAN.package.extraDistr>.
- Wright, L. S. 2025. "limbodeco: a model of macroalgal decomposition (Software)." *Zenodo*.
- Wright, L. S., and A. Foggo. 2021. "Photosynthetic pigments of co-occurring Northeast Atlantic *Laminaria* spp. are unaffected by decomposition." *Marine Ecology Progress Series* 678:227-232. <https://doi.org/10.3354/meps13886>.
- Wright, L. S., and L. Kregting. 2023. "Genus-specific response of kelp photosynthetic pigments to decomposition." *Marine Biology* 170(11): 144. <https://doi.org/10.1007/s00227-023-04289-y>.
- Wright, L. S., A. Pessarrodona, and A. Foggo. 2022. "Climate-driven shifts in kelp forest composition reduce carbon sequestration potential." *Global Change Biology* 28(18): 5514-5531. <https://doi.org/10.1111/gcb.16299>.
- Wright, L. S., T. Simpkins, K. Filbee-Dexter, and T. Wernberg. 2024. "Temperature sensitivity of detrital photosynthesis." *Annals of Botany* 133(1): 17-28. <https://doi.org/10.1093/aob/mcad167>.

Tables

Table 1. Macroalgal decomposition parameter estimates (mean $\pm$ standard deviation) for Equation 10 (μ_m in Equation 12) based on 8×10^4 posterior samples. Units are given in parentheses. For readability the exponential rates α and τ are given as daily percentages, but they are modelled as daily proportions. Pooled estimates within a study are marked with the same symbol.

Species	Treatment	α (% d ⁻¹)	μ (d)	τ (% d ⁻¹)
Brouwer 1996 <i>Desmarestia anceps</i>	Control	0.037 ± 0.017	778 ± 108	$7.2 \pm 0.91^*$
	Pre-killed	-0.6 ± 0.29	28 ± 4.1	$7.2 \pm 0.91^*$
Hamersley <i>et al.</i> 2015 <i>Macrocystis pyrifera</i>	Fresh	-0.31 ± 0.55	5.9 ± 1	12 ± 1.1
	Senescent	-0.19 ± 1	2.3 ± 0.75	9.3 ± 0.54
	Detached	-0.076 ± 1	0.94 ± 0.3	12 ± 0.47
de Bettignies <i>et al.</i> 2020 <i>Laminaria hyperborea</i>	Fresh	1 ± 0.6	10 ± 3.1	1.3 ± 0.11
	Senescent	-0.27 ± 1	3.3 ± 3.7	3.2 ± 0.36
Frontier <i>et al.</i> 2021 <i>Laminaria hyperborea</i> <i>Laminaria ochroleuca</i>	0 m	$0.03 \pm 0.017^*$	259 ± 48	$4.1 \pm 0.49^*$
	15 m	$0.03 \pm 0.017^*$	27 ± 9.2	$4.1 \pm 0.49^*$
	30 m	$0.03 \pm 0.017^*$	1.9 ± 1.2	$4.1 \pm 0.49^*$
	0 m	$0.037 \pm 0.032^\dagger$	57 ± 6.5	$4.6 \pm 0.71^\dagger$
	15 m	$0.037 \pm 0.032^\dagger$	80 ± 18	$4.6 \pm 0.71^\dagger$
	30 m	$0.037 \pm 0.032^\dagger$	14 ± 3.7	$4.6 \pm 0.71^\dagger$
Frontier <i>et al.</i> 2022 <i>Laminaria hyperborea</i> <i>Laminaria ochroleuca</i>	0.5 m	$0.24 \pm 0.042^*$	105 ± 22	$6.1 \pm 2.2^*$
	1.5 m	$0.24 \pm 0.042^*$	94 ± 20	$6.1 \pm 2.2^*$
	3 m	$0.24 \pm 0.042^*$	42 ± 12	$6.1 \pm 2.2^*$
	0.5 m	$0.58 \pm 0.13^\dagger$	43 ± 9.2	$6.1 \pm 2.2^*$
	1.5 m	$0.58 \pm 0.13^\dagger$	35 ± 8.7	$6.1 \pm 2.2^*$
	3 m	$0.58 \pm 0.13^\dagger$	21 ± 3.8	$6.1 \pm 2.2^*$
Bourguès <i>et al.</i> 1996 <i>Ulvaria obscura</i>	Spring	0.43 ± 0.4	14 ± 2.2	15 ± 4.1
	Summer	0.7 ± 0.71	4.4 ± 1.8	12 ± 2.5
	Autumn	0.6 ± 0.33	23 ± 2.8	14 ± 3.4
	Winter	0.29 ± 0.3	24 ± 7.3	9.5 ± 4.9

Figure captions

Figure 1. The mathematical model. Some reasonable parameter values for α , μ and τ and how they independently determine the behaviour of the exponential decay rate k (Equation 7) and the proportion of remaining mass m (Equation 10) over time t .

Figure 2. The statistical model. Example 1: the effect of physiology on the decomposition of *Desmarestia anceps*, shown as the proportion of remaining mass over time t (data from Brouwer, 1996). Data are plotted alongside predictions of new observations ($\tilde{m}$), the precision (v), the mean (μ_m) and the exponential decay rate (k) (Equation 12). Points with error bars denote mean $\pm$ standard deviation of 5 samples. Lines and intervals are medians and the central 50, 80 and 90% of posterior probability.

Figure 3. The statistical model. Example 2: the effect of senescence on the decomposition of *Macrocystis pyrifera* (**a**, data from Hamersley et al., 2015) and *Laminaria hyperborea* (**b**, data from de Bettignies et al., 2020), shown as the proportion of remaining mass over time t . Data are plotted alongside predictions of the mean (μ_m) and the exponential decay rate (k) (Equation 12). Points with error bars denote mean $\pm$ standard deviation of 5 samples. Points without error bars are 1 sample. Lines and intervals are medians and the central 50, 80 and 90% of posterior probability.

Figure 4. The statistical model. Example 3: the effect of light attenuation with depth on the decomposition of *Laminaria hyperborea* and *Laminaria ochroleuca ex situ* (**a**, data from Frontier et al., 2021) and *in situ* (**b**, data from Frontier et al., 2022), shown as the proportion of remaining mass over time t . Data are plotted alongside predictions of new observations ($\tilde{m}$,

Equation 12). Lines and intervals are medians and the central 50, 80 and 90% of posterior probability.

Figure 5. The statistical model. Example 4: the effect of season on the decomposition of *Ulvaria obscura* (data from Bourguès et al., 1996). Data are plotted alongside predictions of new observations ($\tilde{m}$, Equation 12). Lines and intervals are medians and the central 50, 80 and 90% of posterior probability.

For Review Only









