

Ecological Modelling

A model of macroalgal decomposition

--Manuscript Draft--

Manuscript Number:	ECOMOD-25-2203R1
Article Type:	Research Paper
Keywords:	Bayesian statistics; biogeochemistry; carbon cycle; degradation; kelp; ordinary differential equation
Corresponding Author:	Luka Seamus Wright, PhD The University of Western Australia Oceans Institute Perth, Western Australia AUSTRALIA
First Author:	Luka Seamus Wright
Order of Authors:	Luka Seamus Wright
Abstract:	Macroalgae are currently the focus of blue carbon, biological carbon dioxide (CO ₂) removal in the ocean. Carbon exported as seaweed detritus is equivalent to ~7.5% of anthropogenic CO ₂ emissions annually. Macroalgal blue carbon hinges on the fate of this detritus, so a reliable model of its decomposition is key. However, macroalgal decomposition has been neither consistently nor reliably modelled. Here I present a mathematical model of macroalgal decomposition. Informed by the theory of detrital photosynthesis, it treats detritus as an entity in limbo that can initially either grow or decompose. This is achieved by replacing the rate constant of the conventional exponential decay model with a logistic function of time, mirroring the sigmoidal decay of detrital photosynthesis. The resulting mathematical model has three parameters that describe the initial exponential growth rate, the photosynthetic half-life, and the final exponential decay rate once physiology is factored out. The third parameter is equivalent to the rate constant of the conventional model for dead detritus. Using a Bayesian model with a beta prime likelihood implemented in R and Stan, I show that the statistical implementation performs well given empirical data from a variety of contexts. The presented model can describe any macroalgal decomposition trajectory which enables formal comparison across studies for the first time. I anticipate that this will prove instrumental in resolving the debate around macroalgal CO ₂ removal.
Response to Reviewers:	Reviewer #2: I think the paper is very well written, detailed, and well explained. The development of the equations is well structured. I think the manuscript is ideal for publication in the journal. I have minor comment: Line 9: Are you sure this is the world's first mathematical model for macroalgal decomposition? Thank you. This was also noted by Reviewer #3. While to the best of my knowledge this is indeed the first decomposition model specifically designed with macroalgae in mind, I have changed this statement to "a mathematical model of macroalgal decomposition". It is necessary to present a conceptual model that shows the novelty of the work, such as the degradation of macroalgae with existing models and with the modification of the model you are presenting. This was also requested by Reviewer #3. I have fit the conventional model (e^{-kt}) to each example dataset to show that it is ill-fitting in most cases (cf. Figures 2–6 with S2–S6) and formally compare k and τ as well as $t_{1/2}$ and μ (Table 1, 2). Reviewer #3: You present a decomposition model that explicitly accommodates early increases or plateaus in normalised detrital mass $m(t)$ by modelling a time varying rate $k(t)$ that transitions logistically towards a final rate τ. You fit the model in a Bayesian framework, using a beta prime observation model for ratios $m > 0$ (allowing $m > 1$) and a time varying precision $v(t)$. A range of example datasets is used to demonstrate flexibility and interpretability. The paper is clearly structured, and the illustrative examples and

figures are effective in communicating the behaviour of the model and its parameters. The contribution is promising; the main issues below concern (i) parameter identifiability and ecological interpretation, (ii) validation of the observation model and heteroscedastic structure, and (iii) formal comparison with standard approaches to support synthesis oriented claims.

This is the most competent review I have ever received. Thank you. I encourage you to get in touch for further discussions.

Highlights

The submitted highlights are generally relevant, but I recommend revising them to avoid absolute or difficult-to-substantiate claims. In particular, "First mathematical model of macroalgal decomposition" is hard to verify, and "Can describe any macroalgal decomposition trajectory" is too absolute. Likewise, "Enables meta-analysis..." would read more cautiously as "supports/facilitates meta-analysis", unless a formal demonstration is provided in the manuscript. I suggest rephrasing the highlights to emphasise verifiable contributions (e.g., a parsimonious time-varying $k(t)k(t)k(t)$, use of a beta prime likelihood for positive mass ratios including values >1 , and estimation of a terminal decay parameter τ intended to be comparable with the classical exponential k).

This was also requested by Reviewer #2. I have rewritten the highlights accordingly.

Major comments

1. Identifiability and interpretation of μ (transition time / photosynthetic 'death')
 μ governs the timing of the logistic transition in $k(t)$ and is interpreted as the time at which detrital photosynthesis ceases to matter. In at least one key example, μ appears to be weakly informed by the observation window. In the *Desmarestia anceps* example, the manuscript reports μ on the order of many hundreds of days, whilst the time axis shown in the corresponding figure spans a substantially shorter period. Importantly, the manuscript notes that the experimental duration is not long enough to inform τ for the control treatment, which strengthens the concern that μ may also be poorly identified when the transition sits outside the observed window. Please add (i) explicit identifiability diagnostics (pair plots/correlations for μ - τ and μ - α , trace plots, ESS/Rhat), and (ii) a brief sensitivity analysis for the prior on μ (at least two reasonable priors), reporting impact on τ and on predictive trajectories. Where μ clearly exceeds the experimental duration, please clarify how it should be interpreted ecologically (e.g., 'transition not observed / not estimable within study duration') to avoid over interpretation.

Identifiability

Correct, μ is not identifiable if the experimental duration is of insufficient length. Pair plots, trace rank plots and prior-posterior plots for all examples are now available in the repository (github.com/lukaseamus/limbodeco/tree/main/Plots) Diagnostic plots are too numerous and large for inclusion as supplementary figures, but I explicitly refer the reader to the repository folder.

Interpretability

The most intuitive term for μ is "photosynthetic half-life". I was initially reluctant to use this term because mathematically μ is not exactly equivalent to the exponential half-life $t_{1/2} (\ln 2 / k)$ which ecologists are familiar with. After considering your request for better interpretability, I have changed my mind. I believe the term is defensible because it is more intuitive than "photosynthetic death" or "midpoint of photosynthetic decay" and in the case of the underlying fixed intercept logistic function $f(t) = 1/(1 + \exp(-5/\mu * (t - \mu)))$, $f(0) = 0.99$ (essentially 1) and $f(\mu) = 0.5$, so μ is directly analogous to $t_{1/2}$. I believe with the identifier "photosynthetic" this term can be safely used and distinguished from half-life in the conventional sense. I already use this term in another publication under review.

2. Fixed logistic steepness $r = 5/\mu$: justify and test robustness. To reduce non identifiability you fix the logistic steepness via $r\mu = 5$, yielding a three parameter form for $k(t)$. This is a pragmatic choice, but it couples the timing of the transition (μ) to its abruptness, which may not match systems where photosynthesis remains stable then collapses relatively quickly. Please, (i) provide a clearer justification for the constant '5'

(convention vs biological assumption), and (ii) run a small robustness check on 1-2 representative datasets allowing τ to be estimated under a regularising prior (or a hierarchical structure by dataset), and compare predictive performance and parameter stability.

I rewrote the conceptual reasoning behind fixing the logistic intercept at 5, added the alternative constant slope parametrisation as Figure S1 for comparison with Figure 1, and provide performance tests in Table S1. These are discussed in the new section Parameterisation of $k(t)$ in The statistical model.

3. Observation model (beta prime for m): provide posterior predictive checks and a minimal alternative comparison

The beta prime likelihood for the normalised mass m is an elegant choice because it supports m in $(0, \infty)$, allowing $m > 1$. You motivate it via a gamma ratio argument but also acknowledge that strict independence assumptions are questionable in this context. Request: please include posterior predictive checks focused on (i) tail behaviour for $m > 1$, (ii) behaviour near 0, and (iii) the dispersion pattern over time (in combination with $v(t)$). In addition, please compare beta prime against at least one plausible alternative (e.g., lognormal error on m , or modelling absolute mass with explicit measurement error in $M(0)$) on 1-2 datasets using a predictive metric (e.g., PSIS LOO/WAIC) to demonstrate that beta prime is not merely convenient but substantively improves predictive adequacy. Minor but important: please avoid reusing α both as a biological parameter and as a beta prime shape parameter—rename distributional shapes (e.g., a , b) to prevent confusion.

I also added performance tests for the likelihood. These are presented in Table S2 and discussed in the new section Likelihood in The statistical model.

Some common parameter names are unfortunately bound to be repeated in the manuscript, e.g. μ , σ , α , β . There are no other accepted letters for the shape parameters α and β of the beta and beta prime distributions, or for the mean μ , sd σ , variance σ^2 etc. In each case I take care to avoid confusion by redefining what I mean in the given context (e.g. “the sample mean μ ” vs. “the photosynthetic half-life μ ”).

4. Time varying precision $v(t)$: the imposed form may be too restrictive. You model heteroscedasticity through an exponentially declining precision:

$$v(t) = (\epsilon - \theta) e^{(-\lambda t)} + \theta$$

motivated by minimal uncertainty at $t = 0$ because $m(0) = 1$. While plausible in some contexts, this structure forces dispersion to increase monotonically with time and may not reflect situations where variability depends more on remaining mass, fragmentation events, or normalisation choices (particularly when initial mass is not measured per replicate or when data are reconstructed from summary statistics). Please (i) provide posterior predictive checks demonstrating that the model reproduces the observed spread of m through time, and (ii) include a simple sensitivity comparison to a constant precision model (or another minimal alternative) to show whether $v(t)$ materially affects inference on μ and τ and improves predictive performance.

I added performance tests for modelling heteroskedasticity. See Table S3 and new section Heteroskedasticity in The statistical model.

5. Formal validation against standard models to support synthesis/meta analysis claims. A major selling point is that τ is comparable to the standard exponential k , and that ignoring photosynthetic detritus can 'almost invariably' bias decomposition estimates. To support these strong claims (particularly in a synthesis context), the manuscript needs a concise, quantitative model comparison. For 2-3 representative datasets, please fit and compare: (i) the standard exponential model $m(t) = e^{(-k t)}$, (ii) a simple alternative allowing early increase/plateau (even ad hoc), and (iii) the proposed model. Report predictive performance (e.g., PSIS LOO/WAIC and/or out of sample prediction error) and show how k from the exponential compares with τ . Additionally, please include a brief schematic (main text or appendix) showing how τ (and, where identifiable, μ) would be carried forward into a meta analysis/meta regression, including guidance for datasets where μ is not identified within the study duration.

This was also requested by Reviewer #2. An initial test comparison is provided in Table S4. I then fit the conventional model (e^{-kt}) to each example dataset to show that it is ill-fitting in most cases (cf. Figures 2–6 with S2–S6) and formally compare k and τ as well as $t^{1/2}$ and μ (Table 1, 2). I now use multilevel models in several examples, providing a full model description in each case, to show how the proposed model would be used in meta-analysis.

6. Use of simulated observations from summary statistics (when raw data are unavailable). Where raw data are unavailable, you simulate observations from reported means/SDs/sample sizes. This can be pragmatic, but it can understate uncertainty (digitisation, rounding, distributional assumptions) and may create pseudo replication if many points are generated. It would be interesting to propagate this uncertainty (e.g., repeat the simulation several times and show stability of key parameters), or consider a model at the aggregated level that uses means/SDs directly without generating pseudo data.

I now use a measurement error model to avoid simulating data in the case of Brouwer (1996) and received raw data from the authors of the remaining studies upon renewed request.

Minor comments and editorial suggestions

* The figures illustrating predicted new observations and time varying $k(t)$ are helpful; consider adding one sentence to selected captions explicitly reiterating ecological interpretation of μ and τ for non statistical readers.

See comment on the interpretability of μ above. I added the requested detail in the figure captions.

* Where you discuss relevance for blue carbon, please ensure wording distinguishes flux into detrital pools from net long term sequestration.

As far as I know there is no long-term seaweed carbon sequestration independent of the detrital pool. As I explain in the introduction, only seaweed detritus may be buried as allochthonous carbon. I solely use the term “sequestration” in the introduction. My conclusions relate to decomposition affecting estimates of seaweed blue carbon.

* Reproducibility: you note code/data availability; please also provide a specific repository version (commit/tag) associated with the manuscript results.

I have now published the repo and provide a Zenodo DOI:
<https://doi.org/10.5281/zenodo.18654199>.

* Minor editorial corrections (typos and style). The manuscript is generally well written and clearly presented; however, I noted a few small typographical and stylistic issues that could be corrected during revision:

Introduction (lines 74-75): “disproportionally consequential” → “disproportionately consequential”.

Introduction (line 118): “...to growth rather than decompose...” → “...to grow rather than decompose...”.

Computation (lines 361): “numerically stabler” → “numerically more stable”.

Methods (line 388): “not applicable” → “not applicable”.

Punctuation for readability: “Viewed cumulatively this...” (lines 87-88) → add a comma after “Viewed cumulatively,” and “Detritus on the other hand...” (lines 64-65) → “Detritus, on the other hand, ...”.

Style: consider “a relationship that is parameterised...” rather than “...which is parameterised...” (lines 40-42), and rephrase “built in the popular R and Stan languages” to “implemented using R and Stan” (Abstract, lines 20-21).

Capitalisation: “Markov chain Monte Carlo” (lines 264-270).

Thanks, I fixed all these typos as requested and accept your style suggestions.

Sincerely,
Luka Seamus Wright

February 16, 2026

Luka Seamus Wright
Oceans Institute
University of Western Australia
Perth WA 6009
Australia
☎ +61 8 6488 8760
* luka.wright@research.uwa.edu.au

Dear editor,

I wish to resubmit the manuscript *A model of macroalgal decomposition*. Inspired by the reviewers' constructive comments, I have completely reanalysed each case study and rewritten the manuscript. In addition to conceptual justification, I now provide empirical test results for each modelling choice. I have also added a seventh case study and formally compare the new model to the conventional model for each. To prove the utility of this model in meta-analysis, I now explicitly include several multilevel model applications. My response to reviewers is given in italics below. Updated data and code are available in my open-access repository at github.com/lukaseamus/limbodeco and I encourage you to pass these on to the reviewers. I would also be interested to connect with Reviewer #3 for further discussions.

Reviewer #2:

I think the paper is very well written, detailed, and well explained. The development of the equations is well structured. I think the manuscript is ideal for publication in the journal. I have minor comment:

Line 9: Are you sure this is the world's first mathematical model for macroalgal decomposition?

Thank you. This was also noted by Reviewer #3. While to the best of my knowledge this is indeed the first decomposition model specifically designed with macroalgae in mind, I have changed this statement to "a mathematical model of macroalgal decomposition".

It is necessary to present a conceptual model that shows the novelty of the work, such as the degradation of macroalgae with existing models and with the modification of the model you are presenting.

This was also requested by Reviewer #3. I have fit the conventional model (e^{-kt}) to each example dataset to show that it is ill-fitting in most cases (cf. Figures 2–6 with S2–S6) and formally compare k and τ as well as $t_{1/2}$ and μ (Table 1, 2).

Reviewer #3:

You present a decomposition model that explicitly accommodates early increases or plateaus in normalised detrital mass $m(t)$ by modelling a time-varying rate $k(t)$ that transitions logistically towards a final rate τ . You fit the model in a Bayesian framework, using a beta prime observation model for ratios $m > 0$ (allowing $m > 1$) and a time-varying precision $v(t)$. A range of example datasets is used to demonstrate flexibility and interpretability. The paper is clearly structured, and the illustrative examples and figures are effective in communicating the behaviour of the model and its parameters. The contribution is promising; the main issues below concern (i) parameter identifiability and ecological interpretation, (ii) validation of the observation model and heteroscedastic structure, and (iii) formal comparison with standard approaches to support synthesis-oriented claims.

This is the most competent review I have ever received. Thank you. I encourage you to get in touch for further discussions.

Highlights

The submitted highlights are generally relevant, but I recommend revising them to avoid absolute or difficult-to-substantiate claims. In particular, "First mathematical model of macroalgal decomposition" is hard to verify, and "Can describe any macroalgal decomposition trajectory" is too absolute. Likewise, "Enables meta-analysis..." would read more cautiously as "supports/facilitates meta-analysis", unless a formal demonstration is provided in the

manuscript. I suggest rephrasing the highlights to emphasise verifiable contributions (e.g., a parsimonious time-varying $k(t)$, use of a beta prime likelihood for positive mass ratios including values >1 , and estimation of a terminal decay parameter τ intended to be comparable with the classical exponential $k(t)$).

This was also requested by Reviewer #2. I have rewritten the highlights accordingly.

Major comments

1. Identifiability and interpretation of μ (transition time / photosynthetic 'death')

μ governs the timing of the logistic transition in $k(t)$ and is interpreted as the time at which detrital photosynthesis ceases to matter. In at least one key example, μ appears to be weakly informed by the observation window. In the *Desmarestia anceps* example, the manuscript reports μ on the order of many hundreds of days, whilst the time axis shown in the corresponding figure spans a substantially shorter period. Importantly, the manuscript notes that the experimental duration is not long enough to inform τ for the control treatment, which strengthens the concern that μ may also be poorly identified when the transition sits outside the observed window. Please add (i) explicit identifiability diagnostics (pair plots/correlations for μ - τ and μ - α , trace plots, ESS/Rhat), and (ii) a brief sensitivity analysis for the prior on μ (at least two reasonable priors), reporting impact on τ and on predictive trajectories. Where μ clearly exceeds the experimental duration, please clarify how it should be interpreted ecologically (e.g., 'transition not observed / not estimable within study duration') to avoid over-interpretation.

Identifiability

Correct, μ is not identifiable if the experimental duration is of insufficient length. Pair plots, trace rank plots and prior-posterior plots for all examples are now available in the repository (github.com/lukaseamus/limbodeco/tree/main/Plots) Diagnostic plots are too numerous and large for inclusion as supplementary figures, but I explicitly refer the reader to the repository folder.

Interpretability

*The most intuitive term for μ is "photosynthetic half-life". I was initially reluctant to use this term because mathematically μ is not exactly equivalent to the exponential half-life $t_{1/2} (\ln 2 / k)$ which ecologists are familiar with. After considering your request for better interpretability, I have changed my mind. I believe the term is defensible because it is more intuitive than "photosynthetic death" or "midpoint of photosynthetic decay" and in the case of the underlying fixed intercept logistic function $f(t) = 1/(1 + \exp(5/\mu * (t - \mu)))$, $f(0) = 0.99$ (essentially 1) and $f(\mu) = 0.5$, so μ is directly analogous to $t_{1/2}$. I believe with the identifier "photosynthetic" this term can be safely used and distinguished from half-life in the conventional sense. I already use this term in another publication under review.*

2. Fixed logistic steepness $r = 5/\mu$: justify and test robustness. To reduce non-identifiability you fix the logistic steepness via $r\mu = 5$, yielding a three-parameter form for $k(t)$. This is a pragmatic choice, but it couples the timing of the transition (μ) to its abruptness, which may not match systems where photosynthesis remains stable then collapses relatively quickly. Please, (i) provide a clearer justification for the constant '5' (convention vs biological assumption), and (ii) run a small robustness check on 1-2 representative datasets allowing r to be estimated under a regularising prior (or a hierarchical structure by dataset), and compare predictive performance and parameter stability.

I rewrote the conceptual reasoning behind fixing the logistic intercept at 5, added the alternative constant slope parametrisation as Figure S1 for comparison with Figure 1, and provide performance tests in Table S1. These are discussed in the new section Parameterisation of $k(t)$ in The statistical model.

3. Observation model (beta prime for m): provide posterior predictive checks and a minimal alternative comparison. The beta prime likelihood for the normalised mass m is an elegant choice because it supports m in $(0, \infty)$, allowing $m > 1$. You motivate it via a gamma-ratio argument but also acknowledge that strict independence assumptions are questionable in this context. Request: please include posterior predictive checks focused on (i) tail behaviour for $m > 1$, (ii) behaviour near 0, and (iii) the dispersion pattern over time (in combination with $v(t)$). In addition, please compare beta prime against at least one plausible alternative (e.g., lognormal error on m , or modelling absolute mass with explicit measurement error in $M(0)$) on 1-2 datasets using a predictive metric (e.g., PSIS-LOO/WAIC) to demonstrate that beta prime is not merely convenient but substantively improves predictive adequacy. Minor but

important: please avoid reusing α both as a biological parameter and as a beta prime shape parameter—rename distributional shapes (e.g., a , b) to prevent confusion.

I also added performance tests for the likelihood. These are presented in Table S2 and discussed in the new section Likelihood in The statistical model.

Some common parameter names are unfortunately bound to be repeated in the manuscript, e.g. μ , σ , α , β . There are no other accepted letters for the shape parameters α and β of the beta and beta prime distributions, or for the mean μ , sd σ , variance σ^2 etc. In each case I take care to avoid confusion by redefining what I mean in the given context (e.g. “the sample mean μ ” vs. “the photosynthetic half-life μ ”).

4. Time-varying precision $v(t)$: the imposed form may be too restrictive. You model heteroscedasticity through an exponentially declining precision:

$$v(t) = (\epsilon - \theta) e^{(-\lambda t)} + \theta$$

motivated by minimal uncertainty at $t = 0$ because $m(0) = 1$. While plausible in some contexts, this structure forces dispersion to increase monotonically with time and may not reflect situations where variability depends more on remaining mass, fragmentation events, or normalisation choices (particularly when initial mass is not measured per replicate or when data are reconstructed from summary statistics). Please (i) provide posterior predictive checks demonstrating that the model reproduces the observed spread of m through time, and (ii) include a simple sensitivity comparison to a constant-precision model (or another minimal alternative) to show whether $v(t)$ materially affects inference on μ and τ and improves predictive performance.

I added performance tests for modelling heteroskedasticity. See Table S3 and new section Heteroskedasticity in The statistical model.

5. Formal validation against standard models to support synthesis/meta-analysis claims. A major selling point is that τ is comparable to the standard exponential k , and that ignoring photosynthetic detritus can 'almost invariably' bias decomposition estimates. To support these strong claims (particularly in a synthesis context), the manuscript needs a concise, quantitative model comparison. For 2-3 representative datasets, please fit and compare: (i) the standard exponential model $m(t) = e^{(-k t)}$, (ii) a simple alternative allowing early increase/plateau (even ad hoc), and (iii) the proposed model. Report predictive performance (e.g., PSIS-LOO/WAIC and/or out-of-sample prediction error) and show how k from the exponential compares with τ . Additionally, please include a brief schematic (main text or appendix) showing how τ (and, where identifiable, μ) would be carried forward into a meta-analysis/meta-regression, including guidance for datasets where μ is not identified within the study duration.

This was also requested by Reviewer #2. An initial test comparison is provided in Table S4. I then fit the conventional model (e^{-kt}) to each example dataset to show that it is ill-fitting in most cases (cf. Figures 2–6 with S2–S6) and formally compare k and τ as well as $t_{1/2}$ and μ (Table 1, 2). I now use multilevel models in several examples, providing a full model description in each case, to show how the proposed model would be used in meta-analysis.

6. Use of simulated observations from summary statistics (when raw data are unavailable). Where raw data are unavailable, you simulate observations from reported means/SDs/sample sizes. This can be pragmatic, but it can understate uncertainty (digitisation, rounding, distributional assumptions) and may create pseudo-replication if many points are generated. It would be interesting to propagate this uncertainty (e.g., repeat the simulation several times and show stability of key parameters), or consider a model at the aggregated level that uses means/SDs directly without generating pseudo-data.

I now use a measurement error model to avoid simulating data in the case of Brouwer (1996) and received raw data from the authors of the remaining studies upon renewed request.

Minor comments and editorial suggestions

* The figures illustrating predicted new observations and time-varying $k(t)$ are helpful; consider adding one sentence to selected captions explicitly reiterating ecological interpretation of μ and τ for non-statistical readers.

See comment on the interpretability of μ above. I added the requested detail in the figure captions.

* Where you discuss relevance for blue carbon, please ensure wording distinguishes flux into detrital pools from net long-term sequestration.

As far as I know there is no long-term seaweed carbon sequestration independent of the detrital pool. As I explain in the introduction, only seaweed detritus may be buried as allochthonous carbon. I solely use the term “sequestration” in the introduction. My conclusions relate to decomposition affecting estimates of seaweed blue carbon.

* Reproducibility: you note code/data availability; please also provide a specific repository version (commit/tag) associated with the manuscript results.

I have now published the repo and provide a Zenodo DOI: <https://doi.org/10.5281/zenodo.18654199>.

* Minor editorial corrections (typos and style). The manuscript is generally well written and clearly presented; however, I noted a few small typographical and stylistic issues that could be corrected during revision:

- Introduction (lines 74-75): "disproportionally consequential" → "disproportionately consequential".
- Introduction (line 118): "...to growth rather than decompose..." → "...to grow rather than decompose...".
- Computation (lines 361): "numerically stabler" → "numerically more stable".
- Methods (line 388): "not appliable" → "not applicable".
- Punctuation for readability: "Viewed cumulatively this..." (lines 87-88) → add a comma after "Viewed cumulatively," and "Detritus on the other hand..." (lines 64-65) → "Detritus, on the other hand, ...".
- Style: consider "a relationship that is parameterised..." rather than "...which is parameterised..." (lines 40-42), and rephrase "built in the popular R and Stan languages" to "implemented using R and Stan" (Abstract, lines 20-21).
- Capitalisation: "Markov chain Monte Carlo" (lines 264-270).

Thanks, I fixed all these typos as requested and accept your style suggestions.

Sincerely,
Luka Seamus Wright

Ecological Modelling

A model of macroalgal decomposition

Luka Seamus Wright^{1,2*}

¹Oceans Institute, University of Western Australia, Perth, Australia

²School of Biological Sciences, University of Western Australia, Perth, Australia

*luka.wright@research.uwa.edu.au, luka@wright.it, <https://orcid.org/0000-0002-1273-6256>

Highlights

- Specifically tailored to macroalgal decomposition with detrital photosynthesis
- Parametric nonlinear model with time-variant decomposition rate
- Bayesian statistical implementation with gamma and beta prime likelihoods
- Multilevel and measurement error model capability to facilitate meta-analysis
- Validated against seven empirical case studies on seaweeds including kelps

Ecological Modelling

A model of macroalgal decomposition

Luka Seamus Wright^{1,2*}

¹Oceans Institute, University of Western Australia, Perth, Australia

²School of Biological Sciences, University of Western Australia, Perth, Australia

*luka.wright@research.uwa.edu.au, luka@wright.it, <https://orcid.org/0000-0002-1273-6256>

Abstract

Macroalgae are currently the focus of blue carbon, biological carbon dioxide (CO₂) removal in the ocean. Carbon exported as seaweed detritus is equivalent to ~7.5% of anthropogenic CO₂ emissions annually. Macroalgal blue carbon hinges on the fate of this detritus, so a reliable model of its decomposition is key. However, macroalgal decomposition has been neither consistently nor reliably modelled. Here I present a mathematical model of macroalgal decomposition. Informed by the theory of detrital photosynthesis, it treats detritus as an entity in limbo that can initially either grow or decompose. This is achieved by replacing the rate constant of the conventional exponential decay model with a logistic function of time, mirroring the sigmoidal decay of detrital photosynthesis. The resulting mathematical model has three parameters that describe the initial exponential growth rate, the photosynthetic half-life, and the final exponential decay rate once physiology is factored out. The third parameter is equivalent to the rate constant of the conventional model for dead detritus. Using a Bayesian model with a beta prime likelihood implemented in R and Stan, I show that the statistical implementation performs well given empirical data from a variety of contexts. The presented model can describe any macroalgal decomposition trajectory which enables formal comparison across studies for the first time. I anticipate that this will prove instrumental in resolving the debate around macroalgal CO₂ removal.

Keywords

Bayesian statistics, biogeochemistry, carbon cycle, degradation, kelp, ordinary differential equation

Introduction

In a seminal paper on mathematical ecology, Olson (1963) popularised the ordinary differential equation of leaf litter decomposition

$$\frac{dm}{dt} = -km \quad (1)$$

which describes the rate of change in litter mass m proportional to itself, a relationship that is parameterised by the decay constant k . This reflects our ecological understanding that given a theoretically limitless pool of decomposers, decomposition is proportional to the detrital pool. The equation integrates to

$$\ln \frac{m(t)}{m_0} = -kt \quad (2)$$

and simplifies to the familiar exponential decay function

$$m(t) = m_0 e^{-kt} \quad (3)$$

where $m(t)$ is a function of the initial litter mass m_0 , the decay constant k and time t . Often the half-life $t_{1/2} = \frac{\ln 2}{k}$ (Olson 1963) is reported as a more intuitive measure of decay in terms of units of t . This function has prevailed as the conventional model of decomposition.

Interest in decomposition has been rekindled due to the hope that plants can mitigate anthropogenic climate change by fixing carbon dioxide (CO₂), assimilating it into carbohydrates, and sequestering it in their biomass. Carbon sequestration timescales of decades to centuries are being considered the minimum and ten millennia or longer are preferred (Babiker et al. 2022). Plant biomass can be split into two distinct carbon sinks: living plants and the detrital pool, of which the latter is considered the more promising. Often surprising to plant ecologists, plant detritus amounts to a biomass that is generally comparable and often far greater than that of living plants (Cebrián & Duarte 1995, Wright & Kregting 2023). But more importantly, living plants, even when considering stable or increasing populations, are limited by biomass-density constraints (Creed et al. 2019) and lifespans of typically less than a millennium (Thomas 2013). Detritus, on the other hand, is

believed to have the potential of cumulative sequestration over much longer timescales. Fossil fuels were created via the Carboniferous detrital pathway, mostly of scale trees (Hibbett et al. 2016), and the hope of biological CO₂ removal essentially boils down to re-enacting fossil fuel generation.

Biological CO₂ removal in the ocean is referred to by the shorthand blue carbon (Babiker et al. 2022). Macroalgae have recently moved into the focus of blue carbon (Pessarrodona et al. 2023), primarily due to their substantial global area (Duarte et al. 2022), biomass packing (Creed et al. 2019), productivity (Pessarrodona et al. 2022) and detrital export (Pessarrodona et al. 2023, Filbee-Dexter et al. 2024). Notably, decomposition is disproportionately consequential for macroalgal blue carbon. Unlike their fellow blue carbon ecosystems—seagrass meadows, mangrove forests and salt marshes—seaweed forests cannot accumulate carbon in below-ground biomass and soil. There is thus a set limit for the living seaweed carbon sink which can be estimated as the product of 6.1 million km² global cover (Duarte et al. 2022), $1.2 \pm 3 \text{ kg m}^{-2}$ (mean $\pm$ standard deviation) standing biomass (Creed et al. 2019) and $28 \pm 4.5 \%$ carbon content (Pessarrodona et al. 2022). This yields $0.34 \pm 0.87 \text{ kg C m}^{-2}$ and $2.1 \pm 5.3 \text{ Gt C globally}$, or only $19 \pm 49 \%$ of annual anthropogenic CO₂ emissions (Friedlingstein et al. 2025). However, seaweeds are thought to assimilate $0.32 \pm 0.54 \text{ kg C m}^{-2} \text{ y}^{-1}$ (Pessarrodona et al. 2022) and 1.3 Gt C y^{-1} globally (Duarte et al. 2022) of which $61 \pm 4.2 \%$ is exported as detritus (Pessarrodona et al. 2023). Hence $0.19 \pm 0.33 \text{ kg C m}^{-2} \text{ y}^{-1}$ and $0.81 \pm 0.055 \text{ Gt C y}^{-1}$ globally, or $7.5 \pm 0.51 \%$ of annual anthropogenic CO₂ emissions, may enter the detrital pool yearly. Note that these global standard deviations are underestimates because Duarte et al. (2022) report no uncertainty. Viewed cumulatively, this is clearly the more substantial carbon sink. Decomposition alone therefore determines whether macroalgal blue carbon is significant or not, as is illustrated by the central role it takes in the latest macroalgal blue carbon models (Filbee-Dexter et al. 2024). Despite this, macroalgal decomposition remains comparatively poorly understood.

Phycologists have never been able to properly model macroalgal decomposition. Decomposition trajectories rarely match Equation 3 reasonably well (e.g. de Bettignies et al. 2020, Pedersen et al. 2021). Most studies have therefore either refrained from fitting a parametric model (e.g. Birch et al. 1983, Vandendriessche et al. 2007, Hamersley et al. 2015, Frontier et al. 2021) or most commonly fit linear models (e.g. Bedford & Moore 1984, Brouwer 1996, Krumhansl & Scheibling 2012, Filbee-Dexter et al. 2022, Frontier et al. 2022,

Wright et al. 2022). Meta-analyses of macroalgal k (Equation 3) are numerous (Enríquez et al. 1993, Banta et al. 2004, Cebrián & Lartigue 2004, Filbee-Dexter et al. 2024, Kennedy & Blain 2025, White & Norkko 2025) but have generally assumed Equation 3 by default and not assessed the validity of this model in the macroalgal context. It is evident that these meta-analyses are often hampered by the complexity and diversity of macroalgal decomposition trajectories (Kennedy & Blain 2025, White & Norkko 2025). The only serious attempts at modelling diverse trajectories have invariably invoked the logistic function alongside the exponential (Bourguès et al. 1996, Kennedy & Blain 2025). This defeats the purpose of a model since logistic and exponential rate constants cannot be formally compared. Kennedy & Blain (2025) realised this when attempting to compare $t_{1/2}$ across exponential and logistic models. While $t_{1/2}$ for the latter is best estimated as the inflection point of the sigmoid, they ultimately resorted to forcing an exponential model to derive this parameter (Kennedy & Blain 2025), which undermines fitting the logistic model in the first place.

We now know what causes the observed diversity in macroalgal decomposition trajectories. Early phycologists noticed the need to “pre-kill” macroalgal detritus (Josselyn & Mathieson 1980, Rice & Tenore 1981, Birch et al. 1983, Smith & Foreman 1984, Brouwer 1996) if it was to behave according to the assumptions of the conventional model (Equation 3). Recently it became evident that this is because excised macroalgal tissue can continue to photosynthesise for months (de Bettignies et al. 2020), which may allow macroalgal detritus to grow rather than decompose over the short term (Frontier et al. 2021, Pedersen et al. 2021, Filbee-Dexter et al. 2022, Frontier et al. 2022, Wright et al. 2022). To avoid semantic confusion, I hereinafter use “detritus” for all detached material and “detritus *sensu stricto*” for detached *and* dead material. There is now extensive investigation into what has since been termed detrital photosynthesis (Frontier et al. 2021, Wright & Foggo 2021, Frontier et al. 2022, Wright et al. 2022, Wright & Kregting 2023, Wright et al. 2024) and macroalgae have been found to be unique among plants in having detritus that photosynthesises to an extent that is likely to influence its decomposition (Wright 2026b). This knowledge enables the creation of a model that accurately describes macroalgal decomposition by taking into account the effect of detrital photosynthesis. It is probable that, ignorant of detrital photosynthesis, the conventional model has substantially underestimated macroalgal decomposition, leading to overestimated seaweed blue carbon. To remedy this I here present a nonlinear model with three parameters that is tailored to the unique case of macroalgal decomposition. First I describe its mathematical form, then I implement it as a Bayesian

statistical model, and finally I validate it against a variety of empirical examples, comparing its predictive performance to that of the conventional model for each example.

The mathematical model

...tossed and swept like dead leaves from one spot to another, never resting, never giving back their goodness to the earth, never fully dead but in some vast limbo between life and extinction, tossing and tumbling without end... — Philip Pullman (1994) *The Tin Princess*

Macroalgal detritus is essentially in limbo. It may grow or decay, but not indefinitely. The goal is to model this temporary limbo. There are several possible approaches to describing this mathematically. As mentioned above, the logistic function has been the only attempt so far, but its rate constant is not comparable to that of Equation 3 and it cannot describe the full spectrum of possible decomposition trajectories (e.g. an increase followed by a decrease). Hence we must dismiss it. Another approach could be a composite function of a saturating function (e.g. rectangular hyperbola, hyperbolic tangent, exponential) and Equation 3. This may describe all possible trajectories but is somewhat of an artificial construct and more importantly would have more parameters than are practical. The best way forward is to stick with our trusty Equation 3 but rebuild it from its differential form (Equation 1). Instead of treating k as constant, we can let it be a function of time:

$$\frac{dm}{dt} = k(t)m \quad (4)$$

To my knowledge, time-variant exponential decay has only been used twice in the context of decomposition, by Rovira & Rovira (2010) to describe the variety of terrestrial leaf litter decomposition trajectories and by Manojlovic (2021) to model seagrass decomposition (Trevathan-Tackett et al. 2020). Rovira & Rovira (2010) chose the form of $k(t)$ based on context while Manojlovic (2021) used a random walk, the simplest autoregression which defines k only in terms of k at the previous timepoint plus normal error:

$$k(t) \sim N(k(t - \Delta t), \sigma) \quad (5)$$

where σ is the standard deviation of the normal distribution around the mean $k(t - \Delta t)$ and determines the k step size of the random walk. The timestep size Δt is typically set to 1.

While this approach is the most flexible, it requires sophisticated Bayesian state space modelling with an ordinary differential equation solver. Manojlovic (2021) used the Bayesian language LibBi (Murray 2015) and to my knowledge there is currently no way to implement this in the more popular Stan language (Carpenter et al. 2017) (Manojlovic, personal communication, discourse.mc-stan.org/t/random-walk-in-rstan/34308). Furthermore, being the simplest type of order 1 autoregressive process, a random walk's only parameter is σ which has no inherent directionality and precludes informative prediction beyond the range of the data. Finally, no comparison can be made between studies beyond overlaying plots of $k(t)$, and formal comparison with studies that use the conventional model (Equation 3) is out of question. The last two limitations are also inherent to non-parametric nonlinear models. A model with informative parameters which can be formally compared is the way forward. Hence I here opt for the path laid out by Rovira & Rovira (2010).

Now the most important task is to find an appropriate function to describe time-variant k . Since the underlying process affecting k is detrital photosynthesis, it makes sense to start there. The change in detrital photosynthesis with time following detachment, that is detrital age, has mostly been described using various linear models (de Bettignies et al. 2020, Frontier et al. 2021, Wright & Foggo 2021, Wright et al. 2022, Wright & Kregting 2023, Wright et al. 2024). There is no reasonable intrinsic mechanism causing photosynthesis to increase following detachment. Although photosynthesis is often maintained near baseline levels it usually does not exceed this baseline. When an increase is observed it is likely due to an extrinsic factor such as a change in light environment (Frontier et al. 2021, Wright & Kregting 2023) or measurement error. Eventually photosynthesis decays to a minimum beyond which there logically is no further decline. Voilà, logistic decay! Logistic decay with detrital age was inadvertently arrived at in two independent cases: a beta regression describing F_v/F_m (Frontier et al. 2022) and a binomial regression describing the probability of photosynthesis (Wright et al. 2024). Inspired by these cases, I have formalised logistic decay as the model of detrital photosynthesis (Wright 2026b).

In practice, the slope and intercept of the linear term x in the standard logistic function $\frac{1}{1 + e^{-x}}$ are rarely identifiable. Non-identifiability in a linear function is typically combatted by centring the predictor, but this is not sensible when the predictor is time. Instead, the solution is to replace one of the parameters with a constant. Decay functions have a known initial

value. Thus, as with exponential decay, logistic decay must have a fixed intercept. Without a fixed intercept, there is no reference point and the slope or decay rate is no longer directly comparable across cases (e.g. Wright et al. 2024). Take the concept of $t_{1/2}$ which is defined as the time when half of the initial value remains. Without a constant intercept, $t_{1/2}$ is meaningless. The problem is that unlike the exponential function, the standard logistic function is asymptotic at both bounds, so cannot take a value of 1. I believe the closest sensible value to be 0.99, which neatly translates to an intercept of 5 since $\frac{1}{1 + e^{-5}} \approx 0.99$. Because $\frac{1}{1 + e^0} = 0.5$ is approximately half of 0.99, the time when $x = 0$ can be interpreted as the logistic half-life, analogous to the exponential half-life. In Wright (2026b), I have therefore described detrital photosynthesis (p) as

$$p(t) = \frac{\alpha + \tau}{\frac{5(t - \mu)}{1 + e^{-\mu}}} - \tau \quad (6)$$

where α and τ may be read as the maximal CO₂ assimilation and production of the plant and its detritus respectively and μ is the sigmoid inflection point, which I have termed the photosynthetic half-life (Wright 2026b). The naming of parameters is inspired by the story of the golem (McElreath 2020). As an entity in limbo, the golem's state is controlled by the word חַמַּל (graecised $\tau\mu\alpha$), symbolising life ($\aleph$), transition ($\beth$) and death ($\aleph$). In the form of my parameters, these letters similarly control the transition of detritus between physiological states.

In addition to modelling detrital photosynthesis, $p(t)$ can be used to describe time-variant k : $k(t) = p(t)$. I provide empirical support for this decision by comparing the performance of $p(t)$ with alternative parameterisations of the logistic function (see *Parameterisation of $k(t)$* in *The statistical model*). Inserting Equation 6 in place of $k(t)$ in Equation 4 yields

$$\frac{dm}{dt} = m \left(\frac{\alpha + \tau}{\frac{5(t - \mu)}{1 + e^{-\mu}}} - \tau \right) \quad (7)$$

where α is now the initial exponential growth rate (aka relative or specific or instantaneous growth rate) of detritus at detachment and τ is the maximal exponential decay rate of detritus *sensu stricto* as $t \rightarrow \infty$. μ remains the photosynthetic half-life. The influence of each parameter on the trajectory of time-variant k is visualised in Figure 1 (cf. Figure S1).

This equation could be used directly with statistical software that allows solving ordinary differential equations on the fly. But there is a simpler, generally applicable solution. Rovira & Rovira (2010) have shown that the logistic function can be integrated alongside the ordinary differential equation to yield a function like Equation 3. Since we know our function is bounded by $t = 0$, we use an accumulation function. The definite integral of Equation 6 is

$$\int_0^t p(t)dt = t\alpha + \frac{\mu(\alpha + \tau)}{5} \ln \frac{1 + e^{\frac{5(t-\mu)}{\mu}}}{1 + e^{-5}} \quad (8)$$

Thus when Equation 7 is integrated as a whole, we get the final form

$$m(t) = m_0 e^{t\alpha + \frac{\mu(\alpha + \tau)}{5} \ln \frac{1 + e^{\frac{5(t-\mu)}{\mu}}}{1 + e^{-5}}} \quad (9)$$

Decomposition data are best analysed relative to initial mass, i.e. after normalisation. When $m_0 = 1$, the function simplifies to one with only three parameters, the now familiar α , μ and τ . The influence of each parameter on the shape of the curve m describes over time is illustrated in Figure 1 (cf. Figure S1). α , μ and τ may be formally compared within and between studies, including those that used the conventional model (Equation 3). τ is the exponential decay rate of detritus *sensu stricto* once detrital photosynthesis has been factored out. Since conventional decomposition theory assumes that physiology does not play a role in detrital dynamics, k in Equation 3 also represents the exponential decay rate of detritus *sensu stricto*. Thus τ may be directly compared to k to assess the influence of detrital photosynthesis on decomposition. Similarly, μ and $t_{1/2}$ have the same units and may be contrasted (Wright 2026b). I make these formal comparisons in *Examples* below. In the next section I turn Equation 9 into a fully functional statistical model.

The statistical model

Statistical framework

Maximum likelihood solutions to nonlinear modelling invariably demand the tedious task of guessing starting values. Rovira & Rovira (2010) followed this approach using the Levenberg-Marquardt algorithm which is more robust to poor starting values than the usual Gauss-Newton algorithm. Despite this advantage, they still needed to manually approximate

the starting values before passing them to the algorithm. Bayesian inference using Markov chain Monte Carlo (MCMC) is more flexible in that there is no practical difference between fitting linear and parametric nonlinear models (McElreath 2020). Rather than starting values which can make or break the model, a prior probability distribution is provided, which at worst can lead to inefficient sampling and uncertain prediction. With the increase in computing power, every laptop can now run MCMC. For these and other reasons Bayesian inference is quickly becoming the *status quo* in applied statistics across disciplines (McElreath 2020). All models in this study are therefore Bayesian and described using standard notation *sensu* McElreath (2020).

The most popular statistical language is R (R Core Team 2022) and the most popular Bayesian language is Stan (Carpenter et al. 2017) which integrates nicely with R through various packages. Hence I developed the statistical model within this joint framework. Specifically, I use R v4.5.2 with the tidyverse package family v2.0.0 (Wickham et al. 2019) and here v1.0.2 (Müller 2025) within the integrated development environment RStudio v2026.01.0+392 (RStudio Team 2022). All models in this study are written in Stan and sampled with 8 Markov chains in CmdStan v2.38.0 (Lee et al. 2017) via the R interface cmdstanr v0.9.0 (Gabry et al. 2024) with the aid of tidybayes v3.0.7 (Kay 2024). To assess model quality, I check $\hat{R}$ convergence scores and effective sample sizes and visually scrutinise trace rank and pair plots with bayesplot v1.14.0 (Gabry et al. 2019). All reported results are posterior probabilities and derived central tendencies and intervals, which are translated into figures and tables using ggh4x v0.3.1 (van den Brand 2025), patchwork v1.3.2 (Pedersen 2025) and officer v0.7.1 (Gohel et al. 2025). See github.com/lukaseamus/limbodeco for further details.

Parameterisation of $k(t)$

As explained above, there is a strong conceptual reason for choosing Equation 6 to describe time-variant k . Nonetheless, I empirically tested the performance of Equation 6 against two alternative parameterisations of the logistic function: one where the slope instead of the intercept is a constant, and another where neither intercept nor slope are constants. Unlike the intercept, the slope has no reasonable *a priori* value. For simplicity I therefore chose a value of 1 based on simulation (Figure S1). In contrast to the constant intercept parameterisation (Equation 6) which logically slows photosynthetic decay for large μ (Figure 1), the constant

slope parameterisation allows rapid photosynthetic decay independent of μ (Figure S1). This is less intuitive but agrees better with observations of photosynthesis seemingly being unperturbed following detachment but decaying rapidly at some later time (de Bettignies et al. 2020, Frontier et al. 2021, 2022). The full parameterisation with no constants is equivalent to the original naïve beta and binomial models (Frontier et al. 2022, Wright et al. 2024).

Each parameterisation of $k(t)$ was tested on the two largest, most time-resolved decomposition datasets (Vandendriessche et al. 2007, Frontier et al. 2022). α , μ and τ were estimated globally for the smaller dataset ($n = 252$, Frontier et al. 2022) and stratified by replicate for the larger dataset ($n = 1651$, Vandendriessche et al. 2007), in both cases using a normal likelihood with homogenous variance and identical priors across parameterisations. I compared convergence in terms of $\hat{R}$, effective sample size and trace rank plots, predictive performance in terms of leave-one-out cross-validation computed with Pareto-smoothed importance sampling (Vehtari et al. 2017), and identifiability with prior–posterior and pair plots. All statistics are reported in the supplement and plots are available at github.com/lukaseamus/limbodeco/tree/main/Plots. For both datasets, using Equation 6 as $k(t)$ results in marginally better convergence (Table S1). Equation 6 also yields the highest predictive accuracy for the small dataset, albeit the lowest for the large dataset, and the lowest level of overfitting in both cases (Table S1). As discussed in *The mathematical model*, the logistic slope and intercept are usually non-identifiable. Without a constant, the prior therefore essentially determines either slope or intercept. For the smaller dataset, the constant slope parameterisation displays stronger posterior correlation and unstable posterior distributions. While for large datasets slope and intercept may be identifiable and the difference between parameterisations marginal, Equation 6 is not only conceptually but also empirically the strongest parameterisation of $k(t)$ across all cases.

Likelihood function

Several hierarchical functions are needed to describe a full statistical model. The one to rule them all is the likelihood function. It describes the probability distribution of the data. Most likelihoods are described by two parameters, usually the expected value and spread. The mathematical model (Equation 9) describes the expected value of the likelihood. But what likelihood?

Convention would favour the normal distribution but there is almost always a more fitting choice. Normalised decomposition data are ratios of final to initial mass that are positive real numbers and can exceed 1 due to growth caused by detrital photosynthesis. Several distributions are restricted to positive real numbers, such as the popular gamma and lognormal distributions. But perhaps we can do better by thinking generatively. Mass itself can only take positive real numbers and may be thought of as the sum of cells, growth increments etc. Thus the gamma distribution describes mass well. The ratio of two gamma distributions is a beta prime distribution. Technically this is only true if both gamma distributions share the same rate or scale, i.e. size of summand, and are independent. We can assume the former because both measures of mass can be broken down into the same components, e.g. cells. It is difficult to assume the latter because the fundamental assumption of the model is that the change in mass depends on mass (Equation 1). Nonetheless, beta prime remains the best approximation of the true data distribution I can think of.

I empirically tested the validity of this choice by comparing the beta prime likelihood to the normal, lognormal and gamma likelihoods with homogenous variance and identical priors. As before, I compare convergence, predictive performance and identifiability across the two datasets. Convergence is similar across likelihoods (Table S2). Predictive performance is highest for the normal likelihood given the small dataset while the gamma and beta prime likelihoods perform best given the large dataset; the lognormal likelihood performs worst in both cases (Table S2). Since the gamma and beta prime likelihoods perform similarly and obey $m > 0$, either is a reasonable choice.

Heteroskedasticity

If Equation 9 describes the expected value of the likelihood, what describes its spread? Under the typical assumption of homogeneity of variance, this likelihood parameter is usually estimated globally. However, variance rarely being homogenous, it is sensible to model it as a function of the predictors.

In the case of normalised decomposition, $m_0 = 1$ is predetermined, meaning there is no uncertainty at $t = 0$. But with time comes uncertainty and we expect variance to increase. Sometimes remaining mass is calculated relative to mean rather than sample-specific initial mass so there is uncertainty at $t = 0$ (e.g. Brouwer 1996, Krumhansl & Scheibling 2012), but even then it naturally increases with time. The beta prime distribution, typically

parameterised by the two shape parameters α and β , is most intuitively parameterised in terms of mean $\mu = \frac{\alpha}{\beta - 1}$ and precision $\nu = \beta - 2$, where variance $\sigma^2 = \frac{\mu(1 + \mu)}{\nu}$ (Bourguignon et al. 2021). To reparameterise the likelihood, α and β need to be expressed in terms of μ and ν as $\alpha = \mu(1 + \nu)$ and $\beta = 2 + \nu$. Since ν is inversely related to σ^2 , it is expected to decrease over time but cannot violate $\nu > 0$ so an appropriate function is

$$\nu(t) = (\epsilon - \theta)e^{-\lambda t} + \theta \quad (10)$$

where ϵ is ν at $t = 0$, λ is the rate of exponential decay in ν , and θ is ν as $t \rightarrow \infty$, inspired by its representation of Thánatos. This decay function can be used to describe precision in any likelihood while a saturating function would be the logical choice for describing variance.

The improvement of modelling heteroskedasticity over assuming homoskedasticity was empirically tested for the beta prime and gamma likelihoods. The gamma distribution may be parameterised in terms of scale or rate. The rate is inversely related to scale and variance and can therefore be described by Equation 10. In all cases the heteroskedastic models outperform the homoscedastic ones with the gamma likelihood performing marginally better (Table S3).

Optimal model

Empirically speaking, the gamma and beta prime likelihoods are effectively interchangeable with gamma performing better in some cases. Conceptually I prefer the beta prime distribution and use it as the likelihood hereinafter. Combining the beta prime likelihood and Equations 9 and 10 yields the full statistical model

$$m_i \sim \text{Beta}'(\bar{m}_i(1 + \nu_i), 2 + \nu_i) \quad (11)$$

$$\bar{m}_i = e^{t_i \alpha + \frac{\mu(\alpha + \tau)}{5} \ln \frac{1 + e^{\frac{5(t_i - \mu)}{1 + e^{-5}}}}{1 + e^{-5}}}$$

$$\nu_i = (\epsilon - \theta)e^{-\lambda t_i} + \theta$$

$$\alpha, \mu, \tau, \epsilon, \lambda, \theta \sim \text{Priors}$$

where m_i is the ratio of final to initial mass observed at t_i , $\bar{m}_i$ and ν_i being the mean and precision of m_i . $\bar{m}_i$ and ν_i are described by Equations 9 and 10, which for readability I will hereinafter refer to as $m(t, \alpha, \mu, \tau)$ and $\nu(t, \epsilon, \lambda, \theta)$.

Bayesian inference requires a prior probability distribution for each parameter (McElreath 2020). All parameters except α are restricted to positive values and must therefore be assigned a prior that respects this constraint. α may have a normal prior allowing it to take positive (initial growth) or negative (initial decomposition) values but in some cases it may be desirable to enforce initial growth by assigning a prior with that constraint. Beyond simple constraints, priors must be chosen based on existing empirical and/or conceptual knowledge and simulation (McElreath 2020). α , μ and τ are biologically meaningful and can thus be strongly informed by the literature. ϵ , λ and θ are more abstract. Based on simulation I found $\epsilon \sim \text{Gamma}(4, 10^{-4})$, $\lambda \sim \text{Exponential}(1)$ and $\theta \sim \text{Gamma}(4, 0.008)$ to be sensible.

Multilevel modelling is the state of the art because it allows estimation of hyperparameters and prediction for unobserved categories, deals with data sparsity, sampling bias and pseudo-replication, and regularises spurious effects (McElreath 2020). Equation 11 can easily be turned into a multilevel model by partially pooling some or all parameters and can thus be used in meta-analysis. For μ , τ , ϵ , λ and θ , partial pooling is best achieved in log space because the normal distribution is easiest to sample when nested and allows non-centred parameterisation.

In the remainder of this paper, I show that for most macroalgal decomposition data Equation 11 outperforms the conventional model

$$\begin{aligned} m_i &\sim \text{Normal}(\bar{m}_i, \sigma) \\ \bar{m}_i &= e^{-kt_i} \\ k, \sigma &\sim \text{Priors} \end{aligned} \tag{12}$$

where $\bar{m}_i$ is described by Equation 3. A normal likelihood and homogenous variance are conventional assumptions, so they form my baseline. Given the small and large test datasets, Equation 11 respectively has similar and better predictive accuracy than Equation 12 (Table S4). In *Examples*, I apply Equations 11 and 12 to five ecological effects in seven published studies (Bourguès et al. 1996, Brouwer 1996, Vandendriessche et al. 2007, Hamersley et al. 2015, de Bettignies et al. 2020, Frontier et al. 2021, 2022). I either received data from the lead author (Vandendriessche et al. 2007, Hamersley et al. 2015, de Bettignies et al. 2020,

Frontier et al. 2021, 2022) or digitised figures using WebPlotDigitizer v4.6 (Rohatgi 2022). Parameter estimates for both models are presented in Table 1 and contrasts between comparable parameters across models and are shown in Table 2.

Sampling efficiency

There are some final considerations for Equation 11 that are purely computational. Stan uses Hamiltonian Monte Carlo to explore probability space. Some of the expressions I have used in Equations 9 and 10 are easier to read but not friendly to the MCMC sampler. These can be improved for computational efficiency using the quotient and product rules of logarithms:

$$m(t) = e^{t\alpha + \frac{\mu(\alpha + \tau)}{5}(\ln(1 + e^{\frac{5(t-\mu)}{\mu}}) - \ln(1 + e^{-5}))} \quad (13)$$

$$v(t) = e^{\ln(\epsilon - \theta) - \lambda t} + \theta \quad (14)$$

The $\ln(1 + e^x)$ expressions in Equation 12 are further simplified in Stan using the wrapper function `log1p_exp` which is numerically more stable. An equivalent function does not exist in R but can be written as `log1p_exp <- function(x) ifelse( x > 0 , x + log1p( exp(-x) ) , log1p( exp(x) ) )`. Note that Equation 10 performs better than Equation 14 in some cases, e.g. when prior variance for ϵ and/or θ is large, which results in $\epsilon \leq \theta$ at initialisation, Equation 14 causes the log probability to evaluate to $-\infty$.

Examples

Dead or alive

Brouwer (1996) provides a nice example of detrital photosynthesis influencing decomposition of the feathery brown seaweed *Desmarestia anceps* at Signy Island in Antarctica. She pre-killed some of the detritus by heating it to 50°C for 10–15 min prior to *in situ* decomposition. The resulting effect is remarkable but Brouwer (1996) was unable to quantify it due to lack of a suitable model. She only reports an analysis of variance F statistic for change over time in both treatments and describes the divergence in decomposition trajectories. Remaining mass at each timepoint is summarised as mean $\pm$ standard deviation of five samples. To correctly account for summarised uncertainty, I use a measurement error model which gives less predictive weight to means with larger standard deviations (McElreath 2020). Since I assume that these data follow a beta prime distribution, it follows

that measurement error is also beta prime distributed. Directly parameterising beta prime error in terms of sample mean μ and sample standard deviation σ is hard on the MCMC sampler. I therefore pre-calculate the sample precision $v = \frac{\mu(1+\mu)}{\sigma^2}$. The resulting model is

$$\begin{aligned}
 m_{\text{obs},i} &\sim \text{Beta}'(m_i(1 + v_{\text{obs},i}), 2 + v_{\text{obs},i}) \\
 m_i &\sim \text{Beta}'(\bar{m}_i(1 + v_i), 2 + v_i) \\
 \bar{m}_i &= m(t_i, \alpha_{E_i}, \mu_{E_i}, \tau) \\
 v_i &= v(t_i, \epsilon, \lambda_{E_i}, \theta) \\
 \alpha_E &\sim \text{Normal}(-0.01, 0.005) \\
 \mu_E &\sim \text{Gamma}\left(\frac{150^2}{100^2}, \frac{150}{100^2}\right) \\
 \tau &\sim \text{Gamma}\left(\frac{0.1^2}{0.05^2}, \frac{0.1}{0.05^2}\right) \\
 \epsilon &\sim \text{Gamma}(4, 10^{-4}) \\
 \lambda_E &\sim \text{Exponential}(1) \\
 \theta &\sim \text{Gamma}(4, 0.008)
 \end{aligned} \tag{15}$$

where m_i is the latent variable of unobserved true values, subscript obs indicates observed variables and E is the experimental treatment variable. τ and θ are not stratified by E since the experimental duration is not long enough to identify these parameters for the control treatment and it is reasonable to assume that decomposition rates converge once detrital photosynthesis has ceased. Predictions by this model are visualised in Figure 2 (cf. Figure S2). Predictions for the mean and new observations of m are respectively denoted $\bar{m}$ and $\tilde{m}$. Summaries of parameter distributions are given in Table 1.

The model suggests that pre-killed *D. anceps* detritus decomposes much faster than the control (Figure 2). Untreated detritus has an initial relative growth rate of 0.068 ± 0.032 % day⁻¹ (mean $\pm$ standard deviation, $P_{\alpha > 0} = 0.99$) while pre-killed detritus has an initial decay rate of 1.3 ± 0.5 % day⁻¹ ($P_{\alpha < 0} = 0.99$) (Table 1). The photosynthetic half-life of *D. anceps* detritus is 585 ± 100 days and without detrital photosynthesis it decomposes at a rate of 5.6 ± 1.5 % day⁻¹ (Table 1). The conventional model (Figure S2) underestimates decomposition by two orders of magnitude; pre-killing detritus ameliorates this difference, but only partly (Table 1, 2). These results suggest that detrital photosynthesis confounds the conventional

model and even after experimental removal of physiology, it may not yield reliable predictions for detritus *sensu stricto*.

Senescence

Hamersley et al. (2015) and de Bettignies et al. (2020) both studied the effect of tissue senescence on kelp decomposition. Hamersley et al. (2015) collected attached fresh and senescent fronds of the giant kelp *Macrocystis pyrifera* at Catalina Island in California along with naturally detached wrack and raft material which they regarded as detritus *sensu stricto*. de Bettignies et al. (2020) collected old and new blades from above and below the abscission zone of the boreal kelp *Laminaria hyperborea* at Roscoff in Brittany. Their old blade thus represents attached senescing tissue that the plant is in the process of abscising. Hamersley et al. (2015) refrained from fitting a model but de Bettignies et al. (2020) fit Equation 3 using maximum likelihood estimation with the Gauss-Newton algorithm, which passably describes their published data from 2017, and report $k = 3.66$ and $1.07 \% \text{ day}^{-1}$ for old and new blades. Notably, they excluded data from a second experiment on old blades at the same site the following year (2018) which cannot be described by Equation 3. With the permission of the lead author, I include these data here to show that they present no problem for the proposed macroalgal decomposition model. The model for Hamersley et al. (2015) is

$$m_i \sim \text{Beta}'(\bar{m}_i(1 + \nu_i), 2 + \nu_i) \quad (16)$$

$$\bar{m}_i = m(t_i, \alpha_{A_i}, \mu_{A_i}, \tau_{A_i})$$

$$\nu_i = \nu(t_i, \epsilon, \lambda, \theta)$$

$$\alpha_A \sim \text{Normal}(\bar{\alpha}, \sigma_\alpha)$$

$$\ln \mu_A \sim \text{Normal}(\overline{\ln \mu}, \sigma_{\ln \mu})$$

$$\ln \tau_A \sim \text{Normal}(\overline{\ln \tau}, \sigma_{\ln \tau})$$

$$\bar{\alpha} \sim \text{Normal}(0, 0.01)$$

$$\overline{\ln \mu} \sim \text{Normal}(\ln 10, 0.5)$$

$$\overline{\ln \tau} \sim \text{Normal}(\ln 0.2, 0.5)$$

$$\sigma_\alpha \sim \text{Normal}_{0,\infty}(0, 0.01)$$

$$\sigma_{\ln \mu}, \sigma_{\ln \tau} \sim \text{Normal}_{0,\infty}(0, 0.5)$$

$$\epsilon, \lambda, \theta \sim \text{Priors as in Equation 14}$$

where all parameters describing $\bar{m}_i$ are partially pooled across tissue age A . Predictions are shown in Figure 3a (cf. Figure S3a). The model for de Bettignies et al. (2020) is

$$\begin{aligned}
 m_i &\sim \text{Beta}'(\bar{m}_i(1 + \nu_i), 2 + \nu_i) \\
 \bar{m}_i &= m(t_i, \alpha_{A_i}, \mu_{A_i}, \tau_{A_i}) \\
 \nu_i &= \nu(t_i, \epsilon, \lambda, \theta) \\
 \alpha_{A_i} &= \delta_{A_i} - \tau_{A_i} \\
 \ln \delta_A &\sim \text{Normal}(\overline{\ln \delta}, \sigma_{\ln \delta}) \\
 \ln \mu_A &\sim \text{Normal}(\overline{\ln \mu}, \sigma_{\ln \mu}) \\
 \ln \tau_A &\sim \text{Normal}(\overline{\ln \tau}, \sigma_{\ln \tau}) \\
 \overline{\ln \delta} &\sim \text{Normal}(\ln 0.01, 0.2) \\
 \overline{\ln \mu} &\sim \text{Normal}(\ln 30, 0.3) \\
 \overline{\ln \tau} &\sim \text{Normal}(\ln 0.02365, 0.5) \\
 \sigma_{\ln \delta} &\sim \text{Normal}_{0,\infty}(0, 0.2) \\
 \sigma_{\ln \mu} &\sim \text{Normal}_{0,\infty}(0, 0.3) \\
 \sigma_{\ln \tau} &\sim \text{Normal}_{0,\infty}(0, 0.5) \\
 \epsilon, \lambda, \theta &\sim \text{Priors as in Equation 14}
 \end{aligned} \tag{17}$$

where parameters are partially pooled as in Equation 16, but with the added constraint $-\alpha < \tau$ since decomposition rate cannot conceivably decrease according to the theory of detrital photosynthesis. The prior for τ is centred on the mean of reported k . Predictions are shown in Figure 3b (cf. Figure S3b).

Both models suggest that senescent fronds tend to decompose faster than fresh ones (Figure 3), but this effect is variable and not as strong as for pre-killed tissue (cf. Figure 2). Detritus of *M. pyrifera* does not grow regardless of tissue age ($P_{\alpha < 0} = 0.7\text{--}0.74$). New *L. hyperborea* blades grow at detachment ($P_{\alpha > 0} = 0.79$) but old blades either grow ($P_{\alpha > 0} = 0.67$) or decompose ($P_{\alpha < 0} = 1$) (Table 1). Photosynthetic half-life of fresh *M. pyrifera* is 4.1 ± 2 days ($P_{\Delta\mu > 0} = 0.95$) and 5.4 ± 1.2 days ($P_{\Delta\mu > 0} = 1$) longer than for detached and senescent fronds (Figure 3a, Table 1). But detrital photosynthesis of fresh *L. hyperborea* may cease earlier ($\Delta\mu = 55 \pm 25$ days, $P_{\Delta\mu > 0} = 0.99$) than for senescent tissue (Figure 3b, Table 1). These results suggest that senescent kelp typically has impaired photosynthesis and decomposes faster as a consequence. However, the viability of apparently senescent tissue may be highly

unpredictable, as is evidenced by the change in decomposition of old *L. hyperborea* blades from one year to the next. The conventional model (Figure S3) again underestimates decomposition in all cases and conventional estimates are most accurate when detrital photosynthesis is weakest, as for senescent *L. hyperborea* in the 2017 experiment (Table 1, 2).

Season

Bourguès et al. (1996) provide perhaps the only data of macroalgal decomposition across all seasons. They studied decomposition of the ephemeral green seaweed *Ulvaria obscura* in the intertidal Arcachon Bay in France. Bourguès et al. (1996) collected fresh *U. obscura* but buried it below 1 cm of sediment where anoxia apparently persisted and there clearly was no light for detrital photosynthesis. Given these experimental conditions, it would be surprising to see decomposition affected by physiology. Yet Bourguès et al. (1996) were faced with the challenge of having to model decomposition trajectories of sigmoidal appearance for which they chose the logistic function. Excised macroalgal tissue can survive exceptional periods of darkness (Wright et al. 2024) so it is possible that saprotrophs were initially inhibited by chemical defence despite the absence of photosynthesis. Even if this was not the case because anoxia immediately killed the seaweed, and the model therefore is conceptually wrong, it is still useful because it avoids having to fit fundamentally incomparable models as Bourguès et al. (1996) have done. Thus the model is

$$m_i \sim \text{Beta}'(\bar{m}_i(1 + v_i), 2 + v_i) \quad (18)$$

$$\bar{m}_i = m(t_i, \alpha_{M_i}, \mu_{M_i}, \tau_{M_i})$$

$$v_i = v(t_i, \epsilon, \lambda_{M_i}, \theta_{M_i})$$

$$\ln \alpha_M \sim \text{Normal}(\overline{\ln \alpha}, \sigma_{\ln \alpha})$$

$$\ln \mu_M \sim \text{Normal}(\overline{\ln \mu}, \sigma_{\ln \mu})$$

$$\ln \tau_M \sim \text{Normal}(\overline{\ln \tau}, \sigma_{\ln \tau})$$

$$\ln \lambda_M \sim \text{Normal}(\overline{\ln \lambda}, \sigma_{\ln \lambda})$$

$$\ln \theta_M \sim \text{Normal}(\overline{\ln \theta}, \sigma_{\ln \theta})$$

$$\overline{\ln \alpha} \sim \text{Normal}(\ln 0.005, 0.2)$$

$$\overline{\ln \mu} \sim \text{Normal}(\ln 15, 0.5)$$

$$\overline{\ln \tau} \sim \text{Normal}(\ln 0.1, 0.5)$$

$$\epsilon \sim \text{Gamma}(4, 10^{-4})$$

591 $\overline{\ln \lambda} \sim \text{Normal}(\ln 1, 0.4)$
 592 $\overline{\ln \theta} \sim \text{Normal}(\ln 500, 0.4)$
 593 $\sigma_{\ln \alpha} \sim \text{Normal}_{0,\infty}(0, 0.2)$
 594 $\sigma_{\ln \mu}, \sigma_{\ln \tau} \sim \text{Normal}_{0,\infty}(0, 0.5)$
 595 $\sigma_{\ln \lambda}, \sigma_{\ln \theta} \sim \text{Normal}_{0,\infty}(0, 0.4)$

596

597 where parameters describing $\bar{m}_i$ and v_i are partially pooled across months M with the added
 598 constraint $\alpha > 0$ since, unlike the previous cases, Bourguès et al. (1996) used only fresh
 599 tissue. Predictions are shown in Figure 4 (cf. Figure S4).

600

601 According to the model decomposition is generally faster in warmer seasons (Figure 4).
 602 Initial growth rate is not different between any two seasons ($P_{\Delta\alpha > 0} = 0.44\text{--}56$, 0.5 being a
 603 coin toss), but in summer photosynthetic half-life is 8.8 ± 3.9 days ($P_{\Delta\mu > 0} = 0.99$), 19 ± 4.2
 604 days ($P_{\Delta\mu > 0} = 1$) and 20 ± 19 days ($P_{\Delta\mu > 0} = 0.97$) shorter than in spring, autumn and winter
 605 respectively (Table 1). Unless we believe the experimental conditions immediately killed the
 606 seaweed, this agrees well with temperature sensitivity of detrital photosynthesis (Wright et al.
 607 2024) resulting in faster decomposition at higher temperatures (Filbee-Dexter et al. 2022,
 608 Frontier et al. 2022, Wright et al. 2024). As with the previous cases, the conventional model
 609 (Figure S4) underestimates decomposition (Table 1, 2).

610

611 **Light**

612 The accelerating effect of depth on decomposition has repeatedly been evidenced for
 613 *Laminaria* species (Frontier et al. 2021, 2022). Physiological measurements have clarified
 614 that light limitation is behind this effect (Frontier et al. 2021, 2022, Wright et al. 2024) and
 615 detrital photosynthesis has therefore been suggested as a general mechanism opposing
 616 photodegradation (Wright 2026b). Detritus of cold temperate *L. hyperborea* and its warm
 617 temperate congener *L. ochroleuca* was either exposed to three simulated depths *ex situ* at
 618 Roscoff (Frontier et al. 2021) or deployed at three depths *in situ* in Queen Anne's Battery
 619 marina in Plymouth (Frontier et al. 2022). In both studies, samples were repeatedly measured
 620 over time. In the original permutational analysis of variance or linear multilevel model,
 621 pseudo-replication was counteracted by partially pooling replicates (Frontier et al. 2021,
 622 2022). In a Bayesian framework, this strategy can seamlessly be applied to the nonlinear
 623 model

$$\begin{aligned}
& m_i \sim \text{Beta}'(\bar{m}_i(1 + v_i), 2 + v_i) \\
& \bar{m}_i = m(t_i, \alpha_i, \mu_i, \tau_i) \\
& v_i = v(t_i, \epsilon, \lambda, \theta) \\
& \ln \alpha_i = \ln \alpha_{S_i} + \ln \alpha_{R_i} \\
& \ln \mu_i = \ln \mu_{S_i} + \ln \mu_{R_i} - \beta_{S_i} D_i \\
& \ln \tau_i = \ln \tau_{S_i} + \ln \tau_{R_i} \\
& \ln \alpha_S \sim \text{Normal}(\overline{\ln \alpha}, \sigma_{\ln \alpha_S}) \\
& \ln \alpha_R \sim \text{Normal}(0, \sigma_{\ln \alpha_R}) \\
& \ln \mu_S \sim \text{Normal}(\overline{\ln \mu}, \sigma_{\ln \mu_S}) \\
& \ln \mu_R \sim \text{Normal}(0, \sigma_{\ln \mu_R}) \\
& \ln \beta_S \sim \text{Normal}(\overline{\ln \beta}, \sigma_{\ln \beta}) \\
& \ln \tau_S \sim \text{Normal}(\overline{\ln \tau}, \sigma_{\ln \tau_S}) \\
& \ln \tau_R \sim \text{Normal}(0, \sigma_{\ln \tau_R}) \\
& \overline{\ln \alpha} \sim \text{Normal}(\ln 0.004, 0.2) \\
& \overline{\ln \mu} \sim \text{Normal}(\ln 100, 0.15) \text{ or } \text{Normal}(\ln 100, 0.05) \\
& \overline{\ln \beta} \sim \text{Normal}(\ln 0.05, 0.25) \text{ or } \text{Normal}(\ln 0.25, 0.2) \\
& \overline{\ln \tau} \sim \text{Normal}(\ln 0.12, 0.2) \\
& \sigma_{\ln \alpha_S}, \sigma_{\ln \alpha_R}, \sigma_{\ln \tau_S}, \sigma_{\ln \tau_R} \sim \text{Normal}_{0,\infty}(0, 0.2) \\
& \sigma_{\ln \mu_S}, \sigma_{\ln \mu_R} \sim \text{Normal}_{0,\infty}(0, 0.15) \text{ or } \text{Normal}_{0,\infty}(0, 0.05) \\
& \sigma_{\ln \beta} \sim \text{Normal}_{0,\infty}(0, 0.25) \text{ or } \text{Normal}_{0,\infty}(0, 0.2) \\
& \epsilon, \lambda, \theta \sim \text{Priors as in Equation 14}
\end{aligned} \tag{19}$$

where parameters describing $\bar{m}_i$ are partially pooled across species S and replicates R . Additionally, the natural logarithm of μ is modelled to decrease linearly with depth D at a species-specific rate β . D is only included in the equation of μ since it logically cannot affect initial growth or final decay. Again $\alpha > 0$ since fresh tissue was used. I applied Equation 19 to both studies with mostly identical priors; where priors differ, they are listed first for Frontier et al. (2021) and then for Frontier et al. (2022). Mean predictions for each replicate and across replicates are shown in Figure 5 (cf. Figure S5).

Both models quantify how depth accelerates decomposition by reducing photosynthetic half-life. In the first study, the exponential decay rate of photosynthetic half-life with depth (β) is $7.8 \pm 1.5 \% \text{ m}^{-1}$ for *L. hyperborea* and $4.5 \pm 1.1 \% \text{ m}^{-1}$ for *L. ochroleuca*. In the second study, β is greater for both species— $17 \pm 5.5 \% \text{ m}^{-1}$ for *L. hyperborea* and $28 \pm 6.4 \% \text{ m}^{-1}$ for *L. ochroleuca*—likely because light attenuation in the turbid marina (Frontier et al. 2022) is much faster than that simulated by the light filters (Frontier et al. 2021). A species reversal in sensitivity to light limitation is evident from one study to the next. Photosynthetic half-life decays $3.3 \pm 1.9 \% \text{ m}^{-1}$ faster for *L. hyperborea* ($P_{\Delta\beta > 0} = 0.97$) in Roscoff (Figure 5a), while it decays $11 \pm 9.5 \% \text{ m}^{-1}$ faster for *L. ochroleuca* ($P_{\Delta\beta > 0} = 0.9$) in Plymouth (Figure 5b). This reversal may be an effect of latitude, giving each species a thermal “home advantage”. The interspecific contrast seems to be mostly due to detrital photosynthesis, with similar ($\Delta\tau = 0.18 \pm 6.3 \% \text{ day}^{-1}$, $P_{\Delta\tau > 0} = 0.52$) or marginally faster ($\Delta\tau = 8.2 \pm 28 \% \text{ day}^{-1}$, $P_{\Delta\tau > 0} = 0.63$) decay of *L. ochroleuca* detritus *sensu stricto* (Table 1). This aligns with other observations on faster decomposition of *L. ochroleuca* (Wright et al. 2022). The conventional model (Figure S5) underestimates shallow decomposition more than deep decomposition (Table 1, 2), as is expected due to the declining influence of detrital photosynthesis with depth.

Temperature

Several seaweeds float after detachment and therefore decompose in a light environment that is conducive to detrital photosynthesis. Examples include seaweeds with pneumatocysts such as the giant kelp *M. pyrifera* (Rothäusler et al. 2018) and those with a honeycomb thallus such as the bull kelp *Durvillaea antarctica* (Tala et al. 2019). Vandendriessche et al. (2007) report the most resolved, long-term data on floating seaweed decomposition. They conducted an *ex-situ* study on the estuarine egg and bladder wracks *Ascophyllum nodosum* and *Fucus vesiculosus*, which both have pneumatocysts to stay afloat. Vandendriessche et al. (2007) exposed detritus of both species to four temperatures and grazing by the isopod *Idotea balthica*. The data show the broadest spectrum of decomposition trajectories in the published literature, so Vandendriessche et al. (2007) opted for non-parametric local regression to fit trendlines. Naturally this method is not inferential, so they resorted to pre-calculating longevity (days) and linear decomposition rates ($\% \text{ day}^{-1}$) per sample and reporting analysis of variance *F* statistics and Spearman’s rank correlation coefficients for treatment effects. As with the previous example, replicates were repeatedly measured. Since a multilevel solution

to pseudo-replication was not feasible in this case, likely due to highly divergent replicates, I could not partially pool replicates but included them with no pooling as

$$m_i \sim \text{Beta}'(\bar{m}_i(1 + v_i), 2 + v_i) \quad (20)$$

$$\bar{m}_i = m(t_i, \alpha_i, \mu_i, \tau_i)$$

$$v_i = v(t_i, \epsilon, \lambda, \theta)$$

$$\ln \alpha_i = \ln \alpha_{S_i} + \ln \alpha_{R_i}$$

$$\ln \mu_i = \ln \mu_{S_i} + \ln \mu_{T_i} + \ln \mu_{R_i} + \beta_{T_i} G_i$$

$$\ln \tau_i = \ln \tau_{S_i} + \ln \tau_{T_i} + \ln \tau_{R_i} + \gamma_{T_i} G_i$$

$$\ln \alpha_S \sim \text{Normal}(\ln 0.004, 0.2)$$

$$\ln \mu_S \sim \text{Normal}(\ln 30, 0.2)$$

$$\ln \tau_S \sim \text{Normal}(\ln 0.1, 0.2)$$

$$\ln \alpha_R, \ln \mu_T, \ln \mu_R, \ln \tau_T, \ln \tau_R, \beta_T, \gamma_T \sim \text{Normal}(0, 0.2)$$

$$\epsilon, \lambda, \theta \sim \text{Priors as in Equation 14}$$

where both μ and τ are influenced by temperature T and grazing G . Unlike depth, temperature may elicit a nonlinear photosynthetic response, so is here treated categorically. The binary grazing variable G is effect coded as -0.5 for control and $+0.5$ for grazed and stratified by temperature since isopod metabolism is also expected to be temperature-dependent. Exponentiating β and γ therefore yields the multiplicative grazing effects on μ and τ . To highlight the flexibility of the macroalgal model, predictions for each replicate are shown in Figure 6 (cf. Figure S6).

Averaged across replicates and treatments, *A. nodosum* has 0.058 ± 0.043 % day⁻¹ faster initial detrital growth ($P_{\Delta\alpha > 0} = 0.91$) and 9.5 ± 1.6 % day⁻¹ slower final decomposition ($P_{\Delta\tau > 0} = 1$) but a 36 ± 4.5 days shorter photosynthetic half-life ($P_{\Delta\mu > 0} = 1$) than *F. vesiculosus*. Increasing temperature strongly reduces photosynthetic half-life across species (Figure 6). There is a 100% probability that photosynthetic half-life is 13 ± 1.5 times the average at 5°C but only 0.72 ± 0.074 , 0.33 ± 0.036 and 0.39 ± 0.041 times the average at 10, 15 and 18°C respectively. In the absence of detrital photosynthesis, there is practically no accelerating effect of temperature on decomposition ($P_{\Delta \ln \tau > 0} = 0.14\text{--}0.86$). The negative grazing effect on photosynthetic half-life strengthens with increasing temperature: 5°C ($e^\beta = 0.96 \pm 0.12$, $P_{\beta < 0} = 0.66$), 10°C ($e^\beta = 0.6 \pm 0.072$, $P_{\beta < 0} = 1$), 15°C ($e^\beta = 0.29 \pm 0.037$, $P_{\beta < 0} = 1$), 18°C ($e^\beta =$

0.22 $\pm$ 0.026, $P_{\beta < 0} = 1$). In contrast, grazing accelerates final decomposition evenly across temperatures: 5°C ($e' = 2 \pm 0.34$, $P_{\gamma > 0} = 1$), 10°C ($e' = 3.6 \pm 0.47$, $P_{\gamma > 0} = 1$), 15°C ($e' = 2.1 \pm 0.27$, $P_{\gamma > 0} = 1$), 18°C ($e' = 1.4 \pm 0.18$, $P_{\gamma > 0} = 0.99$). These results suggest that the accelerating temperature effect on seaweed decomposition (Filbee-Dexter et al. 2022, Frontier et al. 2022, Wright et al. 2024) acts almost exclusively through detrital photosynthesis rather than decomposer metabolism, which agrees with the conjecture of Wright et al. (2024). The conventional model (Figure S6) underestimates decomposition by up to 3.5 orders of magnitude and is only reasonably accurate for the most photosynthetically impaired detritus exposed to grazing at high temperatures (Table 1, 2).

Conclusion

Macroalgal decomposition has neither been consistently nor reliably modelled. But a realistic model is the key to understanding macroalgal blue carbon. I have presented the first decomposition model tailored to macroalgae by taking into account the theory of detrital photosynthesis. This is made possible by treating the exponential decay rate as a function of time rather than a constant, a function which is informed by the latest understanding of detrital photosynthesis. My model only has three parameters which describe the initial relative growth rate (α), the photosynthetic half-life (μ), and the final exponential decay rate (τ) which is directly comparable to that of the classic model (k). I have shown that these parameters are successfully estimated and ecologically interpretable for empirical data from a variety of contexts. Given a set of parameter estimates, the model can reliably predict new observations, even beyond the duration of the experiment. Although I have written the model in R with Stan and use a beta prime likelihood, it can be as effectively implemented with a gamma or even normal likelihood in any statistical framework that allows robust nonlinear modelling. Failing to account for detrital photosynthesis invariably leads to underestimation of macroalgal decomposition, especially when physiological influence is strong, leading me to believe that the conventional model is inadequate for these plants. I expect that the presented model will increase estimates of macroalgal decomposition in most cases and thus have major implications for seaweed blue carbon. I am already using this model on my own kelp decomposition data and in meta-analysis and hope you may find it useful too.

Acknowledgements

Thanks go to Andy Foggo and Nadia Frontier for inspiring discussions over the years. Nadia Frontier, Robert Hamersley, Sofie Vandendriessche and Florian de Bettignies kindly shared

their data with me. Thanks also to Pere Rovira for the email exchange and Bojana Manojlovic for the chat.

Funding

This work was supported by the Forrest Research Foundation Scholarship and Australian Government Research Training Program Scholarship. Open access publishing was facilitated by The University of Western Australia, as part of the Elsevier–The University of Western Australia agreement via the Council of Australian University Librarians.

Declaration of competing interest

I declare no conflicts of interest. Importantly I am not funded by any private enterprise with an interest in selling carbon credits.

Data and code availability

Data and code to reproduce this paper and apply my model to your own data are available at github.com/lukaseamus/limbodeco. Please cite this paper as well as the model (Wright 2026a).

References

- Babiker M, Berndes G, Blok K, Cohen B, Cowie A, Geden O, Ginzburg V, Leip A, Smith P, Sugiyama M et al. (2022) Cross-sectoral perspectives. Shukla PR, Skea J, Slade R, Al Khourdajie A, van Diemen R, McCollum D, Pathak M, Some S, Vyas P, Fradera R et al. (eds.) *IPCC, 2022: Climate Change 2022: Mitigation of Climate Change. Contribution of Working Group III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, 1245–1354. <https://doi.org/10.1017/9781009157926.005>.
- Banta GT, Pedersen MF, Nielsen SL (2004) Decomposition of marine primary producers: consequences for nutrient recycling and retention in coastal ecosystems. Nielsen SL, Banta GT, Pedersen MF (eds.) *Estuarine nutrient cycling: the influence of primary producers*, 187–216. https://doi.org/10.1007/978-1-4020-3021-5_7.
- Bedford AP, Moore PG (1984) Macrofaunal involvement in the sublittoral decay of kelp debris: the detritivore community and species interactions. *Estuarine, Coastal and Shelf Science* 18, 97–111. [https://doi.org/10.1016/0272-7714\(84\)90009-x](https://doi.org/10.1016/0272-7714(84)90009-x).

787 Birch PB, Gabrielson JO, Hamel KS (1983) Decomposition of *Cladophora*. I. Field studies in
 788 the Peel-Harvey estuarine system, Western Australia. *Botanica Marina* 26, 165–172.
 789 <https://doi.org/10.1515/botm.1983.26.4.165>.

790 Bourguès S, Auby I, de Wit R, Labourg P-J (1996) Differential anaerobic decomposition of
 791 seagrass (*Zostera noltii*) and macroalgal (*Monostroma obscurum*) biomass from
 792 Arcachon Bay (France). *Hydrobiologia* 329, 121–131.
 793 https://link.springer.com/chapter/10.1007/978-94-009-1744-6_10.

794 Bourguignon M, Santos-Neto M, Castro MD (2021) A new regression model for positive
 795 random variables with skewed and long tail. *Metron* 79, 33–55.
 796 <https://doi.org/10.1007/s40300-021-00203-y>.

797 Brouwer PEM (1996) Decomposition in situ of the sublittoral Antarctic macroalga
 798 *Desmarestia anceps* Montagne. *Polar Biology* 16, 129–137.
 799 <https://doi.org/10.1007/bf02390433>.

800 Carpenter B, Gelman A, Hoffman MD, Lee D, Goodrich B, Betancourt M, Brubaker M, Guo
 801 J, Li P, Riddell A (2017) Stan: a probabilistic programming language. *Journal of*
 802 *Statistical Software* 76, 1–32. <https://doi.org/10.18637/jss.v076.i01>.

803 Cebrián J, Duarte CM (1995) Plant growth-rate dependence of detrital carbon storage in
 804 ecosystems. *Science* 268, 1606–1608. <https://doi.org/10.1126/science.268.5217.1606>.

805 Cebrián J, Lartigue J (2004) Patterns of herbivory and decomposition in aquatic and
 806 terrestrial ecosystems. *Ecological Monographs* 74, 237–259. [https://doi.org/10.1890/03-](https://doi.org/10.1890/03-4019)
 807 4019.

808 Creed JC, Vieira VMNCS, Norton TA, Caetano D (2019) A meta-analysis shows that
 809 seaweeds surpass plants, setting life-on-Earth’s limit for biomass packing. *BMC*
 810 *Ecology* 19, 6. <https://doi.org/10.1186/s12898-019-0218-z>.

811 de Bettignies F, Dauby P, Thomas F, Gobet A, Delage L, Bohner O, Loisel S, Davoult D
 812 (2020) Degradation dynamics and processes associated with the accumulation of
 813 *Laminaria hyperborea* (Phaeophyceae) kelp fragments: an in situ experimental
 814 approach. *Journal of Phycology* 56, 1481–1492. <https://doi.org/10.1111/jpy.13041>.

815 Duarte CM, Gattuso J-P, Hancke K, Gundersen H, Filbee-Dexter K, Pedersen MF,
 816 Middelburg JJ, Burrows MT, Krumhansl KA, Wernberg T (2022) Global estimates of
 817 the extent and production of macroalgal forests. *Global Ecology and Biogeography* 0,
 818 1–18. <https://doi.org/10.1111/geb.13515>.

819 Enríquez S, Duarte CM, Sand-Jensen K (1993) Patterns in decomposition rates among
820 photosynthetic organisms: the importance of detritus C:N:P content. *Oecologia* 94,
821 457–471. <https://doi.org/10.1007/bf00566960>.

822 Filbee-Dexter K, Feehan CJ, Smale DA, Krumhansl KA, Augustine S, de Bettignies F,
823 Burrows MT, Byrnes JEK, Campbell J, Davoult D et al. (2022) Kelp carbon sink
824 potential decreases with warming due to accelerating decomposition. *PLoS Biology* 20,
825 e3001702. <https://doi.org/10.1371/journal.pbio.3001702>.

826 Filbee-Dexter K, Pessarrodona A, Pedersen MF, Wernberg T, Duarte CM, Assis J, Bekkby T,
827 Burrows MT, Carlson DF, Gattuso J-P (2024) Carbon export from seaweed forests to
828 deep ocean sinks. *Nature Geoscience* 17, 552–559. [https://doi.org/10.1038/s41561-024-](https://doi.org/10.1038/s41561-024-01449-7)
829 01449-7.

830 Friedlingstein P, O’sullivan M, Jones MW, Andrew RM, Hauck J, Landschützer P, Le Quéré
831 C, Li H, Luijkx IT, Olsen A (2025) Global carbon budget 2024. *Earth System Science*
832 *Data Discussions* 17, 965–1039. <https://doi.org/10.5194/essd-17-965-2025>.

833 Frontier N, de Bettignies F, Foggo A, Davoult D (2021) Sustained productivity and
834 respiration of degrading kelp detritus in the shallow benthos: detached or broken, but
835 not dead. *Marine Environmental Research* 166, 105277.
836 <https://doi.org/10.1016/j.marenvres.2021.105277>.

837 Frontier N, Mulas M, Foggo A, Smale DA (2022) The influence of light and temperature on
838 detritus degradation rates for kelp species with contrasting thermal affinities. *Marine*
839 *Environmental Research* 173, 105529.
840 <https://doi.org/10.1016/j.marenvres.2021.105529>.

841 Gabry J, Češnovar R, Johnson A, Bronder S (2024) cmdstanr: R Interface to ‘CmdStan’. R
842 package version 0.5.3. <https://mc-stan.org/cmdstanr>.

843 Gabry J, Simpson D, Vehtari A, Betancourt M, Gelman A (2019) Visualization in Bayesian
844 workflow. *Journal of the Royal Statistical Society Series A: Statistics in Society* 182,
845 389–402. <https://doi.org/10.1111/rssa.12378>.

846 Gohel D, Moog S, Heckmann M (2025) officer: manipulation of Microsoft Word and
847 PowerPoint documents. <https://ardata-fr.github.io/officeverse>.

848 Hamersley MR, Sohm JA, Burns JA, Capone DG (2015) Nitrogen fixation associated with
849 the decomposition of the giant kelp *Macrocystis pyrifera*. *Aquatic Botany* 125, 57–63.
850 <https://doi.org/10.1016/j.aquabot.2015.05.003>.

851 Hibbett D, Blanchette R, Kenrick P, Mills B (2016) Climate, decay, and the death of the coal
852 forests. *Current Biology* 26, R563–R567. <https://doi.org/10.1016/j.cub.2016.01.014>.

853 Josselyn MN, Mathieson AC (1980) Seasonal influx and decomposition of autochthonous
 854 macrophyte litter in a north temperate estuary. *Hydrobiologia* 71, 197–208.
 855 <https://doi.org/10.1007/bf03216236>.

856 Kay M (2024) tidybayes: tidy data and geoms for Bayesian models v3.0.7. *Zenodo*.
 857 <https://doi.org/10.5281/zenodo.13770114>.

858 Kennedy JR, Blain CO (2025) A systematic review of marine macroalgal degradation:
 859 toward a better understanding of macroalgal carbon sequestration potential. *Journal of*
 860 *Phycology* 61, 399–432. <https://doi.org/10.1111/jpy.70031>.

861 Krumhansl KA, Scheibling RE (2012) Detrital subsidy from subtidal kelp beds is altered by
 862 the invasive green alga *Codium fragile* ssp. *fragile*. *Marine Ecology Progress Series*
 863 456, 73–85. <https://doi.org/10.3354/meps09671>.

864 Lee D, Stan buildbot, seantalts, Carpenter B, Morris M, Kucukelbir A, Betancourt M, Tran D,
 865 Brubaker M, Bales B et al. (2017) stan-dev/cmdstan: v2.17.1. *Zenodo*.
 866 <https://doi.org/10.5281/zenodo.1117248>.

867 Manojlovic B (2021) Data assimilation techniques for aquatic ecological processes. PhD,
 868 University of Technology Sydney. <https://opus.lib.uts.edu.au/handle/10453/150726>.

869 McElreath R (2020) Statistical rethinking²: a Bayesian course with examples in R and Stan,
 870 second edition. Chapman and Hall/CRC. <https://doi.org/10.1201/9780429029608>.

871 Müller K (2025) here: a simpler way to find your files. <https://here.r-lib.org>.

872 Murray LM (2015) Bayesian state-space modelling on high-performance hardware using
 873 LibBi. *Journal of Statistical Software* 67, 1–36. <https://doi.org/10.18637/jss.v067.i10>.

874 Olson JS (1963) Energy storage and the balance of producers and decomposers in ecological
 875 systems. *Ecology* 44, 322–331. <https://doi.org/10.2307/1932179>.

876 Pedersen MF, Filbee-Dexter K, Frisk NL, Sárossy Z, Wernberg T (2021) Carbon
 877 sequestration potential increased by incomplete anaerobic decomposition of kelp
 878 detritus. *Marine Ecology Progress Series* 660, 53–67.
 879 <https://doi.org/10.3354/meps13613>.

880 Pedersen TL (2025) patchwork: the composer of plots. <https://patchwork.data-imaginist.com>.

881 Pessarrodona A, Franco-Santos RM, Wright LS, Vanderklift MA, Howard J, Pidgeon E,
 882 Wernberg T, Filbee-Dexter K (2023) Carbon sequestration and climate change
 883 mitigation using macroalgae: a state of knowledge review. *Biological Reviews* 98,
 884 1945–1971. <https://doi.org/10.1111/brv.12990>.

- Pessarrodona A, Assis J, Filbee-Dexter K, Burrows MT, Gattuso J-P, Duarte CM, Krause-Jensen D, Moore PJ, Smale DA, Wernberg T (2022) Global seaweed productivity. *Science Advances* 8, eabn2465. <https://doi.org/10.1126/sciadv.abn2465>.
- R Core Team (2022) R: a language and environment for statistical computing. R Foundation for Statistical Computing. <https://www.R-project.org/>.
- Rice DL, Tenore KR (1981) Dynamics of carbon and nitrogen during the decomposition of detritus derived from estuarine macrophytes. *Estuarine, Coastal and Shelf Science* 13, 681–690. [https://doi.org/10.1016/s0302-3524\(81\)80049-7](https://doi.org/10.1016/s0302-3524(81)80049-7).
- Rohatgi A (2022) WebPlotDigitizer v4.6. <https://automeris.io>.
- Rothäusler E, Reinwald H, López BA, Tala F, Thiel M (2018) High acclimation potential in floating *Macrocystis pyrifera* to abiotic conditions even under grazing pressure – a field study. *Journal of Phycology* 54, 368–379. <https://doi.org/10.1111/jpy.12643>.
- Rovira P, Rovira R (2010) Fitting litter decomposition datasets to mathematical curves: towards a generalised exponential approach. *Geoderma* 155, 329–343. <https://doi.org/10.1016/j.geoderma.2009.11.033>.
- RStudio Team (2022) RStudio: integrated development environment for R. RStudio, Inc. <http://www.rstudio.com/>.
- Smith BD, Foreman RE (1984) An assessment of seaweed decomposition within a southern Strait of Georgia seaweed community. *Marine Biology* 84, 197–205. <https://doi.org/10.1007/bf00393005>.
- Tala F, López BA, Velásquez M, Jeldres R, Macaya EC, Mansilla A, Ojeda J, Thiel M (2019) Long-term persistence of the floating bull kelp *Durvillaea antarctica* from the South-East Pacific: potential contribution to local and transoceanic connectivity. *Marine Environmental Research* 149, 67–79. <https://doi.org/10.1016/j.marenvres.2019.05.013>.
- Thomas H (2013) Senescence, ageing and death of the whole plant. *New Phytologist* 197, 696–711. <https://doi.org/10.1111/nph.12047>.
- Trevathan-Tackett SM, Jeffries TC, Macreadie PI, Manojlovic B, Ralph P (2020) Long-term decomposition captures key steps in microbial breakdown of seagrass litter. *Science of The Total Environment* 705, 135806. <https://doi.org/10.1016/j.scitotenv.2019.135806>.
- van den Brand T (2025) gg4x: hacks for ‘ggplot2’. <https://github.com/teunbrand/gg4x>.
- Vandendriessche S, Vincx M, Degraer S (2007) Floating seaweed and the influences of temperature, grazing and clump size on raft longevity—a microcosm study. *Journal of Experimental Marine Biology and Ecology* 343, 64–73. <https://doi.org/10.1016/j.jembe.2006.11.010>.

Vehtari A, Gelman A, Gabry J (2017) Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC. *Statistics and Computing* 27, 1413–1432. <https://doi.org/10.1007/s11222-016-9696-4>.

White LJ, Norkko A (2025) A path towards appropriate degradation experiments for assessing carbon sequestration potential of macroalgae. *Journal of Ecology* 113, 2730–2743. <https://doi.org/10.1111/1365-2745.70107>.

Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R, Grolemund G, Hayes A, Henry L, Hester J (2019) Welcome to the Tidyverse. *Journal of Open Source Software* 4, 1686. <https://doi.org/10.21105/joss.01686>.

Wright LS, Foggo A (2021) Photosynthetic pigments of co-occurring Northeast Atlantic *Laminaria* spp. are unaffected by decomposition. *Marine Ecology Progress Series* 678, 227–232. <https://doi.org/10.3354/meps13886>.

Wright LS, Pessarrodona A, Foggo A (2022) Climate-driven shifts in kelp forest composition reduce carbon sequestration potential. *Global Change Biology* 28, 5514–5531. <https://doi.org/10.1111/gcb.16299>.

Wright LS, Kregting L (2023) Genus-specific response of kelp photosynthetic pigments to decomposition. *Marine Biology* 170, 144. <https://doi.org/10.1007/s00227-023-04289-y>.

Wright LS, Simpkins T, Filbee-Dexter K, Wernberg T (2024) Temperature sensitivity of detrital photosynthesis. *Annals of Botany* 133, 17–28. <https://doi.org/10.1093/aob/mcad167>.

Wright LS (2026a) limbodeco: a model of macroalgal decomposition (Software). *Zenodo*. <https://doi.org/10.5281/zenodo.18654199>.

Wright LS (2026b) Plants in limbo — the theory of detrital photosynthesis. *Quantitative Plant Biology* in review.

Tables

Table 1. Macroalgal decomposition parameter estimates. Parameters of the presented macroalgal model (α , μ , τ in Equation 11) are inferred from seven empirical datasets alongside the conventional decay constant k (Equation 12). Estimates are mean $\pm$ standard deviation of 8×10^4 posterior samples; median is given in parentheses when k is highly right-skewed. NB exponential rates are given as percentages.

	k (% day ⁻¹)	α (% day ⁻¹)	μ (days)	τ (% day ⁻¹)
Brouwer 1996				
<i>Desmarestia anceps</i>				
Control	0.06 $\pm$ 0.014	0.068 $\pm$ 0.032	585 $\pm$ 100	5.6 $\pm$ 1.5

Pre-killed	3.4 ± 0.29	-1.3 ± 0.5	30 ± 12	5.6 ± 1.5
Hamersley <i>et al.</i> 2015				
<i>Macrocystis pyrifera</i>				
Fresh	5.7 ± 0.43	-0.3 ± 0.64	7 ± 1.1	11 ± 1.1
Senescent	10 ± 0.9	-0.79 ± 1.6	1.6 ± 0.56	12 ± 0.64
Detached	7.1 ± 0.57	-0.99 ± 1.6	2.9 ± 1.8	9.1 ± 0.8
de Bettignies <i>et al.</i> 2020				
<i>Laminaria hyperborea</i>				
Fresh (2017)	1.1 ± 0.1	0.26 ± 0.38	16 ± 9.5	1.3 ± 0.1
Senescent (2017)	3.7 ± 2.2	-2.4 ± 0.57	34 ± 33	3.6 ± 0.54
Senescent (2018)	0.54 ± 0.06	0.15 ± 0.34	71 ± 23	1.6 ± 0.39
Bourguès <i>et al.</i> 1996				
<i>Ulvaria obscura</i>				
Spring	3.8 ± 1.2	0.5 ± 0.14	13 ± 3.2	14 ± 6.9
Summer	7.7 ± 3.9	0.5 ± 0.14	4 ± 2.9	12 ± 7.5
Autumn	2.5 ± 0.87	0.51 ± 0.14	23 ± 3.4	14 ± 4.8
Winter	2.4 ± 0.97	0.49 ± 0.13	24 ± 19	11 ± 7.9
Frontier <i>et al.</i> 2021				
<i>Laminaria hyperborea</i>				
0 m	0.42 ± 5.8 (0.075)	0.4 ± 0.1	181 ± 190	14 ± 4.6
15 m	2.4 ± 43 (0.56)	0.4 ± 0.1	56 ± 58	14 ± 4.6
30 m	18 ± 136 (4.3)	0.4 ± 0.1	18 ± 22	14 ± 4.6
<i>Laminaria ochroleuca</i>				
0 m	1.5 ± 17 (0.34)	0.42 ± 0.11	169 ± 174	14 ± 4.6
15 m	2.2 ± 30 (0.57)	0.42 ± 0.11	87 ± 88	14 ± 4.6
30 m	4.7 ± 71 (0.98)	0.42 ± 0.11	46 ± 50	14 ± 4.6
Frontier <i>et al.</i> 2022				
<i>Laminaria hyperborea</i>				
0.5 m	0.15 ± 1.7 (0.024)	0.3 ± 0.12	114 ± 20	16 ± 15
1.5 m	0.2 ± 2.5 (0.039)	0.3 ± 0.12	97 ± 16	16 ± 15
3 m	0.4 ± 3.4 (0.081)	0.3 ± 0.12	75 ± 14	16 ± 15
<i>Laminaria ochroleuca</i>				
0.5 m	1.1 ± 25 (0.16)	0.64 ± 0.24	75 ± 11	24 ± 23
1.5 m	1.3 ± 11 (0.25)	0.64 ± 0.24	56 ± 8	24 ± 23
3 m	2.4 ± 16 (0.5)	0.64 ± 0.24	37 ± 6.8	24 ± 23
Vandendriessche <i>et al.</i> 2007				
<i>Ascophyllum nodosum</i>				
5°C Control	0.0074 ± 0.0045	0.42 ± 0.033	262 ± 30	4.2 ± 0.77
5°C Grazed	0.049 ± 0.026	0.42 ± 0.033	248 ± 29	8.3 ± 1.4
10°C Control	1 ± 0.48	0.42 ± 0.033	18 ± 1.9	2.8 ± 0.29
10°C Grazed	3.6 ± 1.6	0.42 ± 0.033	11 ± 1	10 ± 1
15°C Control	1.5 ± 0.73	0.42 ± 0.033	12 ± 1.3	3.7 ± 0.42
15°C Grazed	6.5 ± 3	0.42 ± 0.033	3.5 ± 0.4	7.8 ± 0.76
18°C Control	1.6 ± 0.74	0.42 ± 0.033	16 ± 1.6	5.7 ± 0.63
18°C Grazed	7 ± 3.1	0.42 ± 0.033	3.5 ± 0.38	7.7 ± 0.73
<i>Fucus vesiculosus</i>				
5°C Control	0.0044 ± 0.0027	0.36 ± 0.03	751 ± 96	11 ± 2.1
5°C Grazed	0.029 ± 0.015	0.36 ± 0.03	711 ± 88	22 ± 3.7
10°C Control	0.62 ± 0.29	0.36 ± 0.03	51 ± 5.4	7.2 ± 0.95
10°C Grazed	2.1 ± 0.97	0.36 ± 0.03	31 ± 3.2	26 ± 3.5

15°C Control	0.91 ± 0.45	0.36 ± 0.03	35 ± 3.9	9.7 ± 1.4
15°C Grazed	3.9 ± 1.8	0.36 ± 0.03	9.9 ± 1.2	20 ± 2.5
18°C Control	0.95 ± 0.45	0.36 ± 0.03	46 ± 4.9	15 ± 2.2
18°C Grazed	4.1 ± 1.8	0.36 ± 0.03	9.9 ± 1.1	20 ± 2.3

Table 2. Comparison between macroalgal (Equation 11) and conventional (Equation 12) decomposition parameters. Log ratios indicate by how many orders of magnitude the dead decay rate τ is greater than the conventional decay rate k or the detrital half-life $t_{1/2}$ is greater than the photosynthetic half-life μ . P gives the posterior probability of these differences. Prior $P(\tau > k) = 0.5$, equivalent to a coin toss, but prior $P(t_{1/2} > \mu)$ could not be set in a similar manner because $t_{1/2}$ is derived from k . Estimates are mean ± standard deviation of 8×10^4 posterior samples.

	$\log_{10}(\tau / k)$	$P(\tau > k)$	$\log_{10}(t_{1/2} / \mu)$	$P(t_{1/2} > \mu)$
Brouwer 1996				
<i>Desmarestia anceps</i>				
Control	2 ± 0.14	1	0.32 ± 0.12	0.99
Pre-killed	0.2 ± 0.11	0.98	-0.14 ± 0.16	0.17
Hamersley et al. 2015				
<i>Macrocystis pyrifera</i>				
Fresh	0.3 ± 0.052	1	0.25 ± 0.08	1
Senescent	0.068 ± 0.045	0.93	0.65 ± 0.13	1
Detached	0.11 ± 0.05	0.99	0.58 ± 0.2	0.99
de Bettignies et al. 2020				
<i>Laminaria hyperborea</i>				
Fresh (2017)	0.082 ± 0.053	0.94	0.64 ± 0.15	1
Senescent (2017)	0.027 ± 0.17	0.6	-0.086 ± 0.37	0.42
Senescent (2018)	0.47 ± 0.11	1	0.29 ± 0.17	0.99
Bourguès et al. 1996				
<i>Ulvaria obscura</i>				
Spring	0.55 ± 0.2	1	0.19 ± 0.18	0.88
Summer	0.2 ± 0.19	0.88	0.47 ± 0.31	0.95
Autumn	0.75 ± 0.19	1	0.12 ± 0.16	0.78
Winter	0.61 ± 0.33	0.95	0.2 ± 0.34	0.73
Frontier et al. 2021				
<i>Laminaria hyperborea</i>				
0 m	2.3 ± 0.78	1	0.87 ± 0.85	0.85
15 m	1.4 ± 0.7	0.98	0.5 ± 0.78	0.75
30 m	0.5 ± 0.73	0.76	0.13 ± 0.82	0.56
<i>Laminaria ochroleuca</i>				
0 m	1.6 ± 0.72	0.98	0.24 ± 0.79	0.62
15 m	1.4 ± 0.69	0.98	0.3 ± 0.77	0.65
30 m	1.1 ± 0.73	0.94	0.36 ± 0.8	0.68
Frontier et al. 2022				
<i>Laminaria hyperborea</i>				
0.5 m	2.7 ± 0.87	1	1.4 ± 0.81	0.97
1.5 m	2.5 ± 0.86	1	1.3 ± 0.79	0.96

3 m	2.2 ± 0.88	1	1.1 ± 0.81	0.92
<i>Laminaria ochroleuca</i>				
0.5 m	2 ± 0.84	0.99	0.77 ± 0.77	0.85
1.5 m	1.8 ± 0.83	0.99	0.7 ± 0.75	0.84
3 m	1.6 ± 0.85	0.97	0.59 ± 0.77	0.79
Vandendriessche et al. 2007				
<i>Ascophyllum nodosum</i>				
5°C Control	2.8 ± 0.26	1	1.6 ± 0.25	1
5°C Grazed	2.3 ± 0.23	1	0.81 ± 0.23	1
10°C Control	0.46 ± 0.2	0.99	0.61 ± 0.2	1
10°C Grazed	0.49 ± 0.19	0.99	0.3 ± 0.19	0.94
15°C Control	0.43 ± 0.2	0.98	0.62 ± 0.2	1
15°C Grazed	0.11 ± 0.19	0.72	0.53 ± 0.2	1
18°C Control	0.59 ± 0.2	1	0.47 ± 0.2	0.99
18°C Grazed	0.081 ± 0.19	0.66	0.5 ± 0.19	1
<i>Fucus vesiculosus</i>				
5°C Control	3.5 ± 0.27	1	1.4 ± 0.26	1
5°C Grazed	2.9 ± 0.24	1	0.59 ± 0.23	1
10°C Control	1.1 ± 0.2	1	0.39 ± 0.2	0.97
10°C Grazed	1.1 ± 0.2	1	0.079 ± 0.2	0.65
15°C Control	1.1 ± 0.21	1	0.4 ± 0.21	0.97
15°C Grazed	0.76 ± 0.2	1	0.31 ± 0.2	0.94
18°C Control	1.2 ± 0.21	1	0.24 ± 0.2	0.88
18°C Grazed	0.73 ± 0.19	1	0.28 ± 0.19	0.93

Figures





Figure 1. The mathematical model: the effect of varying α , μ and τ on the trajectory of the exponential decay rate k (Equation 6) and the proportion of remaining mass m (Equation 9) over time t (cf. Figure S1).



966

967 **Figure 2.** Example 1: the effect of physiology on the decomposition of *Desmarestia anceps*,
 968 shown as the proportion of remaining mass (mean $\pm$ standard deviation, $n = 5$) over time t (n
 969 $= 115$, data from Brouwer 1996). Data are plotted alongside predictions of new observations
 970 $\tilde{m}$, the mean $\bar{m}$, the time-variant exponential decay rate k and the photosynthetic half-life μ
 971 (Equation 15, cf. Figure S2). Lines and intervals are medians and the central 50, 80 and 90%
 972 of posterior probability. The distribution is a kernel density estimate of the probability density
 973 function of μ .

974




Figure 3. Example 2: the effect of senescence on the decomposition of *Macrocyctis pyrifera* (a, n = 61, data from Hamersley et al. 2015) and *Laminaria hyperborea* (b, n = 63, data from de Bettignies et al. 2020), shown as the proportion of remaining mass over time t . Data are plotted alongside predictions of the mean $\bar{m}$, the time-variant exponential decay rate k and photosynthetic half-life μ (Equations 16 and 17, cf. Figure S3). Lines and intervals are medians and the central 50, 80 and 90% of posterior probability. Distributions are kernel density estimates of the probability density function of μ .




Figure 4. Example 3: the effect of season on the decomposition of *Ulvaria obscura* ($n = 28$, data from Bourguès et al. 1996). Data are plotted alongside predictions of new observations $\tilde{m}$ (Equation 18, cf. Figure S4). Lines and intervals are medians and the central 50, 80 and 90% of posterior probability.





Figure 5. Example 4: the effect of light attenuation with depth on the decomposition of *Laminaria hyperborea* and *Laminaria ochroleuca* *ex situ* (**a**, $n = 210$, data from Frontier et al. 2021) and *in situ* (**b**, $n = 252$, data from Frontier et al. 2022), shown as the proportion of remaining mass over time t . Data are plotted alongside predictions of the mean $\bar{m}$ where lines are medians per replicate and intervals are the central 90% of posterior probability across replicates (Equation 19, cf. Figure S5).



999

1000 **Figure 6.** Example 5: the effect of temperature and grazing on the decomposition of floating
1001 *Ascophyllum nodosum* and *Fucus vesiculosus* (n = 1651, data from Vandendriessche et al.
1002 2007). Data are plotted alongside predictions of the mean $\bar{m}$ for each replicate (Equation 20,
1003 cf. Figure S6). Lines and intervals are medians and the central 50, 80 and 90% of posterior
1004 probability.





Figure 2

[Click here to access/download;Figure;Fi](#)







Figure 4

[Click here to access/download;Figure;Fi](#)

Spring


Summer


Autumn


Winter

$\mathcal{R}_E$

t (days)











Figure S2

[Click here to access/download;Figure;Fi](#)








Figure S4

[Click here to access/download;Figure;Fi](#)









Click here to access/download
Supplementary material
Supplement.pdf

1 *Ecological Modelling*

2 **A model of macroalgal decomposition**

3 Luka Seamus Wright^{1,2*}

4 ¹Oceans Institute, University of Western Australia, Perth, Australia

5 ²School of Biological Sciences, University of Western Australia, Perth, Australia

6 *luka.wright@research.uwa.edu.au, luka@wright.it, <https://orcid.org/0000-0002-1273-6256>

7

8 **Declaration of competing interest**

9 I declare no conflicts of interest. Importantly I am not funded by any private enterprise with
10 an interest in selling carbon credits.