


Patient Name: D

Patient ID:
Date of Birth:
Scan Date:
Analysis Date:
Image series: CCTA Spiral 0.5 Bv40 3 Best Systolic 340 ms

Plaque Analysis


	LM	LAD	LCx	RCA	Total	Plaque Burden
All Plaques (mm ³)	0	0	55	38	93	12%
 Calcified Plaque	0	0	1	3	4	0%
 Non Calcified Plaque	0	0	54	35	89	12%
 Low Attenuation Plaque	0	0	1	1	2	0%



0°

60°

120°




Remodeling Index	1.19
Total Plaque Volume	35 mm ³
 Calcified Plaque	1 mm ³
 Non Calcified Plaque	34 mm ³
 Low Attenuation Plaque	0 mm ³

0°

60°

120°




Remodeling Index	1.03
Total Plaque Volume	1 mm ³
Calcified Plaque	1 mm ³
Non Calcified Plaque	0 mm ³
Low Attenuation Plaque	0 mm ³

0°

60°

120°



Remodeling Index	1.02
Total Plaque Volume	54 mm ³
 Calcified Plaque	0 mm ³
 Non Calcified Plaque	54 mm ³
 Low Attenuation Plaque	1 mm ³

Summary and Recommendations

Mild amount of plaque (Stage 1) : 1 to 249mm³

Considerations for Medical Therapy of CAD Based on Quantitative Plaque Measurements on Coronary CTA

Table 2 Stages of Coronary Atherosclerosis by Total Plaque Volume and Percent Atheroma Volume

Stage of Atherosclerosis	Angiographic Stenosis Severity	Total Plaque Volume (mm ³)	Percent Atheroma Volume (%)
None	No stenosis	0	0%
Stage 1	1%-49% stenosis 1-vessel CAD >50% stenosis	>0 to 250	>0%-5.0%
Stage 2	2-vessel CAD >50% stenosis	>250 to 750	>5%-15.0%
Stage 3	3-vessel CAD >50% stenosis	>750	>15.0%

CAD = coronary artery disease.

Table 3 Simplified Approach to Medical Therapy Based Upon Stage of Atherosclerosis

Stage	Treatment	Serial CCTA
Stage 0	• GDMT/Shared decision for de-escalation of therapy	4 years
Stage 1	• Statin: (rosuvastatin 10-20 mg QD/atorvastatin 20-40 mg QD) • Ezetimibe 10 mg QD	3 years
Stage 2	• High-intensity statin (rosuvastatin 40 mg QD/atorvastatin 80 mg QD) • Ezetimibe 10 mg QD • Aspirin 81-100 mg QD • Rivaroxaban 2.5 mg BID <i>If diabetic, GLP-1 receptor agonist</i>	2 years

Stage 3	• High-intensity statin (rosuvastatin 40 mg QD/atorvastatin 80 mg QD) • Ezetimibe 10 mg QD • ASA 81-100 mg QD* • Rivaroxaban 2.5 mg BID* • Other lipid-lowering medications: PCSK-9 inhibitors, icosapent ethyl, inclisiran, bempedoic acid • Colchicine 0.6 mg QD • Cardiac rehabilitation or other supervised exercise program (if covered) <i>If diabetic: GLP-1 receptor agonist and SGLT2 inhibitor</i>	1 year
---------	--	--------

Freeman A, et al: Am J Med 2023