

Selective Pressures & Tropic Interactions Affecting Gall Size & Location on Goldenrod Stem


Luke Daniels

1/31/2018

Background

- Galls existing on goldenrod are a model for understanding selection facing flies and plants
- Gall making flies use an ovipositor to stimulate chemical reactions that form a gall
- Parasitic wasps are limited to attacking small galls due to the length of their ovipositor
- Abrahamson W.G 1989 found directional selection towards the larger gall due to parasitic interactions.

“ “




{\includegraphics[height=3cm]{Bird.png}}

Background

- Galls existing on goldenrod are a model for understanding selection facing flies and plants
- Gall making flies use an ovipositor to stimulate chemical reactions that form a gall
- Parasitic wasps are limited to attacking small galls due to the length of their ovipositor
- Abrahamson W.G 1989 found directional selection towards the larger gall due to parasitic interactions.

“ “




{\includegraphics[height=3cm]{Bird.png}}

Background

- Galls existing on goldenrod are a model for understanding selection facing flies and plants
- Gall making flies use an ovipositor to stimulate chemical reactions that form a gall
- Parasitic wasps are limited to attacking small galls due to the length of their ovipositor
- Abrahamson W.G 1989 found directional selection towards the larger gall due to parasitic interactions.

“ “




{\includegraphics{height=3cm}{Bird.png}}

Background

- Galls existing on goldenrod are a model for understanding selection facing flies and plants
- Gall making flies use an ovipositor to stimulate chemical reactions that form a gall
- Parasitic wasps are limited to attacking small galls due to the length of their ovipositor
- Abrahamson W.G 1989 found directional selection towards the larger gall due to parasitic interactions.

“ “




{\includegraphics{height=3cm}}{Bird.png}

Background

- Galls existing on goldenrod are a model for understanding selection facing flies and plants
- Gall making flies use an ovipositor to stimulate chemical reactions that form a gall
- Parasitic wasps are limited to attacking small galls due to the length of their ovipositor
- Abrahamson W.G 1989 found directional selection towards the larger gall due to parasitic interactions.

“ “




{\includegraphics{height=3cm}{Bird.png}}

Background

- Galls existing on goldenrod are a model for understanding selection facing flies and plants
- Gall making flies use an ovipositor to stimulate chemical reactions that form a gall
- Parasitic wasps are limited to attacking small galls due to the length of their ovipositor
- Abrahamson W.G 1989 found directional selection towards the larger gall due to parasitic interactions.

“ “




{\includegraphics{height=3cm}{Bird.png}}

Background

- Galls existing on goldenrod are a model for understanding selection facing flies and plants
- Gall making flies use an ovipositor to stimulate chemical reactions that form a gall
- Parasitic wasps are limited to attacking small galls due to the length of their ovipositor
- Abrahamson W.G 1989 found directional selection towards the larger gall due to parasitic interactions.

“ “




{\includegraphics{height=3cm}}{Bird.png}

Background

- Galls existing on goldenrod are a model for understanding selection facing flies and plants
- Gall making flies use an ovipositor to stimulate chemical reactions that form a gall
- Parasitic wasps are limited to attacking small galls due to the length of their ovipositor
- Abrahamson W.G 1989 found directional selection towards the larger gall due to parasitic interactions.

“ “




{\includegraphics{height=3cm}}{Bird.png}

Background

- Galls existing on goldenrod are a model for understanding selection facing flies and plants
- Gall making flies use an ovipositor to stimulate chemical reactions that form a gall
- Parasitic wasps are limited to attacking small galls due to the length of their ovipositor
- Abrahamson W.G 1989 found directional selection towards the larger gall due to parasitic interactions.

“ “




{\includegraphics{height=3cm}}{Bird.png}

Selective
Pressures &
Tropic
Interactions
Affecting Gail
Size &
Location on
Goldenrod
Stem

Luke Daniels


Trial Figure In L^AT_EX



Selective
Pressures &
Tropic
Interactions
Affecting Gall
Size &
Location on
Goldenrod
Stem


Luke Daniels

Trial Figure In L^AT_EX(2)



Choose a point on the unit circle. Connect it to the origin with a line of length one, and denote the angle between that line and the horizontal coordinate axis by θ .

Trial Figure In $\text{\LaTeX}(3)$



Choose a point on the unit circle. Connect it to the origin with a line of length one, and denote the angle between that line and the horizontal coordinate axis by θ .