

# Statistics of Maximal Independent Sets in Grid-like Graphs

---

Levi Axelrod Nathan Bickel Anastasia Halfpap **Luke Hawranick** Alex Parker Cole Swain

January 27, 2025


Department of Mathematics  
Iowa State University

## Preliminaries

---

## Definition (Independent Sets)


A set  $I \subseteq V(G)$  is called an **independent set** if no two vertices in  $I$  are adjacent.



$G_{3 \times 3}$

## Definition (Independent Sets)


A set  $I \subseteq V(G)$  is called an **independent set** if no two vertices in  $I$  are adjacent.



$G_{3 \times 3}$

## Definition (Independent Sets)


A set  $I \subseteq V(G)$  is called an **independent set** if no two vertices in  $I$  are adjacent.



$G_{3 \times 3}$

## Definition (Independent Sets)

A set  $I \subseteq V(G)$  is called an **independent set** if no two vertices in  $I$  are adjacent.



$G_{3 \times 3}$

## Definition ( $\text{MIS}(G)$ )

Let  $G$  be a graph. We define

$$\text{MIS}(G) := \{M \subseteq V(G) : M \text{ is a maximal independent set.}\}$$

**Definition (MIS( $G$ ))**

Let  $G$  be a graph. We define

$$\text{MIS}(G) := \{M \subseteq V(G) : M \text{ is a maximal independent set.}\}$$

For general graph  $G(n)$ ,

- Enumeration of  $\text{MIS}(G)$  is known to be  $\#P$ -complete



## Definition (MIS( $G$ ))

Let  $G$  be a graph. We define

$$\text{MIS}(G) := \{M \subseteq V(G) : M \text{ is a maximal independent set.}\}$$

For general graph  $G(n)$ ,

- Enumeration of  $\text{MIS}(G)$  is known to be  $\#P$ -complete
- Miller and Muller (1960):

$$\text{MIS}(G) \leq \begin{cases} 3^{n/3} & \text{if } n \equiv 0 \pmod{3} \\ 4 \cdot 3^{n/3-1} & \text{if } n \equiv 1 \pmod{3} \\ 2 \cdot 3^{n/3} & \text{if } n \equiv 2 \pmod{3} \end{cases}$$

## Definition (MIS( $G$ ))

Let  $G$  be a graph. We define

$$\text{MIS}(G) := \{M \subseteq V(G) : M \text{ is a maximal independent set.}\}$$

For general graph  $G(n)$ ,

- Enumeration of  $\text{MIS}(G)$  is known to be  $\#P$ -complete
- Miller and Muller (1960):

$$\text{MIS}(G) \leq \begin{cases} 3^{n/3} & \text{if } n \equiv 0 \pmod{3} \\ 4 \cdot 3^{n/3-1} & \text{if } n \equiv 1 \pmod{3} \\ 2 \cdot 3^{n/3} & \text{if } n \equiv 2 \pmod{3} \end{cases}$$

- Moon and Moser (1965), Erdős (1966): Bounding  $g(n) :=$  the maximum number of different sizes of MIS's

$$n - \log n - H(n) - O(1) \leq g(n) \leq n - \log n$$

## Definition (Grid-like graph)

Let  $V_i := \{(i, j) : 1 \leq j \leq m\}$ . A graph  $G$  is **grid-like** provided that

1.  $G = G_{m \times n} \cup E$  for some edge-set  $E$ .

2.  $\Delta(G) = 4$

3. For any  $v \in V_i$  with  $2 \leq i \leq n-1$ ,

$$N_G(v) \subseteq N_{G_{m \times n}}(v) \cup V_i$$

4. For any  $1 \leq i_1, i_2 \leq n$ ,  $G[V_{i_1}] \cong G[V_{i_2}]$

## Definition (Grid-like graph)

Let  $V_i := \{(i, j) : 1 \leq j \leq m\}$ . A graph  $G$  is **grid-like** provided that

1.  $G = G_{m \times n} \cup E$  for some edge-set  $E$ .

2.  $\Delta(G) = 4$

3. For any  $v \in V_i$  with  $2 \leq i \leq n-1$ ,

$$N_G(v) \subseteq N_{G_{m \times n}}(v) \cup V_i$$


4. For any  $1 \leq i_1, i_2 \leq n$ ,  $G[V_{i_1}] \cong G[V_{i_2}]$

## Definition (Global and Local Structure)


Given a grid-like graph  $G$ , let  $H$  denote the graph to which each subgraph  $G[V_i]$  is isomorphic to. We call  $H$  the **local structure** of  $G$  and each subgraph  $G[V_i]$  to be a **slice** of  $G$ .

## Preliminaries


There are four particular grid-like graphs that we study. They are pictured below for  $m = 3$  and  $n = 4$ :




Fat Cylinder:  $FC_{m \times n}$



Thin Cylinder:  $TC_{m \times n}$



Torus:  $T_{m \times n}$



Möbius Strip:  $M_{m \times n}$

- Asymptotics of  $\text{MIS}(G)$  across several classes of grid-like graphs.

- Asymptotics of  $\text{MIS}(G)$  across several classes of grid-like graphs.
- Enumerating non-isomorphic MIS's for small  $m$ .

- Asymptotics of  $\text{MIS}(G)$  across several classes of grid-like graphs.
- Enumerating non-isomorphic MIS's for small  $m$ .
- Finding the average size of MIS's for small  $m$ .



## Framework

---

Golin et al. (2005) surveyed a set of enumeration problems on grid graphs, grid-cylinders, and grid-tori of fixed height, which can be modeled by the *transfer matrix approach*, including

- Hamiltonian Cycles
- Perfect Matchings
- Spanning Trees
- Cycle Covers

On such a grid-like graph to count  $S(m, n)$  objects, the method finds vectors  $a, b$  and a square matrix  $A$  such that

$$|S(m, n)| = a^\top A^n b$$




## Motivating Example

Let  $G_{2 \times n}$  be formed from  $\{1, 2, \dots, n\} \times \{1, 2\}$ . Consider  $\text{MIS}(G_{2 \times n})$  and an element  $M$ .

## Motivating Example

Let  $G_{2 \times n}$  be formed from  $\{1, 2, \dots, n\} \times \{1, 2\}$ . Consider  $\text{MIS}(G_{2 \times n})$  and an element  $M$ .

1. There can be at most one vertex from each column in  $M$



## Motivating Example


Let  $G_{2 \times n}$  be formed from  $\{1, 2, \dots, n\} \times \{1, 2\}$ . Consider  $\text{MIS}(G_{2 \times n})$  and an element  $M$ .

1. There can be at most one vertex from each column in  $M$

## Motivating Example

Let  $G_{2 \times n}$  be formed from  $\{1, 2, \dots, n\} \times \{1, 2\}$ . Consider  $\text{MIS}(G_{2 \times n})$  and an element  $M$ .

1. There can be at most one vertex from each column in  $M$
2. The first and last columns of  $M$  must include 1 vertex.



## Motivating Example

Let  $G_{2 \times n}$  be formed from  $\{1, 2, \dots, n\} \times \{1, 2\}$ . Consider  $\text{MIS}(G_{2 \times n})$  and an element  $M$ .


1. There can be at most one vertex from each column in  $M$
2. The first and last columns of  $M$  must include 1 vertex.



## Motivating Example

Let  $G_{2 \times n}$  be formed from  $\{1, 2, \dots, n\} \times \{1, 2\}$ . Consider  $\text{MIS}(G_{2 \times n})$  and an element  $M$ .

1. There can be at most one vertex from each column in  $M$
2. The first and last columns of  $M$  must include 1 vertex.
3. For every two adjacent columns, there is at least one vertex in  $M$ .



## Motivating Example

Let  $G_{2 \times n}$  be formed from  $\{1, 2, \dots, n\} \times \{1, 2\}$ . Consider  $\text{MIS}(G_{2 \times n})$  and an element  $M$ .

1. There can be at most one vertex from each column in  $M$
2. The first and last columns of  $M$  must include 1 vertex.
3. For every two adjacent columns, there is at least one vertex in  $M$ .

## Motivating Example

Let  $G_{2 \times n}$  be formed from  $\{1, 2, \dots, n\} \times \{1, 2\}$ . Consider  $\text{MIS}(G_{2 \times n})$  and an element  $M$ .

1. There can be at most one vertex from each column in  $M$
2. The first and last columns of  $M$  must include 1 vertex.
3. For every two adjacent columns, there is at least one vertex in  $M$ .
4.  $M$  has a unique dual, formed by reflecting its choice of vertices over the horizontal axis between the two rows.



## Motivating Example

By (1) and (2), an MIS of  $\text{MIS}(G_{2 \times n})$  contains exactly one of the vertices in the last column. Consider the sets which contain  $(n, 2)$ .


By 3, column  $n - 1$  is either empty or not.

## Motivating Example

By (1) and (2), an MIS of  $\text{MIS}(G_{2 \times n})$  contains exactly one of the vertices in the last column. Consider the sets which contain  $(n, 2)$ .

By 3, column  $n - 1$  is either empty or not.

- Not empty:




## Motivating Example

By (1) and (2), an MIS of  $\text{MIS}(G_{2 \times n})$  contains exactly one of the vertices in the last column. Consider the sets which contain  $(n, 2)$ .

By 3, column  $n - 1$  is either empty or not.

- Empty:



## Motivating Example

By (1) and (2), an MIS of  $\text{MIS}(G_{2 \times n})$  contains exactly one of the vertices in the last column. Consider the sets which contain  $(n, 2)$ .

By 3, column  $n - 1$  is either empty or not.

Thus, by (4),

$$|\text{MIS}(G_{2 \times n})| = 2 \left( \frac{1}{2} |\text{MIS}(G_{2 \times (n-1)})| + \frac{1}{2} |\text{MIS}(G_{2 \times (n-2)})| \right)$$

With the initial conditions

$$|\text{MIS}(G_{2 \times 1})| = 2 \quad , \quad |\text{MIS}(G_{2 \times 2})| = 2$$

$$|\text{MIS}(G_{2 \times n})| = 2F_n$$

If we can use an auxiliary digraph to model our problem, we can borrow the transfer matrix framework for enumeration:  $a^\top A^n b$ .



If we can use an auxiliary digraph to model our problem, we can borrow the transfer matrix framework for enumeration:  $a^\top A^n b$ .

## Definition (State of local structure)

Let  $H$  be the local structure of  $G$ . A **state** of  $H$  is an ordered pair  $(I, D)$  in which

1.  $I$  is an independent set of  $H$  such that  $H[V(H) \setminus N[I]]$  is 2-colorable;
2.  $D$ , the **deficit**, is a color class of a 2-coloring of  $H[V(H) \setminus N[I]]$

We define  $U(I) := V(H) \setminus N[I]$  to be the *uncovered set* of a state  $(I, D)$ .

If we can use an auxiliary digraph to model our problem, we can borrow the transfer matrix framework for enumeration:  $a^\top A^n b$ .

## Definition (State of local structure)

Let  $H$  be the local structure of  $G$ . A **state** of  $H$  is an ordered pair  $(I, D)$  in which

1.  $I$  is an independent set of  $H$  such that  $H[V(H) \setminus N[I]]$  is 2-colorable;
2.  $D$ , the **deficit**, is a color class of a 2-coloring of  $H[V(H) \setminus N[I]]$

We define  $U(I) := V(H) \setminus N[I]$  to be the *uncovered set* of a state  $(I, D)$ .

## Definition (State orderings)

State  $(I', D')$  **follows** state  $(I, D)$  or provided that

1.  $I \cap I' = \emptyset$
2.  $D \subseteq I'$
3.  $D' = U(I') \setminus I$ .

## State Definition Example

In this state,

$$H = P_4$$

$$I = \{b\}$$

$$D = \{d\}$$



## State Definition Example

In this state,

$$H = P_4$$

$$I = \{b\}$$

$$D = \{d\}$$



In this state,


$$H = P_4$$

$$I = \{b\}$$

$$D = \emptyset$$



## State Ordering Example



We construct a digraph that stores information about which states can follow which.

We construct a digraph that stores information about which states can follow which.

### Definition (Map Digraph)

Let  $S(H)$  be the set of states of  $H$ . Let  $M(H)$  be the **map digraph** of  $G$  with

$$V(M(H)) := S(H) \quad , \quad E(M(H)) := \{ \overrightarrow{s_1 s_2} : (s_1, s_2) \in S(H)^2, s_1 \vdash s_2 \}$$

We construct a digraph that stores information about which states can follow which.

### Definition (Map Digraph)

Let  $S(H)$  be the set of states of  $H$ . Let  $M(H)$  be the **map digraph** of  $G$  with

$$V(M(H)) := S(H) \quad , \quad E(M(H)) := \{ \overrightarrow{s_1 s_2} : (s_1, s_2) \in S(H)^2, s_1 \vdash s_2 \}$$

There is a bijection between the number valid state permutations with valid starting and ending states and the number of MISs on  $G$ .



We construct a digraph that stores information about which states can follow which.

### Definition (Map Digraph)

Let  $S(H)$  be the set of states of  $H$ . Let  $M(H)$  be the **map digraph** of  $G$  with

$$V(M(H)) := S(H) \quad , \quad E(M(H)) := \{ \overrightarrow{s_1 s_2} : (s_1, s_2) \in S(H)^2, s_1 \vdash s_2 \}$$

There is a bijection between the number valid state permutations with valid starting and ending states and the number of MISs on  $G$ .

Note that the total number of walks of length  $n$  on this digraph overcount the number of MISs on our base graph  $G$ . To filter out digraph walks with invalid starting and ending states, we create another digraph to store this information.

We construct a digraph that stores information about which states can follow which.

## Definition (Map Digraph)

Let  $S(H)$  be the set of states of  $H$ . Let  $M(H)$  be the **map digraph** of  $G$  with

$$V(M(H)) := S(H) \quad , \quad E(M(H)) := \{\overrightarrow{s_1 s_2} : (s_1, s_2) \in S(H)^2, s_1 \vdash s_2\}$$

There is a bijection between the number valid state permutations with valid starting and ending states and the number of MISs on  $G$ .

Note that the total number of walks of length  $n$  on this digraph overcount the number of MISs on our base graph  $G$ . To filter out digraph walks with invalid starting and ending states, we create another digraph to store this information.

## Definition (Ticket Digraph)

Let  $T$  be the **ticket digraph** of  $G$  with

$$V(T) := S(H) \quad , \quad E(T) := \{\overrightarrow{s_1 s_2} : (s_1, s_2) \in S(H)^2, \text{ an MIS can start in state } s_1 \text{ and end in state } s_2\}$$

## Theorem (Transfer Matrix Application)

Let  $A_{M(H)}$  be the adjacency matrix of  $M(H)$  and  $A_T$  be the adjacency matrix of  $T$ . Then,

$$\tau(n) = |\text{MIS}(G)| = A_T \cdot A_{M(H)}^{n-1}$$

Note that the edges of  $T$  vary with the global structure of  $G$ .

- Global path structure of  $G \implies$

$$E(T) = \overrightarrow{\{(I, D)(I', D') : ((I, D), (I', D')) \in S(H)^2, D = U(I) \text{ and } D' = \emptyset\}}$$

Note that the edges of  $T$  vary with the global structure of  $G$ .

- Global path structure of  $G \implies$


$$E(T) = \{\overrightarrow{(I, D)(I', D')} : ((I, D), (I', D')) \in S(H)^2, D = U(I) \text{ and } D' = \emptyset\}$$

- Global cyclic structure of  $G \implies$

$$E(T) = \{\overrightarrow{s_1 s_2} : (s_1, s_2) \in S(H)^2 : s_2 \vdash s_1\}$$

## Map and Ticket Digraph Example


Let  $G$  have local structure  $P_2$ . The map digraph of  $G$  is



$P_2$  Map Digraph

## Map and Ticket Digraph Example


We consider two global structures of  $G$ : path and cyclic. The corresponding ticket digraphs are




Global Path Structure

## Map and Ticket Digraph Example

We consider two global structures of  $G$ : path and cyclic. The corresponding ticket digraphs are



Global Path Structure



Global Cyclic Structure



Let  $P_m$  or  $C_m$  be the slice of a grid-like graph and let  $M$  be the map digraph of the slice of  $G$ .

Let  $P_m$  or  $C_m$  be the slice of a grid-like graph and let  $M$  be the map digraph of the slice of  $G$ .

- $M$  is strongly connected with diameter at most 4.

Let  $P_m$  or  $C_m$  be the slice of a grid-like graph and let  $M$  be the map digraph of the slice of  $G$ .

- $M$  is strongly connected with diameter at most 4.
- $|V(M)|$  satisfies a linear recurrence and is  $\Theta(\lambda_1^m)$  for  $\lambda_1 \approx 1.75488$ .

Let  $P_m$  or  $C_m$  be the slice of a grid-like graph and let  $M$  be the map digraph of the slice of  $G$ .

- $M$  is strongly connected with diameter at most 4.
- $|V(M)|$  satisfies a linear recurrence and is  $\Theta(\lambda_1^m)$  for  $\lambda_1 \approx 1.75488$ .
- $|E(M)|$  also satisfied a linear recurrence and is  $\Theta(\lambda_2^m)$  for  $\lambda_2 \approx 2.1781$ .

Let  $P_m$  or  $C_m$  be the slice of a grid-like graph and let  $M$  be the map digraph of the slice of  $G$ .

- $M$  is strongly connected with diameter at most 4.
- $|V(M)|$  satisfies a linear recurrence and is  $\Theta(\lambda_1^m)$  for  $\lambda_1 \approx 1.75488$ .
- $|E(M)|$  also satisfied a linear recurrence and is  $\Theta(\lambda_2^m)$  for  $\lambda_2 \approx 2.1781$ .
- For each  $v \in M$ ,  $d^+(v) = 2^k$  for some  $k \in \mathbb{N}_0$ .

Let  $P_m$  or  $C_m$  be the slice of a grid-like graph and let  $M$  be the map digraph of the slice of  $G$ .

- $M$  is strongly connected with diameter at most 4.
- $|V(M)|$  satisfies a linear recurrence and is  $\Theta(\lambda_1^m)$  for  $\lambda_1 \approx 1.75488$ .
- $|E(M)|$  also satisfied a linear recurrence and is  $\Theta(\lambda_2^m)$  for  $\lambda_2 \approx 2.1781$ .
- For each  $v \in M$ ,  $d^+(v) = 2^k$  for some  $k \in \mathbb{N}_0$ .
- $\Delta^+(M(P_m)) = 2^{\lceil \frac{m}{2} \rceil}$ ,  $\Delta^+(M(C_m)) = 2^{\lfloor \frac{m}{2} \rfloor}$

## Enumeration

---

## $\tau(n)$ is a linear recurrence

### Theorem ( $\tau(n)$ is a linear recurrence)

Let  $(M, T)$  be the auxiliary digraphs of  $G$  on  $k$ . Let  $A_M$  and  $A_T$  be the adjacency matrices of  $M$  and  $T$  respectively. Let  $f(x) = x^k + a_{k-1}x^{k-1} + \dots + a_1x + a_0$  be the characteristic polynomial of  $A_M$ . Then,  $\tau$  satisfies the recurrence

$$\tau(n) = -a_{k-1}\tau(n-1) - \dots - a_1\tau(n-k+1) - a_0\tau(n-k)$$



## Theorem ( $\tau(n)$ is a linear recurrence)

Let  $(M, T)$  be the auxiliary digraphs of  $G$  on  $k$ . Let  $A_M$  and  $A_T$  be the adjacency matrices of  $M$  and  $T$  respectively. Let  $f(x) = x^k + a_{k-1}x^{k-1} + \dots + a_1x + a_0$  be the characteristic polynomial of  $A_M$ . Then,  $\tau$  satisfies the recurrence

$$\tau(n) = -a_{k-1}\tau(n-1) - \dots - a_1\tau(n-k+1) - a_0\tau(n-k)$$

## Proof

- By the Cayley-Hamilton Theorem,  $A_M$  satisfies the linear recurrence given by  $f$ .

$$A_M^n = -a_{k-1}A_M^{n-1} - \dots - a_1A_M^{n-k+1} - a_0A_M^{n-k}$$

# $\tau(n)$ is a linear recurrence

## Theorem ( $\tau(n)$ is a linear recurrence)

Let  $(M, T)$  be the auxiliary digraphs of  $G$  on  $k$ . Let  $A_M$  and  $A_T$  be the adjacency matrices of  $M$  and  $T$  respectively. Let  $f(x) = x^k + a_{k-1}x^{k-1} + \dots + a_1x + a_0$  be the characteristic polynomial of  $A_M$ . Then,  $\tau$  satisfies the recurrence

$$\tau(n) = -a_{k-1}\tau(n-1) - \dots - a_1\tau(n-k+1) - a_0\tau(n-k)$$

## Proof

- By the Cayley-Hamilton Theorem,  $A_M$  satisfies the linear recurrence given by  $f$ .

$$A_M^n = -a_{k-1}A_M^{n-1} - \dots - a_1A_M^{n-k+1} - a_0A_M^{n-k}$$

- Matrix dot product is linear.

$$\tau(n) = -a_{k-1}A_T \bullet A_M^{n-1} - \dots - a_1A_T \bullet A_M^{n-k+1} - a_0A_T \bullet A_M^{n-k}$$

Recall that  $|\text{MIS}(G_{2 \times n})| = 2F_n$ . From the theorem above,  $\tau(n) = \tau(n-1) + \tau(n-2)$ , so  $\tau$  grows exponentially with rate  $\varphi = \frac{1+\sqrt{5}}{2}$ . More specifically,

$$\lim_{n \rightarrow \infty} \frac{\tau(n)}{\frac{2}{\sqrt{5}} \cdot \varphi^n} = 1$$

Recall that  $|\text{MIS}(G_{2 \times n})| = 2F_n$ . From the theorem above,  $\tau(n) = \tau(n-1) + \tau(n-2)$ , so  $\tau$  grows exponentially with rate  $\varphi = \frac{1+\sqrt{5}}{2}$ . More specifically,

$$\lim_{n \rightarrow \infty} \frac{\tau(n)}{\frac{2}{\sqrt{5}} \cdot \varphi^n} = 1$$

## Theorem (Existence of $c, r$ )

Let  $m \in \mathbb{N}$ . The sequences  $|\text{MIS}(G_{m \times n})|, |\text{MIS}(FC_{m \times n})|, |\text{MIS}(TC_{m \times n})|, |\text{MIS}(T_{m \times n})|, |\text{MIS}(M_{m \times n})|$  as functions of  $n$  all obey linear recurrences. Moreover, for each sequence  $\tau(n)$ , there exists real numbers  $c > 0$  and  $r > 1$  such that

$$\lim_{n \rightarrow \infty} \frac{\tau(n)}{c \cdot r^n} = 1$$

**Proof:**

## **Theorem (Perron (1907), Frobenius (1912))**

*Let  $A$  be a primitive square matrix. Then,  $A$  has a Perron-Frobenius eigenvalue  $r$ , i.e. an eigenvalue equal to its spectral radius, such that the left and right eigenspaces of  $r$  are generated by single strictly positive vectors  $\vec{w}^\top$  and  $\vec{v}$  respectively. Moreover*

$$\lim_{n \rightarrow \infty} \frac{A^n}{r^n} = \frac{\vec{v} \vec{w}^\top}{\vec{w}^\top \vec{v}}$$

## Proof:

### Theorem (Perron (1907), Frobenius (1912) - application)

$A_M$  is a primitive square matrix. Thus,  $A_M$  has a Perron-Frobenius eigenvalue  $r$ , i.e. an eigenvalue equal to its spectral radius, such that the left and right eigenspaces of  $r$  are generated by single strictly positive vectors  $\vec{w}^\top$  and  $\vec{v}$  respectively. Moreover

$$\lim_{n \rightarrow \infty} \frac{A_M^n}{r^n} = \frac{\vec{v} \vec{w}^\top}{\vec{w}^\top \vec{v}}$$

$$\lim_{n \rightarrow \infty} \frac{\tau(n)}{r^n} = \lim_{n \rightarrow \infty} \frac{A_T \bullet A_M^n}{r^n} = A_T \bullet \lim_{n \rightarrow \infty} \frac{A_M^n}{r^n} = A_T \bullet \frac{\vec{v} \vec{w}^\top}{\vec{w}^\top \vec{v}} = c$$

## Conjecture (Looking for useful Perron-Frobenius eigenvector properties)

Let  $H$  be a graph and let  $\phi$  be a graph automorphism of  $H$ . Let  $r$  be the principle eigenvalue of the map digraph  $M$  of  $H$ . Then,

$$\lim_{n \rightarrow \infty} \frac{|\text{MIS}((H \square P_{n+1})/\phi)|}{r^n} = 1$$

**Efforts:**

## Conjecture (Looking for useful Perron-Frobenius eigenvector properties)

Let  $H$  be a graph and let  $\phi$  be a graph automorphism of  $H$ . Let  $r$  be the principle eigenvalue of the map digraph  $M$  of  $H$ . Then,

$$\lim_{n \rightarrow \infty} \frac{|\text{MIS}((H \square P_{n+1})/\phi)|}{r^n} = 1$$

### Efforts:

- Note that quotienting out by the  $\phi$  does not affect the map digraph. It only affects which pairs of states an MIS can start and end in.



## Conjecture (Looking for useful Perron-Frobenius eigenvector properties)

Let  $H$  be a graph and let  $\phi$  be a graph automorphism of  $H$ . Let  $r$  be the principle eigenvalue of the map digraph  $M$  of  $H$ . Then,

$$\lim_{n \rightarrow \infty} \frac{|\text{MIS}((H \square P_{n+1})/\phi)|}{r^n} = 1$$

### Efforts:

- Note that quotienting out by the  $\phi$  does not affect the map digraph. It only affects which pairs of states an MIS can start and end in.
- When  $\phi = \text{id}$ , an MIS which starts in state  $i$  must end in precisely state  $i$  after  $n$  steps through  $M$ .

## Conjecture (Looking for useful Perron-Frobenius eigenvector properties)

Let  $H$  be a graph and let  $\phi$  be a graph automorphism of  $H$ . Let  $r$  be the principle eigenvalue of the map digraph  $M$  of  $H$ . Then,

$$\lim_{n \rightarrow \infty} \frac{|\text{MIS}((H \square P_{n+1})/\phi)|}{r^n} = 1$$

### Efforts:

- Note that quotienting out by the  $\phi$  does not affect the map digraph. It only affects which pairs of states an MIS can start and end in.
- When  $\phi = \text{id}$ , an MIS which starts in state  $i$  must end in precisely state  $i$  after  $n$  steps through  $M$ .
- Likewise, for any  $\phi$ , an MIS which starts in state  $i$  must end in precisely state  $\phi(i)$  after  $n$  steps.

## Conjecture (Looking for useful Perron-Frobenius eigenvector properties)

Let  $H$  be a graph and let  $\phi$  be a graph automorphism of  $H$ . Let  $r$  be the principle eigenvalue of the map digraph  $M$  of  $H$ . Then,

$$\lim_{n \rightarrow \infty} \frac{|\text{MIS}((H \square P_{n+1})/\phi)|}{r^n} = 1$$

### Efforts:

- Note that quotienting out by the  $\phi$  does not affect the map digraph. It only affects which pairs of states an MIS can start and end in.
- When  $\phi = \text{id}$ , an MIS which starts in state  $i$  must end in precisely state  $i$  after  $n$  steps through  $M$ .
- Likewise, for any  $\phi$ , an MIS which starts in state  $i$  must end in precisely state  $\phi(i)$  after  $n$  steps.
- Thus, only the ticket digraph of  $G$  is affected.

## Conjecture (Looking for useful Perron-Frobenius eigenvector properties)

Let  $H$  be a graph and let  $\phi$  be a graph automorphism of  $H$ . Let  $r$  be the principle eigenvalue of the map digraph  $M$  of  $H$ . Then,

$$\lim_{n \rightarrow \infty} \frac{|\text{MIS}((H \square P_{n+1})/\phi)|}{r^n} = 1$$

### Efforts:

- Sufficient to show that  $c = 1$ .

$$c = A_{T_\phi} \bullet \frac{\vec{v} \vec{w}^\top}{\vec{w}^\top \vec{v}} = \sum_{1 \leq i \leq m} \frac{\vec{v}_{\phi(i)} \vec{w}_i}{\vec{w}^\top \vec{v}}$$

## Conjecture (Looking for useful Perron-Frobenius eigenvector properties)

Let  $H$  be a graph and let  $\phi$  be a graph automorphism of  $H$ . Let  $r$  be the principle eigenvalue of the map digraph  $M$  of  $H$ . Then,

$$\lim_{n \rightarrow \infty} \frac{|\text{MIS}((H \square P_{n+1})/\phi)|}{r^n} = 1$$

### Efforts:

- Sufficient to show that  $c = 1$ .

$$c = A_{T_\phi} \bullet \frac{\vec{v} \vec{w}^\top}{\vec{w}^\top \vec{v}} = \sum_{1 \leq i \leq m} \frac{\vec{v}_{\phi(i)} \vec{w}_i}{\vec{w}^\top \vec{v}}$$

- (More work needed here) Sufficient to show equality to

$$\sum_{1 \leq i \leq m} \frac{\vec{v}_i \vec{w}_i}{\vec{w}^\top \vec{v}} = 1$$



### Conjecture

For sufficiently large  $m$ ,

$$\lim_{n \rightarrow \infty} \frac{|\text{MIS}(G_{m \times n})|}{|\text{MIS}(FC_{m \times n})|} > 1$$

### Conjecture

For sufficiently large  $m$ ,

$$\lim_{n \rightarrow \infty} \frac{|\text{MIS}(G_{m \times n})|}{|\text{MIS}(FC_{m \times n})|} > 1$$

- Generalize the framework to account for more complex local or global structure?



### Conjecture

For sufficiently large  $m$ ,

$$\lim_{n \rightarrow \infty} \frac{|\text{MIS}(G_{m \times n})|}{|\text{MIS}(FC_{m \times n})|} > 1$$

- Generalize the framework to account for more complex local or global structure?

Thank you!



Recall that  $|\text{MIS}(G_{2 \times n})| = 2F_n$ .

- Can we count only those MISs which are distinct in terms of rotation and reflection?

Recall that  $|\text{MIS}(G_{2 \times n})| = 2F_n$ .

- Can we count only those MISs which are distinct in terms of rotation and reflection?
- This is a good place to start in looking to form more precise arguments about symmetries.

Recall that  $|\text{MIS}(G_{2 \times n})| = 2F_n$ .

- Can we count only those MISs which are distinct in terms of rotation and reflection?
- This is a good place to start in looking to form more precise arguments about symmetries.

## Definition (NIMIS( $G$ ))

Two elements  $I, I'$  are *isomorphic* if there exists a graph automorphism  $\varphi : G \rightarrow G$  with  $\varphi(I) = I'$  and *non-isomorphic* if no such  $\varphi$  exists. Denote the set of non-isomorphic MISs on  $G$  by  $\text{NIMIS}(G)$ .

Recall that  $|\text{MIS}(G_{2 \times n})| = 2F_n$ .

- Can we count only those MISs which are distinct in terms of rotation and reflection?
- This is a good place to start in looking to form more precise arguments about symmetries.

### Definition (NIMIS( $G$ ))

Two elements  $I, I'$  are *isomorphic* if there exists a graph automorphism  $\varphi : G \rightarrow G$  with  $\varphi(I) = I'$  and *non-isomorphic* if no such  $\varphi$  exists. Denote the set of non-isomorphic MISs on  $G$  by  $\text{NIMIS}(G)$ .

- In general, we can use the group of symmetries of our graph to act on the set of MISs. The number of distinct orbits of  $\text{MIS}(G)$  counts  $|\text{NIMIS}(G)|$ .

Recall that  $|\text{MIS}(G_{2 \times n})| = 2F_n$ .

Recall that  $|\text{MIS}(G_{2 \times n})| = 2F_n$ .

## Definition (Bit String Map)

Let  $\psi : \text{MIS}(G_{2 \times n}) \rightarrow \{0, 1\}^n$  be defined by

$$\psi(M)(i) = \begin{cases} 1 & \text{if } (i, 1) \in M \text{ or } (i, 2) \in M \\ 0 & \text{if } (i, 1) \notin M \text{ and } (i, 2) \notin M \end{cases}$$

for every  $1 \leq i \leq n$  and every  $M \in \text{MIS}(G_{2 \times n})$



Recall that  $|\text{MIS}(G_{2 \times n})| = 2F_n$ .

## Definition (Bit String Map)

Let  $\psi : \text{MIS}(G_{2 \times n}) \rightarrow \{0, 1\}^n$  be defined by

$$\psi(M)(i) = \begin{cases} 1 & \text{if } (i, 1) \in M \text{ or } (i, 2) \in M \\ 0 & \text{if } (i, 1) \notin M \text{ and } (i, 2) \notin M \end{cases}$$

for every  $1 \leq i \leq n$  and every  $M \in \text{MIS}(G_{2 \times n})$

- Note  $\psi$  maps precisely 2 MISs to some element in its range. These MISs are duals (from before).

Recall that  $|\text{MIS}(G_{2 \times n})| = 2F_n$ .

## Definition (Bit String Map)

Let  $\psi : \text{MIS}(G_{2 \times n}) \rightarrow \{0, 1\}^n$  be defined by

$$\psi(M)(i) = \begin{cases} 1 & \text{if } (i, 1) \in M \text{ or } (i, 2) \in M \\ 0 & \text{if } (i, 1) \notin M \text{ and } (i, 2) \notin M \end{cases}$$

for every  $1 \leq i \leq n$  and every  $M \in \text{MIS}(G_{2 \times n})$

- Note  $\psi$  maps precisely 2 MISs to some element in its range. These MISs are duals (from before).

## Theorem

For  $n \geq 3$ ,

$$|\text{NIMIS}(G_{2 \times n})| = \begin{cases} \frac{1}{2}(F_n + F_{n/2}) & \text{if } n \text{ is even} \\ \frac{1}{2}(F_n + F_{(n+3)/2}) & \text{if } n \text{ is odd} \end{cases}$$

## Theorem

For  $n \geq 3$ ,

$$|\text{NIMIS}(G_{2 \times n})| = \begin{cases} \frac{1}{2}(F_n + F_{n/2}) & \text{if } n \text{ is even} \\ \frac{1}{2}(F_n + F_{(n+3)/2}) & \text{if } n \text{ is odd} \end{cases}$$

**Proof (sketch, odd  $n$ ):**

- $\psi(|\text{MIS}(G_{2 \times n})|) = F_n$

## Theorem

For  $n \geq 3$ ,

$$|\text{NIMIS}(G_{2 \times n})| = \begin{cases} \frac{1}{2}(F_n + F_{n/2}) & \text{if } n \text{ is even} \\ \frac{1}{2}(F_n + F_{(n+3)/2}) & \text{if } n \text{ is odd} \end{cases}$$

**Proof (sketch, odd  $n$ ):**

- $\psi(|\text{MIS}(G_{2 \times n})|) = F_n$
- Size of orbit of an MIS must divide the group of symmetries on a rectangular grid (Orbit-Stabilizer Theorem). An MIS must have either an orbit of size 2 or 4.

## Theorem

For  $n \geq 3$ ,

$$|\text{NIMIS}(G_{2 \times n})| = \begin{cases} \frac{1}{2}(F_n + F_{n/2}) & \text{if } n \text{ is even} \\ \frac{1}{2}(F_n + F_{(n+3)/2}) & \text{if } n \text{ is odd} \end{cases}$$

**Proof (sketch, odd  $n$ ):**

- $\psi(|\text{MIS}(G_{2 \times n})|) = F_n$
- Size of orbit of an MIS must divide the group of symmetries on a rectangular grid (Orbit-Stabilizer Theorem). An MIS must have either an orbit of size 2 or 4.
- Symmetric bit strings are exactly those with orbit 4.

## Theorem

For  $n \geq 3$ ,

$$|\text{NIMIS}(G_{2 \times n})| = \begin{cases} \frac{1}{2}(F_n + F_{n/2}) & \text{if } n \text{ is even} \\ \frac{1}{2}(F_n + F_{(n+3)/2}) & \text{if } n \text{ is odd} \end{cases}$$

**Proof (sketch, odd  $n$ ):**

- $\psi(|\text{MIS}(G_{2 \times n})|) = F_n$
- Size of orbit of an MIS must divide the group of symmetries on a rectangular grid (Orbit-Stabilizer Theorem). An MIS must have either an orbit of size 2 or 4.
- Symmetric bit strings are exactly those with orbit 4.
- Double-count:

$$|\text{NIMIS}(G_{2 \times n})| = \frac{1}{2}(F_n + \text{number of distinct symmetric strings})$$

## Theorem

For  $n \geq 3$ ,

$$|\text{NIMIS}(G_{2 \times n})| = \begin{cases} \frac{1}{2}(F_n + F_{n/2}) & \text{if } n \text{ is even} \\ \frac{1}{2}(F_n + F_{(n+3)/2}) & \text{if } n \text{ is odd} \end{cases}$$

**Proof (sketch, odd  $n$ ):**

- $\psi(|\text{MIS}(G_{2 \times n})|) = F_n$
- Size of orbit of an MIS must divide the group of symmetries on a rectangular grid (Orbit-Stabilizer Theorem). An MIS must have either an orbit of size 2 or 4.
- Symmetric bit strings are exactly those with orbit 4.
- Double-count:

$$|\text{NIMIS}(G_{2 \times n})| = \frac{1}{2}(F_n + \text{number of distinct symmetric strings})$$

- Number of distinct symmetric strings:

$$F_{(n+1)/2} + F_{(n-1)/2}$$