

# Ramsey Numbers of Crossing-Free Matchings in Ordered Graphs

Zachary C. Hammersla<sup>1</sup>

Luke Hawranick<sup>2</sup>

Kevin G. Milans<sup>3</sup>

TCS Seminar

West Virginia University

October 17, 2025

---

<sup>1</sup>West Virginia University

<sup>2</sup>University of South Carolina

<sup>3</sup>West Virginia University

# Ordered Graphs

- ▶ An **ordered graph** is a graph  $G$  whose vertices  $V(G)$  are linearly ordered.

# Ordered Graphs



- ▶ An **ordered graph** is a graph  $G$  whose vertices  $V(G)$  are linearly ordered.

# Ordered Graphs



- ▶ An **ordered graph** is a graph  $G$  whose vertices  $V(G)$  are linearly ordered.
- ▶  $P_3$ ,  $K_{1,2}$ , and  $K_{2,1}$  are distinct ordered graphs.

# Ordered Graphs




- ▶ An **ordered graph** is a graph  $G$  whose vertices  $V(G)$  are linearly ordered.
- ▶  $P_3$ ,  $K_{1,2}$ , and  $K_{2,1}$  are distinct ordered graphs.
- ▶ If  $G$  and  $H$  are ordered graphs, then  $H \subseteq G$  means there is an order respecting injection  $f: V(H) \rightarrow V(G)$  such that  $uv \in E(H)$  implies  $f(u)f(v) \in E(G)$ .

# Ordered Graphs



- ▶ An **ordered graph** is a graph  $G$  whose vertices  $V(G)$  are linearly ordered.
- ▶  $P_3$ ,  $K_{1,2}$ , and  $K_{2,1}$  are distinct ordered graphs.
- ▶ If  $G$  and  $H$  are ordered graphs, then  $H \subseteq G$  means there is an order respecting injection  $f: V(H) \rightarrow V(G)$  such that  $uv \in E(H)$  implies  $f(u)f(v) \in E(G)$ .

# Ordered Graphs



- ▶ An **ordered graph** is a graph  $G$  whose vertices  $V(G)$  are linearly ordered.
- ▶  $P_3$ ,  $K_{1,2}$ , and  $K_{2,1}$  are distinct ordered graphs.
- ▶ If  $G$  and  $H$  are ordered graphs, then  $H \subseteq G$  means there is an order respecting injection  $f: V(H) \rightarrow V(G)$  such that  $uv \in E(H)$  implies  $f(u)f(v) \in E(G)$ .
- ▶ Note:  $K_{1,2}, K_{2,1} \subseteq K_{3,3}$  but  $P_3 \not\subseteq K_{3,3}$ .

# Matchings in Ordered Graphs

- ▶ In an ordered graph, a pair of edges with distinct endpoints falls into one of 3 types.




# Matchings in Ordered Graphs




- ▶ In an ordered graph, a pair of edges with distinct endpoints falls into one of 3 types.

## Matchings in Ordered Graphs




- In an ordered graph, a pair of edges with distinct endpoints falls into one of 3 types.

## Matchings in Ordered Graphs




- In an ordered graph, a pair of edges with distinct endpoints falls into one of 3 types.

# Matchings in Ordered Graphs




- ▶ In an ordered graph, a pair of edges with distinct endpoints falls into one of 3 types.
- ▶ A set of edges  $S$  is **separation-free**, **cross-free**, or **nest-free** if no pair of edges in  $S$  is separated, crossed, or nested.

# Matchings in Ordered Graphs




- ▶ In an ordered graph, a pair of edges with distinct endpoints falls into one of 3 types.
- ▶ A set of edges  $S$  is **separation-free**, **cross-free**, or **nest-free** if no pair of edges in  $S$  is separated, crossed, or nested.
- ▶ Let  $SF_s$ ,  $CF_s$ ,  $NF_s$  be the family of  $2s$ -vertex ordered graphs whose edge sets are separation-free, cross-free, or nest-free matchings of size  $s$ .


# Matchings in Ordered Graphs




- ▶ In an ordered graph, a pair of edges with distinct endpoints falls into one of 3 types.
- ▶ A set of edges  $S$  is **separation-free**, **cross-free**, or **nest-free** if no pair of edges in  $S$  is separated, crossed, or nested.
- ▶ Let  $SF_s$ ,  $CF_s$ ,  $NF_s$  be the family of  $2s$ -vertex ordered graphs whose edge sets are separation-free, cross-free, or nest-free matchings of size  $s$ .



# Matchings in Ordered Graphs



- ▶ In an ordered graph, a pair of edges with distinct endpoints falls into one of 3 types.
- ▶ A set of edges  $S$  is **separation-free**, **cross-free**, or **nest-free** if no pair of edges in  $S$  is separated, crossed, or nested.
- ▶ Let  $SF_s$ ,  $CF_s$ ,  $NF_s$  be the family of  $2s$ -vertex ordered graphs whose edge sets are separation-free, cross-free, or nest-free matchings of size  $s$ .



- ▶  $|CF_s| = \text{sth Catalan number} = \frac{1}{s+1} \binom{2s}{s}$

## Ordered Ramsey Numbers

- ▶ Given families  $\mathcal{F}_1, \dots, \mathcal{F}_k$  of ordered graphs, the Ramsey number  $R(\mathcal{F}_1, \dots, \mathcal{F}_k)$  is the minimum  $n$  such that every  $k$ -edge-coloring of  $K_n$  contains, for some color  $i$ , a color- $i$  copy of some ordered graph in  $\mathcal{F}_i$ .




## Ordered Ramsey Numbers

- ▶ Given families  $\mathcal{F}_1, \dots, \mathcal{F}_k$  of ordered graphs, the **Ramsey number**  $R(\mathcal{F}_1, \dots, \mathcal{F}_k)$  is the minimum  $n$  such that every  $k$ -edge-coloring of  $K_n$  contains, for some color  $i$ , a color- $i$  copy of some ordered graph in  $\mathcal{F}_i$ .
- ▶ Warm-up:  $R(\text{CF}_2, \text{CF}_2) = 5$ .


# Ordered Ramsey Numbers

- ▶ Given families  $\mathcal{F}_1, \dots, \mathcal{F}_k$  of ordered graphs, the **Ramsey number**  $R(\mathcal{F}_1, \dots, \mathcal{F}_k)$  is the minimum  $n$  such that every  $k$ -edge-coloring of  $K_n$  contains, for some color  $i$ , a color- $i$  copy of some ordered graph in  $\mathcal{F}_i$ .
- ▶ Warm-up:  $R(\text{CF}_2, \text{CF}_2) = 5$ .
- ▶  $R(\text{CF}_2, \text{CF}_2) > 4$ :




# Ordered Ramsey Numbers

- ▶ Given families  $\mathcal{F}_1, \dots, \mathcal{F}_k$  of ordered graphs, the **Ramsey number**  $R(\mathcal{F}_1, \dots, \mathcal{F}_k)$  is the minimum  $n$  such that every  $k$ -edge-coloring of  $K_n$  contains, for some color  $i$ , a color- $i$  copy of some ordered graph in  $\mathcal{F}_i$ .
- ▶ Warm-up:  $R(\text{CF}_2, \text{CF}_2) = 5$ .
- ▶  $R(\text{CF}_2, \text{CF}_2) > 4$ :




- ▶  $R(\text{CF}_2, \text{CF}_2) \leq 5$ :



# Ordered Ramsey Numbers

- ▶ Given families  $\mathcal{F}_1, \dots, \mathcal{F}_k$  of ordered graphs, the **Ramsey number**  $R(\mathcal{F}_1, \dots, \mathcal{F}_k)$  is the minimum  $n$  such that every  $k$ -edge-coloring of  $K_n$  contains, for some color  $i$ , a color- $i$  copy of some ordered graph in  $\mathcal{F}_i$ .
- ▶ Warm-up:  $R(\text{CF}_2, \text{CF}_2) = 5$ .
- ▶  $R(\text{CF}_2, \text{CF}_2) > 4$ :




- ▶  $R(\text{CF}_2, \text{CF}_2) \leq 5$ :



- ▶ If  $v_1 v_2 v_3$  is polychromatic, then one of these edges plus  $v_4 v_5$  forms a monochromatic cross-free matching of size 2.

# Ordered Ramsey Numbers

- ▶ Given families  $\mathcal{F}_1, \dots, \mathcal{F}_k$  of ordered graphs, the **Ramsey number**  $R(\mathcal{F}_1, \dots, \mathcal{F}_k)$  is the minimum  $n$  such that every  $k$ -edge-coloring of  $K_n$  contains, for some color  $i$ , a color- $i$  copy of some ordered graph in  $\mathcal{F}_i$ .
- ▶ Warm-up:  $R(\text{CF}_2, \text{CF}_2) = 5$ .
- ▶  $R(\text{CF}_2, \text{CF}_2) > 4$ :




- ▶  $R(\text{CF}_2, \text{CF}_2) \leq 5$ :




- ▶ If  $v_1 v_2 v_3$  is polychromatic, then one of these edges plus  $v_4 v_5$  forms a monochromatic cross-free matching of size 2.
- ▶ So assume  $v_1 v_2 v_3$  is red.

# Ordered Ramsey Numbers

- ▶ Given families  $\mathcal{F}_1, \dots, \mathcal{F}_k$  of ordered graphs, the **Ramsey number**  $R(\mathcal{F}_1, \dots, \mathcal{F}_k)$  is the minimum  $n$  such that every  $k$ -edge-coloring of  $K_n$  contains, for some color  $i$ , a color- $i$  copy of some ordered graph in  $\mathcal{F}_i$ .
- ▶ Warm-up:  $R(\text{CF}_2, \text{CF}_2) = 5$ .
- ▶  $R(\text{CF}_2, \text{CF}_2) > 4$ :




- ▶  $R(\text{CF}_2, \text{CF}_2) \leq 5$ :




- ▶ If  $v_1 v_2 v_3$  is polychromatic, then one of these edges plus  $v_4 v_5$  forms a monochromatic cross-free matching of size 2.
- ▶ So assume  $v_1 v_2 v_3$  is red.
- ▶ By symmetry,  $v_3 v_4 v_5$  is monochromatic and must be blue.

# Ordered Ramsey Numbers

- ▶ Given families  $\mathcal{F}_1, \dots, \mathcal{F}_k$  of ordered graphs, the **Ramsey number**  $R(\mathcal{F}_1, \dots, \mathcal{F}_k)$  is the minimum  $n$  such that every  $k$ -edge-coloring of  $K_n$  contains, for some color  $i$ , a color- $i$  copy of some ordered graph in  $\mathcal{F}_i$ .
- ▶ Warm-up:  $R(\text{CF}_2, \text{CF}_2) = 5$ .
- ▶  $R(\text{CF}_2, \text{CF}_2) > 4$ :



- ▶  $R(\text{CF}_2, \text{CF}_2) \leq 5$ :



- ▶ If  $v_1 v_2 v_3$  is polychromatic, then one of these edges plus  $v_4 v_5$  forms a monochromatic cross-free matching of size 2.
- ▶ So assume  $v_1 v_2 v_3$  is red.
- ▶ By symmetry,  $v_3 v_4 v_5$  is monochromatic and must be blue.
- ▶ Now  $v_1 v_5$  and one of  $\{v_2 v_3, v_3 v_4\}$  wins.

## Prior Work on Graphs

Theorem (Alon–Frankl–Lovász (1986))

*If  $sK_r^{(r)}$  denotes the  $r$ -uniform hypergraph consisting of  $s$  disjoint edges, then  $R(sK_r^{(r)}; k) = (k - 1)(s - 1) + sr$ .*



## Prior Work on Graphs

### Theorem (Alon–Frankl–Lovász (1986))

If  $sK_r^{(r)}$  denotes the  $r$ -uniform hypergraph consisting of  $s$  disjoint edges, then  $R(sK_r^{(r)}; k) = (k - 1)(s - 1) + sr$ .

### Theorem (Cockayne–Lorimer (1975))

For  $s_1 \geq s_2, \dots, s_k$ , we have

$$R(s_1 K_2, \dots, s_k K_2) = \sum_{i=1}^k (s_i - 1) + (s_1 - 1) + 2.$$

## Prior Work on Graphs

### Theorem (Alon–Frankl–Lovász (1986))

If  $sK_r^{(r)}$  denotes the  $r$ -uniform hypergraph consisting of  $s$  disjoint edges, then  $R(sK_r^{(r)}; k) = (k - 1)(s - 1) + sr$ .

### Theorem (Cockayne–Lorimer (1975))

For  $s_1 \geq s_2, \dots, s_k$ , we have

$$R(s_1 K_2, \dots, s_k K_2) = \sum_{i=1}^k (s_i - 1) + (s_1 - 1) + 2.$$

- ▶ Let  $M_s$  be the family of  $2s$ -vertex ordered graphs whose edge sets are matchings of size  $s$ .

## Prior Work on Graphs

### Theorem (Alon–Frankl–Lovász (1986))

If  $sK_r^{(r)}$  denotes the  $r$ -uniform hypergraph consisting of  $s$  disjoint edges, then  $R(sK_r^{(r)}; k) = (k - 1)(s - 1) + sr$ .

### Theorem (Cockayne–Lorimer (1975))

For  $s_1 \geq s_2, \dots, s_k$ , we have

$$R(s_1 K_2, \dots, s_k K_2) = \sum_{i=1}^k (s_i - 1) + (s_1 - 1) + 2.$$

- ▶ Let  $M_s$  be the family of  $2s$ -vertex ordered graphs whose edge sets are matchings of size  $s$ .
- ▶ Let  $s_1 \geq s_2, \dots, s_k$  and let  $\mathcal{F}_{s_i} \in \{\text{NF}_{s_i}, \text{SF}_{s_i}, \text{CF}_{s_i}\}$ . We have

$$\begin{aligned} R(\mathcal{F}_{s_1}, \dots, \mathcal{F}_{s_k}) &\geq R(M_{s_1}, \dots, M_{s_k}) = R(s_1 K_2, \dots, s_k K_2) \\ &= \sum_{i=1}^k (s_i - 1) + (s_1 - 1) + 2. \end{aligned}$$

## Prior Work on Graphs

### Theorem (Alon–Frankl–Lovász (1986))

If  $sK_r^{(r)}$  denotes the  $r$ -uniform hypergraph consisting of  $s$  disjoint edges, then  $R(sK_r^{(r)}; k) = (k - 1)(s - 1) + sr$ .

### Theorem (Cockayne–Lorimer (1975))

For  $s_1 \geq s_2, \dots, s_k$ , we have

$$R(s_1 K_2, \dots, s_k K_2) = \sum_{i=1}^k (s_i - 1) + (s_1 - 1) + 2.$$

- ▶ Let  $M_s$  be the family of  $2s$ -vertex ordered graphs whose edge sets are matchings of size  $s$ .
- ▶ Let  $s_1 \geq s_2, \dots, s_k$  and let  $\mathcal{F}_{s_i} \in \{\text{NF}_{s_i}, \text{SF}_{s_i}, \text{CF}_{s_i}\}$ . We have

$$\begin{aligned} R(\mathcal{F}_{s_1}, \dots, \mathcal{F}_{s_k}) &\geq R(M_{s_1}, \dots, M_{s_k}) = R(s_1 K_2, \dots, s_k K_2) \\ &= \sum_{i=1}^k (s_i - 1) + (s_1 - 1) + 2. \end{aligned}$$

- ▶ We call this the **Graph Lower Bound (GLB)**.

## Prior Work on Ordered Graphs

- ▶ **Graph Lower Bound (GLB):** If  $s_1 = \max\{s_1, \dots, s_k\}$ , then  $R(\mathcal{F}_{s_1}, \dots, \mathcal{F}_{s_k}) \geq \sum_{i=1}^k (s_i - 1) + (s_1 - 1) + 2$ .

## Prior Work on Ordered Graphs

- ▶ **Graph Lower Bound (GLB)**: If  $s_1 = \max\{s_1, \dots, s_k\}$ , then  $R(\mathcal{F}_{s_1}, \dots, \mathcal{F}_{s_k}) \geq \sum_{i=1}^k (s_i - 1) + (s_1 - 1) + 2$ .
- ▶ BGT: Barát–Gyárfás–Tóth (2023)

## Prior Work on Ordered Graphs

- ▶ **Graph Lower Bound (GLB)**: If  $s_1 = \max\{s_1, \dots, s_k\}$ , then  $R(\mathcal{F}_{s_1}, \dots, \mathcal{F}_{s_k}) \geq \sum_{i=1}^k (s_i - 1) + (s_1 - 1) + 2$ .
- ▶ BGT: Barát–Gyárfás–Tóth (2023)
- ▶ KS: Kaiser–Stehlík (2020)

## Prior Work on Ordered Graphs

- ▶ **Graph Lower Bound (GLB)**: If  $s_1 = \max\{s_1, \dots, s_k\}$ , then  $R(\mathcal{F}_{s_1}, \dots, \mathcal{F}_{s_k}) \geq \sum_{i=1}^k (s_i - 1) + (s_1 - 1) + 2$ .
- ▶ BGT: Barát–Gyárfás–Tóth (2023)
- ▶ KS: Kaiser–Stehlík (2020)
- ▶ KPT: Károlyi–Pach–Tóth (1997)



## Prior Work on Ordered Graphs

- ▶ **Graph Lower Bound (GLB):** If  $s_1 = \max\{s_1, \dots, s_k\}$ , then  $R(\mathcal{F}_{s_1}, \dots, \mathcal{F}_{s_k}) \geq \sum_{i=1}^k (s_i - 1) + (s_1 - 1) + 2$ .
- ▶ BGT: Barát–Gyárfás–Tóth (2023)
- ▶ KS: Kaiser–Stehlík (2020)
- ▶ KPT: Károlyi–Pach–Tóth (1997)

	Nest-free	Separation-free	Cross-free
2 colors $k = 2$			
2 edges $s = 2$			
Gen. case			

## Prior Work on Ordered Graphs

- ▶ **Graph Lower Bound (GLB):** If  $s_1 = \max\{s_1, \dots, s_k\}$ , then  $R(\mathcal{F}_{s_1}, \dots, \mathcal{F}_{s_k}) \geq \sum_{i=1}^k (s_i - 1) + (s_1 - 1) + 2$ .
- ▶ BGT: Barát–Gyárfás–Tóth (2023)
- ▶ KS: Kaiser–Stehlík (2020)
- ▶ KPT: Károlyi–Pach–Tóth (1997)

	Nest-free	Separation-free	Cross-free
2 colors $k = 2$	Thm (BGT) $R(\text{NF}_s, \text{NF}_2) = 2s + 1$ $R(\text{NF}_s, \text{NF}_3) = 2s + 2$		
2 edges $s = 2$			
Gen. case			

## Prior Work on Ordered Graphs

- ▶ **Graph Lower Bound (GLB):** If  $s_1 = \max\{s_1, \dots, s_k\}$ , then  $R(\mathcal{F}_{s_1}, \dots, \mathcal{F}_{s_k}) \geq \sum_{i=1}^k (s_i - 1) + (s_1 - 1) + 2$ .
- ▶ BGT: Barát–Gyárfás–Tóth (2023)
- ▶ KS: Kaiser–Stehlík (2020)
- ▶ KPT: Károlyi–Pach–Tóth (1997)

	Nest-free	Separation-free	Cross-free
2 colors $k = 2$	Thm (BGT) $R(\text{NF}_s, \text{NF}_2) = 2s + 1$ $R(\text{NF}_s, \text{NF}_3) = 2s + 2$		
2 edges $s = 2$	Thm (BGT) $R(\text{NF}_2; k) = k + 3$		
Gen. case			

## Prior Work on Ordered Graphs

- ▶ **Graph Lower Bound (GLB):** If  $s_1 = \max\{s_1, \dots, s_k\}$ , then  $R(\mathcal{F}_{s_1}, \dots, \mathcal{F}_{s_k}) \geq \sum_{i=1}^k (s_i - 1) + (s_1 - 1) + 2$ .
- ▶ BGT: Barát–Gyárfás–Tóth (2023)
- ▶ KS: Kaiser–Stehlík (2020)
- ▶ KPT: Károlyi–Pach–Tóth (1997)

	Nest-free	Separation-free	Cross-free
2 colors $k = 2$	Thm (BGT) $R(\text{NF}_s, \text{NF}_2) = 2s + 1$ $R(\text{NF}_s, \text{NF}_3) = 2s + 2$		
2 edges $s = 2$	Thm (BGT) $R(\text{NF}_2; k) = k + 3$		
Gen. case	Conj (BGT) $R(\text{NF}_{s_1}, \dots, \text{NF}_{s_k})$ equals GLB		

## Prior Work on Ordered Graphs

- ▶ **Graph Lower Bound (GLB):** If  $s_1 = \max\{s_1, \dots, s_k\}$ , then  $R(\mathcal{F}_{s_1}, \dots, \mathcal{F}_{s_k}) \geq \sum_{i=1}^k (s_i - 1) + (s_1 - 1) + 2$ .
- ▶ BGT: Barát–Gyárfás–Tóth (2023)
- ▶ KS: Kaiser–Stehlík (2020)
- ▶ KPT: Károlyi–Pach–Tóth (1997)

	Nest-free	Separation-free	Cross-free
2 colors $k = 2$	Thm (BGT) $R(\text{NF}_s, \text{NF}_2) = 2s + 1$ $R(\text{NF}_s, \text{NF}_3) = 2s + 2$	Thm (BGT) $R(\text{SF}_{s_1}, \text{SF}_{s_2}) = \text{GLB}$	
2 edges $s = 2$	Thm (BGT) $R(\text{NF}_2; k) = k + 3$		
Gen. case	Conj (BGT) $R(\text{NF}_{s_1}, \dots, \text{NF}_{s_k})$ equals GLB		

## Prior Work on Ordered Graphs

- ▶ **Graph Lower Bound (GLB):** If  $s_1 = \max\{s_1, \dots, s_k\}$ , then  $R(\mathcal{F}_{s_1}, \dots, \mathcal{F}_{s_k}) \geq \sum_{i=1}^k (s_i - 1) + (s_1 - 1) + 2$ .
- ▶ BGT: Barát–Gyárfás–Tóth (2023)
- ▶ KS: Kaiser–Stehlík (2020)
- ▶ KPT: Károlyi–Pach–Tóth (1997)

	Nest-free	Separation-free	Cross-free
2 colors $k = 2$	Thm (BGT) $R(\text{NF}_s, \text{NF}_2) = 2s + 1$ $R(\text{NF}_s, \text{NF}_3) = 2s + 2$	Thm (BGT) $R(\text{SF}_{s_1}, \text{SF}_{s_2}) = \text{GLB}$	
2 edges $s = 2$	Thm (BGT) $R(\text{NF}_2; k) = k + 3$	Cor (KS) $R(\text{SF}_2; k) = k + 3$	
Gen. case	Conj (BGT) $R(\text{NF}_{s_1}, \dots, \text{NF}_{s_k})$ equals GLB		

## Prior Work on Ordered Graphs

- ▶ **Graph Lower Bound (GLB):** If  $s_1 = \max\{s_1, \dots, s_k\}$ , then  $R(\mathcal{F}_{s_1}, \dots, \mathcal{F}_{s_k}) \geq \sum_{i=1}^k (s_i - 1) + (s_1 - 1) + 2$ .
- ▶ BGT: Barát–Gyárfás–Tóth (2023)
- ▶ KS: Kaiser–Stehlík (2020)
- ▶ KPT: Károlyi–Pach–Tóth (1997)

	Nest-free	Separation-free	Cross-free
2 colors $k = 2$	Thm (BGT) $R(\text{NF}_s, \text{NF}_2) = 2s + 1$ $R(\text{NF}_s, \text{NF}_3) = 2s + 2$	Thm (BGT) $R(\text{SF}_{s_1}, \text{SF}_{s_2}) = \text{GLB}$	
2 edges $s = 2$	Thm (BGT) $R(\text{NF}_2; k) = k + 3$	Cor (KS) $R(\text{SF}_2; k) = k + 3$	
Gen. case	Conj (BGT) $R(\text{NF}_{s_1}, \dots, \text{NF}_{s_k})$ equals GLB	open	

## Prior Work on Ordered Graphs

- ▶ **Graph Lower Bound (GLB):** If  $s_1 = \max\{s_1, \dots, s_k\}$ , then  $R(\mathcal{F}_{s_1}, \dots, \mathcal{F}_{s_k}) \geq \sum_{i=1}^k (s_i - 1) + (s_1 - 1) + 2$ .
- ▶ BGT: Barát–Gyárfás–Tóth (2023)
- ▶ KS: Kaiser–Stehlík (2020)
- ▶ KPT: Károlyi–Pach–Tóth (1997)

	Nest-free	Separation-free	Cross-free
2 colors $k = 2$	Thm (BGT) $R(\text{NF}_s, \text{NF}_2) = 2s + 1$ $R(\text{NF}_s, \text{NF}_3) = 2s + 2$	Thm (BGT) $R(\text{SF}_{s_1}, \text{SF}_{s_2}) = \text{GLB}$	Cor (KPT) $R(\text{CF}_s, \text{CF}_s) = \text{GLB}$
2 edges $s = 2$	Thm (BGT) $R(\text{NF}_2; k) = k + 3$	Cor (KS) $R(\text{SF}_2; k) = k + 3$	
Gen. case	Conj (BGT) $R(\text{NF}_{s_1}, \dots, \text{NF}_{s_k})$ equals GLB	open	



## Prior Work on Ordered Graphs

- ▶ **Graph Lower Bound (GLB)**: If  $s_1 = \max\{s_1, \dots, s_k\}$ , then  $R(\mathcal{F}_{s_1}, \dots, \mathcal{F}_{s_k}) \geq \sum_{i=1}^k (s_i - 1) + (s_1 - 1) + 2$ .
- ▶ BGT: Barát–Gyárfás–Tóth (2023)
- ▶ KS: Kaiser–Stehlík (2020)
- ▶ KPT: Károlyi–Pach–Tóth (1997)

	Nest-free	Separation-free	Cross-free
2 colors $k = 2$	Thm (BGT) $R(\text{NF}_s, \text{NF}_2) = 2s + 1$ $R(\text{NF}_s, \text{NF}_3) = 2s + 2$	Thm (BGT) $R(\text{SF}_{s_1}, \text{SF}_{s_2}) = \text{GLB}$	Cor (KPT) $R(\text{CF}_s, \text{CF}_s) = \text{GLB}$
2 edges $s = 2$	Thm (BGT) $R(\text{NF}_2; k) = k + 3$	Cor (KS) $R(\text{SF}_2; k) = k + 3$	Prop (BGT) $R(\text{CF}_2; k) > k + 3$
Gen. case	Conj (BGT) $R(\text{NF}_{s_1}, \dots, \text{NF}_{s_k})$ equals GLB	open	

## Prior Work on Ordered Graphs

- ▶ **Graph Lower Bound (GLB)**: If  $s_1 = \max\{s_1, \dots, s_k\}$ , then  $R(\mathcal{F}_{s_1}, \dots, \mathcal{F}_{s_k}) \geq \sum_{i=1}^k (s_i - 1) + (s_1 - 1) + 2$ .
- ▶ BGT: Barát–Gyárfás–Tóth (2023)
- ▶ KS: Kaiser–Stehlík (2020)
- ▶ KPT: Károlyi–Pach–Tóth (1997)

	Nest-free	Separation-free	Cross-free
2 colors $k = 2$	Thm (BGT) $R(\text{NF}_s, \text{NF}_2) = 2s + 1$ $R(\text{NF}_s, \text{NF}_3) = 2s + 2$	Thm (BGT) $R(\text{SF}_{s_1}, \text{SF}_{s_2}) = \text{GLB}$	Cor (KPT) $R(\text{CF}_s, \text{CF}_s) = \text{GLB}$
2 edges $s = 2$	Thm (BGT) $R(\text{NF}_2; k) = k + 3$	Cor (KS) $R(\text{SF}_2; k) = k + 3$	Prop (BGT) $R(\text{CF}_2; k) > k + 3$
Gen. case	Conj (BGT) $R(\text{NF}_{s_1}, \dots, \text{NF}_{s_k})$ equals GLB	open	open

## Prior Work on Ordered Graphs

- **Graph Lower Bound (GLB):** If  $s_1 = \max\{s_1, \dots, s_k\}$ , then  $R(\mathcal{F}_{s_1}, \dots, \mathcal{F}_{s_k}) \geq \sum_{i=1}^k (s_i - 1) + (s_1 - 1) + 2$ .
- BGT: Barát–Gyárfás–Tóth (2023)
- KS: Kaiser–Stehlík (2020)
- KPT: Károlyi–Pach–Tóth (1997)

	Nest-free	Separation-free	Cross-free
2 colors $k = 2$	Thm (BGT) $R(\text{NF}_s, \text{NF}_2) = 2s + 1$ $R(\text{NF}_s, \text{NF}_3) = 2s + 2$	Thm (BGT) $R(\text{SF}_{s_1}, \text{SF}_{s_2}) = \text{GLB}$	Cor (KPT) $R(\text{CF}_s, \text{CF}_s) = \text{GLB}$
2 edges $s = 2$	Thm (BGT) $R(\text{NF}_2; k) = k + 3$	Cor (KS) $R(\text{SF}_2; k) = k + 3$	Prop (BGT) $R(\text{CF}_2; k) > k + 3$
Gen. case	Conj (BGT) $R(\text{NF}_{s_1}, \dots, \text{NF}_{s_k})$ equals GLB	open	open

## Our Results

- Thm:  $R(\text{CF}_2; k) = k + \Theta(\sqrt{k})$

## Prior Work on Ordered Graphs

- ▶ **Graph Lower Bound (GLB)**: If  $s_1 = \max\{s_1, \dots, s_k\}$ , then  $R(\mathcal{F}_{s_1}, \dots, \mathcal{F}_{s_k}) \geq \sum_{i=1}^k (s_i - 1) + (s_1 - 1) + 2$ .
- ▶ BGT: Barát–Gyárfás–Tóth (2023)
- ▶ KS: Kaiser–Stehlík (2020)
- ▶ KPT: Károlyi–Pach–Tóth (1997)

	Nest-free	Separation-free	Cross-free
2 colors $k = 2$	Thm (BGT) $R(\text{NF}_s, \text{NF}_2) = 2s + 1$ $R(\text{NF}_s, \text{NF}_3) = 2s + 2$	Thm (BGT) $R(\text{SF}_{s_1}, \text{SF}_{s_2}) = \text{GLB}$	Cor (KPT) $R(\text{CF}_s, \text{CF}_s) = \text{GLB}$
2 edges $s = 2$	Thm (BGT) $R(\text{NF}_2; k) = k + 3$	Cor (KS) $R(\text{SF}_2; k) = k + 3$	Prop (BGT) $R(\text{CF}_2; k) > k + 3$
Gen. case	Conj (BGT) $R(\text{NF}_{s_1}, \dots, \text{NF}_{s_k})$ equals GLB	open	open

## Our Results

- ▶ Thm:  $R(\text{CF}_2; k) = k + \Theta(\sqrt{k})$
- ▶ Prop:  $R(\text{NF}_s, \text{NF}_4) = \text{GLB} = 2s + 3$  for  $s \geq 4$ .

# Lower Bound

## Theorem

$$R(\text{CF}_2; k) \geq k + (1 - o(1))\sqrt{\frac{4}{\pi}k}$$

# Lower Bound

## Theorem

$$R(\text{CF}_2; k) \geq k + (1 - o(1))\sqrt{\frac{4}{\pi}k}$$

- ▶ To prove this, we cover  $E(K_n)$  for  $n = k + (1 - o(1))\sqrt{(4/\pi)k}$  with  $k$  color classes such that each color class avoids graphs in

$\text{CF}_2$ , where  $\text{CF}_2 = \{ \text{graph with 4 vertices and edges } (1,2), (2,3), (3,4), (1,4) \}, \text{graph with 3 vertices and edges } (1,2), (2,3), (1,3) \}, \text{graph with 3 vertices and edges } (1,2), (2,3), (1,3) \} \}.$


# Lower Bound

## Theorem

$$R(\text{CF}_2; k) \geq k + (1 - o(1))\sqrt{\frac{4}{\pi}k}$$


- ▶ To prove this, we cover  $E(K_n)$  for  $n = k + (1 - o(1))\sqrt{(4/\pi)k}$  with  $k$  color classes such that each color class avoids graphs in  $\text{CF}_2$ , where  $\text{CF}_2 = \{ \text{diagram} \}$ .
- ▶ If  $e$  and  $e'$  have the same color, then  $e$  and  $e'$  are not separated and not nested. Hence they are crossed or share an endpoint.

# Lower Bound

$G_1$ : 

## Theorem

$$R(\text{CF}_2; k) \geq k + (1 - o(1))\sqrt{\frac{4}{\pi}k}$$

- ▶ To prove this, we cover  $E(K_n)$  for  $n = k + (1 - o(1))\sqrt{(4/\pi)k}$  with  $k$  color classes such that each color class avoids graphs in  $\text{CF}_2$ , where  $\text{CF}_2 = \{ \text{diagram} \}$ .  

- ▶ If  $e$  and  $e'$  have the same color, then  $e$  and  $e'$  are not separated and not nested. Hence they are crossed or share an endpoint.
- ▶ Our coloring proceeds in stages, starting at stage 1.



## Lower Bound

$G_1$ :    •    •    •    •    •    •    •    •    •    •    •    •    •    •    •    •    •    •    •  
          1    2    3    4    5    6    7    8    9    10   11   12   13   14   15   16   17   18   19

- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .

## Lower Bound

$G_1$ :    •  
          1    2    3    4    5    6    7    8    9    10   11   12   13   14   15   16   17   18   19

- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :

## Lower Bound



- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .

## Lower Bound



- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .
- ▶ We use a color class  $L_t$  for each  $t \in S$ .

## Lower Bound



- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .
- ▶ We use a color class  $L_t$  for each  $t \in S$ .
- ▶ The color class  $L_t$  contains  $uv \in E(G_i)$  when  $uv$  is incident to  $t$  or  $|u - v| \in \{i, i + 1\}$  and  $t \in [u, v]$ .

# Lower Bound



- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .
- ▶ We use a color class  $L_t$  for each  $t \in S$ .
- ▶ The color class  $L_t$  contains  $uv \in E(G_i)$  when  $uv$  is incident to  $t$  or  $|u - v| \in \{i, i + 1\}$  and  $t \in [u, v]$ .

## Lower Bound



- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .
- ▶ We use a color class  $L_t$  for each  $t \in S$ .
- ▶ The color class  $L_t$  contains  $uv \in E(G_i)$  when  $uv$  is incident to  $t$  or  $|u - v| \in \{i, i + 1\}$  and  $t \in [u, v]$ .

## Lower Bound



- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .
- ▶ We use a color class  $L_t$  for each  $t \in S$ .
- ▶ The color class  $L_t$  contains  $uv \in E(G_i)$  when  $uv$  is incident to  $t$  or  $|u - v| \in \{i, i + 1\}$  and  $t \in [u, v]$ .




## Lower Bound




- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .
- ▶ We use a color class  $L_t$  for each  $t \in S$ .
- ▶ The color class  $L_t$  contains  $uv \in E(G_i)$  when  $uv$  is incident to  $t$  or  $|u - v| \in \{i, i + 1\}$  and  $t \in [u, v]$ .

# Lower Bound




- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .
- ▶ We use a color class  $L_t$  for each  $t \in S$ .
- ▶ The color class  $L_t$  contains  $uv \in E(G_i)$  when  $uv$  is incident to  $t$  or  $|u - v| \in \{i, i + 1\}$  and  $t \in [u, v]$ .

## Lower Bound




- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .
- ▶ We use a color class  $L_t$  for each  $t \in S$ .
- ▶ The color class  $L_t$  contains  $uv \in E(G_i)$  when  $uv$  is incident to  $t$  or  $|u - v| \in \{i, i + 1\}$  and  $t \in [u, v]$ .

## Lower Bound




- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .
- ▶ We use a color class  $L_t$  for each  $t \in S$ .
- ▶ The color class  $L_t$  contains  $uv \in E(G_i)$  when  $uv$  is incident to  $t$  or  $|u - v| \in \{i, i + 1\}$  and  $t \in [u, v]$ .

# Lower Bound




- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .
- ▶ We use a color class  $L_t$  for each  $t \in S$ .
- ▶ The color class  $L_t$  contains  $uv \in E(G_i)$  when  $uv$  is incident to  $t$  or  $|u - v| \in \{i, i + 1\}$  and  $t \in [u, v]$ .

## Lower Bound



- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .
- ▶ We use a color class  $L_t$  for each  $t \in S$ .
- ▶ The color class  $L_t$  contains  $uv \in E(G_i)$  when  $uv$  is incident to  $t$  or  $|u - v| \in \{i, i + 1\}$  and  $t \in [u, v]$ .

## Lower Bound



- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .
- ▶ We use a color class  $L_t$  for each  $t \in S$ .
- ▶ The color class  $L_t$  contains  $uv \in E(G_i)$  when  $uv$  is incident to  $t$  or  $|u - v| \in \{i, i + 1\}$  and  $t \in [u, v]$ .
- ▶ Let  $G_{i+1} = G_i - S - \bigcup L_t$ , renaming vertices.

## Lower Bound



- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .
- ▶ We use a color class  $L_t$  for each  $t \in S$ .
- ▶ The color class  $L_t$  contains  $uv \in E(G_i)$  when  $uv$  is incident to  $t$  or  $|u - v| \in \{i, i + 1\}$  and  $t \in [u, v]$ .
- ▶ Let  $G_{i+1} = G_i - S - \bigcup L_t$ , renaming vertices.




## Lower Bound




- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .
- ▶ We use a color class  $L_t$  for each  $t \in S$ .
- ▶ The color class  $L_t$  contains  $uv \in E(G_i)$  when  $uv$  is incident to  $t$  or  $|u - v| \in \{i, i + 1\}$  and  $t \in [u, v]$ .
- ▶ Let  $G_{i+1} = G_i - S - \bigcup L_t$ , renaming vertices.

## Lower Bound




- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .
- ▶ We use a color class  $L_t$  for each  $t \in S$ .
- ▶ The color class  $L_t$  contains  $uv \in E(G_i)$  when  $uv$  is incident to  $t$  or  $|u - v| \in \{i, i + 1\}$  and  $t \in [u, v]$ .
- ▶ Let  $G_{i+1} = G_i - S - \bigcup L_t$ , renaming vertices.

## Lower Bound




- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .
- ▶ We use a color class  $L_t$  for each  $t \in S$ .
- ▶ The color class  $L_t$  contains  $uv \in E(G_i)$  when  $uv$  is incident to  $t$  or  $|u - v| \in \{i, i + 1\}$  and  $t \in [u, v]$ .
- ▶ Let  $G_{i+1} = G_i - S - \bigcup L_t$ , renaming vertices.

## Lower Bound



- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .
- ▶ We use a color class  $L_t$  for each  $t \in S$ .
- ▶ The color class  $L_t$  contains  $uv \in E(G_i)$  when  $uv$  is incident to  $t$  or  $|u - v| \in \{i, i + 1\}$  and  $t \in [u, v]$ .
- ▶ Let  $G_{i+1} = G_i - S - \bigcup L_t$ , renaming vertices.

## Lower Bound



- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .
- ▶ We use a color class  $L_t$  for each  $t \in S$ .
- ▶ The color class  $L_t$  contains  $uv \in E(G_i)$  when  $uv$  is incident to  $t$  or  $|u - v| \in \{i, i + 1\}$  and  $t \in [u, v]$ .
- ▶ Let  $G_{i+1} = G_i - S - \bigcup L_t$ , renaming vertices.

## Lower Bound




- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .
- ▶ We use a color class  $L_t$  for each  $t \in S$ .
- ▶ The color class  $L_t$  contains  $uv \in E(G_i)$  when  $uv$  is incident to  $t$  or  $|u - v| \in \{i, i + 1\}$  and  $t \in [u, v]$ .
- ▶ Let  $G_{i+1} = G_i - S - \bigcup L_t$ , renaming vertices.

## Lower Bound



- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .
- ▶ We use a color class  $L_t$  for each  $t \in S$ .
- ▶ The color class  $L_t$  contains  $uv \in E(G_i)$  when  $uv$  is incident to  $t$  or  $|u - v| \in \{i, i + 1\}$  and  $t \in [u, v]$ .
- ▶ Let  $G_{i+1} = G_i - S - \bigcup L_t$ , renaming vertices.


## Lower Bound



- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .
- ▶ We use a color class  $L_t$  for each  $t \in S$ .
- ▶ The color class  $L_t$  contains  $uv \in E(G_i)$  when  $uv$  is incident to  $t$  or  $|u - v| \in \{i, i + 1\}$  and  $t \in [u, v]$ .
- ▶ Let  $G_{i+1} = G_i - S - \bigcup L_t$ , renaming vertices.




## Lower Bound



- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .
- ▶ We use a color class  $L_t$  for each  $t \in S$ .
- ▶ The color class  $L_t$  contains  $uv \in E(G_i)$  when  $uv$  is incident to  $t$  or  $|u - v| \in \{i, i + 1\}$  and  $t \in [u, v]$ .
- ▶ Let  $G_{i+1} = G_i - S - \bigcup L_t$ , renaming vertices.

## Lower Bound




- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .
- ▶ We use a color class  $L_t$  for each  $t \in S$ .
- ▶ The color class  $L_t$  contains  $uv \in E(G_i)$  when  $uv$  is incident to  $t$  or  $|u - v| \in \{i, i + 1\}$  and  $t \in [u, v]$ .
- ▶ Let  $G_{i+1} = G_i - S - \bigcup L_t$ , renaming vertices.

## Lower Bound




- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .
- ▶ We use a color class  $L_t$  for each  $t \in S$ .
- ▶ The color class  $L_t$  contains  $uv \in E(G_i)$  when  $uv$  is incident to  $t$  or  $|u - v| \in \{i, i + 1\}$  and  $t \in [u, v]$ .
- ▶ Let  $G_{i+1} = G_i - S - \bigcup L_t$ , renaming vertices.

## Lower Bound



- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .
- ▶ We use a color class  $L_t$  for each  $t \in S$ .
- ▶ The color class  $L_t$  contains  $uv \in E(G_i)$  when  $uv$  is incident to  $t$  or  $|u - v| \in \{i, i + 1\}$  and  $t \in [u, v]$ .
- ▶ Let  $G_{i+1} = G_i - S - \bigcup L_t$ , renaming vertices.

## Lower Bound



- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .
- ▶ We use a color class  $L_t$  for each  $t \in S$ .
- ▶ The color class  $L_t$  contains  $uv \in E(G_i)$  when  $uv$  is incident to  $t$  or  $|u - v| \in \{i, i + 1\}$  and  $t \in [u, v]$ .
- ▶ Let  $G_{i+1} = G_i - S - \bigcup L_t$ , renaming vertices.

## Lower Bound



- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .
- ▶ We use a color class  $L_t$  for each  $t \in S$ .
- ▶ The color class  $L_t$  contains  $uv \in E(G_i)$  when  $uv$  is incident to  $t$  or  $|u - v| \in \{i, i + 1\}$  and  $t \in [u, v]$ .
- ▶ Let  $G_{i+1} = G_i - S - \bigcup L_t$ , renaming vertices.
- ▶ The process ends when  $G_i$  is empty.

## Lower Bound



- ▶ At the start of stage  $i$ , we must color an ordered graph  $G_i$  where  $V(G_i) = [n_i]$  and  $uv \in E(G_i)$  implies  $|u - v| \geq i$ .
- ▶ Stage  $i$ :
- ▶ Let  $S = \{t: (i + 1) \mid t\}$ .
- ▶ We use a color class  $L_t$  for each  $t \in S$ .
- ▶ The color class  $L_t$  contains  $uv \in E(G_i)$  when  $uv$  is incident to  $t$  or  $|u - v| \in \{i, i + 1\}$  and  $t \in [u, v]$ .
- ▶ Let  $G_{i+1} = G_i - S - \bigcup L_t$ , renaming vertices.
- ▶ The process ends when  $G_i$  is empty.
- ▶ Num. colors used:  $n - |V(G_\ell)|$ , where  $\ell$  is the last stage.

# Josephus Sieve

Stage:

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28...

- ▶ Start with the list of positive integers and begin stage 1.



# Josephus Sieve

Stage:

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28  $\cdots$

- ▶ Start with the list of positive integers and begin stage 1.
- ▶ At stage  $i$ , delete from the list entries at indices divisible by  $i + 1$ .

# Josephus Sieve

Stage: 1

1 ~~2~~ 3 ~~4~~ 5 ~~6~~ 7 ~~8~~ 9 ~~10~~ 11 ~~12~~ 13 ~~14~~ 15 ~~16~~ 17 ~~18~~ 19 ~~20~~ 21 ~~22~~ 23 ~~24~~ 25 ~~26~~ 27 ~~28~~  $\cdots$

- ▶ Start with the list of positive integers and begin stage 1.
- ▶ At stage  $i$ , delete from the list entries at indices divisible by  $i + 1$ .

# Josephus Sieve

Stage: 1 2

1 ~~2~~ 3 ~~4~~ ~~5~~ ~~6~~ 7 ~~8~~ 9 ~~10~~ ~~11~~ ~~12~~ 13 ~~14~~ 15 ~~16~~ ~~17~~ ~~18~~ 19 ~~20~~ 21 ~~22~~ ~~23~~ ~~24~~ 25 ~~26~~ 27 ~~28~~...

- ▶ Start with the list of positive integers and begin stage 1.
- ▶ At stage  $i$ , delete from the list entries at indices divisible by  $i + 1$ .

# Josephus Sieve

Stage: 1 2 3

1 ~~2~~ 3 ~~4~~ ~~5~~ ~~6~~ 7 ~~8~~ ~~9~~ ~~10~~ ~~11~~ ~~12~~ 13 ~~14~~ 15 ~~16~~ ~~17~~ ~~18~~ 19 ~~20~~ ~~21~~ ~~22~~ ~~23~~ ~~24~~ 25 ~~26~~ ~~27~~ ~~28~~...

- ▶ Start with the list of positive integers and begin stage 1.
- ▶ At stage  $i$ , delete from the list entries at indices divisible by  $i + 1$ .

# Josephus Sieve

Stage: 1 2 3 4

1 ~~2~~ 3 ~~4~~ ~~5~~ ~~6~~ 7 ~~8~~ ~~9~~ ~~10~~ ~~11~~ ~~12~~ 13 ~~14~~ ~~15~~ ~~16~~ ~~17~~ ~~18~~ 19 ~~20~~ ~~21~~ ~~22~~ ~~23~~ ~~24~~ 25 ~~26~~ ~~27~~ ~~28~~...

- ▶ Start with the list of positive integers and begin stage 1.
- ▶ At stage  $i$ , delete from the list entries at indices divisible by  $i + 1$ .

# Josephus Sieve

Stage: 1 2 3 4 5

1 ~~2~~ 3 ~~4~~ ~~5~~ ~~6~~ 7 ~~8~~ ~~9~~ ~~10~~ ~~11~~ ~~12~~ 13 ~~14~~ ~~15~~ ~~16~~ ~~17~~ ~~18~~ 19 ~~20~~ ~~21~~ ~~22~~ ~~23~~ ~~24~~ ~~25~~ 26 ~~27~~ ~~28~~...

- ▶ Start with the list of positive integers and begin stage 1.
- ▶ At stage  $i$ , delete from the list entries at indices divisible by  $i + 1$ .

# Josephus Sieve

Stage: 1 2 3 4 5

1 ~~2~~ 3 ~~4~~ ~~5~~ ~~6~~ 7 ~~8~~ ~~9~~ ~~10~~ ~~11~~ ~~12~~ 13 ~~14~~ ~~15~~ ~~16~~ ~~17~~ 18 ~~19~~ ~~20~~ ~~21~~ ~~22~~ ~~23~~ ~~24~~ ~~25~~ 26 ~~27~~ ~~28~~...

- ▶ Start with the list of positive integers and begin stage 1.
- ▶ At stage  $i$ , delete from the list entries at indices divisible by  $i + 1$ .
- ▶ After stage  $i$ , the first  $i + 1$  entries will never be deleted.

# Josephus Sieve

Stage: 1 2 3 4 5

① 2 ③ 4 5 6 ⑦ 8 9 10 11 12 ⑬ 14 15 16 17 18 ⑲ 20 21 22 23 24 25 26 ⑳ 28...

- ▶ Start with the list of positive integers and begin stage 1.
- ▶ At stage  $i$ , delete from the list entries at indices divisible by  $i + 1$ .
- ▶ After stage  $i$ , the first  $i + 1$  entries will never be deleted.



# Josephus Sieve

Stage: 1 2 3 4 5

① ~~2~~ ③ ~~4~~ ~~5~~ ~~6~~ ⑦ ~~8~~ ~~9~~ ~~10~~ ~~11~~ ~~12~~ ⑬ ~~14~~ ~~15~~ ~~16~~ ~~17~~ ~~18~~ ⑲ ~~20~~ ~~21~~ ~~22~~ ~~23~~ ~~24~~ ~~25~~ ~~26~~ ⑳ ~~28~~...

- ▶ Start with the list of positive integers and begin stage 1.
- ▶ At stage  $i$ , delete from the list entries at indices divisible by  $i + 1$ .
- ▶ After stage  $i$ , the first  $i + 1$  entries will never be deleted.
- ▶ The numbers that survive form a sequence called the **Josephus Sieve**: 1, 3, 7, 13, 19, 27, 39, ...

# Josephus Sieve

Stage: 1 2 3 4 5

① 2 ③ 4 5 6 ⑦ 8 9 10 11 12 ⑬ 14 15 16 17 18 ⑲ 20 21 22 23 24 25 26 ⑲ 28...

- A vertex  $u \in V(G_1)$  survives to the end if and only if  $u$  appears in the Josephus Sieve.

# Josephus Sieve

Stage: 1 2 3 4 5

① 2 ③ 4 5 6 ⑦ 8 9 10 11 12 ⑬ 14 15 16 17 18 ⑲ 20 21 22 23 24 25 26 ⑲ 28...

- ▶ A vertex  $u \in V(G_1)$  survives to the end if and only if  $u$  appears in the Josephus Sieve.
- ▶ Let  $\Psi(n)$  be the number of entries in the Josephus Sieve that are at most  $n$ .

# Josephus Sieve

Stage: 1 2 3 4 5

① 2 ③ 4 5 6 ⑦ 8 9 10 11 12 ⑬ 14 15 16 17 18 ⑲ 20 21 22 23 24 25 26 ⑲ 28...

- ▶ A vertex  $u \in V(G_1)$  survives to the end if and only if  $u$  appears in the Josephus Sieve.
- ▶ Let  $\Psi(n)$  be the number of entries in the Josephus Sieve that are at most  $n$ .
- ▶ Thm: If  $k \geq n - \Psi(n)$ , then  $R(\text{CF}_2; k) > n$ .

# Josephus Sieve

Stage: 1 2 3 4 5

① 2 ③ 4 5 6 ⑦ 8 9 10 11 12 ⑬ 14 15 16 17 18 ⑲ 20 21 22 23 24 25 26 ⑲ 28...

- ▶ A vertex  $u \in V(G_1)$  survives to the end if and only if  $u$  appears in the Josephus Sieve.
- ▶ Let  $\Psi(n)$  be the number of entries in the Josephus Sieve that are at most  $n$ .
- ▶ Thm: If  $k \geq n - \Psi(n)$ , then  $R(\text{CF}_2; k) > n$ .
- ▶ Thm (Andersson (1998)):  $\Psi(n) = \sqrt{(4/\pi)n} + O(n^{1/6})$ .

# Josephus Sieve

Stage: 1 2 3 4 5

① 2 ③ 4 5 6 ⑦ 8 9 10 11 12 ⑬ 14 15 16 17 18 ⑲ 20 21 22 23 24 25 26 ⑲ 28...

- ▶ A vertex  $u \in V(G_1)$  survives to the end if and only if  $u$  appears in the Josephus Sieve.
- ▶ Let  $\Psi(n)$  be the number of entries in the Josephus Sieve that are at most  $n$ .
- ▶ Thm: If  $k \geq n - \Psi(n)$ , then  $R(\text{CF}_2; k) > n$ .
- ▶ Thm (Andersson (1998)):  $\Psi(n) = \sqrt{(4/\pi)n} + O(n^{1/6})$ .
- ▶ Cor:  $R(\text{CF}_2; k) \geq k + (1 - o(1))\sqrt{(4/\pi)k}$ .

## Upper bound

- ▶ An edge-coloring of an ordered graph  $G$  is **good** if every color class avoids subgraphs in  $CF_2$ .

## Upper bound

- ▶ An edge-coloring of an ordered graph  $G$  is **good** if every color class avoids subgraphs in  $\text{CF}_2$ .

### Lemma (Compression)

*Let  $G$  be a good  $k$ -edge-coloring of an  $n$ -vertex ordered graph with  $m$  non-edges. If  $k > 0$ , then there is a good  $(k - 1)$ -edge-coloring of an  $(n - 1)$ -vertex ordered graph with at most  $m + 1$  non-edges.*



# Upper bound

- ▶ An edge-coloring of an ordered graph  $G$  is **good** if every color class avoids subgraphs in  $\text{CF}_2$ .

## Lemma (Compression)

*Let  $G$  be a good  $k$ -edge-coloring of an  $n$ -vertex ordered graph with  $m$  non-edges. If  $k > 0$ , then there is a good  $(k - 1)$ -edge-coloring of an  $(n - 1)$ -vertex ordered graph with at most  $m + 1$  non-edges.*

## Theorem

*If  $\binom{n-k}{2} > k$ , then  $R(\text{CF}_2; k) \leq n$ .*

## Upper bound

- An edge-coloring of an ordered graph  $G$  is **good** if every color class avoids subgraphs in  $\text{CF}_2$ .

### Lemma (Compression)

*Let  $G$  be a good  $k$ -edge-coloring of an  $n$ -vertex ordered graph with  $m$  non-edges. If  $k > 0$ , then there is a good  $(k - 1)$ -edge-coloring of an  $(n - 1)$ -vertex ordered graph with at most  $m + 1$  non-edges.*

### Theorem

*If  $\binom{n-k}{2} > k$ , then  $R(\text{CF}_2; k) \leq n$ .*

### Corollary

$$R(\text{CF}_2; k) \leq k + 1 + \left\lceil \sqrt{2k} \right\rceil$$

# Compression Lemma Illustration



- ▶ Let  $uv$  be a **minimal** edge in  $G$ , meaning  $G$  has no other edge with both endpoints in  $[u, v]$ .

## Compression Lemma Illustration




- ▶ Let  $uv$  be a **minimal** edge in  $G$ , meaning  $G$  has no other edge with both endpoints in  $[u, v]$ .

## Compression Lemma Illustration




- ▶ Let  $uv$  be a **minimal** edge in  $G$ , meaning  $G$  has no other edge with both endpoints in  $[u, v]$ .
- ▶ Let  $\alpha$  be the color of  $uv$ , let  $S$  be the set of edges with color  $\alpha$ , and let  $I = [u, v]$ .

# Compression Lemma Illustration




- ▶ Let  $uv$  be a **minimal** edge in  $G$ , meaning  $G$  has no other edge with both endpoints in  $[u, v]$ .
- ▶ Let  $\alpha$  be the color of  $uv$ , let  $S$  be the set of edges with color  $\alpha$ , and let  $I = [u, v]$ .

## Compression Lemma Illustration



- ▶ Let  $uv$  be a **minimal** edge in  $G$ , meaning  $G$  has no other edge with both endpoints in  $[u, v]$ .
- ▶ Let  $\alpha$  be the color of  $uv$ , let  $S$  be the set of edges with color  $\alpha$ , and let  $I = [u, v]$ .
- ▶ Since  $S$  has no separated or nested edge pairs, every edge in  $S$  besides  $uv$  has exactly one endpoint in  $I$ .


## Compression Lemma Illustration



- ▶ After perhaps recoloring some edges with color  $\alpha$ , we may assume  $N_\alpha(w)$  is a nonempty subinterval of  $I$  for each  $w \notin I$ .





# Compression Lemma Illustration





- After perhaps recoloring some edges with color  $\alpha$ , we may assume  $N_\alpha(w)$  is a nonempty subinterval of  $I$  for each  $w \notin I$ .

# Compression Lemma Illustration





- After perhaps recoloring some edges with color  $\alpha$ , we may assume  $N_\alpha(w)$  is a nonempty subinterval of  $I$  for each  $w \notin I$ .

# Compression Lemma Illustration





- ▶ After perhaps recoloring some edges with color  $\alpha$ , we may assume  $N_\alpha(w)$  is a nonempty subinterval of  $I$  for each  $w \notin I$ .
- ▶ Make a new graph  $G'$  where  $I$  is replaced by an independent set  $J$  with  $|J| = |I| - 1$ .

# Compression Lemma Illustration





- ▶ After perhaps recoloring some edges with color  $\alpha$ , we may assume  $N_\alpha(w)$  is a nonempty subinterval of  $I$  for each  $w \notin I$ .
- ▶ Make a new graph  $G'$  where  $I$  is replaced by an independent set  $J$  with  $|J| = |I| - 1$ .
- ▶ In  $G'$ , we color edges between  $w$  and  $J$  by “deleting” a vertex  $z_w \in N_\alpha(w)$ .

# Compression Lemma Illustration





- ▶ After perhaps recoloring some edges with color  $\alpha$ , we may assume  $N_\alpha(w)$  is a nonempty subinterval of  $I$  for each  $w \notin I$ .
- ▶ Make a new graph  $G'$  where  $I$  is replaced by an independent set  $J$  with  $|J| = |I| - 1$ .
- ▶ In  $G'$ , we color edges between  $w$  and  $J$  by “deleting” a vertex  $z_w \in N_\alpha(w)$ .

# Compression Lemma Illustration





- ▶ After perhaps recoloring some edges with color  $\alpha$ , we may assume  $N_\alpha(w)$  is a nonempty subinterval of  $I$  for each  $w \notin I$ .
- ▶ Make a new graph  $G'$  where  $I$  is replaced by an independent set  $J$  with  $|J| = |I| - 1$ .
- ▶ In  $G'$ , we color edges between  $w$  and  $J$  by “deleting” a vertex  $z_w \in N_\alpha(w)$ .

# Compression Lemma Illustration



- ▶ After perhaps recoloring some edges with color  $\alpha$ , we may assume  $N_\alpha(w)$  is a nonempty subinterval of  $I$  for each  $w \notin I$ .
- ▶ Make a new graph  $G'$  where  $I$  is replaced by an independent set  $J$  with  $|J| = |I| - 1$ .
- ▶ In  $G'$ , we color edges between  $w$  and  $J$  by “deleting” a vertex  $z_w \in N_\alpha(w)$ .
- ▶ Edges incident to  $w$  in color  $\alpha$  besides  $wz_w$  become non-edges.



# Compression Lemma Illustration



- ▶ After perhaps recoloring some edges with color  $\alpha$ , we may assume  $N_\alpha(w)$  is a nonempty subinterval of  $I$  for each  $w \notin I$ .
- ▶ Make a new graph  $G'$  where  $I$  is replaced by an independent set  $J$  with  $|J| = |I| - 1$ .
- ▶ In  $G'$ , we color edges between  $w$  and  $J$  by “deleting” a vertex  $z_w \in N_\alpha(w)$ .
- ▶ Edges incident to  $w$  in color  $\alpha$  besides  $wz_w$  become non-edges.





# Compression Lemma Illustration





- ▶ After perhaps recoloring some edges with color  $\alpha$ , we may assume  $N_\alpha(w)$  is a nonempty subinterval of  $I$  for each  $w \notin I$ .
- ▶ Make a new graph  $G'$  where  $I$  is replaced by an independent set  $J$  with  $|J| = |I| - 1$ .
- ▶ In  $G'$ , we color edges between  $w$  and  $J$  by “deleting” a vertex  $z_w \in N_\alpha(w)$ .
- ▶ Edges incident to  $w$  in color  $\alpha$  besides  $wz_w$  become non-edges.
- ▶ Edges incident to  $w$  in other colors are copied in  $G'$ .

# Compression Lemma Illustration





- ▶ After perhaps recoloring some edges with color  $\alpha$ , we may assume  $N_\alpha(w)$  is a nonempty subinterval of  $I$  for each  $w \notin I$ .
- ▶ Make a new graph  $G'$  where  $I$  is replaced by an independent set  $J$  with  $|J| = |I| - 1$ .
- ▶ In  $G'$ , we color edges between  $w$  and  $J$  by “deleting” a vertex  $z_w \in N_\alpha(w)$ .
- ▶ Edges incident to  $w$  in color  $\alpha$  besides  $wz_w$  become non-edges.
- ▶ Edges incident to  $w$  in other colors are copied in  $G'$ .

## Compression Lemma Illustration





- ▶ After perhaps recoloring some edges with color  $\alpha$ , we may assume  $N_\alpha(w)$  is a nonempty subinterval of  $I$  for each  $w \notin I$ .
- ▶ Make a new graph  $G'$  where  $I$  is replaced by an independent set  $J$  with  $|J| = |I| - 1$ .
- ▶ In  $G'$ , we color edges between  $w$  and  $J$  by “deleting” a vertex  $z_w \in N_\alpha(w)$ .
- ▶ Edges incident to  $w$  in color  $\alpha$  besides  $wz_w$  become non-edges.
- ▶ Edges incident to  $w$  in other colors are copied in  $G'$ .

## Compression Lemma Illustration





- ▶ After perhaps recoloring some edges with color  $\alpha$ , we may assume  $N_\alpha(w)$  is a nonempty subinterval of  $I$  for each  $w \notin I$ .
- ▶ Make a new graph  $G'$  where  $I$  is replaced by an independent set  $J$  with  $|J| = |I| - 1$ .
- ▶ In  $G'$ , we color edges between  $w$  and  $J$  by “deleting” a vertex  $z_w \in N_\alpha(w)$ .
- ▶ Edges incident to  $w$  in color  $\alpha$  besides  $wz_w$  become non-edges.
- ▶ Edges incident to  $w$  in other colors are copied in  $G'$ .

# Compression Lemma Illustration





- ▶ Another vertex  $w' \notin I$  may delete a different  $z_{w'} \in N_\alpha(w')$ .

# Compression Lemma Illustration





- ▶ Another vertex  $w' \notin I$  may delete a different  $z_{w'} \in N_\alpha(w')$ .

# Compression Lemma Illustration



- ▶ Another vertex  $w' \notin I$  may delete a different  $z_{w'} \in N_\alpha(w')$ .



# Compression Lemma Illustration



- ▶ Another vertex  $w' \notin I$  may delete a different  $z_{w'} \in N_\alpha(w')$ .





# Compression Lemma Illustration





- ▶ Another vertex  $w' \notin I$  may delete a different  $z_{w'} \in N_\alpha(w')$ .

# Compression Lemma Illustration





- ▶ Another vertex  $w' \notin I$  may delete a different  $z_{w'} \in N_\alpha(w')$ .

# Compression Lemma Illustration



- ▶ Another vertex  $w' \notin I$  may delete a different  $z_{w'} \in N_\alpha(w')$ .
- ▶ All color classes in  $G'$  still avoid subgraphs in  $\text{CF}_2$ .

# Compression Lemma Illustration



- ▶ Another vertex  $w' \notin I$  may delete a different  $z_{w'} \in N_\alpha(w')$ .
- ▶ All color classes in  $G'$  still avoid subgraphs in  $\text{CF}_2$ .
- ▶ New non-edges in  $G'$  corresponding to  $\alpha$  edges in  $G$  are almost offset by replacing the near-independent set  $I$  in  $G$  with the independent set  $J$  in  $G'$  since  $|J| = |I| - 1$ .

## Open Problems

- Improve the constant on the  $\sqrt{k}$  term in

$$k + (1 - o(1))\sqrt{(4/\pi)k} \leq R(\text{CF}_2; k) \leq k + \left\lceil \sqrt{2k} \right\rceil + 1.$$

## Open Problems

- ▶ Improve the constant on the  $\sqrt{k}$  term in

$$k + (1 - o(1))\sqrt{(4/\pi)k} \leq R(\text{CF}_2; k) \leq k + \left\lceil \sqrt{2k} \right\rceil + 1.$$

- ▶  $R(\text{CF}_s; k)$  for fixed  $s$  when  $s \geq 3$ :

## Open Problems

- ▶ Improve the constant on the  $\sqrt{k}$  term in

$$k + (1 - o(1))\sqrt{(4/\pi)k} \leq R(\text{CF}_2; k) \leq k + \left\lceil \sqrt{2k} \right\rceil + 1.$$

- ▶  $R(\text{CF}_s; k)$  for fixed  $s$  when  $s \geq 3$ :

- ▶ Prop:

$$(s-1)k + (1 - o(1))\sqrt{\frac{4}{\pi}(s-1)k} \leq R(\text{CF}_s; k) \leq 2(s-1)k + 2$$

# Open Problems

- ▶ Improve the constant on the  $\sqrt{k}$  term in

$$k + (1 - o(1))\sqrt{(4/\pi)k} \leq R(\text{CF}_2; k) \leq k + \lceil \sqrt{2k} \rceil + 1.$$

- ▶  $R(\text{CF}_s; k)$  for fixed  $s$  when  $s \geq 3$ :

- ▶ Prop:

$$(s-1)k + (1 - o(1))\sqrt{\frac{4}{\pi}(s-1)k} \leq R(\text{CF}_s; k) \leq 2(s-1)k + 2$$

- ▶ Conj: for each fixed  $s$ , we have  $R(\text{CF}_s; k) = (s-1)k + \Theta(\sqrt{k})$ .



## Open Problems

- ▶ Improve the constant on the  $\sqrt{k}$  term in

$$k + (1 - o(1))\sqrt{(4/\pi)k} \leq R(\text{CF}_2; k) \leq k + \lceil \sqrt{2k} \rceil + 1.$$

- ▶  $R(\text{CF}_s; k)$  for fixed  $s$  when  $s \geq 3$ :

- ▶ Prop:

$$(s-1)k + (1 - o(1))\sqrt{\frac{4}{\pi}(s-1)k} \leq R(\text{CF}_s; k) \leq 2(s-1)k + 2$$

- ▶ Conj: for each fixed  $s$ , we have  $R(\text{CF}_s; k) = (s-1)k + \Theta(\sqrt{k})$ .
  - ▶ If  $s = 3$ , then  $2k + (1 - o(1))\sqrt{(8/\pi)k} \leq R(\text{CF}_3; k) \leq 4k + 2$

## Open Problems

- ▶ Improve the constant on the  $\sqrt{k}$  term in

$$k + (1 - o(1))\sqrt{(4/\pi)k} \leq R(\text{CF}_2; k) \leq k + \lceil \sqrt{2k} \rceil + 1.$$

- ▶  $R(\text{CF}_s; k)$  for fixed  $s$  when  $s \geq 3$ :

- ▶ Prop:

$$(s-1)k + (1 - o(1))\sqrt{\frac{4}{\pi}(s-1)k} \leq R(\text{CF}_s; k) \leq 2(s-1)k + 2$$

- ▶ Conj: for each fixed  $s$ , we have  $R(\text{CF}_s; k) = (s-1)k + \Theta(\sqrt{k})$ .
  - ▶ If  $s = 3$ , then  $2k + (1 - o(1))\sqrt{(8/\pi)k} \leq R(\text{CF}_3; k) \leq 4k + 2$
  - ▶ Our techniques might show that  $R(\text{CF}_3; k) \leq 3k + O(\sqrt{k})$

## Open Problems

- ▶ Improve the constant on the  $\sqrt{k}$  term in

$$k + (1 - o(1))\sqrt{(4/\pi)k} \leq R(\text{CF}_2; k) \leq k + \lceil \sqrt{2k} \rceil + 1.$$

- ▶  $R(\text{CF}_s; k)$  for fixed  $s$  when  $s \geq 3$ :

- ▶ Prop:

$$(s-1)k + (1 - o(1))\sqrt{\frac{4}{\pi}(s-1)k} \leq R(\text{CF}_s; k) \leq 2(s-1)k + 2$$

- ▶ Conj: for each fixed  $s$ , we have  $R(\text{CF}_s; k) = (s-1)k + \Theta(\sqrt{k})$ .

- ▶ If  $s = 3$ , then  $2k + (1 - o(1))\sqrt{(8/\pi)k} \leq R(\text{CF}_3; k) \leq 4k + 2$

- ▶ Our techniques might show that  $R(\text{CF}_3; k) \leq 3k + O(\sqrt{k})$

- ▶ Let  $s_1 \geq s_2$ . Improve bounds on

$$2s_1 + s_2 - 1 \leq R(\text{NF}_{s_1}, \text{NF}_{s_2}) \leq 2(s_1 + s_2 - 1) - O(1).$$

# Open Problems

- ▶ Improve the constant on the  $\sqrt{k}$  term in

$$k + (1 - o(1))\sqrt{(4/\pi)k} \leq R(\text{CF}_2; k) \leq k + \lceil \sqrt{2k} \rceil + 1.$$

- ▶  $R(\text{CF}_s; k)$  for fixed  $s$  when  $s \geq 3$ :

- ▶ Prop:

$$(s-1)k + (1 - o(1))\sqrt{\frac{4}{\pi}(s-1)k} \leq R(\text{CF}_s; k) \leq 2(s-1)k + 2$$

- ▶ Conj: for each fixed  $s$ , we have  $R(\text{CF}_s; k) = (s-1)k + \Theta(\sqrt{k})$ .

- ▶ If  $s = 3$ , then  $2k + (1 - o(1))\sqrt{(8/\pi)k} \leq R(\text{CF}_3; k) \leq 4k + 2$

- ▶ Our techniques might show that  $R(\text{CF}_3; k) \leq 3k + O(\sqrt{k})$

- ▶ Let  $s_1 \geq s_2$ . Improve bounds on

$$2s_1 + s_2 - 1 \leq R(\text{NF}_{s_1}, \text{NF}_{s_2}) \leq 2(s_1 + s_2 - 1) - O(1).$$

- ▶ Conj (BGT):

$$R(\text{NF}_{s_1}, \dots, \text{NF}_{s_k}) = \text{GLB} = \sum_{i=1}^k (s_i - 1) + (s_1 - 1) + 2$$

## Open Problems

- ▶ Improve the constant on the  $\sqrt{k}$  term in

$$k + (1 - o(1))\sqrt{(4/\pi)k} \leq R(\text{CF}_2; k) \leq k + \lceil \sqrt{2k} \rceil + 1.$$

- ▶  $R(\text{CF}_s; k)$  for fixed  $s$  when  $s \geq 3$ :

- ▶ Prop:

$$(s-1)k + (1 - o(1))\sqrt{\frac{4}{\pi}(s-1)k} \leq R(\text{CF}_s; k) \leq 2(s-1)k + 2$$

- ▶ Conj: for each fixed  $s$ , we have  $R(\text{CF}_s; k) = (s-1)k + \Theta(\sqrt{k})$ .

- ▶ If  $s = 3$ , then  $2k + (1 - o(1))\sqrt{(8/\pi)k} \leq R(\text{CF}_3; k) \leq 4k + 2$

- ▶ Our techniques might show that  $R(\text{CF}_3; k) \leq 3k + O(\sqrt{k})$

- ▶ Let  $s_1 \geq s_2$ . Improve bounds on

$$2s_1 + s_2 - 1 \leq R(\text{NF}_{s_1}, \text{NF}_{s_2}) \leq 2(s_1 + s_2 - 1) - O(1).$$

- ▶ Conj (BGT):

$$R(\text{NF}_{s_1}, \dots, \text{NF}_{s_k}) = \text{GLB} = \sum_{i=1}^k (s_i - 1) + (s_1 - 1) + 2$$

- ▶  $R(\text{SF}_{s_1}, \dots, \text{SF}_{s_k})$ ?

# Open Problems

- ▶ Improve the constant on the  $\sqrt{k}$  term in

$$k + (1 - o(1))\sqrt{(4/\pi)k} \leq R(\text{CF}_2; k) \leq k + \lceil \sqrt{2k} \rceil + 1.$$

- ▶  $R(\text{CF}_s; k)$  for fixed  $s$  when  $s \geq 3$ :

- ▶ Prop:

$$(s-1)k + (1 - o(1))\sqrt{\frac{4}{\pi}(s-1)k} \leq R(\text{CF}_s; k) \leq 2(s-1)k + 2$$

- ▶ Conj: for each fixed  $s$ , we have  $R(\text{CF}_s; k) = (s-1)k + \Theta(\sqrt{k})$ .

- ▶ If  $s = 3$ , then  $2k + (1 - o(1))\sqrt{(8/\pi)k} \leq R(\text{CF}_3; k) \leq 4k + 2$

- ▶ Our techniques might show that  $R(\text{CF}_3; k) \leq 3k + O(\sqrt{k})$

- ▶ Let  $s_1 \geq s_2$ . Improve bounds on

$$2s_1 + s_2 - 1 \leq R(\text{NF}_{s_1}, \text{NF}_{s_2}) \leq 2(s_1 + s_2 - 1) - O(1).$$

- ▶ Conj (BGT):

$$R(\text{NF}_{s_1}, \dots, \text{NF}_{s_k}) = \text{GLB} = \sum_{i=1}^k (s_i - 1) + (s_1 - 1) + 2$$

- ▶  $R(\text{SF}_{s_1}, \dots, \text{SF}_{s_k})$ ?

- ▶ Mixed families, e.g.  $R(\text{CF}_s, \text{NF}_t)$ ?

# Open Problems

- ▶ Improve the constant on the  $\sqrt{k}$  term in

$$k + (1 - o(1))\sqrt{(4/\pi)k} \leq R(\text{CF}_2; k) \leq k + \lceil \sqrt{2k} \rceil + 1.$$

- ▶  $R(\text{CF}_s; k)$  for fixed  $s$  when  $s \geq 3$ :

- ▶ Prop:

$$(s-1)k + (1 - o(1))\sqrt{\frac{4}{\pi}(s-1)k} \leq R(\text{CF}_s; k) \leq 2(s-1)k + 2$$

- ▶ Conj: for each fixed  $s$ , we have  $R(\text{CF}_s; k) = (s-1)k + \Theta(\sqrt{k})$ .

- ▶ If  $s = 3$ , then  $2k + (1 - o(1))\sqrt{(8/\pi)k} \leq R(\text{CF}_3; k) \leq 4k + 2$

- ▶ Our techniques might show that  $R(\text{CF}_3; k) \leq 3k + O(\sqrt{k})$

- ▶ Let  $s_1 \geq s_2$ . Improve bounds on

$$2s_1 + s_2 - 1 \leq R(\text{NF}_{s_1}, \text{NF}_{s_2}) \leq 2(s_1 + s_2 - 1) - O(1).$$

- ▶ Conj (BGT):

$$R(\text{NF}_{s_1}, \dots, \text{NF}_{s_k}) = \text{GLB} = \sum_{i=1}^k (s_i - 1) + (s_1 - 1) + 2$$

- ▶  $R(\text{SF}_{s_1}, \dots, \text{SF}_{s_k})$ ?

- ▶ Mixed families, e.g.  $R(\text{CF}_s, \text{NF}_t)$ ?

Thank You