

On a Rainbow Extremal Number in the Hypercube

Kenny Adeogun¹ Luke Hawranick² Muchen Ju³ Henry Zhan⁴

¹Carnegie Mellon University

²West Virginia University

³Fudan University

⁴Middlebury College

USS XML Presentation Session
IAS Park City Mathematics Institute, Park City, UT
July 22, 2025

Preliminaries

Definition (Hypercube Q_k)

The **hypercube** is a graph with vertex set $\{0, 1\}^k$ such that two k -tuples are adjacent if and only if they differ in exactly one position.




Figure: Q_3 .

Preliminaries

Definition (Proper Edge-coloring)

A **proper edge-coloring** of a graph is a labeling of its edges such that incident edges receive different colors.

Preliminaries

Definition (Proper Edge-coloring)

A **proper edge-coloring** of a graph is a labeling of its edges such that incident edges receive different colors.




Figure: Proper Edge-coloring of Q_3 .

Preliminaries

Definition (Rainbow Edge-coloring)

A **rainbow edge-coloring** of a graph is a labeling of its edges such that each edge receives a different color.

Preliminaries

Definition (Rainbow Edge-coloring)

A **rainbow edge-coloring** of a graph is a labeling of its edges such that each edge receives a different color.



Figure: Rainbow Edge-colored P_4 .

Preliminaries

Definition (The Rainbow Extremal Number of F)

Given a graph H , let $ex^*(n, F)$ be maximum number of edges in an n -vertex graph which admits a proper edge coloring with no rainbow copy of F .

Preliminaries

Definition (The Rainbow Extremal Number of F)

Given a graph H , let $\text{ex}^*(n, F)$ be maximum number of edges in an n -vertex graph which admits a proper edge coloring with no rainbow copy of F .

Theorem (Keevash–Mubayi–Sudakov–Verstraëte, 2007)

Let G be a fixed graph. Then,

$$\text{ex}(n, G) \leq \text{ex}^*(n, G) \leq \text{ex}(n, G) + o(n^2).$$

Relative Rainbow Extremal Numbers

Definition (The Rainbow Extremal Number of F with respect to G)

Given a graph F , let $\text{ex}^*(G, F)$ be the maximum number of edges in a subgraph of G which admits a proper coloring with no rainbow copy of F .

Relative Rainbow Extremal Numbers

Definition (The Rainbow Extremal Number of F with respect to G)

Given a graph F , let $\text{ex}^*(G, F)$ be the maximum number of edges in a subgraph of G which admits a proper coloring with no rainbow copy of F .

Question

Determine $\text{ex}^(Q_n, P_{n+1})$.*

Existing Bounds

Theorem (Rombach, unpublished)

$$ex^*(Q_n, P_{n+1}) < n2^{n-1} = |E(Q_n)|.$$

Existing Bounds

Theorem (Rombach, unpublished)

$$ex^*(Q_n, P_{n+1}) < n2^{n-1} = |E(Q_n)|.$$

Proof (sketch).

- Let $s(v_i, v_j) = v_i + v_j \in \{e_i\}_{i \in [n]}$ for $(v_i, v_j) \in E(Q_n)$.

Existing Bounds

Theorem (Rombach, unpublished)

$$ex^*(Q_n, P_{n+1}) < n2^{n-1} = |E(Q_n)|.$$

Proof (sketch).

- Let $s(v_i, v_j) = v_i + v_j \in \{e_i\}_{i \in [n]}$ for $(v_i, v_j) \in E(Q_n)$.
- Greedily build a rainbow path with an arbitrary start $v_0 v_1$.

Existing Bounds

Theorem (Rombach, unpublished)

$$ex^*(Q_n, P_{n+1}) < n2^{n-1} = |E(Q_n)|.$$

Proof (sketch).

- Let $s(v_i, v_j) = v_i + v_j \in \{e_i\}_{i \in [n]}$ for $(v_i, v_j) \in E(Q_n)$.
- Greedily build a rainbow path with an arbitrary start $v_0 v_1$.
- At v_k , there are at least $n - k$ incident colors not on the path. Choose v_{k+1} such that $s(v_k, v_{k+1})$ is distinct from the previous $\leq (n - k)$ -many choices.

Existing Bounds

Theorem (Rombach, unpublished)

$$ex^*(Q_n, P_{n+1}) < n2^{n-1} = |E(Q_n)|.$$

Proof (sketch).

- Let $s(v_i, v_j) = v_i + v_j \in \{e_i\}_{i \in [n]}$ for $(v_i, v_j) \in E(Q_n)$.
- Greedily build a rainbow path with an arbitrary start $v_0 v_1$.
- At v_k , there are at least $n - k$ incident colors not on the path. Choose v_{k+1} such that $s(v_k, v_{k+1})$ is distinct from the previous $\leq (n - k)$ -many choices.
- Greedy choice does not induce a cycle.



Existing Bounds

Theorem (Quail, unpublished)


$$ex^*(Q_n, P_{n+1}) \geq (n-1)2^{n-1}.$$

Existing Bounds

Theorem (Quail, unpublished)

$$ex^*(Q_n, P_{n+1}) \geq (n-1)2^{n-1}.$$

Proof (sketch).




Existing Bounds

Theorem (Quail, unpublished)

$$ex^*(Q_n, P_{n+1}) \geq (n-1)2^{n-1}.$$

Proof (sketch).



Contribution

Theorem

$$ex^*(Q_n, P_{n+1}) < n2^{n-1} - 1.$$

Proof.

Contribution

Theorem

$$ex^*(Q_n, P_{n+1}) < n2^{n-1} - 1.$$

Proof.

- Suppose uv is deleted from $E(Q_n)$.

Contribution

Theorem

$$\text{ex}^*(Q_n, P_{n+1}) < n2^{n-1} - 1.$$

Proof.

- Suppose uv is deleted from $E(Q_n)$.
- Restore the edge and assign it a color not used in the coloring. This yields a properly edge-colored Q_n .

Contribution

Theorem

$$ex^*(Q_n, P_{n+1}) < n2^{n-1} - 1.$$

Proof.

- Suppose uv is deleted from $E(Q_n)$.
- Restore the edge and assign it a color not used in the coloring. This yields a properly edge-colored Q_n .
- Find a rainbow P_{n+1} using Rombach's Algorithm starting from uv' for $v' \neq v$.

Contribution

Theorem

$$\text{ex}^*(Q_n, P_{n+1}) < n2^{n-1} - 1.$$

Proof.

- Suppose uv is deleted from $E(Q_n)$.
- Restore the edge and assign it a color not used in the coloring. This yields a properly edge-colored Q_n .
- Find a rainbow P_{n+1} using Rombach's Algorithm starting from uv' for $v' \neq v$.
- The path will not include the edge uv , as the algorithm doesn't create a cycle by construction.



Open Problems

Conjecture

$$ex^*(Q_n, P_{n+1}) = (n-1)2^{n-1}.$$

- Verified for $n = 3, 4$. Confirmed independently by Crawford, King, and Spiro (2025).

Open Problems

Conjecture

$$ex^*(Q_n, P_{n+1}) = (n-1)2^{n-1}.$$

- Verified for $n = 3, 4$. Confirmed independently by Crawford, King, and Spiro (2025).

Thank You