

Manuscript type: Article

Title: **Climate warming, resource availability, and the metabolic meltdown of ectotherms**

Authors:

Running Head: Metabolic meltdown

Online enhancements: Appendices

* Corresponding author:

ORCIDs:


1 ABSTRACT: Climate warming may lower environmental resource levels, growth, and fitness of
2 many ectotherms. In a classic experiment, Brett and colleagues (1969, 1971) documented that
3 growth rates of salmon depended strikingly on both temperature and food levels. Here we
4 develop a simple bioenergetic model that explores how fixed temperatures and food jointly alter
5 the thermal sensitivity of net energy gain. The model incorporates differing thermal sensitivities
6 of energy intake and metabolism. In qualitative agreement with Brett's results, it predicts that
7 decreased food intake reduces growth rates, lowers optimal temperatures for growth, and lowers
8 the highest temperatures sustaining growth (upper thermal limit). Consequently, ectotherms
9 facing reduced food intake in warm environments should restrict activity to times when low body
10 temperatures are biophysically feasible, but – in a warming world – that will force ectotherms to
11 shorten activity times and thus further reduce food intake. This 'metabolic meltdown' is a
12 consequence of declining energy intake coupled with accelerating metabolic costs at high
13 temperatures, and with warming-imposed restrictions on activity. Next, we extend the model to
14 explore how increasing mean environmental temperatures alter the thermal sensitivity of growth:
15 when food intake is reduced, optimal temperatures and upper thermal limits for growth are
16 lowered. We discuss our model's key assumptions and caveats, and its relationship to a recent
17 model for phytoplankton. Both models illustrate that the deleterious impacts of climate warming
18 on ectotherms will be amplified if food intake is also reduced, either because warming reduces
19 standing food resources or because it restricts foraging time.

20
21 *Keywords:* climate warming, energetics, resource availability, growth, metabolism, temperature
22

Introduction

Net energy gain influences an organism's growth and reproduction, and ultimately its fitness. An ectotherm's net energy gain depends primarily on its body temperature (T_b) and its intake of available food resources. Climate warming could potentially affect net energy gain by altering T_b distributions, by restricting foraging opportunities, or by reducing the availability of food in the environment (Dunham 1993; Kearney et al. 2009; Sinervo et al. 2010; Sears et al. 2016; Lister and Garcia 2018).

Here we develop a basic bioenergetic model that examines how the combined effects of climate warming, T_b , and food intake alter the energetics and thermoregulatory strategies of ectotherms. Our project was inspired by J. R. Brett and colleagues (1969; 1971), who discovered that the thermal sensitivity of growth rates of young salmon depended strikingly on temperature and food ration (fig. 1, redrawn from data in Brett et al., 1969). When Brett provided food in excess, salmon grew fastest at an intermediate temperature ($\sim 13^\circ\text{C}$) but began to lose mass at $\sim 24^\circ\text{C}$, which thus represented the upper thermal limit for growth. But when Brett restricted food: (1) growth rate declined at all temperatures, roughly in proportion to level of food restriction, (2) the upper thermal limit for growth dropped, and (3) the optimum temperature (T_{opt}) for growth also dropped. In effect, growth rate became less heat tolerant under restricted food. Similar patterns were later documented for other ectotherms (e.g., Elliott 1976; Gerard 1997; Peck et al. 2003; Thomas et al. 2017) and suggests the negative effects of climate warming will be amplified if warming results in reduced food intake (Thomas et al. 2017).



The thermal sensitivity of growth physiology (fig. 2A, C) provides a conceptual basis for the patterns in fig. 1. Food consumption (gross energy intake) under unlimited food typically increases with temperature up to an optimum temperature and then drops precipitously (fig. 2a, Waldschmidt et al. 1986; Koskela et al. 1997). In contrast, total metabolic expenses (maintenance, foraging, and digestion) typically increase exponentially with temperature, but ultimately drop at stressfully high T_b (Lighton and Turner 2004; Marshall and McQuaid 2011). Net energy gain (consumption gain minus metabolic expenses) thus is maximal at an intermediate but relatively warm temperature (see fig. 2B, D). Using this logic, Hainsworth and Wolf (fig. 2 in Hainsworth and Wolf 1978) and Huey (fig. 5 in Huey 1982) developed graphic models suggesting that lowered food rations will not merely reduce net energy gain (thus growth), but will also reduce T_{opt} for growth, consistent with Brett's (1969; 1971) findings.




Figure 2: When T_b and food intake are held constant, thermal dependence of net energy gain is sensitive to the breadth of the thermal performance curve for energy intake (generalists, specialists) as well as to the steepness (Q_{10} , or the factorial increase in metabolic rate over $+10^{\circ}\text{C}$ shift) of the metabolic-temperature curve. A) Gross energy gain (declining from black to gray) under three different food levels and metabolic losses (red) at three Q_{10} values (2, 2.5, 3) for a thermal generalist and C) for a thermal specialist. Net energy gain for a generalist (B) and a specialist (D) under three food levels and with Q_{10} s of 2 (solid line) or 3 (dotted line). T_{opt} for each (red dots) and the upper temperature threshold for growth both decline with food ration, especially for generalists with steep Q_{10} .

We begin by building a simple energy balance model that formalizes the graphical models of Hainsworth and Wolf (1978) and of Huey (1982). Our basic model mimics Brett's experiment

design – and thus assumes fixed T_b and food ration – but also adds comparisons involving thermal generalists versus specialists as well as for metabolic-rate curves with different steepness (Q_{10} , fig. 2). However, because ectotherms in nature have variable not fixed T_b , we next recast the model to allow for T_b distributions that can have different means and variances. We can thereby determine the T_b distribution that maximizes net energy gain under different food intake levels: in effect, this approach explores how the combined effects of climate change and food level alter the mean net energy gain.

Net energy gain will of course decline if climate warming forces ectotherms to be active at T_b above optimal physiological levels (fig. 2B, D). But if food intake is simultaneously reduced, then net energy gain and growth are now maximized at low T_b . In other words, a warming environment may push ectotherms towards accepting activity at high T_b , but energetics would push ectotherms to try to be active at low T_b . Thus, the selective forces are in opposite directions. In the DISCUSSION we will argue that declining food intake is in fact a likely consequence of climate warming, at least in environments that are already warm or hot. Thus, warming plus the associated food restriction could potentially amplify negative effects of climate warming, effectively lead to a “metabolic meltdown,” and thereby accentuate any negative energetic consequences of warming (see also Thomas et al. 2017).

Growth Rates in Constant Thermal Environments: Basic Model

We use a basic bioenergetic model (Kitchell et al. 1977; Hainsworth and Wolf 1978; Huey 1982), where the growth rate G of an organism is determined by its energy flux balance:

$$G = (A - fC) - R \quad (1)$$

In this model, C = rate of energy intake of ingested food (ingestion rate); R = rate of energy expenditure associated with metabolism and digestion (metabolic rate); A = the conversion efficiency of ingested food; and f = food level (see below). In general, C , R , and A will all depend on body temperature, T_b . For simplicity, we will assume that conversion efficiency A is constant (Brett 1971; Kingsolver and Woods 1997; Angilletta 2009) (see DISCUSSION).

The temperature dependence of ingestion rate ($C(T_b)$) and of metabolic rate ($R(T_b)$) have been examined in many ectotherms (Irlich et al. 2009; Dell et al. 2011). A thermal performance curve (“TPC,” Huey and Kingsolver 1989) for ingestion rate and other aspects of organism performance has a characteristic, asymmetric shape (figs. 1, 2), in which performance is maximal at some intermediate to warm ‘optimal’ temperature (T_{opt}) and declines more rapidly at higher than at lower temperatures (Gilchrist 1995; Dell et al. 2011). Here we model the thermal performance curve for ingestion rate (and gross energy intake rate = $A * C$) as a Gaussian multiplied by a Gompertz function (eq. 2), with parameters T_{opt} , ρ , and σ , where σ determines the steepness of the left portion of the TPC and ρ determines the steepness of the right portion (for details, see Frazier et al. 2006)}:

$$C = C_{max} \exp(-\exp((\rho \exp(-(T_b - T_{opt})/\sigma) - 6) - \sigma (T_b - T_{opt})^2)). \quad (2)$$

In contrast, metabolic rate typically increases exponentially over a wide range of temperatures (Savage 2004) until very high T_b (Lighton and Turner 2004; Marshall and McQuaid 2011). Consequently, we used an exponential equation to quantify the thermal dependence of metabolic rate for three typical Q_{10} values (2.0, 2.5, 3.0) and assume that R is independent of food intake (but see DISCUSSION). The net rate of growth (or net energy gain) then is the difference between gross energy gain and metabolic rate (eqn. 1 and 2, fig. 2C, E). All analyses were performed using the R statistical environment v. 3.5.3 (R Core Team 2019).

How might food limitation alter the thermal sensitivity of ingestion and net energy gain? We will explore another scenario below, but the simplest approach is to allow food level to be altered proportionately at all temperatures: i.e., that ingestion rate = $f * C$, where f is the proportional food level relative to unlimited food, when $f = 1$. In this case reductions in food will basically flatten the intake curve (e.g., fig. 2A vs 2B). Importantly, reduced food intake not only reduces maximum growth, but also reduces both T_{opt} and the maximum T_b for positive growth, especially for steep Q_{10} s (fig. 2B, D). This simple model, which formalizes prior graphical models (Hainsworth and Wolf 1978; Huey 1982), generates results that qualitatively match those of Brett's (1969; 1971) experiments (fig. 1).

The quantitative effects of food limitation on net growth are also influenced by the shape of thermal performance curve for ingestion ($C(T_b)$) and of the thermal sensitivity of metabolic rate ($R(T_b)$). To illustrate this, we modeled two hypothetical cases: a thermal generalist with a broad performance breadth ($\rho = 0.9$, $\sigma = 0.008$) (fig. 2 A,B, and a thermal specialist with a narrow performance breadth ("thermal specialist", $\rho = 0.9$, $\sigma = 0.02$) (fig. 2C, D). We set T_{opt} to 30°C for both cases, and incorporate a generalist-specialist tradeoff where the area under the TPC for energy ingestion is the same for the two cases (Levins 1968; Huey and Slatkin 1976; Gilchrist 1995), resulting in a lower maximum energy gain for the generalist than the specialist (compare fig. 2B, C). We also consider two levels (3.0, 2.0) of Q_{10} (fig. 2B, C).

The basic patterns (fig. 1) in Brett (1969; 1971) hold qualitatively for each of these variations (fig. 2). Specifically, reduced food levels reduce T_{opt} as well as the maximum temperature allowing positive net growth of both generalists and specialists. However, reduced food lowers both traits more for generalists than for specialists (fig. 2), and a steep Q_{10} also accentuates the drops in these traits.

Mean Growth Rates in Variable Thermal Environments

The above model quantifies ingestion, metabolic and net growth rate for a given (fixed) temperature T_b and a fixed food level f , thus matching Brett's (1971) experimental conditions. However, natural thermal environments vary in both time and space (Waldschmidt and Tracy 1983; Grant and Dunham 1990; Potter et al. 2013; Sears et al. 2016), and so ectotherms generally experience a range of environmental and body temperatures. Our model can predict mean growth rate during some time interval or life stage, by integrating over the distribution of body temperatures experienced during the interval or stage ($p(T_b)$, Huey and Slatkin 1976; Deutsch et al. 2008; Vasseur et al. 2014). Body temperatures distributions of active ectotherms are typically left-skewed and unimodal, and can be broad or narrow (Huey and Pianka 2017). We arbitrarily selected activity T_b of two desert lizards (*Pedioplanus lineoocellata*, from Kalahari desert; *Phrynosoma playrhinos*, from northern USA deserts) as exemplars of T_b distributions (Huey and Pianka 2017) of ectotherms with a relatively narrow versus broad distributions, respectively. We fit a skewed normal distribution (R package = fGARCH, Wuertz et al. 2017) to field T_b data for each species. Skewness was similar for both species (0.53, 0.41, respectively), but their standard deviations differed (2.1°C, 3.7°C). In our simulations, we set skewness = 0.5 for both, and set standard deviations of 2°C and 4°C, respectively. For simplicity, in our simulations the skew and standard deviation do not change with mean environmental temperature θ . We then calculate mean growth rate, \bar{G} for a temperature distribution, $p(T_b)$:

$$\bar{G} = \int G(T_b) p(T_b) dT_b \quad (3)$$

Equation (3) shows that mean growth rate will depend on the distribution of body temperatures, the thermal sensitivities of ingestion and metabolic rates, and the food level (see eq. 1). For simplicity, suppose environmental temperatures directly determine T_b of an ectotherm (a

“thermoconformer,” see DISCUSSION). As mean environmental temperature θ (and thus T_b) increases during climate warming, mean growth rate will also change. In the simplest case (food level remains constant and high during climate warming), mean growth rate declines rapidly at high mean environmental temperatures, because ingestion declines while metabolic rate accelerates (see fig. 4, below).

Climate change may also have direct or indirect effects on the food resources available to ectotherms. As described in the DISCUSSION, global warming or other aspects of anthropogenic environmental change may decrease available food levels (Deutsch et al. 2008; Hallmann et al. 2017; Bestion et al. 2018; Lister and Garcia 2018) as well as reduce opportunities to forage (Dunham 1993; Sinervo et al. 2010; Sears et al. 2011). We consider two cases: 1) reduced food levels that are independent of mean environmental temperatures (fig. 3, solid black lines); or 2) food levels that decline as mean environmental temperature increases (fig. 3, gray line), which might occur if resources decline with climate warming, or if foraging becomes restricted at high environmental temperatures. We characterize this decline in food level f using a sigmoid function (Borchers 2018):

$$f = f_{\max} - \frac{(f_{\max} - f_{\min})}{1 + \exp(-a(\theta - \theta_{\text{mid}}))} \quad (4)$$

where f_{\max} and f_{\min} are the maximum and minimum food levels, θ_{mid} is the midpoint environmental temperature at which food level is midway between f_{\max} and f_{\min} , and a determines the slope of the decline in food level. Food level f varies between 0 and 1, where $f = 1$ indicates the case of unlimited food (see above). Note that at $\theta = \theta_{\text{mid}}$, the slope of eq (4) is $df/d\theta = -a(f_{\max} - f_{\min})/4$ (see fig. 3).





Figure 3: Food levels at different environmental temperatures. Case 1: food level is independent of temperature, and solid lines indicate constant food levels of 1.0 (black), 0.75 (gray), and 0.5 (light gray). Case 2: food level declines with increasing mean environmental temperatures (dashed black line, $a = 0.3$). Line thickness is scaled to food level.

Effects on growth rates of various levels of food ration and of mean environmental temperature show that Brett's basic results (fig. 1) hold for shifting temperature distributions (fig. 4) well as at fixed temperatures (fig. 2B, D). When food declines proportionately at all temperatures (solid black or gray lines in fig. 4), growth declines, T_{opt} shifts to lower temperatures, and the highest temperature permitting growth all decline, and do so more strongly for thermal generalists (fig. 4A, B vs. C, D). When food declines (fig. 3) mainly at high environmental temperature (θ), T_{opt} drops dramatically (dashed black lines, fig. 4). In this case, the growth curve converges on that of excess food at low θ , but on that of restricted food at high θ . Increasing the magnitude of thermal environmental variation (SD in the temperature distribution $p(T_b)$) generally decreases maximum mean growth rates, but causes only a modest increase ($< 0.5^\circ\text{C}$) in the magnitude of the response to (uniformly) decreased food intake

(compare left and right columns in fig. 4). Changing the midpoint temperature (θ_{id}) or slope (a) of the relationship between food level and environmental temperature (eqn 4, fig. 3) does not qualitatively alter these effects. Thus, declining resources will amplify the negative consequences of climate warming for mean growth, and will lower the environmental temperatures at which climate warming will start to have negative impacts on growth rate.



DISCUSSION

Climate warming can have disruptive effects on the performance and fitness of organisms, and these effects will be mediated in part on physiological processes such as food consumption and

metabolism as well as by behavioral thermoregulation (Kearney et al. 2009). For ectotherms, high environmental temperatures can lead to reduced rates of growth, at least if T_b are elevated above optimal levels. But if food resources are declining simultaneously, either directly as a result of warming or of other concurrent environmental changes, or indirectly as a result of behavioral restrictions on activity time during warm periods (Stevenson 1985; Dunham 1993; Sinervo et al. 2010), then maximal growth rates will decline, the highest temperature permitting growth will decline, and the optimal temperature for growth will also decline. Thus, although climate warming can itself be energetically deleterious, but warming plus reduced food will be multiplicatively deleterious: Ectotherms will be becoming energetically less heat tolerant while living in a warming environment, a phenomenon we call “metabolic meltdown.” Below we discuss the biological justification underlying key assumptions of our simple model, and also consider some implications of metabolic meltdown.

Will T_b Increase with Warming?

Our model assumes that T_b will increase one to one with changes in environmental temperatures (θ). Thus, our model is most appropriate for thermoconformers, which typically have limited mobility or live in thermally homogeneous environments, and thus have limited opportunity for behavioral adjustments (Huey 1974; Willmer 1982; Helmuth et al. 2002; Chown and Nicholson 2004). However, our model should still be relevant for thermoregulators, even though their T_b do not change as fast as T_e (Huey 1974; Kearney et al. 2009). Lizards that are careful thermoregulators typically have elevated T_b at in warm seasons (Huey et al. 1977; Christian et al. 1983; Gunderson and Leal 2012); and some even stay active at very high T_b during hot times

(McGinnis and Dickson 1967; Grant and Dunham 1988; Gunderson and Leal 2012). Thus, $p(T_b)$ – especially that of thermoconformers – is likely to increase as climates warm.

Importantly, the primary behavioral adjustments used by thermoregulators often involve restricting activity and foraging time (i.e., by withdrawing to a thermal retreat during hot periods) (Porter et al. 1973; Stevenson 1985; Kearney et al. 2009) or restricting space use (i.e., to cool microenvironments, Stevenson 1985; Gunderson and Leal 2015; Sears et al. 2016): both adjustments should effectively reduce food consumption and reproduction (Dunham 1993; Sinervo et al. 2010), even if environmental food levels are unchanged. In effect, the optimal behavioral solution (activity only at lower T_b) will further reduce net energy gain by restricting foraging. Therefore, thermoregulators may be able to reduce metabolic meltdown in warming environments (Kearney et al. 2009; Huey et al. 2012), but they cannot escape it.

Will Food Availability Decline with Warming?

The effects of climate change on primary productivity and resource levels will vary across geographic regions and ecosystems, but recent analyses suggest that warming temperatures and climate change will negatively impact resources in many temperate and tropical systems. For example, decreasing mean and increased variability in precipitation can reduce primary productivity in grassland ecosystems (Pan et al. 2017). Phytoplankton productivity may decline in response to climate change in many marine ecosystems, particularly in tropical shelf and upwelling areas (Blanchard et al. 2012; Thomas et al. 2017).

Warming temperatures potentially reduce standing food levels in several ways. Warming can reduce primary productivity, thereby reducing the food base for herbivores and higher trophic

levels (Hallmann et al. 2017; Lister and Garcia 2018). Warming plus drought can cause plant dieback (Ciais et al. 2005), which will not only reduce the food base (for herbivores), but also reduce habitat shading, increasing operative temperatures, and thus either increase T_b or reduce foraging time and opportunity (or both).

Warming temperatures can also alter rates of population growth (fitness) of organisms serving as competitors or as prey for higher trophic levels (Buckley 2013; Bestion et al. 2018). In cold environments, of course, warming may accelerate prey population growth; but in warm environments, warming may depress rates of population growth (Vasseur and McCann 2005; Deutsch et al. 2008; Kingsolver et al. 2011). In particular, climate change is predicted to reduce reproductive rates and survival of insects in tropical and many temperate regions, (Vasseur and McCann 2005; Deutsch et al. 2008; Kingsolver et al. 2011). In addition, increasing mean temperatures can reduce population carrying capacity (Savage et al. 2004; Vasseur and McCann 2005; Woodworth-Jefcoats et al. 2017; Bernhardt et al. 2018). For example, a recent experimental study showed that increasing temperature reduced mean cell size, carrying capacity, and total biomass of a mobile phytoplankton (Bernhardt et al. 2018). These results suggest that climate warming – at least in already warm environments – may reduce the density of resources available for many ectothermic predators and herbivores.

Field studies confirm that food levels have declined in recent decades in some terrestrial ecosystems. For example, abundances of arthropods have declined markedly over several decades in Puerto Rico and in México, with apparent repercussions on higher trophic levels: climate warming is the likely cause (Lister and Garcia 2018). Declines in insect pollinators can potentially reduce plant reproduction and also food supply to predators of pollinators (Biesmeijer et al. 2006). Of course, food can decline for reasons other than climate warming (Zhao and

Running 2010; Hallmann et al. 2017). For example, the biomass of flying insects in protected sites in Germany has declined 75% in 27 years; but this decline shows no obvious imprint of climate change (Hallmann et al. 2017). In any case, what matters is whether food is declining in parallel with climate warming, not whether climate warming is the driving factor. Interestingly, diverse ectotherms lower their preferred body temperatures under reduced (or fasting) food regimes (reviewed in Angilletta 2009, pp. 102-105), but lowered preferred temperatures will constrain activity time in warm environments, further reducing energy gain (Dunham 1993; Sinervo et al. 2010). Alternatively, ectotherms could reduce metabolic rates in response to warming (Marshall and McQuaid 2011), which will reduce metabolic meltdown but likely slow growth and reproduction.

Our discussion has focused on the consequences of warming in already warm environments. However, many ectotherms routinely experience low, suboptimal temperatures – especially in seasonal environments (fig. 6b in Huey et al. 2012). For them, warming may enhance growth rates (Deutsch et al. 2008; Kingsolver et al. 2013), but that enhancement will be blunted under reduced food ration (fig. 4). Nevertheless, metabolic meltdown is mainly a high temperature effect.

Can Other Mechanisms Lead to ‘Meltdown’?

The predictions from our model emerge from three basic elements: maximal ingestion rate varies with temperature, where ingestion rate is maximal at some intermediate (optimal) temperature; ingestion rate at any given temperature increases with food level; and metabolic rate accelerates with increasing temperature. Our simple model assumes that ingestion rate is

linearly related to food level, but non-linear relationships between ingestion (intake) and food level (nutrient concentration) are widespread. In particular, individual or population growth rate is a saturating (decelerating) function of the availability of a limiting resource in many bacteria, plankton and animals (Monod 1949; Holling 1959; Eppley et al. 1969; Thomas et al. 2017).

Thomas et al. (2017) recently modeled interactions between temperature and nutrient concentration in phytoplankton. In their model, population growth rate is a function of both temperature (where population growth rate is maximal at some intermediate (optimal) temperature) and of nutrient concentration (where population growth rate is a decelerating function of concentration that saturates at some maximal growth rate). This model predicts that declining nutrient concentrations will reduce maximal (population) growth rate, the optimal temperature and the upper thermal limit (temperature at which population growth rate is zero). Thus predictions from our energetic model of metabolic meltdown qualitatively match those a population one (Thomas et al. 2017). Laboratory experiments with a marine diatom confirm that the optimal temperature for population growth declines with declining nutrient (phosphate and nitrate) concentrations; the decline in optimal temperatures is particularly pronounced at very low nutrient concentrations (Thomas et al. 2017). Note that their model does not explicitly consider metabolic or respiration rates (which are incorporated via their effects on population growth rates), nor does it consider behavioral responses to warming.

Mathematically, the similar qualitative predictions from these two models emerge from the decelerating function relating growth rate to resource concentration (Thomas et al. 2017) versus the accelerating function relating respiration rate to temperature (the current model; fig. 2). More generally, suppose that net performance or fitness results from the difference between two functions of temperature: a decelerating ‘gain’ function and an accelerating ‘loss’ function (eqns

1-2). Reductions in resources (or other environment factors) that reduce gains or magnify losses will have similar effects on shifting optimal and maximal temperatures as described here. In particular, the effects of resource levels on birth and death functions could also alter the thermal dependence of fitness in ectotherms (Amarasekare and Savage 2012). These considerations suggest there are multiple mechanisms by which temperature and resources may interact to exacerbate the negative consequences of climate changes for ectothermic organisms (Thomas et al. 2017).

Caveats

Our model is intentionally simple and summarizes key factors involving food, temperature, and energetics that are universal for ectotherm growth. In the future, more complex models (e.g., incorporating nutrients, energetics, biophysical models of heat exchange: Kooijman 2010; Clissold and Simpson 2015; Sears et al. 2016; Thomas et al. 2017; Malishev et al. 2018) can explore fine-scaled aspects of metabolic meltdown. Here we highlight a few known complications.

First, our model belongs to a family of models (e.g., Huey and Slatkin 1976; Vasseur et al. 2014) that integrate performance or fitness over a distribution of T_b (eq. 2). However, physiologists have long appreciated that the effect of T_b on performance or fitness can change with acclimation (Bernard 1865; Fry and Hart 1948), ontogeny (Brett 1970), and time-at-temperature (Wilhoft 1958; Kingsolver and Woods 2016). Such effects – when major – will alter the conclusions, especially at high temperatures (Brakefield and Kesbeke 1996; Schulte et al. 2011; Niehaus et al. 2012; Colinet et al. 2014; Kingsolver and Woods 2016; Sinclair et al. 2016; Kremer et al. 2018; Bernhardt et al. 2018).

Second, we assumed that metabolic rate is independent of food intake. However, many ectotherms (and even humans) reduce metabolic rate under reduced food (Secor and Diamond 2000; Auer et al. 2015). Reduced metabolic rate under low food intake should reduce – but not eliminate – the declines in T_{opt} and in the highest temperature for growth (compare results for the two Q_{10} s in Figure 2B, D).

Third, our model assumes that the distribution of T_b is unimodal and left skewed: this is valid for ectotherms such as diurnal desert lizards when active (Huey and Pianka 2017). However, growth is sensitive to T_b over 24 hours, not just during activity. When compiled over 24 h (or longer), T_b distributions typically have relatively broad distributions, at least at low T_b ; and sometimes they are even bimodal (Stevenson et al. 1985). However, our simulations with two breadth levels (standard deviations) suggest that the breadth of the T_b distribution will have limited effect on the magnitude of the drop in T_{opt} caused by food restriction. For example, the drops are 1.8°C and 2.2°, respectively, for the two generalist cases (fig. 4a, b) and 0.7°C and 0.9°C the specialist cases (fig. 4c, d).

To explore in brief the impact of bimodal T_b distribution, we ran the model with two modes set at + 4°C and - 4°C of the mean T_b (standard deviations of 1 and of 2, respectively). The resultant growth curves (not shown) are very similar qualitatively to those the unimodal cases in fig. 4, but have reduced growth (reflecting time at low T_b) and slightly left shifted T_{opt} and upper threshold temperatures. Also, differences in maximum growth rates between generalists and specialists are reduced relative to the unimodal cases.

Fourth, ectotherms live in communities; and the associated exploitative competition and predation can result in reduced food intake (Huey and Slatkin 1976; Gilman et al. 2010; Thakur et al. 2017). Moreover, micro-environments are often heterogeneous such that individuals may

differ in exposure to food intake and to heat stress, even on a local scale (Denny et al. 2011; Sears et al. 2011; Potter et al. 2013). Modeling such complexities will be challenging, given that the responses of interacting ectotherms to climate change may be species specific and context dependent.

Fifth, our model considers the effects of food (resource) quantity on ingestion and net growth rates, but assumes that resource quality (as reflected in assimilation efficiency—see eq. 1) is constant and independent of environmental conditions. However, climate change can reduce resource quality. For example, increasing atmospheric CO₂ reduces protein content and protein:carbohydrate ratios, and increases concentrations of secondary defense chemicals, in leaves of most terrestrial plants (Mooney 1972; Chapin 1980). Numerous studies demonstrate that these effects reduce rates of growth, development, and fitness in insect and other herbivores as well as alter plant defenses against insect herbivores (see Lincoln et al. 1993). In addition, increasing temperatures during development reduces final (adult) body size in most ectotherms, and reductions in mean body size during climate warming during recent decades has been widely (but not universally) reported (Atkinson 1994; Kingsolver and Huey 2008; Daufresne et al. 2009; Gardner et al. 2011) in ectotherms. If smaller prey size reduces the efficiency of search, handling or assimilation by predators, this will decrease net energy gain. Reductions in food quality or conversion efficiency with climate warming should have similar consequences for the thermal-dependence of growth as modeled here for declining food levels (see eq. 1).

Sixth, we assumed a Gompertz X Gaussian function to describe the thermal growth curve (eq. 2), and a simple exponential for metabolic rate versus temperature. Other functional relationships are reasonable (Izem and Kingsolver 2005; Angilletta 2006; Sheth and Angert

2014): substitution of biologically reasonable functional forms are unlikely to change the qualitative patterns described here.

In summary, our conceptual extensions of Brett's pioneering experiments (Brett et al. 1969; Brett 1971) and a recent model of Thomas and colleagues (Thomas et al. 2017) suggest that the combination of climate warming plus food decline will exaggerate the negative consequences of climate change alone or of food decline alone – at least in areas that are already warm. If only the environment warms, ectotherms may need to accept activity at supra-optimal T_b to maintain sufficient activity time, potentially at a cost of stress. If only food levels decline, ectotherms should accept active at sub-optimal T_b , which are energetically favorable but reduce maximal growth. But if both occur together, ectotherms face antagonistic thermoregulatory pressures and thus will be in an ecological 'double bind' or "Catch-22." Accepting high T_b will maintain activity time but greatly reduce net energetic gain. Accepting low T_b will be energetically advantageous, but will greatly reduce activity time, further lowering food intake, and further lowering growth.

Acknowledgments

Our paper honors J. R. Brett, who was a pioneer of fish physiological ecology, on the 50th anniversary year of his 1969 paper. We thank L. Buckley, M. Denny, D. Miles, B. Sinervo, D. Vasseur, and an anonymous reviewer for constructive discussions and suggestions. The Whiteley Center provided an ideal environment for collaborating on model simulations and drafting the manuscript.

Literature Cited

- Amarasekare, P., and V. Savage. 2012. A framework for elucidating the temperature dependence of fitness. *The American Naturalist* 179:178-191.
- Angilletta, M. J., Jr. 2006. Estimating and comparing thermal performance curves. *Journal of Thermal Biology* 31:541-545.
- . 2009. *Thermal adaptation: a theoretical and empirical synthesis*. Oxford, U.K., Oxford University Press.
- Atkinson, D. 1994. Temperature and organism size - a biological law for ectotherms? *Advances in Ecological Research* 25:1-58.
- Auer, S. K., K. Salin, A. M. Rudolf, G. J. Anderson, and N. B. Metcalfe. 2015. Flexibility in metabolic rate confers a growth advantage under changing food availability. *Journal of Animal Ecology* 85:1405-1411.
- Bernard, C. 1865, *Introduction à l'étude de la médecine expérimentale*. Paris, Bailliere.
- Bernhardt, J. F., J. M. Sunday, and M. I. O'Connor. 2018. Metabolic theory and the temperature-size rule explain the temperature dependence of population carrying capacity. *The American Naturalist* 192:687-697.
- Bernhardt, J. R., J. M. Sunday, P. L. Thompson, and M. I. O'Connor. 2018 Nonlinear averaging of thermal experience predicts population growth rates in a thermally variable environment. *Proceedings of the Royal Society B* 285:20181076.
- Bestion, E., B. García-Carreras, C.-E. Schaum, S. Pawar, and G. Yvon-Durocher. 2018. Metabolic traits predict the effects of warming on phytoplankton competition. *Ecology Letters* 21:655-664.

Biesmeijer, J. C., S. P. M. Roberts, M. Reemer, R. Ohlemüller, M. Edwards, T. Peeters, A. P.

Schaffers et al. 2006. Parallel declines in pollinators and insect-pollinated plants in Britain and the Netherlands. *Science* 313:351-354.

Blanchard, J. L., S. Jennings, R. Holmes, J. Harle, G. Merino, J. I. Allen, J. Holt et al. 2012.

Potential consequences of climate change for primary production and fish production in large marine ecosystems. *Philosophical Transactions of the Royal Society B* 367:2979-2989.

Borchers, H. W. 2018. *pracma: Practical Numerical Math Functions*, version R package version 2.2.2.

Brakefield, P. M., and P. Kesbeke. 1996. Genotype x environment interactions for insect

growth in constant and fluctuating temperature regimes. *Proceedings of the Royal Society B* 264:717-723.

Brett, J. R. 1970. Temperature, fishes., Pages 515-560 in O. Kinne, ed. *Marine Ecology* vol. 1. New York, N. Y., John Wiley & Sons.

—. 1971. Energetic responses of salmon to temperatures. A study of some thermal relations in the physiology and freshwater ecology of sockeye salmon (*Oncorhynchus nerka*). *American Zoologist* 11:99-113.

Brett, J. R., J. E. Shelbourn, and C. T. Shoop. 1969. Growth rate and body composition of fingerling sockeye salmon, *Oncorhynchus nerka*, in relation to temperature and ration size. *Journal of the Fisheries Research Board of Canada* 26:2363-2394.

Buckley, L. B. 2013. Get real: putting models of climate change and species interactions in practice. *Annals of the New York Academy of Sciences* 1297:126-138.

- Chapin, F. S., III. 1980. The mineral nutrition of wild plants. *Annual Review of Ecology and Systematics* 11:233-260.
- Chown, S. L., and S. Nicholson. 2004. *Insect physiological ecology: mechanisms and patterns*. Oxford, U.K., Oxford University Press.
- Christian, K. A., C. R. Tracy, and W. P. Porter. 1983. Seasonal shifts in body temperature and use of microhabitats by Galapagos land iguanas (*Conolophus pallidus*). *Ecology* 64:463-468.
- Ciais, P., M. Reichstein, N. Viovy, A. Granier, J. Ogee, V. Allard, M. Aubinet et al. 2005. Europe-wide reduction in primary productivity caused by the heat and drought in 2003. *Nature* 437:529-533.
- Clissold, F. J., and S. J. Simpson. 2015. Temperature, food quality and life history traits of herbivorous insects. *Current Opinion in Insect Science*. 11:63-70.
- Colinet, H., B. Sinclair, P. Vernon, and Renault. 2014. Insects in fluctuating thermal environments. *Annual Review of Entomology* 60:123-140.
- Daufresne, M., K. Lengfellner, and U. Sommer. 2009. Global warming benefits the small in aquatic ecosystems. *Proceedings of the National Academy of Sciences, USA* 106:12788-12793.
- Dell, A. I., S. Pawar, and V. M. Savage. 2011. Systematic variation in the temperature dependence of physiological and ecological traits. *Proceedings of the National Academy of Sciences, USA* 108:10591-10596.
- Denny, M. W., W. W. Dowd, L. Bilir, and K. J. Mach. 2011. Spreading the risk: Small-scale body temperature variation among intertidal organisms and its implications for species persistence. *Journal of Experimental Marine Biology and Ecology* 400:175-190.

- Deutsch, C. A., J. J. Tewksbury, R. B. Huey, K. S. Sheldon, C. K. Ghalambor, D. C. Haak, and P. R. Martin. 2008. Impacts of climate warming on terrestrial ectotherms across latitude. *Proceedings of the National Academy of Sciences, USA* 105:6668–6672.
- Dunham, A. E. 1993. Population responses to environmental change: physiologically structured models, operative environments, and population dynamics, Pages 95-119 in P. M. Kareiva, J. G. Kingsolver, and R. B. Huey, eds. *Biotic Interactions and Global Change*. Sunderland, MA, Sinauer Associates.
- Elliott, J. M. 1976. The energetics of feeding, metabolism and growth of brown trout (*Salmo trutta* L.) in relation to body weight, water temperature and ration size. *Journal of Animal Ecology* 45:923-948.
- Eppley, R. W., J. N. Rogers, and J. J. McCarthy. 1969. Half-saturation constants for uptake of nitrate and ammonium by marine phytoplankton. *Limnology and Oceanography* 14:912-920.
- Frazier, M. R., R. B. Huey, and D. Berrigan. 2006. Thermodynamics constrains the evolution of insect population growth rates: "warmer is better". *The American Naturalist* 168:512-520.
- Fry, F. E. J., and J. S. Hart. 1948. Cruising speed of goldfish in relation to water temperature. *Journal of the Fisheries Research Board of Canada* 7:169-175.
- Gardner, J. L., A. Peters, M. R. Kearney, L. Joseph, and R. Heinsohn. 2011. Declining body size: a third universal response to warming? *Trends in Ecology & Evolution* 26:285-291.
- Gerard, V. A. 1997. The role of nitrogen nutrition in high-temperature tolerance of the kelp *Laminaria saccharina* (Chromophyta). *Journal of Phycology* 33:800–810.

- Gilchrist, G. W. 1995. Specialists and generalists in changing environments. I. Fitness landscapes of thermal sensitivity. *The American Naturalist* 146:252-270.
- Gilman, S. E., M. C. Urban, J. Tewksbury, G. W. Gilchrist, and R. D. Holt. 2010. A framework for community interactions under climate change. *Trends in Ecology & Evolution* 25:325-331.
- Grant, B. W., and A. E. Dunham. 1988. Thermally imposed time constraints on the activity of the desert lizard *Sceloporus merriami*. *Ecology* 69:167-176.
- . 1990. Elevational covariation in environmental constraints and life histories of the desert lizard *Sceloporus merriami*. *Ecology* 71:1765-1776.
- Gunderson, A. R., and M. Leal. 2012. Geographic variation in vulnerability to climate warming in a tropical Caribbean lizard. *Functional Ecology* 26:783–793.
- . 2015. Patterns of thermal constraint on ectotherm activity. *The American Naturalist* 185:653-664.
- Hainsworth, F. R., and L. F. Wolf. 1978. The economics of temperature regulation and torpor in nonmammalian organisms *in* L. C. H. Wang, and J. W. Hudson, eds. *Strategies in cold: natural torpidity and thermogenesis*. New York, Academic Press.
- Hallmann, C. A., M. Sorg, E. Jongejans, H. Siepel, N. Hofland, H. Schwan, W. Stenmans et al. 2017. More than 75 percent decline over 27 years in total flying insect biomass in protected areas. *PLoS ONE* 12:e0185809.
- Helmuth, B., C. D. G. Harley, P. M. Halpin, M. O'Donnell, G. E. Hofmann, and C. A. Blanchete. 2002. Climate change and latitudinal patterns of intertidal thermal stress. *Science* 298:1015-1017.

- Holling, C. S. 1959. The components of predation as revealed by a study of small-mammal predation of the European Pine Sawfly. *The Canadian Entomologist* 91:293-320.
- Huey, R. B. 1974. Behavioral thermoregulation in lizards: importance of associated costs. *Science* 184:1001-1003.
- . 1982. Temperature, physiology, and the ecology of reptiles, Pages 25-91 *in* C. Gans, and F. H. Pough, eds. *Biology of the Reptilia*, Vol. 12, Physiology (C). London, Academic Press.
- Huey, R. B., M. R. Kearney, A. Krockenberger, J. A. M. Holthum, M. Jess, and S. E. Williams. 2012. Predicting organismal vulnerability to climate warming: roles of behaviour, physiology and adaptation. *Philosophical Transactions of the Royal Society B* 367:1665-1679.
- Huey, R. B., and J. G. Kingsolver. 1989. Evolution of thermal sensitivity of ectotherm performance. *Trends in Ecology & Evolution* 4:131-135.
- Huey, R. B., and E. R. Pianka. 2017. Body temperature distributions of active diurnal lizards in three deserts: skewed up or skewed down? *Functional Ecology* 32:334-344.
- Huey, R. B., E. R. Pianka, and J. Hoffmann. 1977. Seasonal variation in thermoregulatory behavior and body temperatures of diurnal Kalahari lizards. *Ecology* 58:1066-1075.
- Huey, R. B., and M. Slatkin. 1976. Cost and benefits of lizard thermoregulation. *Quarterly Review of Biology* 51:363-384.
- Irlich, U. M., J. S. Terblanche, T. M. Blackburn, and S. L. Chown. 2009. Insect rate-temperature relationships: environmental variation and the metabolic theory of ecology. *The American Naturalist* 174:819-835.

- Izem, R., and J. G. Kingsolver. 2005. Variation in continuous reaction norms: quantifying directions of biological interest. *The American Naturalist* 166:277-289.
- Kearney, M., R. Shine, and W. P. Porter. 2009. The potential for behavioral thermoregulation to buffer "cold-blooded" animals against climate warming. *Proceedings of the National Academy of Sciences, USA* 106:3835-3840.
- Kingsolver, J. G., S. E. Diamond, and L. B. Buckley. 2013. Heat stress and the fitness consequences of climate change for terrestrial ectotherms. *Functional Ecology* 27:1415-1423.
- Kingsolver, J. G., and R. B. Huey. 2008. Size, temperature, and fitness: Three rules. *Evolutionary Ecology Research* 10:251-268.
- Kingsolver, J. G., and H. A. Woods. 1997. Thermal sensitivity of feeding and digestion in *Manduca* caterpillars. *Physiological Zoology* 70:631-638.
- Kingsolver, J. G., and H. A. Woods. 2016. Beyond thermal performance curves: modeling time-dependent effects of thermal stress on ectotherm growth rates. *The American Naturalist* 187:283-294.
- Kingsolver, J. G., H. A. Woods, L. B. Buckley, K. A. Potter, H. J. MacLean, and J. K. Higgins. 2011. Complex life cycles and the responses of insects to climate change. *Integrative and Comparative Biology* 51:719-732.
- Kitchell, J. F., D. J. Stewart, and D. Weininger. 1977. Application of a bioenergetics model of yellow perch (*Perca flavescens*) and walleye (*Stizostedion vitreum vitreum*). *Journal of Fisheries Research Board of Canada* 34:1922– 1935.
- Kooijman, S. A. L. M. 2010, *Dynamic energy budget theory for metabolic organization*. Cambridge, U.K., Cambridge University Press.

- Koskela, J., J. Pirhonen, and M. Jobling. 1997. Feed intake, growth rate and body composition of juvenile Baltic salmon exposed to different constant temperatures. *Aquaculture International* 5:351-360.
- Kremer, C. T., S. B. Fey, A. A. Arellano, and D. A. Vasseur. 2018. Gradual plasticity alters population dynamics in variable environments: thermal acclimation in the green alga *I. Proceedings of the Royal Society B* 285:20171942.
- Levins, R. 1968, *Evolution in changing environments*. Princeton, N.J., Princeton University Press.
- Lighton, J. R. B., and R. J. Turner. 2004. Thermolimit respirometry: an objective assessment of critical thermal maxima in two sympatric desert harvester ants, *Pogonomyrmex rugosus* and *P. californicus*. *Journal of Experimental Biology* 207:1903 -1913.
- Lincoln, D. E., E. D. Fajer, and R. H. Johnson. 1993. Plant-insect herbivore interactions in elevated CO₂ environments. *Trends in Ecology & Evolution* 8:64-68.
- Lister, B. C., and A. Garcia. 2018. Climate-driven declines in arthropod abundance restructure a rainforest food web. *Proceedings of the National Academy of Sciences, USA* 115:E10397–E10406.
- Malishev, M., C. M. Bull, and M. R. Kearney. 2018. An individual-based model of ectotherm movement integrating metabolic and microclimatic constraints. *Methods in Ecology and Evolution* 9:472-489.
- Marshall, D. J., and C. D. McQuaid. 2011. Warming reduces metabolic rate in marine snails: adaptation to fluctuating high temperatures challenges the metabolic theory of ecology. *Proceedings of the Royal Society B* 278:281-288.

- McGinnis, S. M., and L. L. Dickson. 1967. Thermoregulation in the desert iguana *Dipsosaurus dorsalis*. *Science* 156:1757-1759.
- Monod, J. 1949. The growth of bacterial cultures. *Annual Review of Microbiology* 3:371-394.
- Mooney, H. A. 1972. The carbon balance of plants. *Annual Review of Ecology and Systematics* 3:315-346.
- Niehaus, A. C., M. J. Angilletta, M. W. Sears, C. E. Franklin, and R. S. Wilson. 2012. Predicting the physiological performance of ectotherms in fluctuating thermal environments. *Journal of Experimental Biology* 215:694-701.
- Pan, S., S. R. S. Dangal, B. Tao, J. Yang, and H. Tian. 2017. Recent patterns of terrestrial net primary production in Africa influenced by multiple environmental changes. *Ecosystem Health and Sustainability* 1:1-15.
- Peck, M. A., L. J. Buckley, E. M. Caldarone, and D. A. Bengtson. 2003. Effects of food consumption and temperature on growth rate and biochemical-based indicators of growth in early juvenile Atlantic cod *Gadus morhua* and haddock *Melanogrammus aeglefinus*. *Marine Ecology Progress Series* 251:233–243.
- Porter, W. P., J. W. Mitchell, W. A. Beckman, and C. B. DeWitt. 1973. Behavioral implications of mechanistic ecology. Thermal and behavioral modeling of desert ectotherms and their microenvironment. *Oecologia* 13:1-54.
- Potter, K. A., H. A. Woods, and S. Pincebourde. 2013. Microclimatic challenges in global change biology. *Global Change Biology* 19:2932-2939.
- R Core Team. 2019. R: A language and environment for statistical computing, Vienna, Austria.

- Savage, V. M. 2004. Improved approximations to scaling relationships for species, populations, and ecosystems across latitudinal and elevational gradients. *Journal of Theoretical Biology* 227:525-534.
- Savage, V. M., J. F. Gillooly, J. H. Brown, G. B. West, and E. L. Charnov. 2004. Effects of body size and temperature on population growth. *The American Naturalist* 163:429-441.
- Schulte, P. M., T. M. Healy, and N. A. Fangue. 2011. Thermal performance curves, phenotypic plasticity, and the time scales of temperature exposure. *Integrative and Comparative Biology* 51:1-12.
- Sears, M. W., M. J. Angilletta, M. S. Schuler, J. Borchert, K. F. Dilliplane, M. Stegman, T. Rusch et al. 2016. Configuration of the thermal landscape determines thermoregulatory performance of ectotherms. *Proceedings of the National Academy of Sciences U.S.A.* 113:10595–10600.
- Sears, M. W., E. Raskin, and M. J. Angilletta, Jr. 2011. The world is not flat: defining relevant thermal landscapes in the context of climate change. *Integrative and Comparative Biology* 5:666-675.
- Secor, S. M., and J. M. Diamond. 2000. Evolution of regulatory responses to feeding in snakes. *Physiological and Biochemical Zoology* 73:123-141.
- Sheth, S. N., and A. L. Angert. 2014. The evolution of environmental tolerance and range size: a comparison of geographically restricted and widespread *Mimulus*. *Evolution* 68:2917-2931.
- Sinclair, B. J., K. E. Marshall, M. A. Sewell, D. L. Levesque, C. S. Willett, C. D. G. Harley, D. J. Marshall et al. 2016. Can we predict ectotherm responses to climate change using thermal performance curves and body temperatures? *Ecology Letters* 19:1372-1375.

- Sinervo, B., F. Méndez-de-la-Cruz, D. B. Miles, B. Heulin, E. Bastiaans, M. Vallagrán-Santa Cruz, R. Lara-Resendiz et al. 2010. Erosion of lizard diversity by climate change and altered thermal niches. *Science* 328:894-899.
- Stevenson, R. D. 1985. The relative importance of behavioral and physiological adjustments controlling body temperature in terrestrial ectotherms. *The American Naturalist* 126:362-386.
- Stevenson, R. D., C. R. Peterson, and J. S. Tsuji. 1985. The thermal dependence of locomotion, tongue flicking, digestion and oxygen consumption in the wandering garter snake. *Physiological Zoology* 58:46-57.
- Thakur, M. P., T. Künne, J. N. Griffin, and N. Eisenhauer. 2017. Warming magnifies predation and reduces prey coexistence in a model litter arthropod system. *Proceedings of the Royal Society B* 284: 20162570.
- Thomas, M. K., Aranguren-Gassis, C. T. Kremer, M. R. Gould, K. Anderson, C. A. Klausmeier, and E. Litchman. 2017. Temperature-nutrient interactions exacerbate sensitivity to warming in phytoplankton. *Global Change Biology* 23:3169-3280.
- Vasseur, D. A., J. P. DeLong, B. Gilbert, H. S. Greig, C. D. G. Harley, K. S. McCain, V. Savage et al. 2014. Increased temperature variation poses a greater risk to species than climate warming. *Proceedings of the Royal Society B* 281:20132612.
- Vasseur, D. A., and K. S. McCann. 2005. A mechanistic approach for modeling temperature-dependent consumer-resource dynamics. *The American Naturalist* 166:184-198.
- Waldschmidt, S., and C. R. Tracy. 1983. Interactions between a lizard and its thermal environment: implications for sprint performance and space utilization in the lizard *Uta stansburiana*. *Ecology* 64:476-484.

- Waldschmidt, S. R., S. M. Jones, and W. P. Porter. 1986. The effect of body temperature and feeding regime on activity, passage time, and digestive coefficient in the lizard *Uta stansburiana*. *Physiological Zoology* 59:376-383.
- Wilhoft, D. C. 1958. The effect of temperature on thyroid histology and survival in the lizard, *Sceloporus occidentalis*. *Copeia* 1958:265-276.
- Willmer, P. G. 1982. Microclimate and the environmental physiology of insects. *Advances in Insect Physiology* 16:1-57.
- Woodworth-Jefcoats, P. A., J. J. Polovina, and J. C. Drazen. 2017. Climate change is projected to reduce carrying capacity and redistribute species richness in North Pacific pelagic marine ecosystems. *Global Change Biology* 23:1000-1008.
- Wuertz, D., T. Setz, Y. Chalabi, C. Boudt, P. Chausse, and M. Miklovac. 2017. fGarch: Rmetrics - Autoregressive Conditional Heteroskedastic Modelling.
- Zhao, M., and S. W. Running. 2010. Drought-induced reduction in global terrestrial net primary productivity from 2000 through 2009. *Science* 329:940-943.

Figure Legends

Figure 1: Growth rate of fingerling salmon at fixed temperatures and food rations (from 0% (starvation) to excess, units = “% dry weight/day”). Inverted triangles indicate estimated T_{opt} for each food-ration level: T_{opt} and the upper threshold temperature for growth both decline with decreasing food ration. Curves are loess smoothers. Redrawn from data in Brett et al. (1969; 1971).

Figure 2: When T_b and food intake are held constant, thermal dependence of net energy gain is sensitive to the breadth of the thermal performance curve for energy intake (generalists, specialists) as well as to the steepness (Q_{10} , or the factorial increase in metabolic rate over +10°C shift) of the metabolic-temperature curve. A) Gross energy gain (declining from black to gray) under three different food levels and metabolic losses (red) at three Q_{10} values (2, 2.5, 3) for a thermal generalist and C) for a thermal specialist. Net energy gain for a generalist (B) and a specialist (D) under three food levels and with Q_{10} s of 2 (solid line) or 3 (dotted line). T_{opt} for each (red dots) and the upper temperature threshold for growth both decline with food ration, especially for generalists with steep Q_{10} .

Figure 3: Food levels at different environmental temperatures. Case 1: food level is independent of temperature, and solid lines indicate constant food levels of 1.0 (black), 0.75 (gray), and 0.5 (light gray). Case 2: food level declines with increasing mean environmental temperatures (dashed black line, $a = 0.3$). Line thickness is scaled to food level.

Figure 4: Net energy gain as a function of mean environmental temperature (θ) for thermal generalists (A, B) and specialists (C, D) during climate warming. Two levels of thermal environmental variation are considered: SD = 2°C (left column, A and C) and SD = 4°C (right column, B and D). The solid black lines net energy gain under excess food, and solid gray lines indicate food levels of 0.75 (darker) or 0.5 (lighter). The dashed black line indicates a sigmoidal decline in food with temperature (1.0 to 0.5). Solid red circles indicate T_{opt} for different food levels.

