

CH 10 - Complex Numbers

Luke Lu • 2025-11-19

Standard Form

Info – Definition of Complex Numbers

A **complex number** z in **standard form** is an expression of the form $z = x + yi$ where $x, y \in \mathbb{R}$.

The real number x is called the **real part** of z , and is written $\Re(x)$.

The real number y is called the **imaginary part** of z , and is written $\Im(z)$.

The set of complex numbers is

$$\mathbb{C} = \{x + yi : x, y \in \mathbb{R}\}$$

The complex number $z = x + yi$ and $w = u + vi$ are equal ($z = w$) if and only if $x = u, y = v$

A complex number z is said to be purely real if $y = 0$ (i.e. $1 = 1 + 0i$)

A complex number z is said to be purely imaginary if $x = 0$ (i.e. $i = 0 + 1i$)

0 is purely real and purely imaginary (i.e. $0 = 0 + 0i$)

Info – Complex Arithmetics

Let $z = a + bi$ and $w = c + di$ be complex numbers. Then the

Addition is defined as

$$z + w = (a + c) + (b + d)i$$

Multiplication is defined as

$$zw = (ac - bd) + (ad + bc)i$$

Examples:

Let $z = 2 + 3i, w = -1 + 7i$

1. $z + w = (2 - 1) + (3 + 7)i = 1 + 10i$

2. $zw = (-2 - 21) + (14 - 3)i = -23 + 11i$

3. $i^2 = ii = (0 + 1i) \cdot (0 + 1i) = (0 - 1) + (0 + 0)i = -1$

$\therefore i^2 = -1$

From (3), we can derive a easier way of multiplication

 **Tip** – Multiplication trick

$$(a + bi)(c + di) = ac + adi + bci + bdi^2 = (ac - bd) + (ad + bc)i$$

Properties of Complex Arithmetics

1. $z + 0 = 0 + z = z$
2. $z0 = 0z = 0$
3. $z + (-1)z = (-1)z + z = 0$
4. $z1 = 1z = z$


 **Info** – Multiplicative Inverse

For all complex numbers z , the multiplicative inverse of z exists if and only if $z \neq 0$. Moreover, for $a + bi \neq 0$, the multiplicative inverse is unique given by

$$z^{-1} = \frac{a}{a^2 + b^2} - \frac{b}{a^2 + b^2}i = \frac{a - bi}{a^2 + b^2}$$

Examples:

1. $\frac{(1-2i)-(3+4i)}{(5-6i)} = [(1-2i) - (3+4i)] \cdot (5-6i)^{-1} = (-2-6i) \cdot \frac{1}{25+36} \cdot (5+6i)$
 $= \frac{1}{61} \cdot (-10 - 30i - 12i + 36) = \frac{26-42i}{61} = \frac{26}{61} - \frac{42}{61}i$
2. $i^{2025} = (i^4)^{506} \cdot i = (-1)^{506} \cdot i = i$

 **Tip** – i^n Solutions

$$i^n = \begin{cases} 1, n \equiv 0 \pmod{4} \\ i, n \equiv 1 \pmod{4} \\ -1, n \equiv 2 \pmod{4} \\ -i, n \equiv 3 \pmod{4} \end{cases}$$

 **Info** – Properties of Complex Arithmetics

In complex arithmetic, the following properties are valid for $u, v, z \in \mathbb{C}$

1. Associativity of addition: $(u + v) + z = u + (v + z)$
2. Commutativity of addition: $u + v = v + u$
3. Additive identity: $0 = 0 + 0i$ has the property that $z + 0 = z$
4. Additive inverses: If $z = a + bi$, then there exists an additive inverse of z , written $-z$, with the property that $z + (-z) = 0$. The additive inverse of $z = a + bi$ is $-z = -a - bi$
5. Associativity of multiplication: $(uv)z = u(vz)$
6. Commutativity of multiplication: $uv = vu$
7. Multiplicative identity: $1 = 1 + 0i$ has property that $z1 = z$
8. Multiplicative inverses: If $z = a + bi \neq 0$, then there exists a multiplicative inverse of z , written z^{-1} , with the property that $zz^{-1} = 1$. The multiplicative inverse of $z = a + bi \neq 0$ is $z^{-1} = \frac{a-bi}{a^2+b^2}$
9. Distributivity: $z(u + v) = zu + zv$

Example:

Proof of PCA Part 5: $(uv)z = u(vz) \quad \forall u, v, z \in \mathbb{C}$

Let $u = a + bi, v = c + di, z = x + yi$ where $a, b, c, d, x, y \in \mathbb{R}$ so that $u, v, z \in \mathbb{C}$

$$(uv)z = [(a + bi)(c + di)](x + yi) = (ac - bd + (ad + bc)i)(x + yi) =$$

$$= ((ac - bd)x - (ad + bc)y) + ((ac - bd)y + (ad + bc)x)i$$

$$= (axc - bdx - ady - bcy) + (acy - bdy + adx + bcx)i$$

$$u(vz) = (a + bi)[(c + di)(x + yi)] = (a + bi)(cx - dy) + (cy + dx)i$$

$$= (a(cx - dy) - b(cy + dx)) + (a(cy + dx) + b(cx - dy))i$$

$$= (acx - ady - bcy - bdx) + (acy + adx + bcx - bdy)i$$

$$\text{Thus } (uv)z = u(vz) = (axc - bdx - ady - bcy) + (acy - bdy + adx + bcx)i$$

□

Info – Other Arithmetic of Complex Numbers

For $z \in \mathbb{C}$

1. $z^0 = 1$
2. $z^1 = z$
3. $z^{k+1} = z^k z \quad \forall k \in \mathbb{N}$
4. $(z^n)^m = z^{nm}$ and $z^n z^m = z^{n+m} \quad \forall n, m \in \mathbb{N} \cup \{0\}$

For $n \notin \mathbb{N} \cup \{0\}$ will be discussed in later lectures

Examples:

$$\text{Find a real solution to } 6z^3 + (1 + 3\sqrt{2}i)z^2 - (11 - 2\sqrt{2})i - 6 = 0$$

Let $z = x \in \mathbb{R}$, that is $z = x + 0i$

$$\implies 6x^3 + (1 + 3\sqrt{2}i)x^2 - (11 - \sqrt{2}i)x - 6 = 0$$

$$\implies (6x^3 + x^2 - 11x - 6) + (3\sqrt{2}x^2 + 2\sqrt{2}x)i = 0 + 0i$$

$$\implies (6x^3 + x^2 - 11x - 6) = 0 \text{ and } 3\sqrt{2}x^2 + 2\sqrt{2}x = 0$$

$x = 0$ or $x = -\frac{2}{3}$ for the imaginary part.

However, $x = 0$ does not satisfy the real part.

$$\therefore x = -\frac{2}{3} \implies z = -\frac{2}{3} + 0i$$

Conjugate and Modulus

Info – Complex Conjugate

The complex **conjugate** of a complex number $z = x + yi$ written \bar{z} is the complex number

$$\bar{z} = x - yi$$



Info – Properties of Conjugate

For the complex conjugate, the following properties hold $\forall z, w \in \mathbb{C}$

1. $\overline{(\bar{z})} = z$
2. $\overline{z + w} = \bar{z} + \bar{w}$
3. $z + \bar{z} = 2\Re(z)$ and $z - \bar{z} = 2\Im(z)i$
4. $\overline{zw} = \bar{z} \cdot \bar{w}$
5. If $z \neq 0$, $\overline{(z^{-1})} = (\bar{z})^{-1}$
6. If $w \neq 0$, $\overline{(\frac{z}{w})} = \frac{\bar{z}}{\bar{w}}$

Proof

Part 3

$$z + \bar{z} = 2\Re(z) \text{ and } z - \bar{z} = 2\Im(z)i$$

Let $z = x + yi$

$$z + \bar{z} = (x + yi) + (x - yi) = 2x + 0i = 2\Re(z)$$

$$z - \bar{z} = (x + yi) - (x - yi) = 0x + 2i = 2\Im(z)i$$

Part 4

$$\overline{zw} = \bar{z} \cdot \bar{w}$$

Let $z = x + yi, w = a + bi$

$$\overline{zw} = \overline{(x + yi)(a + bi)} = \overline{(xa - yb) + (xb + ya)i} = (xa - yb) - (xb + ya)i$$

$$\begin{aligned} \overline{zw} &= \overline{x + yia + bi} = (x - yi)(a - bi) = (xa - (-y)(-b)) + (x(-b) + (-y)a)i \\ &= (xa - yb) - (xb + ya)i \end{aligned}$$

$$\text{Thus } \overline{zw} = \bar{z} \cdot \bar{w}$$

□

Examples:

1. Prove $z \in \mathbb{R} \iff z = \bar{z}$

$$\text{Let } z = x + yi \quad \forall x, y \in \mathbb{R}$$

\implies

Suppose $z \in \mathbb{R}$, then $y = 0$, so that $z = x + 0i = x \in \mathbb{R}$.

$$\text{T. hen } \bar{z} = x - 0i = x$$

$$\therefore z = \bar{z}$$

\longleftarrow

$$\text{Suppose } z = \bar{z}, \text{ then } x + yi = x - yi$$

This implies $x = x, y = -y$. Thus $y = 0$

$$\therefore z = x + 0i = x \in \mathbb{R}$$

2. Prove that z is purely imaginary $\Leftrightarrow z = -\bar{z}$

3. Solve $z^2 = i\bar{z}$

Let $z = x + yi$. Then $z^2 = (x + yi)^2 = x^2 - y^2 + 2xyi$

and $i\bar{z} = i(x - yi) = ix - i^2y = xi + y$

Then the equation becomes $x^2 - y^2 + 2xyi = xi + y$

$$\Rightarrow \begin{cases} x^2 - y^2 = y \\ 2xy = x \end{cases} \Rightarrow x = 0 \wedge y = \frac{1}{2} \Rightarrow \begin{cases} x^2 - \frac{1}{4} = \frac{1}{2} \\ -y^2 = 0 \end{cases} \Rightarrow z = \left\{ 0, -i, \pm \left(\frac{\sqrt{3}}{2} \right) + \frac{1}{2}i \right\}$$

Modulus of Complex Numbers

Info – Modulus of Complex Number

The **modulus** of the complex number $z = x + yi$, written $|z|$, is the non-negative real number

$$|z| = \sqrt{x^2 + y^2}$$

Info – Properties of Modulus

For the modulus, the following properties $\forall z, w \in \mathbb{C}$:

1. $|z| = 0 \Leftrightarrow z = 0$

2. $|\bar{z}| = |z|$

3. $\bar{z}z = |z|^2$

4. $|zw| = |z||w|$

5. If $z \neq 0$ then $|z^{-1}| = |z|^{-1}$

Side note: for $z \neq 0$, $z^{-1} = \frac{\bar{z}}{|z|^2}$

6. If $w \neq 0$, $\frac{z}{w} = z \cdot \frac{\bar{w}}{|w|^2}$

7. $|z + w|^2 + |z - w|^2 = 2(|z|^2 + |w|^2)$

8. $\overline{\sum z_i} = \sum \overline{(z_i)}$

9. $\overline{\prod z_i} = \prod \overline{(z_i)}$

10. $|\prod z_i| = \prod (|z_i|)$

Proof

Part 3

$$\bar{z}z = |z|^2$$

Let $z = a + bi$, $\forall a, b \in \mathbb{R}$

$$\text{then } \bar{z}z = (a - bi)(a + bi) = (a^2 - (-b)(b)) + (ab + (-b)a)i$$

$$= a^2 + b^2 + 0i = \left(\sqrt{a^2 + b^2} \right)^2 = |z|^2$$

The Complex Plane and Polar Form

💡 **Tip** — Graphical Properties

$z \wedge \bar{z}$ are reflection of each other over real axis