

## CH 2 — Sequence and Limits

Luke Lu • 2025-09-15

---

### Triangle Inequality

$$|x - y| \leq |x - z| + |z - y|$$

for  $x, y, z \in \mathbb{R}$

Idea: the straight-line distance is shortest.

Without loss of generality assume  $x \leq y$ ; swapping  $x, y$  preserves the statement.

Number-line proof by cases:

- Case 1  $z \leq x \leq y$ :  $|x - y| \leq |z - y| \leq |x - z| + |z - y|$
- Case 2  $x \leq z \leq y$ :  $|x - y| = |x - z| + |z - y|$
- Case 3  $x \leq y \leq z$ :  $|x - y| \leq |x - z| + |z - y|$

### Triangle Inequality 2

For all  $a, b \in \mathbb{R}$

$$|a + b| \leq |a| + |b|$$

Proof:

apply the triangle inequality to  $x = a, y = -b, z = 0$ .

### Quick check

Is  $|a - b| \leq |a| - |b|$  for all  $a, b$ ?

No

Example:

$a = 10, b = -9$  gives  $|10 - (-9)| = 19$ , while  $|10| - |-9| = 1$

Hence this statement is false.

### Interval translations

1.  $|x - a| < \delta \Rightarrow x \in (a - \delta, a + \delta)$
2.  $|x - a| \leq \delta \Rightarrow x \in [a - \delta, a + \delta]$
3.  $0 \leq |x - a| \leq \delta \Rightarrow x \in (a - \delta, a) \cup (a, a + \delta)$

### Practice

1) Solve  $|2x - 5| < 3$

$$-3 < 2x - 5 < 3 \Rightarrow 1 < x < 4$$

**Answer:**  $x \in (1, 4)$

2) Solve  $2 < |x + 7| \leq 3$

Split into  $|x + 7| > 2$  and  $|x + 7| \leq 3$

**Answer:**  $x \in [-10, -9] \cup (-5, -4]$

3) Solve  $\frac{|x+2|}{|x-2|} > 5$

Consider regions  $(-\infty, -2)$ ,  $(-2, 2)$ ,  $(2, \infty)$  and track signs of  $x + 2$  and  $x - 2$

**Answer:**  $x \in (\frac{4}{3}, 2) \cup (2, 3)$

## Infinite Sequences

A sequence is an ordered list  $a_1, a_2, a_3, \dots$ ; write  $(a_n)_{n=1}^{\infty}$

A subsequence chooses indices  $n_1 < n_2 < \dots$ , yielding  $a_{\{n_1\}}, a_{\{n_2\}}, \dots$

The tail with cutoff  $k$  is  $a_k, a_{\{k+1\}}, a_{\{k+2\}}, \dots$

## Convergence (definition)

### IMPORTANT

We say  $\lim_{n \rightarrow \infty} a_n = L$  if for every  $\varepsilon > 0$  there exists  $N$  such that  $n > N \Rightarrow |a_n - L| < \varepsilon$

Equivalent formulations:

- Every interval  $(L - \varepsilon, L + \varepsilon)$  contains a tail of  $(a_n)$
- Only finitely many terms lie outside  $(L - \varepsilon, L + \varepsilon)$
- More generally, any open interval  $(a, b)$  containing  $L$  contains a tail

### Examples

1) Show  $\lim_{n \rightarrow \infty} \frac{1}{\sqrt[3]{n}} = 0$

Choose  $N = \frac{1}{\varepsilon^3}$

Then  $n > N \Rightarrow \left| \frac{1}{\sqrt[3]{n}} \right| < \varepsilon$

2) Show  $\lim_{n \rightarrow \infty} \frac{3n^2+2n}{4n^2+n+1} = \frac{3}{4}$

Estimate  $\left| \frac{3n^2+2n}{4n^2+n+1} - \frac{3}{4} \right| \leq \frac{5}{16n+4}$

Pick  $N > \frac{5}{16\varepsilon} - \frac{1}{4}$

## Theorem (Equivalent definitions of the limit of a sequence)

### IMPORTANT

For a sequence  $(a_n)$  and a number  $L$ , the following are equivalent

- 1)  $\lim_{n \rightarrow \infty} a_n = L$
- 2) For every  $\varepsilon > 0$ , the interval  $(L - \varepsilon, L + \varepsilon)$  contains a tail of  $(a_n)$
- 3) For every  $\varepsilon > 0$ , only finitely many  $n$  satisfy  $|a_n - L| \geq \varepsilon$
- 4) Every interval  $(a, b)$  containing  $L$  contains a tail of  $(a_n)$
- 5) Given any interval  $(a, b)$  containing  $L$ , only finitely many terms of  $(a_n)$  lie outside  $(a, b)$

### Example 1

Show  $\lim_{n \rightarrow \infty} \frac{1}{\sqrt[3]{n}} = 0$

Side work:

$$\left| \frac{1}{\sqrt[3]{n}} \right| < \varepsilon \Rightarrow \frac{1}{\sqrt[3]{n}} < \varepsilon \Rightarrow \sqrt[3]{n} > \frac{1}{\varepsilon} \Rightarrow n > \frac{1}{\varepsilon^3}$$

### Proof

Let  $\varepsilon > 0$  and choose  $N = \frac{1}{\varepsilon^3}$

If  $n > N$  then  $\left| \frac{1}{\sqrt[3]{n}} \right| < \frac{1}{\sqrt[3]{N}} = \frac{1}{\sqrt[3]{\frac{1}{\varepsilon^3}}} = \varepsilon$

### Example 2)

Prove  $\lim_{n \rightarrow \infty} \frac{3n^2+2n}{4n^2+n+1} = \frac{3}{4}$

Rough work

$$\left| \frac{3n^2+2n}{4n^2+n+1} - \frac{3}{4} \right| = \frac{|5n-3|}{16n^2+4n+4} \leq \frac{5n}{16n^2+4n} = \frac{5}{16n+4}$$

### Proof

Given  $\varepsilon > 0$ , pick  $N > \frac{5}{16\varepsilon} - \frac{1}{4}$

Then for  $n > N$

$$\left| \frac{3n^2+2n}{4n^2+n+1} - \frac{3}{4} \right| < \frac{5}{16n+4} \leq \frac{5}{16N+4} < \varepsilon$$

### Limits

Thinking question: can a sequence converge to two different limits  $L \neq M$ ?

No, we are saying then  $\varepsilon < \frac{|L-M|}{2}$

If  $a_n \rightarrow L$  a tail of the sequence lies in  $(L - \varepsilon, L + \varepsilon)$  so only finite many terms can lie in the interval  $(M - \varepsilon, M + \varepsilon)$ , that is  $a_n \nrightarrow M$

### IMPORTANT

#### Theorem (Uniqueness of Limits):

Let  $\{a_n\}$  be a sequence. If  $\{a_n\}$  has limit  $L$ , then the value  $L$  is unique.

We say that a sequence **diverges to  $\infty$**  if for every  $m > 0$ , there exists  $N \in \mathbb{N}$  such that for all  $n > N$ ,  $a_n > m$ .

We say that a sequence **diverges to  $\infty$**  if any interval of the form  $(m, \varepsilon)$  for some  $m > 0$  contains a tail of  $\{a_n\}$ . We write that  $\lim_{n \rightarrow \infty} a_n = \infty$

We say that a sequence **diverges to  $-\infty$**  if for every  $m < 0$ , there exists  $N \in \mathbb{N}$  such that for all  $n > N$ ,  $a_n < m$

We say that a sequence **diverges to  $-\infty$**  if any interval of the form  $(m, \varepsilon)$  for some  $m < 0$  contains a tail of  $\{a_n\}$ . We write that  $\lim_{n \rightarrow \infty} a_n = -\infty$

Thinking questions:

1. If a sequence consists of non-negative terms, is the limit non-negative?

ANS: YES Suppose not, then  $a_n \rightarrow L$ ,  $a_n > 0, \forall n$ . Consider  $\varepsilon < \frac{|L|}{2}$ . Then  $(L - \varepsilon, L + \varepsilon)$  only contains negative numbers, so it can't include a tail of  $a_n$ , contradiction.

2. If a sequence consists of positive terms, is the limit positive?

ANS: NO, consider the sequence  $\{\frac{1}{n}\}$ ,  $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$

3.  $\lim_{n \rightarrow \infty} a_n + b_n = a + b$

4.  $\lim_{n \rightarrow \infty} a_n b_n = ab$

5.  $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{a}{b}, b \neq 0$

6. If  $a_n \geq 0 \forall n, \alpha > 0$ , then  $\lim_{\{n \rightarrow \infty\}} a_n^\alpha = a^\alpha$

7. For any  $k \in \mathbb{N}$ ,  $\lim_{n \rightarrow \infty} a_n = a$

Examples: Prove that  $\lim_{\{n \rightarrow \infty\}}$  Let  $m > 0$  and consider the interval  $m, \infty$ . If  $n > \sqrt[3]{m}$  then  $n^3 > m$  and so  $n^3 \in (m, \infty)$ . So choose  $k = \lceil \sqrt[3]{m} \rceil + 1$ , then the tails lies in  $(m, \infty)$

## Limit Laws

### IMPORTANT

Let  $\{a_n\}, \{b_n\}$  be sequences with  $\lim_{n \rightarrow \infty} a_n = a, \lim_{n \rightarrow \infty} b_n = b$  for some  $a, b \in \mathbb{R}$  then:

1. For any  $c \in \mathbb{R}$ , if  $a_n = c$  for all  $n$  then  $c = a$

2. For any  $c \in \mathbb{R}$ , if  $\lim_{n \rightarrow \infty} ca_n = ca$

Prove the SUM of Sequences Rule

### Proof

$$a_n \rightarrow a, b_n \rightarrow b$$

$$\forall \varepsilon > 0, \exists M, N \in \mathbb{R}, \forall n > M, n > N, |a_n - a| < \varepsilon, |b_n - b| < \varepsilon$$

$$|(a_n + b_n) - (a + b)| = |(a_n - a) + (b_n - b)| \leq |a_n - a| + |b_n - b| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon$$

### IMPORTANT

Tandem Convergence Theorem:

If  $\lim_{n \rightarrow \infty} \frac{a_n}{b_n}$  exists and  $\lim_{n \rightarrow \infty} b_n = 0$ , then  $\lim_{n \rightarrow \infty} a_n = 0$