CH 2 — Linear Transformations

Luke Lu • 2026-10-02

2.1 Linear Transformations Between Abstract Vectors

📘 **Definition — Linear Map**

Let V, W be vector spaces over $\mathbb{F}$

A **linear map** from V to W is a function $L : V \to W$ satisfiying

$$L(\vec{x} + \vec{y}) = L(\vec{x}) + L(\vec{y})$$
$$L(c\vec{x}) = cL(\vec{x})$$

Remarks:

1. L is linear if and only if $L(c\vec{x} + \vec{y}) = cL(\vec{x}) + L(\vec{y}), \quad \forall \vec{x}, \vec{y} \in V, c \in \mathbb{F}$
2. $L\left(\vec{0}\right) = \vec{0}$

Proof

$L\left(\vec{0}\right) = L\left(\vec{0} + \vec{0}\right) = L\left(\vec{0}\right) + L\left(\vec{0}\right) \implies \vec{0} = L\left(\vec{0}\right)$

□

Examples:

1. Zero map: $\vec{z} : V \to W$ defined by $\vec{z}(x) = \vec{0}_W, \quad \forall \vec{x} \in V$
2. Identity map: $\mathrm{Id} : V \to W$ defined by $\vec{x} \to \vec{x}$
3. B-coordinates. $[]_B : V \to \mathbb{F}^n$ is linear defined by $\vec{x} \to [\vec{x}]_B$
4. Let $A \in M_{m\times n}(\mathbb{F})$, let $L_A : \mathbb{F}^n \to \mathbb{F}^m$ defined by $\vec{x} \to A\vec{x}$
5. Evaulation is linear. Let $t_0 \in \mathbb{F}$, $\mathrm{ev}_0 : P_n(\mathbb{F}) \to \mathbb{F}$ defined by $p(x) \to p(t_0)$ $\mathrm{ev}_0 : P_2(\mathbb{F}) \to \mathbb{F} \implies p(x) \to p(0) \implies a + bx + cx^2 \to a)$
6. Differentiation and integration are linear.

   Consider $D : P_n(\mathbb{F}) \to P_{n-1}(\mathbb{F})$ defined by $p(x) \to p'(x)$

   This is linear because

   $$D(p + q) = (p + q)' = p' + q' = D(p) + D(q)$$
   $$D(cp) = cp' = cD(p)$$

   Similarly, $L : P_{n(\mathbb{R})} \to \mathbb{R}$ defined by $p(x) \to \int_0^1 p(x)\,\mathrm{d}x$
7. Trace map is linear. $M_{n\times n}(\mathbb{F}) : A \to \mathrm{tr}(A)$

💡 **Tip — Combining Linear Maps**
Suppose $L, T : V \to W$.

1. $L + T$ is linear
2. cL is linear $\forall c \in \mathbb{F}$
3. If $S : W \to U$ is linear, then $S \circ T : V \to W$ is linear

Proof

If $S : W \to U$ is linear, then $S \circ T : V \to W$ is linear

$$\begin{aligned}(S \circ T)(c\vec{x} + \vec{y}) &= S(T(c(\vec{x}) + \vec{y})) \\ &= S(cT(\vec{x}) + T(\vec{y})) = cS(T(\vec{x})) + S(T(\vec{y})) \\ &= c(S \circ T)(\vec{x}) + (S \circ T)(\vec{y})\end{aligned}$$

□

💡 **Tip — Non-Linear Maps**(Rule of thumb)

1. Anything with $L\left(\vec{0}\right) \neq \vec{0}$ is probably non-linear
2. Anything involving muliplicators scaling probably is non-linear

Examples of non-linear maps:

1. $L(p(x)) = p(x) + 1$ is non-linear
2. $L(p(x)) = [p(x)]^2$ is non-linear
3. $L(A) = L(\det A)$ is non-linear
4. $L(p(x)) = xp(x)$ IS LINEAR

2.2 Rank-Nullity

IMPORTANT:

1. State and Prove the Fundamental of Linear Algebra(Rank-Nullity Theorem)
2. Show linear maps $\iff$ matrices

📘 **Definition — Kernel and Range**

Let $L : V \to W$ be a linear map.

1. $\text{Ker}(L) = \{\vec{v} \in V : L\left(\vec{v} = \vec{0}_W\right\}$ is a subspace of V
2. $\text{Range}(L) = \{\vec{w} \in W : \vec{w} = L(\vec{v}))\}$ is a subspace of W

Proof

$\text{Ker}(L) = \{\vec{v} \in V : L\left(\vec{v} = \vec{0}_W\right\}$ is a subspace of V

$L\left(\vec{0}_V\right) = \vec{0}_W, \vec{0}_V \in \text{Ker}(L)$

Let $\vec{x}, \vec{y} \in \text{Ker}(L), c \in \mathbb{F}$

$L(c\vec{x} + \vec{y}) = cL(\vec{x}) + L(\vec{y}) = \vec{0}_W \in W$, thus $c\vec{x} + \vec{y} \in \text{Ker}(L)$

□

Example:

$L = []_B \implies L : V \to \mathbb{F}^n$

$\text{Ker}(L) = \left\{ \vec{x} \in V : [\vec{x}]_B = \begin{bmatrix} 0 \\ \dots \\ 0 \end{bmatrix} \right\} = \left\{\vec{0}_V\right\}$, by the Uniqueness of B-coordinates

$\text{Range}(L) = \{[\vec{x}]_B : \vec{x} \in V\} = \mathbb{F}^n$

Definition — Nullity-Rank Theorem

The **nuliity** of L is

$$\text{nullity } L = \dim(\text{Ker } L)$$

The **rank** of L is

$$\text{rank } L = \dim(\text{Range } L)$$

Let $L : V \to W$ be linear mapping.

$$\dim V = \text{nullity } L + \text{rank } L$$
$$\dim V - \text{nullity } L = \text{rank } L$$

Total degrees - lost degrees = output degrees

Examples:

1. Let $A \in M_{m \times n}(\mathbb{F}), L_A : \mathbb{F}^n \to \mathbb{F}^m, \vec{x} \to A\vec{x}$

   a. Ker $L_A = \text{Null} A$ is the nullspace

   b. Range $L_A = \text{Col } A$ is the column space

   Proof

   $$\begin{aligned} \text{Ker } L_A &= \{\vec{x} \in \mathbb{F}^n : L_A(\vec{x}) = 0\} \\ &= \left\{\vec{x} \in \mathbb{F}^n : A\vec{x} = \vec{0}\right\} \\ &= \text{Null } A \end{aligned}$$

$$\begin{aligned}\text{Range } L_A &= \{\vec{x} \in \mathbb{F}^n : \vec{y} = L_A(\vec{x})\} \\ &= \{\vec{x} \in \mathbb{F}^n : \vec{y} = 0A\vec{x}\} \\ &= \{y \in \mathbb{F}^m : \vec{y} = [\vec{a}_1, ..., \vec{a}_n]\vec{x}\} = \{y \in \mathbb{F}^m : \vec{y} = \vec{a}_1 x_1 + ...\vec{a}_n x_n, (x_1, ..., x_n) \in \mathbb{F}\} \\ &= \text{Span } \{\vec{a}_1, ..., \vec{a}_n\} = \text{Col } A\end{aligned}$$

□

This example will tell us eventually how to compute $\text{Ker}(L)$ and Range (L) in general

This result gives useful interpretation for the rank $A = \dim\ (\text{Range}(L))$

2. Let $D : P_n(\mathbb{R}) \to P_{n-1}(\mathbb{R}), p(x) \to p'(x)$ be differentiation

   $P_2(\mathbb{R}) \to P_1(\mathbb{R}) \iff a + bx + cx^2 \to b + 2cx \iff \begin{bmatrix} a \\ b \\ c \end{bmatrix} \iff \begin{bmatrix} 0 \\ b \\ 2c \end{bmatrix}$

   Educated guess: nullity $D = 1, \text{rank } D = (1 + n) - 1 = n$

   Proof

   $$\begin{aligned}\text{Ker } D &= \left\{p(x) \in P_n(\mathbb{F}) : D(p(x)\} = \vec{0}\right\} = \{a_1 + a_2 x + ... + a_n x^n\} = \{0 + 0x + ... + 0x^n\} \quad (a_1 = a_2 = ... = \\ &= \left\{a_1 + 0x^2 + ... + 0x^n : a_i \in \mathbb{F}\right\} = \text{span } \{1\} \in P_n(\mathbb{F})\end{aligned}$$

   Since $\{1\}$ is a basis for Ker $D \implies \dim(\text{Ker } D) = 1$
   $\text{Range}(D) = \{q(x) \in P_{n-1}(\mathbb{R}) : q(x) = D(p(x)), \text{for some } p(x) \in P_n(\mathbb{R})\}$

   Claim Range D $= P_{n-1}(\mathbb{R})$
   Range D $\subseteq P_{n-1}(\mathbb{R})$ is trivial

   $P_{n-1}(\mathbb{R}) \subseteq$ Range D
   Let $q(x) = a_0 + a_1 x + ... + a_{n-1} x^{n-1}$
   Then $q(x) = D\big(a_0 + \frac{a_1}{2} x + ... + \frac{a_{n-1}}{n} x^{n-1}\big) \in P_n(\mathbb{R})$
   $q(x) \in \text{Range}(D) \implies \text{rank } L = \dim \text{ Range } L = \dim\ P_{n-1}(\mathbb{R}) = n$

□

Proof

Rank-Nullity Theorem

Let $B = \{v_1, ..., v_k\}$ be a basis for Ker L, extend $B.$ to obtain $C = \{v_1, ..., v_k, v_{k+1}, ..., v_n\}$

Claim $D = \{L(v_{k+1}), ..., L(v_n)\}$ is a basis for Range L, meaning that

rank $L = \dim$ Range $L = |D| = n - k = \dim V - \text{nullity} L$

Consider $c_{k+1}L(v_{k+1}) + ... + c_k L(v_n) = \vec{0}, \quad \forall c_i \in \mathbb{F}$

$$\begin{aligned}L(c_{k+1}v_{k+1} + ... + c_n v_n) = \vec{0} \implies c_{k+1}v_{k+1} + ... + c_n v_n &\in \text{Ker } L \\ = c_{k+1}v_{k+1} + ... + c_n v_n &= a_1 v_1 + ... + a_k v_k \\ = c_{k+1}v_{k+1} + ... + c_n v_n - a_1 v_1 - ... - a_k v_k &= \vec{0}\end{aligned}$$

Since C is linearly independent, thus $c_i = 0$

We need to prove D spans Range L : span$D$ = Range$L$ to show the claim is true

span$D \subseteq$ Range$L$ is trivial

Range$L \subseteq$ span$D$

Given $w \in$ Range L, show $w \in$ span D.

We can write $a_1v_1 + ... + a_kv_k + a_{k+1}v_{k+1} + ... + a_nv_n$

So $w = L(v) = L(\sum a_iv_i) = a_{k+1}v_{k+1} + ... + a_nv_n \in$ span D

□

📘 **Definition — Isomorphism**
Let $L : V \to W$ be linear mapping

1. L is **injective**(one-to-one) $\iff$ Ker$L = \{0\} \iff$ nullity$L = 0$
2. L is **surjective**(onto) $\iff$ Range$L = W \iff$ rank $L =$ dim$W$
3. L is an **isomorphism** if and only if Ker$L = \{0\}$ and Range$L = W$

Example:

1. Let B be a basis for V, then the coordinate map $[]_B : V \to \mathbb{F}^n$ is an isomorphism

   $\because$ Ker$[]_B = \{0\}$ thus is injective, Range$[]_B = \mathbb{F}^n$ thus is surjective
2. $P_2(\mathbb{R}) \to \mathbb{R}^3$ is an isomorphism
3. $D : P_n(\mathbb{R}) \to P_{n-1}(\mathbb{R}), p(x) \to p'(x)$ be differentiation

   nullity $D = 1 \neq 0$, thus not injective, rank$D =$ dimP_{n-1} is surjective
4. $L : \mathbb{F}^2 \to M_{2\times 2}(\mathbb{F}), \begin{bmatrix} a \\ b \end{bmatrix} \to \begin{bmatrix} a+b & a \\ b & a-b \end{bmatrix}$

   We suspect that L cannot be surjective

   Range $L = \left\{ \begin{bmatrix} a+b & a \\ b & a-b \end{bmatrix} : a, b \in \mathbb{F} \right\} = \left\{ a \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \right\} + b \begin{bmatrix} 1 & 0 \\ 1 & -1 \end{bmatrix} : a, b \in \mathbb{F} \} = \text{span } \left\{ \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 1 & -1 \end{bmatrix} \right\}$

   rank$L \neq$ dim Range L, so L is not surjective

   By rank-nullity theorem, nullity $L =$ dim$\mathbb{F}^2 -$ rank $L = 0$