CH 8 - Modular Arithmetics

Luke Lu • 2026-10-19

Basic Modular Arithmetics

> **Info — Congruence and Modular Expression**
> Let m be a fixed positive integer. For integers a and b, we say that a is **congruent** to b **modulo** m, and write
>
> $$a \equiv b \pmod{m}$$
>
> if and only if $m \mid (a-b)$. For integers a and b such that $m \nmid (a-b)$, we write $a \not\equiv b \pmod{m}$. We refer to $\equiv$ as **congruence**, and m as its **modulus**.
>
> $$a \equiv b \pmod{m} \iff m \mid (a-b) \iff \exists k \in \mathbb{Z}, a-b=km \iff \exists k \in \mathbb{Z}, a=km+b$$

Examples:

1. $6 \equiv 18 \pmod{12} : 6-18=-12, 12 \mid -12$
2. $73 \equiv 1 \pmod{2} : 13-1=72, 2 \mid 72$
3. $5 \equiv 1 \pmod{4} : 5-1=4, 4 \mid 4$
4. $24 \equiv 0 \pmod{24} : 24-0=24, 24 \mid 24$
5. $-5 \equiv 7 \pmod{12} : -5-7=-12, 12 \mid -12$

> **Info — Equality Properties**
> 1. Reflexivity: $\forall a \in \mathbb{Z}, a=a$
> 2. Symmetry: $\forall a, b \in \mathbb{Z}, a=b \implies b=a$
> 3. Transitivity: $\forall a, b, c \in \mathbb{Z}, a=b \land b=c \implies a=c$

> **Info — Congruence Relations**
> $\forall a, b, c \in \mathbb{Z}$
> 1. $a \equiv a \pmod{m}$
> 2. $a \equiv b \pmod{m} \implies b \equiv a \pmod{m}$
> 3. $a \equiv b \pmod{m} \land b \equiv c \pmod{m} \implies a \equiv c \pmod{m}$

Info – Basic Modular Operations

$\forall a_1, a_2, b_1, b_2 \in \mathbb{Z}$ and $\forall n \in \mathbb{N}$, if $a_1 \equiv b_1 \pmod m$ and $a_2 \equiv b_2 \pmod m$ then

1. $a_1 + a_2 \equiv b_1 + b_2 \pmod m$
2. $a_1 - a_2 \equiv b_1 - b_2 \pmod m$
3. $a_1 a_2 \equiv b_1 b_2 \pmod m$
4. $a_1 + a_2 + ... + a_n \equiv b_1 + b_2 + ... + b_n \pmod m$
5. $a_i \equiv b_i \implies a_1 a_2 ... a_n \equiv b_1 b_2 ... b_n \pmod m$
6. $\forall a, b \in \mathbb{Z}$ if $a \equiv b \pmod m$ then $a^n \equiv b^n \pmod m$
7. $\forall a, b, c \in \mathbb{Z}$, if $ac \equiv bc \pmod m$ and $\gcd(c, m) = 1$, then $a \equiv b \pmod m$

Proof

$\forall a_1, a_2, b_1, b_2 \in \mathbb{Z}$ where $a_1 \equiv b_1 \pmod m$ and$a_2 \equiv b_2 \pmod m$

1. $a_1 + a_2 - b_1 - b_2 = a_1 - b_1 + a_2 - b_2 \pmod m$.

   Since $a_1 \equiv b_1 \pmod m$ and $a_2 \equiv b_2$, therefore $m \mid (a_1 - b_1)$ and $m \mid (a_2 - b_2)$.

   By DIC $m \mid (a_1 - b_1 + a_2 - b_2) \equiv m \mid (a_1 + a_2 - (b_1 + b_2))$.

   By definition of Congruence, $a_1 + a_2 \equiv b_1 + b_2 \pmod m$

2. $a_1 - a_2 - b_1 + b_2 = a_1 - b_1 + a_2 - b_2 \pmod m$.

   Since $a_1 \equiv b_1 \pmod m$ and $a_2 \equiv b_2 \pmod m$, therefore $m \mid (a_1 - b_1)$ and $m \mid (a_2 - b_2)$.

   By DIC $m \mid (a_1 - b_1 - a_2 + b_2) \equiv m \mid (a_1 - a_2 - (b_1 - b_2))$.

   By definition of Congruence, $a_1 - a_2 \equiv b_1 - b_2 \pmod m$

3. Since $a_1 \equiv b_1 \pmod m$ and $a_2 \equiv b_2 \pmod m$,

   therefore $\exists k, l \in \mathbb{Z}$ s.t. $a_1 = km + b_1; a_2 = lm + b_2$.

   $a_1 b_1 - b_1 b_2 = (km + b_1)(lm + b^2) - b_1 b_2 = klm^2 + kmb_2 + b_1 lm + b_1 lm + b_1 b_2$

   $(klm + kb_2 + b_1 l) \cdot m \implies m \mid (klm + kb_2 + b_1 l)$.

   Hence, $a_1 a_2 \equiv b_1 b_2 \pmod m$

□

Examples:

1. Is $5^9 + 62^{2000} - 14$ divisible by 7

   $5^9 + 62^{2000} - 14 \equiv 0 \pmod 7$

   $5^9 + 62^{2000} \equiv 0 \pmod 7$ since $14 \equiv 0 \pmod 7$

   $(5^2)^4 \cdot 5 + (-1)^{2000} \equiv 0 \pmod 7$ since $62 \equiv -1 \pmod 7$ because $62 - (-1) = 63, 7 \mid 63$

   $4^4 \cdot 5 + 1 \equiv 0 \pmod 7$ since $25 \equiv 4 \pmod 7$

   $2^2 \cdot 5 + 1 \equiv 0 \pmod 7$ since $7 \mid (16 - 2)$

   $21 \equiv 0 \pmod 7$ since $7 \mid 21$

   $\therefore 5^9 + 62^{2000} - 14 \equiv 0 \pmod 7$ since $7 \mid 5^9 + 62^{2000} - 14$, meaning, $5^9 + 62^{2000} - 14$ is divisible by 7.

2. Illustration of Congruence Divide
   $3 \equiv 27 \pmod 6$
   $3 \cdot 1 \equiv 3 \cdot 9 \pmod 6, 1 \not\equiv 9 \pmod 6$ since $\gcd(3,6) \neq 1$

Congruence and Remaidners

> 🐳 **Info — Congruent Iff Same Remainder**
>
> $\forall a, b \in \mathbb{Z}, a \equiv b \pmod m$ if and only if a and b have the same remainder when divided by m

> 🐳 **Info — Congruent to Remainder**
>
> $\forall a, b \in \mathbb{Z}$ with $0 \leq b < m, a \equiv b \pmod m$ if and only if a has a remainder b when divided by m

Examples:

1. What is the remaidner when $77^{100} \cdot 999 - 6^{83}$ divided by 4?
   $77 \equiv 1 \pmod 4$
   $999 \equiv -1 \pmod 4$
   $6 \equiv 2 \pmod 4$
   $\equiv 1^{100} \cdot -1 - 2^{83} \pmod 7$
   $\equiv -1 - 2^{82} \cdot 2 \equiv -1 - 2(4)^{41} \equiv -1 - 2(0) \equiv -1 \pmod 4$

   By CTR $3 \equiv -1 \pmod 4$, the remainder is 3

> 💡 **Tip — Divisibility by 3**
>
> For all non-negative integers a, a is divisible by 3 if and only if the sum of the digits in the decimal representation of a is divisible by 3

Proof

Let a be non-negative integer and expressed as

$a = d_k 10^k + d_{k-1} 10^{d-1} + ... + d_1 10 + d_0$ where $0 \leq d_i \leq 9$ are the digit $\forall i \in \mathbb{N} \cup \{0\}$

Notice $10 \equiv 1 \pmod 3$

$a \equiv d_k 1^k + d_{k-1} 1^{d-1} + ... + d_1 1^1 + d_0 \pmod 3$

$a \equiv \sum_{i=0}^{k} d_i \pmod 3$

Assume a is divisible by 3, then $3 \mid (a - 0) \iff a \equiv 0 \pmod 3$.

Since $a \equiv \sum_{i=0}^{k} d_i \pmod 3 \overset{\text{by CER}}{\iff} \sum_{i=0}^{k} d_i \equiv 0 \pmod 3$

Hence $3 \mid \sum_{i=0}^{k} d_i$

□

💡 **Tip — Divisbility by 11**
For all non-negative integers a, $11 \mid a$ if and only if $11 \mid (S_e - S_o)$ where

- S_e is the sum of all even digits of a in the decimal representation
- S_o is the sum of all odd digits of a in the decimal representation

💡 **Tip — Mod 7 or 13**
7. Remove last digit d, subtract $2d$, repeat.
13. Remove last digit d, add $4d$, repeat.

Linear Congruences

🐳 **Info — Definition or Linear Congruences**
A relation of the form

$$ax \equiv c \pmod{m}$$

is called a **linear congruence** in the variable x. A solution to such linear congruence is an integer x_0 s.t.

$$ax_0 \equiv c \pmod{m}$$

📐 **Theorem — Linear Congruence Theorem**
For all integers a, c where $a \neq 0$, the linear congruence

$$ax \equiv c \pmod{m}$$

has a solution if and only if $d \mid c$, where $d = \gcd(a, m)$. Moreover, if $x = x_0$ is one particular solution to this congruence, then the set of all solutions is given by

$$\left\{x \in \mathbb{Z} : x \equiv x_0 \pmod{\frac{m}{d}}\right\}$$

or alternatively

$$\left\{x \in \mathbb{Z} : x \equiv x_0, x_0 + \frac{m}{d}, x_0 + 2\frac{m}{d}, ..., x_0 + (d-1)\frac{m}{d} \pmod{m}\right\}$$

Examples:

1. $4x \equiv 5 \pmod{3}$
   $\gcd(a, m) = \gcd(4, 3) = 1, 1 \mid 5$
   By LCT, there is a solution
   $4x \equiv 5 \equiv 2 \equiv 8 \pmod{3} \implies 4x = 8 \implies x = 2$
   and $3 \mid (8 - 5)$
   By LCT, all solutions are $\{x \in \mathbb{Z}, x \equiv 2 \pmod{3}\}$.
2. $4x \equiv 8 \pmod{12}$
   $\gcd(a, m) = \gcd(4, 12) = 4, 4 \mid 8$

By LCT, there is a solution
$4x \equiv 8 \equiv 4(2) \pmod{12} \implies 4x = 8 \implies x = 2$
and $4 \mid 4 \mid (8-8)$
By LCT, all solutions are $\{x \in \mathbb{Z}, x \equiv 2 \pmod 3\}$
or $\{x \in \mathbb{Z}, x \equiv 2, 5, 8, 11 \pmod{12}\}$

Non-linear Congruences

> 💡 **Tip** — Non-linear congruences do not have theorems that directly helps solving. The solutiosn generally are by brute force

Examples:

$x^2 \equiv 6 \pmod{10}$

$x \pmod{10}$	0	1	2	3	4	5	6	7	8	9
$x^2 \pmod{10}$	0	1	2	9	6	5	6	9	4	1

Hence $x \equiv 4, 6 \pmod{10}$

Congruence Classes and Modular Arithmetic

> 🐳 **Info — Congruence class**
>
> The **congruence class** modulo m of the integer a is the set of integers
>
> $$[a] = \{x \in \mathbb{Z} : x \equiv a \pmod m\}$$

> 🐳 **Info — Modular Arithmetic**
>
> We define $\mathbb{Z}_m$ to be the set of m congruence classes
>
> $$\mathbb{Z}_m = \{[0], [1], [2], ..., [m-1]\}$$
>
> and we define two operations on $\mathbb{Z}_m$, **addition** and **multiplication**, as follows:
>
> $$[a] + [b] = [a + b)]$$
>
> $$[a][b] = [ab]$$
>
> When we apply these operations on the set $\mathbb{Z}_m$, we are doing that is known as **modular arithmetic**

> 🐳 **Info — Basic Properties of Congruence Classes**
>
> For all $[a] \in \mathbb{Z}_m$
>
> 1. $[a] + [0] = [a]$
> 2. $[a][0] = [0]$
> 3. $[a] + [-a] = [0]$
> 4. $[a][1] = [a]$

Example:

Construct a table for $\mathbb{Z}_4$
That is $\mathbb{Z}_4 = \{[0], [1], [2], [3]\}$

Addition table

$+$	$[0]$	$[1]$	$[2]$	$[3]$
$[0]$	$[0]$	$[1]$	$[2]$	$[3]$
$[1]$	$[1]$	$[2]$	$[3]$	$[0]$
$[2]$	$[2]$	$[3]$	$[0]$	$[1]$
$[3]$	$[3]$	$[0]$	$[1]$	$[2]$

Multiplicaiton table

$*$	$[0]$	$[1]$	$[2]$	$[3]$
$[0]$	$[0]$	$[0]$	$[0]$	$[0]$
$[1]$	$[0]$	$[1]$	$[2]$	$[3]$
$[2]$	$[0]$	$[2]$	$[0]$	$[2]$
$[3]$	$[0]$	$[3]$	$[2]$	$[1]$

📐 **Theorem — Modular Arithmetic Theorem**

For all integers a and c, with a non-zero, the equation

$$[a][x] = [c]$$

in $\mathbb{Z}_m$ has a solution if and only if $d \mid c$, where $d = \gcd(a, m)$. Moreover, when $d \mid c$, there are d solutions, given by

$$[x_0], \left[x_0 + \frac{m}{d}\right], \left[x_0 + 2\frac{m}{d}\right], ..., \left[x_0 + (d-1)\frac{m}{d}\right]$$

where $x = [x_0]$ is one particular solution

🐳 **Info — Inverse to $\mathbb{Z}_m$**

Let a be an integer with $1 \leq a \leq m - 1$. The element $[a]$ in $\mathbb{Z}_m$ has a multiplicative inverse if and only if $\gcd(a, m) = 1$. Moreover, when $\gcd(a, m) = 1$, the multiplicative inverse is unique.

🐳 **Info — Inverse to $\mathbb{Z}_p$**

For all prime numbers p and non-zero element $[a]$ $\mathbb{Z}_p$ the multiplicative inverse $[a]^-1$ exists and is unique

Examples:

In $\mathbb{Z}_{10}$, solve the following:

1. $[12][x]+[3]=[8]=[2][x]+[3]=[8] \implies [2][x]=[5] \overset{\text{by MAT}}{\implies}$ No solutions
2. $[15][x]+[7]=[12]=[5][x]=[5] \overset{\text{by MAT}}{\implies} [x]=[1]$

   Also by MAT, there are 5 solutions: $\{[1],[3],[5],[7],[9]\}$

> 📐 **Theorem — Fermat's Little Theorem**
>
> For all prime numbers p, integers a not divisible by p and $a \neq 0$, we have
>
> $$a^{p-1} \equiv 1 \pmod{p}$$

> 💡 **Tip** — Additional Corollaries
>
> 1. By FℓT, $[a]^{-1}=[a]^{p-2}$
> 2. For all prime numbers p and integers a, we have
>
> $$a^p \equiv a \pmod{p}$$

Examples:

1. Find the remainder when 7^{92} is divided by 11
   In $\mathbb{Z}_{11}$, $[7]^{10}=[1]$ by FℓT

   $7^{92}=(7^{10})^9 7^2=[1]^9[7^2]=[49]=[5]$

   So the reimainder upon dividing 7^{92} by 11 is 5
2. If p is prime, $p \nmid a$ and $r \equiv s \pmod{(p-1)}$, then $a^r \equiv a^s \pmod{p}$

   Let p be prime, $a \in \mathbb{Z}$ with $p \nmid a$

   Suppose $r \equiv s \pmod{(p-1)} \implies r-s=k(p-1)$ for some $k \in \mathbb{Z}$

   $r=k(p-1)+s \implies a^r \equiv a^{k(p-1)+s} \equiv a^s\left(a^{p-1}\right)^k \overset{\text{by F}\ell\text{T}}{\equiv} a^s(1)^k \equiv a^s \pmod{p}$
3. If $r=s+kp$, then $a^r \equiv a^{s+k} \pmod{p}$

Chinese Remainder Theorem

> 📐 **Theorem — Chinese Remainder Theorem**
>
> For all integers a_1 and a_2, and positive integers m_1 and m_2, if $\gcd(m_1,m_2)=1$, then the simultaneous linear congruences
>
> $$n \equiv a_1 \pmod{m_1}$$
>
> $$n \equiv a_2 \pmod{m_2}$$
>
> have a unique solution modulo $m_1 m_2$. Thus, if $n=n_0$ is one particular solution, then the solutions are given by the set of all integers n such that
>
> $$n=n_0 \pmod{m_1 m_2}$$

Info — Generalized CRT

For all positive integers k and $m_1, m_2, ..., m_k$ and integers $a_1, a_2, ..., a_k$, if $\gcd(m_i, m_j) = 1$ for all $i \neq j$, then the simultaneous congruences

$$n \equiv a_1 \pmod{m_1}$$

$$n \equiv a_2 \pmod{m_2}$$

$$...$$

$$n \equiv a_k \pmod{m_k}$$

have a unique solution modulo $m_1 m_2 ... m_k$. Thus, if $n = n_0$ is one particular solution, then the solutions are given by the set of all integers n s.t.

$$n \equiv n_0 \pmod{m_1 m_2 ... m_k}$$

Examples:

1. Find all x s.t. $x \equiv 2 \pmod{13}$ and $x \equiv 17 \pmod{29}$

   From $x \equiv 17 \pmod{29}$, we have $x - 17 = 29k$ for some $k \in \mathbb{Z}$. Then $x = 17 + 29k$

   So, $17 + 29k \equiv 2 \pmod{13} \implies 4 + 3k \equiv 2 \pmod{13}$

   $3k \equiv 11 \pmod{13}$ because 2-4 = −2, which is congruent to 11 (mod 13)

   $3k - 11 = 13j$ for some $j \in \mathbb{Z}, \implies 3k - 13j = 11 \implies k_0 = 8, j_0 = 1.$

   $x = 17 + 29(8) = 249, \{x \in \mathbb{Z} : x \equiv 249 (\mathrm{mod}\, 377)\}$

2. Find all x satisfying
   1. $x \equiv 5 \pmod{6}$
   2. $x \equiv 2 \pmod{7}$
   3. $x \equiv 3 \pmod{11}$

   $x \equiv 58 \pmod{77} \implies x \equiv 443 \pmod{462}$

3. Solve
   1. $3x \equiv 2 \pmod{5}$
   2. $2x \equiv 6 \pmod{7}$

   ANS: Multiply (1) by 2 and (2) by 3.
   - $6x \equiv 4 \pmod{5}$
   - $6x \equiv 18 \equiv 4 \pmod{7}$

   Let $y = 6x$

   Then $y \equiv 4 \pmod{5}, y \equiv 4 \pmod{7}$
   By CRT, $6x \equiv 4 \pmod{35} \implies x \equiv 24 \pmod{35}$

4. Solve

1. $x \equiv 4 \pmod 6$
2. $x \equiv 2 \pmod 8$

ANS: From (2), $x - 2 = 8k$ for some $k \in \mathbb{Z} \implies x = 2 + 8k$
(1) becomes $2 + 8k \equiv 4 \pmod 6 \implies 2k \equiv 2 \pmod 6$.
By inspection, $k = 1$ is one solution, so $x = 2 + 8(1) = 10$ is a solution.
Since $\gcd(6, 8) = 2 \neq 1$. We then have $x \equiv 10 \pmod 6$ and $x \equiv 10 \pmod 8 \implies x \equiv 10 \pmod{(\text{lcm}(6, 8))} \implies x \equiv 10 \pmod{24}$.

5. Solve $x^{12} \equiv 5 \pmod{55}$

We apply CRT in reverse to obtain the following simultaneous congruences:

1. $x^{12} \equiv 5 \pmod{11} \land x^{12} \equiv 0 \pmod 5$
2. $x^{12} \equiv 5 \pmod{11} \overset{\text{by CF}\mathscr{C}\text{T}}{\implies} x^2 \equiv 5 \pmod{11}$

We got $x \equiv 4 \land x \equiv 7 \pmod{11}$

Table mod 11

x	0	1	2	3	4	5	6	7	8	9	10
x^2	0	1	4	9	5	3	3	5	9	4	1

For (1), $x \equiv 0 \pmod 5$

We have one of these two congruences
3. $x \equiv 4 \pmod{11} \land x \equiv 0 \pmod 5$
4. $x \equiv 7 \pmod{11} \land x \equiv 0 \pmod 5$

$x \equiv 15 \ (55)$, so by CRT it is all solutions to (3) .
$x \equiv 40 \pmod{55}$, so by CRT is all soluions to (4).

$\therefore x \equiv 15, 40 \pmod{55}$ are solutions to $x^{12} \equiv 5 \pmod{55}$

> 📐 **Theorem — Splitting Modulus Theorem**
>
> For all integers a and positive integers m_1 and m_2, if $\gcd(m_1, m_2) = 1$, then the simultaneous congruences
>
> $$n \equiv a \pmod{m_1}$$
>
> $$n \equiv a \pmod{m_2}$$
>
> has exactly the same solutions as the single congruence $n \equiv a \pmod{m_1 m_2}$

Example:

For which integers x. is $x^5 + x^3 + 2x^2 + 1$ is divisible by 6
ANS: No solutions. Idea: Table of Congruences