CH 3 - Matrices and Linear Mapping

Luke Lu • 2026-10-19

Info — Matrix Definition

A $m \times n$ **matrix** A is a rectangular array with m rows and n columns. We denote the entry in the i-th row and j-th column by a_{ij} or $(A)_{ij}$.

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1j} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2j} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots & \dots & \dots \\ a_{i1} & a_{i2} & \dots & a_{ij} & \dots & a_{jn} \\ \dots & \dots & \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & a_{mj} & \dots & a_{mn} \end{bmatrix}$$

Two $m \times n$ matrices A, B are equal if $a_{ij} = b_{ij},\ \forall 1 \le i \le m, 1 \le j \le n$.

The set of all $m \times n$ matrices with real entries is denoted by $M_{m\times n}(\mathbb{R})$

Let $A, B \in M_{m\times n}(\mathbb{R}), c \in \mathbb{R}$

$$(A+B)_{ij} = (A)_{ij} + (B)_{ij}$$
$$(cA)_{ij} = c(A)_{ij}$$

Example:

$$\begin{bmatrix} 2 & 1 \\ 0 & 5 \\ 6 & -3 \end{bmatrix} + 7\begin{bmatrix} \pi & -1 \\ 6 & 3 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 2+7\pi & -6 \\ 42 & 26 \\ 6 & -3 \end{bmatrix}$$

Info — Matrix Properties

If $A, B, C \in M_{m\times n}(\mathbb{R}), c, d \in \mathbb{R}$ then

1. $A + B \in M_{m\times n}(\mathbb{R})$
2. $(A+B)+C = A+(B+C)$
3. $A + B = B + A$
4. There exists a matrix $O_{m,n} \in M_{m\times n}(\mathbb{R})$, s.t. $A + O_{m,n} = A\ \forall A$
5. For every $A \in M_{m\times n}(\mathbb{R}), \exists(-A) \in M_{m\times n}(\mathbb{R})$ s.t. $A + (-A) = O_{m,n}$
6. $cA = M_{m\times n}(\mathbb{R})$
7. $c(dA) = (cd)A$
8. $(c+d)A = cA + dA$
9. $c(A+B) = cA + cB$
10. $1A = A$

Info – Transpose

The **transpose** of a $m \times n$ matrix A is the $n \times m$ matrix A^T whose ij-th entry is the ji-th entry of A. $(A^T)_{ij} = (A)_{ji}$

If $A, B \in M_{m\times n}(\mathbb{R}), c \in \mathbb{R}$

1. $(A^T)^T = A$
2. $(A+B)^T = A^T + B^T$
3. $(cA)^T = cA^T$

Proof

$((cA)^T)_{ij} = (cA)_{ji} = c(A)_{ji} = c(A^T)_{ij} = (cA^T)_{ij}$

□

Note: we can regard vectors in $\mathbb{R}^n$ as matrices in $M_{n\times 1}(\mathbb{R})$

Examples:

1. $A = \begin{bmatrix} 2 & 1 \\ 0 & 5 \\ 6 & -3 \end{bmatrix}$

   $A^T = \begin{bmatrix} 2 & 0 & 6 \\ 1 & 5 & -3 \end{bmatrix}$
2. $\vec{x} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}$

   $\vec{x}^T = \begin{bmatrix} 2 & 1 & 3 \end{bmatrix}$

Matrix Multiplications

Note that

- $A = \begin{bmatrix} \overrightarrow{a_1}^T \\ \overrightarrow{a_2}^T \\ \dots \\ \overrightarrow{a_m}^T \end{bmatrix}$
- $\vec{a_i} = \vec{x} = \begin{bmatrix} a_{i1} \\ a_{i2} \\ \dots \\ 1_{in} \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \\ \dots \\ x_n \end{bmatrix} = a_{i1}x_1 + a_{i2}x_2 + \dots + a_{in}x_n$

Thus we can the linear system with augmented matrix $\left[A \mid \vec{b}\right]$ as $\begin{bmatrix} \overrightarrow{a_1}\cdot\vec{x} \\ \overrightarrow{a_2}\cdot\vec{x} \\ \dots \\ \overrightarrow{a_m}\cdot\vec{x} \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \dots \\ b_m \end{bmatrix}$

Info — Matrix-Vector Multiplication

1. Let A be an $m \times n$ matrix whose rows are denoted $\vec{a}_i^T$ for $1 \leq i \leq m$. For any $\vec{x} \in \mathbb{R}^n$

$$A\vec{x} = \begin{bmatrix} \vec{a}_1 \cdot \vec{x} \\ \vec{a}_2 \cdot \vec{x} \\ \dots \\ \vec{a}_m \cdot \vec{x} \end{bmatrix}$$

2. Let $A = [\vec{a}_1, \vec{a}_2, ..., \vec{a}_n]$ be $m \times n$ matrix , $\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \dots \\ x_n \end{bmatrix} \in \mathbb{R}^n$

$$A\vec{x} = \vec{a}_1 x_1 + \vec{a}_2 x_2 + ... + \vec{a}_n x_n$$

Note that $\vec{a}_i$ refers to the i —th column of the matrix A

Properties

If $A \in M_{m\times n}(\mathbb{R}), \vec{x}, \vec{y} \in \mathbb{R}^n, c \in \mathbb{R}$

1. $A(\vec{x} + \vec{y}) = A\vec{x} + A\vec{y}$
2. $A(c\vec{x}) = c(A\vec{x}$

Examples:

1. Let $A = \begin{bmatrix} 2 & 1 \\ 5 & 7 \\ 0 & -3 \end{bmatrix}, \vec{x} = \begin{bmatrix} 3 \\ -1 \end{bmatrix}$

$$A\vec{x} = \begin{bmatrix} \begin{bmatrix} 2 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ -1 \end{bmatrix} \\ \begin{bmatrix} 5 \\ 7 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ -1 \end{bmatrix} \\ \begin{bmatrix} 0 \\ -3 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ -1 \end{bmatrix} \end{bmatrix} = \begin{bmatrix} 5 \\ 8 \\ 3 \end{bmatrix}$$

2. Compute E1 in another way:

$$3\begin{bmatrix} 2 \\ 5 \\ 0 \end{bmatrix} - 1\begin{bmatrix} 1 \\ 7 \\ -3 \end{bmatrix} = \begin{bmatrix} 5 \\ 8 \\ 3 \end{bmatrix}$$

3. Let $A = \begin{bmatrix} 2 & 2 & 7 \\ 1 & 6 & 3 \\ 4 & 0 & 3 \\ 0 & 0 & 1 \end{bmatrix}, \vec{x} = \begin{bmatrix} 4 \\ 2 \\ -3 \end{bmatrix}$

$$A\vec{x} = \begin{bmatrix} -9 \\ -1 \\ 7 \\ -3 \end{bmatrix}$$

Theorem — Column Extraction Theorem

If $\vec{e}_i$ is the i-th standard basis and $A = [\vec{a_1} \ \dots \ \vec{a_n}]$ then

$$A\vec{e}_i = \vec{a}_i$$

Example:

Let A be the coefficient matrix of the homogeneous linear system $\left[A \mid \vec{0}\right]$ or $A\vec{x} = \vec{0}$

Let S be the solution set. Prove that S is a subspace of $\mathbb{R}^n$

Proof

1. $\vec{x} = \vec{0}$ is a solution to $\left[A \mid \vec{0}\right], \vec{0} \in S$
2. For $\vec{x}, \vec{y} \in S$, we have $A\vec{x} = \vec{0}, A\vec{y} = \vec{0}$. Then $A(\vec{x} + \vec{y}) = A\vec{x} + A\vec{y} = \vec{0} + \vec{0} = \vec{0} \in S$
3. For $\vec{x} \in S, c \in \mathbb{R}$, $A(c\vec{x}) = c(A\vec{x}) = c\vec{0} = \vec{0} \in S$

By subspace test, S is a subspace of $\mathbb{R}^n$

□

📐 **Theorem — Transpose - Dot Product Theorem**
If $\vec{x}, \vec{y} \in \mathbb{R}^n$

$$\vec{x}^T\vec{y} = \vec{x} \cdot \vec{y}$$

🐳 **Info — Matrix-Marix Multiplication**

Let A be a $m \times n$ matrix, B be $n \times p$ matrix. Then AB is the $m \times p$ matrix defined by

$$AB = \left[A\vec{b}_1 \ A\vec{b}_2 \ \dots \ A\vec{b}_p\right]$$

where $B = \left[\vec{b}_1 \ \vec{b}_2 \ \dots \ \vec{b}_p\right], A = \left[\vec{a}_1 \ \vec{a}_2 \ \dots \ \vec{a_p}\right]$

Consider the ijth entry of AB, that is the entry of AB in the i-th row and j-th column, that is, $(AB)_{ij}$

1. The j-th column of AB is $A\vec{b}_j$, and thus its i-th entry will be the ij-th entry of AB
2. If we write $A = \begin{bmatrix} \vec{a}_1^T \\ \vec{a}_2^T \\ \dots \\ \vec{a}_m^T \end{bmatrix}$, then $A\vec{b}_j = \begin{bmatrix} \vec{a}_1^T \\ \vec{a}_2^T \\ \dots \\ \vec{a}_m^T \end{bmatrix} \vec{b}_j = \begin{bmatrix} \vec{a}_1 \cdot \vec{b}_1 \\ \vec{a}_2 \cdot \vec{b}_2 \\ \dots \\ \vec{a}_m \cdot \vec{b}_j \end{bmatrix}$

   Then the i-th entry of this vector is $(AB)_{ij} = \vec{a}_i \cdot \vec{b}_j$

Properties

If A, B, C be matrices of the correct size so that the required products are defined, $t \in \mathbb{R}$,

1. $A(B + C) = AB + AC$
2. $(A + B)C = AC + BC$
3. $t(AB) = (tA)(B) = A(tB)$
4. $A(BC) = (ABC)$
5. $(AB)^T = B^T A^T$

Note that AB is not BA generally speaking. There might be special cases.

Examples:

1. Let $A = \begin{bmatrix} 2 & 1 \\ 1 & -2 \\ 3 & 4 \end{bmatrix}, B = \begin{bmatrix} 3 & 5 & 6 \\ 0 & 1 & -1 \end{bmatrix}$

   Then $AB = \begin{bmatrix} 2 & 1 \\ 1 & -2 \\ 3 & 4 \end{bmatrix} \begin{bmatrix} 3 & 5 & 6 \\ 0 & 1 & -1 \end{bmatrix} = \begin{bmatrix} (2)(3)+(1)(0) & (2)(5)+(1)(1) & (2)(6)+(1)(-1) \\ (1)(3)+(-2)(0) & (1)(5)+(-2)(1) & (1)(6)+(-2)(-1) \\ (3)(3)+(4)(0) & (3)(5)+(4)(1) & (3)(5)+(4)(-1) \end{bmatrix} = \begin{bmatrix} 6 & 11 & 11 \\ 3 & 4 & 8 \\ 9 & 19 & 14 \end{bmatrix}$

   $BA = \begin{bmatrix} 3 & 5 & 6 \\ 0 & 1 & -1 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ 1 & -2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} (3)(2)+(5)(1)+(6)(3) & (3)(1)+(5)(-2)+(6)(4) \\ (0)(2)+(1)(1)+(-1)(3) & (0)(1)+(-1)(-2)+(1)(4) \end{bmatrix} = \begin{bmatrix} 29 & 17 \\ -2 & -6 \end{bmatrix}$

2. Let$A = \begin{bmatrix} 5 & 1 \\ 2 & 6 \end{bmatrix}, B = \begin{bmatrix} 0 & 1 \\ -1 & 3 \end{bmatrix}$

   Then $AB = \begin{bmatrix} 5 & 1 \\ 2 & 6 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ -1 & 3 \end{bmatrix} = \begin{bmatrix} -1 & 8 \\ -6 & 20 \end{bmatrix}$

   $BA = \begin{bmatrix} 0 & 1 \\ -1 & 3 \end{bmatrix} \begin{bmatrix} 5 & 1 \\ 2 & 6 \end{bmatrix} = \begin{bmatrix} 2 & 6 \\ 1 & 17 \end{bmatrix}$

📐 **Theorem — Matrix Equality Theorem**
If A and B are $m \times n$ matrices s.t. $A\vec{x} = B\vec{x}$ for every $\vec{x} \in \mathbb{R}^n$, then $A = B$

Proof

If $A\vec{x} = B\vec{x},\ \forall \vec{x} \in \mathbb{R}^n$, then $A\vec{e}_i = B\vec{e}_i,\ \forall 1 \leq i \leq n$.

By Column Extraction Theorem, $A\vec{e}_i = \vec{a}_i, B\vec{e}_i = \vec{b}_i$ where $\vec{a}_i = \vec{b}_i$, are also the i-th column of A and B respectively. Hence $A = B$

□

🐳 **Info — Identity Matrix**
The $n \times n$ **identity matrix**, denoted by I or I_n, is the matrix s.t. $(I)_{ii} = 1$ for $1 \leq i \leq n, (I)_{ij} = 0, \forall i \neq j$.

$$I = [\vec{e}_1\ \ldots\ \vec{e}_n] = \begin{bmatrix} 1 & 0 & 0 & \ldots & 0 \\ 0 & 1 & 0 & \ldots & 0 \\ \ldots & \ldots & \ldots & \ldots & \ldots \\ 0 & 0 & 0 & \ldots & 1 \end{bmatrix}$$

If $I = [\vec{e}_1\ \ldots\ \vec{e}_n]$, thenfor any $n \times n$ matrix A:

$$AI = A = IA$$

Example:

$\begin{bmatrix} 6 & 0 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 6 & 0 \\ 1 & 2 \end{bmatrix}$

Linear Mappngs

🐳 **Info — Matrix Mappings**

If A is an $m \times n$ matrix, then we define a function $f : \mathbb{R}^n \to \mathbb{R}^m$ by $f(\vec{x}) = A\vec{x}$ called a matrix mapping.

Note that we can write $f(x_1, ..., x_n) = (y_1, ..., y_m)$

Example:

Let $A = \begin{bmatrix} 6 & 0 \\ 1 & 2 \end{bmatrix}$. Then the matrix mapping $f : \mathbb{R}^n \to \mathbb{R}^m$, given by $f(\vec{x}) = A\vec{x}$ can be expressed as $f(\vec{x}) = f\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = \begin{bmatrix} 6 & 0 \\ 1 & 2 \end{bmatrix}\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 6x_1 \\ x_1+2x_2 \end{bmatrix}$

Info — Distrbutive Matrix Property

If A is an $m \times n$ matrix and $f : \mathbb{R}^n \to \mathbb{R}^m$ is defined by $f(\vec{x}) = A\vec{x}$, then $\forall \vec{x}, \vec{y} \in \mathbb{R}^n$ and $b, c \in \mathbb{R}$ we have

$$f(b\vec{x} + c\vec{y}) = bf(\vec{x}) + cf(\vec{y})$$

Info — Linear Mapping

A function $L : \mathbb{R}^n \to \mathbb{R}^m$ is said to be **linear mapping** if for every $\vec{x}, \vec{y} \in \mathbb{R}^n; b, c \in \mathbb{R}$ we have

$$L(b\vec{x} + c\vec{y}) = bL(\vec{x}) + cL(\vec{y})$$

Two linear mappings $L : \mathbb{R}^n \to \mathbb{R}^m$ and $M : \mathbb{R}^n \to \mathbb{R}^m$ are said to be **equal** if $L(\vec{x}) = M(\vec{x}),\ \forall \vec{x} \in \mathbb{R}^n$, that is $L = M$.

A linear mapping $L : \mathbb{R}^n \to \mathbb{R}^n$ is somtimes called a **linear operator**. This is often done when we wish to stress the fact that the domain and codomain of the linear mapping are thhe same.

If $L : \mathbb{R}^n \to \mathbb{R}^m$ is a linear mapping, then $L\left(\vec{0}\right) = \vec{0}$

Tip — Linear Mapping and Matrix Mapping

Every linear mapping can be represented as matrix mapping

Every matrix mapping is a linear mapping

Observe: for an arbitrary $\vec{x} \in \mathbb{R}^n$, we can write $\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \dots \\ x_n \end{bmatrix} = x_1\vec{e}_1 + x_2\vec{e}_2 + \dots + x_n\vec{e}_n$.

$$\begin{aligned} L(\vec{x}) &= x_1\vec{e}_1 + x_2\vec{e}_2 + \dots + x_n\vec{e}_n \\ &= x_1L(\vec{e}_1) + x_2L(\vec{e}_2) + \dots + x_nL(\vec{e}_n) \\ &= [L(\vec{e}_1)\ L(\vec{e}_2)\ \dots\ L(\vec{e}_n)]\begin{bmatrix} x_1 \\ x_2 \\ \dots \\ x_n \end{bmatrix} \\ &= [L]\vec{x}] \end{aligned}$$

Let $L : \mathbb{R}^n \to \mathbb{R}^m$ be a linear mapping, then

Example:

Let $L : \mathbb{R}^2 \to \mathbb{R}^2$ be defined by $L\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = \begin{bmatrix} x_2 \\ -x_1 \end{bmatrix}$. Is L linear.

Let $\vec{x}, \vec{y} \in \mathbb{R}^n; b, c \in \mathbb{R}$. Then $L(b\vec{x} + c(\vec{y})) = L\left(\begin{bmatrix} bx_1+cy_1 \\ bx_2+cy_2 \end{bmatrix}\right) = \begin{bmatrix} bx_2+cy_2 \\ -bx_1-cy_1 \end{bmatrix} = b\begin{bmatrix} x_2 \\ -x_1 \end{bmatrix} + c\begin{bmatrix} y_2 \\ -y_1 \end{bmatrix} = bL(\vec{x}) + cL(\vec{y})$.

Hence L is linear.

Info — Standard Matrix

Every linear mapping $L : \mathbb{R}^n \to \mathbb{R}^m$ can be represented as a matrix mapping with matrix whose i-th column is the image of the i-th standard basis vector of $\mathbb{R}^n$ under L for all $1 \leq i \leq n$. That is, $L(\vec{x}) = [L]\vec{x}$ where

$$[L] = [L(\vec{e}_1) \ \ldots \ L(\vec{e}_n)]$$

Let $L : \mathbb{R}^n \to \mathbb{R}^m$ a linear mapping. The matrix $[L] = [L(\vec{e}_1) \ \ldots \ L(\vec{e}_n)]$ is called the **standard matrix** of L. It satisfies $L(\vec{x}) = [L]\vec{x}$

Example:

Let $L : \mathbb{R}^3 \to \mathbb{R}^3$ be the mapping given by $L(\vec{x}) = \text{proj}_{\vec{a}}(\vec{x}), \vec{a} = \begin{bmatrix} 1 \\ 3 \\ 6 \end{bmatrix}$. Find $[L]$

$L(\vec{e}_1) = \text{proj}_{\vec{a}}(\vec{e_1}) = \frac{\vec{a}\cdot\vec{e}_1}{\|\vec{a}\|^2}\vec{a} = \frac{1}{46}\begin{bmatrix} 1 \\ 3 \\ 6 \end{bmatrix}$

$L(\vec{e}_2) = \text{proj}_{\vec{a}}(\vec{e_2}) = \frac{\vec{a}\cdot\vec{e}_2}{\|\vec{a}\|^2}\vec{a} = \frac{3}{46}\begin{bmatrix} 1 \\ 3 \\ 6 \end{bmatrix}$

$L(\vec{e}_3) = \text{proj}_{\vec{a}}(\vec{e_3}) = \frac{\vec{a}\cdot\vec{e}_3}{\|\vec{a}\|^2}\vec{a} = \frac{6}{46}\begin{bmatrix} 1 \\ 3 \\ 6 \end{bmatrix}$

Therefore $L = \frac{1}{46}\begin{bmatrix} 1 & 3 & 6 \\ 3 & 9 & 18 \\ 6 & 18 & 36 \end{bmatrix}$

Rotation


Info — Rotations in $\mathbb{R}^2$

Let $R_\theta = \mathbb{R}^2 \to \mathbb{R}^2$ denote the funciton that rotates a vector $\vec{x} \in \mathbb{R}^2$ about the orgin through an angle θ. We can show:

$$R_\theta(x_1, x_2) = (x_1 \cos\theta - x_2 \sin\theta, x_1 \sin\theta +_2 \cos\theta)$$

We can get the standard matrix of R_θ:

$$R_\theta = \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix}$$



If $R_\theta : \mathbb{R}^2 \to \mathbb{R}^2$ is a rotation with rotation matrix $A = [R_\theta]$, then column of A are orthogonal unit vectors.

Example:

$\vec{x} = \begin{bmatrix} 4 \\ 2 \end{bmatrix}$ rotated about the origine by angle $\theta = \frac{\pi}{6}$


$R_{\frac{\pi}{6}}(\vec{x}) = \begin{bmatrix} \cos\frac{\pi}{6} & -\sin\frac{\pi}{6} \\ \sin\frac{\pi}{6} & \cos\frac{\pi}{6} \end{bmatrix} \begin{bmatrix} 4 \\ 2 \end{bmatrix} = \begin{bmatrix} \sqrt{3} & -\frac{1}{2} \\ \frac{1}{2} & \sqrt{3} \end{bmatrix} \begin{bmatrix} 4 \\ 2 \end{bmatrix} = \begin{bmatrix} 2\sqrt{3}-1 \\ 2+\sqrt{3} \end{bmatrix}$

Reflection

Info — Reflections

Let $\text{refl}_P : \mathbb{R}^n \to \mathbb{R}^n$ denote the mapping that sends a vector $\vec{x}$ to its mirror image in the hyperplane P with normal vector $\vec{n}$.

$$\text{refl}_P(\vec{x}) = \vec{x} - 2\text{proj}_{\vec{n}}(\vec{x})$$



Special Subspaces

Info — Range

Let $L : \mathbb{R}^n \to \mathbb{R}^n$ be a linear mapping. The **range** of L is defined by

$$\text{Range}(L) = \{L(\vec{x}) \in \mathbb{R}^m \mid \vec{x} \in \mathbb{R}^n\}$$

If $L : \mathbb{R}^n \to \mathbb{R}^m$ is a linear mapping, then $\text{Range}(L)$ is a subspace of $\mathbb{R}^m$

Example:

Let $L : \mathbb{R}^2 \to \mathbb{R}^3$ be given by $L(x_1, x_2) = (x_1 + 2x_2, x_2 - x_1, 0)$

It is linear.

1. Is $\vec{v} = \begin{bmatrix} 2 \\ 2 \\ 0 \end{bmatrix} \in \text{Range}(L)$? (i.e. Does there exist $\vec{x} \in \mathbb{R}^2$ s.t. $\vec{v} = L(\vec{x})$?)

   Set $\begin{bmatrix} 2 \\ 2 \\ 0 \end{bmatrix} = \begin{bmatrix} x_1+2x_2 \\ x_2-x_1 \\ 0 \end{bmatrix} \implies \begin{cases} 2=x_1+2x_2 \\ 2=x_2-x_1 \end{cases} \implies \left[\begin{array}{cc|c} 1 & 2 & 2 \\ -1 & 1 & 2 \\ 0 & 0 & 0 \end{array}\right] \sim \left[\begin{array}{cc|c} 1 & 0 & -\frac{2}{3} \\ 0 & 1 & \frac{4}{3} \\ 0 & 0 & 0 \end{array}\right]$

   Therefore there exists $\vec{x} \in \mathbb{R}^2$ s.t. $L(\vec{x}) = \vec{v} \implies \vec{v} \in \text{Range}(L)$

2. Is $\vec{u} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix} \in \text{Range}(L)$

   No, because $3 \neq 0$

Info — Onto(Surjective)

A linear mapping $L : \mathbb{R}^n \to \mathbb{R}^m$ is called **onto** if $\text{Range}(L) = \mathbb{R}^m$

We can say that L is not onto since $\vec{u} \notin \text{Range}(L)$

Kernel

> **Info — Kernel**
>
> Let $L : \mathbb{R}^n \to \mathbb{R}^m$ be a linear mapping. The **kernel** (nullspace) of L is the set of all vectors in the domain which are mapped to the zero vector in the codomain.
>
> $$\text{Ker}(L) = \left\{\vec{x} \in \mathbb{R}^n \mid L(\vec{x}) = \vec{0}\right\}$$
>
> If $L : \mathbb{R}^n \to \mathbb{R}^m$ is a linear mapping, then $\text{Ker}(L)$ is a subspace of $\mathbb{R}^n$
>
> A linear mapping L is called **one-to-one** if $\text{Ker}(L) = \left\{\vec{0}\right\}$

Example:

1. Let $L : \mathbb{R}^2 \to \mathbb{R}^3$ be given by $L(x_1, x_2) = (x_1 + 2x_2, x_2 - x_1, 0)$

   Find $\text{Ker}(L)$

   We set $\begin{bmatrix}0\\0\\0\end{bmatrix} = \begin{bmatrix}x_1+2x_2\\-x_1+x_2\\0\end{bmatrix} \implies \left[\begin{array}{cc|c}1&2&0\\-1&1&0\\0&0&0\end{array}\right] \sim \left[\begin{array}{cc|c}1&0&0\\0&1&0\\0&0&0\end{array}\right]$

$\implies x_1 = 0, x_2 = 0$, so the only solution, is $\text{Ker}(L) = \vec{0}$

2. Let $M(x, x_2) = (x_1 + 2x_2, 2x_1 + 4x_2, 0)$

   $\left[\begin{array}{cc|c}1&2&0\\2&4&0\\0&0&0\end{array}\right] \sim \left[\begin{array}{cc|c}1&2&0\\0&0&0\\0&0&0\end{array}\right] \implies \vec{x} = t\begin{bmatrix}-2\\1\end{bmatrix},\ t \in \mathbb{R}$

   $\implies \text{Ker}(M) = \left\{t\begin{bmatrix}-2\\1\end{bmatrix} \mid t \in \mathbb{R}\right\} = \text{Span}\left\{\begin{bmatrix}-2\\1\end{bmatrix}\right\} \implies \left\{\begin{bmatrix}-2\\1\end{bmatrix}\right\}$ is a basis for $\text{Ker}(M)$

3. A kernel can never be empty, it has at least $\vec{0}$

4. M from E2 is not "one-to-one", but L from E1 is "one-to-one"

> **Info — One-To-One(Injective)**
>
> Let $L : \mathbb{R}^n \to \mathbb{R}^m$ be a linear mapping. L is one-to-one if and only if for every $\vec{u}, \vec{v} \in \mathbb{R}^n$ s.t. $L(\vec{u}) = L(\vec{v})$, we must have $\vec{u} = \vec{v}$
>
> If a mapping is both injective and surjective, it is bijective

> **Info — The Column Space and Nullspace of a Matrix**
>
> Let $L : \mathbb{R}^n \to \mathbb{R}^m$ be a linear mapping with standard marix $[L]$. Then, $\vec{x} \in \text{Ker}(L)$ if and only if $[L]\vec{x} = \vec{0}$
>
> Let $A \in M_{m\times n}(\mathbb{R})$. The set $\left\{\vec{x} \in \mathbb{R}^n \mid A\vec{x} = \vec{0}\right\}$ is a subspace of $\mathbb{R}^n$

Info — Nullspace

1. Let A be an $m \times n$ matrix. The **nullspace** (kernel) of A is defined by
$$\text{Null}(A) = \left\{\vec{x} \in \mathbb{R}^n \mid A\vec{x} = \vec{0}\right\}$$
2. Let A be an $m \times n$ matrix. Suppose that the vector equation of the solution set of $A\vec{x} = \vec{0}$ as determined by the Gauss-Jordan algorithm is given by
$$\vec{x} = t_1\vec{v}_1 + t_2\vec{v}_2 + ... + t_k\vec{v}_k, \quad t_1, ..., t_k \in \mathbb{R}$$
Then $\{\vec{v}_1, ..., \vec{v}_k\}$ is a basis for $\text{Null}(A)$
3. If A be an $m \times n$ matrix, then
$$\dim \text{Null}(A) = n - \text{rank}(A)$$
4. Let A be an $m \times n$ matrix. The **nullity** of A is defined by
$$\text{nullity } A = \dim \text{Null}(A)$$

If $L : \mathbb{R}^n \to \mathbb{R}^m$ is a linear mapping with standard matrix $[L] = A = [\vec{a_1} \ ... \ \vec{a}_n]$, then
$$\text{Range}(L) = \text{Span}\{\vec{a}_1, ..., \vec{a}_n\}$$

Info — Column Space

Let $A = [\vec{a}_1 \ ... \ \vec{a}_n] \in M_{m\times n}(\mathbb{R})$. The **column space** of A, denoted by $\text{Col}(A)$, is the subspace of $\mathbb{R}^m$ spanned by the columns of A. That is,
$$\text{Col}(A) = \text{Span}\{\vec{a}_1, ..., \vec{a}_n\}$$

If $L : \mathbb{R}^n \to \mathbb{R}^m$ bis a linear mapping with standard matrix $[L]. = A[\vec{a}_1 \ ... \ \overrightarrow{a_n}]$, then
$$\text{Range}(L) = \text{Col}([L])$$

Let $A = [\vec{a}_1 \ ... \ \vec{a}_n]$ be an $m \times n$ matrix. Suppose that $\text{rank}A = r$ and that the RREF of A has leading ones in columns $j_1, ..., j_r$. Then $\left\{\vec{a}_{j_1}, ..., \vec{a}_{j_r}\right\}$ is a basis for $\text{Col}(A)$
$$\dim \text{Col}(A) = \text{rank}(A)$$

Let A be a $m \times n$ matrix and let $\vec{b} \in \mathbb{R}^m$. Then $\vec{b} \in \text{Col}(A)$ if and only if the system $\left[A \mid \vec{b}\right]$ is consistent.

Example:

Let $A = \begin{bmatrix} 1 & 2 & 0 \\ 3 & 1 & 2 \end{bmatrix}$

1. Find a basis for $\text{Null}(A)$

$\begin{bmatrix} 1 & 2 & 0 \\ 3 & 1 & 2 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & \frac{4}{5} \\ 0 & 1 & -\frac{2}{5} \end{bmatrix} \implies x_1 = -\frac{4}{5}x_3, x_2 = \frac{2}{5}x_3 \implies t = x_3 \implies \vec{x} = \begin{bmatrix} -\frac{4}{5} \\ \frac{2}{5} \\ 1 \end{bmatrix} t, t \in \mathbb{R}$

$B = \left\{ \begin{bmatrix} -\frac{4}{5} \\ \frac{2}{5} \\ 1 \end{bmatrix} \right\}$ is a basis for $\text{Null}(A)$, nullity of $A = 1$

2. Find a basis for $\text{Col}(A)$

$\begin{bmatrix}1 & 2 & 0\\ 3 & 1 & 2\end{bmatrix} \sim \begin{bmatrix}1 & 0 & \frac{4}{5}\\ 0 & 1 & -\frac{2}{5}\end{bmatrix}$ Columns 1 and 2 of A forms a basis for $\text{Col}(A) : e = \left\{\begin{bmatrix}1\\3\end{bmatrix}, \begin{bmatrix}2\\1\end{bmatrix}\right\}$ is a basis.

📐 **Theorem — Rank-Nullity Theorem**

If A is a $m \times n$ matrix, then

$$\text{rank}(A) + \text{nullity}(A) = n$$

📐 **Theorem** — Theorem

Let A be a $m \times n$ matrix

rank A = rank A^T

Operations on Linear Mappings

🐳 **Info — Operations on Linear Mappings**

Addition and Scalar Multiplication

Let $L, M \in \mathbb{L}(\mathbb{R}^n, \mathbb{R}^m)$. We define $L + M : \mathbb{R}^n \to \mathbb{R}^m$ and $cL : \mathbb{R}^n \to \mathbb{R}^m, c \in \mathbb{R}$

$$(L + M)(\vec{x}) = L(\vec{x}) + M(\vec{x})$$

$$(cL)(\vec{x}) = cL(\vec{x})$$

Composition

Let $L : \mathbb{R}^n \to \mathbb{R}^m$ and $M : \mathbb{R}^m \to \mathbb{R}^p$ be linear mappings. The **composition** of M, L is the function $M \circ L : \mathbb{R}^n \to \mathbb{R}^p$

$$(M \circ L)(\vec{x}) = M(L(\vec{x})) = ([M][L])\vec{x} \implies [M \circ L] = [M][L]$$

🐳 **Info — Identity Mapping**

The linear mapping $\text{Id} : \mathbb{R}^n \to \mathbb{R}^m$ defined by $\text{Id}(\vec{x}) = \vec{x}$