CH 2 — Differential Equations

Luke Lu • 2026-10-19

Introduction to Differential Equations

> **Info — Differential Equations**
>
> A differential equation (DE), is an equation involving an unknown function and its derivatives. The term ordinary differential equation (ODE) refers to a differential equation involving single-variable functions, whereas the term partial differential equation (PDE) refers to a differential equation involving multivariable functions (i.e., functions with multiple inputs).
>
> An ODE is expressed
>
> $$F(x, y, y', y'', ..., y^n) = 0$$
>
> for some $n \in \mathbb{N}$
>
> The order of a DE is the order of the **highest derivative** that appears in the equation.
>
> A function $y = \varphi(x)$ is a solution to the differential equation $F(x, y, y', y'', ..., y^n) = 0$ if
>
> $$F(x, \varphi(x), \varphi'(x), \varphi''(x), ..., \varphi^n(x)) = 0$$
>
> The graph of a solution to a DE is called a **solution curve**
>
> The complete collection of solutions to a DE, including any arbitrary constants, is called its general solution. A particular solution to a DE is one in which all arbitrary constants have been specified.
>
> A differential equation together with one or more initial conditions is known as an initial value problem (IVP)

Examples:

1. First order differential equation $\frac{\mathrm{d}y}{\mathrm{d}x} = x + y$
2. Second order differential equation $\frac{\mathrm{d}y}{\mathrm{d}x} + x^2\frac{\mathrm{d}^2y}{\mathrm{d}x^2} = \ln x + y$
3. Solve $\frac{\mathrm{d}y}{\mathrm{d}x} = \sin x \implies y = -\cos x + C, c \in \mathbb{R}$
4. $\frac{\mathrm{d}y}{\mathrm{d}x} = \sin x, y(0) = 2 \implies -\cos x + C \implies -\cos x + 3$
5. $y' = x + y$

   $y = -1 - x + 2e^x \implies y' = -1 + 2e^x \implies (x + y) = x + (-1 - x + 2e^x) = -1 + 2e^x = y'$
6. $y = -1 - x - 5e^x$

   $y' = -1 - 5e^x \implies x + (-1 - x - 5e^x) = -1 - 5e^x = y'$
7. $y = -5 - x + 2e^x$

   $y' = -1 + 2e^x \implies x + (-5 - x + 2e^x) = -5 + 2e^x \neq y'$

Thus $y = -1 - x + Ce^x$, for $C \in \mathbb{R}$ is always a solution

8. Determine all real numbers k s.t. $x(t) = \sin(kt)$ is a solution to the second-order differential equation $\frac{\mathrm{d}^2y}{\mathrm{d}x^2} = -2x$

$$\begin{cases} x'(t) = k\cos(kt) \\ x''(t) = -k^2\sin(kt) \end{cases} \implies -k^2 \sin kt = 02\sin kt$$

$$(k^2 - 2)\sin(kt) = 0$$

$$k = \pm\sqrt{2}, k = 0$$


Direction Fields

> **Info — Direction Field**
>
> A direction field for the different equation $y' = F(x)$ displays short line segments of slope $F(x, y)$ at various points in the Cartesian plane

> **Tip** — Direction field plotter for DE: https://www.desmos.com/calculator/p7vd3cdmei

Examples:

1. $\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{x}{y}$




We can obtain this when plugging-in numbers.


At $(-2, 1)$, $\left.\frac{\mathrm{d}y}{\mathrm{d}x}\right|_{(x,y)=(-2,1)} = \left.\frac{x}{y}\right|_{(x,y)=(-2,1)} \implies -\frac{2}{1} = -2$

Hence a slope of -2 at $(-2, 1)$ and other points have the same method


2. $y'x = x + y$



3. $y' = xy$



4. $y' = y$



Separable Differential Equations

> **Info — Separable Differential Equation**
>
> A first order differential equation is said to be separable if it is written in format
>
> $$\frac{\mathrm{d}y}{\mathrm{d}x} = g(x)h(y)$$

Examples:

1. $y' = \sin x$
2. $y' = \frac{x}{y}$
3. $y' = 5y$
4. $y^2 y' = 2y + xy$
5. $y' = x + y$ is not separable but linear, see in Linear First Order DEs

💡 **Tip — Solving Separable DE**

We have two cases

1. Determine any solution y with $h(y) = 0$
2. Find the solutions y where $h(y) \not\equiv 0$ by evaluating

$$\int \frac{1}{h(y)}\,\mathrm{d}y = \int g(x)\,\mathrm{d}x$$

   If possible, isolate y as a function of x in the resulting equation.

The general solution is the collection of all solutions obtained in Case 1 together with all solutions obtained in Case 2.

Examples:

1. $y' = y$

   Case 1: $y = 0$

   Case 2:

$$\begin{aligned}\int \frac{1}{y}\,\mathrm{d}y &= \int 1\,\mathrm{d}x \\ \ln|y| &= x + C_1 \\ |y| &= e^{x+C_1} y = \pm e^{C_1} e^x = \pm C e^x,\ C \in \mathbb{R}\end{aligned}$$

   The general solution is $y = Ae^x,\ A \in \mathbb{R}$

2. $y' = \frac{x}{y}$

   Case 1: $h(y) = \frac{1}{y} \neq 0$

   Case 2:

$$\begin{aligned}\int y\,\mathrm{d}y &= \int x\,\mathrm{d}x \\ \frac{y^2}{2} &= \frac{x^2}{2} + C \\ y^2 &= x^2 + 2C \\ y &= \pm\sqrt{x^2 + 2C}\end{aligned}$$

   The general solution $y = \pm\sqrt{x^2 + D},\ D \in \mathbb{R}$

3. $y' = xy^2, y(0) = 3$

   Case 1: $h(y) = 0 \Longrightarrow y \equiv 0$

   Case 2:

$$\int \frac{1}{y^2}\,\mathrm{d}y = \int x\,\mathrm{d}x \Longrightarrow -\frac{1}{y} = \frac{x^2}{2} + C \Longrightarrow y = -2\frac{2}{x^2 + 2C}$$

   The general solution is $y \equiv 0$ or $y = \frac{-2}{x^2+D},\ D \in \mathbb{R}$

   For $y(0) = 3 \Longrightarrow -\frac{2}{D} = 3 \Longrightarrow D = -\frac{2}{3}$ ==> the unique solution is $y = \frac{-2}{x^2-\frac{2}{3}}$

4. $y' - x^2 - y^2 - 2xy + 1 = 0 \implies y' = (x+y)^2 - 1 \overset{v=x+y}{\implies} v' = v^2$

   Case 1: $h(v) = v^2 \equiv 0 \implies v \equiv 0$

   Case 2: $\int \frac{1}{v^2}\,\mathrm{d}v = x \implies -\frac{1}{v} = x + C \implies v = -\frac{1}{x+C}, C \in \mathbb{R}$

   The general solution $v \equiv 0$ or $v = -\frac{1}{x+C}, C \in \mathbb{R}$

   $y = -x$ or $y = -x - \frac{1}{x+C}, C \in \mathbb{R}$

Linear First-Order Differential Equations

> 🐳 **Info — First-Order Linear Differential Equation**
>
> First order Linear DE has the form
>
> $$A_0(x)y + A_1(x)y' = B(x) \quad A_1(x) \not\equiv 0$$
>
> Such an equation can be written inthe form $y' + P(x)y = Q(x)$ called the standard form.

> 🐳 **Info — Integrating Factor**
>
> Given a DE of the form $y' + P(x)y = Q(x)$,
>
> $$\mu = e^{\int P(x)\,\mathrm{d}x}$$
>
> is called the integrating factor for the DE.

> 💡 **Tip — Solving First-Order Linear DE**
>
> Given $A_1(x)y' + A_0(x)y = B(x)$
>
> 1. Divide by $A_1(x)$ to rewrite the DE in standard form
> 2. Multiply the equation by the integrating factor.
> 3. Rewrite LHS as $(\mu(x)y)'$
> 4. Integrate in respext to x
> 5. Isolate y

Example:

$$y' + \frac{3}{x}y = 1$$

$$\overset{\mu(x)=e^{3\ln(x)}}{\implies} \quad x^3y' + 3x^2y = x^3$$

$$\frac{\mathrm{d}}{\mathrm{d}x}x^3y = x^3$$

$$\int \frac{\mathrm{d}}{\mathrm{d}x}x^3y\,\mathrm{d}x = \int x^3\,\mathrm{d}x$$

1. $$y = \frac{1}{4}x + \frac{C}{x^3}$$

$y' = x + y \implies y' - y = x$

$$\overset{\mu(x)=e^{-x}}{\implies} \quad e^{-x}y' - e^{-x}y = e^{-x}x$$

$$\frac{\mathrm{d}}{\mathrm{d}x}e^{-x}y = xe^{-x}$$

2. $$y = -x - 1 + Ce^x, C \in \mathbb{R}$$

$$y' + 2xy = 1$$

$$\overset{\mu(x)=e^{x^2}}{\implies} \quad e^{x^2}y' + 2xye^{x^2} = e^{x^2}$$

$$\frac{\mathrm{d}}{\mathrm{d}x}e^{x^2}y = e^{x^2}$$

3. $$y = e^{-x^2}\int_0^1 e^{t^2}\,\mathrm{d}t + Ce^{-x^2}, C \in \mathbb{R}$$

Q5 CUTOFF

Applications

Mixing Problem

> 💡 **Tip — Formula - Mixing Problem**
>
> $$\frac{\mathrm{d}A}{\mathrm{d}t} = (\text{rate of substance in}) - (\text{rate of substance out})$$

Q1:

Suppose that a tank has 1000L of salt water at an initial concentration of 0.1kg/L. Salt water at a concentration of 0.3kg/L flows into tht tank at a rate of 10 L/min. The solution is kept mixed and drain out at a rate of 10 L/min.

Determine the amount of salt, in kg, in thte tank at time t

Let $A(t)$ represent the amount of salt(kg) in the tank at time t(min)

$\frac{\mathrm{d}A}{\mathrm{d}t} = \frac{\mathrm{d}s_i}{\mathrm{d}t} - \frac{\mathrm{d}s_o}{\mathrm{d}t}$ in kg/min

$\frac{\mathrm{d}s_i}{\mathrm{d}t} = 0.3 \cdot 10 = 3$kg/min

$\frac{\mathrm{d}s_0}{\mathrm{d}t} = \frac{A(t)}{1000} \cdot 10 = \frac{A}{100}$kg/min

$A(0) = 100$kg

$\frac{\mathrm{d}A}{\mathrm{d}t} = 3 - \frac{A}{100} = -\frac{A-300}{100} \implies \int \frac{1}{A-300}\,\mathrm{d}A = \int -\frac{1}{100}\,\mathrm{d}t \implies \ln|A-300| = -\frac{t}{100} + C$

$A = 300 \pm e^{-\frac{t}{100}+C}$

General solution $A = 300 + Be^{-\frac{t}{100}}, B \in \mathbb{R}$

$B = -200$ since $A(0) = 100$

Particular solution $A = 300 - 200e^{-\frac{t}{100}}$

Newton's Law of Heating/Cooling

> 💡 **Tip — Formula Newton's Law**
>
> $T(t)$ represents the temperature of an object at time t, and T_s is the constant temperature of tis surroundings, then theres exists a constant $k > 0$
>
> $$\frac{\mathrm{d}T}{\mathrm{d}t} = -k(T - T_s)$$
>
> The general solution to this DE is given
>
> $$T(t) = T_s + Ae^{-kt}, \ A \in \mathbb{R}$$
>
> If $T(0) > T_s$, the object will cool.
>
> An equality suggests no change.

$\frac{\mathrm{d}T}{\mathrm{d}t} = -k(T - T_s)$

1. Constant solution: $T \equiv T_s$
2. $\int \frac{1}{T-T_s}\,\mathrm{d}T = \int -k\,\mathrm{d}t \implies \ln|T - T_s| = -kt + C$
   $= T = \pm e^{-kt+C} + T_s = \pm e^{-kt}e^{C} + T_s = Ae^{-kt} + T_s, \ A \in \mathbb{R}$

General Solution:

$$T_s + Ae^{-kt}, \ A \in \mathbb{R}$$

Q2:

A cup of coffee has a temperature of 98°C in a room of 20°C. After 1 minute, the temperature of the coffee is 96°C. How long will it reach 80°C?

$T_s = 20, \exists k > 0$ s.t. $\frac{\mathrm{d}T}{\mathrm{d}t} = -k(T - 20)$ holds.

Initial conditions, $T(0) = 98, T(1) = 96$

The general solution will be $T = 20 + Ae^{-kt}$

With initial condition, we obtain $T = 20 + 78e^{\ln\frac{38}{39}t}$

$80 = 20 + 78e^{\ln(\frac{38}{39})t} \implies t = \frac{\ln(\frac{10}{13})}{\ln(\frac{38}{39})} \approx 10.1$ min

Population Growth

> 💡 **Tip — Exponential Growth/Decay**
>
> The general solution to the exponential growth DE $\frac{\mathrm{d}P}{\mathrm{d}t} = kP$ is given
>
> $$P(t) = Ae^{kt}, \ A = P(0)$$
>
> - $k > 0$ for growth
> - $k < 0$ for decay

💡 **Tip — Logistic Growth**

- If P is significantly smaller than M, $\frac{\mathrm{d}P}{\mathrm{d}t} \approx kP$
- If $P \approx M$, $\frac{\mathrm{d}P}{\mathrm{d}t} \approx 0$
- If $P > M$, $\frac{\mathrm{d}P}{\mathrm{d}t} < 0$

Formula

$$\frac{\mathrm{d}P}{\mathrm{d}t} = kP\left(1 - \frac{P}{M}\right)$$

General Solution:

$$P(t) = \frac{M}{1 + Ae^{-kt}},\ A \in \mathbb{R}$$

.

If $P_0 = P(0)$ is the initial population, then $A = \frac{M - P_0}{P_0}$

Q3:Scientists placed 100 geese in a nature preserve. After 1 year, the population increased to 150.

1. Assume that the population grows according to the law of natural growth.
   - Determine the population function, P (t).
   - After how many years will the population reach 500 geese?
2. Assume that the population grows according to the logistic model with a carrying capacity of 1500.
   - Determine the population function, P (t).
   - After how many years will the population reach 500 geese?

1. IVP: $\frac{\mathrm{d}P}{\mathrm{d}t} = kP, P(0) = 100, P(1) = 150$
   - The general solution is $P(t) = Ae^{kt}, A \in \mathbb{R}$

     $P(0) = Ae^0 = A = 100$

     $P(1) = 100e^k = 150 \Longrightarrow e^k = \frac{3}{2} \Longrightarrow k = \ln\frac{3}{2}$

     $P(t) = 100e^{\ln(\frac{3}{2})t} = 100\left(\frac{3}{2}\right)^t$
   - $P(t) = 100\left(\frac{3}{2}\right)^t = 500 \Longrightarrow t = \log_{\frac{3}{2}} 5 \approx 4.0$

     Therefore, the population reaches 500 after around 4 years.
2. IVP: $\frac{\mathrm{d}P}{\mathrm{d}t} = kP\left(1 - \frac{P}{1500}\right)$
   - The general solution is $P(t) = \frac{1500}{1 + Ae^{-kt}}, A \in \mathbb{R}$

     $A = \frac{M - P(0)}{P(0)} = 14$

     $P(1) = \frac{1500}{1 + 14e^{-k}} = 150 \Longrightarrow e^{-k} = \frac{9}{14} \Longrightarrow k = \ln\left(\frac{9}{14}\right)$

     Particular solution: $P(t) = \frac{1500}{1 + 14\left(\frac{9}{14}\right)^t}$
   - $P(t) = \frac{1500}{1 + 14\left(\frac{9}{14}\right)^t} = 500 \Longrightarrow t = \log_{\frac{9}{14}} \frac{1}{7} \approx 4.4$

   Therefore, the population reaches 500 after around 4.4 years